Prolonged XPO1 inhibition is essential for optimal anti-leukemic activity in *NPM1*-mutated AML

Giulia Pianigiani¹, Andrea Gagliardi¹, Federica Mezzasoma¹, Francesca Rocchio¹,
Valentina Tini¹, Barbara Bigerna¹, Paolo Sportoletti^{1,2}, Simona Caruso³, Andrea
Marra¹, Giulio Spinozzi¹, Sharon Shacham⁴, Yosef Landesman⁴, Concetta
Quintarelli^{3,5}, Franco Locatelli^{3,6}, Maria Paola Martelli^{1,2}, Brunangelo Falini^{1,2*},
Lorenzo Brunetti^{1,2*}

¹ Hematology, Department of Medicine and Surgery, Center for Hemato-Oncological
Research (CREO), University of Perugia, Perugia, Italy

² Institute of Hematology and Bone Marrow Transplantation, Santa Maria della
Misericordia Hospital, Perugia, Italy

³ Department of Hematology and Oncology, Cell and Gene Therapy, IRCCS
Bambino Gesù Children's Hospital, Rome, Italy

⁴ Karyopharm Therapeutics, Newton, MA, USA

⁵ Department of Translational Medical Sciences, Federico II University, Naples, Italy

⁶ Department of Gynecology/Obstetrics & Pediatrics, Sapienza University of Rome,
Rome, Italy

* Co-senior authors

Correspondence to: lorenzo.brunetti@unipg.it; brunangelo.falini@unipg.it

Text word count: 1244

Abstract word count: 195

Figures and tables: 2 figures

References: 22

Abstract

34 *NPM1* encodes for a nucleolar multifunctional protein and is the most frequently
 35 mutated gene in adult acute myeloid leukemia (AML). *NPM1* mutations cause the
 36 aberrant accumulation of mutant NPM1 (NPM1c) in the cytoplasm of leukemic cells,
 37 that is mediated by the nuclear exporter Exportin-1 (XPO1). Recent work has
 38 demonstrated that the interaction between NPM1c and XPO1 promotes high
 39 homeobox (HOX) genes expression, which is critical for maintaining the leukemic
 40 state of *NPM1*-mutated cells. However, the XPO1 inhibitor Selinexor administered
 41 once or twice/week in early-phase clinical trials did not translate into clinical benefit
 42 for *NPM1*-mutated AML patients. Here, we demonstrate that this dosing strategy
 43 results in only temporary disruption of the XPO1-NPM1c interaction and transient
 44 HOX genes downregulation, limiting the efficacy of Selinexor in the context of *NPM1*-
 45 mutated AML. Since second-generation XPO1 inhibitors can be administered more
 46 frequently, we compared intermittent (twice/week) versus prolonged (5 days/week)
 47 XPO1 inhibition in *NPM1*-mutated AML models. Integrating *in vitro* and *in vivo* data,
 48 we show that only prolonged XPO1 inhibition results in stable HOX downregulation,
 49 cell differentiation and remarkable anti-leukemic activity. This study lays the
 50 groundwork for the accurate design of clinical trials with second-generation XPO1
 51 inhibitors in *NPM1*-mutated AML.

52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67

Introduction

NPM1-mutated acute myeloid leukemia (AML) accounts for about one third of AML in adults^{1,2}. The most distinguishing feature of *NPM1*-mutated cells is the aberrant localization of mutant NPM1 (NPM1c) in the cytoplasm¹, caused by the loss of a nucleolar localization signal and the gain of a nuclear export signal within the C-terminal end of NPM1^{3,4}. Both NPM1c nuclear export and cytoplasmic accumulation are dependent on its interaction with the nuclear exporter Exportin-1 (XPO1)^{3,4}. Another unique property of *NPM1*-mutated AML is the high expression of homeobox (HOX) genes and their cofactors *MEIS1* and *PBX3* (hereafter referred as to HOX/MEIS)^{5,6}. We recently found that high HOX/MEIS levels are required to maintain the undifferentiated state of leukemic cells⁷ and that HOX/MEIS expression is directly dependent on the interaction between NPM1c and XPO1⁷.

The selective inhibitors of nuclear export Selinexor and Eltanexor covalently bind XPO1 and disrupt the interaction with its cargo proteins⁸, including NPM1c⁷. Preclinical studies have demonstrated that XPO1 inhibition cause NPM1c nuclear relocation, loss of HOX expression, differentiation, and growth arrest of *NPM1*-mutated cells^{7,9,10}. However, patients with *NPM1*-mutated AML showed suboptimal responses to Selinexor in early-phase clinical trials¹¹⁻¹⁴. As Selinexor has a half-life of 6 hours¹¹ and was administered once or twice/week, we hypothesized that intermittent dosing may not stably inhibit the NPM1c-XPO1 interaction, limiting its efficacy. In contrast, Eltanexor, currently tested in early-phase trials, can be administered more frequently (i.e. 5 days/week)¹⁵. Therefore, we asked whether prolonged XPO1 inhibition by Eltanexor could elicit a more pronounced anti-leukemic activity in *NPM1*-mutated cells.

Methods

Selinexor and Eltanexor were evaluated in cellular and animal models of *NPM1*-mutated AML. Parental OCI-AML3, NPM1c-GFP OCI-AML3 (in-frame knock-in of GFP at the NPM1c endogenous locus)⁷ and NPM1c-FKBP(F36V)-GFP OCI-AML3 (in-frame knock-in of FKBP(F36V) and GFP at the NPM1c endogenous locus)⁷ cells were used for *in vitro* experiments. Patient-derived xenograft (PDX) of two *NPM1*-mutated AML patients were used for *in vivo* and *in vitro* experiments. RNA-sequencing data were analyzed applying the ARPIR pipeline and are available at GEO (GSE181176)¹⁶. Detailed methods are provided in the supplemental materials.

Results and discussion

We first addressed the impact of intermittent XPO1 inhibition on the NPM1c-XPO1 interaction. As NPM1c subcellular localization is dependent on its binding to XPO1 (i.e. cytoplasmic when interacting with XPO1, nuclear when XPO1 is inhibited), we tracked the subcellular localization of endogenous NPM1c fused to GFP (NPM1c-GFP) in OCI-AML3 cells upon intermittent XPO1 inhibition. As expected, NPM1c-GFP was completely relocated to the nucleus after 12-hour Selinexor incubation. However, drug withdrawal caused cytoplasmic relocation over the following 24 hours, demonstrating quick recovery of the NPM1c-XPO1 interaction after transient XPO1 inhibition (Figure 1A).

As in *NPM1*-mutated cells high HOX/MEIS expression depends on the NPM1c-XPO1 interaction⁷, we hypothesized that early loss of XPO1 inhibition may result in inefficient HOX/MEIS downregulation. We determined *HOXA9*, *HOXA10* and *MEIS1* expression at 24 and 72 hours in cells treated with Selinexor for 24 hours (short treatment, ST) or for 72 hours (continuous treatment, CT). While ST caused only transient downregulation of HOX/MEIS expression, CT resulted in stable loss of these targets (supplemental Figure S1A). Next, to determine the impact of transient and stable XPO1 inhibition on the transcriptome, we performed RNA-sequencing in parental OCI-AML3 cells applying the same treatment strategy. After 24-hour incubation with Selinexor, the transcriptome was clearly perturbed, including downregulated HOX/MEIS (Figures 1B, supplemental Figures 1B, 1C, supplemental Table 1). However, drug withdrawal reduced transcriptional perturbation in the following 48 hours with only 57 residual differentially expressed genes (Figures 1B, 1C, supplemental Figures 1B, 1C, supplemental Table 1). Conversely, CT for 72 hours caused persistent downregulation of HOX/MEIS, combined with upregulation of genes related to myeloid differentiation and TP53 downstreams (Figures 1B, 1C, supplemental Figures 1C, 1D, supplemental Table 1). As XPO1 interacts with multiple cargo proteins⁸, to corroborate the hypothesis that the changes observed were mainly due to the loss of NPM1c-XPO1 interaction, we tested the effects of intermittent (2 days/week, e.g. Monday and Thursday) and prolonged (5 days/week, e.g. Monday to Friday) selective NPM1c degradation in CRISPR-engineered OCI-AML3 cells with endogenous NPM1c fused to the FKBP(F36V) degron tag and GFP⁷. Only prolonged NPM1c degradation caused stable HOX/MEIS downregulation

at 72 hours and significant differentiation, mimicking what observed upon XPO1 inhibition (Figures 1D, 1E, supplemental Figures 1F, 1G). Altogether, these results clearly indicate that only prolonged loss of the NPM1c-XPO1 interaction can induce stable HOX/MEIS downregulation and differentiation in *NPM1*-mutated AML cells.

Next, we compared the ability of intermittent and prolonged XPO1 inhibition to induce differentiation of *NPM1*-mutated AML cells *in vitro*. Prolonged XPO1 inhibition with 50 nM of either Selinexor or Eltanexor induced differentiation of OCI-AML3 and *NPM1*-mutated PDX cells (PDX2)⁷, while 2 days/week treatment resulted in negligible changes (Figures 1F-1G, supplemental Figure S1H). Doubling the concentration of Selinexor to 100 nM 2 days/week did not significantly increase differentiation (Figure 1G). Importantly, ectopic expression of *HOXA9* and *MEIS1* significantly rescued differentiation upon prolonged XPO1 inhibition (Figure 1I, supplemental Figure S1I), confirming that persistent HOX/MEIS downregulation is required to achieve differentiation of *NPM1*-mutated cells.

Finally, we assessed the anti-leukemic potential of intermittent and prolonged XPO1 inhibition *in vivo* using two highly aggressive *NPM1/FLT3/DNMT3A* triple-mutated luciferase-expressing PDX models (PDX2 and PDX3). First, we treated NSG mice engrafted with PDX2 cells with either vehicle, Selinexor 2 days/week, Selinexor 5 days/week and Eltanexor 5 days/week (Figure 2A). HOX/MEIS expression and differentiation in sorted leukemic cells was analyzed after one week of treatment. While Selinexor 2 days/week did not induce changes of *HOXA9*, *HOXA10* and *MEIS1* nor of CD11b levels, both Selinexor and Eltanexor 5 days/week caused remarkable HOX/MEIS downregulation and differentiation (Figure 2B and 2C). Next, we investigated the leukemic engraftment of PDX2 cells by flow-cytometry and immunohistochemistry in the bone marrow following two weeks of treatment. Both 5 days/week regimens caused significant engraftment reduction, while 2 days/week Selinexor did not (Figure 2D and 2E), demonstrating that only prolonged XPO1 inhibition is effective against *NPM1*-mutated cells *in vivo*. Finally, to test the impact of prolonged XPO1 inhibition on AML growth *in vivo* and on survival, we treated both PDX2 and PDX3 mice with Eltanexor 5 days/week for 4 consecutive weeks. Eltanexor resulted in significant reduction of bioluminescence and prolonged survival compared to vehicle in both PDX models (Figures 2F-2I). Treatment was well tolerated with no progressive weight loss reported (supplemental Figures 2C and 2D).

NPM1-mutated AML is genetically well-characterized and is now included as distinct entity in the World Health Organization classification of myeloid neoplasms¹⁷. Therefore, there is growing interest in developing molecular targeted therapies for this AML variant¹⁸, including drugs interfering with HOX/MEIS expression, such as Menin-MLL¹⁹⁻²¹ and XPO1 inhibitors⁷. This study clearly demonstrates that 5 days/week XPO1 inhibition stably downregulates HOX/MEIS, induces differentiation and results into prolonged survival of *NPM1*-mutated PDX mice. In contrast, twice/week XPO1 inhibition does not elicit robust antileukemic activity *in vitro* and *in vivo* in *NPM1*-mutated AML, likely explaining the lack of benefit of Selinexor in patients with this leukemia. How XPO1 inhibition results in HOX/MEIS downregulation remains unclear. Possible hypotheses include nuclear relocation of NPM1c interactors with transcriptional repressive properties² and displacement of NPM1c from XPO1 bound at HOX/MEIS loci^{2,22}. Furthermore, mechanisms other than those mediated by NPM1c, e.g. TP53 activation (supplemental Figure 1D and ⁹), may contribute to the anti-leukemic effects of XPO1 inhibitors in *NPM1*-mutated cells. In conclusion, as Phase 1 data of Eltanexor have shown it can be safely administered 5 days/week¹⁵, this study lays the groundwork for the appropriate design of new clinical trials with XPO1 inhibitors in *NPM1*-mutated AML.

Acknowledgments

This work has been supported by the Associazione Italiana per la Ricerca sul Cancro (AIRC Start-Up Grant 2019 n. 22895) and the European Research Council (ERC Adv grant 2016 n. 740230 and ERC Cons Grant 2016 n. 725725).

Authorship contribution

G.P., B.F. and L.B. conceived the study. G.P., A.G., F.R. and A.M. performed *in vitro* experiments. B.B. performed IHC analysis. G.P. and F.M. performed *in vivo* experiments. S.C. and C.Q. performed RNA-sequencing. V.T. and G.S. performed bioinformatic analysis. M.P.M., P.S., F.L., S.S. and Y.L. provided reagents and critical inputs. G.P., B.F. and L.B. analyzed the results and wrote the manuscript with the input from all the authors.

Conflict-of-interest

L.B. declares consultancy at scientific advisory boards for Abbvie and Amgen. B.F. licensed a patent on NPM1 mutants (n. 102004901256449). B.F. and M.P.M. declare honoraria from Rasna Therapeutics, Inc for scientific advisor activities. M.P.M. also declares honoraria/consultancy at scientific advisory board for Abbvie, Amgen, Celgene, Janssen, Novartis, Pfizer, Jazz Pharmaceuticals. P.S. declares honoraria/consultancy at scientific advisory board for Abbvie, Janssen, Novartis, AstraZeneca, Incyte. Y.L and S.S. are employees and stockholders of Karyopharm Therapeutics Inc. F.L. reported receiving personal fees from Amgen Speakers' Bureau and advisory board membership, Novartis Speakers' Bureau and advisory board membership, Bellicum Pharmaceuticals advisory board membership, Miltenyi Speakers' Bureau, Jazz Pharmaceutical Speakers' Bureau, Takeda Speakers' Bureau, Neovii advisory board membership, and Medac Speakers' Bureau outside the submitted work.

References

1. Falini B, Mecucci C, Tiacci E, et al. Cytoplasmic nucleophosmin in acute myelogenous leukemia with a normal karyotype. *The New England journal of medicine*. 2005;352(3):254-266.
2. Falini B, Brunetti L, Sportoletti P, Martelli MP. NPM1-mutated acute myeloid leukemia: from bench to bedside. *Blood*. 2020;136(15):1707-1721.
3. Falini B, Bolli N, Shan J, et al. Both carboxy-terminus NES motif and mutated tryptophan(s) are crucial for aberrant nuclear export of nucleophosmin leukemic mutants in NPMc+ AML. *Blood*. 2006;107(11):4514-4523.
4. Falini B, Bolli N, Liso A, et al. Altered nucleophosmin transport in acute myeloid leukaemia with mutated NPM1: molecular basis and clinical implications. *Leukemia*. 2009;23(10):1731-1743.
5. Alcalay M, Tiacci E, Bergomas R, et al. Acute myeloid leukemia bearing cytoplasmic nucleophosmin (NPMc+ AML) shows a distinct gene expression profile characterized by up-regulation of genes involved in stem-cell maintenance. *Blood*. 2005;106(3):899-902.
6. Verhaak RG, Goudswaard CS, van Putten W, et al. Mutations in nucleophosmin (NPM1) in acute myeloid leukemia (AML): association with other gene abnormalities and previously established gene expression signatures and their favorable prognostic significance. *Blood*. 2005;106(12):3747-3754.
7. Brunetti L, Gundry MC, Sorcini D, et al. Mutant NPM1 Maintains the Leukemic State through HOX Expression. *Cancer Cell*. 2018;34(3):499-512 e499.
8. Azizian NG, Li Y. XPO1-dependent nuclear export as a target for cancer therapy. *J Hematol Oncol*. 2020;13(1):61.
9. Ranganathan P, Yu X, Na C, et al. Preclinical activity of a novel CRM1 inhibitor in acute myeloid leukemia. *Blood*. 2012;120(9):1765-1773.
10. Gu X, Ebrahim Q, Mahfouz RZ, et al. Leukemogenic nucleophosmin mutation disrupts the transcription factor hub that regulates granulomonocytic fates. *J Clin Invest*. 2018;128(10):4260-4279.
11. Garzon R, Savona M, Baz R, et al. A phase 1 clinical trial of single-agent selinexor in acute myeloid leukemia. *Blood*. 2017;129(24):3165-3174.
12. Bhatnagar B, Zhao Q, Mims AS, et al. Selinexor in combination with decitabine in patients with acute myeloid leukemia: results from a phase 1 study. *Leuk Lymphoma*. 2020;61(2):387-396.

13. Fiedler W, Chromik J, Amberg S, et al. A Phase II study of selinexor plus cytarabine and idarubicin in patients with relapsed/refractory acute myeloid leukaemia. *Br J Haematol*. 2020;190(3):e169-e173.
14. Sweet K, Bhatnagar B, Dohner H, et al. A 2:1 randomized, open-label, phase II study of selinexor vs. physician's choice in older patients with relapsed or refractory acute myeloid leukemia. *Leuk Lymphoma*. 2021:1-12.
15. Lee S, Mohan S, Knupp J, et al. Updated overall survival of eltanexor for the treatment of patients with hypomethylating agent refractory myelodysplastic syndrome. *Hemasphere*. 2021(5):431.
16. Spinozzi G, Tini V, Adorni A, Falini B, Martelli MP. ARPIR: automatic RNA-Seq pipelines with interactive report. *BMC Bioinformatics*. 2020;21(Suppl 19):574.
17. Swerdlow SH, World Health Organization, International Agency for Research on Cancer. WHO classification of tumours of haematopoietic and lymphoid tissues (ed Revised 4th edition.). Lyon: International Agency for Research on Cancer; 2017.
18. Falini B, Brunetti L, Martelli MP. How I diagnose and treat NPM1-mutated AML. *Blood*. 2020.
19. Kuhn MW, Song E, Feng Z, et al. Targeting Chromatin Regulators Inhibits Leukemogenic Gene Expression in NPM1 Mutant Leukemia. *Cancer Discov*. 2016;6(10):1166-1181.
20. Uckelmann HJ, Kim SM, Wong EM, et al. Therapeutic targeting of preleukemia cells in a mouse model of NPM1 mutant acute myeloid leukemia. *Science*. 2020;367(6477):586-590.
21. Klossowski S, Miao H, Kempinska K, et al. Menin inhibitor MI-3454 induces remission in MLL1-rearranged and NPM1-mutated models of leukemia. *J Clin Invest*. 2020;130(2):981-997.
22. Oka M, Mura S, Otani M, et al. Chromatin-bound CRM1 recruits SET-Nup214 and NPM1c onto HOX clusters causing aberrant HOX expression in leukemia cells. *Elife*. 2019;8.

Figure legends

Figure 1. Prolonged XPO1 inhibition is necessary to elicit significant antileukemic activity in *NPM1*-mutated AML *in vitro*.

A) Fluorescence microscopy of NPM1c-GFP OCI-AML3 treated for 12 hours with Selinexor 100nM. After 12 hours, Selinexor was removed from medium and cells left in culture for the following 24 hours taking pictures at 2, 4, 6, 20 and 24 hours after drug washout. Hoechst 33342 was used to stain nuclei of the cells. 100X magnification. Scale bar, 20 μ m. **B)** Principal component analysis (PCA) plot derived from the means (N=2) of the FPKM values of parental OCI-AML3 cells collected at 24 hours treated with either DMSO or Selinexor 50 nM, and OCI-AML3 cells collected at 72 hours treated with either DMSO, Selinexor 50 nM short treatment (ST, 24h Selinexor + 48h fresh medium) or Selinexor 50 nM continuous treatment (CT, 72h Selinexor 50 nM). **C)** Volcano plots depicting differentially expressed genes in parental OCI-AML3 cells treated with 72h Selinexor continuous treatment (CT, 72h Selinexor 50 nM) and 72h Selinexor short treatment (ST, 24h Selinexor 50 nM + 48h fresh medium), compared to DMSO. Log₂FC and Log₁₀ p values are shown on the X and Y axis, respectively. Genes belonging to the HOXA (red) HOXB (blue) and MEIS/PBX (green) families are highlighted. **D)** *HOXA9*, *HOXA10* and *MEIS1* expression by qPCR in NPM1c-FKBP(F36V)-GFP OCI-AML3 cells after 24 hours treatment with either DMSO or dTAG and 72h treatment with either DMSO, dTAG-13 ST (short treatment, 24h dTAG-13 + 48h fresh medium) or dTAG-13 CT (continuous treatment, 72h dTAG-13). N=3. Mean $\pm$ SEM. Tukey multiple comparison test. **E)** Flow-cytometry quantification of CD11b, expressed as MFI fold change relative to DMSO, in NPM1c-FKBP(F36V)-GFP OCI-AML3 cells at day 11 following treatment with either DMSO, dTAG-13 2 days/week or dTAG-13 5 days/week. N=4. Mean $\pm$ SEM. Tukey multiple comparison test. **F)** May-Grünwald Giemsa staining of OCI-AML3 cells at day 11 following treatment with either DMSO, Selinexor 2 days/week, Selinexor 5 days/week or Eltanexor 5 days/week. 40X magnification. **G)** Flow-cytometry quantification of CD11b, expressed as MFI fold change relative to DMSO, in OCI-AML3 cells at day 11 of treatment with either DMSO, Selinexor 50 nM 2 days/week, Selinexor 100 nM 2 days/week, Selinexor 50 nM 5 days/week and Eltanexor 50 nM 5 days/week. N=3. Mean $\pm$ SEM. Tukey multiple comparison test. **H)** Flow-cytometry quantification of CD11b, expressed as MFI fold change relative to

DMSO, in PDX2 cells at day 11 of treatment with either DMSO, Selinexor 50 nM 2 days/week, Selinexor 100 nM 2 days/week, Selinexor 50 nM 5 days/week or Eltanexor 50 nM 5 days/week. N=3 for DMSO and Eltanexor, N=2 for the other groups. Mean $\pm$ SEM. Tukey multiple comparison test. **I)** Flow-cytometry quantification of CD11b, expressed as MFI fold change relative to DMSO, in untransduced and HOXA9/MEIS1-transduced OCI-AML3 cells after 7 days of continuous treatment with either DMSO or Selinexor 50nM. N=4. Mean $\pm$ SEM. Tukey multiple comparison test.

H33342, Hoechst 33342; PC, principal component; FC, fold change; padjust, adjusted p value, MFI, Median Fluorescence Intensity; H9/M1, HOXA9/MEIS1; Seli, Selinexor. Elta, Eltanexor

Figure 2: Prolonged XPO1 inhibition is necessary to elicit significant antileukemic activity in *NPM1*-mutated AML *in vivo*.

A) Schematic overview of the *in vivo* experiments. Each NSG mouse was transplanted with 1×10^6 GFP-Luc positive PDX cells. **B)** *HOXA9*, *HOXA10*, *MEIS1* and *PBX3* expression by qPCR in sorted PDX2 cells after 7 days of treatment with either vehicle, Selinexor 5 mg/kg 2 days/week, Selinexor 5 mg/kg 5 days/week and Eltanexor 10 mg/kg 5 days/week. N=4 mice per group. Mean $\pm$ SEM. Dunnett multiple comparison test. **C)** Flow-cytometry quantification of human CD11b, expressed as MFI fold change relative to vehicle, in sorted PDX2 cells after 7 days of treatment with either vehicle, Selinexor 5 mg/kg 2 days/week, Selinexor 5 mg/kg 5 days/week and Eltanexor 10 mg/kg 5 days/week. N=4 mice per group. Mean $\pm$ SEM. Tukey multiple comparison test. **D)** Bone marrow engraftment of PDX2 cells measured as human CD45 percent of positive cells after two weeks of treatment with either vehicle, Selinexor 5 mg/kg 2 days/week, Selinexor 5 mg/kg 5 days/week and Eltanexor 10 mg/kg 5 days/week. N=3 mice per group. Mean $\pm$ SEM. Tukey multiple comparison test. **E)** Representative images of BM histological sections stained for human CD45 after 14 days of treatment with either vehicle, Selinexor 5 mg/kg 2 days/week, Selinexor 5 mg/kg 5 days/week and Eltanexor 10 mg/kg 5 days/week. 40X magnification. Scale bars, 20 μ m. **F)** Representative bioluminescence images of NSG mice transplanted with PDX2 cells treated with either vehicle (N=6) or Eltanexor 10 mg/kg (N=7) 5 days/week for 4 weeks. **G)** Kaplan-Meier curves of

371 PDX2 mice treated with either vehicle (N=6) or Eltanexor 10 mg/kg (N=7) 5
 372 days/week for 4 weeks. Treatment time is shown in light grey. Long-rank (Mantel-
 373 Cox) test. **H)** Representative bioluminescence images of NSG mice transplanted
 374 with PDX3 cells and treated with either vehicle (N=6) or Eltanexor 10 mg/kg (N=7) 5
 375 days/week for 4 weeks. **I)** Kaplan-Meier curves of PDX3 mice treated with either
 376 vehicle (N=6) or Eltanexor 10 mg/kg (N=7) 5 days/week for 4 weeks. Treatment time
 377 is shown in light grey. Long-rank (Mantel-Cox) test.
 378 BM, bone marrow; Seli, Selinexor; Elta, Eltanexor; HR; hazard ratio.

A

B

C

D

E

F

G OCI-AML3

H PDX2

I

Figure 1

Figure 2