1 **Chromosome-scale haplotype-phased genome assemblies of the male and female lines of wild**
2 **asparagus (*Asparagus kiusianus*), a dioecious plant species**

3
4 Kenta Shirasawa^{1,*†}, Saki Ueta^{2,†}, Kyoko Murakami², Mostafa Abdelrahman^{3,4,5}, Akira Kanno³, Sachiko
5 Isobe¹

6
7 ¹Kazusa DNA Research Institute, Kisarazu, Chiba 292-0818, Japan

8 ²Kagawa Prefectural Agricultural Experiment Station, Ayagawa, Kagawa 761-2306, Japan

9 ³Graduate School of Life Sciences, Tohoku University, Sendai, Miyagi 980-8577, Japan

10 ⁴Biotechnology Program, Faculty of Science, Galala University, Suze, Galala 43511, Egypt

11 ⁵Botany Department, Faculty of Science, Aswan University, Aswan 81528, Egypt

12
13 [†]These authors contributed equally to the work.

14
15 ^{*}To whom correspondence should be addressed:

16 Kenta Shirasawa

17 Kazusa DNA Research Institute

18 2-6-7 Kazusa-kamatari, Kisarazu, Chiba 292-0818, Japan

19 Tel.: +81-438-52-3935.

20 Fax.: +81-438-52-3934.

21 E-mail: shirasaw@kazusa.or.jp

22
23 **Running title:**

24 Genome assemblies of wild asparagus

25

26 Abstract

27 *Asparagus kiusianus* is a disease-resistant dioecious plant species and a wild relative of garden asparagus
 28 (*A. officinalis*). To enhance *A. kiusianus* genomic resources, advance plant science, and facilitate asparagus
 29 breeding, we determined the genome sequences of the male and female lines of *A. kiusianus*. Genome
 30 sequence reads obtained with a linked-read technology were assembled into four haplotype-phased contig
 31 sequences (~1.6 Gb each) for the male and female lines. The contig sequences were aligned onto the
 32 chromosome sequences of garden asparagus to construct pseudomolecule sequences. Approximately
 33 55,000 potential protein-encoding genes were predicted in each genome assembly, and ~70% of the
 34 genome sequence was annotated as repetitive. Comparative analysis of the genomes of the two species
 35 revealed structural and sequence variants between the two species as well as between the male and female
 36 lines of each species. Genes with high sequence similarity with the male-specific sex determinant gene in
 37 *A. officinalis*, *MSE1/AoMYB35/AspTDF1*, were presented in the genomes of the male line but absent from
 38 the female genome assemblies. Overall, the genome sequence assemblies, gene sequences, and structural
 39 and sequence variants determined in this study will reveal the genetic mechanisms underlying sexual
 40 differentiation in plants, and will accelerate disease-resistance breeding in garden asparagus.

41

42 **Keywords:** dioecy, genome assembly, haplotype phasing sequence, linked reads, structural variation

43

44 Introduction

45 *Asparagus kiusianus* is a wild relative of garden asparagus (*Asparagus officinalis*). While garden asparagus
 46 is a cultivated species belonging to the Asparagaceae family and is consumed as a vegetable crop around
 47 the world, *A. kiusianus* is native to the coastal regions of Japan¹. Therefore, *A. kiusianus* might exhibit
 48 tolerances and/or resilience to abiotic and biotic stresses. Although *A. kiusianus* has been identified as a
 49 potential donor of stem-blight disease resistance in asparagus breeding programs², neither the genetic
 50 mode nor the genetic loci of resistance have been elucidated to date.

51 *A. officinalis* is a dioecious species and is widely recognized as a model for sex determination in

plants. Recent studies indicate that the male-specific *MYB*-like gene, *MSE1/AoMYB35/AspTDF1*, located at the masculinization-promoting *M* locus of the Y-specific region in asparagus, functions in sex determination in asparagus³⁻⁵. Since *A. kiusianus* is also a dioecious plant species, like garden asparagus, it is possible that both species share the same system of sex determination. Therefore, comparative genome sequence and structure analyses between the two species could provide insights into the molecular mechanisms underlying sex determination in Asparagaceae and the evolutionary processes involved therein.

Advances in sequencing technologies have enabled the whole-genome sequencing of various plant species, thus providing fundamental information required for understanding the plant biology and accelerating breeding programs. Nevertheless, while the genome sequence data of garden asparagus³ and transcriptome data of *A. kiusianus*^{6,7} have been made publicly available, no whole-genome sequence data have been released for *A. kiusianus* to date. Owing to the dioecious nature of *A. kiusianus*, which leads to allogamy, its genome is predicted to be highly heterozygous. Therefore, haplotype-phased genome sequence data would be useful for dissecting the allelic sequence and structural variations in *A. kiusianus*. In this study, we employed a linked-read technology (10X Genomics, Pleasanton, CA, USA) to construct haplotype-based genome sequence assemblies of the male and female lines of *A. kiusianus*. The genome sequence assemblies were then used for gene prediction and sequence and structural variant discovery. Overall, the genome sequence information of *A. kiusianus* obtained in this study could accelerate studies on plant sex determination and facilitate asparagus breeding programs.

71

72 **Materials and methods**

73 *Plant materials*

Male (K1) and female (K2) lines of *A. kiusianus* cultivated at Kagawa Prefectural Agricultural Experiment Station (Kagawa, Japan) were used in this study. Genomic DNA was extracted from the stems of young seedlings using the modified cetyltrimethylammonium bromide (CTAB) method⁸.

77

78 *Genome sequencing and assembly*

79 Genomic DNA libraries of male and female lines were prepared using the Chromium Genome Library Kit
80 v2 (10X Genomics), and sequenced on NovaSeq 6000 (Illumina, San Diego, CA, USA) in paired-end, 150
81 bp mode. The sequence reads were assembled with Supernova (10X Genomics) to construct the contig
82 sequences, scaffold the contigs, and resolve haplotype phases. DNA library preparation, sequencing, and
83 assembly were conducted by Takara Bio (Shiga, Japan) as an outsourcing service.

84 The genome sizes of male and female lines were estimated based on short reads using Jellyfish. To
85 construct pseudomolecule sequences at the chromosome level, the assembled contigs were aligned against
86 the sequence of 10 *A. officinalis* chromosomes (reference) using RaGoo.

87 The software tools used for data analyses are listed in Supplementary Table S1.

88

89 *Repetitive sequence analysis and gene prediction*

90 Repetitive sequences in the assemblies were identified with RepeatMasker, using repeat sequences
91 registered in Repbase and a *de novo* repeat library built with RepeatModeler.

92 RNA-Seq reads of *A. kiusianus* and *A. officinalis* were obtained from a public DNA database
93 (GenBank Sequence Read Archive accession number: SRA1003110)^{6,7}. The RNA-Seq reads, from which
94 adapter sequences were trimmed with fastx_clipper in the FASTX-Toolkit, were aligned against the
95 assembled sequences with HISAT2. Gene prediction was performed with BREAKER2 using the positional
96 information of the repeats, RNAs, and peptide sequences of the predicted genes of *A. officinalis* (V1.1)³
97 released in Phytozome.

98

99 *Comparative genome structure analysis of A. kiusianus and A. officinalis*

100 Chromosome-level genome sequence assemblies of *A. kiusianus* (this study) and *A. officinalis* (V1.1)³
101 were compared with Minimap2, and the resultant Pairwise Mapping Format (PAF) files were visualized
102 with pafv.

103

Results and data description

Haplotype-phased genome assembly

Short-read sequences of the male (143.7 Gb) and female (140.0 Gb) lines of *A. kiusianus* were obtained in this study, and their genome sizes were estimated at 1,563.8 and 1,729.4 kb, respectively (Figure 1).

The short-read sequences of the male line were assembled into raw contigs (total length = 3,724.6 kb, N50 = 7.5 kb), which included gaps and all homologous sequences of the diploid genome (Supplementary Table S2). Then, the homologous sequences were flattened, and the gaps were filled by joining the sequence to its flanking sequence, thus producing megabubble sequences (total length = 1,811.6 kb, N50 = 170.6 kb) (Supplementary Table S2). Finally, two haplotype-phased genome assemblies (each containing 111,443 sequences) were generated from the megabubble sequences (Table 1). Haplotype 1 spanned 1,618.9 Mb in total with an N50 value of 155.5 kb, while haplotype 2 spanned 1,618.5 Mb in total with an N50 length of 155.3 kb. Complete Benchmarking Single-Copy Orthologs (BUSCO) scores were 88.4% and 88.6% for haplotypes 1 and 2, respectively (Table 1). The male genome assemblies for haplotype 1 and 2 were designated as AKIK1p1 and AKIK1p2, respectively.

Similarly, short-read sequences of the female line were assembled into raw contigs (total length = 3,780.7 kb, N50 = 7.0 kb) and megabubble sequences (total length = 1,703.6 Gb, N50 = 76.2 kb) (Supplementary Table S2). The resultant haplotype-phased assemblies (each containing 125,241 sequences) spanned 1,567.6 Mb in total with an N50 value of 59.7 kb for “haplotype 1”, and 125,241 sequences of 1,567.3 Mb in total with an N50 value of 59.6 kb for “haplotype 2” (Table 1). Complete BUSCO scores were 88.7% and 88.6% for haplotypes 1 and 2, respectively (Table 1). The female genome assemblies for haplotype 1 and 2 were designated as AKIK2p1 and AKIK2p2, respectively.

The four sets of genome sequence assemblies of *A. kiusianus* (haplotypes 1 and 2 of male and female lines) were aligned against the chromosome-scale genome assembly of *A. officinalis*. A total of 96,224 sequences (1,535.9 kb) for haplotype 1 and 96,224 sequences for haplotype 2 (1,535.3 kb) in the male line, and 107,875 sequences (1,491.7 kb) for haplotype 1 and 107,864 sequences (1,489.7 kb) for haplotype 2 in the female line, could be aligned to the 10 chromosome sequences of *A. officinalis* (Table 2).

Repetitive sequence analysis and gene prediction

Repeat sequences occupied 66.8% (AKIK1p1), 66.8% (AKIK1p2), 68.1% (AKIK2p1), and 68.1% (AKIK2p2). The most abundant repetitive sequences were long terminal repeats (LTRs) (45.4–46.3%), followed by unclassified repeats (14.9–13.7%) and DNA transposons (4.7–4.8%) (Table 3).

A total of 404.2 million RNA reads for 18 samples were mapped to the genome sequences. The mapping rates of *A. kiusianus* RNA-Seq reads were 93.7–94.1%, while those of *A. officinalis* reads were 82.4–82.6%. Based on the positions of RNA-Seq reads on the genome sequences, a total of 59,208, 56,706, 57,523, and 58,694 potential protein-coding genes were predicted in AKIK1p1, AKIK1p2, AKIK2p1, and AKIK2p2, respectively (Table 2), of which 365, 380, 472, and 505 genes contained premature termination codons in their internal sequences. Complete BUSCO scores ranged from 90.1% (AKIK2p1) to 91.4% (AKIK1p2).

Next, we compared the sequences of predicted genes with *MSE1/AoMYB35/AspTDF1*, which has been reported as the male-specific sex determinant gene in *A. officinalis*³⁻⁵. Two genes, K1p1ch01g28074 and K1p2ch01g47751, identified in the genomes of the male line exhibited high sequence similarity with the query; however, none of the genes in the female genome assemblies showed significant sequence similarity with the query.

Genome sequence and structural variations between *A. kiusianus* and *A. officinalis*

Haplotype sequences within the male and female lines were compared. A total of 386,723 single nucleotide polymorphisms (SNPs; transition [Ts]/transversion [Tv] ratio = 3.0) and 46,007 insertions/deletions (indels) were identified between the two haplotypes of the male line (Table 4). On the other hand, 293,196 SNPs (Ts/Tv = 3.0) and 35,732 indels were identified between the two haplotype sequences of the female line (Table 4). The haplotype sequences of male and female lines were also compared, and 321,334 SNPs and 49,921 indels on average were identified across the four haplotype combinations (Table 4).

While the chromosome structures were conserved within *A. kiusianus* lines and between *A. kiusianus*

and *A. officinalis* (Figure 2), genomic rearrangements were observed at the local level. For instance, at the sex-related region including the male-specific gene *MSE1/AoMYB35/AspTDF1* of *A. officinalis*, sequence collinearity was disrupted by inversions and translocations between the male and female lines (Figure 3). Although sequence similarity was low between the male haplotypes of *A. kiusianus* and *A. officinalis*, sequence collinearity was moderately conserved (Figure 3).

Conclusion and future perspectives

We present the chromosome-level haplotype-phased genome assemblies of the male and female lines of *A. kiusianus*, a wild relative of garden asparagus. The genome size of *A. kiusianus* was estimated to be approximately 1.6 Gb (Figure 1), which was 300 Mb larger than that of garden asparagus (ca. 1.3 Gb)³. This estimation was reflected in the difference between the assembly sizes of *A. kiusianus* (1.6 Gb; Table 1) and garden asparagus (1.2 Gb). Of the 1.6 Gb assembly, 1.5 Gb could be aligned to the pseudomolecule sequence of asparagus, without any structural rearrangements (Figure 2 and Table 2). Since we determined haplotype-phased genome sequences for the male and female lines of *A. kiusianus*, it was possible to compare the sequence and structure of the *M* locus between the Y-specific region of the male line and the corresponding region of the female line (Figure 3). The result suggested dynamic genome rearrangements between the male and female lines, similar to that reported in *jojoba*⁹, which might lead to presence/absence variation of the male-specific gene *MSE1/AoMYB35/AspTDF1* between the male and female lines³⁻⁵.

The genome of *A. kiusianus* harbors valuable genes that could be used for the breeding of elite garden asparagus cultivars. Because of cross-compatibility between the two species^{2,10}, important genetic loci such as those imparting resistance to stem blight², which causes considerable production losses, could be transferred from *A. kiusianus* into garden asparagus. However, while DNA markers linked to the genes would facilitate the selection of disease-resistant lines in breeding programs, the genetic loci responsible for disease resistance have not been reported so far. The chromosome-level genome sequence of *A. kiusianus* presented in this study could serve as a reference for genetic mapping and the identification of

resistance genes, as well as for transcriptome analysis and the determination of gene functions and mechanisms underlying the resistance and susceptible phenotypes^{6,7}.

Although the plant genomics era started with the whole-genome sequencing of *Arabidopsis thaliana*¹², an undomesticated species, the advanced approaches of plant genomics have been applied more frequently to agronomically important crops rather than to wild plant species¹¹. Wild plants have the potential to accelerate the pace of breeding programs and to further the field of plant science^{13,14}. The genome sequence information of *A. kiusianus* generated in this study will help to reveal the genetic mechanisms underlying sexual differentiation in plants and will accelerate disease-resistance breeding in asparagus.

Data availability

Sequence reads generated in this study are available from the Sequence Read Archive (DRA) of the DNA Data Bank of Japan (DDBJ) under accession number: DRA012987. The DDBJ accession numbers of the assembled genome sequences are BQKN01000001-BQKN01111443 (AKIK1p1), BQKO01000001-BQKO01111443 (AKIK1p2), BQKP01000001-BQKP01125241 (AKIK2p1), and BQKQ01000001-BQKQ01125241 (AKIK2p2). The genome sequence information generated in this study is available at Plant GARDEN (<https://plantgarden.jp>).

Supplementary data

Supplementary Table S1 Software tools used for genome assembly and gene prediction.

Supplementary Table S2 Statistics of the primary genome sequence assemblies of the male and female lines of *Asparagus kiusianus*.

Acknowledgements

We thank T. Ikeuchi and T. Nakamura (Kagawa Prefectural Agricultural Experiment Station) for providing the plant material, and A. Watanabe (Kazusa DNA Research Institute) for technical assistance.

Funding

This work was partially supported by the Kazusa DNA Research Institute Foundation.

Conflict of interest

None declared.

References

1. Ohwi, J. 1965, Asparagus L. *Flora of Japan*.
2. Iwato, M., Kosaza, M., Takeuchi, Y., et al. 2014, Stem blight resistance of Asparagus kiusianus and its hybrid with A. officinalis. *Advances in Horticultural Science*, **28**, 202–207.
3. Harkess, A., Zhou, J., Xu, C., et al. 2017, The asparagus genome sheds light on the origin and evolution of a young Y chromosome. *Nat Commun*, **8**, 1279.
4. Murase, K., Shigenobu, S., Fujii, S., et al. 2017, MYB transcription factor gene involved in sex determination in Asparagus officinalis. *Genes Cells*, **22**, 115–123.
5. Tsugama, D., Matsuyama, K., Ide, M., Hayashi, M., Fujino, K. and Masuda, K. 2017, A putative MYB35 ortholog is a candidate for the sex-determining genes in Asparagus officinalis. *Sci Rep*, **7**, 41497.
6. Abdelrahman, M., Nakabayashi, R., Mori, T., et al. 2020, Comparative Metabolome and Transcriptome Analyses of Susceptible Asparagus officinalis and Resistant Wild A. kiusianus Reveal Insights into Stem Blight Disease Resistance. *Plant Cell Physiol*, **61**, 1464–1476.
7. Abdelrahman, M., Suzumura, N., Mitoma, M., et al. 2017, Comparative de novo transcriptome profiles in Asparagus officinalis and A. kiusianus during the early stage of Phomopsis asparagi infection. *Sci Rep*, **7**, 2608.
8. Porebski, S., Bailey, L. G. and Baum, B. R. 1997, Modification of a CTAB DNA extraction

234 protocol for plants containing high polysaccharide and polyphenol components. *Plant*
235 *Molecular Biology Reporter*, **15**, 8–15.

236 9. Al-Dossary, O., Alsubaie, B., Kharabian-Masouleh, A., Al-Mssallem, I., Furtado, A. and
237 Henry, R. J. 2021, The jojoba genome reveals wide divergence of the sex chromosomes in
238 a dioecious plant. *Plant J.*

239 10. Ito, T., Konno, I., Kubota, S., et al. 2011, Production and characterization of
240 interspecific hybrids between *Asparagus kiusianus* Makino and *A. officinalis* L. *Euphytica*,
241 **182**.

242 11. Shirasawa, K., Harada, D., Hirakawa, H., Isobe, S. and Kole, C. 2021, Chromosome-level
243 de novo genome assemblies of over 100 plant species. *Breed Sci*, **71**, 117–124.

244 12. The Arabidopsis Genome Initiative 2000, Analysis of the genome sequence of the flowering
245 plant *Arabidopsis thaliana*. *Nature*, **408**, 796–815.

246 13. Bohra, A., Kilian, B., Sivasankar, S., et al. 2021, Reap the crop wild relatives for
247 breeding future crops. *Trends Biotechnol.*

248 14. Migicovsky, Z. and Myles, S. 2017, Exploiting Wild Relatives for Genomics-assisted
249 Breeding of Perennial Crops. *Front Plant Sci*, **8**, 460.

250

251 **Table 1** Statistics of the genome assemblies of the male and female lines of *Asparagus kiusianus*.

	AKIK1p1_r1.0	AKIK1p2_r1.0	AKIK2p1_r1.0	AKIK2p2_r1.0
Total contig size (bp)	1,618,866,837	1,618,519,636	1,567,596,515	1,567,276,205
Number of sequences	111,443	111,443	125,241	125,241
Sequence N50 length (bp)	155,490	155,269	59,656	59,610
Longest sequence size (bp)	2,375,321	2,375,321	1,231,068	1,222,162
Gap size (bp)	168,294,890	168,291,350	132,162,460	132,161,290
Single-copy complete BUSCOs	82.4%	82.6%	83.5%	83.4%
Duplicated complete BUSCOs	6.0%	6.0%	5.2%	5.2%
Fragmented BUSCOs	5.8%	5.5%	5.6%	5.6%
Missing BUSCOs	5.8%	5.9%	5.7%	5.8%

252

253 **Table 2** Statistics of the pseudomolecule sequences of *A. kiusianus*.

	AKIK1p1_r1.0				AKIK1p2_r1.0			
	Total length (bp)	Gap (%)	Number of contigs	Number of genes	Total length (bp)	Gap (%)	Number of contigs	Number of genes
ch01	179,686,458	11.8	10,537	6,938	179,663,709	11.8	10,543	6,706
ch02	94,379,980	12.5	6,056	3,631	94,286,185	12.5	6,066	3,427
ch03	160,690,220	10.4	10,792	6,135	161,086,385	10.5	10,791	5,855
ch04	222,888,549	11.1	13,146	7,547	222,850,935	11.1	13,139	7,251
ch05	175,875,079	12.0	10,573	6,878	176,035,479	12.0	10,582	6,668
ch06	112,254,750	9.4	7,553	3,696	112,182,592	9.4	7,553	3,504
ch07	231,128,771	11.0	13,973	8,285	230,178,869	11.0	13,928	7,953
ch08	209,954,455	10.5	13,763	7,167	209,720,314	10.5	13,796	6,882
ch09	73,375,337	10.6	4,435	2,759	73,241,043	10.7	4,437	2,638
ch10	85,316,880	10.8	5,396	3,353	85,678,944	10.9	5,389	3,229
Subtotal	1,545,550,479	11.0	96,224	56,389	1,544,924,455	11.0	96,224	54,113
ch00	84,496,458	10.4	15,223	2,819	84,775,281	10.6	15,223	2,593
Total	1,630,046,937	11.0	111,447	59,208	1,629,699,736	11.0	111,447	56,706

	AKIK2p1_r1.0				AKIK2p2_r1.0			
	Total length (bp)	Gap (%)	Number of contigs	Number of genes	Total length (bp)	Gap (%)	Number of contigs	Number of genes
ch01	167,748,676	9.5	11,500	6,772	167,821,022	9.5	11,527	6,835
ch02	88,515,862	10.3	6,726	3,557	87,905,216	10.2	6,698	3,651

ch03	159,691,030	9.0	12,193	5,956	159,408,608	9.0	12,181	6,030
ch04	217,008,733	9.0	15,036	7,377	217,068,856	9.0	15,080	7,655
ch05	172,460,497	9.9	11,974	6,762	171,835,019	9.9	11,943	6,970
ch06	107,102,574	8.1	8,296	3,565	107,106,266	8.1	8,278	3,621
ch07	222,804,915	9.2	15,815	7,998	221,855,507	9.2	15,811	8,158
ch08	213,448,917	8.6	15,408	7,141	213,032,411	8.6	15,401	7,250
ch09	71,157,753	9.4	5,055	2,517	71,672,067	9.5	5,054	2,573
ch10	82,583,459	9.3	5,872	3,262	82,780,279	9.2	5,891	3,270
Subtotal	1,502,522,416	9.2	107,875	54,907	1,500,485,251	9.2	107,864	56,013
ch00	77,633,199	8.8	17,366	2,616	79,350,054	8.9	17,377	2,681
Total	1,580,155,615	9.2	125,241	57,523	1,579,835,305	9.2	125,241	58,694

254

255 **Table 3** Repetitive sequences in the *A. kiusianus* genomes.

Repeat type	AKIK1p1			AKIK1p2			AKIK2p1			AKIK2p2		
	Number of elements	Length occupied (bp)	%	Number of elements	Length occupied (bp)	%	Number of elements	Length occupied (bp)	%	Number of elements	Length occupied (bp)	%
SINEs	296	100,129	0.0	295	100,876	0.0	260	88,942	0.0	269	88,784	0.0
LINEs	36,139	22,836,398	1.4	36,171	22,868,973	1.4	36,022	22,848,619	1.5	36,257	22,865,141	1.5
LTR elements	534,586	740,043,346	45.4	534,419	739,629,919	45.4	538,917	731,783,197	46.3	538,772	731,702,512	46.3
DNA transposons	119,624	77,129,754	4.7	119,369	77,003,389	4.7	117,911	76,193,533	4.8	118,264	76,216,494	4.8
Unclassified	426,795	242,401,476	14.9	426,435	242,359,956	14.9	422,640	239,811,052	15.2	422,494	239,544,071	15.2
Small RNA	447	231,343	0.0	451	233,377	0.0	435	232,256	0.0	442	228,647	0.0
Satellites	1,945	843,970	0.1	1,894	847,909	0.1	2,155	901,555	0.1	2,171	901,844	0.1
Simple repeats	197,902	17,199,114	1.1	198,009	17,248,715	1.1	195,639	16,992,727	1.1	195,634	17,024,257	1.1
Low complexity	32,476	1,822,571	0.1	32,548	1,822,378	0.1	32,388	1,816,175	0.1	32,482	1,817,483	0.1

256

257 **Table 4** Completeness of gene predictions, based on BUSCO assessment.

	Within a line		Between male and female lines			
	AKIK1p1	AKIK2p1	AKIK1p1	AKIK1p1	AKIK1p2	AKIK1p2
	vs	vs	vs	vs	vs	vs
	AKIK1p2	AKIK2p2	AKIK2p1	AKIK2p2	AKIK2p1	AKIK2p2
Base substitutions	386,723	293,196	324,361	327,460	314,704	318,810
Ts/Tv ratio	3.0	3.0	2.9	2.9	2.9	2.9
1bp indels	23,833	17,283	20,344	20,379	19,692	19,970
2bp indels	5,799	4,477	5,722	5,763	5,645	5,691
[3,50) indels	11,100	8,641	13,931	13,897	13,647	13,777
[50,1000) indels	1,871	1,472	7,141	7,127	7,096	7,096
>=1000 indels	3,404	3,859	3,199	3,229	3,160	3,179

258

Figure legends

Figure 1 Estimation of the genome sizes of the male and female lines of *Asparagus kiusianus*, based on k -mer analysis ($k = 17$) with the given multiplicity values.

Figure 2 Comparative analysis of the genome sequence and structure of *Asparagus kiusianus* and *A. officinalis*.

Dots represent similarities between the genome sequence and structure of the two species. Genomes of the male and female lines of *A. kiusianus* are indicated as AKIK1 and AKIK2, respectively. Haplotype phases 1 and 2 are indicated as p1 and p2, respectively. *AsparagusV1* indicates *A. officinalis* genome.

Figure 3 Comparison between the *M* locus in the Y-specific region of the male line and the corresponding region of the female line.

Gray bars indicate similarities in the genome sequence and structure. Black arrows indicate the position and orientation of *MSE1/AoMYB35/AspTDF1*.

AKIK1 **AKIK2**

AKIK2p2 (♀)

AKIK2p1 (♀)

AKIK1p2 (♂)

AKIK1p1 (♂)

AsparagusV1.1 (♂)

0 Mb

2 Mb

4 Mb

6 Mb

8 Mb