A method to estimate the cellular composition of the mouse brain from heterogeneous datasets

Authors:

Dimitri Rodarie¹ (CRediT abbreviations: C, DC, FoA, I, M, So, Va, Vi, WrD, WrR),

Csaba Verasztó¹ (C, DC, I, M, So, Va, WrD, WrR),

Yann Roussel¹ (C, DC, I, M, Va, WrD, WrR),

Michael Reimann¹ (Va, WrD, WrR),

Daniel Keller¹ (Va, WrD, WrR),

Srikanth Ramaswamy¹ (Va, Pa, WrD, WrR),

Henry Markram¹ (Su, Pa, Va, WrD, WrR)

Marc-Oliver Gewaltig¹ (Su, Va, WrD, WrR),

¹ Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Campus Biotech,
1202 Geneva, Switzerland

Corresponding author:

Dimitri Rodarie

Blue Brain Project / EPFL

Campus Biotech

Chemin des Mines, 9

CH-1202 Geneva

E-mail: dimitri.rodarie@epfl.ch

Tel: +41 21 69 55 180

Abstract (233 words)

The mouse brain contains a rich diversity of inhibitory interneuron types that have been characterized by their patterns of gene expression. However, it is still unclear how these cell types are distributed across the mouse brain. We developed a computational method to estimate the densities of different interneuron types across the mouse brain. Our method allows the unbiased integration of diverse and disparate datasets into one framework to predict interneuron densities for uncharted brain regions. We constrained our estimates based on previously computed brain-wide neuron densities, gene expression data from *in situ* hybridization image stacks together with a wide range of values reported in the literature. Using constrained optimization, we derived coherent estimates of cell densities for the different interneuron types. We estimate that 20.3% of all neurons in the mouse brain are inhibitory. Among all inhibitory neurons, 18% predominantly express parvalbumin (PV), 16% express somatostatin (SST), 3% express vasoactive intestinal peptide (VIP), and the remainder 63% belong to the residual GABAergic population. We find that our density estimations improve as more literature values are integrated. Our pipeline is extensible, allowing new cell types or data to be integrated as they become available. The data, algorithms, software, and results of our pipeline are publicly available and update the Blue Brain Cell Atlas. This work therefore leverages the research community to collectively converge on the numbers of each cell type in each brain region.

Author summary (194 words)

Obtaining a global understanding of the cellular composition of the brain is a very complex task, not only because of the great variability that exists between reports of similar counts but also because of the numerous brain regions and cell types that make up the brain. Previously, we presented a model of a cell atlas, which provided an estimate of the densities of neurons, glia and their subtypes for each region in the mouse brain. Here, we describe an extension of this model to include more inhibitory neuron types. We collected estimates of inhibitory neuron counts from literature and built a framework to combine them into a consistent cell atlas. Using brain slice images, we also estimated inhibitory neuron density in regions where no literature data are available. We estimated that in the mouse brain 20.3% of all neurons are inhibitory. Among all inhibitory neurons, 18% predominantly express parvalbumin (PV), 16% express somatostatin (SST), 3% express vasoactive intestinal peptide (VIP), and the remainder 63% belong to the residual GABAergic population. Our approach can be further extended to any other cell type and provides a resource to build tissue-level models of the rodent brain.

Key words: Cell atlas, inhibitory neurons, mouse brain, cellular composition, gene expression

1. Introduction

Over the past century, numerous studies have reported a great variety of cells in the mouse brain according to their morphological, electrical, and molecular features. Several groups such as the BRAIN Initiative® have launched ambitious projects to undertake a comprehensive census of all brain cells, determining their molecular, structural and functional properties [1–4]. Building a catalogue of the vast diversity of neuron types in the brain is a challenge for modern neuroscience. Characterizing this variety is exacerbated by the fact that distinct cell types are localized to specific brain regions - for example, pyramidal cells in the cerebral cortex or Purkinje cells in the cerebellum [5,6]. Therefore, studies tend to simplify this task by focusing on specific brain regions only. Moreover, the variety of methods used results in considerable variability in the outcomes, even when the same region is considered [7–9]. The cellular composition problem can only be conclusively solved by integrating disparate datasets from literature into a coherent framework. Obtaining a complete and comprehensive cell atlas of the mouse brain is, therefore, a monumental task, which needs to be tackled to enable the in-silico reconstruction of multi-scale neural circuits [6,10–13].

1.1. First version of the mouse brain cell atlas

Erö and colleagues [14] were able to estimate the cellular composition of each brain region defined by the Allen Mouse Brain Atlas [15] with respect to excitatory (glutamatergic) and inhibitory (GABAergic) neurons, astrocytes, oligodendrocytes, and microglia. The authors aligned multiple datasets, including genetic marker expression from *in situ* hybridization (ISH) experiments from the Allen Institute for Brain Science (AIBS), to the Allen Reference Nissl volume from Lein et al. [16] to determine the positions of all cells within the annotation volume. This resulted in the first version of the Blue Brain Mouse Cell Atlas pipeline (BBCAv1) as described in Erö et al. [14].

Erö et al. argued that literature values are often inconsistent and are therefore unreliable [9,14]. Hence, the authors used only whole-brain values from literature (e.g., the total number of cells

and neurons in the mouse brain from Herculano-Houzel et al. [17]) to constrain their estimates of cell densities in every brain region, avoiding any bias toward a particular region. However, there are also several regions for which a consensus on cell composition has been reached (e.g., for the cerebellar Purkinje layer, see review from Keller et al. [9]). For instance, in all regions of the isocortex (also called neocortex), it is well-known that layer 1 regions should only be composed of inhibitory neurons [5]. To take this aspect into account, the BBCAv1 had to manually integrate those regional constraints *a posteriori* (e.g., force each neuron in isocortex L1 to be inhibitory) which in turn contradicted the original global constraints. Moreover, the BBCAv1 strategy yields regional estimates that sometimes do not match their respective literature values (e.g. cell estimates for cortical subregions compared to Herculano-Houzel et al. [18]). Finally, to obtain a more fine-grained classification, the BBCAv1 strategy required brain-wide estimates for every new cell type to be integrated. These constraints may not yet exist. We therefore decided to refine the BBCAv1 strategy to obtain inhibitory neuron subtypes densities.

1.2. A novel approach to estimate cell densities

We extended the workflow from Erö et al. [14], using estimates of cell densities for individual brain regions from the literature instead of whole-brain (global) estimates for specific cell types. Specifically, using additional ISH experiment datasets from the AIBS, we were able to further refine the GABAergic neuron densities from Erö et al. [14] and estimate the densities of GABAergic subclasses. This includes parvalbumin (PV+), somatostatin (SST+), vasoactive intestinal peptide (VIP+) immunoreactive neurons, and the remaining inhibitory neurons. PV+ neurons are fast spiking, resulting in fast and efficient suppression of their surrounding neurons' activity [19,20]. A deficit of PV+ neurons seems to be linked to neural diseases such as schizophrenia [21], Alzheimer's [22] and autism [23,24]. SST+ neurons contribute to long range inhibition [20]. Their population decreases with age and seems to contribute to the development of Alzheimer's disease [25–27]. VIP+ neurons, on the other hand, tend to target

114 other inhibitory neurons in the isocortex, playing the role of local disinhibitors [28]. SST+ and
 115 VIP+ neurons are also involved in the circadian clock which regulates daily brain activity [29].
 116 We also highlighted the fact that the alignment of multiple datasets to a common coordinate
 117 system presents a significant source of uncertainty that we measured and addressed with deep
 118 learning techniques. We combined all these new tools and methods to produce a second
 119 version of the Blue Brain Mouse Cell Atlas pipeline (BBCAv2). We further extended the
 120 BBCAv2 in a companion paper [30], mapping well identified morpho-electrical types to our
 121 GABAergic neuron subclasses. These refinements to the model will help neuroscientists to get
 122 a better understanding at cellular composition in the brain and will pave the way for more
 123 accurate *in silico* reconstructions of brain tissues.

2. Materials and methods

2.1. Overview of the pipeline

2.1.1. Original pipeline

In this paper we refined several steps of the BBCAv1 to facilitate the integration of more and more specific cell types without having to rely on the total number or ratio of these cell types. We started with the original Blue Brain Mouse Cell Atlas workflow (BBCAv1) from Erö et al. [14] which consisted of six steps:

1. The Nissl-stained brain slices from Dong [15], and the corresponding mouse brain Annotation Volume (AV) were manually realigned along the sagittal axis. Every ISH dataset [16] used in the following steps was also manually realigned to match the Nissl volume.
2. The annotated Nissl volume from the AIBS was combined with a total number of cells in the mouse brain from Herculano-Houzel et al. [17] to estimate the number of cells in each brain region.
3. A combination of genetic marker datasets and a global ratio of glial cells in the brain from Herculano-Houzel et al. [17] was used to distinguish neurons from glial cells.
4. Glial cells were labelled as astrocytes, oligodendrocytes or microglia based on whole brain ratios together with their respective ISH datasets.
5. GAD67 (glutamic acid decarboxylase) marker experiment from Lein et al. [16], associated with a global number of inhibitory neurons from Kim et al. (7) was used to distinguish the excitatory from the inhibitory neurons. This number stands for the sum of PV+, SST+ and VIP+ reacting cells in the brain and was assumed in this pipeline to represent the entire inhibitory population.
6. Manual correction of purely inhibitory regions: molecular layer regions of the cerebellum, layer 1 of the isocortex, the reticular nucleus of the thalamus, and the

striatum were considered to be fully inhibitory, which means that all their neurons are labeled as GABAergic.

2.1.2. Adjustments for improved pipeline

Step 1 of the BBCAv1 involved manual selection of landmark points on each slice of the AV, and ISH data to realign to the Nissl volume. This step was not only labor intensive but also error prone. In our updated workflow, we replaced the manual realignment by a novel deep learning based alignment algorithm by Krepl et al. [31].

In step 5, the total number of inhibitory neurons in the mouse brain was a global constraint. Its biological value was taken from Kim et al. [8] and represents the sum of SST+, PV+, and VIP+ neurons. However, these three types do not represent all inhibitory neurons (e.g., LAMP5 cells in the isocortex [32]). This means that the total number of inhibitory cells in BBCAv1 was underestimated, despite the corrections applied in step 6.

We therefore reworked steps 5 and 6 by integrating more literature estimates of neuron type densities as well as by using additional ISH datasets. This not only improved our estimates of the excitatory to inhibitory ratios in the brain, but also provided more precise estimates of GABAergic neuron types, including PV+, SST+, and VIP+ cells. However, as we will discuss later, the spatial resolution and the whole-brain coverage of these datasets are not sufficient to estimate the density of all inhibitory subtypes in each voxel of the mouse brain, as it was done in BBCAv1. Therefore, we computed the mean density of each inhibitory type for each region of the AV, instead of estimating the density per voxel.

Since several AV and Nissl volumes were recently released [33], we also developed a method to choose the best combination to estimate cell densities (see Section 2.2).

2.1.3. New Pipeline

Fig. 1 shows the new pipeline for the Blue Brain Mouse Cell Atlas version 2 (BBCAv2). It consumes data from the AIBS in the form of image stacks of stained coronal brain slices, as well as a combination of annotation and Nissl reference atlases.

Fig 1. New workflow for the Blue Brain Mouse Cell Atlas (BBCAv2).

Steps of the pipeline are displayed in rows with the data consumed in the left column, the method used in the middle column, and the produced results on the right. Each step of this pipeline builds on the results of the previous steps.

The BBCAv2 has four main steps.

1. In the first step, the different image stacks from ISH experiments from the AIBS are automatically aligned and registered to a reference volume. We describe this step in Section 2.3.
2. The second step generates cell, neuron, and glial densities for each region of the AV. It corresponds to steps 2 and 3 of the BBCAv1 pipeline.
3. We devise assumptions to estimate densities of different inhibitory neuron types in a coherent framework (see Section 2.4), and apply them for step 3 and 4. In step 3, we assume a correlation between genetic marker expression and cell type density gathered from literature (see Section 2.5) and the previously realigned and filtered ISH datasets. The details of this step are described in Section 2.6.
4. Step 4 integrates the results of step 3 into a consistent estimate of the cell densities for each inhibitory subtype, according to our assumptions (see Section 2.7). Finally, the cells are placed in the brain volume to produce the updated BBCAv2 model (see Section 2.8).

2.2. Selection of the reference brain atlas based on the resulting cell density distribution

Step 2 of the BBCAv2 (see Fig. 1) uses a whole-brain volume, derived from Nissl-stained coronal brain slices from Dong [15], together with an annotation volume (AV) which maps the voxels of the brain volume to a brain region. Brain regions are organized in a hierarchical tree, with leaves representing the finest subdivisions captured by the AIBS. These reference volumes are registered by the AIBS to a common vector space: the Common Coordinate Framework (CCF).

The AIBS has released multiple versions of the Allen Brain Nissl volume and the related annotation volumes.

The first version (Nissl1 + AV1) was published in 2011 and was based on a single mouse (CCFv1). Nissl1 + AV1 were well aligned to each other at the level of individual slices, however adjacent slices were not. This led to rather a jagged Nissl brain volume and corresponding annotations. Therefore Erö et al. [14] realigned adjacent Nissl and AV slices along the sagittal axis with a non-rigid algorithm to create a new CCF (CCFbbp).

The AIBS used a comparable technique on the same dataset to obtain Nissl2 + AV2 (CCFv2). It still represents one individual, but this time aligned to an averaged mouse brain volume that was derived from more than a thousand adeno-associated virus (AAV) tracer experiments [34,35].

A third AV version (AV3) was completely derived from this average brain, using the pipeline described in Wang et al. [36]. AV3 is currently the smoothest AV, but it is no longer derived and aligned to a Nissl volume (see Fig. 2A); it is paired with the closest Nissl volume available (Nissl2). Moreover, some subregions such as the molecular and granular layers of the cerebellum are merged in the AV3.

Clearly, for the purpose of estimating cell densities, different combinations of Nissl volumes and annotation atlases yield different cell density estimates (see Fig. 2CD). In this section, we therefore present our analysis to identify the best reference dataset pair (referred to as CCF version) for density estimation. We show that the quality of cell density estimates is strongly correlated to the alignment of the Nissl volume to the AV.

Poor alignment means that parts of one region in the Nissl volume are mislabeled in the AV as being part of a neighboring region. This means also that parts of the cell estimates (step 2 of Fig. 1) are assigned to wrong regions. To assess the quality of the alignment we look at parts of the brain where the boundary between two regions is clearly visible in the Nissl volume, specifically the isocortex L1/L2 separation and the cerebellar cortex molecular/granular layers. Fig. 2AB illustrates the misalignment between Nissl and AV volumes for the different reference atlases for these regions.

Fig 2. Local impact of the annotation atlas on the estimation of cell densities.

Panels (A) and (B) show the misalignment between the Nissl datasets in grey and their respective annotation atlases overlaid as colors (coronal slices). Boundaries between annotated regions, when available, are displayed in yellow. Different versions are shown from left to right: CCFbbp, CCFv2 from the AIBS, and CCFv3 from the AIBS. (C) and (D) compare the estimated neuron densities from the BBCAv2 pipeline (step 1 and 2 of Fig. 1) based on each reference atlas version to literature (bars in brick red with black outline). Each reference atlas version appears in different colors: dark blue for CCFbbp, orange for CCFv2, light blue for CCFv3, and green for the manually realigned AV2 and Nissl2.

(A) Focus on the isocortex L1/L2 boundary. On the right of each image, L1 and L2 annotations are shown respectively in blue and orange, on top of the Nissl expression. The density difference between L1 and L2 creates a visible boundary (see left part of the images). However, in each CCF version, parts of the denser L2 are annotated

as L1, raising the estimated density of neurons for L1. The magnifications shown for CCFv3 highlight some examples of this problem.

(B) Focus on the cerebellar cortex, granular/molecular boundary (Purkinje layer not represented). Each subregion of the cerebellar cortex in the annotation atlas is displayed on top of the Nissl dataset with a different color. Molecular (ML) and granular (GL) layers are shown in different shades of the same color, except for CCFv3 which does not distinguish these layers anymore. Here also, the annotation boundaries do not follow the visible density changes between these layers in the Nissl data.

(C) Focus on different regions of the isocortex L1. For these regions, the neuron counts produced with the pipeline using each CCF version are greater than their literature counterpart. Using the CCFv3 version (light blue bars) yields the worst estimates. Regions shown: Primary somatosensory area, barrel field (BF); Primary visual area (VA); Primary somatosensory area (SS); Prelimbic area (PV).

(D) Focus on lingula (Ling) and flocculus (Floc) regions of the cerebellum. Results of the manual alignment of the AV2 with Nissl2 are shown with green bars. We observe that after manual realignment, estimates tend to be closer to literature values than for the other CCF versions. CCFv3 estimates are not shown, because its AV lacks the separation between molecular and granular layers.

We apply the first two steps of our pipeline (see Fig. 1), for the three pairs of annotation/Nissl volumes:

- the CCFbbp version derived from the Nissl1 and AV1. which we name Nissl1a and AV1a.
- CCFv2 version (Nissl2 + AMBA2) from the AIBS also derived from the Nissl1 and AV1. These datasets are publicly available on the AIBS website [37,38].
- CCFv3 version (Nissl2 + AMBA3) from the AIBS [33].

For each pair of reference atlases, we first estimate the densities of cells and neurons per voxel in the corresponding brain volume, based on the total number of neurons from Herculano-Houzel et al. [17]. We then compare the resulting distribution of densities to each other and to the literature (see Fig. 2CD) [39–41]. Despite an overall agreement (see S1 Fig.), we observe important differences for smaller regions. For regions with low cell density such as isocortex L1, this difference can scale up to two times more cells for CCFv3 as compared to CCFv2 (see Fig. 2C), and even more compared to some literature values. To quantify more

precisely the impact of such misalignments, we manually realign AV2 to Nissl2 for the lingula and flocculus regions of the cerebellum. We find that this indeed leads to density estimates closer to their literature counterparts (see Fig. 2D). We reproduce the figure 7A from Erö et al. [14], comparing the literature values against the corresponding estimates, using the different pairs of reference atlases. However, the results are not conclusive enough to select a reference atlas version because most of the generated cell counts agree with their literature counterparts within their range of variability (see S1 Fig.). Our earlier observations indicated that misalignments lead to overestimation of densities for low density regions and to underestimation in high density regions, reducing the differences between regions and thus, making their values more homogeneous. We quantify this by calculating the standard deviation of the neuron count distribution for each CCF. We find that the standard deviation is the largest for the version of CCFbbp ($\text{CCFbbp } 2.82 \times 10^6$, $\text{CCFv2 } 2.26 \times 10^6$, $\text{CCFv3 } 2.57 \times 10^5$) and therefore least affected by misalignment. We therefore adopt CCFbbp throughout the rest of this paper.

2.3. Alignment of *in situ* hybridization coronal brain slices

For step 3 of the BBCAv2, we consume ISH datasets from the AIBS [16] which are sectioned in the coronal plane (one 25 μm thick section every 200 μm , see Fig. 3D). However, the individual images are not aligned to the same Nissl volume that we use to estimate neuron densities. In the BBCAv1, ISH images slices were manually realigned to the Nissl volume with a non-rigid landmark-based algorithm. This task was very tedious and time consuming because it required an expert to manually choose common landmarks between adjacent sections. For our new pipeline, we therefore replace this step with a fully automatic machine learning based method.

Fig 3. AIBS GAD67 ISH experiment data.

(A) raw image, (B) filtered image and (C) realigned binarized image. This experiment highlights inhibitory cells of the mouse brain. The raw slice image (A) contains artefacts and a minimum expression value offsets the dataset. The filtered image (B) is derived from the raw image (A) by the AIBS. Somas reacting to GAD67 are detected in the raw image and isolated from the background. The different colors of the cells in (B) represent the different levels of expression (AIBS 2015 white paper). The filtered images are realigned and thresholded to obtain the binarized image shown in (C). (D) Positions of the coronal stained slices, shown in blue on a sagittal slice of the CCFbbp brain volume.

The details of this automatic alignment method are described in a separate paper [31]. In brief, this algorithm constructs a displacement field by finding pixel by pixel correspondences between an image and a reference. Applying these displacement fields as geometric transformations, images are aligned to the same reference and then converted to a 3D volume at the same resolution using downsampling.

The different steps of the automatic realignment of every ISH dataset from the AIBS to the reference atlas are as follows:

- Original ISH images (or raw images - see Fig. 3A) and their thresholded versions, later called filtered images (see Fig. 3B), are downloaded from the AIBS website. The filtered images isolate the somas reacting to their specific marker from the brain background in the raw image.
- Each image is scaled, positioned, and rotated according to the reference brain volume, using the provided metadata.
- For each dataset, every coronal section is automatically realigned to the corresponding anatomical section of the Nissl-stained mouse brain using the Krepl algorithm [31]. Here, the raw *in situ* images are used as they provide more landmarks for the algorithm.
- Resulting displacement fields are then applied to the filtered images of the corresponding ISH experiment.
- A 3D volume for each gene is created by downsampling the images to match the final voxel dimensions of the reference atlas. Missing sections between the 2D coronal slices are generated with a linear intensity-based interpolation.

The resulting 3D voxel-based datasets are consumed by our pipeline to estimate inhibitory neuron subtype densities (see Section 2.6).

2.4. Assumptions underlying the estimation of inhibitory subtypes densities

We wish to obtain the number of inhibitory neurons and their subtypes for each region of the AV. The regions are indexed by region ids $\mathbf{R} = \{0, r_1, r_2, \dots, r_n\} \in \mathbb{N}^{861}$, with 0 denoting the outside of the mouse brain. We are particularly interested in inhibitory subtypes expressing the markers GAD67, PV, SST and VIP.

First, the number of neurons can be expressed as the sum of the different subtypes in each region r , $\forall r \in \mathbf{R}$:

$$nNeu_r = nInh_r + nExc_r + nOther_r \quad (1)$$

where $nNeu$, $nInh$, and $nExc$ are respectively the total, the inhibitory, and the excitatory neuron numbers. As reported by Zeisel et al. [42], these populations seem to be mutually exclusive. The term $nOther$ represents the number of neurons that are neither excitatory nor inhibitory, meaning they could be purely modulatory neurons.

The number $nInh$ can be estimated by the number of neurons, reacting to the markers GAD67 and GAD65 [42] which implies that $nInh = nGAD$.

We further subdivide $nGAD$ into inhibitory subtypes positively reacting to the markers PV, SST and VIP. This yields the following sum, $\forall r \in \mathbf{R}$:

$$nGAD_r = nPV_r + nSST_r + nVIP_r + nRest_r \quad (2)$$

Here nPV , $nSST$ and $nVIP$ are respectively the number of PV+, SST+ and VIP+ immunoreactive neurons in the brain region r . The term $nRest$ corresponds to the number of inhibitory neurons which react neither to PV, SST nor to VIP (InhR neurons), including, for instance, the LAMP5 cells in the isocortex [32]. InhR neurons might define various populations of GABAergic cells, reacting to a great variety of markers [42], which makes it more difficult to estimate $nRest$. It is therefore easier to estimate $nGAD$ from literature and the marker GAD67 (for regions not covered by literature). Similarly, nPV , $nSST$ and $nVIP$ can be estimated from literature and the markers PV, SST and VIP, respectively. Finally, we can estimate $nRest$ by subtraction in Equation (2). The remaining neuronal populations ($nExc + nOther$) can be deduced from $nGAD$ (Equation (1)) and the neuron distribution ($nNeu$) obtained by step 2 of our BBCAv2 pipeline (see Fig. 1).

In this estimation, we rely on the following assumptions:

1. Every GABAergic neuron expresses GAD67 and every GAD67 reacting cell is a GABAergic neuron. This genetic marker is indeed responsible for over 90% of the synthesis of GABA [43,44]. Additionally, no cells expressing GAD65 without expressing GAD67 have been reported in the RNA-sequence study from Zeisel et al. [42].
2. GAD67, PV, SST and VIP are only expressed in neurons [29,42,45,46].

3. PV+, SST+ and VIP+ populations are non-overlapping i.e., there are no cells in the mouse brain that co-express a combination of these markers. This assumption is supported in the isocortex and hippocampus by transcriptomic studies such as Huang and Paul [32] and Zeisel et al. [42] and we extrapolate these findings to other areas.
4. Every PV+, SST+ and VIP+ neuron also expresses GAD67 as observed by Celio [47] and Tasic et al. [48].
5. Neuronal composition is homogeneous within subregions at the lowest level of the AIBS region hierarchy.

We will also consider the cell, glia and neuron density distributions obtained in step 2 of the pipeline (see Fig. 1) to be correct. These numbers have been validated against literature by Erö et al. [14] and can be refined in the future.

Based on these assumptions, BBCAv2 will provide estimates of the densities of GAD67+, PV+, SST+, and VIP+ neurons for each region of the brain. In contrast with BBCAv1, this new version will present estimates that are as close as possible to available literature values.

2.5. Literature review of estimates of the mouse inhibitory neuron densities

For the wild-type adult mouse, the AV1a defines 861 regions. Ideally, we would need at least one literature value for the cell density of each marker in each of these regions. We should also have multiple estimates of the same quantity to increase the reliability of the estimation. The extensive review of Keller et al. [9] provided us with references for inhibitory neuron numbers in the mouse brain (e.g., [21,49–52]). Among them, the extensive dataset of Kim et al. [8] gives a complete measurement of the distribution of the PV+, SST+ and VIP+ cells in the mouse brain, covering 97% of the regions of our selected AV. We also made a comprehensive literature search for cell densities or cell counts of GAD67, PV, SST, and VIP immunoreactive neurons in the mouse brain (wild-type species). In total, we found 32 references providing values for GAD67, 29 for PV, 14 for SST and 9 for VIP (see Table 1).

These estimates are in the form of density of neurons or cell counts within either a volume or a slice. For counts with no volume provided, we used the corresponding volume within the AV to estimate a mean density within the respective region. Finally, some literature sources report percentages of labelled neurons or cells within a region volume. For these estimates, we used the distributions obtained in step 2 of our pipeline (see Fig. 1).

	Number of papers	Number of regions covered (out of 861)	Coefficient of variation	
			Median CV	CV Skew.
GAD67	32	151 (17.5%)	0.20	1.19
PV	29	836 (97.1%)	0.24	2.94
SST	14	849 (98.6%)	0.13	3.74
VIP	9	849 (98.6%)	0.22	1.90

Table 1. Summary of papers reviewed by marker.

The number of papers from which we extract density values as input for our model are listed in the first column. The table also shows how many regions of the brain are covered by these values. For each region where we obtain an estimate from at least two different papers, we can compute a coefficient of variation. From the list of these coefficients of variation, we then extract the median values and the skewness of the distribution, defined as:

$$s = \frac{1}{n} \sum_{r \in R} \left[\frac{CV_r - \text{mean}(CV_r)}{\text{std}(CV_r)} \right]^3$$

which are displayed for each marker in the last two columns.

Erö et al. [14] considered the neurons of some regions to be inhibitory only (step 6 of the BBCAv1 workflow). Similarly, we imposed this rule on layer 1 of the isocortex, the molecular layer of the cerebellar cortex and the reticular nucleus of the thalamus. This means that for these regions, we assume that the density of GAD67+ neurons corresponds to its neuron density. Thus, in total we collected GAD67+ density estimates for about 17.5% of all regions of the AV (see Fig. 4A).

Fig 4. Overview of published data on inhibitory neuron densities in the mouse brain.

(A) Illustration of regions for which information on densities or absolute numbers of GAD67+ neurons were published. The disk is divided into rings and sectors. Each ring represents a hierarchical level in the AV and sectors represent the contained regions. The center of the disk represents the entire brain, each surrounding ring then represents the next hierarchy level. Colored areas represent regions where at least one study reports cell numbers or densities. The size of the sectors corresponds to the number of neurons relative to the number of neurons in the brain. The colors correspond to regions according to the AV such as the cerebellar cortex in yellow and cortical areas in green.

(B) Variability of published cell density estimates. Each panel shows the distribution of reported cell densities (log scale). For each region, data points are depicted in blue, the average in orange. The minimum and maximum values are indicated by the whiskers. Regions shown: Dentate Gyrus (DG); Field CA1 (CA1); Field CA3 (CA3); PreLimbic area (PL); Primary SomatoSensory area (SSp); Primary VISual area (VISp); STRiatum (STR); SUBiculum (SUB);

Primary somatosensory area, barrel field (BF); Lateral Amygdalar nucleus (LA); Cerebellar cortex, Purkinje layer (CBXpu).

For some regions such as the CA1 field of the hippocampus, we observe a large variability of the literature estimates (see Fig. 4B): spanning from 317 GAD67+ cells per mm³ computed from Han et al. [53] to 7166 GAD67+ cells per mm³ according to Jinno et al. [54] (see the complete list of literature data in the review S1 Excel). In addition to the inter-subject variability, these large variations can arise from a combination of different factors, such as differences in the staining methods, the counting methods, or the methods to determine subregion boundaries (annotations) [7,8].

Our literature review highlights the current lack of knowledge and consensus on cell composition in the mouse brain, emphasized by the large variability of published data. In the following section, we will describe a method to estimate cell densities in regions for which no literature values were published.

2.6. Transfer functions from marker expression to cell density

We use the realigned ISH datasets obtained in Section 2.3 to estimate the densities of inhibitory cell types in regions where no literature values were reported.

Different cell types express genetic markers at different levels [42]. Nonetheless, we count every cell reacting to the markers. We do this by applying an all-or-none filter to the realigned filtered image stacks (Fig. 3B) using Otsu's method [55] to define the threshold. This forces each pixel showing a cell reacting to the marker to the value 1 and all other pixels to 0 (binarized image stacks see Fig. 3C). The more pixels in a region are set to 1, the more densely the region is filled with marker-positive neurons. We therefore expect, for a specific brain region, that its mean pixel intensity correlates with the density of marker positive cells. We use this relationship to construct transfer functions linking densities of GAD67+, PV+, SST+, and VIP+ cells reported in the literature to the mean intensity in each region of the AV (see Fig. 5).

We leverage every literature density value available for each subregion of the brain, assigning the same weight to each point. We compare separately the cerebellum, isocortex and the rest of the brain. We choose this division because in the cerebellum, cell densities are very high compared to the rest of the brain [17]. And, inhibitory cell densities and cell type composition across layers of the isocortex are similar in all of its subregions [8]. The results of the comparison between region mean pixel intensity and cell type density are shown in Fig. 5.

We expect a region with no inhibitory neurons to have a null mean intensity and conversely a region with null mean intensity should not host any inhibitory neurons. However, we do not observe this property in many cases. This observation can be linked, on the one hand, to the misalignment between the annotation atlas and the ISH datasets (cells reacting appear in the wrong region) or on the other hand, to the insufficient coverage of the brain by the ISH data (region of interest does not appear on the ISH slices, see Fig. 3D). Therefore, these points are excluded from Fig. 5. Additionally, as we discussed in Section 2.2, the density estimation in fully inhibitory regions tends to be highly impacted by misalignment. The points linked to these regions are therefore also excluded from the figure.

At the first glance, there seems to be no clear relationship. However, as we have seen above there is a large variability in the cell densities reported by literature (see Fig. 4B). This explains the spreading of the points along the y axis in the different panels of Fig. 5 (see Fig. 5, different point colors). Similarly, points along the x axis are also spread out because of the sparseness of the ISH datasets (see Fig. 3D) and its misalignment to the AV (see Fig. 2). This variability might also be linked to the different cell sizes which impacts the region mean intensity. These two sources of error make it difficult to estimate a transfer function that maps region mean intensity to cell type density. Logically, we expect that there is a monotonically increasing relationship between region mean intensity and cell type density — one for each marker and each main division of the brain. The simplest assumption is that these relations are linear and can be calculated by fitting (black diagonals in Fig. 5).

To evaluate the quality of the fit from region mean intensity to densities of cells, we compute the coefficient of determination R^2 for each linear function f , which fits the cloud of points (x, y) where x is the region mean intensity and y is the literature density:

$$R^2 = \frac{\sum_{i \in x} (f(i) - \bar{y})^2}{\left(\sum_{i \in x} (f(i) - \bar{y})^2 + \sum_{i \in x} (y(i) - f(i))^2 \right)} \quad (3)$$

The low values of R^2 especially for GAD67 indicates that the great variability of (x, y) could not be fully accounted for (roughly 25% of the variability accounted for GAD67, see Table 2).

	Cerebellum		Isocortex		Rest	
	alpha	R^2	alpha	R^2	alpha	R^2
PV	4.557×10^5	43.1%	2.964×10^5	46.4%	1.182×10^5	47.9%
SST	1.417×10^5	33.0%	2.523×10^5	50.6%	3.158×10^5	34.3%
VIP	7.903×10^4	26.4%	2.311×10^5	37.4%	1.154×10^4	56.5%
GAD	5.783×10^5	27.6%	3.458×10^5	37.0%	5.265×10^5	13.5%

Table 2: Results of the linear fitting

The slope factor, alpha, of the linear fitting, and its coefficient of determination R^2 for each major region of the brain and for each genetic marker.

Thus, we use the fitted relation only to predict densities of GAD67+, PV+, SST+ and VIP+ in regions where no literature data are available. The standard deviations of these estimates are derived from the standard deviation of the alpha value of the fit (see Table 2 and S2 Excel). For regions covered by the literature we instead use the average of all means and standard deviations of the reported values (see previous section). We assume indeed that the different literature values scatter around a common average.

Therefore, at this stage we have obtained estimates of density (orig) and their standard deviation (σ) for PV+, SST+, VIP+ and GA67+ neurons for all regions, i.e. $\forall r \in \mathbf{R}$:

$$orig_r = \{nPv_r; nSST_r; nVIP_r; nGAD_r\} \quad (4)$$

$$\sigma_r = \{std(nPv_r); std(nSST_r); std(nVIP_r); std(nGAD_r)\} \quad (5)$$

orig and σ are lists of 3444 values: one per neuron type (PV, SST, VIP, GAD67) and per region of the brain (861 in AV1a).

2.7. Combination of neuron type densities

Since our original estimates of inhibitory neuron (orig and σ) densities are derived from the transfer functions or independent literature sources, their combination can lead to incongruent results. For instance, the original estimates of PV+ neurons might be greater than the estimates of GAD67+ neurons, which contradicts our assumption 4 (see Section 2.4). We therefore want to ensure that orig and σ match our assumptions from Section 2.4. To this end, we deduce a list of linear constraints based on equations (1) and (2), $\forall r \in \mathbf{R}$:

$$0 \leq nPV_r \leq nNeu_r \quad (6)$$

and similarly for $nSST_r$, $nVIP_r$ and $nGAD_r$.

$$-nNeu_r \leq nPV_r + nSST_r + nVIP_r - nGAD_r \leq 0 \quad (7)$$

We also ensure that the consistency of the region hierarchy is maintained for every parent region ($R_m \in \mathbf{R}$). The number of inhibitory neurons (and their subtypes) in R_m is equal to the sum of the corresponding estimates of R_m 's direct child regions in the region hierarchy ($children_{R_m}$) plus the estimates in voxels of the AV labeled as belonging to R_m but none of its children ($R_m \setminus child$):

$$nPV_{R_m \setminus child} + \sum_{r \in children_{R_m}} nPV_r = nPV_{R_m} \quad (8)$$

and similarly for $nSST_{R_m}$, $nVIP_{R_m}$ and $nGAD_{R_m}$.

Thus, whenever one of the constraints (6), (7) or (8) is violated, we rescale our original estimates (orig).

We minimize the amount of corrections required to find a solution for the BBCAv2 model through optimization. The amount of corrections is defined as the sum of the distances between the original (orig) and corrected values counterparts (x) divided by the standard deviation of the original value (σ):

$$\min. \sum_{r \in \mathbf{R}} \frac{|x_r - orig_r|}{\sigma_r} \quad (9)$$

This nonlinear function is convex which guarantees a global minimum. We convert it to a linear problem, by introducing a slack variable z so that $\forall r \in \mathbf{R}$:

$$-z_r \leq x_r - orig_r \leq z_r \quad (10)$$

Hence, our problem becomes:

$$\min. \sum_{r \in \mathbf{R}} \frac{z_r}{\sigma_r}$$

(11)

st. Equations (6), (7), (8), (10)

The solution x is a vector with 3444 values and matches a total of 8613 linear constraints. An initial solution can also be found for each region, correcting original estimates to match our constraints starting from leaf regions in the region hierarchy to the top-level brain regions (see supplementary methods S2 Fig. and S3 Fig.). We use the simplex algorithm from the scipy python library [56].

We find that for 16% of the regions of the brain (shown in colors in Fig. 6AB), their original estimates are incongruent with the rest of the brain. Their corrected values x are therefore not falling within the range $orig \pm \sigma$. In some regions of the striatum (including caudoputamen), the number of GAD67 neurons from orig overshoots the estimated total number of neurons. Our optimization makes these regions fully inhibitory. This is in line with literature findings on the mouse striatum, which describe it as almost fully inhibitory [57]. Among the rest of the regions where significant corrections are needed, almost one-third are subregions of the hindbrain, for which we collected very few literature values. Conversely, few inconsistent first estimates are found in subregions of the isocortex, for which more literature data are available. The remaining inconsistencies for orig may have different sources:

- They can be explained by a poor estimate of the densities from the transfer function.
- Divergent estimates from the literature for the same region might violate equation (7). These regions would appear in yellow on Fig. 6A).
- Our assumptions may not be accurate in some regions of the brain.

- The densities of neurons from the step 2 of the pipeline (see Fig. 1), used in equation (6) and (7), diverge locally from the literature findings.

Fig 6. Results of the density pipeline.

(A) Sagittal and (B) coronal view of the Nissl reference atlas used for BBCAv2, showing cells of the mouse brain. Regions with high cell density appear in dark grey. Regions that required significant corrections (see Section 2.7) for their neuron subtype density estimates are colorized (Neu = neuron). The coronal slice chosen is displayed as an orange line on the sagittal slices. Arrow in black shows the location of the lateral ventricle of the mouse brain. The blue line behind the sagittal slice highlights the drop of Nissl expression coming from the original Nissl experiment from Dong [15].

(C). Sagittal and (D). coronal view of the BBCAv2, showing the positions of the different types of neurons. PV+, SST+ and VIP+ cells appear respectively in yellow, dark orange and blue. The rest of the GABAergic population is color-coded in green and the remaining neurons in gray.

The variation of the distribution of neurons is following the original distribution of Nissl expression. The Nissl expression drops significantly after the blue line which leads to a similar decrease of cell counts in the resulting BBCAv2.

After our corrections, we obtain consistent density estimates for PV+, SST+, VIP+, and GAD67+ neurons for each region of the AV.

2.8. Placement of the cells within the mouse brain volume

At step 4 of the pipeline (see Fig. 1), we convert our density estimates into a list of cell positions for each region, using an acceptance-rejection algorithm. For each region, the algorithm samples voxels, according to the cell density distribution (computed at step 2), and assigns cells to them, until it reaches the target total number of cells in that region. The same algorithm can then be applied to label cells as neurons or glial cells as described in Erö et al. [14]. We further subdivide neurons into PV+, SST+, VIP+, and GAD67+ neurons, following our density estimates. In brief, $\forall r \in \mathbf{R}$, we select uniformly neurons within the voxels of region r , that we label as inhibitory, until the region contains the target number of $n\text{GAD}_r$ inhibitory neurons. Similarly, we distribute the $n\text{PV}_r$ neurons, in voxels of r containing GAD67 neurons, $n\text{SST}_r$ in voxels of r containing non-PV inhibitory neurons, and $n\text{VIP}_r$ in voxels of r containing the remaining inhibitory neurons. The non-inhibitory neurons are labelled as excitatory or purely modulatory neurons following Equation (1). Finally, we assign to each cell a uniformly random 3D position within the boundaries of its voxel. The result of this procedure is shown in Fig. 6CD.

3. Results

3.1. An improved computational method to build an atlas of inhibitory neurons in the mouse brain

In this study, we present a substantially improved workflow to produce the second version of the Blue Brain Mouse Cell Atlas workflow (BBCAv2). The new workflow replaces the manual alignment of ISH datasets with the image alignment algorithm developed by Krepl et al. [31]. The pipeline also includes a new method to estimate cell densities from a large body of literature data. In total, we integrated values from 56 different literature sources.

All steps of this process are guided by a set of assumptions about the cellular composition of the mouse brain. Using our improved workflow, we are not only able to re-estimate the densities of inhibitory neurons in the mouse brain, but also to estimate the densities of their various subtypes. The result is an updated mouse cell atlas which provides a 3D position and a type label for each cell within the CCFbbp AV (see Fig. 6CD). The same methods can also be applied to other cell types to deepen our knowledge on cellular composition of the mouse brain. The new version of the Blue Brain Mouse Cell Atlas is available on the Blue Brain Portal website (<https://bbp.epfl.ch/nexus/cell-atlas>). The code used to produce these results can also be downloaded from <https://bbpgitlab.epfl.ch/nse/atlas-building-tools>.

3.2. Distribution of inhibitory neuron subtypes for the whole mouse brain

The new cell atlas has updated estimates for the densities and numbers of inhibitory neurons, based on more literature data. In addition, the new atlas differentiates PV+, SST+ and VIP+ neurons and provides their densities and absolute numbers for each region of the mouse brain. An overview of the results is shown in Fig. 7 and Table 3. The complete list of densities for all cell types in all regions of the Allen Mouse Brain Atlas is available as supporting information (see S2. Excel). We estimate that the mouse brain holds approximately 14.55 million of

inhibitory neurons in total (see Table 3). GABAergic neurons make up 20.27% of the total neuron population, which is significantly larger than the previous estimate of 15.68%, using the BBCAv1 workflow. We find that 3.57% of neurons are PV positive, 3.20% are SST positive, and 0.63% are VIP positive (see Fig. 7A and S2 Excel). This leaves a residual GABAergic neuron population of 12.87%, which represents almost two-thirds of all inhibitory neurons.

Fig 7. Distribution of the GABAergic neurons in the mouse brain.

(A) Ratios of PV+, SST+, VIP+, GAD67+, and remaining neurons of the mouse brain. InhR cells appear in green, PV+ in yellow, SST+ in orange, and VIP+ in blue. The remaining neurons appear in grey.

(B), (C) Circular distribution of GAD67+ neurons in different regions of the mouse brain according to different parcellation schemes ranging from coarse at the center to fine-grained at the periphery (colors and disposition similar to Fig. 4A). (B) displays results of BBCAv1 and (C) results of BBCAv2.

Brain Region	BBCAv1	BBCAv2					
	Inh.	Inh.	$\frac{\text{Inh}}{\text{Neu}}$ (%)	$\frac{\text{PV}}{\text{Inh}}$ (%)	$\frac{\text{SST}}{\text{Inh}}$ (%)	$\frac{\text{VIP}}{\text{Inh}}$ (%)	$\frac{\text{Rest}}{\text{Inh}}$ (%)
Isocortex	1.44	2.20	24.7	23.2	25.0	12.6	39.2
Olfactory areas	1.45	2.05	25.1	1.3	13.8	4.7	80.2
Hippocampal form	0.51	0.49	9.9	12.6	33.9	9.0	44.5
Cortical subplate	0.12	0.22	43.8	2.6	28.3	3.7	65.4
Striatum	1.72	1.40	81.4	1.2	17.5	0.2	81.1
Pallidum	0.18	0.23	91.8	6.9	28.7	0.2	64.2
Thalamus	0.19	0.26	18.9	15.2	32.7	0.0	52.1
Hypothalamus	0.42	0.50	41.4	3.0	29.0	0.4	67.6
Midbrain	0.39	0.99	76.4	12.6	37.4	0.9	49.1
Hindbrain	0.26	0.66	62.5	23.9	38.2	0.6	37.3
Cerebellum	4.79	5.47	13.0	28.8	0.8	0.1	70.3
Whole Brain	11.25	14.55	20.3	17.6	15.8	3.1	63.5

Table 3: Estimates of inhibitory neurons in the mouse brain in millions and percentage (%) subtypes

Inhibitory neuron counts (in millions) extracted from the cell atlas generated using the BBCAv1 and BBCAv2 pipelines. The second column of the BBCAv2 corresponds to the ratio of inhibitory neurons according to the total number of neurons in this region. The following columns display the proportion of each inhibitory neuron subtype according to the inhibitory neuron population.

Compared to BBCAv1 (Fig. 7B), the number of inhibitory neurons in BBCAv2 (Fig. 7C) is larger in almost all regions of the brain (except notably, the hippocampal formation and the striatum). The estimated number of inhibitory neurons in the cerebellar cortex granular layer increases from 203,000 in the previous pipeline to 867,000. Similarly in the isocortex, this number rises roughly by 53% (see S2 Excel). Finally, the BBCAv2 estimates 2.05 million inhibitory neurons in the olfactory areas.

3.3. New Excitatory/Inhibitory ratios for the isocortex

The BBcAv2 also provides new estimates for the ratio of inhibitory/excitatory neurons (see Fig. 8). For example, we predict that 25% of the neurons of the mouse's isocortex to be inhibitory (see Table 3 and Fig. 8B). Thus, we estimate a higher proportion of inhibitory neurons than previously reported. For rodents, literature estimates are around 20% inhibitory neurons [11,58].

Fig 8. Cortical excitatory/inhibitory ratios.

(A) Relative cortical layer cell densities from the primary somatosensory cortex lower limb (SSp-II), primary auditory cortex (AUDp) and temporal association areas (TEa). Blue lines correspond to excitatory neuron densities (Exc) and the orange lines inhibitory neuron densities (Inh) in cells/ mm^3 . Final confidence intervals are also displayed for each layer as vertical lines.

(B) Ratios of inhibitory/excitatory neurons (Inh/Exc) across anatomical regions of the isocortex, expressed as percentage and arranged in five subnetworks based on Kim et al. [8]. Acronyms correspond to the AV naming convention [36] (see S1 Table). The portion of inhibitory neurons that belongs to L1 is shown with a black contour.

(C) Somatosensory barrel field neuron distributions according to pia. Each layer is represented by a different color and the total density according to distance to pia appears in black.

(D) Somatosensory barrel field excitatory/Inhibitory neuron distributions (Exc/Inh) according to pia compared to the results extracted from Fig. 2EFG from Meyer et al. [58]. AV mean layer limits are represented as dash lines and aligned to the ones provided by Meyer et al.

The poor alignment of the AV to the Nissl dataset counterpart (including the layer 1 layer 2 boundary, see Section 2.2) may explain this result. It is artificially raising the density of neurons in layer 1, which in turn, increases the ratio of inhibitory neurons in the isocortex (black contour in Fig. 8B). Additionally, the densities of PV+, SST+ and VIP+ neurons, mostly constrained by Kim et al. [8], are providing a lower limit for the total number of inhibitory neurons in each region (green bars in Fig. 8B). To improve on the ratios of inhibitory neurons in the isocortex we should therefore check its densities of neurons and our predictions of GAD67 neurons in those regions.

We also notice that the estimated densities of inhibitory neurons are quite similar from layers 2 to 5, and across isocortex subregions (Fig. 8A and S6 Fig.). We further investigate the neuron density profiles according to the distance of their soma position according to the pia (see additional methods and Fig. 8CD). We manage to reproduce the findings of Meyer et al. on rats in the barrel cortex (see Fig. 8D) [58].

3.4. Validation of the pipeline

In our revised workflow for BBCAv2, we used a linear transfer function to map ISH expression levels to literature cell density to then estimate the cell densities in regions where no literature value is available. The fidelity of the transfer function depends on both the quality of the ISH images and the quality of the underlying literature estimates of the cell densities. As we highlighted in Section 2.3, the genetic marker datasets are prone to artefacts due to slicing.

Also, their coverage of the mouse brain is not perfect (see Fig. 3D). Combining multiple datasets of the same genetic marker can blur the spatial details of brain structures. For each marker, the AIBS website hosts similar ISH experiments in sagittal (Fig. 10A) and coronal (Fig. 3A) slices. We compare the results of our pipeline, using two different coronal ISH datasets for GAD67 and find very similar results in terms of fitting, distribution of inhibitory neurons and remaining errors after corrections (see S2 Excel). To blur the individual artefacts of the ISH datasets, we combine these datasets after realignment, taking the mean of their expressions. We then linearly fit the marker expression from the resulting dataset to literature densities but we find no significant improvement. This result can be explained by the slicing of these two experiments which, in both cases, is not fully covering the tips of the olfactory bulbs and the cerebellum as illustrated in Fig. 3D.

Next, we explore the impact of the amount of literature values available on the quality of the transfer function.

We do this experimentally by running our workflow with a fraction f of the available literature points (70% up to 95%). For each fraction f , we perform 20 separate experiments, each with a different random subset of literature values. Thus, for each fraction f , we obtain 20 different estimates of the cell density in each brain region $r \in \mathbf{R}$. We also compute the standard deviation of the resulting density values for each region ($\sigma_{(r, f)}$). As we increase the number of literature values that are contributing, we expect the standard deviation of the density estimates to decrease. We observe that the average of all standard deviations for each marker gets smaller as more literature values are integrated (see Fig. 9A).

Fig 9. Impact of the amount of literature on the BBCAv2.

(A) Evolution of the mean standard deviation of each genetic marker (shown in different colors) using different percentages of the total amount of literature values available.

(B) Comparison of the generated density values of the BBCAv2 with their literature counterparts using 90% of the total amount of literature values available (20 trials). The color encodes the brain regions according to the AV, while the shapes of the points encode for cell types. The middle line delimits equal quantities, while the dashed line shows the average deviation of 2.7-fold between literature values reporting on the same region for PV, SST and VIP neurons.

(C) Evolution of the coefficient of determination R^2 . Each line corresponds to the R^2 value of the fitting performed using different percentages of the total amount of literature values available. The color of these lines is linked to the gene chosen and their color shade corresponds to their fitting group (here Isocortex and Rest, as in Fig. 5).

For each region r , we then use all $\sigma_{(r, f)}$ to extrapolate the standard deviation of the density for $f=100\%$. This standard deviation $\sigma_{(r, 100\%)}$ represents the confidence of our pipeline in our final predicted densities for each region (see S2 Excel).

Surprisingly, the quality of the fitting does not significantly improve with more literature values added to the pipeline (see Fig. 9C). This result indicates that the coefficient of determination is mostly influenced by the neuron distribution obtained at step 2 (see Fig. 1), the ISH

experiments or the method to group the regions (here cerebellum, isocortex and rest, see Fig. 5).

We also test the capacity of the BBCAv2 to predict values of literature. We select 90% of the literature points available, and then compare the generated densities to the remaining points (see Fig. 9B). Most of the density values produced with this method fall within the confidence interval of their literature counterparts. This workflow is therefore capable of predicting densities of cells within range of what literature provides.

As an additional qualitative validation, we compare our generated GABAergic neurons positions with a sagittal slice of GAD67 expression from the AIBS (see Fig. 10AB) and find a similar whole brain distribution. Some of the visual differences between Fig. 10A and Fig. 10B can be explained by the fact that in the image from the BBCAv2 (Fig. 10B), the soma sizes are uniform across all cells, while in the brain, soma sizes differ considerably between cell types and brain regions (Fig. 10A). Moreover, the AV1a is based on the coronal Nissl experiment from the AIBS (see Section 2.2) which is not well covering the tips of the cerebellum and the olfactory bulbs (see Fig. 6A) and deforming them. However, this comparison highlights the fact that in our model the thalamus and the cerebellum have too many inhibitory neurons. Additionally, the granule layer of the dentate gyrus is clearly visible on the original experiment but not in the BBCAv2 brain slice (center black arrow). The original slice in panel A shows spatial detail with higher resolution than our reconstruction in panel B. For example, in the olfactory bulb, panel A shows the layering of cells, as we do in the cerebellum and hippocampus. The spatial resolution of our cell atlas is lower, because the density is assumed uniform within regions at the lowest level of the AIBS region hierarchy (assumption 5), following the estimated cell density.

Fig 10. Qualitative validation of the BBCAv2 and comparison with BBCAv1.

GAD67 sagittal brain slices comparison between (A) the filtered image from an ISH experiment by the AIBS at the top and (B) the BBCAv2 predicted GABAergic neurons positions at the bottom. Cells reacting to the marker are displayed in purple. Blank regions in our model correspond to fiber tracts regions which host no neuron. This slice does not take regional soma sizes into account. Most of the density features of the real experiment can be found in the BBCAv2 slices. Black arrows point to the thalamus (left arrow), molecular layer of the dentate gyrus (center arrow) and granular layer of the cerebellum (right arrow) of each brain.

(C) Circular distributions of the densities of PV+, SST+ and VIP+ neurons computed according to the Erö et al. method on the left [14] and our method on the right (colors and disposition as in Fig. 4A).

Distribution of PV+, SST+ and VIP+ neuron density values reported in literature against generated densities, using the BBCAv1 method (D) and the BBCAv2 method (E), for similar regions. The color encodes the brain regions according to the AV, while the shapes of the points encode for cell types. The middle line delimits equal quantities. In (E) the model is built on the regional literature references used in the Figure which explain the numerous points on the middle line.

We also compare the estimates of PV, SST and VIP densities from the BBCAv2 against similar estimates with the BBCAv1 [14]. For the BBCAv1, we use the global number of PV, SST and VIP neurons in the mouse brain from Kim et al. and the PV, SST, and VIP realigned datasets (see Section 2.3, Fig. 10C and S2 Excel) [8]. The BBCAv1 maintains the global number of inhibitory subtypes expected in the whole brain at the cost of losing the local distribution (see Fig. 10D). For instance, Kim et al. [8] found that the cerebral nuclei regions contain 62 VIP neurons per mm³, while the BBCAv1 predicts it to contain 963 VIP neurons per mm³ (see Fig. 10C, cerebral nuclei regions appear in light blue). The BBCAv2 method, on the other hand, aims to maintain local estimates from literature in a coherent atlas (see Fig. 10E). It is therefore usually underestimating the densities of each neuron type according to literature, since the neuron and inhibitory neuron densities act as maximum constraints.

Our model integrates literature values in the coherent framework of the cell atlas, with constraints and assumptions that we control and that we can refine whenever more data become available about the mouse brain.

4. Discussion

In this study, we show that our new method is capable of estimating inhibitory neuron counts for each region of the mouse brain, including PV+, SST+, VIP+ neurons, and remaining inhibitory neuron subtypes, in a consistent framework: The Blue Brain Mouse Cell Atlas (BBCAv2). The results of our method are constrained by literature findings (see Section 2.5), ISH experiment data from the AIBS (see Section 2.3) and simple assumptions (see Section 2.4). BBCAv2 extends the results of BBCAv1 and deepens our knowledge of the cellular composition of the mouse brain. Additionally, the BBCAv2 can be easily adapted to estimate the densities of other cell types, providing more details about the composition of the brain in terms of morphological, molecular, or electrical properties.

4.1. A new approach to integrate literature findings into the Blue Brain Cell Atlas

We presented a new approach to reach a consensus on cellular composition in the mouse brain based on local literature findings instead of global constraints. To compensate for the great variability of cell density measurements in literature and to avoid a bias towards any particular finding, we decided to use as many data sources as possible (see Section 2.5). The BBCAv2 therefore reflects our current knowledge about cell composition in the mouse brain. However, this makes it also more difficult to validate, since only limited data or studies remain available to compare our results to. Moreover, any additional literature value that we could use to validate our counts might itself only be as trustworthy as the literature values used to build our model. On the other hand, qualitative validations can highlight inconsistencies in our results as shown in Fig. 10AB.

4.2. Reference datasets' impact on cell densities

Both Blue Brain Mouse Cell Atlas pipelines are based on a pair of reference atlas datasets from the AIBS which exist in multiple versions. We assessed newer versions of the AV and analyzed their impact on cell densities calculation.

We found that the best cell density estimates can be obtained if the pair of reference datasets (Nissl volume and AV) is perfectly aligned to each other. For example, if the border between two regions with significantly differing densities (e.g., isocortex L1 and L2, see Fig. 2AC) is not perfectly aligned with the corresponding Nissl, then the densities in the particular regions will be over- or underestimated. Hence, the CCFv3 AV, while depicting a smooth average brain volume, provides the worst density results when combined with Nissl2.

The Nissl reference volumes are stretched and sometimes torn by the slicing, which yields a very noisy dataset. The derived annotation atlas had to follow this roughness with sometimes discontinuous region borders. The resulting AV are therefore less attractive for circuit modeling. We can still benefit from the best of each dataset, computing the densities using the CCFv2 or CCFbbp and then populate the CCFv3 smooth brain volume with these estimates. As for our assumption 5, cell distribution in the lowest level of the region hierarchy will be considered homogenous. Potential inner region gradient can be kept if a common vector base is defined in each AV version. Another solution could involve the realignment of the AV2 to the AV3. With this work, we will recover the subregions of the AV2 that were not found in AV3 (e.g., cerebellar cortex, hippocampus CAs).

In Fig. 6A (blue line), we can also see another default of the Nissl dataset in the form of a drop of expression in slices between the olfactory bulbs and the thalamus which seems to be unrelated to the regions they are crossing (see also S4 Fig. ABC for more details). This drop of expression is then clearly visible in the resulting distribution of neurons in Fig. 6B. Yet, the neuron counts estimates are used in the rest of the pipeline as a top constraint for our inhibitory neuron distribution, which can partially explain some of the errors we encounter at the final stage. Finding other Nissl datasets to create an average Nissl volume or exploring image

processing techniques such as histogram matching and blurring should be part of future works to improve the model.

4.3. Challenges linked to ISH marker expression datasets

We applied Krepl's algorithm [31] to realign the filtered ISH images of the mouse brain from the AIBS at step 1 of our method. The resulting datasets are crucial for the BBCAv2 pipeline as they allow cell density estimation in the regions missing literature values.

The ISH experiments cover partially the whole mouse brain, with one slice stained per 200 μm in the rostro-caudal axis (see Fig. 3D)[16]. Additionally, they represent different mouse individuals whose neuron distribution might vary a lot with respect to one another. As we discussed in Section 3.4, the combination of similar ISH datasets can improve their coverage of the brain and blur out the remaining artefacts. Furthermore, ISH experiments performed with a sagittal slicing covers regions not included in the coronal experiments and their addition will improve the ISH whole brain coverage. Using these datasets requires the realignment of the sagittal images of the inhibitory markers to the Nissl volume, using the deep learning algorithm described from Krepl et al. [31]. The usage of these datasets is currently investigated for a future update of the pipeline.

The marker datasets, despite having been processed by the AIBS, are still subject to artefacts and noise which will artificially raise our cell counts estimates (see Fig. 3A). To counter these effects, Erö et al. [14] used the *Nrn1* marker, as it is expressed in excitatory neurons, to limit the expression of GAD67. The GAD65 marker has also been reported to be expressed in GABAergic cells [59]. Similarly, we can extend our approach, knowing the density of neurons expressing *Nrn1* and GAD65 in regions of interest, to correct our estimates of GABAergic neurons.

We also chose to use an all-or-none filter on the filtered images because we were interested in counting all the cells expressing the marker. However, we could deduce counts of

subpopulations of neurons which express their marker at specific levels. These levels might depend on the regions or the cell types but it is an opportunity to improve the BBCAv2 locally. Future analysis will include estimating cell density from the ISH images using automatic point-detection algorithms as presented in Erö et al. [14]. These algorithms will provide new cell counts to rectify our estimates. However, cell counting requires significant resources to be applied to the whole brain, and will also be biased by cell overlapping, especially in highly dense regions such as for Cerebellum, as discussed in Erö et al. [14].

4.4. Limitations of the assumptions of the pipeline

The BBCAv2 estimates of the different cell densities are based on the five assumptions described in Section 2.4. These assumptions come from findings mostly published on the isocortex and represent the rules we used to categorize all cells in the mouse brain [29,32,42–47]. However, for each of our first four assumptions, we found exceptions reported in literature:

1. A small portion of astrocytes from the visual cortex have been reported to express GAD67 [60].
2. Szabolcsi and Celio [61] observed that ependymal glial cells express PV. These cells are surrounding the ventricles of the mouse brain which regions are present in the AV. No neurons were placed within these regions as can be seen in Fig 6; these cells should therefore have a minor impact on our results.
3. Zeisel et al. [42] reported that PV, SST, and VIP populations overlap in most regions of the brainstem. There are many types of cells co-expressing at least two of our markers with different levels of expression and we found no automatic solution to detect each of these cells using the data available, since the ISH data from the AIBS comes from different species. Lee et al. [62] also pointed out, using double ISH staining, that a population of neurons in the isocortex is co-expressing PV and SST.

4. Zeisel [42] also found that PV and SST are expressed in excitatory neurons and VIP in modulatory and non-inhibitory neurons in some regions of the brainstem such as the colliculus and medulla, which were detected as errors in our reports (see Fig 6A).

If validated by additional studies of density estimates in the regions where these exceptions have been reported, we could justifiably subtract a proportional amount from the deduced numbers of GABAergic cells in future versions of the model.

As we describe in Section 4.2, some of the inner region variation of cell densities can be lost applying the assumption 5. This concerns gradients within the neuronal population since the neuron distribution is derived from the Nissl volume. For instance, in the hippocampus CAs, Jinno et al. [54] have shown a dorsal to ventral distinction in the density of the inhibitory neurons, which is not reflected by the subdivision of the AV. Additional subregions for the AV have to be created to include these findings.

4.5. Reliability of quantitative stereological studies

One of our objectives was to match regional estimates from literature with our neuron types' density distribution. To do so, we performed a literature review of quantitative stereological studies on PV+, SST+, VIP+, and GAD67+ neurons in the mouse brain (see Section 2.5).

We found that literature does not provide a complete coverage of the whole mouse brain for GAD67. This can be explained by the difficulty to perform cell counting or cell detection on large tissue volumes, or by the fact that some regions are more frequently studied than others (e.g., barrel cortex, hippocampus CAs, see Fig. 4B). Moreover, GAD67, PV, SST and VIP are extensively used genetic markers in literature, this is not the case for most of the markers available (e.g., LAMP5). This means that the more precise our cell composition becomes the more difficult it will be to gather enough data points from literature and obtain meaningful transfer functions (see section 2.6).

We also found that studies reporting similar measurements of neuron densities sometimes yielded very disparate results (see Fig. 4B and Table 1). Literature density estimates can be

biased by the age, sex, staining technique, neuron counting method, or measurement of the volume of the region of interest [7,8].

Some of the densities used to build our transfer functions were computed according to cell counts from literature divided by the volumes of the regions of the AV, and potentially prone to errors as we pointed out in Section 2.2. Some regions are also known to be purely inhibitory (e.g., layer 1 of the isocortex) and we therefore used the Cell Atlas neuron densities, which are themselves subject to errors as discussed previously. Future work on the reference atlases should improve the reliability of the literature estimates.

4.6. Variability of the transfer functions

Fitting a transfer function from region mean intensity to cell type density is a required process to obtain the density of inhibitory cell type in regions where no literature data are available (see Section 2.6). However, we found that the fitting of the transfer functions from marker expression to cell densities is sensitive to errors linked to the datasets used as input which can explain some of the BBCAv2 results and the fitting low coefficient of determination.

In future work on the BBCA, the transfer functions can be improved by creating more subregion groups for which separate fittings will be done. These subgroups could correspond to different expected soma sizes for the cell type of interest. However, this would reduce the number of points available to fit the function and might result in an overfit.

Other monotonically increasing functions can also be tested to fit the cloud of points of Fig. 5 such as sigmoid functions to integrate any threshold dynamics for low expression values or saturation for high expression values.

4.7. Consolidation of neuron density estimates

We described a method in Section 2.7 to reconcile disparate density estimates into the consistent framework of the BBCAv2. This method aims to minimize the amount of correction needed to obtain a coherent solution.

In our minimization function (11), we chose to divide the differences measured according to the standard deviation of our first estimates (σ). This parameter accomplishes two purposes: first, it reduces the impact of dense regions on the global minimization score, and second, it gives more weight to estimates where either literature or the fitting are more confident. Other σ values can be tested in the future, such as the total number of neurons in the region to limit the impact of neuron density and in turn, give the same relative weight to each region. Similarly, we can measure the impact of adding an extra weight to maintain the proportion of each cell type within a region. This new constraint will have to be balanced with the previous one. We assumed that the distribution of neurons in the mouse brain using the BBCAv1 method is correct, which means that whenever there was a contradiction between this distribution and the BBCAv2 inhibitory neuron estimates (coming from literature or the fitting), the error will be attributed to the latter. BBCAv2 results are therefore biased by BBCAv1. The results of Erö et al. have been validated against literature counterparts [14] but might be incorrect in some specific regions (as depicted in Fig. 10C). The BBCAv2 seems however to be more in agreement with literature than the BBCAv1 (see Fig. 10DE). In future work, we will therefore test our new approach to estimate cell, neuron and glia densities in the mouse brain (step 2 of the pipeline - see Fig. 1) based on literature and the fitting. To this end, we will extend the literature review from Keller et al. [9]. The optimization process from step 4 will have to include these new constraints and variables to solve them all at once.

4.8. Changes in counts of inhibitory neuron in the brain

We obtained a new distribution of inhibitory neurons in the mouse brain, including a new distribution of PV+, SST+, and VIP+ neurons. The ratios of our inhibitory neuron subtypes counts divided by the neuron counts of the BBCAv2 fall near the numbers reported by Kim et al.: 4.03% PV+, 3.29% SST+, 0.63% VIP+ neurons [8]. The differences can be explained by the integration of other sources of literature to estimate these neuron population densities in the brain.

Our new method also accounts for the Rest population. Inhibitory neuron counts in BBCAv2 are therefore higher compared to BBCAv1. The three regions with the most pronounced updates of inhibitory neuron densities are: isocortex, olfactory areas, and cerebellum (see Table 2 and S2 Excel). In the cerebellum, only the granular layer was updated, since the molecular layer was already marked as purely inhibitory in both versions of the BBCA. As we observe in Fig. 10AB, this BBCAv2 model seems to display too many inhibitory neurons in the granular layer when compared to the GAD67 ISH experiment. Some interneurons of the cerebellum might co-express PV and SST [42]. As we discussed in Section 4.4, we can implement this specificity in our assumptions. In the olfactory areas, we have collected density estimates from literature [63,64]. These estimates compensate for the poor coverage of the olfactory bulb by the GAD67 marker experiment (see Fig. 3D) and result in larger inhibitory cell density estimates in BBCAv2. In the isocortex, the BBCAv2 estimates an overall higher proportion of inhibitory neurons. In some regions, such as the barrel field or the auditory areas (see Fig. 8B), the ratio of inhibitory neurons in the isocortex seems to be close to the 20% reported in Markram et al. [11] and Meyer et al. [58]. In other regions, such as the motor areas, the ratio of inhibitory neurons averages above 30%. For most of the isocortex regions, we gathered at least one density estimate for PV, SST and VIP expressing neurons from the literature; the ratio of inhibitory neurons in these regions is therefore minimized by the sum of these literature values divided by the counts of neurons from the BBCAv2. We studied the distribution of these ratios in the isocortex and found its average value to be 17.5% and its maximum to be 34.4%. The distribution of the ratios of GAD67 counts from literature divided by the counts of neurons from the BBCAv2 spans from 16.9% to 34.4% with a mean value at 28.5% (see S4 Fig.). This indicates either that the expected 20% ratio of inhibitory neurons is incorrect in some regions of the isocortex or that the distribution of neurons from the BBCAv2 obtained at step 2 is incorrect (see Fig. 1).

Conclusion

The updated Blue Brain Mouse Cell Atlas pipeline estimates the numbers and densities of GAD67+, PV+, SST+ and VIP+ neurons in each region of the mouse brain according to a methodology employing assumptions and literature values. It also captures a set of brain regions in which our assumptions cause contradictions between estimates (from literature or the transfer functions) that we have to correct with an optimization algorithm. Our future improvements will therefore target to reduce the errors that our analyses detected at the end of the pipeline. Future work will involve: integrating more literature points as they become available, improving the input data from the AIBS, refining our assumptions locally, and finally considering new methods for the fitting. Further analyses can study ratios and correlations of the different cell types of the brain. The workflow can be applied to any pair of reference volumes and to any genetic marker dataset from the AIBS website, which means that this pipeline can be extended to other cell types, if enough cell density estimates from literature are available to allow a fitting of mean expression data.

This project aims to involve the scientific community to contribute with open access to data, software, and tools. Our results will be released at <http://bbp.epfl.ch/nexus/cell-atlas> (Blue Brain Cell Atlas website interface) and the code accessible at <https://bbpgitlab.epfl.ch/nse/atlas-building-tools>. The BBCAv2 end product is expected to be of use for many purposes. For instance, in a companion paper [30], we use our inhibitory neuron density estimations in the isocortex to estimate the densities of morpho-electrical neuron subtypes. Our model will allow experimentalists to understand regional composition and permit computational neuroscientists to place defined cell types in their simulations. Furthermore, it sets the stage for further subdividing of inhibitory interneurons into more fine-grained subclasses and allows the neuroscience community to identify areas where current knowledge can be enhanced by additional constraints.

Acknowledgements

We would like to thank the Allen Institute for Brain Science for the large array of publicly available data.

Nissl CCFv2:

http://download.alleninstitute.org/informatics-archive/current-release/mouse_ccf/ara_nissl/

Annotation atlas CCFv2:

http://download.alleninstitute.org/informatics-archive/current-release/mouse_ccf/annotation/ccf_2015/

Annotation atlas CCFv3:

http://download.alleninstitute.org/informatics-archive/current-release/mouse_ccf/annotation/ccf_2017/

The AIBS ISH dataset can be found at <http://mouse.brain-map.org/search/index>. The corresponding experiment identifiers are listed below:

- Parvalbumin: Experiment id #868
- Somatostatin: #1001
- Vasoactive Intestinal Peptide: #77371835
- GAD1: #479 (equivalent of GAD67 see Fig. 3) and #79556706 (used in Section 3.4)

Conflicts of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Funding

This study was supported by funding to the Blue Brain Project, a research center of the École polytechnique fédérale de Lausanne (EPFL), from the Swiss government's ETH Board of the Swiss Federal Institutes of Technology.

References

1. Jorgenson LA, Newsome WT, Anderson DJ, Bargmann CI, Brown EN, Deisseroth K, et al. The BRAIN Initiative: developing technology to catalyse neuroscience discovery. *Philos Trans R Soc B Biol Sci.* 2015;370: 20140164. doi:10.1098/rstb.2014.0164
2. Muñoz-Castañeda R, Zingg B, Matho KS, Chen X, Wang Q, Foster NN, et al. Cellular anatomy of the mouse primary motor cortex. *Nature.* 2021;598: 159–166. doi:10.1038/s41586-021-03970-w
3. BRAIN Initiative Cell Census Network (BICCN), BRAIN Initiative Cell Census Network (BICCN) Corresponding authors, Callaway EM, Dong H-W, Ecker JR, Hawrylycz MJ, et al. A multimodal cell census and atlas of the mammalian primary motor cortex. *Nature.* 2021;598: 86–102. doi:10.1038/s41586-021-03950-0
4. National Institutes of Health. The BRAIN Initiative ®. 2021. Available: <https://braininitiative.nih.gov/>
5. Ramaswamy S, Markram H. Anatomy and physiology of the thick-tufted layer 5 pyramidal neuron. *Front Cell Neurosci.* 2015;9: 233. doi:10.3389/fncel.2015.00233
6. Casali S, Marenzi E, Medini C, Casellato C, D'Angelo E. Reconstruction and Simulation of a Scaffold Model of the Cerebellar Network. *Front Neuroinformatics.* 2019;13: 37. doi:10.3389/fninf.2019.00037
7. Todtenkopf MS, Vincent SL, Benes FM. A cross-study meta-analysis and three-dimensional comparison of cell counting in the anterior cingulate cortex of schizophrenic and bipolar brain. *Schizophr Res.* 2005;73: 79–89. doi:10.1016/j.schres.2004.08.018
8. Kim Y, Yang GR, Pradhan K, Venkataraju KU, Bota M, García del Molino LC, et al. Brain-wide Maps Reveal Stereotyped Cell-Type-Based Cortical Architecture and Subcortical Sexual Dimorphism. *Cell.* 2017;171: 456-469.e22. doi:10.1016/j.cell.2017.09.020
9. Keller D, Erö C, Markram H. Cell Densities in the Mouse Brain: A Systematic Review. *Front Neuroanat.* 2018;12: 83. doi:10.3389/fnana.2018.00083
10. Lang S, Dercksen VJ, Sakmann B, Oberlaender M. Simulation of signal flow in 3D reconstructions of an anatomically realistic neural network in rat vibrissa cortex. *Neural Netw.* 2011;24: 998–1011. doi:10.1016/j.neunet.2011.06.013
11. Markram H, Muller E, Ramaswamy S, Reimann MW, Abdellah M, Sanchez CA, et al. Reconstruction and Simulation of Neocortical Microcircuitry. *Cell.* 2015;163: 456–492. doi:10.1016/j.cell.2015.09.029
12. Lindahl M, Hellgren Kotaleski J. Untangling Basal Ganglia Network Dynamics and Function: Role of Dopamine Depletion and Inhibition Investigated in a Spiking Network Model. *eneuro.* 2016;3: ENEURO.0156-16.2016. doi:10.1523/ENEURO.0156-16.2016
13. Billeh YN, Cai B, Gratiy SL, Dai K, Iyer R, Gouwens NW, et al. Systematic Integration of Structural and Functional Data into Multi-scale Models of Mouse Primary Visual Cortex. *Neuron.* 2020;106: 388-403.e18. doi:10.1016/j.neuron.2020.01.040

- 1074 14. Erö C, Gewaltig M-O, Keller D, Markram H. A Cell Atlas for the Mouse Brain. *Front*
1075 *Neuroinformatics*. 2018;12: 84. doi:10.3389/fninf.2018.00084
- 1076 15. Dong HW. Allen reference atlas: a digital color brain atlas of the C57Black/6J male
1077 mouse. Hoboken, N.J.: Wiley; 2007.
- 1078 16. Lein ES, Hawrylycz MJ, Ao N, Ayres M, Bensinger A, Bernard A, et al. Genome-wide
1079 atlas of gene expression in the adult mouse brain. *Nature*. 2007;445: 168–176.
1080 doi:10.1038/nature05453
- 1081 17. Herculano-Houzel S, Ribeiro P, Campos L, Valotta da Silva A, Torres LB, Catania KC,
1082 et al. Updated Neuronal Scaling Rules for the Brains of Glires (Rodents/Lagomorphs).
1083 *Brain Behav Evol*. 2011;78: 302–314. doi:10.1159/000330825
- 1084 18. Herculano-Houzel S, Watson C, Paxinos G. Distribution of neurons in functional areas
1085 of the mouse cerebral cortex reveals quantitatively different cortical zones. *Front*
1086 *Neuroanat*. 2013;7: 35. doi:10.3389/fnana.2013.00035
- 1087 19. Hu H, Gan J, Jonas P. Fast-spiking, parvalbumin+ GABAergic interneurons: From
1088 cellular design to microcircuit function. *Science*. 2014;345: 1255263–1255263.
1089 doi:10.1126/science.1255263
- 1090 20. Wall NR, De La Parra M, Sorokin JM, Taniguchi H, Huang ZJ, Callaway EM. Brain-Wide
1091 Maps of Synaptic Input to Cortical Interneurons. *J Neurosci*. 2016;36: 4000–4009.
1092 doi:10.1523/JNEUROSCI.3967-15.2016
- 1093 21. Schmalbach B, Lepsveridze E, Djogo N, Papashvili G, Kuang F, Leshchyns'ka I, et al.
1094 Age-dependent loss of parvalbumin-expressing hippocampal interneurons in mice
1095 deficient in CHL1, a mental retardation and schizophrenia susceptibility gene. *J*
1096 *Neurochem*. 2015;135: 830–844. doi:10.1111/jnc.13284
- 1097 22. Fasulo L, Brandi R, Arisi I, La Regina F, Berretta N, Capsoni S, et al. ProNGF Drives
1098 Localized and Cell Selective Parvalbumin Interneuron and Perineuronal Net Depletion
1099 in the Dentate Gyrus of Transgenic Mice. *Front Mol Neurosci*. 2017;10: 20.
1100 doi:10.3389/fnmol.2017.00020
- 1101 23. Lauber E, Filice F, Schwaller B. Dysregulation of Parvalbumin Expression in the
1102 *Cntnap2*^{-/-} Mouse Model of Autism Spectrum Disorder. *Front Mol Neurosci*. 2018;11:
1103 262. doi:10.3389/fnmol.2018.00262
- 1104 24. Pirone A, Alexander JM, Koenig JB, Cook-Snyder DR, Palnati M, Wickham RJ, et al.
1105 Social Stimulus Causes Aberrant Activation of the Medial Prefrontal Cortex in a Mouse
1106 Model With Autism-Like Behaviors. *Front Synaptic Neurosci*. 2018;10: 35.
1107 doi:10.3389/fnsyn.2018.00035
- 1108 25. Ramos B, Baglietto-Vargas D, Rio JC del, Moreno-Gonzalez I, Santa-Maria C, Jimenez
1109 S, et al. Early neuropathology of somatostatin/NPY GABAergic cells in the
1110 hippocampus of a PS1×APP transgenic model of Alzheimer's disease. *Neurobiol Aging*.
1111 2006;27: 1658–1672. doi:10.1016/j.neurobiolaging.2005.09.022
- 1112 26. Trujillo-Estrada L, Dávila JC, Sánchez-Mejias E, Sánchez-Varo R, Gomez-Arboledas A,
1113 Vizuite M, et al. Early Neuronal Loss and Axonal/Presynaptic Damage is Associated
1114 with Accelerated Amyloid- β Accumulation in A β PP/PS1 Alzheimer's Disease Mice
1115 Subiculum. *J Alzheimers Dis*. 2014;42: 521–541. doi:10.3233/JAD-140495

- 1116 27. Sanchez-Mejias E, Nuñez-Diaz C, Sanchez-Varo R, Gomez-Arboledas A, Garcia-Leon
1117 JA, Fernandez-Valenzuela JJ, et al. Distinct disease-sensitive GABAergic neurons in
1118 the perirhinal cortex of Alzheimer's mice and patients. *Brain Pathol.* 2020;30: 345–363.
1119 doi:10.1111/bpa.12785
- 1120 28. Prönneke A, Scheuer B, Wagener RJ, Möck M, Witte M, Staiger JF. Characterizing VIP
1121 Neurons in the Barrel Cortex of VIPcre/tomato Mice Reveals Layer-Specific
1122 Differences. *Cereb Cortex.* 2015;25: 4854–4868. doi:10.1093/cercor/bhv202
- 1123 29. Passemard S, Rabat CA, Gressens P, Lelièvre V. Chapter 128 - VIP. 2nd ed.
1124 Handbook of Biologically Active Peptides. 2nd ed. Cambridge, Massachusetts: Elsevier;
1125 2013. pp. 966–974. doi:10.1016/B978-0-12-385095-9.00128-7
- 1126 30. Roussel Y, Verasztó C, Rodarie D, Damart T, Reimann M, Ramaswamy S, et al.
1127 Mapping of morpho-electric features to molecular identity of cortical inhibitory neurons.
1128 *Neuroscience*; 2021 Nov. doi:10.1101/2021.11.24.469815
- 1129 31. Krepl J, Casalegno F, Delattre E, Erö C, Lu H, Keller D, et al. Supervised Learning With
1130 Perceptual Similarity for Multimodal Gene Expression Registration of a Mouse Brain
1131 Atlas. *Front Neuroinformatics.* 2021;15: 37. doi:10.3389/fninf.2021.691918
- 1132 32. Huang ZJ, Paul A. The diversity of GABAergic neurons and neural communication
1133 elements. *Nat Rev Neurosci.* 2019;20: 563–572. doi:10.1038/s41583-019-0195-4
- 1134 33. Allen Institute for Brain Science. CCFv3 Adult Mouse Annotation Atlas. 2017. Available:
1135 [http://download.alleninstitute.org/informatics-archive/current-](http://download.alleninstitute.org/informatics-archive/current-release/mouse_ccf/annotation/ccf_2017/)
1136 [release/mouse_ccf/annotation/ccf_2017/](http://download.alleninstitute.org/informatics-archive/current-release/mouse_ccf/annotation/ccf_2017/)
- 1137 34. Oh SW, Harris JA, Ng L, Winslow B, Cain N, Mihalas S, et al. A mesoscale connectome
1138 of the mouse brain. *Nature.* 2014;508: 207–214. doi:10.1038/nature13186
- 1139 35. Kuan L, Li Y, Lau C, Feng D, Bernard A, Sunkin SM, et al. Neuroinformatics of the Allen
1140 Mouse Brain Connectivity Atlas. *Methods.* 2015;73: 4–17.
1141 doi:10.1016/j.ymeth.2014.12.013
- 1142 36. Wang Q, Ding S-L, Li Y, Royall J, Feng D, Lesnar P, et al. The Allen Mouse Brain
1143 Common Coordinate Framework: A 3D Reference Atlas. *Cell.* 2020;181: 936-953.e20.
1144 doi:10.1016/j.cell.2020.04.007
- 1145 37. Allen Institute for Brain Science. CCFv2 Adult Mouse Annotation Atlas. 2015. Available:
1146 [http://download.alleninstitute.org/informatics-archive/current-](http://download.alleninstitute.org/informatics-archive/current-release/mouse_ccf/annotation/ccf_2015/)
1147 [release/mouse_ccf/annotation/ccf_2015/](http://download.alleninstitute.org/informatics-archive/current-release/mouse_ccf/annotation/ccf_2015/)
- 1148 38. Allen Institute for Brain Science. Adult Mouse Allen Reference Atlas. 2015. Available:
1149 [http://download.alleninstitute.org/informatics-archive/current-](http://download.alleninstitute.org/informatics-archive/current-release/mouse_ccf/ara_nissl/)
1150 [release/mouse_ccf/ara_nissl/](http://download.alleninstitute.org/informatics-archive/current-release/mouse_ccf/ara_nissl/)
- 1151 39. Gourfinkel-An I, Parain K, Hartmann A, Mangiarini L, Brice A, Bates G, et al. Changes
1152 in GAD67 mRNA expression evidenced by in situ hybridization in the brain of R6/2
1153 transgenic mice. *J Neurochem.* 2003;86: 1369–1378. doi:https://doi.org/10.1046/j.1471-
1154 4159.2003.01916.x
- 1155 40. Lefort S, Tómm C, Floyd Sarria J-C, Petersen CCH. The Excitatory Neuronal Network
1156 of the C2 Barrel Column in Mouse Primary Somatosensory Cortex. *Neuron.* 2009;61:
1157 301–316. doi:10.1016/j.neuron.2008.12.020

- 1158 41. Xu X, Roby KD, Callaway EM. Immunochemical characterization of inhibitory mouse
1159 cortical neurons: Three chemically distinct classes of inhibitory cells. *J Comp Neurol*.
1160 2010;518: 389–404. doi:10.1002/cne.22229
- 1161 42. Zeisel A, Hochgerner H, Lönnerberg P, Johnsson A, Memic F, van der Zwan J, et al.
1162 Molecular Architecture of the Mouse Nervous System. *Cell*. 2018;174: 999-1014.e22.
1163 doi:10.1016/j.cell.2018.06.021
- 1164 43. Asada H, Kawamura Y, Maruyama K, Kume H, Ding R-G, Kanbara N, et al. Cleft palate
1165 and decreased brain -aminobutyric acid in mice lacking the 67-kDa isoform of glutamic
1166 acid decarboxylase. *Proc Natl Acad Sci*. 1997;94: 6496–6499.
1167 doi:10.1073/pnas.94.12.6496
- 1168 44. Kash SF, Johnson RS, Tecott LH, Noebels JL, Mayfield RD, Hanahan D, et al. Epilepsy
1169 in mice deficient in the 65-kDa isoform of glutamic acid decarboxylase. *Proc Natl Acad*
1170 *Sci*. 1997;94: 14060–14065. doi:10.1073/pnas.94.25.14060
- 1171 45. Celio MR, Heizmann CW. Calcium-binding protein parvalbumin as a neuronal marker.
1172 *Nature*. 1981;293: 300–302. doi:10.1038/293300a0
- 1173 46. Emson PC. Somatostatin and Receptors. In: Squire LR, editor. *Encyclopedia of*
1174 *Neuroscience*. Oxford: Academic Press; 2009. pp. 121–127. doi:10.1016/B978-
1175 008045046-9.01473-X
- 1176 47. Celio M. Parvalbumin in most gamma-aminobutyric acid-containing neurons of the rat
1177 cerebral cortex. *Science*. 1986;231: 995–997. doi:10.1126/science.3945815
- 1178 48. Tasic B, Yao Z, Graybuck LT, Smith KA, Nguyen TN, Bertagnolli D, et al. Shared and
1179 distinct transcriptomic cell types across neocortical areas. *Nature*. 2018;563: 72–78.
1180 doi:10.1038/s41586-018-0654-5
- 1181 49. Irintchev A, Rollenhagen A, Troncoso E, Kiss JZ, Schachner M. Structural and
1182 Functional Aberrations in the Cerebral Cortex of Tenascin-C Deficient Mice. *Cereb*
1183 *Cortex*. 2005;15: 950–962. doi:10.1093/cercor/bhh195
- 1184 50. Ransome MJ, Turnley AM. Analysis of neuronal subpopulations in mice over-expressing
1185 suppressor of cytokine signaling-2. *Neuroscience*. 2005;132: 673–687.
1186 doi:10.1016/j.neuroscience.2004.12.041
- 1187 51. Neddens J, Buonanno A. Selective populations of hippocampal interneurons express
1188 ErbB4 and their number and distribution is altered in ErbB4 knockout mice.
1189 *Hippocampus*. 2010;20: 724–744. doi:https://doi.org/10.1002/hipo.20675
- 1190 52. Schmid JS, Bernreuther C, Nikonenko AG, Ling Z, Mies G, Hossmann K-A, et al.
1191 Heterozygosity for the mutated X-chromosome-linked L1 cell adhesion molecule gene
1192 leads to increased numbers of neurons and enhanced metabolism in the forebrain of
1193 female carrier mice. *Brain Struct Funct*. 2013;218: 1375–1390. doi:10.1007/s00429-
1194 012-0463-9
- 1195 53. Han L-C, Zhang H, Wang W, Wei Y-Y, Sun X-X, Yanagawa Y, et al. The Effect of
1196 Sevoflurane Inhalation on GABAergic Neurons Activation: Observation on the GAD67-
1197 GFP Knock-In Mouse. *Anat Rec Adv Integr Anat Evol Biol*. 2010;293: 2114–2122.
1198 doi:10.1002/ar.21113

- 1199 54. Jinno S, Aika Y, Fukuda T, Kosaka T. Quantitative analysis of GABAergic neurons in
1200 the mouse hippocampus, with optical disector using confocal laser scanning
1201 microscope. *Brain Res.* 1998;814: 55–70. doi:10.1016/S0006-8993(98)01075-0
- 1202 55. Otsu N. A Threshold Selection Method from Gray-Level Histograms. *IEEE Trans Syst*
1203 *Man Cybern.* 1979;9: 62–66. doi:10.1109/TSMC.1979.4310076
- 1204 56. Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al.
1205 SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat Methods.*
1206 2020;17: 261–272. doi:10.1038/s41592-019-0686-2
- 1207 57. Gerfen CR, Wilson CJ. Chapter II The basal ganglia. *Handbook of Chemical*
1208 *Neuroanatomy.* Elsevier; 1996. pp. 371–468. doi:10.1016/S0924-8196(96)80004-2
- 1209 58. Meyer HS, Schwarz D, Wimmer VC, Schmitt AC, Kerr JND, Sakmann B, et al. Inhibitory
1210 interneurons in a cortical column form hot zones of inhibition in layers 2 and 5A. *Proc*
1211 *Natl Acad Sci.* 2011;108: 16807–16812. doi:10.1073/pnas.1113648108
- 1212 59. Trifonov S, Yamashita Y, Kase M, Maruyama M, Sugimoto T. Glutamic acid
1213 decarboxylase 1 alternative splicing isoforms: characterization, expression and
1214 quantification in the mouse brain. *BMC Neurosci.* 2014;15: 114. doi:10.1186/1471-
1215 2202-15-114
- 1216 60. Wang X, Allen WE, Wright MA, Sylwestrak EL, Samusik N, Vesuna S, et al. Three-
1217 dimensional intact-tissue sequencing of single-cell transcriptional states. *Science.*
1218 2018;361: eaat5691. doi:10.1126/science.aat5691
- 1219 61. Szabolcsi V, Celio MR. De novo expression of parvalbumin in ependymal cells in
1220 response to brain injury promotes ependymal remodeling and wound repair. *Glia.*
1221 2015;63: 567–594. doi:https://doi.org/10.1002/glia.22768
- 1222 62. Lee S, Hjerling-Leffler J, Zagha E, Fishell G, Rudy B. The Largest Group of Superficial
1223 Neocortical GABAergic Interneurons Expresses Ionotropic Serotonin Receptors. *J*
1224 *Neurosci.* 2010;30: 5926–5934. doi:10.1523/JNEUROSCI.1869-10.2010
- 1225 63. Parrish-Aungst S, Shipley MT, Erdelyi F, Szabo G, Puche AC. Quantitative analysis of
1226 neuronal diversity in the mouse olfactory bulb. *J Comp Neurol.* 2007;501: 825–836.
1227 doi:10.1002/cne.21205
- 1228 64. Suzuki N, Bekkers JM. Inhibitory neurons in the anterior piriform cortex of the mouse:
1229 Classification using molecular markers. *J Comp Neurol.* 2010;518: 1670–1687.
1230 doi:https://doi.org/10.1002/cne.22295

1231

Supporting information

S1 Fig. Global impact of the annotation atlas on cells and neurons densities

(A), (B), (C) Distribution of cell and neuron density values reported in literature against generated densities for similar regions, using the 3 pairs of annotations and Nissl volumes (from left to right: CCFbbp, CCFv2 from the AIBS, CCFv3 from the AIBS). When multiple literature sources are available for the exact same region, they are both shown as a data point and are linked together. The color encodes the brain regions according to the AV, while the shapes of the points encode for cell types. The middle line delimits equal quantities, while the dashed line shows the average deviation of 2.7-fold between literature values reporting on the same region. Some subregions of the brain are not represented in CCFv3 which explains the different numbers of points.

(D), (E), (F) Histogram of the brain regions in terms of neuron density values for the 3 pairs of AV and Nissl volumes (from left to right: CCFbbp, CCFv2 from the AIBS, CCFv3 from the AIBS). Each region is represented with a single-color patch of the same size.

S2 Fig. Algorithm 1: Cap inhibitory densities to number of neurons.

The estimated counts of GAD67+ neurons and the sum of PV+, SST+, VIP+ neurons counts are limited by the previously-computed neuron counts (step 2 of the BBCAv2 pipeline - see Fig. 1) to ensure that assumptions 2 and 3 are fulfilled (see Section 2.4). Recall that $\mathbf{R}$ is the set of brain regions, $\forall r \in \mathbf{R}$ inversely ordered according to their depth in the region hierarchy of the AV ($\mathbf{R}_o$), the algorithm checks if the conditions $n\text{Neu}_r \geq n\text{GAD}_r$ and $n\text{Neu}_r \geq n\text{PV}_r + n\text{SST}_r + n\text{VIP}_r$ are satisfied. If not, it finds a solution which tries to match the following properties, ordered by priority: (1): Remain in range of the standard deviation of each value (confidence intervals), (2): Maintain the proportion of PV, SST and VIP within the region. If a solution exists within the confidence intervals, then the number of extra neurons (diff variable at line 19) is subtracted proportionally to the ratio of each neuron type (computed at lines 16-18). Then, if one of the neuron type estimates reaches the minimum of its confidence interval, then it is no more reduced (lines 20-22) and the remaining extra neurons are subtracted from the other neuron type estimates (lines 25, 28, 31).

S3 Fig. Algorithm 2: Maintain inhibitory densities coherence

This algorithm corrects the estimated densities of PV+, SST+, VIP+ and GAD67+ neurons in the model so that assumption 4 of Section 2.4 is fulfilled. $\mathbf{R}$ is the set of brain regions, $\forall r \in \mathbf{R}$ inversely ordered according to their depth in the region hierarchy of the AV ($\mathbf{R}_o$), the algorithm checks if the condition $n\text{GAD}_r \geq n\text{PV}_r + n\text{SST}_r + n\text{VIP}_r$ is satisfied. If not, it finds a solution which tries to match the following properties, ordered by priority: (1): Remain in range of the standard deviation of each value (confidence intervals), (2): Maintain the proportion (ratios) of PV, SST and VIP. When a set of value exists so that the property (1) is fulfilled, there is a correction factor $q \in [0,1]$ which

corresponds to the fraction of standard deviation needed to guarantee that the sum of the inhibitory subtypes remains under the estimated count of inhibitory neurons and that each value remains within its confidence interval.

S4 Fig. Explaining the isocortex inhibitory densities from BBCAv2.

(A) Coronal view of the Nissl reference atlas used for BBCAv2, showing cells of the mouse brain. Regions with high cell density appear in dark grey. The blue, orange and green lines behind the sagittal slice highlight the rapid changes of Nissl expression coming from the original Nissl experiment from Dong [15].
(B) (C) Evolution of the median expression level along the sagittal axis, in the Nissl volume realigned in Erö et al. [14], for the whole brain (B) and the isocortex (C). The rapid changes of Nissl expression that were detected in (A) are also visible in (B) and (C).
(D) isocortex ratio of inhibitory neurons according to literature. The distributions show to ratios literature cell type counts in isocortex divided by the counts of neurons of the BBCAv2 in their corresponding region. The bottom and top distributions correspond to the proportion of respectively the sum of the reported values of PV, SST and VIP counts, and the reported values of GAD67. The mean value of each distribution is shown in orange. The minimum and maximum values are indicated by the whiskers

S5 Fig. Orientations field and depth computation of the barrel cortex

(A) Coronal slice of the AV showing the barrel cortex and its different sublayers.
(B) Coronal slice of the scalar field of the barrel cortex. A weight is assigned to every voxel of the AV. These weights follow the order of crossing the isocortex by its fibers from the corpus callosum to layer 1. The borders of the barrel field are highlighted in orange. The weight assigned to the surrounding voxels of the region corresponds to their closest layer's. The weight of the voxels outside the region beyond layer 1 increases as moving away from the barrel cortex.
(C) Coronal slice of the orientation field of the barrel cortex. To each voxel of the AV, a 3D direction normalized vector is computed corresponding to the main axis of the axons in the region. Colors represent the orientation vectors norm on their respective plane, black lines their projected axis.
(D) Coronal slice of the depth according to pia in the barrel cortex expressed in micrometers.

S6 Fig. Cortical inhibitory density distribution

(A) Normalized distribution of cortical inhibitory neuron densities across the isocortex subregions grouped by layers. For each layer, the fitted normal distribution associated is displayed as a line on top of it. The normal distributions of L2/3 (mean 1.6×10^4 , std. 3.8×10^3), L4 (mean 1.9×10^4 , std. 4.1×10^3) and L5 (mean 1.8×10^4 , std. 3.1×10^3) are overlapping significantly.

(B) Distribution of inhibitory neuron densities according to cortical layers for each subregion of the isocortex. For most of the cortical subregions, the density is constant from L2 to L5 and drops for L6.

S1 Excel. Literature review of PV+, SST+, VIP+, and GAD67+ neuron densities in the mouse brain

This file lists papers reporting quantification of neurons reacting to GAD67 (first sheet) and PV, SST, VIP (second sheet). For each paper and for each matching brain region of the AV, a mean density of neurons and its corresponding standard deviation is extracted. Assumptions on annotations and preprocessing steps are described in the comments' column. The age of the mice used by the papers is also indicated. The data extracted from the BBCAv2 model, used to convert reported counts or proportions into densities are stored into the last column. The last sheet contains the list of regions known containing only excitatory or inhibitory neurons and their deduced corresponding inhibitory densities of neurons according to the BBCAv2.

S2 Excel. Results of the BBCAv2 pipeline and additional analysis

The first sheet of this file contains the final densities and standard deviation of each neuron type and for each region of the AV based on the BBCAv2 pipeline. The final column stores the volume of each region of the brain.

The second sheet contains the result of the fitting of the transfer functions from region mean expression to cell density for each genetic marker (see Section 2.6). Two different ISH experiments from the AIBS were used for the GAD67 (see Section 2.4 and 3.4). The results of the fitting for each experiment and for the combined dataset are stored in different columns.

The final sheet contains the densities of each neuron types for each region of the AV obtained with the BBCAv1 pipeline. The process to obtain these values is described in Section 2.1.1 and 3.4.