The aphid BCR4 structure and activity uncover a new defensin peptide superfamily

Karine Loth^{1,2#}, Nicolas Parisot^{3#}, Françoise Paquet¹, Catherine Sivignon⁴, Isabelle Rahioui⁴,
Mélanie Ribeiro Lopes³, Karen Gaget⁴, Gabrielle Duport⁴, Agnès F. Delmas¹, Vincent
Aucagne¹, Abdelaziz Heddi³, Federica Calevro⁴ and Pedro da Silva^{3*}

¹: Centre de Biophysique Moléculaire, CNRS UPR 4301, Orléans, France.

²: UFR Sciences et Techniques, Université d'Orléans, Orléans, F-45071, France

³: Univ Lyon, INSA Lyon, INRAE, BF2I, UMR 203, F-69621, Villeurbanne, France

⁴: Univ Lyon, INRAE, INSA Lyon, BF2I, UMR 203, F-69621, Villeurbanne, France

These authors contributed equally

*To whom correspondence should be addressed: Pedro Da Silva,

e-mail: pedro.da-silva@insa-lyon.fr

ABSTRACT

Aphids (Hemiptera: Aphidoidea) are among the most injuring insects for agricultural plants and their management is a great challenge in agronomical research. A new class of proteins, called Bacteriocyte-specific Cysteine-Rich (BCR), provides an alternative to chemical insecticides for pest control. BCRs have been initially identified in the pea aphid *Acyrtosiphon pisum*. They are small disulfide bond-rich proteins expressed exclusively in aphid bacteriocytes, the insect derived cells that host intracellular symbiotic bacteria. Here, we show that one out of the *A. pisum* BCRs, BCR4, displays an outstanding insecticidal activity against the pea aphid, impairing insect survival and nymphal growth, providing evidence for its potential use as a new biopesticides. Our comparative genomics and phylogenetic analysis indicate that BCRs seem restricted to the aphid lineage. The 3D structure of the BCR4 reveals that this peptide belongs to a yet unknown structural class of peptides and defines a new superfamily of defensins.

INTRODUCTION

Insects are among the most important pests of cultured plants and stored products, causing an estimated yearly loss of hundreds of millions of dollars worldwide (Oerke & Dehne, 2004; Bradshaw et al., 2016; Diagne et al., 2020). Aphids, in particular, hold a prominent place among insect pests as they represent up to 26% of the pests found on the main food crops grown in temperate climate (maize, wheat, potatoes, sugar beet, barley and tomatoes) (Dedryver et al., 2010; Calevro et al., 2019). Aphid control strategies rely almost exclusively on chemical treatments, which cause persistent environmental pollution and lead to the emergence of insect resistance (Goulson, 2018). New ecologically-friendly solutions are therefore required to control aphids and other phloem-feeders.

Small Disulfide-Rich Proteins (DRP) extracted from plants or arthropods are promising alternative biopesticide molecules (Gressent et al., 2011; Huang et al., 2019; King, 2019). Those naturally occurring molecules display a very broad range of biological activities, mainly related to host-defense processes (Shafee et al., 2017), and show a large structural and chemical diversity. Their exceptional stability is particularly appealing for drug discovery purposes (Gonzalez-Castro et al., 2021). In addition to rather rigid three-dimensional conformations imposed by their polycyclic architecture, DRP exhibit a strong resistance toward *in vivo* enzymatic degradation. Altogether, these features have contributed in establishing DRP as emerging lead compounds for the development of novel peptide-based drugs and, more recently, as potential biopesticides in agronomical research (Rahoui et al., 2014; Huang et al., 2019; King, 2019; Bell et al., 2021). For instance, a knottin DRP extracted from pea seeds, PA1b (Pea Albumin 1, subunit b, 37 amino acids, three disulfide bonds) (Delobel et al., 1998; Gressent et al., 2011; Rahoui et al., 2014), is toxic to numerous insects, including aphids, cereal weevils, mosquitos and moths (Rahoui et al., 2014).

A new class of DRP, called Bacteriocyte-specific Cysteine Rich (BCR) peptides, has recently been identified in a major crop pest, the pea aphid, *Acyrtosiphon pisum* (Shigenobu & Stern, 2013). As many other crop pest insects that thrive on unbalanced diets, aphids have evolved long-lasting relationships with endosymbiotic bacteria and almost all aphids are found in association with the γ -proteobacterium *Buchnera aphidicola* (Shigenobu et al., 2000; Baumann, 2005; Douglas, 2015). This bacterium supplements the host diet with nutrients lacking or being limited in their habitats, therefore allowing insects to proliferate and cause major economic, social, and health damage. Neither the host nor the endosymbionts can survive independently from each other. *B. aphidicola* is non-culturable and insects artificially deprived of their endosymbionts (aposymbiotic) cannot survive nor reproduce (Brinza et al., 2009). The maintenance of this association relies on the compartmentalization of endosymbionts in specialized insect cells, called bacteriocytes (Simonet et al., 2018). BCRs are encoded by seven orphan genes, and are all expressed exclusively in bacteriocytes of both embryonic and adult aphids (Shigenobu & Stern, 2013). This suggests that BCRs may play a role in bacteriocyte homeostasis, presumably in endosymbiont control as previously in the cereal-weevil endosymbiosis (Login et al. 2011). Consistent with this hypothesis, it has been shown that BCR1, BCR2, BCR3, BCR4, BCR5 and BCR8 exhibit antimicrobial activity or can permeabilize the membrane of *E. coli* cells (Uchi et al., 2019).

Each BCR peptide consists of a secretory signal peptide and a mature peptide, composed of 44-84 amino acids and containing from six in the case of BCR1 to 5 and BCR8, to eight, in the case of BCR6, cysteine residues (Shigenobu & Stern, 2013). Between BCR1, BCR2, BCR4 and BCR5, the cysteine-rich region is highly divergent, but the six cysteines have almost identical spacing in the predicted proteins. Three of these genes, BCR1, BCR4 and BCR5, are found within a genomic region of 20 kbp, suggesting that they may have arisen by a recent tandem gene duplication. No similarity could be found between any of the other BCR family genes

(Shigenobu & Stern, 2013). Intriguingly, the pea aphid BCRs show no significant sequence similarity with genes in species outside of the aphid lineage. DRPs have undergone extensive divergent evolution in their sequence structure and function. They are classified based on their secondary structure orientation, cysteine distribution across the sequence and cysteine bonding pattern tertiary structure similarities and precursor gene sequence (Shafee et al., 2016; Shafee et al., 2017). Based on their sequence analysis, it was assumed that they could have evolved from defensin-type antimicrobial peptides (AMPs), small proteins containing six to eight cysteines, which are universally found in both animal and plants (Shigenobu & Stern, 2013; Uchi et al., 2019).

In this study, we focused on the BCR4 of *A. pisum*, a benchmark example of peptide with antimicrobial activities (Uchi et al., 2019). We discovered that BCR4 displays an outstanding insecticidal activity against the pea aphid. Taking advantage of the recent sequencing of several aphid genomes, which enables the study of gene families' diversification through comparative and evolutionary analyses (Ribeiro Lopes et al., 2020; Huygens et al., 2021), we conducted a comparative genomic analysis across 22 aphid species for which sequence information was available. This allowed the identification of 76 new BCR sequences, all restricted to aphid species and not related to any known defensins. Finally, we determined BCR4 3D structure and showed that it belongs to a new structural class of disulfide-rich proteins. Overall, the biochemical analyses, the evolutionary history and the 3D structure of the BCR4 have highlighted significant insights into the biological and structural properties of BCRs, and have provided evidence for the use of this new defensin superfamily as a potential new biopesticides.

RESULTS

Total synthesis of BCR4

To investigate the biological activity of members of the BCR peptide family, a pure sample of synthetic BCR4 was produced through total chemical synthesis. As standard automated Fmoc/*t*Bu solid phase peptide synthesis (SPPS) was unsuccessful (SI appendix Figures S1 and S2, table S1), we turned to a native chemical ligation (NCL)-based approach (Dawson et al., 1994), relying on the assembly of two medium-sized peptide segments. Using a recently developed methodology (Lelievre et al., 2016; Terrier et al., 2016; Terrier et al., 2017; Abboud & Aucagne, 2020; Abboud, et al., 2021a; Abboud, et al., 2021b), we coupled a 20 amino acid (aa) *N*-2-hydroxy-5-nitrobenzylcysteine (*N*-Hnb-Cys) crypto-thioester with a 30 amino acid cysteinyl peptide and obtained the full length reduced form of BCR4 at a high purity (Figure 1). Oxidative folding under thermodynamic control was achieved using a standard protocol (Derache et al., 2012; Martinez et al., 2016), leading to one major compound featuring three disulfide bridges as evidenced by HRMS analysis (SI appendix Figures S3 to S11).

Figure 1. Chemical synthesis of BCR4. (A) Schematic representation of the BCR4 peptide chemical synthesis process. (B) RP-HPLC chromatogram and ESI-HRMS mass spectrum of the purified folded peptide.

Antimicrobial activities of BCR4

To assess the antimicrobial activity of BCR4, we tested the effect of various concentrations (ranging from 5 μ M to 80 μ M) of this peptide on the growth of the Gram-negative bacteria *Escherichia coli* (strain NM522) and the Gram-positive bacteria *Micrococcus luteus*. Consistent with previous observations (Uchi et al., 2019), a slight inhibition of *E.coli* growth was detected at 5 μ M of BCR4. This antimicrobial activity increased with BCR4 concentration and we determined a minimal inhibitory concentration (MIC) of 17.0 ± 2.4 μ M for which no bacterial growth was detected. Comparatively, no antimicrobial activity was detected against the Gram-positive bacteria *Micrococcus luteus* (Table 1).

Table 1. Antimicrobial activities of BCR4 peptide

	MIC ^a (μ M)
<i>Escherichia coli</i> NM522	17.0 ± 2.4
<i>Micrococcus luteus</i>	> 80

^a Minimal inhibitory concentration

Insect bioassays

The insecticidal potential of BCR4 was assayed by oral administration of various concentration of this peptide to the pea aphid and monitoring of its survival. For all BCR4 concentrations tested (5-80 μ M), aphid mortality rate was always significantly greater in the BCR4-aphids compared to the control group, demonstrating a specific effect of BCR4 ingestion on aphid survival (Figure 2). This effect was dependent on both BCR4 dosage and duration of treatment. The highest concentration tested (80 μ M) also had the strongest lethal effect with a Lethal Time 50% (LT50) of 1.16 days (Table 2) and no surviving aphids after two days of BCR4 treatments (Figure 2).

Table 2. Toxicity of BCR4 peptide on the pea aphid *Acyrtosiphon pisum* (means Lethal Time 50 (LT50) in days, with confidence intervals), analyzed by survival analysis with a log-normal fit.

	Concentration (μM)						
BCR4 peptide	80	40	35	27.5	20	10	5
LT50 (days)	1.16 [0.98 – 1.37]	1.76 [1.27 – 2.42]	1.94 [1.47 – 2.57]	3.24 [2.22 – 4.72]	3.48 [2.28 – 5.31]	11.3 [4.16 – 30.8]	> 20

Even for concentration as low as 20 μM , aphids, on average, did not survive more than 4 days and only one third reached the adulthood. Apart from this effect on aphid survival, the most striking phenotypical effect observed with BCR4 treatment was a statistically significant growth inhibition of the surviving aphids for doses higher than 5 μM , with, for instance, a 60% weight reduction at 7 days after ingestion of BCR4 at 27.5 μM (Figure 2B).

A

B

Figure 2. (A) Survival curves of the aphid *Acyrtosiphon pisum* reared on artificial diets containing different concentrations of BCR4 peptide. Means Lethal Time 50 (LT50), in days, are indicated above each curve. **(B)** Mass (mg) of 7-day old pea aphid *Acyrtosiphon pisum* subjected to BCR4 treatment. Concentrations labelled with different letters are significantly different ($P < 0.05$); sample size of BCR4 35 and 40 μM concentrations too small for statistics.

Phylogeny of the BCR family

To decipher the molecular evolution of the BCR peptides, we aimed at identifying the whole set of proteins homologous to the seven *A. pisum* BCR proteins. Through an exhaustive search of the NCBI nucleotide and protein databases, coupled with a specific inspection of the sequenced aphid species available in the AphidBase (Legeai et al., 2010) database, we retrieved a total of 76 new BCR sequences across 20 aphid species (Table S2) among the 22 for which sequences are available in those publicly available databases. We also found one additional sequence from *A. pisum*, shifting the total number of BCRs in this insect to eight. Interestingly, all these sequences belong to member of the Aphidoidea super-family, thus confirming BCR proteins are restricted to the aphid (*s.l.*) lineage (Shigenobu & Stern, 2013). The complete set of 83 BCR sequences was used for amino acid sequences alignment and phylogenetic tree reconstruction (Figure 3A). Based on these, the BCR sequences can be grouped into four

subfamilies including homologs of the BCR1-2-4-5, BCR3, BCR6 and BCR8 sequences of *A. pisum*, respectively. The majority of subfamilies has six cysteine residues with the exception of the BCR6 subfamily which has 8 cysteines. The spacing between cysteines within the BCR sequences is fully conserved within subfamilies, which ensures secure within-family homologies and phylogenetic placement (Table 3).

Nineteen aphid species have full genome sequences available (*Acyrtosiphon pisum*, *Aphis craccivora*, *Aphis glycines*, *Aphis gossypii*, *Aulacorthum solani*, *Cinara cedri*, *Diuraphis noxia*, *Eriosoma lanigerum*, *Macrosiphum rosae*, *Melanaphis sacchari*, *Myzus cerasi*, *Myzus persicae*, *Pentalonia nigronervosa*, *Rhopalosiphum maidis*, *Rhopalosiphum padi*, *Schizaphis graminum*, *Sipha flava*, *Sitobion avenae*, *Sitobion miscanthi*). This permits to predict the complete set of BCR peptides encoded by those aphid genomes. In this group, we observed a high variability in the number and distribution of BCR peptides among the four subfamilies (Table 3). No BCR were found in the *E. lanigerum* (Eriosomatinae subfamily) and *S. flava* (Chaitophorinae subfamily) genomes and only one in *C. cedri* 's (Lachninae subfamily) (Figure 3B). Comparatively, all members of the Aphidinae subfamily appear to possess at least two BCR genes and *M. rosae* has the largest repertoire of BCR sequences, with 13 distinct sequences distributed across the four BCR subfamilies. While members of the BCR3 and BCR8 subfamilies are present in all the Aphidinae species included in this study, members of the BCR1-2-4-5 and BCR6 subfamilies appear restricted to the Macrosiphini aphid tribe suggesting that the genes encoding those BCRs arose from duplication of the former.

Table 3. Characteristics of the four BCR subfamilies in 19 aphid species with fully sequenced genomes.

	Number of cysteine residues	Consensus motif in <i>A. pisum</i> ^a	Number of orthologous sequences in aphid species ^b
BCR1-2-4-5	6	C (3) C (3-5) C (4-6) C (1) C (11-13) C	Apis (5), Dnox (1), Mros (5), Smis (1)
BCR3	6	C (3) C (6) C (21-25) C (6) C (3) C	Acra (3), Agly (1), Agos (2), Apis (1), Asol (2), Cced (1), Dnox (2), Mcer (1), Mper (1), Mros (3), Msac (3), Pnig (2), Rmai (3), Rpad (1), Sgra (2), Smis (2),
BCR6	8	C (15) C (3) C (3) C (10) C (10) C (1) CC	Apis (1), Asol (1), Dnox (2), Mcer (1), Mper (1), Mros (3), Pnig (5), Save (1), Smis (2)
BCR8	6	C (3-4) C (10) C (15-17) C (1) CC	, Acra (3), Agly (1), Agos (1), Apis (1), Asol (1), Dnox (1), Mcer (1), Mper (1), Msac (1), Mros (2), Pnig (1), Rmai (1), Rpad (1), Smis (2)

^a The number of residues between cysteines are noted in parenthesis.

^b Abbreviations: Acra, *Aphis craccivora*; Agly, *Aphis glycines*; Agos, *Aphis gossypii*; Apis, *Acyrtosiphon pisum*; Asol, *Aulacorthum solani*; Cced, *Cinara cedri*; Dnox, *Diuraphis noxia*; Mcer, *Myzus cerasi*; Mper, *Myzus persicae*; Msac, *Melanaphis sacchari*; Mros, *Macrosiphum rosae*; Pnig, *Pentalonia nigronervosa*; Rmai, *Rhopalosiphum maidis*; Rpad, *Rhopalosiphum padi*; Save, *Sitobion avenae*; Sgra, *Schizaphis graminum*; Smis, *Sitobion miscanthi*. No BCR were found in *Eriosoma lanigerum* and *Sipha flava*.

Figure 3. (A) Phylogenetic trees of BCR proteins. Maximum-likelihood phylogenetic tree reconstruction was performed using PhyML (Guindon et al., 2010) with an LG 4-rate class model. Branch-support values were calculated using the bootstrap method, with 1000 replicates. Poorly supported branches (<50%) were collapsed using TreeCollapseCL4 (Hodcroft, 2021). The sequences used for the phylogenetic analysis are listed in SI appendix Table S2. Sequences labelled in red reflect those identified by the Shigenobu and Stern (Shigenobu & Stern, 2013) study in the pea aphid (*A. pisum* genome V3.0). Abbreviations: Acra, *Aphis craccivora*; Agly, *Aphis glycines*; Agos, *Aphis gossypii*; Akon, *Acyrtosiphon kondoi*; Apis, *Acyrtosiphon pisum*; Cced, *Cinara cedri*; Dnox, *Diuraphis noxia*; Dvit, *Daktulosphaira vitifoliae*; Masc, *Myzus ascalonicus*; Mcer, *Myzus cerasi*; Mper, *Myzus persicae*; Msac, *Melanaphis sacchari*; Rmai, *Rhopalosiphum maidis*; Rpad, *Rhopalosiphum padi*; Save, *Sitobion avenae*; Sgra, *Schizaphis graminum*; Tcit, *Toxoptera citricida*.

(B) Repartition of BCR peptides in aphid taxonomic groups. Phylogeny of the 14 members of the aphid group whose genome is annotated and available in databases (among the 19 with sequenced genomes). Their distribution in subfamilies and tribes are indicated in red and black, respectively [adapted from Calevro et al. (2019)]. Colored dots near each species name indicate if a member of each BCR subfamily is present in the genome. Colored dots near each species name indicate if a member of each BCR subfamily is present in the genome of this aphid or not.

BCR4 solution structure

To better characterise the structure evolution of the BCR4 peptide, we have determined its 3D structure by NMR spectroscopy. The ^1H NMR and the natural-abundance ^1H - ^{15}N sofast-HMQC spectra (Schanda et al., 2005) of the protein showed a good dispersion of the amide chemical shifts, indicative of highly structured peptides. Following standard procedures, the analysis of the set of 2D-TOCSY and NOESY spectra allowed a complete assignment of ^1H chemical shifts. This assignment was facilitated by heteronuclear ^1H - ^{15}N and ^1H - ^{13}C NMR spectra, particularly in crowded regions of the ^1H TOCSY and NOESY spectra corresponding to side chains (BRMB entry 34197). The 3D structures were calculated by considering a total of 923 distance restraints, 16 hydrogen bonds, 88 dihedral angles and 3 ambiguous disulfide bridges (Table 4).

Table 4. NMR constraints and structural statistics.

NMR restraints	
<i>Distance restraints</i>	
Total NOE	923
Unambiguous	763
Ambiguous	160
Hydrogen bonds	16
<i>Dihedral Angle Restraints</i>	88
<i>Disulfide bridges^a</i>	3
Structural Statistics (7PQW.pdb)	
<i>Average violations per structure</i>	
NOEs $\geq 0.3 \text{ \AA}$	0
Hydrogen bonds $\geq 0.3 \text{ \AA}$	0
Dihedrals $\geq 15^\circ$	3
Dihedrals $\geq 10^\circ$	8
Average pairwise rmsd ($\text{\AA}$)	0.206
Ramachandran Analysis	
Most favored region and allowed region	96.9
Generously allowed	0.7
Disallowed	2.4
Energies (kcal.mol^{-1}) ^b	
Electrostatic	-1631.48 ± 29.53
Van der Waals	-334.18 ± 11.14
Total energy	-1196.56 ± 31.80
Residual NOE energy	56.38 ± 5.28

^a Introduced as ambiguous; ^b Values are given as mean $\pm$ standard deviation (n=20)

The 200 water-refined structures of BCR4 possess three disulfide bridges with an identical pairing: C17-C34, C21-C32 and C25-C48. Among them, 20 structures were selected in agreement with all NMR experimental data and the standard covalent geometry. Coordinates were deposited as PDB entry 7PQW. Analysis of the 20 final structures with PROCHECK-NMR (Laskowski et al., 1996) showed that almost all of the residues (96.9%) were in the most favored or additionally allowed regions of the Ramachandran diagram (Table 4).

BCR4 is folded into a compact globular unit consisting of an N-terminal tail (D1-T13) and an α -helix (T14-V24) followed by an antiparallel β -sheet composed of two short β -strands, β_1 C34-A37 and β_2 Q43-P46 (Figure 4). The 3D structure is stabilized by three disulfide bridges linking the α -helix to β_1 (C17-C34), to the loop C25-Y33 (C21-C32) and to the C-terminal tail (C25-C48), respectively.

Figure 4. (A) Primary structure of BCR4 peptide. (B) Three-dimensional structure of BCR4. (C) Topology diagram of BCR4. α -helix, β -sheet, and random coil are represented in red, cyan and black, respectively. Disulfide bonds are colored in yellow.

DISCUSSION

Thus far, most pest control strategies have relied on the use of systemic chemical pesticides, which are increasingly stigmatized because of their persistence in the environment and their toxicity to non-target organisms (Goulson, 2018). This creates the need for the development of new pest management solutions, which will undoubtedly be built upon a better investigation of new biological molecules that may be used as potential biopesticides. Small Disulfide Rich Proteins (DRPs) extracted from plants or arthropods are promising alternative biopesticide molecules (Rahoui et al., 2014; Huang et al., 2019; King, 2019). Genomic plasticity of DRP-encoding sequences is known to foster the adaptability of organisms and to enable the acquisition of new functions (Shafee et al., 2017). In this work, we successfully produced in sufficient amount and purity the folded form of BCR4 (Figure 1), a DRP encoded by the pea aphid genome and part of the BCR family of peptides, that was used to explore the insecticidal properties of this protein. We showed that BCR4 strongly interferes with survival and growth of *A. pisum* in a dose-dependent manner (Figure 2). Its range of activity (5-80 μ M, Figure 2) is similar to that displayed by PA1b, a promising plant biopesticide (Gressent et al., 2011; Rahoui et al., 2014) active on the same insect target (Gressent et al., 2007). Importantly, from a functional perspective and as previously reported (Uchi et al., 2019), BCR4 has a significant bactericidal effect on *E. coli* (Table 1), a free-living relative of *B. aphidicola*, the obligatory aphid endosymbiont (Uchi et al., 2019). Based on these results, we propose that the ingestion of exogenous BCR may block nymphal development and induce aphid death by interfering with the population density of this endosymbiont, essential for nymphal growth and survival (Brinza et al., 2009).

From an evolution perspective, a large-scale homology searches through genomic, transcriptomic and proteomic databases was performed to complete the repertoire of BCR peptides, up to now limited to seven identified sequences in the pea aphid genome. Importantly,

the 76 additional sequences we found (bringing to 83 the total number of BCR sequences identified) are all encoded by aphid genomes (Figure 3). Their evolutionary analysis showed that aphid BCRs are organized into four subfamilies including BCR1-2-4-5, BCR3, BCR6 and BCR8 sequences (Figure 3), respectively. BCRs from those four subfamilies all bear 6 or 8 cysteine residues and we observed a very clear intra-group conservation of cysteine topology (e. g. distribution in the sequence and disulfide pairing). But, as for other DRP, BCRs present high sequence diversity in their inter-cysteine loops preventing the detection of any firm homology with arthropod defensins at this moment (Shafee et al., 2016; Shafee et al., 2017). While BCR subfamilies had been previously reported (Shigenobu & Stern, 2013), we were able to enrich each of them with many new members (Figure 3). We also showed that (i) aphids present varying numbers of BCR-encoding genes, from 1 to 13 and (ii) while BCR3 and BCR8 homologs are widely spread in the aphid lineage, genes from the BCR1-2-4-5 and BCR6 subfamilies are restricted to the Macrosiphini tribe, which includes many major agricultural pests. This suggests a complex evolutionary history involving several events of gene duplications and losses with possible functional diversification of the resulting homologs.

To study the structure evolution of the BCR4 peptide, its three-dimensional structure was determined by NMR spectroscopy and careful protein fold analysis revealed that BCR4 belongs to a yet unknown structural class of defensin proteins (Figure 4). Defensins are a well-characterized group of DRPs present in all eukaryotes' genomes (Shafee et al., 2016). Two superfamilies have been described to date, each derived from independent evolutionary event: the so-called trans-defensin superfamily is uniquely composed of the CS $\alpha\beta$ (cysteine-stabilized α -helix β -sheet) family and the cis-defensin superfamily is composed of the α -, β -, θ -, and big defensins families (Shafee et al., 2016; Shafee et al., 2017). Contrary to the CS $\alpha\beta$ family (Figure 5), the CXC motif (X denoting any amino acid residue) of BCR4 is in the first β -strand, leading to a different cysteine bonding pattern, the one corresponding to the β -defensin fold

(the second family of defensin well-known as antimicrobial peptides) (Shafee et al., 2016; Shafee et al., 2017). The fold of BCR4 is therefore a new type of defensin peptide suggesting that BCRs constitute a new class of antimicrobial cysteine-rich proteins. Shafee's phylogenetic work showed that the defensins consist of two independent and convergent superfamilies (Shafee et al., 2016; Shafee et al., 2017). We here hypothesize that the defensins consist of at least three independent and convergent superfamilies: cis-, trans- and BCR defensins

Figure 5. Disulfide connectivities in BCR4 peptide and in C α β and β defensin groups. α -helices indicated by rectangle, β -strands represented by arrows, and disulfides in black lines.

BCRs are orphan genes exclusively expressed in aphid bacteriocytes. This gene family has probably evolved in aphid lineages to ensure several functions related to endosymbiosis, including bacteriocyte homeostasis and endosymbiont control. Related studies on cereal weevils *Sitophilus* have shown that the coleopteracin A (ColA) AMP selectively targets endosymbionts within bacteriocytes and impairs bacterial cytokinesis, thereby regulating bacterial cell division and preventing bacterial exit from weevil bacteriocytes (Login et al., 2011; Login & Heddi, 2013). Similar results were obtained in the *Rhizobium*-legumes symbiosis, where nodule-specific secreted peptides called NCRs were shown to target bacteria and induce an irreversible elongation of bacteria, rendering them metabolically active but unable to multiply *in vitro* outside plant nodules (Van de Velde et al., 2010). In the actinorhizal plant *Alnus glutinosa*, nodules express defensin-like peptides, including the Ag5 peptide that

targets symbiont vesicles and increases permeability of vesicle membranes (Carro et al., 2015). Small peptides are becoming emerging molecules that target a variety of functions in host-symbiont interaction, including symbiont control and physiology (Mergaert et al., 2017). Such a convergent evolution pattern is most interesting in its applied perspectives.

Overall, the susceptibility of aphids to BCR peptides may lead to the development of effective strategies for controlling such sucking pests. The exploration of this work may end up in a new protein family lead for the specific control of aphids, which include some of the most important pests of global agriculture.

Materials and Methods

Peptide synthesis

BCR4 was synthesized through native chemical ligation (NCL) of two peptide segments, followed by oxidative folding to form the three disulfide bridges, using already described protocols (Lelievre et al., 2016; Terrier et al., 2016; Terrier et al., 2017; Abboud & Aucagne, 2020; Abboud et al., 2021a; Abboud et al., 2021b). Briefly, a N-terminal cysteinyl peptide segment (sequence: H-²¹CAVVCNYTSRPCYCV EAAKERDQWFPYCY⁵⁰D-OH) and a C-terminal crypto-thioester segment (sequence: H-¹DFDPTEFKGPFPPTIEICK²⁰Y-(Hnb)C(StBu)G-NH₂) were synthesized on a Prelude peptide synthesizer (Protein Technologies, Tucson, AZ, USA) using standard Fmoc/tBu chemistry at a 25 µmol scale and starting from a Tentagel resin equipped with a Rink's amide linker. The *N*-2-hydroxy-5-nitrobenzyl (Hnb) group was automatically introduced through resin reductive amination. Both peptide segments were purified by C₁₈ reverse phase (RP)-HPLC on a Nucleosil C₁₈ column (300 Å, 250 × 10 mm) (Macherey-Nagel, Düren, Germany), maintained at 25°C. Solvent A was 0.1% trifluoroacetic acid (TFA) in water and B was 0.1% TFA in acetonitrile. A linear gradient, from 20% B to 55% B, was applied for 30 min at a flow rate of 3 ml/min and afforded the pure

reduced form of BCR4 through collection of the peak identified as the target compound. The purified peptides were then chemoselectively coupled under standard NCL conditions (2 mM peptide segments, 50 mM tris-carboxyethylphosphine, 200 mM 4-mercaptophenylacetic acid, 6 M guanidinium chloride, 200 mM phosphate buffer, pH 6.5, 37 °C, 24 h) (Lelievre et al., 2016; Terrier et al., 2016) and purified as previously to retrieve the pure reduced form of BCR4 (sequence:

H-¹DFDPTEFKGPFPTIEICKSKYCAVVCNYTSRPCYCVEAAKERDQWFPYCY⁵⁰D-OH).

The purified compound was further analyzed by ESI-HRMS on a Bruker maXisTM 22 ultra-high-resolution Q-TOF mass spectrometer (Bruker Daltonics, Bremen, Germany), in positive mode and a [M+H]⁺ *m/z* ratio of 5897.5883 was found (theoretical monoisotopic *m/z* calculated for C₂₆₆H₃₇₉N₆₂O₇₉S₆: 5897.5869). Subsequently, the pure reduced form of BCR4 was oxidatively folded *in vitro* (10 μM peptide concentration, 0.1 mM oxidized glutathione, 1 mM glutathione, 1 mM EDTA, 100 mM TRIS, pH 8.5, 20°C, 48 h) (Silva et al., 2009; Derache et al., 2012) before purification to homogeneity by C₁₈ RP-HPLC. Analysis of the purified folded product *via* ESI-HRMS analysis gave results consistent with the formation of three disulfide bridges ([M+H]⁺ *m/z* obtained: 5891.5437; theoretical monoisotopic *m/z* calculated for C₂₆₆H₃₇₃N₆₂O₇₉S₆: 5891.5400).

Antimicrobial assays

E. coli and *M. luteus* were grown in Lysogeny Broth (LB) and Terrific Broth (TB) medium (Sigma-Aldrich, Saint-Louis, MO, USA), respectively. Antimicrobial assays were performed in sterile 96-well plates with a final volume of 100 μl per well, composed of 50 μl of culture and 50 μl of serially diluted peptides (5-80 μM). *E. coli* and *M. luteus* were added at an OD₆₀₀ of 0.006 and 0.05, respectively. Plates were incubated at 30 °C for 24 h and growth was measured at 600 nm using a Power wave XS-Biotek plate reader (Biotek Instrument, Colmar, France). The lowest concentration of BCR4 peptide showing complete inhibition was taken as

minimal inhibitory concentration (MIC). All analyses were performed in triplicate, with the results expressed as mean $\pm$ standard deviation of mean (SEM).

Insects and insect assays

The aphid clone used was *A. pisum* LL01, a long-established alfalfa-collected clone containing only the primary endosymbiont *B. aphidicola*. Aphids were maintained on young broad bean plants (*Vicia faba* L. cv. Aguadulce) at 21 °C, with a photoperiod of 16 h light - 8 h dark to obtain strictly parthenogenetic aphid matriline, that were reared and synchronized as previously described (Simonet et al., 2018).

For toxicity analyses, three groups of 10^{1st} instar nymphs (aged between 0 and 24 hours) were collected and placed in *ad hoc* feeding chambers containing an AP3 artificial diet (Febvay et al., 1988) supplemented with different experimental doses (5 μ M to 80 μ M) of solubilised BCR4 peptide. Toxicity was evaluated by scoring survival daily over the whole nymphal life of the pea aphid (7 days) (Simonet et al., 2016). Growth was also measured by weighing adult aphids on a Mettler AE163 analytical microbalance (Mettler Toledo, Columbus, OH, USA) at the closest 10 μ g, following the protocol described previously (Rahbe & Febvay, 1993; Loth et al., 2015).

Aphid mortality data for all BCR4 concentrations were analysed separately in a parametric survival analysis with a log-normal fit. Aphid weights were analysed by Anova followed by Tukey-Kramer HSD test for comparing multiple means. All analyses were made with JMP software version 11 (SAS Institute Cary USA, MacOS version).

Sequence analysis and phylogeny

Homologous BCR protein sequences were retrieved using a combination of TBLASTN and BLASTP (Altschul et al., 1997; Boratyn et al., 2012) against the aphid genomes available from (i) AphidBase (Legeai et al., 2010), (ii) the NCBI genome database and (iii) the whole NCBI

non-redundant protein, nucleotide and EST databases (with *A. pisum* BCR as blast seeds; see SI appendix Table S2 for a complete list of BCR protein sequences used for the phylogenetic analysis). BCR protein sequences were subjected to multiple sequence alignments using the MUSCLE program (Edgar, 2004). Subsequently, a phylogenetic tree was constructed, using the PhyML method (Guindon et al., 2010) implemented in the Seaview software (v5.0.4) (Gouy et al., 2010; Guindon et al., 2010) (LG model with 4 rate classes), and the reliability of each branch was evaluated using the bootstrap method, with 1000 replicates. Poorly supported branches (<50%) were collapsed using TreeCollapseCL4 (Hodcroft, 2021). Graphical representation and editing of the phylogenetic tree were performed with FigTree (v1.4.3) (Drummond & Rambaut, 2007).

NMR Experiments

Prior to NMR analysis, the synthesized BCR4 peptides were dissolved in H₂O:D₂O (9:1 ratio) at a concentration of 0.6 mM and pH was adjusted to 4.8. Then, 2D ¹H NOESY, 2D ¹H TOCSY, a sofast-HMQC (Schanda et al., 2005) (¹⁵N natural abundance) and a ¹³C-HSQC (¹³C natural abundance) were performed at 298 K on an Avance III HD BRUKER 700 MHz spectrometer equipped with a cryoprobe. ¹H chemical shifts were referenced to the water signal (4.77 ppm at 298 K). NMR data were processed using the Topspin software version 3.2TM (Bruker, Billerica, MA, USA) and analyzed with CCPNMR version 2.2.2 (Vranken et al., 2005).

Structure calculations

Structures were calculated using the Crystallography and NMR System (Brunger et al., 1998; Brunger, 2007) through the automatic assignment software ARIA2 version 2.3 (Rieping et al., 2007) with NOE derived distances, hydrogen bonds (in accordance with the observation of typical long or medium distance NOE cross peaks network for β-sheets and α-helices respectively – HN/HN, HN/Hα, Hα/Hα), backbone dihedral angle restraints (determined with

the DANGLE program (Cheung et al., 2010) and three ambiguous disulfide bridges. The ARIA2 protocol, with default parameters used, simulated annealing with torsion angle and Cartesian space dynamics. The iterative process was repeated until the assignment of the NOE cross peaks was complete. The last run for BCR4 was performed with 1000 initial structures and 200 structures were refined in water. 20 structures were selected on the basis of total energies and restraint violation statistics, to represent the structure of BCR4 in solution. The quality of final structures was evaluated using PROCHECK-NMR (Laskowski et al., 1996) and PROMOTIF (Hutchinson & Thornton, 1996). The figures were prepared with PYMOL (De Lano, 2002).

REFERENCES

- Abboud, S. A., Amoura, M., Madinier, J. B., Renoux, B., Papot, S., Piller, V., & Aucagne, V. (2021a). Enzyme-cleavable linkers for protein chemical synthesis through solid-phase ligations. *Angew Chem Int Ed Engl*, 60(34), 18612-18618. doi: 10.1002/anie.202103768
- Abboud, S. A., & Aucagne, V. (2020). An optimized protocol for the synthesis of N-2-hydroxybenzyl-cysteine peptide crypto-thioesters. *Org Biomol Chem*, 18(40), 8199-8208. doi: 10.1039/d0ob01737j
- Abboud, S. A., Cisse, E. H., Doudeau, M., Benedetti, H., & Aucagne, V. (2021b). A straightforward methodology to overcome solubility challenges for N-terminal cysteinyl peptide segments used in native chemical ligation. *Chemical Science*, 12(9), 3194-3201. doi: 10.1039/d0sc06001a
- Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J. H., Zhang, Z., Miller, W., & Lipman, D. J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research*, 25(17), 3389-3402. doi: 10.1093/nar/25.17.3389

- 459 Baumann, P. (2005). Biology bacteriocyte-associated endosymbionts of plant sap-sucking
460 insects. *Annu Rev Microbiol*, 59, 155-189. doi:
461 10.1146/annurev.micro.59.030804.121041
- 462 Bell, J., Sukiran, N. A., Walsh, S., & Fitches, E. C. (2021). The insecticidal activity of
463 recombinant nemertide toxin alpha-1 from *Lineus longissimus* towards pests and
464 beneficial species. *Toxicon*, 197, 79-86. doi: 10.1016/j.toxicon.2021.04.003
- 465 Boratyn, G. M., Schaffer, A. A., Agarwala, R., Altschul, S. F., Lipman, D. J., & Madden, T. L.
466 (2012). Domain enhanced lookup time accelerated BLAST. *Biology Direct*, 7. doi: Artn
467 1210.1186/1745-6150-7-12
- 468 Bradshaw, C. J. A., Leroy, B., Bellard, C., Roiz, D., Albert, C., Fournier, A., . . . Courchamp,
469 F. (2016). Massive yet grossly underestimated global costs of invasive insects. *Nature*
470 *Communications*, 7, 12986. doi: 10.1038/ncomms12986
- 471 Brinza, L., Viñuelas, J., Cottret, L., Calevro, F., Rahbé, Y., Febvay, G., . . . Charles, H. (2009).
472 Systemic analysis of the symbiotic function of *Buchnera aphidicola*, the primary
473 endosymbiont of the pea aphid *Acyrtosiphon pisum*. *Comptes Rendus Biologies*,
474 332(11), 1034-1049. doi: <https://doi.org/10.1016/j.crvi.2009.09.007>
- 475 Brunger, A. T. (2007). Version 1.2 of the Crystallography and NMR system. *Nature Protocols*,
476 2(11), 2728-2733. doi: 10.1038/nprot.2007.406
- 477 Brunger, A. T., Adams, P. D., Clore, G. M., DeLano, W. L., Gros, P., Grosse-Kunstleve, R. W.,
478 . . . Warren, G. L. (1998). Crystallography & NMR system: A new software suite for
479 macromolecular structure determination. *Acta Crystallographica Section D-Biological*
480 *Crystallography*, 54, 905-921. doi: Doi 10.1107/S09074444998003254
- 481 Calevro, F., Tagu, D., & Callaerts, P. (2019). *Acyrtosiphon pisum*. *Trends Genet*, 35(10), 781-
482 782. doi: 10.1016/j.tig.2019.07.003

- 483 Carro, L., Pujic, P., Alloisio, N., Fournier, P., Boubakri, H., Hay, A. E., . . . Normand, P. (2015).
484 Alnus peptides modify membrane porosity and induce the release of nitrogen-rich
485 metabolites from nitrogen-fixing *Frankia*. *ISME J*, 9(8), 1723-1733. doi:
486 10.1038/ismej.2014.257
- 487 Cheung, M. S., Maguire, M. L., Stevens, T. J., & Broadhurst, R. W. (2010). DANGLE: A
488 Bayesian inferential method for predicting protein backbone dihedral angles and
489 secondary structure. *Journal of Magnetic Resonance*, 202(2), 223-233. doi:
490 10.1016/j.jmr.2009.11.008
- 491 Da Silva, P., Strzepa, A., Jouvensal, L., Rahioui, I., Gressent, F., & Delmas, A. F. (2009). A
492 folded and functional synthetic PA1b, an interlocked entomotoxic miniprotein.
493 *Biopolymers*, 92. doi: 10.1002/bip.21217
- 494 Dawson, P. E., Muir, T. W., Clark-Lewis, I., & Kent, S. B. (1994). Synthesis of proteins by
495 native chemical ligation. *Science*, 266(5186), 776-779. doi: 10.1126/science.7973629
- 496 De Lano, W. L. (2002). Pymol. *De Lano Scientific, South San Francisco, CA*.
- 497 Dedryver, C. A., Le Ralec, A., & Fabre, F. (2010). The conflicting relationships between aphids
498 and men: A review of aphid damage and control strategies. *Comptes Rendus Biologies*,
499 333(6-7), 539-553. doi: 10.1016/j.crv.2010.03.009
- 500 Delobel, B., Grenier, A. M., Gueguen, J., Ferrasson, E., & Mbaiguinam, M. (1998). Use of
501 polypeptide derived from a palb legume albumen as insecticide. *Paris Patent*
502 No:Brevet-98/05877
- 503 Derache, C., Meudal, H., Aucagne, V., Mark, K. J., Cadene, M., Delmas, A. F., . . . Landon, C.
504 (2012). Initial insights into structure-activity relationships of avian defensins. *J Biol*
505 *Chem*, 287(10), 7746-7755. doi: 10.1074/jbc.M111.312108

- Diagne, C., Leroy, B., Gozlan, R. E., Vaissiere, A. C., Assailly, C., Nuninger, L., . . . Courchamp, F. (2020). InvaCost, a public database of the economic costs of biological invasions worldwide. *Sci Data*, 7(1), 277. doi: 10.1038/s41597-020-00586-z
- Douglas, A. E. (2015). Multiorganismal insects: diversity and function of resident microorganisms. *Annu Rev Entomol*, 60, 17-34. doi: 10.1146/annurev-ento-010814-020822
- Drummond, A. J., & Rambaut, A. (2007). BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evol Biol*, 7. doi: Artn 21410.1186/1471-2148-7-214
- Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*, 32(5), 1792-1797. doi: 10.1093/nar/gkh340
- Febvay, G., Delobel, B., & Rahbe, Y. (1988). Influence of the amino-Acid balance on the improvement of an artificial diet for a biotype of *Acyrtosiphon pisum* (Homoptera, Aphididae). *Canadian Journal of Zoology-Revue Canadienne De Zoologie*, 66(11), 2449-2453. doi: 10.1139/Z88-362
- Gonzalez-Castro, R., Gomez-Lim, M. A., & Plisson, F. (2021). Cysteine-rich peptides: hyperstable scaffolds for protein engineering. *Chembiochem*, 22(6), 961-973. doi: 10.1002/cbic.202000634
- Goulson, D. (2018). Call to restrict neonicotinoids. *Science*, 360(6392), 973-973. doi: 10.1126/science.aau0432
- Gouy, M., Guindon, S., & Gascuel, O. (2010). SeaView version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol*, 27. doi: 10.1093/molbev/msp259
- Gressent, F., Da Silva, P., Eyraud, V., Karaki, L., & Royer, C. (2011). Pea Albumin 1 subunit b (PA1b), a promising bioinsecticide of plant origin. *Toxins (Basel)*, 3(12), 1502-1517. doi: 10.3390/toxins3121502

- Gressent, F., Duport, G., Rahioui, I., Pauchet, Y., Bolland, P., Specty, O., & Rahbe, Y. (2007). Biological activity and binding site characteristics of the PA1b entomotoxin on insects from different orders. *J Insect Sci*, 7. doi: 10.1673/031.007.1201
- Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W., & Gascuel, O. (2010). New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology*, 59. doi: 10.1093/sysbio/syq010
- Hodcroft, E. TreeCollapseCL 4. from <http://emmahodcroft.com/TreeCollapseCL.html>, accessed July 2021
- Huang, Y. H., Du, Q., & Craik, D. J. (2019). Cyclotides: disulfide-rich peptide toxins in plants. *Toxicon*, 172, 33-44. doi: 10.1016/j.toxicon.2019.10.244
- Hutchinson, E. G., & Thornton, J. M. (1996). PROMOTIF - A program to identify and analyze structural motifs in proteins. *Protein Science*, 5(2), 212-220.
- Huygens, C., Ribeiro Lopes M., Gaget K., Duport G., Peignier S., De Groef S., Parisot N., Calevro F., & Callaerts P. (2021). Evolutionary diversification of insulin-related peptides (IRPs) in aphids and spatiotemporal distribution in *Acyrtosiphon pisum*. *Insect Biochem Mol Biol*, 103670. doi: 10.1016/j.ibmb.2021.103670.
- King, G. F. (2019). Tying pest insects in knots: the deployment of spider-venom-derived knottins as bioinsecticides. *Pest Management Science*, 75(9), 2437-2445. doi: 10.1002/ps.5452
- Laskowski, R. A., Rullmann, J. A. C., MacArthur, M. W., Kaptein, R., & Thornton, J. M. (1996). AQUA and PROCHECK-NMR: Programs for checking the quality of protein structures solved by NMR. *Journal of Biomolecular Nmr*, 8(4), 477-486. doi: 10.1007/Bf00228148

- 554 Legeai, F., Shigenobu, S., Gauthier, J. P., Colbourne, J., Rispe, C., Collin, O., . . . Tagu, D.
555 (2010). AphidBase: a centralized bioinformatic resource for annotation of the pea aphid
556 genome. *Insect Mol Biol*, 19 Suppl 2, 5-12. doi: 10.1111/j.1365-2583.2009.00930.x
- 557 Lelievre, D., Terrier, V. P., Delmas, A. F., & Aucagne, V. (2016). Native chemical ligation
558 strategy to overcome side reactions during Fmoc-Based Synthesis of C-terminal
559 cysteine-containing peptides. *Org Lett*, 18(5), 920-923. doi:
560 10.1021/acs.orglett.5b03612
- 561 Login, F. H., Balmand, S., Vallier, A., Vincent-Monegat, C., Vigneron, A., Weiss-Gayet, M., .
562 . . Heddi, A. (2011). Antimicrobial peptides keep insect endosymbionts under control.
563 *Science*, 334(6054), 362-365. doi: 10.1126/science.1209728
- 564 Login, F. H., & Heddi, A. (2013). Insect immune system maintains long-term resident bacteria
565 through a local response. *Journal of Insect Physiology*, 59(2), 232-239. doi:
566 10.1016/j.jinsphys.2012.06.015
- 567 Loth, K., Costechareyre, D., Effantin, G., Rahbe, Y., Condemine, G., Landon, C., & da Silva,
568 P. (2015). New Cyt-like delta-endotoxins from *Dickeya dadantii*: structure and
569 aphicidal activity. *Sci Rep*, 5, 8791. doi: 10.1038/srep08791
- 570 Martinez, G., Hograindleur, J. P., Voisin, S., Abi Nahed, R., Abd El Aziz, T. M., Escoffier, J.,
571 . . . Arnoult, C. (2016). Spermaurin, an La1-like peptide from the venom of the scorpion
572 *Scorpio maurus palmatus*, improves sperm motility and fertilization in different
573 mammalian species. *Mol Hum Reprod*. doi: 10.1093/molehr/gaw075
- 574 Mergaert, P., Kikuchi, Y., Shigenobu, S., & Nowack, E. C. M. (2017). Metabolic Integration
575 of Bacterial Endosymbionts through Antimicrobial Peptides. *Trends Microbiol*, 25(9),
576 703-712. doi: 10.1016/j.tim.2017.04.007

- Oerke, E. C., & Dehne, H. W. (2004). Safeguarding production - losses in major crops and the role of crop protection. *Crop Protection*, 23(4), 275-285. doi: 10.1016/j.cropro.2003.10.001
- Rahbe, Y., & Febvay, G. (1993). Protein toxicity to aphids - an *in vitro* test on *Acyrtosiphon pisum*. *Entomologia Experimentalis Et Applicata*, 67(2), 149-160. doi: 10.1111/j.1570-7458.1993.tb01663.x
- Rahoui, I., Eyraud, V., Karaki, L., Sasse, F., Carre-Pierrat, M., Qin, A., . . . Gressent, F. (2014). Host range of the potential biopesticide Pea Albumin 1b (PA1b) is limited to insects. *Toxicon*, 89, 67-76. doi: 10.1016/j.toxicon.2014.07.004
- Ribeiro Lopes, M., Parisot N., Gaget K., Huygens C., Peigner S., Duport G., . . . , Calevro F. (2020). Evolutionary novelty in the apoptotic pathway of aphids. *Proc Natl Acad Sci U S A*, 117(51):32545-32556. doi: 10.1073/pnas.2013847117
- Rieping, W., Habeck, M., Bardiaux, B., Bernard, A., Malliavin, T. E., & Nilges, M. (2007). ARIA2: Automated NOE assignment and data integration in NMR structure calculation. *Bioinformatics*, 23(3), 381-382. doi: 10.1093/bioinformatics/btl589
- Schanda, P., Kupce, E., & Brutscher, B. (2005). SOFAST-HMQC experiments for recording two-dimensional heteronuclear correlation spectra of proteins within a few seconds. *Journal of Biomolecular Nmr*, 33(4), 199-211. doi: 10.1007/s10858-005-4425-x
- Shafee, T. M., Lay, F. T., Hulett, M. D., & Anderson, M. A. (2016). The defensins consist of two independent, convergent protein superfamilies. *Mol Biol Evol*, 33(9), 2345-2356. doi: 10.1093/molbev/msw106
- Shafee, T. M., Lay, F. T., Phan, T. K., Anderson, M. A., & Hulett, M. D. (2017). Convergent evolution of defensin sequence, structure and function. *Cell Mol Life Sci*, 74(4), 663-682. doi: 10.1007/s00018-016-2344-5

- Shigenobu, S., & Stern, D. L. (2013). Aphids evolved novel secreted proteins for symbiosis with bacterial endosymbiont. *Proc Biol Sci*, 280(1750), 20121952. doi: 10.1098/rspb.2012.1952
- Shigenobu S., Watanabe H., Hattori M., Sakaki Y., & Ishikawa H. (2000). Genome sequence of the endocellular bacterial symbiont of aphids Buchnera sp. APS. *Nature*, 407(6800):81-6. doi: 10.1038/35024074.
- Simonet, P., Gaget, K., Balmand, S., Ribeiro Lopes, M., Parisot, N., Buhler, K., . . . Calevro, F. (2018). Bacteriocyte cell death in the pea aphid/Buchnera symbiotic system. *Proc Natl Acad Sci U S A*, 115(8), E1819-E1828. doi: 10.1073/pnas.1720237115
- Simonet, P., Gaget, K., Parisot, N., Duport, G., Rey, M., Febvay, G., . . . Calevro, F. (2016). Disruption of phenylalanine hydroxylase reduces adult lifespan and fecundity, and impairs embryonic development in parthenogenetic pea aphids. *Sci Rep*, 6, 34321. doi: 10.1038/srep34321
- Terrier, V. P., Adihou, H., Arnould, M., Delmas, A. F., & Aucagne, V. (2016). A straightforward method for automated Fmoc-based synthesis of bio-inspired peptide crypto-thioesters. *Chemical Science*, 7(1), 339-345. doi: 10.1039/c5sc02630j
- Terrier, V. P., Delmas, A. F., & Aucagne, V. (2017). Efficient synthesis of cysteine-rich cyclic peptides through intramolecular native chemical ligation of N-Hnb-Cys peptide crypto-thioesters. *Org Biomol Chem*, 15(2), 316-319. doi: 10.1039/c6ob02546c
- Uchi, N., Fukudome, M., Nozaki, N., Suzuki, M., Osuki, K. I., Shigenobu, S., & Uchiumi, T. (2019). Antimicrobial activities of cysteine-rich peptides specific to bacteriocytes of the pea aphid *Acyrtosiphon pisum*. *Microbes Environ*, 34(2), 155-160. doi: 10.1264/jsme2.ME18148

Van de Velde, W., Zehirov, G., Szatmari, A., Debreczeny, M., Ishihara, H., Kevei, Z., . . . Mergaert, P. (2010). Plant peptides govern terminal differentiation of bacteria in symbiosis. *Science*, 327(5969), 1122-1126. doi: 10.1126/science.1184057

Vranken, W. F., Boucher, W., Stevens, T. J., Fogh, R. H., Pajon, A., Llinas, P., . . . Laue, E. D. (2005). The CCPN data model for NMR spectroscopy: Development of a software pipeline. *Proteins-Structure Function and Bioinformatics*, 59(4), 687-696. doi: 10.1002/prot.20449

ACKNOWLEDGEMENTS

This work was supported by INRAE (Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement), INSA Lyon (Institut National des Sciences Appliquées Lyon) and the French ANR-119-CE11-0004-01 (Biofamily) program grant. We thank Philippe Marceau and Jean-Baptiste Madinier for assistance in solid phase peptide synthesis.

We declare no competing interests.

AUTHORS' CONTRIBUTIONS

Karine Loth: investigation, formal analysis, data curation, validation, visualisation; Nicolas Parisot: investigation, formal analysis, data curation, validation, visualisation; Françoise Paquet: investigation, formal analysis, data curation, validation, visualisation, Catherine Sivignon: investigation; Isabelle Rahioui : investigation; Mélanie Ribeiro Lopes: formal analysis, visualization; writing—review & editing; Karen Gaget: resources; Gabrielle Duport : resources, Agnès F. Delmas : conceptualisation, writing—review & editing, Vincent Aucagne: conceptualisation, investigation, formal analysis, data curation, validation, visualisation, writing—review & editing, Abdelaziz Heddi writing—review & editing, Federica Calevro conceptualisation, supervision, writing—review & editing, funding Acquisition and Pedro da

647 Silva : conceptualisation, supervision, project administration, writing—original draft, Funding
648 Acquisition

649 **COMPETING FINANCIAL INTERESTS**

650 The authors declare no competing financial interests

651