1 The interaction of climate history and evolution impacts alpine biodiversity assembly **2 differently in freshwater and on land**

3 Luiz Jardim-deQueiroz^{1,2}, Carmela J. Doenz^{1,2}, Florian Altermatt^{3,4}, Roman Alther^{3,4}, Špela
4 Borko⁵, Jakob Brodersen^{1,2}, Martin M. Gossner^{6,7}, Catherine Graham⁶, Blake Matthews^{1,2},
5 Ian R. McFadden^{6,7}, Loïc Pellissier^{6,7}, Thomas Schmitt^{8,9}, Oliver M. Selz^{1,*}, Soraya Villalba¹,
6 Lukas Rüber^{2,10}, Niklaus Zimmermann^{6,7}, Ole Seehausen^{1,2,‡}

7 ¹ Department of Fish Ecology and Evolution, Swiss Federal Institute of Aquatic Science and
8 Technology (Eawag), Switzerland.

9 ² Institute of Ecology and Evolution, University of Bern, Switzerland.

10 ³ Department of Aquatic Ecology, Swiss Federal Institute of Aquatic Science and
11 Technology (Eawag), Switzerland.

12 ⁴ Department of Evolutionary Biology and Environmental Studies, University of Zurich,
13 Switzerland.

14 ⁵ SubBio Lab, Department of Biology, Biotechnical Faculty, University of Ljubljana,
15 Slovenia.

16 ⁶ Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Switzerland.

17 ⁷ Department of Environmental Systems Science, Swiss Federal Institute of Technology in
18 Zürich (ETHZ), Switzerland.

19 ⁸ Senckenberg German Entomological Institute, Germany.

20 ⁹ Institute of Biochemistry and Biology, Faculty of Science, University of Potsdam,
21 Germany.

22 ¹⁰ Naturhistorisches Museum Bern (NMBE), Switzerland.

23 ORCIDs: LJQ: 0000-0002-5524-6281; CJD: 0000-0003-4143-7623; FA: 0000-0002-4831-
24 6958; RA: 0000-0001-7582-3966; ŠP: 0000-0002-8383-8778; JB: 0000-0003-2060-6379;
25 MMG: 0000-0003-1516-6364; CG: 0000-0001-9267-7948; BM: 0000-0001-9089-704X;
26 IRM: 0000-0002-4508-7272; LP: 0000-0002-2289-8259; TS: 0000-0002-1389-8396; OMS:
27 0000-0002-2210-5909; LR: 0000-0003-0125-008X; NZ: 0000-0003-3099-9604; OS: 0000-
28 0001-6598-1434.

29 * Authorship alphabetically ordered from FA to SV.

30 ‡ Corresponding author: Ole.Seehausen@eawag.ch; Ole.Seehausen@iee.unibe.ch.

31 **Abstract**

32 Quaternary climate fluctuations can affect biodiversity assembly through speciation in two
 33 non-mutually-exclusive ways: a glacial species pump, where isolation in glacial refugia
 34 accelerates allopatric speciation, and adaptive radiation during ice-free periods. Here we
 35 detected biogeographic and genetic signatures associated with both mechanisms in the
 36 generation of the European Alps biodiversity. Age distributions of endemic and widespread
 37 species within aquatic and terrestrial taxa (amphipods, fishes, amphibians, butterflies and
 38 flowering plants) revealed that endemic fish evolved only in lakes, are highly sympatric and
 39 mainly of Holocene age, consistent with adaptive radiation. Endemic amphipods are ancient,
 40 suggesting preglacial radiation with limited range expansion and local Pleistocene survival,
 41 perhaps facilitated by a groundwater-dwelling lifestyle. Terrestrial endemics are mostly of
 42 Pleistocene age, and are thus more consistent with the glacial species pump. The lack of
 43 evidence for Holocene adaptive radiation in the terrestrial biome may be attributable to a
 44 faster range expansion of these taxa after glacial retreats, though fewer stable environments
 45 may also have contributed to differences between terrestrial areas and lakes. The high
 46 proportion of young, endemic species make the Alps vulnerable to climate change, but the
 47 mechanisms and consequences of species loss will likely differ between biomes because of
 48 their distinct histories.

49 **Keywords:** Time for speciation, Allopatric speciation, Adaptive radiation, Pleistocene
 50 refugia, Glacial species pump, European Alps

51 **1. Background**

52 Immigration, speciation and extinction are the three main processes underlying the assembly
 53 of biodiversity in island-like habitats [1–4]. The relative contribution of these processes
 54 depends on size, isolation and fragmentation of the region, ecosystem or habitat. For instance,
 55 immigration rates decrease with increasing isolation, extinction rates decrease with increasing
 56 area, and rates of in situ speciation increase with both area, isolation, and fragmentation
 57 [1,2,5–7]. The occurrence and interaction of these processes over geological history leave
 58 strong imprints in the contemporary structure of regional and local species assemblages,
 59 including phylogenetic structure and relatedness. The species age distribution, and the nature
 60 and degree of endemism are some of the resulting biodiversity features [8].

61 Some of the mechanisms that can lead to endemism are through cladogenetic speciation, i.e.,
 62 when an ancestral species diverges into two or more derived species within an island,

63 archipelago or geographic region, or through anagenetic speciation, when a local or regional
 64 population or set of populations diverges from its progenitors outside the island, archipelago
 65 or region [3]. Recent cladogenetic and anagenetic speciation both result in neoendemic
 66 species, which are young species with geographically restricted distributions. Moreover, if
 67 cladogenetic and anagenetic speciation are the main processes behind regional biodiversity
 68 assembly, a regional biota can be composed of many relatively young and closely related
 69 species. Non-endemic species, in turn, are generally more widespread because they either
 70 have immigrated to the focal region from the outside after range expansion or they have
 71 arisen in the focal region and had time to spread beyond it. Such non-endemic species are
 72 also expected to be older, according to the ‘age and area’ hypothesis [9], which predicts that
 73 older species have had more time to disperse and hence become geographically more
 74 widespread, whereas younger species often are still confined to smaller ranges. However, the
 75 ‘age and area’ hypothesis assumes biome stability (including climatic stability) and does not
 76 consider factors other than age that could in fact have strong effects on species range sizes.
 77 For instance, population extirpation or local extinction [10], the presence and movement of
 78 physical or climatic barriers in space and time [11], changes in habitat size through time,
 79 variation between lineages in the ecological versatility and evolutionary adaptability of
 80 species [12], variation in species dispersal ability [13,14] and ecological interactions [15] can
 81 all be important predictors of species range size in addition to species age. The interaction of
 82 these factors can explain, for instance, why some species, despite being so old, are
 83 geographically narrowly confined in the present time (geographical relicts or paleoendemics)
 84 [16].

85 Physically rugged mountain landscapes at lower and intermediate latitude, such as those of
 86 the Alpine bioregion of Europe (hereafter the European Alps), are hotspots of biodiversity
 87 and endemism [17–19]. In such environments, endemism and species radiations arise through
 88 the interaction of dispersal limitation with steep ecological gradients and often archipelago-
 89 like habitat structures [20–23]. In the European Alps, multiple terrestrial taxa have undergone
 90 local radiations leading to the emergence of endemic clades in several groups such as
 91 flowering plants [24–26] or butterflies [27,28]. Furthermore, some of the largest endemic
 92 radiations in European freshwater habitats also took place in or around the Alps, especially
 93 for amphipods [29–32] and fish [33,34]. Importantly, these radiations occurred in habitats
 94 that are geographically isolated from similar habitats elsewhere, but surrounded by less
 95 isolated habitats, containing diverse assemblages of widely distributed taxa. For example,

96 mountain-tops surrounded by lowlands, or permanently cold, deep lakes isolated from other
97 such lakes by the seasonally relatively warm, shallow flowing water of rivers.

98 The climatic and habitat instability driven by the Quaternary climate fluctuations [35] has
99 interacted evolutionary and ecological processes to shape biodiversity in the Alps [34,36–41].

100 This includes influences on speciation, extinction and immigration of lineages, and the
101 reshaping of species abundance, range distribution, richness and genetic diversity patterns
102 [42–44]. An important fraction of biodiversity in the Alps is due to recently colonizing
103 species that immigrated into the region from far away (such as central Asia), or expanded
104 their range from adjacent regions but have not yet speciated in the area. On the other hand,
105 endemic biodiversity may have emerged through two alternative, non-exclusive mechanisms
106 both driven by the succession of glacial–interglacial cycles: (1) the glacial species pump
107 [45,46], and (2) adaptive radiation during interglacial periods (hereafter adaptive radiation)
108 [47] (figure 1). The glacial species pump is a process in which allopatric speciation is
109 accelerated via the isolation of small populations in glacial refugia. It operates when the
110 expansion of glaciers makes large areas of a species’ range unavailable, but leaves isolated
111 pockets of suitable habitat (figure 1b; Hewitt 2000; Hewitt 2004; Holderegger and Thiel-
112 Egenter 2009; April et al. 2013). It can therefore be expected that glacial pump creates
113 assemblages composed of many species that originally emerged in allopatry (figure 1c) but
114 might have come into secondary contact more recently (figure 1e). The role of refugia in
115 promoting species persistence and in glacial vicariant speciation has been widely reported for
116 multiple extant European taxa, both animals and plants [36,37,e.g. 50–53].

117 After each glacial maximum, the retreat of glaciers opens up new, unoccupied habitat in both
118 terrestrial and aquatic ecosystems, which may lead early colonists to diversify *in situ* via
119 adaptive radiations [54] (figure 1b). Adaptive radiation takes place when an ancestor
120 diversifies ecologically and phenotypically, giving rise in a relatively short period of time, to
121 multiple species that are ecologically distinct [55]. Emblematic examples of adaptive
122 radiation during inter- or postglacial periods are associated the emergence of many lakes in
123 the Holarctic realm by the end of the last glacial period, providing opportunities for many
124 radiations of freshwater fish that occurred in parallel [33,34,47,56–59]. Such process would
125 be expected to generate sympatric or parapatric species by *in situ* cladogenetic speciation
126 [23]. In addition, postglacial expansion of populations also has the potential to bring together
127 lineages that had previously diverged in Pleistocene refugia [60]. Secondary contact of
128 lineages can also be important in adaptive radiation, either through causing ecological

character displacement in sympatry occurring in response to competition [55] (Schluter 2000, or through the occurrence of hybridization which may facilitate the onset of adaptive radiations upon colonization of new environments [61,62]. This ‘hybrid swarm origin’ hypothesis of adaptive radiation posits that functional genetic variation, which becomes enriched in hybrid zones, elevates the evolvability and response to natural selection in hybrid populations [61,63].

Despite its large information content for investigating hypothesis regarding biodiversity assembly and to understand biogeographic patterns in species distributions [64], few studies have made use of the species age distribution (SAD) [41,65,66], and fewer had investigated SAD from a multi-taxon perspective [8,67]. We conducted a comparative phylogenetic analysis to quantify SADs and the extent and type of endemism in aquatic and terrestrial ecosystems of the European Alps. Our work focuses on several taxonomic groups: amphipods, fish, butterflies and amphibians with nearly complete taxon sampling, as well as 15 nearly completely sampled representative genera of perennial flowering angiosperms plants (henceforth “plants”). We predicted that SADs support a scenario with dominance of the glacial species pump for the origin of endemism in terrestrial groups, with species dating to the Pleistocene; whereas SADs may indicate a prominent role of postglacial adaptive radiation for groups that depend on open water habitats, i.e., fish. This is because high altitude ranges in the terrestrial habitats became fragmented but not completely erased during glacial maxima, whereas year-round open water bodies as habitat for fish were entirely absent during glacial maxima, with a few exceptions at the edge of the southern Alps [68]. We also predicted that SADs of both amphipods and amphibians will resemble fully terrestrial taxa more than those of fish. Amphipods occupy both open and subterranean freshwater habitats, and some of the subterranean habitats persisted during the glacial maxima [69], and most amphibians require open water bodies only during spring and summer and are terrestrial for the remainder of the year, while some lack an aquatic life stage altogether [70]. Therefore, some amphibian species likely found Pleistocene refugia within the Alpine region during the glacial maxima [71,72]. Regarding non-endemic species, we did not expect large differences in age structure among taxa because non-endemics tend to be widespread, are probably mostly older and diversified on a wider geographical scale drive by processes that may have been decoupled from the climate dynamics of the Alps. Therefore, we predicted that non-endemic species are older on average than endemics. Because dispersal ability, distance between sink (new habitats that become available after glacial maxima) and sources of

colonization, historical stability and heterogeneity of habitat occupied vary between the taxonomic groups, the magnitude of the difference in species ages between non-endemic and endemic species will be taxon-dependent.

2. Methods

Our work focuses on the European Alps, following previous delimitations of European high mountain systems [73] but including the peripheral lowland areas, where most of the perialpine glacial lakes are located. To select the taxonomic groups to be included in this work, we looked for lineages with reliable distribution data, robust dated phylogenetic trees that include most of the diversity of the given lineage, and/or genetic data for most of the recognized species, so that we could estimate and calibrate phylogenetic trees where none existed. Then, we chose five major taxonomic groups to represent the terrestrial and aquatic alpine and pre-alpine biomes of the region: nearly all known regional species of amphipods, fishes, amphibians and butterflies, and 14 nearly completely sampled clades of flowering plants (*Homogyne*, *Petasites*, *Tussilago*, *Campanula*, *Jasione*, *Phyoplexis*, *Knautia*, *Androsace*, *Primula*, *Soldanella*, *Gentiana*, *Saxifraga*, *Carex* and *Festuca*). Only species native to the European Alps were considered. To assemble species age distributions (stem age, i.e., time since divergence from closest relative in million years, Ma), we combined published time-trees and our own estimates (electronic Supplementary Methods).

(a) Endemism and speciation mode

Species were considered endemic if they naturally occur only in the alpine and/or perialpine regions of the Alps. When a species was classified as endemic, we assigned its speciation mode to anagenetic speciation (Alpine species diverged from its non-Alpine sister species, but did not undergo *in situ* diversification) or cladogenetic speciation (Alpine species emerged through *in situ* diversification, with either the non-endemic sister being native to the region too, or the two or more sister species all being Alpine endemics), following Rosindell and Phillimore (2011) (electronic supplementary material, tables S1). To apply these definitions, we first identified the position of each of the endemic species within the phylogeny (electronic supplementary material, file S2). Then, if the endemic species was nested within a clade composed mainly of species that occur outside of the Alps, and its direct sister species occurred only outside the Alps, we assumed anagenetic speciation. If the species was nested within a group mostly of species native to the Alps, we assumed cladogenetic speciation.

(b) Comparisons of species age distributions

We performed multiple permutation tests on the distribution of age estimates assembled from all the species to identify differences in the age distributions among and within major clades. Here, we asked 1) whether age distributions differed between endemic and non-endemic species, overall and within each taxonomic group independently; 2) whether species age distributions differed among taxonomic groups; 3) whether non-endemic species age distributions differed among groups; and 4) whether endemic species age distributions differed among groups. When a difference was significant, if a significant result was found, we performed post-hoc pairwise permutations to identify which distribution and taxonomic group (or groups) were distinct from one another, using Bonferroni correction for each group of analyses. These analyses were performed using the function ‘oneway_test’ of the library ‘coin’ [74] in R 4.0.2 1106 [75] with 10,000 resamples with a distribution approximated via Monte Carlo resampling.

3. Results

A total of 617 species were included in our analyses: 39 amphipods, 124 fishes, 31 amphibians, 245 butterfly and 178 plant species (electronic supplementary material, table S1). Approximately half of all fish and amphipod species were found to be endemic to the European Alps (45.2 % and 49 %, respectively), whereas smaller fractions of 12 %, 13 % and 30 % of the butterflies, amphibians and plants, respectively, were found to be endemic (figure 2). Our analysis of speciation mode suggests that approximately half of the endemic amphipods and plants, and one third of butterfly species emerged by cladogenesis (53 %, 52 % and 36 %, respectively). Cladogenesis was also inferred to be the mode of speciation for the majority of the endemic amphibian and fish species (75 % and 98 %, respectively) (figure 2).

We found that 94 % of the extant species that now occupy the Alps, irrespective of endemism status, emerged over the past 15 million years (from the Middle Miocene) (figure 2). Endemics were overall younger than non-endemics (p-value < 0.0001), which was also true for each taxonomic group when analysed individually (electronic supplementary material, table S9). We found the species age distributions among non-endemic species to be similar between taxonomic groups (figure 2), with most of the species ages spanning the Late Pleistocene to Early Miocene (90 % fell between 0.3 and 20 Mya). The only exception was the comparison of plants vs. amphibians, as we found non-endemic amphibians older than

non-endemic plants ($p < 0.0001$; electronic supplementary material S10). SADs of endemic species were also similar among taxa (90 % fell between 0.25 and 8 Ma), except for fish, which are younger than any other group (90 % fell between 0.6 and 114 Ky; $p < 0.0001$; figure 2; electronic supplementary material S11).

We observed the following patterns in the SAD: i) endemics are younger than non-endemics in all taxonomic groups, and ii) there is a high degree of similarity among taxa in the age distributions when endemics and non-endemics are analysed separately. However, we found on one hand that while most endemic fish species arose in the Holocene through cladogenesis, very few endemic butterflies and no endemic species in the other groups arose in the Holocene. On the other hand, only very few endemic fish (all from the southern periphery of the Alps) date to the Pleistocene whereas most of the endemic butterflies and many other endemic species date do.

4. Discussion

Inferring past evolutionary process from the structure of current biodiversity is one of the goals of macroecology, macroevolution and biogeography. Species age distributions among regional biota is one informative aspect of this structure [64]. We combined estimates of species origination times with information on endemism and mode of speciation (cladogenetic vs. anagenetic) to investigate the dominant mechanisms of biodiversity assembly in different taxa and major biomes of the European Alps. We found that most of the species diversity in fish, amphipods, amphibians, butterflies and plants is relatively young, emerging at the beginning in the Middle Miocene, which coincided with the period of maximum geological uplift in the region that culminated with the formation of the Alps [76].

Our multi-taxon comparative analysis had the advantage of directly comparing independently evolving and ecologically distinct clades in the same region. It showed that speciation timing was dramatically different between terrestrial taxa and those aquatic taxa that require permanent open surface water (i.e., fish). While 80 % of the Alp's endemics in the terrestrial groups originated between the Late Miocene and Late Pleistocene, most endemic fish species arose only after the final retreat of the glaciers and re-establishment of permanent open water bodies in the formerly glaciated areas. Combined with the observation that the vast majority of endemic fish are products of cladogenetic speciation, this suggests that the assembly process of the fish fauna of the Alps is dominated by an interaction between colonization from outside the region and adaptive radiation during the last postglacial period, but similar

processes may have cyclically repeated themselves in previous interglacial periods, alternating with extinction during glacial maxima. In contrast, for the terrestrial groups, our results suggest that colonization from outside the region and the glacial species pump are the dominant mechanisms, as anagenetic speciation was more important in these taxa, and endemic richness assembled throughout the Pleistocene. Interestingly, we observed some postglacial speciation in butterflies, coinciding with the major mode in fish, but being dwarfed in butterflies by the much larger Pleistocene mode. We suggest that general, non-exclusive mechanisms underlay these contrasting patterns: 1) Quaternary climate fluctuations that accelerate allopatric speciation during cold stages but open up new ecological opportunities for adaptive radiation during interglacial periods; 2) variation among groups in their dispersal ability and associated rate of range expansion, and finally 3) the influence of variation in seasonal and inter-annual habitat stability in either constraining or promoting adaptive radiation. Below we discuss each of these mechanisms and how they may have affected diversification in the different taxa studied here.

(a) The role of Quaternary climate fluctuations

We suggest our finding that most endemic fish are of postglacial origin, while endemics in other groups arose in the Pleistocene or earlier, is explained by the different effects that the Quaternary climate oscillations had on freshwater versus terrestrial habitat [68]. Permanent open surface water habitats, as required by fishes, were absent in the glaciated parts of the Alps during the Last Glacial Maximum (LGM), because all lakes and river valleys were covered by thick glaciers. Therefore, the complete lack of endemic fish species older than 20,000 years on the northern and western flanks of the Alps is likely due to local extirpation of all fish populations across the region during the LGM. With the progressive retreat of glaciers, which achieved their modern configuration in the Late Holocene, fish would then have returned to the region from areas located in downstream sections of the large rivers, that were often far from the alpine region especially on the North face of the Alps [e.g. 77]. This Holocene recolonization by older widespread species explains the large fraction of old, widespread and non-endemic fish species in the northern and western Alpine region.

The first fish to colonize after the LGM were probably species adapted to cold water conditions, such as salmonids and sculpins, that would have lived nearby in the rivers of the Pleistocene tundra downstream of the Alpine glacier shield. Salmonids are indeed known for their remarkable colonizing ability and rapid establishment in postglacial freshwater habitats

[78]. These fish would have encountered ecological opportunities in the emerging large and deep lakes of the region and radiated into many distinct species as they adapted to the vacant niches associated with distinct lacustrine zones. This process likely generated the young endemic species, nowadays predominantly in three lineages, whitefish (*Coregonus* [79]), chars (*Salvelinus* [80]) and sculpins (*Cottus* [34]), that have rapidly radiated in perialpine lakes. The very few old, relic endemic fish species in the region that date to prior to the Holocene, are the lake herring *Alosa agone*, and two trouts of the genus *Salmo* (*S. carpio* and *S. sp.* ‘Blackspot’). These three species are endemic to lakes in Northern Italy and southern Switzerland, a region where probably not all lakes were fully covered by ice sheets during the LGM [68]. Therefore, these species likely originated during earlier interglacials, when southern perialpine lakes would have become extensive, and then found refugia during the LGM to persist to the present day [81]. That there are no young postglacial species among the non-endemic fish is perhaps due to insufficient time and connectivity between lakes to allow new species to arise in deep lakes elsewhere in Europe (e.g., in northern Germany and Scandinavia) in the Holocene prior to expanding their range into the Alps or vice-versa. It is important to mention that adaptive radiation of fish are far more frequent in deep lakes [82,83], while riverine adaptive radiations are rare (but see [84–86]).

Unlike for fish, our analysis revealed that all endemic (and all nonendemic) amphipods, the second fully aquatic taxon in our data, emerged during or before the Pleistocene. This could be because amphipods can persist in smaller water bodies than fish, and many species in this group are ice-associated, being able to survive under ice cover and in its immediate forefield [87]. Both factors would have allowed some species to persist in the region throughout the glaciations. Additionally, as suggested for other invertebrates [69], some species likely survived in subterranean refugia, such as caves or groundwaters, habitats occupied by many freshwater amphipod species today, notably species of *Niphargus*, the most species-rich group in the region [29,32].

We found that endemic species in the terrestrial groups are much older than endemic fish species. Pleistocene refugia are hypothesized for terrestrial taxa in many geographically restricted areas on nunataks, i.e, mountain peaks that have never been glaciated [88], or surrounding the Alps [41,89–92], such that extinctions during the glacial cycles did not wipe out the terrestrial fauna entirely. It is very likely that terrestrial organisms, particularly butterflies [41,93], plants [94–96] and amphibians [71,72] survived in glacial refugia in the Alps and at its periphery. Therefore, differential impacts of Quaternary climate fluctuations

and the resulting glaciations on different habitats and taxa go a long way helping to explain extant patterns of diversity and endemism in the region.

(b) Dispersal ability to explain postglacial radiation

Dispersal ability often negatively correlates with rapid niche evolution. The evolutionary response to local environmental and ecological conditions tends to be faster in taxa with limited dispersal, leading to faster niche shifts and higher rates of speciation and adaptive radiation [97,98]. Therefore, the intriguingly few cases of postglacial speciation in fully terrestrial species and amphibians could be related to the dispersal rates imposed by the environments they occupy. Terrestrial taxa experience, in general, less dispersal limitation than freshwater taxa. For example, many species have acquired adaptations for aerial dispersal, such as active flying in butterflies [99,100] or passive airborne propagation in many plants [101], allowing such taxa to disperse virtually in all directions. Conversely, freshwater-bound taxa need to navigate the dendritic landscapes of rivers and lakes to disperse, making it a lot more difficult to reach isolated habitat patches [102,103]. Given these limitations to dispersal for many freshwater taxa, postglacial dispersal may have happened at a much slower pace in fish than for most terrestrial taxa. Terrestrial species may, hence, have expanded their ranges faster after glacial retreat, also likely facilitated by the proximity of refugia to the Alps (including inner-alpine refugia), resulting in faster recolonization of the newly open landscape through long-distance dispersal and range expansion. Some recent studies have shown that many plant species rapidly and substantially expanded their range during the recent postglacial period [104–106]. Faster filling of emerging terrestrial habitats through range expansion left fewer opportunities/less time for the first colonists to undergo ecological speciation and adaptive radiation in response to ecological opportunity among terrestrial groups than among aquatic taxa. To test the relative importance of dispersal limitation versus other aquatic/terrestrial differences, future work could investigate mainly aquatic taxa with strong aerial dispersal abilities, such as Odonata and other insects that spent most of their life cycle in freshwater and have short but highly dispersive terrestrial adult phases.

(c) Seasonal and interannual environmental variation limiting ecological speciation

Habitat stability in the Postglacial Era may have been an additional factor explaining the larger number of Holocene speciation events in fish, but not in terrestrial groups in our study. Theory and models suggest that environmental fluctuations and stochasticity can reduce or

even inhibit ecological speciation in unstable habitats [107,108]. Rapid variation in environmental conditions, both seasonal and interannual, make adaptation difficult and ecological speciation nearly impossible.

Environmental conditions in terrestrial ecosystems are much more variable than aquatic ecosystems [109,110], especially large and deep lakes, both in terms of seasonal and year-to-year variation. For instance, whereas seasonal variations in solar irradiance, temperature and snow cover make the high mountain terrestrial habitat extremely seasonal, with large year-to-year variation in the onset and duration of seasons [111,112], they are nearly constant through the year in the deeper parts of lakes [113,114]. The longer growing and reproductive season, despite low productivity, and the much more stable environment in deep lakes may create increased opportunities for ecological speciation and adaptive radiation compared to the alpine terrestrial ecosystems.

In addition, despite their greater temporal stability, deep lakes also have much steeper environmental gradients because pressure, light and temperature all change much faster with depth in water than with elevation in the terrestrial realm. This unique property of water may explain the high frequency of ecological speciation in deep lakes, with sister-species being spatially very close to each other but occupying different water depths [58], as observed among East African cichlids [115] and Alpine whitefish [116].

5. Final considerations

We suggest that the formation of the unique biota of the European Alps was driven by interacting mechanisms: non-random Pleistocene survival, postglacial immigration, vicariant speciation during glacial maxima and adaptive radiation in the Postglacial. These interacting mechanisms left distinct imprints on the age structure of regional assemblages in different biomes and associated taxon groups. Historical factors (Quaternary climate fluctuations and Pleistocene refuge availability) impacted freshwater and terrestrial biomes in different ways, and contemporary ecological factors such as environmental stochasticity and dispersal limitations also vary between these biomes, shaping them very differently through ecological and evolutionary processes. *In situ* speciation and adaptive radiation were prominent in fish, but occurred mainly after the LGM, and only in deep lakes, likely due to the unavailability of suitable freshwater habitat during the LGM and the stable conditions within habitats after the LGM. Amphipods and all terrestrial clades have much older endemic species, perhaps because their ecology (i.e., cold-resistant and groundwater-dwelling in amphipods) and the

availability of Pleistocene refugia within the region allowed many species to survive the LGM. At the same time, none of the terrestrial groups generated many young postglacial species, likely because higher Pleistocene survival and faster postglacial niche filling through range expansion left fewer ecological opportunities and because larger seasonal variation in the terrestrial environment places constraints on ecological speciation.

Knowing the history of biodiversity formation is crucial to establish effective strategies of conservation [117]. For the Alps we show a high fraction of endemism in many groups, with endemic species having survived in some taxa and ecosystems through repeated glacial cycles, while those in others are due to prolific speciation after the retreat of the glaciers. These results improve our understanding of how the Alpine hotspot of species diversity and endemism emerged, and they reinforce that biodiversity in this region is fragile. Endemic species are often range-restricted, show limited population size and are hence much more vulnerable to climate change and other environmental changes than non-endemic species [118], and because of that, they are of high concern for conservation. Even a comparatively small and transient disturbance of an ecosystem can lead to the extinction of young species that evolved in adaptation to specific ecological conditions as has already been observed in the recent past for adaptive radiations of fish in Swiss lakes [116]. The sharp increase of extinction rates driven by human activity in the Anthropocene threatens the biodiversity of the European Alps, and especially that of endemic species [119]. Therefore, this region deserves greater attention to conserve both the regional biodiversity, as well as the evolutionary processes that gave rise to it and that are required to continue operating if biodiversity is to be maintained.

References

1. MacArthur RH, Wilson EO. 1963 An equilibrium theory of insular zoogeography. *Evolution* **17**, 373–387. (doi:10.2307/2407089)
2. MacArthur RH, Wilson EO. 1967 *The theory of island biogeography*. New Jersey: Princeton Univ. Press.
3. Rosindell J, Phillimore AB. 2011 A unified model of island biogeography sheds light on the zone of radiation. *Ecology Letters* **14**, 552–560. (doi:10.1111/j.1461-0248.2011.01617.x)
4. Warren BH *et al.* 2015 Islands as model systems in ecology and evolution: prospects fifty years after MacArthur-Wilson. *Ecology Letters* **18**, 200–217. (doi:10.1111/ele.12398)

- 420 5. Losos JB, Schluter D. 2000 Analysis of an evolutionary species–area relationship.
421 *Nature* **408**, 847–850. (doi:10.1038/35048558)
- 422 6. Rosenzweig ML. 2001 Loss of speciation rate will impoverish future diversity. *PNAS*
423 **98**, 5404–5410. (doi:10.1073/pnas.101092798)
- 424 7. Valente L *et al.* 2020 A simple dynamic model explains the diversity of island birds
425 worldwide. *Nature* **579**, 92–96. (doi:10.1038/s41586-020-2022-5)
- 426 8. Merckx VSFT *et al.* 2015 Evolution of endemism on a young tropical mountain. *Nature*
427 **524**, 347–350. (doi:10.1038/nature14949)
- 428 9. Willis JC. 1918 The age and area hypothesis. *Science* **47**, 626–628.
- 429 10. Gaston KJ. 1998 Species-range size distributions: products of speciation, extinction and
430 transformation. *Philosophical Transactions of the Royal Society of London. Series B:*
431 *Biological Sciences* **353**, 219–230. (doi:10.1098/rstb.1998.0204)
- 432 11. Ribas CC, Aleixo A, Nogueira ACR, Miyaki CY, Cracraft J. 2012 A
433 palaeobiogeographic model for biotic diversification within Amazonia over the past
434 three million years. *Proceedings of the Royal Society B: Biological Sciences* **279**, 681–
435 689. (doi:10.1098/rspb.2011.1120)
- 436 12. Brown JH. 1984 On the relationship between abundance and distribution of species. *The*
437 *American Naturalist* **124**, 255–279.
- 438 13. Svenning J-C, Skov F. 2004 Limited filling of the potential range in European tree
439 species. *Ecology Letters* **7**, 565–573. (doi:https://doi.org/10.1111/j.1461-
440 0248.2004.00614.x)
- 441 14. Smith BT *et al.* 2014 The drivers of tropical speciation. *Nature* **515**, 406–409.
442 (doi:10.1038/nature13687)
- 443 15. Louthan AM, Doak DF, Angert AL. 2015 Where and when do species interactions set
444 range limits? *Trends in Ecology & Evolution* **30**, 780–792.
445 (doi:10.1016/j.tree.2015.09.011)
- 446 16. Grandcolas P, Nattier R, Trewick S. 2014 Relict species: a relict concept? *Trends in*
447 *Ecology & Evolution* **29**, 655–663. (doi:10.1016/j.tree.2014.10.002)
- 448 17. Körner C. 2004 Mountain biodiversity, its causes and function. *ambi* **33**, 11–17.
449 (doi:10.1007/0044-7447-33.sp13.11)
- 450 18. Rahbek C *et al.* 2019 Building mountain biodiversity: Geological and evolutionary
451 processes. *Science* **365**, 1114–1119. (doi:10.1126/science.aax0151)
- 452 19. Rahbek C, Borregaard MK, Colwell RK, Dalsgaard B, Holt BG, Morueta-Holme N,
453 Nogues-Bravo D, Whittaker RJ, Fjeldså J. 2019 Humboldt’s enigma: What causes
454 global patterns of mountain biodiversity? *Science* **365**, 1108–1113.
455 (doi:10.1126/science.aax0149)

- 456 20. Hughes C, Eastwood R. 2006 Island radiation on a continental scale: Exceptional rates
457 of plant diversification after uplift of the Andes. *PNAS* **103**, 10334–10339.
458 (doi:10.1073/pnas.0601928103)
- 459 21. Hughes CE, Atchison GW. 2015 The ubiquity of alpine plant radiations: from the Andes
460 to the Hengduan Mountains. *New Phytologist* **207**, 275–282.
461 (doi:https://doi.org/10.1111/nph.13230)
- 462 22. Bertuzzo E, Carrara F, Mari L, Altermatt F, Rodriguez-Iturbe I, Rinaldo A. 2016
463 Geomorphic controls on elevational gradients of species richness. *PNAS* **113**, 1737–
464 1742.
- 465 23. Gillespie RG *et al.* 2020 Comparing adaptive radiations across space, time, and taxa.
466 *Journal of Heredity* **111**, 1–20. (doi:10.1093/jhered/esz064)
- 467 24. Comes HP, Kadereit JW. 2003 Spatial and temporal patterns in the evolution of the flora
468 of the European Alpine System. *TAXON* **52**, 451–462.
469 (doi:https://doi.org/10.2307/3647382)
- 470 25. Roquet C, Boucher FC, Thuiller W, Lavergne S. 2013 Replicated radiations of the
471 alpine genus *Androsace* (Primulaceae) driven by range expansion and convergent key
472 innovations. *Journal of Biogeography* **40**, 1874–1886.
473 (doi:https://doi.org/10.1111/jbi.12135)
- 474 26. Zhang L-B, Peter Comes H, W. Kadereit J. 2015 The temporal course of Quaternary
475 diversification in the European high mountain endemic *Primula* sect. *Auricula*
476 (Primulaceae). *International Journal of Plant Sciences* (doi:10.1086/380747)
- 477 27. Peña C, Witthauer H, Klečková I, Fric Z, Wahlberg N. 2015 Adaptive radiations in
478 butterflies: evolutionary history of the genus *Erebia* (Nymphalidae: Satyrinae). *Biol J*
479 *Linn Soc* **116**, 449–467. (doi:10.1111/bij.12597)
- 480 28. Pitteloud C *et al.* 2017 Climatic niche evolution is faster in sympatric than allopatric
481 lineages of the butterfly genus *Pyrgus*. *Proceedings of the Royal Society B: Biological*
482 *Sciences* **284**, 20170208. (doi:10.1098/rspb.2017.0208)
- 483 29. Altermatt F, Alther R, Fišer C, Jokela J, Konec M, Küry D, Mächler E, Stucki P,
484 Westram AM. 2014 Diversity and distribution of freshwater amphipod species in
485 Switzerland (Crustacea: Amphipoda). *PLOS ONE* **9**, e110328.
486 (doi:10.1371/journal.pone.0110328)
- 487 30. Fišer C, Delić T, Luštrik R, Zagmajster M, Altermatt F. 2019 Niches within a niche:
488 ecological differentiation of subterranean amphipods across Europe's interstitial waters.
489 *Ecography* **42**, 1212–1223. (doi:https://doi.org/10.1111/ecog.03983)
- 490 31. Zakšek V, Delić T, Fišer C, Jalžić B, Trontelj P. 2019 Emergence of sympatry in a
491 radiation of subterranean amphipods. *Journal of Biogeography* **46**, 657–669.
492 (doi:https://doi.org/10.1111/jbi.13514)
- 493 32. Borko Š, Trontelj P, Seehausen O, Moškrič A, Fišer C. 2021 A subterranean adaptive
494 radiation of amphipods in Europe. *Nat Commun* **12**, 3688. (doi:10.1038/s41467-021-
495 24023-w)

- 496 33. Hudson AG, Vonlanthen P, Seehausen O. 2011 Rapid parallel adaptive radiations from
497 a single hybridogenetic ancestral population. *Proc. R. Soc. B.* **278**, 58–66.
498 (doi:10.1098/rspb.2010.0925)
- 499 34. Lucek K, Keller I, Nolte AW, Seehausen O. 2018 Distinct colonization waves underlie
500 the diversification of the freshwater sculpin (*Cottus gobio*) in the Central European
501 Alpine region. *Journal of Evolutionary Biology* **31**, 1254–1267. (doi:10.1111/jeb.13339)
- 502 35. Fiebig M, Preusser F. 2008 Pleistocene glaciations of the northern Alpine Foreland.
503 *Geographica Helvetica* **63**, 145–150. (doi:10.5194/gh-63-145-2008)
- 504 36. Hewitt GM. 1999 Post-glacial re-colonization of European biota. *Biological Journal of*
505 *the Linnean Society* **68**, 87–112. (doi:10.1111/j.1095-8312.1999.tb01160.x)
- 506 37. Hewitt G. 2000 The genetic legacy of the Quaternary ice ages. *Nature* **405**, 907–913.
507 (doi:10.1038/35016000)
- 508 38. Hänfling B, Hellemans B, Volckaert F a. M, Carvalho GR. 2002 Late glacial history of
509 the cold-adapted freshwater fish *Cottus gobio*, revealed by microsatellites. *Molecular*
510 *Ecology* **11**, 1717–1729. (doi:https://doi.org/10.1046/j.1365-294X.2002.01563.x)
- 511 39. Stehlik I. 2003 Resistance or emigration? Response of alpine plants to the ice ages.
512 *TAXON* **52**, 499–510. (doi:https://doi.org/10.2307/3647448)
- 513 40. Neuenschwander S, Lurgiader CR, Ray N, Currat M, Vonlanthen P, Excoffier L. 2008
514 Colonization history of the Swiss Rhine basin by the bullhead (*Cottus gobio*): inference
515 under a Bayesian spatially explicit framework. *Molecular Ecology* **17**, 757–772.
516 (doi:10.1111/j.1365-294X.2007.03621.x)
- 517 41. Menchetti M *et al.* 2021 Two ways to be endemic. Alps and Apennines are different
518 functional refugia during climatic cycles. *Molecular Ecology* **30**, 1297–1310.
519 (doi:https://doi.org/10.1111/mec.15795)
- 520 42. Médail F, Diadema K. 2009 Glacial refugia influence plant diversity patterns in the
521 Mediterranean Basin. *Journal of Biogeography* **36**, 1333–1345.
522 (doi:https://doi.org/10.1111/j.1365-2699.2008.02051.x)
- 523 43. Holm SR, Svenning J-C. 2014 180,000 years of climate change in Europe: Avifaunal
524 responses and vegetation implications. *PLOS ONE* **9**, e94021.
525 (doi:10.1371/journal.pone.0094021)
- 526 44. Ordonez A, Svenning J-C. 2017 Consistent role of Quaternary climate change in
527 shaping current plant functional diversity patterns across European plant orders. *Sci Rep*
528 **7**, 42988. (doi:10.1038/srep42988)
- 529 45. Schoville SD, Roderick GK, Kavanaugh DH. 2012 Testing the ‘Pleistocene species
530 pump’ in alpine habitats: lineage diversification of flightless ground beetles (Coleoptera:
531 Carabidae: *Nebria*) in relation to altitudinal zonation. *Biological Journal of the Linnean*
532 *Society* **107**, 95–111. (doi:10.1111/j.1095-8312.2012.01911.x)

- 533 46. April J, Hanner RH, Dion-Côté A-M, Bernatchez L. 2013 Glacial cycles as an allopatric
534 speciation pump in north-eastern American freshwater fishes. *Molecular Ecology* **22**,
535 409–422. (doi:<https://doi.org/10.1111/mec.12116>)
- 536 47. Schluter D, Rambaut A, Clarke BC, Grant PR. 1996 Ecological speciation in postglacial
537 fishes. *Philosophical Transactions of the Royal Society of London. Series B: Biological*
538 *Sciences* **351**, 807–814. (doi:10.1098/rstb.1996.0075)
- 539 48. Hewitt GM. 2004 Genetic consequences of climatic oscillations in the Quaternary.
540 *Philosophical Transactions of the Royal Society of London. Series B: Biological*
541 *Sciences* **359**, 183–195. (doi:10.1098/rstb.2003.1388)
- 542 49. Holderegger R, Thiel-Egenter C. 2009 A discussion of different types of glacial refugia
543 used in mountain biogeography and phylogeography. *Journal of Biogeography* **36**, 476–
544 480. (doi:<https://doi.org/10.1111/j.1365-2699.2008.02027.x>)
- 545 50. Bermingham E, Rohwer S, Freeman S, Wood C. 1992 Vicariance biogeography in the
546 Pleistocene and speciation in North American wood warblers: a test of Mengel’s model.
547 *PNAS* **89**, 6624–6628.
- 548 51. Culling MA, Janko K, Boron A, Vasil’ev VP, Côté IM, Hewitt GM. 2006 European
549 colonization by the spined loach (*Cobitis taenia*) from Ponto-Caspian refugia based on
550 mitochondrial DNA variation. *Molecular Ecology* **15**, 173–190.
551 (doi:<https://doi.org/10.1111/j.1365-294X.2005.02790.x>)
- 552 52. Provan J, Bennett KD. 2008 Phylogeographic insights into cryptic glacial refugia.
553 *Trends in Ecology & Evolution* **23**, 564–571. (doi:10.1016/j.tree.2008.06.010)
- 554 53. Wallis GP, Waters JM, Upton P, Craw D. 2016 Transverse alpine speciation driven by
555 glaciation. *Trends in Ecology & Evolution* **31**, 916–926.
556 (doi:10.1016/j.tree.2016.08.009)
- 557 54. Stroud JT, Losos JB. 2016 Ecological opportunity and adaptive radiation. *Annu. Rev.*
558 *Ecol. Evol. Syst* **47**, 507–532.
- 559 55. Schluter D. 2000 *The ecology of adaptive radiation*. OUP Oxford.
- 560 56. Bernatchez L, Chouinard A, Lu G. 1999 Integrating molecular genetics and ecology in
561 studies of adaptive radiation: whitefish, *Coregonus* sp., as a case study. *Biological*
562 *Journal of the Linnean Society* **68**, 173–194. (doi:10.1111/j.1095-8312.1999.tb01165.x)
- 563 57. Bolnick DI, Lau OL. 2008 Predictable patterns of disruptive selection in stickleback in
564 postglacial lakes. *The American Naturalist* **172**, 1–11. (doi:10.1086/587805)
- 565 58. Seehausen O, Wagner C. 2014 Speciation in freshwater fishes. *Annual review of*
566 *ecology, evolution, and systematics* **45**, 621–651. (doi:10.1146/annurev-ecolsys-120213-
567 091818)
- 568 59. Doenz CJ, Krähenbühl AK, Walker J, Seehausen O, Brodersen J. 2019 Ecological
569 opportunity shapes a large Arctic charr species radiation. *Proceedings of the Royal*
570 *Society B: Biological Sciences* **286**, 20191992. (doi:10.1098/rspb.2019.1992)

- 571 60. Kangas V-M, Kvist L, Kholodova M, Nygrén T, Danilov P, Panchenko D, Fraimout A,
572 Aspi J. 2015 Evidence of post-glacial secondary contact and subsequent anthropogenic
573 influence on the genetic composition of Fennoscandian moose (*Alces alces*). *Journal of*
574 *Biogeography* **42**, 2197–2208. (doi:<https://doi.org/10.1111/jbi.12582>)
- 575 61. Seehausen O. 2004 Hybridization and adaptive radiation. *Trends in Ecology &*
576 *Evolution* **19**, 198–207. (doi:10.1016/j.tree.2004.01.003)
- 577 62. Meier JJ, Marques DA, Mwaiko S, Wagner CE, Excoffier L, Seehausen O. 2017
578 Ancient hybridization fuels rapid cichlid fish adaptive radiations. *Nat Commun* **8**,
579 14363. (doi:10.1038/ncomms14363)
- 580 63. Seehausen O. 2013 Conditions when hybridization might predispose populations for
581 adaptive radiation. *J Evol Biol* **26**, 279–281. (doi:10.1111/jeb.12026)
- 582 64. Goldberg EE, Roy K, Lande R, Jablonski D. 2005 Diversity, endemism, and age
583 distributions in macroevolutionary sources and sinks. *The American Naturalist* **165**,
584 623–633. (doi:10.1086/430012)
- 585 65. Pellissier L *et al.* 2014 Quaternary coral reef refugia preserved fish diversity. *Science*
586 **344**, 1016–1019. (doi:10.1126/science.1249853)
- 587 66. Ding W-N, Ree RH, Spicer RA, Xing Y-W. 2020 Ancient orogenic and monsoon-
588 driven assembly of the world's richest temperate alpine flora. *Science* **369**, 578–581.
589 (doi:10.1126/science.abb4484)
- 590 67. Rull V. 2008 Speciation timing and neotropical biodiversity: the Tertiary–Quaternary
591 debate in the light of molecular phylogenetic evidence. *Molecular Ecology* **17**, 2722–
592 2729. (doi:<https://doi.org/10.1111/j.1365-294X.2008.03789.x>)
- 593 68. Seguinot J, Ivy-Ochs S, Jouvet G, Huss M, Funk M, Preusser F. 2018 Modelling last
594 glacial cycle ice dynamics in the Alps. *The Cryosphere* **12**, 3265–3285.
595 (doi:<https://doi.org/10.5194/tc-12-3265-2018>)
- 596 69. Christian E. 2002 Distribution patterns of cavernicolous Collembola in Austria.
597 *Pedobiologia* **46**, 261–266. (doi:10.1078/0031-4056-00132)
- 598 70. Wells KD. 2010 *The ecology and behavior of amphibians*. University of Chicago Press.
- 599 71. Vörös J, Mikulíček P, Major Á, Recuero E, Arntzen JW. 2016 Phylogeographic analysis
600 reveals northerly refugia for the riverine amphibian *Triturus dobrogicus* (Caudata:
601 Salamandridae). *Biological Journal of the Linnean Society* **119**, 974–991.
602 (doi:10.1111/bij.12866)
- 603 72. Lucati F *et al.* 2020 Multiple glacial refugia and contemporary dispersal shape the
604 genetic structure of an endemic amphibian from the Pyrenees. *Molecular Ecology* **29**,
605 2904–2921. (doi:10.1111/mec.15521)
- 606 73. Schmitt T. 2017 Molecular biogeography of the high mountain systems of Europe: An
607 overview. In *High Mountain Conservation in a Changing World* (eds J Catalan, JM
608 Ninot, MM Aniz), pp. 63–74. Cham: Springer International Publishing.
609 (doi:10.1007/978-3-319-55982-7_3)

- 610 74. Hothorn T, Hornik K, van de Wiel MA, Zeileis A. 2008 Implementing a class of
611 permutation Tests: The coin package. *J. Stat. Softw.* **28**, 1–23.
- 612 75. R Core Team. 2020 *R: A language and environment for statistical computing*. Vienna: R
613 Foundation for Statistical Computing. See <https://www.R-project.org/>.
- 614 76. Ager DV. 1975 The geological evolution of Europe. *Proceedings of the Geologists’*
615 *Association* **86**, 127–154. (doi:10.1016/S0016-7878(75)80097-6)
- 616 77. Seifertová M, Bryja J, Vyskočilová M, Martínková N, Šimková A. 2012 Multiple
617 Pleistocene refugia and post-glacial colonization in the European chub (*Squalius*
618 *cephalus*) revealed by combined use of nuclear and mitochondrial markers. *Journal of*
619 *Biogeography* **39**, 1024–1040. (doi:<https://doi.org/10.1111/j.1365-2699.2011.02661.x>)
- 620 78. Klemetsen A, Amundsen P-A, Dempson JB, Jonsson B, Jonsson N, O’Connell MF,
621 Mortensen E. 2003 Atlantic salmon *Salmo salar* L., brown trout *Salmo trutta* L. and
622 Arctic charr *Salvelinus alpinus* (L.): a review of aspects of their life histories. *Ecology of*
623 *Freshwater Fish* **12**, 1–59. (doi:10.1034/j.1600-0633.2003.00010.x)
- 624 79. Hudson AG, Lundsgaard Hansen B, Lucek K, Vonlanthen P, Seehausen O. 2017
625 Managing cryptic biodiversity: Fine-scale intralacustrine speciation along a benthic
626 gradient in Alpine whitefish (*Coregonus* spp.). *Evolutionary Applications* **10**, 251–266.
627 (doi:<https://doi.org/10.1111/eva.12446>)
- 628 80. Freyhof J, Kottelat M. 2007 *Handbook of European freshwater fishes*. See
629 <https://portals.iucn.org/library/node/9068>.
- 630 81. Splendiani A, Ruggeri P, Giovannotti M, Pesaresi S, Occhipinti G, Fioravanti T,
631 Lorenzoni M, Nisi Cerioni P, Caputo Barucchi V. 2016 Alien brown trout invasion of
632 the Italian peninsula: the role of geological, climate and anthropogenic factors. *Biol*
633 *Invasions* **18**, 2029–2044. (doi:10.1007/s10530-016-1149-7)
- 634 82. Vonlanthen P, Roy D, Hudson AG, Largiadèr CR, Bittner D, Seehausen O. 2009
635 Divergence along a steep ecological gradient in lake whitefish (*Coregonus* sp.). *Journal*
636 *of Evolutionary Biology* **22**, 498–514. (doi:[https://doi.org/10.1111/j.1420-](https://doi.org/10.1111/j.1420-9101.2008.01670.x)
637 [9101.2008.01670.x](https://doi.org/10.1111/j.1420-9101.2008.01670.x))
- 638 83. McGee MD *et al.* 2020 The ecological and genomic basis of explosive adaptive
639 radiation. *Nature* **586**, 75–79. (doi:10.1038/s41586-020-2652-7)
- 640 84. Levin BA *et al.* 2019 Phylogeny, phylogeography and hybridization of Caucasian
641 barbels of the genus *Barbus* (Actinopterygii, Cyprinidae). *Molecular Phylogenetics and*
642 *Evolution* **135**, 31–44. (doi:10.1016/j.ympev.2019.02.025)
- 643 85. Levin BA, Simonov E, Dgebuadze YY, Levina M, Golubtsov AS. 2020 In the rivers:
644 Multiple adaptive radiations of cyprinid fishes (*Labeobarbus*) in Ethiopian Highlands.
645 *Sci Rep* **10**, 7192. (doi:10.1038/s41598-020-64350-4)
- 646 86. Levin B, Simonov E, Franchini P, Mugue N, Golubtsov A, Meyer A. 2021 Adaptive
647 radiation and burst speciation of hillstream cyprinid fish *Garra* in African river. *bioRxiv*
648 (doi:<https://doi.org/10.1101/2021.05.04.442598>)

- 649 87. Vader W, Tandberg AHS. 2019 Gammarid amphipods (Crustacea) in Norway, with a
650 key to the species. *Fauna Norvegica* (doi:10.5324/fn.v39i0.2873)
- 651 88. Nordal I. 1987 Tabula rasa after all? Botanical evidence for ice-free refugia in
652 Scandinavia reviewed. *Journal of Biogeography* **14**, 377–388. (doi:10.2307/2844945)
- 653 89. Taberlet P, Fumagalli L, Wust-Saucy A-G, Cosson J-F. 1998 Comparative
654 phylogeography and postglacial colonization routes in Europe. *Molecular Ecology* **7**,
655 453–464. (doi:https://doi.org/10.1046/j.1365-294x.1998.00289.x)
- 656 90. Schmitt T. 2009 Biogeographical and evolutionary importance of the European high
657 mountain systems. *Frontiers in Zoology* **6**, 9. (doi:10.1186/1742-9994-6-9)
- 658 91. Saladin B, Pellissier L, Graham CH, Nobis MP, Salamin N, Zimmermann NE. 2020
659 Rapid climate change results in long-lasting spatial homogenization of phylogenetic
660 diversity. *Nat Commun* **11**, 4663. (doi:10.1038/s41467-020-18343-6)
- 661 92. Pan D, Hülber K, Willner W, Schneeweiss GM. 2020 An explicit test of Pleistocene
662 survival in peripheral versus nunatak refugia in two high mountain plant species.
663 *Molecular Ecology* **29**, 172–183. (doi:10.1111/mec.15316)
- 664 93. Schmitt T, Louy D, Zimmermann E, Habel JC. 2016 Species radiation in the Alps:
665 multiple range shifts caused diversification in Ringlet butterflies in the European high
666 mountains. *Org Divers Evol* **16**, 791–808. (doi:10.1007/s13127-016-0282-6)
- 667 94. Willis KJ. 1996 Where did all the flowers go? The fate of temperate European flora
668 during glacial periods. *Endeavour* **20**, 110–114. (doi:10.1016/0160-9327(96)10019-3)
- 669 95. Schönswetter P, Stehlik I, Holderegger R, Tribsch A. 2005 Molecular evidence for
670 glacial refugia of mountain plants in the European Alps. *Molecular Ecology* **14**, 3547–
671 3555. (doi:https://doi.org/10.1111/j.1365-294X.2005.02683.x)
- 672 96. Szövényi P, Arroyo K, Guggisberg A, Conti E. 2009 Effects of Pleistocene glaciations
673 on the genetic structure of *Saxifraga florulenta* (Saxifragaceae), a rare endemic of the
674 Maritime Alps. *TAXON* **58**, 532–543. (doi:https://doi.org/10.1002/tax.582017)
- 675 97. Patterson TB, Givnish TJ. 2004 Geographic cohesion, chromosomal evolution, parallel
676 adaptive radiations, and consequent floral adaptations in *Calochortus* (Calochortaceae):
677 evidence from a cpDNA phylogeny. *New Phytologist* **161**, 253–264.
678 (doi:https://doi.org/10.1046/j.1469-8137.2003.00951.x)
- 679 98. Price JP, Wagner WL. 2004 Speciation in Hawaiian Angiosperm lineages: cause,
680 consequence, and mode. *evol* **58**, 2185–2200. (doi:10.1554/03-498)
- 681 99. Stevens VM, Turlure C, Baguette M. 2010 A meta-analysis of dispersal in butterflies.
682 *Biological Reviews* **85**, 625–642. (doi:10.1111/j.1469-185X.2009.00119.x)
- 683 100. Stevens VM, Trochet A, Dyck HV, Clobert J, Baguette M. 2012 How is dispersal
684 integrated in life histories: A quantitative analysis using butterflies. *Ecology Letters* **15**,
685 74–86. (doi:10.1111/j.1461-0248.2011.01709.x)

- 686 101. Frei T. 1997 Pollen distribution at high elevation in Switzerland: Evidence for medium
687 range transport. *Grana* **36**, 34–38. (doi:10.1080/00173139709362587)
- 688 102. Brown BL, Swan CM. 2010 Dendritic network structure constrains metacommunity
689 properties in riverine ecosystems. *Journal of Animal Ecology* **79**, 571–580.
690 (doi:10.1111/j.1365-2656.2010.01668.x)
- 691 103. Altermatt F. 2013 Diversity in riverine metacommunities: a network perspective. *Aquat*
692 *Ecol* **47**, 365–377. (doi:10.1007/s10452-013-9450-3)
- 693 104. Magri D *et al.* 2006 A new scenario for the Quaternary history of European beech
694 populations: palaeobotanical evidence and genetic consequences. *New Phytologist* **171**,
695 199–221. (doi:10.1111/j.1469-8137.2006.01740.x)
- 696 105. Tinner W, Lotter AF. 2006 Holocene expansions of *Fagus sylvatica* and *Abies alba* in
697 Central Europe: where are we after eight decades of debate? *Quaternary Science*
698 *Reviews* **25**, 526–549. (doi:10.1016/j.quascirev.2005.03.017)
- 699 106. Magri D. 2008 Patterns of post-glacial spread and the extent of glacial refugia of
700 European beech (*Fagus sylvatica*). *Journal of Biogeography* **35**, 450–463.
701 (doi:10.1111/j.1365-2699.2007.01803.x)
- 702 107. Johansson J, Ripa J. 2006 Will sympatric speciation fail due to stochastic competitive
703 exclusion? *The American Naturalist* **168**, 572–578. (doi:10.1086/507996)
- 704 108. Ripa J, Dieckmann U. 2013 Mutant invasions and adaptive dynamics in variable
705 environments. *Evolution* **67**, 1279–1290. (doi:https://doi.org/10.1111/evo.12046)
- 706 109. Halley JM. 2005 Comparing aquatic and terrestrial variability: at what scale do
707 ecologists communicate? *Marine Ecology Progress Series* **304**, 274–280.
- 708 110. Steele JH, Brink KH, Scott BE. 2019 Comparison of marine and terrestrial ecosystems:
709 suggestions of an evolutionary perspective influenced by environmental variation. *ICES*
710 *Journal of Marine Science* **76**, 50–59. (doi:10.1093/icesjms/fsy149)
- 711 111. Häder D-P, Barnes PW. 2019 Comparing the impacts of climate change on the
712 responses and linkages between terrestrial and aquatic ecosystems. *Science of The Total*
713 *Environment* **682**, 239–246. (doi:10.1016/j.scitotenv.2019.05.024)
- 714 112. Liu J, Liu J, Linderholm HW, Chen D, Yu Q, Wu D, Haginoya S. 2012 Observation and
715 calculation of the solar radiation on the Tibetan Plateau. *Energy Conversion and*
716 *Management* **57**, 23–32. (doi:10.1016/j.enconman.2011.12.007)
- 717 113. Fenocchi A, Rogora M, Sibilla S, Dresti C. 2017 Relevance of inflows on the
718 thermodynamic structure and on the modeling of a deep subalpine lake (Lake Maggiore,
719 Northern Italy/Southern Switzerland). *Limnologia* **63**, 42–56.
720 (doi:10.1016/j.limno.2017.01.006)
- 721 114. Arai T. 1981 Attenuation of Incident Solar Radiation in Lake Water. *Jap. J. Limnol.* **42**,
722 92–99. (doi:10.3739/rikusui.42.92)

115. Wagner CE, Harmon LJ, Seehausen O. 2014 Cichlid species-area relationships are shaped by adaptive radiations that scale with area. *Ecology Letters* **17**, 583–592. (doi:<https://doi.org/10.1111/ele.12260>)
116. Vonlanthen P *et al.* 2012 Eutrophication causes speciation reversal in whitefish adaptive radiations. *Nature* **482**, 357–362. (doi:[10.1038/nature10824](https://doi.org/10.1038/nature10824))
117. Posadas P, Esquivel DRM, Crisci JV. 2001 Using phylogenetic diversity measures to set priorities in conservation: An example from southern South America. *Conservation Biology* **15**, 1325–1334. (doi:[10.1111/j.1523-1739.2001.99404.x](https://doi.org/10.1111/j.1523-1739.2001.99404.x))
118. Manes S *et al.* 2021 Endemism increases species' climate change risk in areas of global biodiversity importance. *Biological Conservation* **257**, 109070. (doi:[10.1016/j.biocon.2021.109070](https://doi.org/10.1016/j.biocon.2021.109070))
119. Johnson CN, Balmford A, Brook BW, Buettel JC, Galetti M, Guangchun L, Wilmschurst JM. 2017 Biodiversity losses and conservation responses in the Anthropocene. *Science* **356**, 270–275. (doi:[10.1126/science.aam9317](https://doi.org/10.1126/science.aam9317))

Acknowledgements

We would like to thank Salome Mwaiko and Rosi Siber for laboratory and GIS support, respectively, and Timothy Alexander for his work in the execution of the *Projet Lac*. Info Fauna provided distribution data of Swiss fishes. For sharing published phylogenetic trees, we thank Natalia Tkach, Adrien Favre, Józef Mitka, Carmen Benitez-Benitez, Markus Dillenberger, Andrew Cowl and Bozko Frajman. We are also grateful to Kay Lucek, Hanna ten Brink and Catalina Chaparro for thoughtful and constructive discussion. Funding from the ETH Board through the Blue-Green Biodiversity (BGB) Initiative (BGB 2020) is acknowledged. ŠB was supported by Slovenian Research Agency through funding programme P1-0184, project J1-2464 and PhD grant.

Figure captions

Figure 1. Evolutionary and ecological history of a hypothetical biodiversity assembly in an alpine-like system. A) Biodiversity in a preglacial phase. B) Early glacial phase: glacial periods erase freshwater habitats and fragment the terrestrial biome. Some populations survive in refugia and C) can diverge into distinct species through allopatric speciation. D) The retreat of glaciers opens up new, unoccupied habitats offering ecological opportunities for colonizers. E) Some colonizers undergo adaptive radiation and niche space is filled up again.

Figure 2. Species age distribution (SAD) of endemic and non-endemic species of, from top to bottom, amphibians, amphipods, butterflies, fish and flowering plants. Pie charts show the proportion of endemic and non-endemic species as well as the proportion of endemic species that have emerged through cladogenetic or anagenetic speciation.