dynamAedes: a unified modelling framework for invasive *Aedes* mosquitoes

Daniele Da Re^{1,*}, Wim Van Bortel², Friederike Reuss^{3,4},
Ruth Müller², Sebastien Boyer⁵, Fabrizio Montarsi⁶,
Silvia Ciocchetta⁷, Daniele Arnoldi⁸, Giovanni Marini⁸,
Annapaola Rizzoli⁸, Gregory L'Amber⁹,
Guillaume Lacour¹⁰, Constantianus J.M. Koenraadt¹¹,
Sophie O. Vanwambeke^{1,†}, Matteo Marcantonio^{12,†}

21st December 2021

¹ Georges Lemaître Center for Earth and Climate Research, Earth and Life Institute, UCLouvain, Louvain-la-Neuve, Belgium

² Unit Entomology and the Outbreak Research Team, Tropical Medicine Institute, Antwerp, Belgium

³ Senckenberg Biodiversity and Climate Research Centre, Frankfurt am Main, Germany

⁴ Institute of Occupational, Social and Environmental Medicine, Goethe University, Frankfurt am Main, Germany

⁵ Medical and Veterinary Entomology Unit, Institute Pasteur du Cambodge, Cambodia

⁶ Laboratory of Parasitology, National reference centre/OIE collaborating centre for diseases at the animal-human interface, Istituto Zooprofilattico Sperimentale delle Venezie, Legnaro, Italy

⁷ The University of Queensland, School of Veterinary Science, Gatton, Australia

⁸ Department of Biodiversity and Molecular Ecology, Research and Innovation Centre, Fondazione Edmund Mach, San Michele all'Adige, Trento, Italy

⁹ EID Méditerranée, Direction Technique, Montpellier, France

¹⁰ Altopictus, Pérols, France

¹¹ Wageningen University & Research, Department of Plant Sciences, Laboratory of Entomology, Wageningen, The Netherlands.

¹² Evolutionary Ecology and Genetics Group, Earth and Life Institute, UCLouvain, Louvain-la-Neuve, Belgium

* corresponding author: danielle.dare@uclouvain.be

† SOV and MM equally supervised the project

Abstract

1. Mosquito species belonging to the genus *Aedes* have attracted the interest of scientists and public health officers for their invasive species traits and efficient capacity of transmitting viruses affecting humans. Some of these species were brought outside their native range by human activities such as trade and tourism, and colonised new regions thanks to a unique combination of eco-physiological traits.
2. Considering mosquito physiological and behavioural traits to understand and predict the spatial and temporal population dynamics is thus a crucial step to develop strategies to mitigate the local densities of invasive *Aedes* populations.
3. Here, we synthesised the life cycle of four invasive *Aedes* species (*Ae. aegypti*, *Ae. albopictus*, *Ae. japonicus* and *Ae. koreicus*) in a single multi-scale stochastic modelling framework which we coded in the R package `dynamAedes`. We designed a stage-based and time-discrete stochastic model driven by temperature, photo-period and inter-specific larval competition that can be applied to three different spatial scales: punctual, local and regional. These spatial scales consider different degrees of spatial complexity and data availability, by accounting for both active and passive dispersal of mosquito species as well as for the heterogeneity of the input temperature data.
4. Our overarching aim was to provide a flexible, open-source and user-friendly tool rooted in the most updated knowledge on species biology which could be applied to the management of invasive *Aedes* populations as well as for more theoretical ecological inquiries.

Keywords: Biological invasions; Invasion ecology; Process-based models; Spatial epidemiology; Dispersal; Vector-borne pathogens

1 Introduction

Some mosquito species within the *Aedes* taxon have a unique combination of biological traits such as: 1) efficient transmission of viruses debilitating for humans and animals (Gratz, 2004; Hurk et al., 2011; Souza-Neto et al., 2019), 2) eco-physiological plasticity that allows for rapid adaptation (Kramer et al., 2021) and exploitation of novel environments created by humans (McBride et al., 2014a), 3) egg stage with high resistance to dry and cold conditions which facilitate displacements over long ecological and geographical distances (Thomas et al., 2012; Versteirt et al., 2012; Kaufman and Fonseca, 2014). Some of these species were accidentally brought outside their native areas by human activities and colonised new regions thanks to a unique combination of eco-physiological traits. These mosquitoes, often referred as "*Aedes* invasive mosquitoes" (AIM), have attracted the interest of scientists and public health officers and much effort has been done to unravel their physiological and behavioural traits. Among these species, *Ae. aegypti*, *Ae. albopictus*, *Ae. japonicus* and *Ae. koreicus* showed a rapid expansion of their geographical range, with the first two species often causing an important burden on public health. As a consequence, large experimental and observational datasets on the relationship between water or air temperature and physiological parameters have been collected and used to develop mechanistic models that reproduce the basic life cycle of these four species (e.g., for *Ae. aegypti* Focks et al., 1993a,b; Otero et al., 2006; Da Re et al., 2021; Caldwell et al., 2021; for *Ae. albopictus* Tran et al., 2013; Erguler et al., 2016; Metelmann et al., 2019; Pasquali et al., 2020; Tran et al., 2020; for *Ae. japonicus* Wieser et al., 2019; for *Ae. koreicus* Marini et al., 2019b). The inclusion of such functions, which describe physiological and developmental rates, into modelling frameworks allow for more reliable model extrapolations, as the chances of biological unrealistic outcome may be lower compared to pure correlative model approaches (Kearney, 2006; Kearney and Porter, 2009). Comparisons between modelled and observed population trends showed that such mechanistic models can be used, for example, to understand population dynamics in space and time, and thus can enhance the pest control strategies against AIM (Baldacchino et al., 2015).

Models targeting AIM developed so far aimed to simulate the population dynamics of only one species at the time and for a single qualitative (i.e. "individual", "container" or "household") or quantitative (cell in a lattice grid) spatial scale. Moreover, only a few of these models have been made readily operational, for example by organising them in open-access, user-friendly software or libraries with sufficient documentation for practical applications (Stallman, 1985). SkeeterBuster, a container-level population dynamical model for *Ae. aegypti*, has been the first agent-based model for mosquitoes made available as a free (but not open-source) software (Magori et al., 2009). Concerning *Ae. albopictus*, Erguler et al. (2016) made available a model developed as a Python library that was after-

wards wrapped into the R package `albopictus`. More recently, the European Centre for Disease Control (ECDC) has provided a free and open-source adaptation of a model initially developed by Tran et al. (2013), making it accessible via the R Shiny application `AedesRisk`¹. Similarly, two generic age and stage-structured discrete-time population dynamics models, also applicable to mosquitoes, were proposed in the last few years: `stagePop` (Kettle and Nutter 2015; applied to *Ae. japonicus* in Wieser et al. 2019) and `sPop` (Erguler, 2018). Despite the current availability of models applicable to invasive *Aedes*, none of them can be directly generalised while retaining biological credibility, e.g. species-specific models work only for a single species whereas generic models may oversimplify the life cycle structure or are not equipped with species-specific physiological parameters. Hence, if users decide to use generic models, they need to screen the scientific literature, filter and manipulate experimental data (often scarce and non-standardised) to inform models on the species of interest. Moreover, often models do not consider mosquito dispersal or even completely lack spatial structure.

Here, we synthesised the life cycle of four AIM species: *Ae. aegypti*, *Ae. albopictus*, *Ae. japonicus* and *Ae. koreicus* in a single modelling framework, which we coded in the R package `dynamAedes`. We designed a stage-based and time discrete stochastic model informed by temperature and photoperiod that can be applied to three different spatial scales: punctual, local and regional. These spatial scales were thought to meet different degrees of spatial complexity and data availability, by accounting for both active and human-mediated passive dispersal of the modelled mosquito species as well as for the heterogeneity of temperature data. Our overarching aim was to provide a flexible and open-source tool which could be used for applications related to the management of AIM populations but also for more theoretical ecological inquiries. We described and assessed the model using observational mosquito data and then showed how to use the R package with coding examples and relevant case studies.

2 Materials and methods

2.1 A summary of invasive *Aedes* species ecology

2.1.1 *Aedes aegypti*

Aedes (Stegomyia) aegypti (Linnaeus, 1762), commonly referred to as the "yellow fever mosquito", was progressively brought outside sub-Saharan Africa by human trade. It was first introduced in the Americas during the 16th century and afterwards to tropical and temperate regions of Asia and Oceania (Powell et al., 2013). Its invasion was likely favoured

¹AedesRisk v1.0: <https://shinyapps.ecdc.europa.eu/shiny/AedesRisk/>

by a series of functional traits, such as egg desiccation-resistance, that allows them to withstand dry conditions for months, and egg moderate resistance to cold temperatures (Juliano et al., 2002; Thomas et al., 2012; Kramer et al., 2020). *Aedes aegypti* efficiently transmit several viruses to humans, including yellow fever, dengue, chikungunya, Zika, Rift Valley, Mayaro and eastern equine encephalitis viruses (Leta et al., 2018; Näslund et al., 2021; da Silva Neves et al., 2021). This is the result of several eco-evolutionary traits that are specific to the species: i) high preference for human hosts (anthropophily), which is channelled by genetic traits linked to behavioural and physiological evolutionary advantages (Harrington et al., 2001; McBride et al., 2014b), ii) exploitation of human dwellings and architectures as shelter, hide and resting indoor sites (endophily) to avoid unfavourable environmental conditions (Dzul-Manzanilla et al., 2017; Gloria-Soria et al., 2018), and iii) selection of artificial containers for oviposition and subsequent larval development (eusynantrophy; Christophers, 1960).

2.1.2 *Aedes albopictus*

Aedes (Stegomyia) albopictus (Skuse, 1895), commonly referred as the "Asian tiger mosquito", is native of tropical and subtropical regions of Southern-East Asia and Indonesia (Watson, 1967; Hawley, 1988). It is a competent vector of several viruses, including dengue, chikungunya, Zika, West Nile, eastern equine encephalitis and La Crosse viruses (Koch et al., 2016; McKenzie et al., 2019; Takken and van den Berg, 2019) and it was implicated as the vector species causing local transmission of dengue, chikungunya or Zika virus, even at temperate latitudes outside its native distributional range (Effler et al., 2005; Rezza et al., 2007; Delatte et al., 2008; Venturi et al., 2017; Brady and Hay, 2019; Giron et al., 2019; Barzon et al., 2021). This species is a more opportunistic feeder compared to *Ae. aegypti* (Cebrián-Camisón et al., 2020). It prefers sub-urban habitats with the presence of vegetation, dispersing bites among several species, a behaviour that might decrease the probability of pathogen transmission to humans (Turell et al., 1994; Lounibos and Kramer, 2016). Populations of this species located at temperate latitudes show: i) an adaptation to temperate climatic conditions (Marini et al., 2020) and ii) a stronger tendency to laying diapausing eggs at the end of summer (Hawley et al., 1989; Lacour et al., 2015). Diapausing eggs have been found to be resistant to below-freezing temperatures and probably allowed *Ae. albopictus* populations to overwinter and spread towards higher latitudes than *Ae. aegypti* (Hawley et al., 1989; Thomas et al., 2012).

2.1.3 *Aedes japonicus japonicus*

Aedes (Hulecoeteomyia) japonicus japonicus (Theobald, 1901) [*Hulecoeteomyia japonica*], the "Asian bush mosquito", originated in an area comprised between East China,

131 East Russia and Japan (Tanaka et al., 1979a). This species may be competent for the trans-
 132 mission of pathogens of medical importance for humans, such as dengue, West Nile, Zika
 133 and Usutu viruses, but only experimental evidences of its role as vector exist (Takashima
 134 and Rosen, 1989; Scott, 2003; Schaffner et al., 2011; Westby et al., 2015; Veronesi et al.,
 135 2018; Jansen et al., 2018; Martinet et al., 2019; De Carlo et al., 2020; Abbo et al., 2020;
 136 Hopkins et al., 2020; but see Kilpatrick et al., 2005 for an estimated risk of transmitting
 137 WNV by this species). Its likely lesser role as a vector for human pathogens may also be
 138 assumed from the tendency to feed on other species than humans as well as the preference
 139 for more natural over urbanised areas. Established populations of *Ae. japonicus* were de-
 140 tected in North America from 1998 and more recently in European countries (Scott, 2003;
 141 Versteirt et al., 2009; Seidel et al., 2016; Eritja et al., 2019; Müller et al., 2020; ECDC,
 142 2021). This species is well adapted to cold climates, overwintering either as larvae in the
 143 warmer areas, or as diapausing eggs (Krupa et al., 2021) in areas where larval habitats
 144 freeze completely (Scott, 2003; Reuss et al., 2018; Day et al., 2020).

145 **2.1.4 *Aedes koreicus***

146 *Aedes (Hulecoeteomyia) koreicus* (Edwards, 1917) [*Hulecoeteomyia koreica*] commonly
 147 referred to as the "Korean bush mosquito" is native to temperate areas of Northeast Asia
 148 comprising Russia, the Korean peninsula, Japan and north-east China (Tanaka et al., 1979b).
 149 This species is a suspected vector of *Dirofilaria immitis*, Japanese encephalitis and chikun-
 150 gunya viruses, but it has not yet been directly implicated in transmission events of zoonotic
 151 pathogens (Tanaka et al., 1979b; Montarsi et al., 2015a; Ciocchetta et al., 2018). *Aedes*
 152 *koreicus* is adapted to temperate climates (Versteirt et al., 2012) and has recently colonised
 153 areas of Central Europe while continuing its range expansion (Capelli et al., 2011; Versteirt
 154 et al., 2012; Montarsi et al., 2015a; Marcantonio et al., 2016; Werner et al., 2016; Negri
 155 et al., 2021; Horváth et al., 2021; Andreeva et al., 2021; Gradoni et al., 2021; ECDC, 2021).
 156 *Aedes koreicus* seems to prefer rural over highly urbanised habitats and has been found to
 157 feed on other species than humans (Montarsi et al., 2013, 2014; Cebrián-Camisón et al.,
 158 2020). In areas where *Ae. koreicus* lives in sympatry with other invasive *Aedes* species,
 159 the Korean bush mosquito is able to colonise higher altitudes and its development can start
 160 earlier in the season with respect to other AIM (Montarsi et al., 2015a; Marcantonio et al.,
 161 2016). This trait may give them a competitive advantage over other container-breeding
 162 mosquitoes whose adults emerge later in the season.

2.2 The theoretical structure of the model

The basic structure of `dynamAedes` has been described in Da Re et al. (2021). We amended some components of the model to generalise its structure. Thus, we provide here a short recap of model structure while describing the new model features.

`dynamAedes` is composed of three main compartments (life stages) that represent a simplified version of *Aedes* the mosquito life cycle: egg, juvenile and adult stages (Fig. 1). Larval and pupal stages, which can be assumed to have somewhat similar physiological requirements, are fused in a unique "juvenile" compartment. Each compartment is divided into sub-compartments to account for the different physiological states for individuals in the three main compartments (e.g. 1-day old adult females that are not sexually mature). The number of sub-compartments into each compartment is dictated by the known minimum number of days needed by each species to pass to the next stage or complete the gonotrophic cycle (for adults). Thus, the minimum duration of development in each compartment varies among developmental stages as well as among species. As an example, the whole duration of the developmental cycle (i.e. from egg-laying to adult emergence) has a minimum duration of 11 days for *Ae. aegypti* and *Ae. albopictus*, whereas 21 days for *Ae. koreicus* and *Ae. japonicus* (see Tab S3 in SM for generic model assumptions).

In the model, time is treated as a discrete quantity and "day" is the fundamental temporal unit. Therefore, each event in the simulated life cycle occurs once per day and always in the same order. The model can be run with or without a spatial structure. If the model is spatially explicit, space is treated as a discrete quantity. In this case, the fundamental spatial unit is a (user defined) cell of a lattice grid into which the species life cycle takes place and, if relevant (see below), among which adult mosquitoes disperse.

Adult female mosquitoes lay non-diapausing eggs, E, in the summer months or diapausing eggs, Ed, at the end of the season. The embryonic development and hatching of diapausing eggs are activated by increasing daily temperature and photoperiod (typically at the end of winter or early spring). All the developmental and reproductive events considered in the model were treated as stochastic processes with probabilities derived from temperature(or photoperiod)-dependent functions by following the generic formulation:

$$X_{s,t}^{event} \sim \text{Binomial}(X_{s,t-1}, \pi_X) \quad (1)$$

where $X_{s,t-1}$ may represent eggs, juveniles or adults that undergo one of the following events in the life cycle: lay eggs, hatch, emerge or survive in cell s , at the end of the day $t - 1$. π_X is the temperature-dependent (or photoperiod dependent for the hatching of diapausing eggs) daily probability of any of the life cycle events X . All the temperature-dependent functions were calibrated using data from the scientific literature (see Tab S4 in SM) fitted using exponential, polynomial equations, and non-linear Beta density functions,

Figure 1: Graphical representation of dynamAedes model structure, adapted from Da Re et al. (2021): a) the life cycle of a generic simulated mosquito species, while in b) a representation of active and passive dispersal processes happening within the Adult (A) compartment at local scale. E: egg compartment; Ed: diapause egg compartment (available for all species except *Ae. aegypti*); J: juvenile compartment; A: adult compartment; Aad: adult active dispersal; Apdd: adult passive dispersal; Apd: adult probability of being caught in a car.

using a combination of `drc` (non-linear models) and `aomisc` (Beta function self-starters) R packages (Ritz et al., 2015; Onofri, 2020). The beta function derives from the beta density function and it has been adapted to describe phenomena taking place only within a minimum and a maximum threshold value (threshold model), such as physiological rates with respect to temperatures in the mosquito life cycle (Onofri, 2020). In a similar fashion, adult active dispersal was modelled as species-specific log-normal decaying functions of distances derived from dispersal estimates from field observations for *Ae. aegypti* and *Ae. albopictus* (Roche et al. 2015; Marcantonio et al. 2019; Marini et al. 2019a; Müller et al. 2020; see Tab S5 in SM for dispersal parameters). In addition to active dispersal, the model also considers dispersal aided by cars along the main road network (a matrix containing the coordinates of the grid cells of the landscape intersecting the road network must be provided, see the "Spatial scales of the model and temperature data sources" section), defined as the "hitchhiking" probability of a female to enter in a car and to be driven and released further away. This probability has been defined for all species by estimates measured for *Ae. albopictus* (Eritja et al., 2017), while the average distance covered by a single car trip was taken from Pasaoglu et al. (2012). This type of dispersal is thought to be amongst the main drivers of medium-range geographical expansion for invasive *Aedes* mosquitoes, especially for *Ae. aegypti* and *Ae. albopictus* (Marcantonio et al., 2016; Eritja et al., 2017; Müller et al., 2020).

Density-dependent survival is an important regulatory factor of mosquito population dynamics (Gilpin and McClelland, 1979). Its regulatory effect for juvenile stages appears to be more common in mosquitoes breeding in container or highly ephemeral habitats (Juliano, 2007), such as invasive *Aedes*. In `dynamAedes` we parameterised a density-dependent function by extracting observations on *Ae. aegypti* from Figures 2a and 2b in Hancock et al. (2016) using the Webplot Digitizer (Rohatgi, 2020). We considered the proportion of juveniles that survived through the juvenile stage (in a 2 L container) reported by these authors as an estimate of juvenile survival probability at different densities. Mortality probabilities ($1 - \text{proportion surviving}$) were converted into rates, which were scaled to a daily time step dividing by the corresponding immature development time (in days) at different densities. Finally, we regressed the natural logarithms of these daily mortality rates on the corresponding densities. The fitted daily survival rate at different densities was then summed to the temperature-dependent juvenile mortality. The resulting probability was then used to inform a binomial random draw (see equation 1) describing overall juvenile daily survival.

Some invasive *Aedes* can lay eggs resistant to low temperature commonly referred to as "diapausing eggs" (Thomas et al., 2012; Lacour et al., 2015; Krupa et al., 2021). Diapause describes the evolutionary adaptation exploited by insect species to overcome unfavourable environmental conditions by passing through an alternative and dormant physiological stage. In *Ae. albopictus*, maternal photoperiod is the environmental stimulus implied to induce oviposition of "diapausing eggs" (Pumpuni et al., 1992; Lacour et al., 2015). In `dynamAedes`, the oviposition of diapausing/non-diapausing eggs was integrated as a species-specific exponential function on the incidence of diapausing eggs given photoperiod length (and thus geographically-dependent; Urbanski et al. 2012; Petrić et al. 2021). The function is based on data from Lacour et al. (2015) for *Ae. albopictus* and Krupa et al. (2021) for *Ae. japonicus*. We applied the same diapausing function developed for *Ae. japonicus* to *Ae. koreicus* due to the close phylogeny of these species and the lack of data for *Ae. koreicus* (see Tab. S6). The daily survival of diapausing eggs was set to be constant (0.99) only for *Ae. japonicus* and *Ae. koreicus*, while for *Ae. albopictus* we used the exponential function described in Metelmann et al. (2019). The hatching rate of diapause eggs was triggered by an increasing photoperiod regime (spring) from 11.44 hours of light for *Ae. albopictus* (95th percentile estimated from Petrić et al. 2021) and 10.71 hours for *Ae. japonicus* or *Ae. koreicus* (Krupa et al., 2021).

2.3 Overview of the R package

The function `dynamAedes.m` calls the model and allows to customise the simulated scenario through a suite of options. As for the simplest application of the model (no explicit

253 spatial dimension, `scale="ws"`, see next paragraph for further details), the user has to
 254 define what species to model through the argument `species` (default `"aegypti"`), the
 255 "type" and number of introduced propagules through `intro.eggs`, `intro.juvenile`
 256 or `intro.adults` (default `intro.eggs=100`, `intro.juvenile=0`, `intro.adults=0`),
 257 and the volume (L) of water habitats wanted in each spatial unit with the argument `lhvw`
 258 (larval-habitat water volume, parameterised from Hancock et al. 2016; default `lhvw=2`;
 259 see Fig S13 for a sensitivity analysis of this parameter). Moreover, the argument `temps.matrix`
 260 takes the matrix of daily average temperature (in Celsius degree) used to fit the life cycle
 261 rates. This matrix must be organised with the daily temperature observations as columns
 262 and the geographic position of the *i*-grid cell as rows (it follows that the matrix will have
 263 only one row when `scale="ws"`). The day of start, end and number of iterations are
 264 defined by `startd`, `endd` and `iter`, respectively. The model has been optimised for
 265 parallel computing and the number of parallel processes can be specified through the op-
 266 tion `n.clusters`. If the modelled species is *Ae. albopictus*, *Ae. japonicus* or *Ae.*
 267 *koreicus* (e.g., `species="albopictus"`) then the arguments defining latitude (`lat`),
 268 longitude (`long`) and year of introduction (`intro.year`) should be adequately defined
 269 to allow a correct switch to and from the egg diapausing stage.

270 The default output of `dynamAedes` consists of a list of numerical matrices containing,
 271 for each iteration, the number of individuals in each life stage per day (and for each grid
 272 cell of the study area if `scale="lc"` or `"rg"`). If the argument `compressed.output=FALSE`
 273 (default `TRUE`), the model returns the daily number of individuals in each life stage sub-
 274 compartment. The model, coded in the R statistical language (R Core Team, 2021),
 275 and adapted for parallel computation, is available on the the following link [https:](https://github.com/mattmar/dynamAedes)
 276 [//github.com/mattmar/dynamAedes](https://github.com/mattmar/dynamAedes) (it has meanwhile been submitted to the
 277 CRAN).


```

#model parameters
cl=47 #number of cluster
it=100 #number of iterations
ie=500 #number of introduced eggs
str=1 #day intro
endr=ncol(wvl) #day end simulations

aeg.cal <- dynamAedes(species="aegypti",
                      scale="rg",
                      temps.matrix=wvl,
                      startd=str, endd=endr,
                      n.clusters=cl,
                      iter=it,
                      intro.eggs=ie,
                      ihwv=100,
                      verbose=FALSE)

albo.cal <- dynamAedes(species="albopictus",
                      lat=37, long=-120,
                      intro.year=2015,
                      scale="rg",
                      temps.matrix=wvl,
                      startd=str, endd=endr,
                      n.clusters=cl,
                      iter=it,
                      intro.eggs=ie,
                      ihwv=100,
                      verbose=FALSE)

```

278

279 2.3.1 Spatial scales of the model and input temperature data

280 The selection of the geographical scale for population dynamics is a crucial aspect of
 281 the whole package and the temperature dataset provided to dynamAedes function must
 282 reflect this decision. Along with the photoperiod, temperature is the other only environ-
 283 mental driver of our model, which is dictated by its central role in mosquito develop-
 284 ment and activity. The measurement of temperature is inevitably scale-dependent, thus we
 285 structured the model to allow for temperature datasets relevant for different measurement
 286 spatial scales (Fig. 2) and to match the different hypotheses that users may want to test.

287 The punctual or "weather station" scale (scale="ws") is the smallest geographic
 288 scale (i.g., no spatial dimension) available in dynamAedes and the environment mod-

Figure 2: dynamAedes allows to simulate *Aedes* mosquitoes population dynamics at three different spatial scales: a) regional, b) local, and c) punctual (weather station). Passive and active dispersal is enabled only at local spatial scale.

elled is assumed to be what represented by the chosen weather station (or any other data-loggers). In this case, the model has no spatial structure, thus dispersal is not considered: the model will return the temporal trend of population dynamics given the chosen temperature, larval-habitat water volume and photoperiod conditions.

The "local" scale (`scale= "lc"`) represents those scenarios and spatial resolutions at which species dispersal and local microclimate variability are relevant for users. We suggest to keep the resolution of the matrix of temperatures equal or smaller than the maximum daily dispersal range of the mosquito species (i.e., usually under 1 km for *Aedes* species; Guerra et al., 2014). The optional arguments `cellsize`, `dispbins` and `maxdisp` are available to fine tune the dispersal kernel which drives the spatial behaviour of the simulated mosquito populations. The argument `cellsize` (default `"cellsize=250"` meters) sets the minimal distance of the dispersal kernel and should

match the size of the cell to avoid inconsistencies (i.e., mosquitoes dispersing at a finer or bigger grain than the arena), `maxdisp` sets the maximum daily dispersal (default `maxdisp=600` meters), and `dispbins` the resolution of the dispersal kernel (default `dispbins=10`). Passive dispersal is also implemented and it requires i) a matrix containing the coordinates of the grid cells of the landscape intersecting the road network (argument `road.dist.matrix`), and ii) to specify the average car trip distance through the argument `country`, which can be defined by the user or considering estimates for the following countries: France, Germany, Italy, Poland, Spain and the United Kingdom Pasaoglu et al. (2012). An extensive example of model application at local spatial scales is described in Da Re et al. (2021).

The rationale behind the third spatial scale considered in the model, the "regional" scale (`scale= "rg"`) was to return an overview of invasive *Aedes* population dynamics over large extents (i.e., larger than 1 km). The model in regional scale does not account for species dispersal, introductions happen separately (but at the same time) in each grid cell which hence are closed systems. The output of the model at "regional" scale can be compared to those produced by correlative species distribution models (SDMs), with the advantage of mechanistic rather than purely correlative model foundations.

The amount of water available for larval development in each spatial unit(s) (at any model spatial scale) was set as 2 L that is the water volume considered in the experiments we used to parametrise model functions Hancock et al. (2016). It is likely that for many real-world model applications, the relative availability of breeding habitats is much higher, and we encourage users to set a value based on their scenarios and hypotheses (i.e. through the model option `lhvw`; Hartemink et al. 2015).

2.3.2 Auxiliary functions

Several auxiliary functions are available to analyse model outputs. The function `psi` returns the proportion of model iterations that resulted in a viable population for the given date. It works for all spatial scales and the output can reflect either the overall grid or each single cell. Likewise, summaries of mosquito abundance at each life stage for each day can be obtained through `adci`, which by default returns the inter-quartile range abundance of each life stage. Similarly, `icci` returns a summary of the number of invaded cells over model iterations. Estimates of dispersal spread (in km²) of the simulated mosquito populations is provided by the function `dici`, which is available only for model results computed at the local scale (the only scale which integrates dispersal). Finally, the function `get_rates_spatial`, returns the output of the temperature-dependent physiological functions used by `dynamAedes` to derive the daily rates. It can be used to better understand the outcome of model simulations, by highlighting those areas where the predicted

values of the temperature-dependent functions are maximised or minimised, or to derive causal-based physiological estimations that, for example, could be used as inputs for correlative SDMs (e.g., Kearney and Porter, 2009; Mathewson et al., 2017).

3 Case studies and model validation

We applied the model to three case studies representing different geographical scales and areas, species and invasion trajectories. The case studies were chosen considering the availability of optimal mosquito data-sets to show model strength and weakness. We did not report any example for the "local scale" as it had already been provided in Da Re et al. (2021), who applied a previous version of *dynamAedes*.

3.1 Regional scale models

3.1.1 Likelihood of successful introductions of *Ae. albopictus* and *Ae. aegypti* across California, USA

We used *dynamAedes* at "regional" scale to assess the likelihood of successful introductions for two invasive *Aedes* across California: *Ae. aegypti* and *Ae. albopictus* which are considered established from 2011 and 2013, respectively (Fujioka, 2012; Gloria-Soria et al., 2014). California is among the few areas where established populations of these two species were detected during the last decade and their progressive spread was documented in great detail². We downloaded daily minimum and maximum temperature data from the NASA Daily Surface Weather Data on a 1-km Grid for North America (DAYMET), Version 4 (Thornton et al., 2020) from 1 January 2011 to the 31 December 2018. These two sets of data (in netCDF raster format) were clipped to the boundary of California and aggregated at a spatial resolution of 2.5 km by using a combination of GDAL (GDAL/OGR contributors, 2021) and Climate Data Operators (CDO; Schulzweida, 2019) software. The two sets of raster layers were then imported in R 4.0.4 (R Core Team, 2021), transformed in matrices, averaged and converted to integers to obtain a single average daily temperature integer matrix with cell id as observations (rows) and days as variables (columns). This dataset was used as the input temperature matrix for the model. We run 80 model iterations for five years, introducing 500 eggs in each cell of the gridded landscape on 15 May 2011 and 2013 for *Ae. aegypti* and *Ae. albopictus* respectively. By using the auxiliary function `psi`, we then derived a map showing the proportion of iterations with viable population of both *Ae. aegypti* and *Ae. albopictus* at the end of the simulated period (15

²See <https://maps.vectorsurv.org/>

May 2016 and 2018). The photoperiod was set to match conditions in the geographical center of California (Lat 37° N, Lo -120° W) and the amount of breeding habitat was set to 100 L (per cell), which we assumed to be representative of the potential water larval habitats available given cell size and overall regional climate.

We validated model prediction using maps of cities in California with known species presence updated to 2021 by the California Department of Public Health (CDPH³). We derived the average predicted probability of successful introduction for each city and computed the Area Under the ROC Curve (AUC) score which defines the probability that a randomly chosen positive city will be ranked higher than a randomly chosen negative city. An AUC score > 0.5 indicates that the model is performing better than random, while a score of 1 indicates perfect prediction. In addition, we calculated the percentage of positive cities that fell into a grid cell that had a probability of establishment higher or equal to 1% (e.g., at least 1 out of the n_{th} iterations reported a viable mosquito population in that cell).

The predicted spatial pattern of areas with a high likelihood of *Ae. aegypti* and *Ae. albopictus* successful introduction is in general consistent with updated information on the presence of this species (Fig. 3). Predictions show moderate-to-high probabilities of a successful introduction in all counties with known presence of these species, except for the extreme South-East coastal part of the state, where *Ae. aegypti* was predicted to have a low probability of successful introduction whereas being well established (Fig. 3). This may be due to the high micro-climatic variability that characterises coastal areas of California which may not be resolved by the temperature datasets that we have considered in this case study. On the contrary, areas predicted to be suitable for *Ae. albopictus* exceeded by far the known current distribution of this species. Factors other than temperature and photoperiod, more nuanced aspects of species invasion history and the extremely low humidity during the dry season in the Central Valley of California or the Inland Empire, may hinder species establishment in these areas. Nevertheless, recently *Ae. albopictus* was found as north as Redding, Shasta County, thus it is not unlikely that this species is also present (perhaps at low densities), but not yet detected, at southern latitudes in California.

Both models had over 75% of successful introduction scores (calculated as the proportion of pixels with species observations and simulated proportion of invasion > 1%) when validated at city levels (Tab. 1). Still, only the prediction for *Ae. aegypti* validated at city levels reported an AUC score bigger than 0.5 (Tab. 1), whereas validating the model by averaging the predictions at county level resulted in an AUC bigger than 0.5 for both species (0.892 and 0.717 for *Ae. aegypti* and *Ae. albopictus* respectively; Fig. S14).

³California Department of Public Health, "Map and City List of *Aedes aegypti* and *Aedes albopictus* Mosquitoes in CA, 2011-2021" (accessed on 28th October 2021): <https://www.cdph.ca.gov/Programs/CID/DCDC/Pages/VBDS.aspx>

Figure 3: Predicted percentage of established introductions of *Ae. aegypti* *Ae. albopictus* in California (USA) for the years 2011-2016 and 2013-2018, respectively. The red dots represent the centroids of the Californian municipalities with established populations as reported by the Californian Department of Public Health (CDPH).

3.1.2 Likelihood of successful *Aedes albopictus* introductions in France

Aedes albopictus was first detected in metropolitan France in 1999 (Schaffner et al., 2000) and since 2004 it has established populations in the southern part of the country while still expanding its distribution range (ECDC, 2021). We used *dynamAedes* model at "regional" scale to assess the success of introductions of *Ae. albopictus* for the whole metropolitan France. We processed ERA5-Land (Muñoz-Sabater et al., 2021) hourly air temperature measured at 2 m above surface from January 1st 2015 to December 31st 2020 in the Climate Data Store (CDS) Toolbox⁴ to get the daily mean temperature of the period considered for the whole metropolitan France, at the spatial resolution of ~ 9 km. The netCDF file obtained was imported in R 4.0.4 (R Core Team, 2021), where was clipped to the extent of metropolitan France, converted from degrees Kelvin to Celsius and converted to integer to obtain a single average daily temperature integer matrix with cell id

⁴CDS Toolbox: <https://cds.climate.copernicus.eu/toolbox/doc/index.html>

as observations (rows) and days as variables (columns). This dataset was then used as the input temperature matrix for the model. We run 100 simulations for five years, introducing 100 eggs in each cell of the gridded landscape on 15 May 2015. By using the auxiliary function `psi`, we derived a probability map of the areas showing the proportions of iterations that produced a viable population of *Ae. albopictus* five years after the simulated introductions.

We validate the model predictions using the list of the 3419 French municipality reporting established *Ae. albopictus* until 2020 provided by the French Health Ministry (data collected thanks to active monitoring from French mosquito operators and passive surveillance⁵). Similarly to the previous case study, we computed both the AUC as well as the proportion of positive location falling into a grid cell that had a percentage of established introductions higher than or equal to 1%. The spatial pattern of the areas predicted to have a high likelihood of successful *Ae. albopictus* introduction (Fig. 4) is consistent with updated observational data (*Ae. albopictus* map; ECDC, 2020) as well as with the results of other mechanistic models (see for instance Metelmann et al., 2019; Pasquali et al., 2020). The Mediterranean French coast and the Rhone valley are the areas where our model predicted the highest percentage of successful introduction. Similarly, the Aquitaine region on the Atlantic coast and the Alsace region in the North-East part of France showed relatively high predicted percentage of successful introduction. The northern and the central part of France, as well as the Pyrenees areas, show low percentage of successful introduction under the current climatic conditions. However, the resolution of the pixel, approximately 10 km, may have played a role influencing the model outcomes especially in topographically complex areas such as the Pyrenees or the French Alps, where the microclimate of the valleys may be underestimated. Similarly, the model was not able to predict the successful introduction of the species in areas such as Paris, where *Ae. albopictus* is established and probably favoured by i) local climatic factors such as the urban heat island effect, and ii) the continuous inflow of *Ae. albopictus* propagules to Paris from areas where the species is already established. Indeed, most railways, flights and highways have a connection with Paris, and the Paris-Lyon-Mediterranée axis is the main artery of France, with an average of >60,000 cars/day on the highway⁶ and 240 trains per day⁷, thus the quantity of imported mosquitoes can compensate for the less favorable climatic conditions of Paris compared to the Mediterranean region (recent phylogeographic findings support this hypothesis, see Sherpa et al., 2019). All the model performances metrics assessed support the capacity of the model to discriminate between areas where the species can or cannot be established

⁵https://signalement-moustique.anses.fr/signalement_albopictus/

⁶<https://www.data.gouv.fr/fr/datasets/trafic-moyen-journalier-annuel-sur-le-reseau-routier/>

⁷<https://www.leparisien.fr/economie/l-europe-fait-passer-la-lgv-paris-lyon-a-l-heure-20190910>
php

449 (Tab. 1).

Figure 4: Percentage of successful introduction for *Ae. albopictus* in France for the years 2015-2020: a) Model prediction, b) Model prediction and in red the centroids of the French Municipalities with established population of *Ae. albopictus* reported by the French Health Ministry (SI-LAV).

450 **3.2 Punctual population dynamics temporal trends**

451 **3.2.1 *Aedes albopictus* population dynamics in Nice, South-East France**

452 *Aedes albopictus* is established mainly in the southern part of metropolitan France where
 453 more than 40% of municipalities are colonized⁸ and it is still expanding in other areas. We
 454 computed the population dynamics of the species informing dynamAedes model at punc-
 455 tual spatial scale using temperature observations downloaded from the National Oceanic
 456 and Atmospheric Administration (NOAA) network via the R package rnoaa. The obser-
 457 vations of the NOAA weather station located in Nice area (usaf code = 076900, Lat 43°
 458 42' 00" N, Lon 7° 13' 12" E, 3.7 m a.s.l.), spanning from January 1st 2013 to December
 459 31st 2018, were linearly interpolated to fill missing values at hourly and daily level. Af-
 460 terwards, the daily average for all observations was computed. We run 100 simulations
 461 for five years, introducing 500 eggs on 15 May 2013. By using the auxiliary function
 462 adci, we then derived the daily abundance inter-quantile range for each life stage and the
 463 abundance of newly-laid eggs per day.

464 The model was validated comparing the simulated newly laid eggs per day to egg
 465 counts from mosquito ovitrap data, following the validation approach presented in Tran
 466 et al. (2013). During the years 2014-2018, fifty ovitraps were installed in the Nice area and
 467 inspected fortnightly from April-May to November-December (data collected and kindly
 468 provided by EID Méditerranée). We computed the Spearman's ρ correlation coefficient
 469 between the weekly-aggregated simulated newly-laid eggs and the mean observed eggs
 470 per day.

471 Results showed that the model was able to correctly reproduce the seasonal dynamics
 472 of the new-laid eggs over five years (Spearman's $\rho = 0.753$, p.value < 0.001). The first
 473 simulated eggs were laid during the late spring each year, confirming the fact that the first
 474 overwintering eggs hatch at the end of the winter season or at the beginning of spring.
 475 The ovipositing season seem to last until the late autumn accordingly to the observations,
 476 while our model seems to predict a shorter length of the ovipositing period. Nevertheless
 477 the model is able to correctly infer the peak of the ovipositing season during the summer
 478 months (Fig. 5).

479 **3.2.2 *Aedes koreicus* population dynamics in Trento, North-East Italy**

480 *Aedes koreicus* was first detected in Trento Autonomous Province (NE Italy) in 2013, soon
 481 after the first Italian detection in the neighboring Belluno province (Capelli et al., 2011).

⁸<https://solidarites-sante.gouv.fr/sante-et-environnement/risques-microbiologiques-physiques-et-chimiques/especes-nuisibles-et-parasites/article/cartes-de-presence-du-moustique-tigre-aedes-albopictus-en-france-metropolitaine>

Figure 5: Temporal trend reporting the simulated and observed new-laid eggs of *Ae. albopictus* years 2014-2018 in Nice, SE France. The light blue bands represent the winter seasons, while the orange bands the summer seasons. The simulated data were rescaled for graphical purpose using as rescaling factor the ratio between the maximum observed value and the maximum median simulated values.

482 We computed the population dynamics of the species informing `dynamAedes` model
 483 at punctual spatial scale using temperature observations downloaded from the local net-
 484 work of weather stations⁹. The daily average temperature observations from the "Trento
 485 Laste" weather station (Lat 46° 04' 18.5" N, Lon 11° 08' 08.5" E, 312 m a.s.l.) spanning
 486 from 1 January 2015 to 31 December 2018 were linearly interpolated to fill missing val-
 487 ues. We run 100 simulations for five years, introducing 500 eggs on 15 May 2015. Using
 488 the auxiliary function `adci`, we then derived the daily inter-quartile range abundance for
 489 each life stage and for the daily host-seeking female sub-compartment.

490 The model was validated computing Spearman's ρ correlation coefficient between
 491 the monthly-aggregated simulated host-seeking females and the observations gathered

⁹www.meteotrentino.it

from four BG-Sentinel traps installed in Trento municipality from April to November during the years 2016, 2017, and 2018 (data obtained from Marini et al. 2019b). In order to compare observed and simulated data, the whole simulated host-seeking females abundance was multiplied by a BG-sentinel catching rate equal to 0.157, as estimated by Marini et al. (2019b) and similar to what already reported for *Ae. albopictus* in previous studies (Guzzetta et al., 2017).

The simulated population dynamics showed that *Ae. koreicus* could be successfully introduced in the study area. The model correctly predicted the start of the seasonal activity in early spring, while the higher abundance of female host-seeking mosquitoes was predicted to be in late summer. Considering the three years together, our model was able to reproduce the observed seasonal population dynamics, where the 76.2% (47.6%) of the observed captures lie within the 95% (50%) credible intervals of model predictions (Tab S7). Similarly, the Spearman's rho for the three years was 0.735 (p.value < 0.001) (Tab. 1)

Figure 6: Temporal trend reporting the boxplot of simulated and observed host-seeking *Ae. koreicus* females for the years 2016-2018 in Trento, NE Italy.

Species	Geographic location	Scale	AUC	Specificity	Sensitivity	Sensitivity 1%	Spearman's ρ
<i>Ae. aegypti</i>	California	Regional	0.658 (0.401-0.914)	0.8	0.642	0.795	-
<i>Ae. albopictus</i>	California	Regional	0.435 (0.367-0.502)	0.241	0.641	1	-
<i>Ae. albopictus</i>	France	Regional	0.874 (0.867-0.880)	0.788	0.802	0.848	-
<i>Ae. albopictus</i>	S-E France	Punctual	-	-	-	-	0.753***
<i>Ae. koreicus</i>	N-E Italy	Punctual	-	-	-	-	0.735***

Table 1: Model validation. The column "Sensitivity 1%" reports the proportion of cities predicted to have at least one successful introduction over the total number of iterations (predicted introduction success equal or over 1%. *** p.value < 0.001).

4 Discussion

From an ecological perspective, our modelling approach focuses on the species fundamental thermal niche (*sensu* Hutchinson, 1957), since we considered temperature as the main driver of population growth and dynamics. In light of this, our model was able to infer spatial and temporal population dynamics of different species, at different geographic scales and locations. The model showed overall good validation performance, and the areas predicted to likely support *Aedes* mosquito populations largely matched what reported by observational studies and other existing models (Kraemer et al., 2015, 2019; Oliveira et al., 2021).

Nevertheless, we raise the attention on two aspects that should be considered when applying dynamAedes and interpreting model results. First, good quality information on survival and developmental rates is largely available for *Ae. aegypti* and *Ae. albopictus*, whereas much less for the other two species. Similarly, mosquito observational datasets with sufficient longitudinal depth for model validation are scarce for *Ae. koreicus* and presently absent, to the best of our knowledge, for *Ae. japonicus*. Thus, whilst we have built the foundations for an open-source modelling framework that can be progressively expanded, life cycle functions and thus outputs for *Ae. japonicus* and *Ae. koreicus* should be interpreted more carefully.

Secondly, we recognise that pixel size may influence model outcome because of the aggregating effect of the Modifiable Unit Areal Problem (Jelinski and Wu, 1996; Da Re et al., 2020). While the consequences of this artifact are well known in SDMs applications, they are rarely mentioned or addressed (but see Peterson, 2014). Thus, the correct choice of temperature datasets is crucial to investigate species population dynamics and interpret model results (Bütikofer et al., 2020). Climatic reanalysis and Global/Regional Circulation Models are reliable data sources with high temporal resolution for present climatic conditions and robust future projections, though they have a coarse spatial resolution that may underestimate the effect of microclimate on species biology (Metelmann et al., 2019;

Liu-Helmersson et al., 2019). Coarse resolution of input temperature data is rather common since temperature is often estimated over large extents, and can become an issue in topographically complex regions where the effect of microclimatic variation on population dynamics may pass undetected. The results showed in Fig. 3–4 may be better interpreted considering the coarse resolution of temperature data which may have caused lower proportion of established populations in topographically complex areas such as the Pyrenees or the Peninsular Ranges. Interpolated local micro-climatic conditions, for example estimated with the `microclima` R package (Maclean et al., 2019), have the advantage of providing fine spatial and temporal resolution datasets. But, they need to be first properly validated with field data which are typically difficult to gather also over small geographical extents. Temperature measured with classical weather stations may be considered as the most accurate available observations of local climatic conditions. Though it is not always easy to deal with such data due to the limited number of weather stations and gaps in the time series, they are suited for statistical downscaling or bias adjustment for climate change projections (Bedia et al., 2020).

4.1 Model assumptions

Model structure has been designed to be as ecologically relevant as possible considering available data, however, when data were limited, we relied on a set of "expert-based" assumptions that must be clearly stated.

The interplay of multiple environmental factors drives the population dynamics of *Aedes* mosquitoes but we chose to mold our model framework just on established information available for temperature and photo-period (Pumpuni et al., 1992; Waddock et al., 2013; Eisen et al., 2014). This choice was suggested by a generalised lack of clear relationships between other environmental factors and *Aedes* population dynamics. For example, concerning the role of precipitations, different studies report contrasting results (Koenraadt and Harrington, 2008; Tran et al., 2013; Caldwell et al., 2021). Moreover, invasive *Aedes* mosquitoes mostly thrive in urban or suburban landscapes where the presence of standing water is often independent from precipitations (except for extreme rainfall events (Roiz et al., 2015)). We suggest that, at the present stage, `dynamAedes` it is better suited for applications in temperate climates, where temperature seasonality is a more important limiting factor than in tropical climates, where other factors may limit mosquito life cycle (Lega et al. (2017)).

On the one hand, we did not consider in the model biotic interactions such as prey-predator or food and space competition with other mosquito taxa during the larval stages, despite this is another factor that influences the trajectory of introduced populations (Armistead et al., 2008; Tripet et al., 2011; Reiskind and Lounibos, 2013; Montarsi et al., 2013,

2015b; Müller et al., 2018). On the other hand, we considered the effect of intra-specific competition on larval survival (but not development). We generalised the information available for *Ae. aegypti* to the other *Aedes* species due to the lack of species-specific experiments (Hancock et al., 2016). We recognise that this is not optimal under many facets, but intra-specific larval interactions were a key driver of mosquito-population dynamics that could not be excluded by model structure.

Finally, we did not consider evolutionary processes in our model which may affect invasion success over medium-long time spans. Given the reproductive strategy of *Aedes* mosquitoes, rapid evolutionary processes may take place over relatively short temporal periods (e.g. decades), making introduced populations able to extend their original niche (McBride et al., 2014b).

4.2 Proposed research directions

dynamAedes is an open-source tool for testing ecological hypothesis and/or to support management plans concerning AIMs. Selecting areas at risk of AIM establishment or periods when abundances are more likely to peak should be considered as facets of AIM surveillance. The importance of such early information becomes fundamental for protecting human health when treating AIM involved in pathogen transmission, as early information on new trajectories of AIM populations becomes critical in the current climate change era. Mosquitoes are affected by temperature changes in, often, predictable ways, though changes in population dynamics can be extremely rapid. Modelling population dynamics under climate change scenarios may thus provide information for anticipating both AIM population changes in space and time and human health risks.

The conceptualisation and design of *dynamAedes* required the review of up-to-date ecological and physiological literature available on four *Aedes* species, which was integrated with feedback from expert ecologists and medical entomologists. It emerged that knowledge on some ecological aspects of these species is highly fragmented or poor (e.g., Cebrián-Camisón et al. 2020), and often dependant on experimental settings and lab strain (e.g., Kramer et al. 2020). Thus, the exploitation of such sets of information for process-based models and, hence, for AIM management would greatly benefit from a standardised review effort and possibly centralised repositories, as already done in other scientific fields such as plant functional traits (Kattge et al., 2020) or systems biology (Tsigkinopoulou et al., 2018). Moreover, experiments on *Ae. japonicus* and *Ae. koreicus* life cycles are just starting to unravel these species eco-physiological rates (*Ae. koreicus*: (Ciocchetta et al., 2017; Marini et al., 2019b); *Ae. japonicus*: (Scott, 2003; Reuss et al., 2018; Wieser et al., 2019) and much work remain to be done on this species. On the contrary, there is large information concerning *Ae. aegypti* and *Ae. albopictus* physiological

rates which anyhow have been shown to be highly heterogeneous (Eisen et al., 2014) likely due to non-standardised experimental designs and ecological plasticity of *Aedes* populations (*sensu* Kramer et al. 2021). Information regarding mosquito dispersal is even scarcer and available only for *Ae. aegypti* and *Ae. albopictus*, while passive dispersal through auto-vehicles has been estimated only for *Ae. albopictus* (Eritja et al., 2017) despite the worldwide spread of *Aedes* species was most likely caused by means of passive dispersal.

From a biological perspective, future developments of `dynamAedes` may consider also the addition of a `strain` argument, where the physiological temperature-dependent function can be fitted on geographically different mosquito strains, such as tropical, mediterranean or temperate (Marini et al., 2020; Kramer et al., 2021). Moreover, if observational data are available, calibration of some parameters values, such as the juvenile density-dependent mortality rate, might be implemented following for instance a Bayesian approach (Marini et al., 2019b).

We believe that a closer interaction between modelers and experimenters will motivate the collection of standardised data on unknown eco-physiological AIM rates that would lead to more accurate model predictions. This project was inspired by such interactions and, in this spirit, `dynamAedes` was meant to be modified or extended to relax its assumptions and limitations with new available information by anyone having basic R programming skills.

5 Conclusion

In this study, we presented `dynamAedes`, a mechanistic process-based model to infer invasive *Aedes* mosquito spatio-temporal population dynamics. This first version of the model showed to be often reliable in terms of both biological realism and statistical accuracy. The open-source nature and programming language accessibility and flexibility of the project offers great potential to further develop the model, allowing to better tune the temperature-dependent functions when new physiological observations and findings become available. Abundance estimations derived from `dynamAedes` could be used to inform epidemiological models (e.g., SIR or SEIR) and thus obtain estimations on the risk of pathogens transmission. Finally, it does not seem unrealistic to extend the model application to other species of the genus *Aedes* such as *Ae. notoscriptus* or to species belonging to other genus of medical interest belonging to the *Culicidae* family, such as *Anopheles*, or even to other blood-sucking insects belonging to different taxa such as *Culicoides*.

6 Acknowledgements

No conflict of interest has been declared by the author(s).

We thank the French Ministry of Health for providing the confirmed *Ae. albopictus* established population data (SI-LAV), and the EID Méditerranée for providing us the ovitraps data from Nice. Computational resources have been provided by the DART research group at the University of California, Davis as well as supercomputing facilities of the Université catholique de Louvain (CISM/UCL) and the Consortium des Équipements de Calcul Intensif en Fédération Wallonie Bruxelles (CÉCI) funded by the Fond de la Recherche Scientifique de Belgique (F.R.S.-FNRS) under convention 2.5020.11 and by the Walloon Region.

DDR and MM are supported by the FRS-FNRS. MM is also supported by the “Action de Recherche concertée” grant number 17/22-086. FR and RM were supported by the Hessian Centre on Climate Change (FZK) of the Hessian Agency for Nature Conservation, Environment and Geology (HLNUG). Research of RM was funded through the 2018-2019 BiodivERsA joint call for research proposals, under the BiodivERsA3 ERA-Net COFUND programme, and with the funding organisation FWO (BiodivERsA2018-A-323). The Outbreak Research Team of the Institute of Tropical Medicine is financially supported by the Department of Economy, Science and Innovation of the Flemish government.

This project was done within the framework of AIM-COST Action CA17108 (www.aedescost.eu).

7 Authors contribution

MM and DDR conceived the ideas, designed the methodology and analysed the data; WVB, FR, RM, SB, FM, SC, DA, GM, AP, GLA, GL and CJMK provided expert opinion on mosquito biology as well as observational datasets to validate the model; DDR, MM and SOV led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

References

- Abbo, S. R., Visser, T. M., Wang, H., Göertz, G. P., Fros, J. J., Abma-Henkens, M. H., Geertsema, C., Vogels, C. B., Koopmans, M. P., Reusken, C. B., et al. (2020). The invasive Asian bush mosquito *Aedes japonicus* found in the Netherlands can experimentally transmit Zika virus and Usutu virus. *PLoS neglected tropical diseases*, 14(4):e0008217.
- Andreeva, Y. V., Khrabrova, N. V., Alekseeva, S. S., Abylkassymova, G. M., Simakova, A. V., and Sibataev, A. K. (2021). First record of the invasive mosquito species *Aedes koreicus* (Diptera, Culicidae) in the Republic of Kazakhstan. *Parasite*, 28:52.
- Armistead, J., Arias, J., Nishimura, N., and Lounibos, L. (2008). Interspecific larval competition between *Aedes albopictus* and *Aedes japonicus* (Diptera: Culicidae) in Northern Virginia. *Journal of medical entomology*, 45(4):629–637.
- Baldacchino, F., Caputo, B., Chandre, F., Drago, A., della Torre, A., Montarsi, F., and Rizzoli, A. (2015). Control methods against invasive *Aedes* mosquitoes in Europe: a review. *Pest management science*, 71(11):1471–1485.
- Barzon, L., Gobbi, F., Capelli, G., Montarsi, F., Martini, S., Riccetti, S., Sinigaglia, A., Pacenti, M., Pavan, G., Razu, M., et al. (2021). Autochthonous dengue outbreak in Italy 2020: clinical, virological and entomological findings. *Journal of travel medicine*.
- Bedia, J., Baño-Medina, J., Legasa, M. N., Iturbide, M., Manzanar, R., Herrera, S., Casanueva, A., San-Martín, D., Cofiño, A. S., and Gutiérrez, J. M. (2020). Statistical downscaling with the downscaler package (v3. 1.0): contribution to the VALUE intercomparison experiment. *Geoscientific Model Development*, 13(3):1711–1735.
- Brady, O. J. and Hay, S. I. (2019). The first local cases of Zika virus in Europe. *The Lancet*, 394(10213):1991–1992.
- Bütikofer, L., Anderson, K., Bebbier, D. P., Bennie, J. J., Early, R. I., and Maclean, I. M. (2020). The problem of scale in predicting biological responses to climate. *Global Change Biology*, 26(12):6657–6666.
- Caldwell, J. M., LaBeaud, A. D., Lambin, E. F., Stewart-Ibarra, A. M., Ndenga, B. A., Mutuku, F. M., Krystosik, A. R., Ayala, E. B., Anyamba, A., Borbor-Cordova, M. J., et al. (2021). Climate predicts geographic and temporal variation in mosquito-borne disease dynamics on two continents. *Nature communications*, 12(1):1–13.

- 692 Capelli, G., Drago, A., Martini, S., Montarsi, F., Soppelsa, M., Delai, N., Ravagnan, S.,
693 Mazzon, L., Schaffner, F., Mathis, A., et al. (2011). First report in Italy of the exotic
694 mosquito species *Aedes (Finlaya) koreicus*, a potential vector of arboviruses and filariae.
695 *Parasites & vectors*, 4(1):1–5.
- 696 Cebrián-Camisón, S., Martínez-de la Puente, J., and Figuerola, J. (2020). A literature
697 review of host feeding patterns of invasive *Aedes* mosquitoes in Europe. *Insects*,
698 11(12):848.
- 699 Christophers, S. R. (1960). *Aedes aegypti (L.) The Yellow Fever Mosquito: Its Life*
700 *History, Bionomics and Structure*. Cambridge University Press. Google-Books-ID:
701 TBcf5DTKL1oC.
- 702 Ciocchetta, S., Darbro, J. M., Frentiu, F. D., Montarsi, F., Capelli, G., Aaskov, J. G., and
703 Devine, G. J. (2017). Laboratory colonization of the European invasive mosquito *Aedes*
704 (*Finlaya*) *koreicus*. *Parasites & vectors*, 10(1):1–6.
- 705 Ciocchetta, S., Prow, N. A., Darbro, J. M., Frentiu, F. D., Savino, S., Montarsi, F., Capelli,
706 G., Aaskov, J. G., and Devine, G. J. (2018). The new European invader *Aedes (Fin-*
707 *laya)* *koreicus*: a potential vector of chikungunya virus. *Pathogens and Global Health*,
708 112(3):107–114.
- 709 Da Re, D., Gilbert, M., Chaiban, C., Bourguignon, P., Thanapongtharm, W., Robinson,
710 T. P., and Vanwambeke, S. O. (2020). Downscaling livestock census data using mul-
711 tivariate predictive models: Sensitivity to modifiable areal unit problem. *PloS one*,
712 15(1):e0221070.
- 713 Da Re, D., Montecino-Latorre, D., Vanwambeke, S. O., and Marcantonio, M. (2021). Will
714 the yellow fever mosquito colonise Europe? Assessing the re-introduction of *Aedes*
715 *aegypti* using a process-based population dynamical model. *Ecological Informatics*,
716 61:101180.
- 717 da Silva Neves, N. A., da Silva Ferreira, R., Morais, D. O., Pavon, J. A. R., de Pinho, J. B.,
718 and Shlessarenko, R. D. (2021). Chikungunya, Zika, Mayaro, and Equine encephalitis
719 virus detection in adult *Culicinae* from South Central Mato Grosso, Brazil, during the
720 rainy season of 2018. *Brazilian Journal of Microbiology*, pages 1–8.
- 721 Day, C. A., Lewandowski, K., Vonesh, J. R., and Byrd, B. D. (2020). Phenology of
722 rock pool mosquitoes in the southern Appalachian mountains: Surveys reveal apparent
723 winter hatching of *Aedes japonicus* and the potential for asymmetrical stage-specific
724 interactions. *Journal of the American Mosquito Control Association*, 36(4):216–226.

- 725 De Carlo, C. H., Campbell, S. R., Bigler, L. L., and Mohammed, H. O. (2020). *Aedes*
726 *japonicus* and West Nile Virus in new york. *Journal of the American Mosquito Control*
727 *Association*, 36(4):261–263.
- 728 Delatte, H., Paupy, C., Dehecq, J., Thiria, J., Failloux, A., and Fontenille, D. (2008). *Aedes*
729 *albopictus* , vecteur des virus du chikungunya et de la dengue à la Réunion : biologie et
730 contrôle. *Parasite*, 15(1):3–13.
- 731 Dzul-Manzanilla, F., Ibarra-López, J., Bibiano Marín, W., Martini-Jaimes, A., Leyva,
732 J. T., Correa-Morales, F., Huerta, H., Manrique-Saide, P., and Vazquez-Prokopec, G. M.
733 (2017). Indoor Resting Behavior of *Aedes aegypti* (Diptera: Culicidae) in Acapulco,
734 Mexico. *Journal of Medical Entomology*, 54(2):501–504.
- 735 ECDC (2021). European Centre for Disease Prevention and Control and
736 European Food Safety Authority. mosquito maps [internet]. stockholm:
737 Ecdc; 2021. [https://ecdc.europa.eu/en/disease-vectors/](https://ecdc.europa.eu/en/disease-vectors/surveillance-and-disease-data/mosquito-maps)
738 [surveillance-and-disease-data/mosquito-maps](https://ecdc.europa.eu/en/disease-vectors/surveillance-and-disease-data/mosquito-maps). Accessed: 2021-12-
739 01.
- 740 Effler, P. V., Pang, L., Kitsutani, P., Vorndam, V., Nakata, M., Ayers, T., Elm, J., Tom, T.,
741 Reiter, P., Rigau-Perez, J. G., Hayes, J. M., Mills, K., Napier, M., Clark, G. G., Gubler,
742 D. J., and Hawaii Dengue Outbreak Investigation Team (2005). Dengue fever, Hawaii,
743 2001-2002. *Emerging Infectious Diseases*, 11(5):742–749.
- 744 Eisen, L., Monaghan, A. J., Lozano-Fuentes, S., Steinhoff, D. F., Hayden, M. H., and
745 Bieringer, P. E. (2014). The impact of temperature on the bionomics of *Aedes* (*Ste-*
746 *gomyia*) *aegypti* , with special reference to the cool geographic range margins. *Journal*
747 *of medical entomology*, 51(3):496–516.
- 748 Erguler, K. (2018). sPop: Age-structured discrete-time population dynamics model in C,
749 Python, and R. *F1000Research*, 7:1220.
- 750 Erguler, K., Chandra, N. L., Proestos, Y., Lelieveld, J., Christophides, G. K., and Parham,
751 P. E. (2017). A large-scale stochastic spatiotemporal model for *Aedes albopictus*-borne
752 chikungunya epidemiology. *PloS one*, 12(3):e0174293.
- 753 Erguler, K., Smith-Unna, S. E., Waldock, J., Proestos, Y., Christophides, G. K., Lelieveld,
754 J., and Parham, P. E. (2016). Large-scale modelling of the environmentally-driven pop-
755 ulation dynamics of temperate *Aedes albopictus* (skuse). *PloS one*, 11(2):e0149282.
- 756 Eritja, R., Palmer, J. R., Roiz, D., Sanpera-Calbet, I., and Bartumeus, F. (2017). Direct
757 evidence of adult *Aedes albopictus* dispersal by car. *Scientific Reports*, 7(1):1–15.

- 758 Eritja, R., Ruiz-Arrondo, I., Delacour-Estrella, S., Schaffner, F., Álvarez-Chachero, J.,
759 Bengoa, M., Puig, M.-Á., Melero-Alcázar, R., Oltra, A., and Bartumeus, F. (2019).
760 First detection of *Aedes japonicus* in Spain: an unexpected finding triggered by citizen
761 science. *Parasites & vectors*, 12(1):1–9.
- 762 Focks, D., Haile, D., Daniels, E., and Mount, G. (1993a). Dynamic life table model
763 for *Aedes aegypti* (Diptera: Culicidae): simulation results and validation. *Journal of*
764 *medical entomology*, 30(6):1018–1028.
- 765 Focks, D. A., Haile, D., Daniels, E., and Mount, G. A. (1993b). Dynamic life table model
766 for *Aedes aegypti* (Diptera: Culicidae): analysis of the literature and model develop-
767 ment. *Journal of medical entomology*, 30(6):1003–1017.
- 768 Fujioka, K. (2012). Discovery of *Aedes albopictus* (Skuse) in the city of El Monte and the
769 initial response. *Proc. Pap. Mosq. Vect. Control Assoc. Calif.*, 80:27–29.
- 770 GDAL/OGR contributors (2021). *GDAL/OGR Geospatial Data Abstraction software Li-*
771 *brary*. Open Source Geospatial Foundation.
- 772 Gilpin, M. E. and McClelland, G. A. (1979). Systems analysis of the yellow fever
773 mosquito *Aedes aegypti*. *Fortschritte Der Zoologie*, 25(2-3):355–388.
- 774 Giron, S., Franke, F., Decoppet, A., Cadiou, B., Travaglini, T., Thirion, L., Durand, G.,
775 Jeannin, C., L’ambert, G., Grard, G., et al. (2019). Vector-borne transmission of Zika
776 virus in Europe, southern France, August 2019. *Eurosurveillance*, 24(45):1900655.
- 777 Gloria-Soria, A., Brown, J. E., Kramer, V., Yoshimizu, M. H., and Powell, J. R. (2014).
778 Origin of the Dengue Fever mosquito, *Aedes aegypti*, in California. *PLOS Neglected*
779 *Tropical Diseases*, 8(7):e3029.
- 780 Gloria-Soria, A., Lima, A., Lovin, D. D., Cunningham, J. M., Severson, D. W., and Powell,
781 J. R. (2018). Origin of a High-Latitude Population of *Aedes aegypti* in Washington, DC.
782 *The American Journal of Tropical Medicine and Hygiene*, 98(2):445–452.
- 783 Gradoni, F., Bertola, M., Carlin, S., Accordi, S., Toniolo, F., Visentin, P., Patregnani, T.,
784 Adami, S., Terzo, L., Dal Pont, M., et al. (2021). Geographical data on the occurrence
785 and spreading of invasive *Aedes* mosquito species in Northeast Italy. *Data in Brief*,
786 36:107047.
- 787 Gratz, N. G. (2004). Critical review of the vector status of *Aedes albopictus*. *Medical and*
788 *veterinary entomology*, 18(3):215–227. 00498 PMID: 15347388.

- 789 Guerra, C. A., Reiner, R. C., Perkins, T. A., Lindsay, S. W., Midega, J. T., Brady, O. J.,
790 Barker, C. M., Reisen, W. K., Harrington, L. C., Takken, W., Kitron, U., Lloyd, A. L.,
791 Hay, S. I., Scott, T. W., and Smith, D. L. (2014). A global assembly of adult fe-
792 male mosquito mark-release-recapture data to inform the control of mosquito-borne
793 pathogens. *Parasites & Vectors*, 7:276.
- 794 Guzzetta, G., Trentini, F., Poletti, P., Baldacchino, F. A., Montarsi, F., Capelli, G., Rizzoli,
795 A., Rosà, R., Merler, S., and Melegaro, A. (2017). Effectiveness and economic as-
796 sessment of routine larviciding for prevention of chikungunya and dengue in temperate
797 urban settings in Europe. *PLoS neglected tropical diseases*, 11(9):e0005918.
- 798 Hancock, P. A., Linley-White, V., Callahan, A. G., Godfray, H. C. J., Hoffmann, A. A.,
799 and Ritchie, S. A. (2016). Density-dependent population dynamics in *Aedes aegypti*
800 slow the spread of wMel *Wolbachia*. *Journal of Applied Ecology*, 53(3):785–793.
- 801 Harrington, L. C., Edman, J. D., and Scott, T. W. (2001). Why do female *Aedes aegypti*
802 (Diptera: Culicidae) feed preferentially and frequently on human blood? *Journal of*
803 *Medical Entomology*, 38(3):411–422.
- 804 Hartemink, N., Vanwambeke, S. O., Purse, B. V., Gilbert, M., and Van Dyck, H. (2015).
805 Towards a resource-based habitat approach for spatial modelling of vector-borne disease
806 risks. *Biological Reviews*, 90(4):1151–1162.
- 807 Hawley, W. A. (1988). The biology of *Aedes albopictus*. *Journal of the American Mosquito*
808 *Control Association. Supplement*, 1:1–39.
- 809 Hawley, W. A., Pumpuni, C. B., Brady, R. H., and Craig, G. B. (1989). Overwintering
810 Survival of *Aedes albopictus* (Diptera: Culicidae) Eggs in Indiana. *Journal of Medical*
811 *Entomology*, 26(2):122–129.
- 812 Hopkins, M. C., Zink, S. D., Paulson, S. L., and Hawley, D. M. (2020). Influence of forest
813 disturbance on La Crosse virus risk in Southwestern Virginia. *Insects*, 11(1):28.
- 814 Horváth, C., Cazan, C. D., and Mihalca, A. D. (2021). Emergence of the invasive Asian
815 bush mosquito, *Aedes (Finlaya) japonicus japonicus*, in an urban area, Romania. *Para-*
816 *sites & vectors*, 14(1):1–7.
- 817 Hurk, A. F. v. d., McElroy, K., Pyke, A. T., McGee, C. E., Hall-Mendelin, S., Day, A.,
818 Ryan, P. A., Ritchie, S. A., Vanlandingham, D. L., and Higgs, S. (2011). Vector Com-
819 petence of Australian Mosquitoes for Yellow Fever Virus. *The American Journal of*
820 *Tropical Medicine and Hygiene*, 85(3):446–451. Publisher: The American Society of
821 Tropical Medicine and Hygiene.

- Hutchinson, G. (1957). Concluding remarks cold spring harbor symposia on quantitative biology, 22: 415–427. *GS SEARCH*.
- Iwamura, T., Guzman-Holst, A., and Murray, K. A. (2020). Accelerating invasion potential of disease vector *Aedes aegypti* under climate change. *Nature Communications*, 11(1):1–10.
- Jansen, S., Heitmann, A., Lühken, R., Jöst, H., Helms, M., Vapalahti, O., Schmidt-Chanasit, J., and Tannich, E. (2018). Experimental transmission of Zika virus by *Aedes japonicus japonicus* from southwestern Germany. *Emerging microbes & infections*, 7(1):1–6.
- Jelinski, D. E. and Wu, J. (1996). The modifiable areal unit problem and implications for landscape ecology. *Landscape ecology*, 11(3):129–140.
- Juliano, S. A. (2007). Population Dynamics. *Journal of the American Mosquito Control Association*, 23(2 Suppl):265–275.
- Juliano, S. A., O’Meara, G. F., Morrill, J. R., and Cutwa, M. M. (2002). Desiccation and thermal tolerance of eggs and the coexistence of competing mosquitoes. *Oecologia*, 130(3):458–469.
- Kattge, J., Bönsch, G., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., Tautenhahn, S., Werner, G. D. A., Aakala, T., Abedi, M., and Acosta, A. T. R. (2020). TRY plant trait database – enhanced coverage and open access. *Global Change Biology*, 26(1):119–188.
- Kaufman, M. G. and Fonseca, D. M. (2014). Invasion biology of *Aedes japonicus japonicus* (Diptera: Culicidae). *Annual Review of Entomology*, 59:31–49.
- Kearney, M. (2006). Habitat, environment and niche: what are we modelling? *Oikos*, 115(1):186–191.
- Kearney, M. and Porter, W. (2009). Mechanistic niche modelling: combining physiological and spatial data to predict species’ ranges. *Ecology letters*, 12(4):334–350.
- Kettle, H. and Nutter, D. (2015). stagePop: modelling stage-structured populations in R. *Methods in Ecology and Evolution*, 6(12):1484–1490. _eprint: <https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/2041-210X.12445>.
- Kilpatrick, A. M., Kramer, L. D., Campbell, S. R., Alleyne, E. O., Dobson, A. P., and Daszak, P. (2005). West Nile Virus risk assessment and the bridge vector paradigm. 11(3):425–429.

- 854 Koch, L. K., Cunze, S., Werblow, A., Kochmann, J., Dörge, D. D., Mehlhorn, H., and
855 Klimpel, S. (2016). Modeling the habitat suitability for the arbovirus vector *Aedes*
856 *albopictus* (Diptera: Culicidae) in Germany. *Parasitology research*, 115(3):957–964.
- 857 Koenraadt, C. J. M. and Harrington, L. C. (2008). Flushing effect of rain on container-
858 inhabiting mosquitoes *Aedes aegypti* and *Culex pipiens* (Diptera: Culicidae). 45(1):28–
859 35.
- 860 Kraemer, M. U., Reiner, R. C., Brady, O. J., Messina, J. P., Gilbert, M., Pigott, D. M., Yi,
861 D., Johnson, K., Earl, L., Marczak, L. B., et al. (2019). Past and future spread of the
862 arbovirus vect*Aedes aegypti* and *Aedes albopictus*. *Nature microbiology*, 4(5):854–863.
- 863 Kraemer, M. U., Sinka, M. E., Duda, K. A., Mylne, A., Shearer, F. M., Brady, O. J.,
864 Messina, J. P., Barker, C. M., Moore, C. G., Carvalho, R. G., et al. (2015). The global
865 compendium of *Aedes aegypti* and *Ae. albopictus* occurrence. *Scientific data*, 2(1):1–8.
- 866 Kramer, I. M., Kreß, A., Klingelhöfer, D., Scherer, C., Phuyal, P., Kuch, U., Ahrens, B.,
867 Groneberg, D. A., Dhimal, M., and Müller, R. (2020). Does winter cold really limit the
868 dengue vector *Aedes aegypti* in Europe? *Parasites & vectors*, 13(1):1–13.
- 869 Kramer, I. M., Pfeiffer, M., Steffens, O., Schneider, F., Gerger, V., Phuyal, P., Braun,
870 M., Magdeburg, A., Ahrens, B., Groneberg, D. A., et al. (2021). The ecophysiological
871 plasticity of *Aedes aegypti* and *Aedes albopictus* concerning overwintering in cooler
872 ecoregions is driven by local climate and acclimation capacity. *Science of The Total*
873 *Environment*, 778:146128.
- 874 Krupa, E., Henon, N., and Mathieu, B. (2021). Diapause characterisation and seasonality
875 of *Aedes japonicus japonicus* (Diptera, Culicidae) in the Northeast of France. *Parasite*,
876 28.
- 877 Kurucz, K., Manica, M., Delucchi, L., Kemenesi, G., and Marini, G. (2020). Dynamics
878 and distribution of the invasive mosquito *Aedes koreicus* in a temperate European city.
879 *International journal of environmental research and public health*, 17(8):2728.
- 880 Lacour, G., Chanaud, L., L’Ambert, G., and Hance, T. (2015). Seasonal synchronization of
881 diapause phases in *Aedes albopictus* (Diptera: Culicidae). *PloS one*, 10(12):e0145311.
- 882 Lega, J., Brown, H. E., and Barrera, R. (2017). *Aedes aegypti* (Diptera: Culicidae) abun-
883 dance model improved with relative humidity and precipitation-driven egg hatching.
884 *Journal of medical entomology*, 54(5):1375–1384.

- 885 Leta, S., Beyene, T. J., De Clercq, E. M., Amenu, K., Kraemer, M. U., and Revie, C. W.
886 (2018). Global risk mapping for major diseases transmitted by *Aedes aegypti* and *Aedes*
887 *albopictus*. *International Journal of Infectious Diseases*, 67:25–35.
- 888 Liu-Helmersson, J., Rocklöv, J., Sewe, M., and Brännström, Å. (2019). Climate change
889 may enable *Aedes aegypti* infestation in major european cities by 2100. *Environmental*
890 *research*, 172:693–699.
- 891 Lounibos, L. P. and Kramer, L. D. (2016). Invasiveness of *Aedes aegypti* and *Aedes al-*
892 *bopictus* and vectorial capacity for chikungunya virus. *The Journal of infectious dis-*
893 *eases*, 214(suppl_5):S453–S458.
- 894 Maclean, I. M., Mosedale, J. R., and Bennie, J. J. (2019). Microclima: An R package for
895 modelling meso-and microclimate. *Methods in Ecology and Evolution*, 10(2):280–290.
- 896 Magori, K., Legros, M., Puente, M. E., Focks, D. A., Scott, T. W., Lloyd, A. L., and Gould,
897 F. (2009). Skeeter Buster: A Stochastic, Spatially Explicit Modeling Tool for Studying
898 *Aedes aegypti* Population Replacement and Population Suppression Strategies. *PLOS*
899 *Neglected Tropical Diseases*, 3(9):e508.
- 900 Marcantonio, M., Metz, M., Baldacchino, F., Arnoldi, D., Montarsi, F., Capelli, G., Carlin,
901 S., Neteler, M., and Rizzoli, A. (2016). First assessment of potential distribution and
902 dispersal capacity of the emerging invasive mosquito *Aedes koreicus* in Northeast Italy.
903 *Parasites & Vectors*, 9(1):63.
- 904 Marcantonio, M., Reyes, T., and Barker, C. M. (2019). Quantifying *Aedes aegypti* disper-
905 sal in space and time: a modeling approach. *Ecosphere*, 10(12):e02977.
- 906 Marini, F., Caputo, B., Pombi, M., Travaglio, M., Montarsi, F., Drago, A., Rosà, R.,
907 Manica, M., and Torre, A. d. (2019a). Estimating Spatio-Temporal Dynamics of *Aedes*
908 *albopictus* Dispersal to Guide Control Interventions in Case of Exotic Arboviruses in
909 Temperate Regions. *Scientific Reports*, 9(1):1–9.
- 910 Marini, G., Arnoldi, D., Baldacchino, F., Capelli, G., Guzzetta, G., Merler, S., Montarsi,
911 F., Rizzoli, A., and Rosà, R. (2019b). First report of the influence of temperature on the
912 bionomics and population dynamics of *Aedes koreicus*, a new invasive alien species in
913 Europe. *Parasites & vectors*, 12(1):1–12.
- 914 Marini, G., Manica, M., Arnoldi, D., Inama, E., Rosà, R., and Rizzoli, A. (2020). Influence
915 of temperature on the life-cycle dynamics of *Aedes albopictus* population established at
916 temperate latitudes: A laboratory experiment. *Insects*, 11(11):808.

- 917 Martinet, J.-P., Ferté, H., Failloux, A.-B., Schaffner, F., and Depaquit, J. (2019).
918 Mosquitoes of north-western Europe as potential vectors of arboviruses: a review.
919 *Viruses*, 11(11):1059.
- 920 Mathewson, P. D., Moyer-Horner, L., Beever, E. A., Briscoe, N. J., Kearney, M., Yahn,
921 J. M., and Porter, W. P. (2017). Mechanistic variables can enhance predictive models
922 of endotherm distributions: the American pika under current, past, and future climates.
923 *Global Change Biology*, 23(3):1048–1064.
- 924 McBride, C. S., Baier, F., Omondi, A. B., Spitzer, S. A., Lutomiah, J., Sang, R., Ignell, R.,
925 and Vosshall, L. B. (2014a). Evolution of mosquito preference for humans linked to an
926 odorant receptor. *Nature*, 515(7526):222–227.
- 927 McBride, C. S., Baier, F., Omondi, A. B., Spitzer, S. A., Lutomiah, J., Sang, R., Ignell, R.,
928 and Vosshall, L. B. (2014b). Evolution of mosquito preference for humans linked to an
929 odorant receptor. *Nature*, 515(7526):222–227.
- 930 McKenzie, B. A., Wilson, A. E., and Zohdy, S. (2019). *Aedes albopictus* is a competent
931 vector of Zika virus: A meta-analysis. *PLOS ONE*, 14(5):e0216794. Publisher: Public
932 Library of Science.
- 933 Metelmann, S., Caminade, C., Jones, A. E., Medlock, J. M., Baylis, M., and Morse, A. P.
934 (2019). The UK’s suitability for *Aedes albopictus* in current and future climates. *Journal*
935 *of The Royal Society Interface*, 16(152):20180761. Publisher: Royal Society.
- 936 Montarsi, F., Ciocchetta, S., Devine, G., Ravagnan, S., Mutinelli, F., Frangipane Di Re-
937 galbono, A., Otranto, D., and Capelli, G. (2015a). Development of *Dirofilaria immitis*
938 within the mosquito *Aedes (Finlaya) koreicus*, a new invasive species for Europe. *Par-*
939 *asites and Vectors*, 8(1).
- 940 Montarsi, F., Drago, A., Dal Pont, M., Delai, N., Carlin, S., Cazzin, S., Ciocchetta, S.,
941 Arnoldi, D., Baldacchino, F., Rizzoli, A., et al. (2014). Current knowledge on the
942 distribution and biology of the recently introduced invasive mosquito *Aedes koreicus*
943 (Diptera: Culicidae). *Firenze (Italy): Atti Accademia Nazionale Italiana di Entomolo-*
944 *gia*, 62:169–174.
- 945 Montarsi, F., Drago, A., Martini, S., Calzolari, M., De Filippo, F., Bianchi, A., Mazzucato,
946 M., Ciocchetta, S., Arnoldi, D., Baldacchino, F., et al. (2015b). Current distribution of
947 the invasive mosquito species, *Aedes koreicus* [*Hulecoeteomyia koreica*] in Northern
948 Italy. *Parasites & vectors*, 8(1):1–5.

- 949 Montarsi, F., Martini, S., Dal Pont, M., Delai, N., Milone, N. F., Mazzucato, M., Sop-
950 pelsa, F., Cazzola, L., Cazzin, S., Ravagnan, S., et al. (2013). Distribution and habitat
951 characterization of the recently introduced invasive mosquito *Aedes koreicus* [*Hulecoe-*
952 *teomyia koreica*], a new potential vector and pest in north-eastern Italy. *Parasites &*
953 *vectors*, 6(1):1–10.
- 954 Müller, P., Engeler, L., Vavassori, L., Suter, T., Guidi, V., Gschwind, M., Tonolla, M.,
955 and Flacio, E. (2020). Surveillance of invasive *Aedes* mosquitoes along Swiss traffic
956 axes reveals different dispersal modes for *Aedes albopictus* and *Ae. japonicus*. *PLoS*
957 *neglected tropical diseases*, 14(9):e0008705.
- 958 Müller, R., Knautz, T., Vollroth, S., Berger, R., Kreß, A., Reuss, F., Groneberg, D. A.,
959 and Kuch, U. (2018). Larval superiority of *Culex pipiens* to *Aedes albopictus* in a
960 replacement series experiment: prospects for coexistence in Germany. *Parasites &*
961 *vectors*, 11(1):1–16.
- 962 Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo,
963 G., Boussetta, S., Choulga, M., Harrigan, S., Hersbach, H., et al. (2021). ERA5-Land:
964 A state-of-the-art global reanalysis dataset for land applications. *Earth System Science*
965 *Data Discussions*, pages 1–50.
- 966 Näslund, J., Ahlm, C., Islam, K., Evander, M., Bucht, G., and Lwande, O. W. (2021).
967 Emerging mosquito-borne viruses linked to *Aedes aegypti* and *Aedes albopictus*: Global
968 status and preventive strategies. *Vector-Borne and Zoonotic Diseases*, 21(10):731–746.
- 969 Negri, A., Arnoldi, I., Brilli, M., Bandi, C., Gabrieli, P., and Epis, S. (2021). Evidence for
970 the spread of the alien species *Aedes koreicus* in the Lombardy region, Italy. *Parasites*
971 *& Vectors*, 14(1):1–6.
- 972 Oliveira, S., Rocha, J., Sousa, C. A., and Capinha, C. (2021). Wide and increasing suit-
973 ability for *Aedes albopictus* in Europe is congruent across distribution models. *Scientific*
974 *reports*, 11(1):1–9.
- 975 Onofri, A. (2020). *aomisc: Statistical methods for the agricultural sciences*. R package
976 version 0.64.
- 977 Otero, M., Solari, H. G., and Schweigmann, N. (2006). A stochastic population dynamics
978 model for *Aedes aegypti*: formulation and application to a city with temperate climate.
979 *Bulletin of mathematical biology*, 68(8):1945–1974.

- 980 Pasaoglu, G., Fiorello, D., Martino, A., Scarcella, G., Alemanno, A., Zubaryeva, A., and
981 Thiel, C. (2012). Driving and parking patterns of European car drivers-a mobility sur-
982 vey. *Luxembourg: European Commission Joint Research Centre*.
- 983 Pasquali, S., Mariani, L., Calvitti, M., Moretti, R., Ponti, L., Chiari, M., Sperandio, G.,
984 and Gilioli, G. (2020). Development and calibration of a model for the potential estab-
985 lishment and impact of *Aedes albopictus* in Europe. *Acta tropica*, 202:105228.
- 986 Peterson, A. T. (2014). *Mapping disease transmission risk: enriching models using bio-*
987 *geography and ecology*. JHU Press.
- 988 Petrić, M., Ducheyne, E., Gossner, C. M., Marsboom, C., Nicolas, G., Venail, R., Hen-
989 drickx, G., and Schaffner, F. (2021). Seasonality and timing of peak abundance of *Aedes*
990 *albopictus* in Europe: Implications to public and animal health. *Geospatial Health*,
991 16(1).
- 992 Powell, J. R., Tabachnick, W. J., Powell, J. R., and Tabachnick, W. J. (2013). History of
993 domestication and spread of *Aedes aegypti* - A Review. *Memórias do Instituto Oswaldo*
994 *Cruz*, 108:11–17.
- 995 Pumpuni, C. B., Knepler, J., and Craig, G. B. (1992). Influence of temperature and larval
996 nutrition on the diapause inducing photoperiod of *Aedes albopictus*. *Journal of the*
997 *American Mosquito Control Association*, 8(3):223–227.
- 998 R Core Team (2021). *R: A Language and Environment for Statistical Computing*. R
999 Foundation for Statistical Computing, Vienna, Austria.
- 1000 Reiskind, M. and Lounibos, L. (2013). Spatial and temporal patterns of abundance of
1001 *Aedes aegypti* l.(*Stegomyia aegypti*) and *Aedes albopictus* (skuse)[*Stegomyia albopictus*
1002 (*Skuse*)] in southern florida. *Medical and veterinary entomology*, 27(4):421–429.
- 1003 Reuss, F., Wieser, A., Niamir, A., Bálint, M., Kuch, U., Pfenninger, M., and Müller, R.
1004 (2018). Thermal experiments with the asian bush mosquito (*Aedes japonicus japoni-*
1005 *cus*)(Diptera: Culicidae) and implications for its distribution in Germany. *Parasites &*
1006 *vectors*, 11(1):1–10.
- 1007 Rezza, G., Nicoletti, L., Angelini, R., Romi, R., Finarelli, A. C., Panning, M., Cordioli,
1008 P., Fortuna, C., Boros, S., Magurano, F., Silvi, G., Angelini, P., Dottori, M., Ciufolini,
1009 M. G., Majori, G. C., Cassone, A., and CHIKV study group (2007). Infection with
1010 Chikungunya virus in Italy: an outbreak in a temperate region. *Lancet*, 370(9602):1840–
1011 1846.

- 1012 Ritz, C., Baty, F., Streibig, J. C., and Gerhard, D. (2015). Dose-response analysis using r.
1013 *PLOS ONE*, 10(e0146021).
- 1014 Roche, B., Léger, L., L' Ambert, G., Lacour, G., Foussadier, R., Besnard, G., Barré-Cardi,
1015 H., Simard, F., and Fontenille, D. (2015). The spread of *Aedes albopictus* in metropoli-
1016 tan France: contribution of environmental drivers and human activities and predictions
1017 for a near future. *PloS one*, 10(5):e0125600.
- 1018 Rohatgi, A. (2020). *Webplotdigitizer: Version 4.4*.
- 1019 Roiz, D., Boussès, P., Simard, F., Paupy, C., and Fontenille, D. (2015). Autochthonous
1020 chikungunya transmission and extreme climate events in southern France. *PLoS Ne-*
1021 *glected Tropical Diseases*, 9(6):e0003854.
- 1022 Schaffner, F., Karch, S., et al. (2000). First record of *Aedes albopictus* (Skuse, 1894) in
1023 metropolitan France. *COMPTES RENDUS-ACADEMIE DES SCIENCES PARIS SERIE*
1024 *3*, 323(4):373–376.
- 1025 Schaffner, F., Vazeille, M., Kaufmann, C., Failloux, A.-B., and Mathis, A. (2011). Vec-
1026 tor competence of *Aedes japonicus* for chikungunya and dengue viruses. *European*
1027 *Mosquito Bulletin*, (29):141–142.
- 1028 Schulzweida, U. (2019). Cdo user guide.
- 1029 Scott, J. (2003). *The Ecology of the Exotic Mosquito Ochlerotatus (Finlaya) japonicus*
1030 *japonicus (Theobald 1901) (Diptera: Culicidae) and an Examination of its Role in the*
1031 *West Nile Virus Cycle in New Jersey*. Ph.D. thesis, Rutgers University, New Brunswick,
1032 NJ.
- 1033 Seidel, B., Montarsi, F., Huemer, H. P., Indra, A., Capelli, G., Allerberger, F., and
1034 Nowotny, N. (2016). First record of the Asian bush mosquito, *Aedes japonicus japon-*
1035 *icus*, in Italy: invasion from an established Austrian population. *Parasites & vectors*,
1036 9(1):1–4.
- 1037 Sherpa, S., Blum, M. G., Capblancq, T., Cumer, T., Rioux, D., and Després, L. (2019).
1038 Unravelling the invasion history of the Asian tiger mosquito in Europe. *Molecular*
1039 *Ecology*, 28(9):2360–2377.
- 1040 Souza-Neto, J. A., Powell, J. R., and Bonizzoni, M. (2019). *Aedes aegypti* vector compe-
1041 tence studies: A review. *Infection, Genetics and Evolution*, 67:191–209.
- 1042 Stallman, R. (1985). The GNU Manifesto. *j-DDJ*, 10(3):30–35.

- 1043 Takashima, I. and Rosen, L. (1989). Horizontal and vertical transmission of Japanese
1044 encephalitis virus by *Aedes japonicus* (Diptera: Culicidae). *Journal of Medical Ento-*
1045 *mology*, 26(5):454–458.
- 1046 Takken, W. and van den Berg, H. (2019). Manual on prevention of establishment and
1047 control of mosquitoes of public health importance in the WHO European region (with
1048 special reference to invasive mosquitoes).
- 1049 Tanaka, K., Mizusawa, K., and Saugstad, E. S. (1979a). A revision of the adult and larval
1050 mosquitoes of Japan (including the Ryukyu archipelago and the Ogasawara islands) and
1051 Korea (Diptera: Culicidae). Technical report, ARMY MEDICAL LAB PACIFIC APO
1052 SAN FRANCISCO 96343.
- 1053 Tanaka, T., Mizusawa, K., and Saugstad, E. (1979b). *Mosquitoes of Japan and Korea*,
1054 volume 16 of *Contributions of the American Entomological Institute*. the American
1055 Entomological Institute.
- 1056 Thomas, S. M., Obermayr, U., Fischer, D., Kreyling, J., and Beierkuhnlein, C. (2012).
1057 Low-temperature threshold for egg survival of a post-diapause and non-diapause Eu-
1058 ropean aedine strain, *Aedes albopictus* (Diptera: Culicidae). *Parasites & Vectors*,
1059 5(1):100.
- 1060 Thornton, M., Shrestha, R., Wei, Y., Thornton, P., Kao, S., and Wilson, B. (2020). Daymet-
1061 Daymet: Daily Surface Weather Data on a 1-km Grid for North America, Version 4.
1062 Artwork Size: 0 MB Medium: netCDF Publisher: ORNL Distributed Active Archive
1063 Center Version Number: 4.
- 1064 Tran, A., l’Ambert, G., Lacour, G., Benoît, R., Demarchi, M., Cros, M., Cailly, P., Aubry-
1065 Kientz, M., Balenghien, T., and Ezanno, P. (2013). A rainfall-and temperature-driven
1066 abundance model for *Aedes albopictus* populations. *International journal of environ-*
1067 *mental research and public health*, 10(5):1698–1719.
- 1068 Tran, A., Mangeas, M., Demarchi, M., Roux, E., Degenne, P., Haramboure, M., Le Goff,
1069 G., Damiens, D., Gouagna, L.-C., Herbreteau, V., et al. (2020). Complementarity of
1070 empirical and process-based approaches to modelling mosquito population dynamics
1071 with *Aedes albopictus* as an example—application to the development of an operational
1072 mapping tool of vector populations. *PloS one*, 15(1):e0227407.
- 1073 Tripet, F., Lounibos, L. P., Robbins, D., Moran, J., Nishimura, N., and Blosser, E. M.
1074 (2011). Competitive reduction by satyrization? evidence for interspecific mating in

1075 nature and asymmetric reproductive competition between invasive mosquito vectors.
1076 *The American journal of tropical medicine and hygiene*, 85(2):265–270.

1077 Tsigkinopoulou, A., Hawari, A., Uttley, M., and Breitling, R. (2018). Defining informative
1078 priors for ensemble modeling in systems biology. *Nature Protocols*, 13(11):2643.

1079 Turell, M. J., Beaman, J. R., and Neely, G. W. (1994). Experimental transmission of
1080 eastern equine encephalitis virus by strains of *Aedes albopictus* and *A. taeniorhynchus*
1081 (Diptera: Culicidae). *Journal of Medical Entomology*, 31(2):287–290.

1082 Urbanski, J., Mogi, M., O'Donnell, D., DeCotiis, M., Toma, T., and Armbruster, P. (2012).
1083 Rapid adaptive evolution of photoperiodic response during invasion and range expan-
1084 sion across a climatic gradient. *The American Naturalist*, 179(4):490–500.

1085 Venturi, G., Luca, M. D., Fortuna, C., Remoli, M. E., Riccardo, F., Severini, F., Toma,
1086 L., Manso, M. D., Benedetti, E., Caporali, M. G., Amendola, A., Fiorentini, C., Lib-
1087 erato, C. D., Giammattei, R., Romi, R., Pezzotti, P., Rezza, G., and Rizzo, C. (2017).
1088 Detection of a chikungunya outbreak in Central Italy, August to September 2017. *Eu-
1089 rosurveillance*, 22(39):17–00646.

1090 Veronesi, E., Paslaru, A., Silaghi, C., Tobler, K., Glavinic, U., Torgerson, P., and Mathis,
1091 A. (2018). Experimental evaluation of infection, dissemination, and transmission rates
1092 for two West Nile virus strains in European *Aedes japonicus* under a fluctuating temper-
1093 ature regime. *Parasitology research*, 117(6):1925–1932.

1094 Versteirt, V., De Clercq, E. M., Fonseca, D. M., Pecor, J., Schaffner, F., Coosemans, M.,
1095 and Van Bortel, W. (2012). Bionomics of the Established Exotic Mosquito Species
1096 *Aedes koreicus* in Belgium, Europe. *Journal of Medical Entomology*, 49(6):1226–1232.

1097 Versteirt, V., Schaffner, F., Garros, C., Dekoninck, W., Coosemans, M., and Van Bor-
1098 tel, W. (2009). Introduction and establishment of the exotic mosquito species *Aedes*
1099 *japonicus japonicus* (Diptera: Culicidae) in Belgium. *Journal of Medical Entomology*,
1100 46(6):1464–1467.

1101 Waldock, J., Chandra, N. L., Lelieveld, J., Proestos, Y., Michael, E., Christophides, G.,
1102 and Parham, P. E. (2013). The role of environmental variables on *Aedes albopictus*
1103 biology and chikungunya epidemiology. *Pathogens and global health*, 107(5):224–241.

1104 Watson, M. S. (1967). *Aedes (Stegomyia) albopictus* (Skuse): A Literature Review. *Mis-
1105 cellaneous publication*, 22:42.

- 1106 Werner, D., Zielke, D., and Kampen, H. (2016). First record of *Aedes koreicus* (Diptera:
1107 Culicidae) in Germany. *Parasitology Research*, 115(3):1331–1334.
- 1108 Westby, K. M., Fritzen, C., Paulsen, D., Poindexter, S., and Moncayo, A. C. (2015). La
1109 Crosse encephalitis virus infection in field-collected *Aedes albopictus*, *Aedes japoni-*
1110 *cus*, and *Aedes triseriatus* in Tennessee. *Journal of the American Mosquito Control*
1111 *Association*, 31(3):233–241.
- 1112 Wieser, A., Reuss, F., Niamir, A., Müller, R., O’Hara, R. B., and Pfenninger, M. (2019).
1113 Modelling seasonal dynamics, population stability, and pest control in *Aedes japonicus*
1114 *japonicus* (Diptera: Culicidae). *Parasites & vectors*, 12(1):1–12.

1115 **A Review of mechanistic model for invasive *Aedes***

Name	Year	Species	Biological resolution	Time	Space	Programming language	Reference
SkeeterBooster	2009	<i>Ae. aegypti</i>	Agent-based			C++	Magori et al. (2009)
stagePop	2009		Process-based			R	Kettle and Nutter (2015)
sPop	2018					C++, R, Python	Erguler (2018)
albopictus	2019	<i>Ae. albopictus</i>	Process-based			C++, R, Python 4	Erguler et al. (2016, 2017)
	2019	<i>Ae. albopictus</i>	Process-based			Octave v4.2.1, Runge-Kutta 4	Metelmann et al. (2019)
	2019	<i>Ae. albopictus</i>	Process-based			R, Ocelet	Tran et al. (2013, 2020)
	2020	<i>Ae. aegypti</i>	Process-based			R	Iwamura et al. (2020)
	2020	<i>Ae. koreicus</i>	Process-based			R	Kurucz et al. (2020)

Table 2: Description of mechanistic codes for invasive *Aedes* made available as software or scripts.

1116 **B Parameters used in dynamAedes**

Species	Stage	Parameters	Main source	data	Origin	Value	Comments
<i>Ae. aegypti</i> and <i>Ae. al- bopictus</i>	Eggs	min number of days for eggs development	Christophers 1960; Delatte et al 2009		Literature therein; La Reunion (France)	4	
<i>Ae. aegypti</i> and <i>Ae. al- bopictus</i>	Juveniles	min number of days for immature development	Christophers 1960; Delatte et al 2009		Literature therein; La Reunion (France)	6	
<i>Ae. aegypti</i> and <i>Ae. al- bopictus</i>	Adults	min number of days for gonotrophic cycle	Christophers 1960; Delatte et al 2009		Literature therein; La Reunion (France)	4	
<i>Ae. koreicus</i> and <i>Ae. japon- icus</i>	Eggs	min number of days for eggs development	Dr Daniele Arnoldi unpublished results for <i>Ae. koreicus</i>		Trento (NE Italy)	8	
<i>Ae. koreicus</i> and <i>Ae. japon- icus</i>	Juveniles	min number of days for immature development	Dr Daniele Arnoldi unpublished results for <i>Ae. koreicus</i> ; Wieser et al. (2019)		Trento (NE Italy)	9	min 10 days for <i>Ae. koreicus</i> , min 8 days for <i>Ae. japonicus</i> . Assumed 9 for both species
<i>Ae. koreicus</i> and <i>Ae. japon- icus</i>	Adults	min number of days for gonotrophic cycle	Assumed the same of <i>Ae. aegypti</i>		-	4	Since the probability of complete the gonothropic cycle is roughly 1/3 of <i>Ae. aegypti</i> , we kept the same leght the same lenght of <i>Ae. aegypti</i>
<i>All species</i>	Juveniles	Survival rate (density-dependent)	Hancock et al. 2016		Tropical (NE Australia)	Log-normal distribution	1-density dependant survival rate is additive to 1-temperature-dependent survival rate; the estimates made for <i>Ae. aegypti</i> were extended to the other three species
	Adults	Sex ratio 1:1	-		-	-	General modelling assumption widely taken in process-based modelling literature

Table 3: Other model features

Species	Stage	Model parameter	Data source	Origin	Function	Comments
<i>Ae. aegypti</i>	Egg	Survival rate	Thomas et al. 2012, Eisen et al. 2014	Laboratory colony(Tropical Asia origin); Literature review	Beta	Data for T>0 taken from Eisen 2014, data for T<0 taken from Thomas et al. 2012
		Hatching rate	Christophers 1960; Farnesi et al 2009; Mohammed and Chadee 2011	Literature therein; Rock-feller strain; Trinidad and Tobago	Beta	48h estimates from Mohammed and Chadee 2011 divided by two to get a daily rate. Additional data: 0 hatching a 7°C from Christophers and 0.025 at 12°C daily hatching from Farnesi et al. (2009)
	Juvenile	Survival rate (temperature-dependent)	Yang et al. 2009	NW of São Paulo State (Brazil)	Beta	Table 8 from Yang et al. 2009 Appendix I
		Emergence rate	Yang et al. 2009; Grech et al (2015)	NW of São Paulo State (Brazil); Cordoba (Argentina)	Beta	Data from Yang et al. (2009) refer to all immature stages, we corrected them to match only pupa using information present in Grech et al (2015) by taking the ratio between the minimum pupation time of the two experiment as a scaling factor
		Survival rate	Yang et al. 2009	NW of São Paulo State (Brazil)	Beta	Table 4
<i>Ae. albopictus</i>	Adult	Gonotrophic cycle	Yang et al. 2009	NW of São Paulo State (Brazil)	Beta	Table 5
		Oviposition: number of eggs	Christophers 1960; Yang et al. 2009	Literature therein; NW of São Paulo State (Brazil)	Beta	Table 5 from Yang et al (2009) rescaled using the average number of eggs/female reported in Christophers (1960)
	Egg	Survival rate (non-diapausing)	Metelmann et al. 2019; Expert based	Literature therein	nonlinear	Polynomial function taken from the SM manually adapted to get no survival at T>40°C
		Survival rate (diapausing)	Metelmann et al. 2019	Literature therein	nonlinear	Polynomial function taken from the SM
		Hatching rate ((non-diapausing)	Delatte et al. 2009	La Reunion (France)	Beta	Table 1; column Egg-L1
	Juvenile	Hatching rate (diapausing)	-	-	-	Assumed the same as for non-diapausing eggs
		Survival rate (temperature-dependent)	Metelmann et al. 2019	Literature therein	nonlinear	Polynomial function taken from the SM
		Emergence rate	Delatte et al. 2009	La Reunion (France)	Beta	Table 1; column Pupae-adult
	Adult	Survival rate	Metelmann et al. 2019	Literature therein	nonlinear	Polynomial function taken from the SM
		Gonotrophic cycle	Delatte et al. 2009	La Reunion (France)	Beta	Table 6

Species	Stage	Model parameter	Data source	Origin	Function	Comments
<i>Ae. japonicus</i>	Egg	Oviposition: number of eggs	Delatte et al. 2009	La Reunion (France)	Beta	Table 6
		Survival rate	Wieser et al. 2019	Biberach Germany	Beta	Observation taken from Fig. 1 in Wieser et al (2019)
		Hatch rate	Wieser et al. 2019	Biberach Germany	Beta	Reuss et al., unpublished. Hatching observed also below a thin ice layer (T<0)
	Juvenile	Survival rate	Wieser et al. 2019	Biberach Germany	Beta	Reuss et al., unpublished
		Emergence rate	Reuss et al. 2018 SM	Biberach Germany	Beta	Table S2; survival upper temperature limit was adapted by accounting for expert opinion.
	Adult	Survival rate	Reuss et al. 2018 SM	Biberach Germany	Beta	Fig. 2
		Gonotrophic cycle	-	-	Beta	No information available, assumed the same of <i>Ae. koreicus</i> due to their phylogenetic and biogeographic similarity
<i>Ae. koreicus</i>	Egg	Oviposition: number of eggs	Reuss et al. 2018	Biberach Germany	Beta	Number of eggs estimated by mean female wing length, using the formula taken from Armistead et al. (2009), their Fig. 5; we divided the estimated eggs per female per gonotrophic cycle by a factor of two, due the two ovipositing days we are accounting in the model.
		Survival rate	Marini et al. 2019; Expert Based	Trento (NE Italy)	nonlinear	Polynomial function F3 and parameters values taken from Tab. 5 in Marini et al (2019) adjusted using observations provided in Arnoldi et al., unpublished observations
		Hatching rate	Marini et al. 2019; Expert Based	Trento (NE Italy)	Beta	Table 1 scaled with unpublished data to account for non-embryonated eggs; Arnoldi et al., unpublished observations
	Juvenile	Survival rate (temperature-dependent)	Marini et al. 2019; Expert Based	Trento (NE Italy)	Beta	Table 1 averaged from instar 1 to 4; the the upper limit has been adapted accordingly to expert opinions by Ciocchetta et al., unpublished observations and Arnoldi et al., unpublished observations
		Emergence rate	Marini et al. 2019	Trento (NE Italy)	Beta	Table 2
	Adult	Survival rate	Marini et al. 2019; Expert Based	Trento (NE Italy)	Beta	Table 3
		Gonotrophic cycle	Marini et al. 2019	Trento (NE Italy)	Beta	Table 3
		Oviposition: number of eggs	Expert Based	Trento (NE Italy)	Beta	Arnoldi et al., unpublished observations

Table 4: Species-specific temperature-dependent physiological parameters

Species	Stage	Parameters	Main source	Origin	Distribution	Comments
<i>Ae. aegypti</i>	Adults	Active dispersal	Marcantonio et al. 2019	California (USA)	Log-normal	Table 3
<i>Ae. albopictus</i>	Adults	Active dispersal	Marini et al. 2019B	Rome (Italy)	Log-normal	Figure 2a; Data from the 3 MRR were fit with several distributions, best distribution chosen using AIC
<i>Ae. japonicus</i>	Adults	Active dispersal	-	-	Log-normal	No information available, assumed the same of <i>Ae. albopictus</i>
<i>Ae. koreicus</i>	Adults	Active dispersal	-	-	Log-normal	No information available, assumed the same of <i>Ae. albopictus</i>
All species	All species	Passive dispersal (average trip distance)	Pasaoglu et al. 2012	-	-	Trip distance weighted average for ITA, DEU, FRA, ESP, POL, UK. Data taken from Figure 4.4 Average trip distance (km) by trip purpose using WebPlotDigitalizer
All species	Adults	Passive dispersal (Hitchhiking probability)	Ertijia et al. 2017	Mediterranean (Catalonia, Spain)	gamma distribution	0.0051 probability of a female mosquito to enter a car; Assumed to be the same for all the species as estimated by Ertijia et al. 2017 for <i>Ae. albopictus</i>

Table 5: Species specific dispersal parameters

Species	Stage	Model parameter	Data source	Origin	Function	Comments
<i>Ae. aegypti</i>	Diapause Egg	Diapause incidence	Lacour et al. 2015	Ovitrap (Provence, Southern France)	Exponential	
<i>Ae. japonicus</i>	Diapause Egg	Diapause incidence	Krupa et al. 2021	Ovitrap (Bas-Rhin, Northeast France)	Exponential	
<i>Ae. kor-eicus</i>	Diapause Egg	Diapause incidence	Krupa et al. 2021	Ovitrap (Bas-Rhin, Northeast France)	Exponential	We used the same exponential function used for <i>Ae. japonicus</i> due to the close phylogenetic relationship between these two species

Table 6: Species specific photoperiod parameters

1117 **A** *Aedes sp.* response curve

Figure 7: Overview of the temperature-dependent functions used in the model for the four *Aedes* species

Figure 8: Overview of the temperature-dependent functions used in the model for *Ae. aegypti*

Figure 9: Overview of the temperature-dependent functions used in the model for *Ae. albopictus*

Figure 10: Overview of the temperature-dependent functions used in the model for *Ae. japonicus*

Figure 11: Overview of the temperature-dependent functions used in the model for *Ae. koreicus*

Figure 12: Overview of the photoperiod-dependent diapause incidence function used to in the model for *Ae. albopictus* and *Ae. japonicus*. The *Ae. japonicus* function was used for *Ae. koreicus* as well.

1118 **B Larval habitat water volume parameter sensitivity**

Figure 13: Sensitivity analysis on the effect of the $1hvw$ parameter on the estimated number of individuals. In this example, we used the temperature observations of the Nice weather station used in the case study and varied the amount of water volume. The seasonal trend remained the same but, as expected, the simulated number of individuals increase as the $1hvw$ parameter increase.

C *Aedes aegypti* and *Ae. albopictus* regional scale case study

Figure 14: Predicted percentage of establishment of *Ae. aegypti* *Ae. albopictus* in California (USA) for the years 2011-2016 and 2013-2018, respectively. Only pixels having a probability of successful introduction >0 are shown. The red dots represent the counties where the species have been found.

1121 **D** *Aedes koreicus* population dynamics punctual scale case 1122 **study**

Year	Month	CI 2.5%	CI 25%	CI 50%	CI 75%	CI 97.5%	Observed <i>Ae. Koreicus</i>
2016	May	0.00	1.45	5.26	13.74	31.22	2
2016	June	0.00	1.37	6.36	14.40	40.64	5
2016	July	0.00	2.32	11.85	32.62	101.76	3
2016	August	0.70	24.57	66.65	128.03	266.00	9
2016	September	0.18	14.52	37.37	88.27	175.66	37
2016	October	0.00	0.24	1.18	3.14	10.17	0
2017	May	0.02	0.51	2.67	7.85	18.44	9
2017	June	0.29	2.75	10.60	28.57	55.28	8
2017	July	3.67	10.91	30.62	85.53	186.93	10
2017	August	7.24	32.50	74.18	159.67	283.74	24
2017	September	0.61	8.79	31.48	88.08	156.47	2
2017	October	0.08	0.82	2.20	4.98	11.06	2
2017	November	0.00	0.00	0.00	0.00	0.14	0
2018	April	1.66	5.26	11.54	22.25	34.30	0
2018	May	1.73	3.22	5.73	9.18	13.21	1
2018	June	1.57	5.81	10.52	22.29	32.89	6
2018	July	13.74	29.99	48.04	74.34	98.01	7
2018	August	37.83	71.67	109.27	157.55	201.00	29
2018	September	54.55	82.03	112.57	147.58	179.09	24
2018	October	2.13	2.90	3.77	5.89	7.80	6
2018	November	0.00	0.00	0.00	0.08	0.15	0

Table 7: Model validation for *Aedes koreicus* model in Trento (NE Italy)