Title

Intrinsic apoptosis is evolutionary divergent among metazoans

Authors

Gabriel Krasovec ^{1,2,3*}, Éric Quéinnec ^{1**}, Jean-Philippe Chambon ^{2,4**}

* Authors for correspondence: GK, gabriel.krasovec@nuigalway.ie.

** These authors contributed equally to this work.

Affiliations

¹ISYEB, Institut de Systématique, Evolution et Biodiversité, Sorbonne Université, CNRS, MNHN, F-75252 Paris Cedex 05, France.

²Sorbonne Université, CNRS, Institut de Biologie Paris Seine, IBPS, Evolution Paris Seine, F-75252 Paris Cedex 05, France.

³Centre for Chromosome Biology, School of Natural Sciences, National University of Ireland Galway, Galway, Ireland.

⁴Centre de Recherche de Biologie Cellulaire de Montpellier (CRBM), Montpellier Univ., CNRS, 34000 Montpellier, France.

Summary

Apoptosis is characterised by an analogous set of morphological features¹ that depend on a proteolytic multigenic family, the caspases^{2,3}. Each apoptotic signalling pathway involves a specific initiator caspase, upstream of the pathway regulation, which finally converges to common executioner caspases. Intrinsic apoptosis, previously known as the mitochondrial apoptotic pathway, is often considered as ancestral and evolutionary conserved among animals^{2,4–8}. First identified in the nematode *Caenorhabditis elegans*, intrinsic apoptosis was next characterised in fruit fly *Drosophila melanogaster* and mammals. Intrinsic apoptosis depends on the key initiator caspase-9 (named Ced-3 and Dronc in *Caenorhabditis* and *Drosophila*, respectively), the activator Apaf-1 and the Bcl-2 multigenic family^{2,6,9}. Many functional studies have led to a deep characterisation of intrinsic apoptosis based on those classical models. Nevertheless, the biochemical role of mitochondria, the pivotal function of cytochrome c and the modality of caspases activation remain highly heterogeneous and hide profound molecular divergences among apoptotic pathways in animals^{8,10}. Independent of functional approaches, the phylogenetic history of the signal transduction actors, mostly the caspase family, is the Rosetta Stone to shed light on intrinsic apoptosis evolution. Here, after exhaustive research on CARD-caspases, we demonstrate by phylogenetic analysis that the caspase-9, the fundamental key of intrinsic apoptosis, is deuterostomes-specific, while it is the caspase-2 which is ancestral and common to bilaterians. Our analysis of Bcl-2 family and Apaf-1 confirm the high heterogeneity in apoptotic pathways elaboration in animals. Taken together, our results support convergent emergence of distinct intrinsic apoptotic pathways during metazoan evolution.

Keywords

Intrinsic apoptosis, initiator CARD-caspase, apoptotic network evolution, phylogeny

RESULTS AND DISCUSSION

Apoptosis, a regulated cell death, occurs during metazoan development, tissue homeostasis and regeneration^{11–13}. Pioneering works in *Caenorhabditis* established a molecular network of apoptosis decision, execution, and engulfment-degradation^{2,4,14}. Next, investigation of apoptotic cascade key components from *Drosophila* and mammals imposed the paradigm that the intrinsic apoptotic “molecular program” is conserved throughout animal evolution^{2,4–7}. However, recent researches revealed evolutionary divergences between these models, with major functional implications^{15,16}. Based on evolutionarily conserved or divergent features of the cell death machinery among metazoans, we investigated the evolutionary history of several major actors of intrinsic apoptosis. We conducted exhaustive research of Apaf-1 and extensive phylogenetic analyses of initiator CARD-caspases and Bcl-2 multigenic family proteins from major animal phyla and describe their evolutionary patterns. It is apparent that in spite of common functional similarities, each actor of these pathways has a distinct evolutionary history that led us to consider the structural and functional organisation of intrinsic apoptotic components as the result of evolutionary convergence among animals. Thus, the three animal models (nematode, fly, mouse) which were used to elaborate a unified concept of intrinsic apoptosis machinery, not only present major functional divergences, but are markedly distinct in their protein architecture, whose origin and evolutionary history followed very different molecular pathways.

Initiator caspases of the intrinsic apoptosis pathways are not homologous

Initiation and execution of apoptotic signalling pathways are fundamentally linked to the complex diversification of the caspases multigenic family, having a widely extended repertoire encompassing all metazoan phyla^{17–20}.

Caspases are a class of proteases composed of three protein domains; the pro-domain, the small P10, and the large P20^{9,21}. Initiator caspases, specific to an apoptotic pathway, have a specific Caspase Recruiting Domain (CARD) pro-domain or two Death Effector Domains (DED) pro-domains. The two distinct intrinsic and extrinsic apoptosis involve specifically the caspase-9 (CARD pro-domain) and the caspases-8-10 (DED pro-domains), respectively. Both pathways converge on common executioner caspases activation, triggering apoptosis execution^{2,3}.

Due to the pivotal role of caspase-9 as initiator in intrinsic apoptosis, our main analyses focused on CARD-caspases. We confirmed their distribution in most animal phyla and reconstructed their phylogenetic relationships (Figure 1, Table S1). Regardless of the phylogenetic methodology used for the analysis, three strongly supported (PP> 0.99) monophyletic groups were identified: caspase-9, caspase-2 and a more heterogeneous group named here [Inflammatory Caspases + Caspase-Y], respectively. Although relationships among these three clades differ depending on the methodology employed, their monophyly remains robust and together they form a clade strictly corresponding to bilaterian animals (PP = 0.87; BS>90), while the sequences of [Cnidaria + Ctenophora + Placozoa] form a divergent paraphyletic group named Caspase-X (Figure 1, Supplementary Figure 1).

Caspase-2 appears to be widely distributed among bilaterians and topology of the gene is congruent with the three major groups of eumetazoan [Deuterostomia + Ecdysozoa + Lophotrochozoa/Spiralia], reflecting an ancestral origin in bilaterians. However caspase-9 is restricted to deuterostome animals. Conversely, the clade [Caspase-Y + Inflammatory

Caspase] appears to be restricted to [Vertebrata and Lophotrochozoa]. The robustness of the deutostomes-specific caspase-9 clade was explored (Supplementary Figure 2), and its strict diversification among [Vertebrata + Cephalochordata + Echinodermata + Hemichordata] was confirmed (the relative position of the cephalochordate *Branchiostoma belcheri* 9A paralogous gene remains unstable). However, consistent with previous studies on the ascidian *Ciona intestinalis*²², the five representatives of ascidian genomes studied are devoid of any caspase-9, probably confirming the loss of this gene in urochordates, the sister-group of vertebrates²³. Within bilaterians, the clade-specific acquisition of caspase-9 is a major event, probably leading to the specific functionalities observed in mammals, such as the mitochondrial outer membrane permeabilisation (MOMP) allowing cytochrome c release.

Unexpectedly, *Drosophila* Dronc and *Caenorhabditis* Ced-3 are distinctly identified as orthologous to caspase-2 of vertebrates, and not to caspase-9, as was previously reported and largely accepted^{2,6}. All Dronc/Ced3 proteins of insects, horseshoe crab (Xiphosura), and nematodes form a well-defined, strongly supported (PP = 1), monophyletic clade showing only one orthologous gene per species (except horseshoe crab). Thus, the caspase-2 clade of ecdysozoans is the sister group of caspase-2 of [Lophotrochozoa + Deuterostomia]. The absence of identifiable caspase-2 in echinoderms or hemichordates can probably be interpreted as a specific loss in the latter group (i.e. Ambulacraria).

Taken globally, our analysis shown that the usually considered conserved and ancestral initiator caspases of intrinsic apoptosis are not orthologs but divergent.

The caspase activation regulator apoptosome is highly divergent in metazoans

In intrinsic apoptosis, initiator caspase activation depends on their recruitment by a pivotal and considered shared component, the apoptosome platform^{2,15}. Apoptosome

formation relies on critical protein interactions which are *Caenorhabditis* Ced-4, *Drosophila* Dark, and human Apaf-1 with their respective initiator caspases, Ced-3, Dronk, and procaspase-9¹⁵. CARD and other protein domains (NOD, arm) are highly conserved in Apaf-1, Dark and Ced-4^{24–26}. However, excepted the majority of nematode species¹⁶, Apaf-1 possesses WD40 repeats at its C-terminus to bind to the cytochrome c. This binding is required in mammals for Apaf-1 oligomerisation and apoptosome formation^{26,27}, while Ced-4 and Dark do not require cytochrome c for their assembly into an apoptosome^{15,28}. Taken with major structural and regulating assembly differences between the octameric Dark, tetrameric Ced-4 and heptameric Apaf-1 apoptosomes, it reveal evolutionary divergences between animal apoptosomes formation and procaspases activation mechanisms, probably differentially modulating the cell death execution pathway²⁹.

Here we conducted exhaustive analysis by reciprocal BLAST and phylogenetic analyses of Apaf-1 homologs (data not show) and confirmed its ubiquity in the majority of metazoan phyla (Table S2). Remarkably, Apaf-1 orthologs were absent from all urochordates and molluscs in accordance with previous work^{22,25,30–33}, and from *Capitella teleta* (Annelida), suggesting independent losses in those animals or, a less parsimonious hypothesis, multiple independent evolution of modular proteins from the common ancestor of metazoans²⁵ (Figure 2). In non-bilaterians, Apaf-1 gene family has not been found in *Pleurobrachia pileus* genome (Ctenophora), but seems present in other early diverging metazoan phyla [Porifera, Cnidaria and Placozoa] (^{25,34,35}, our data). However, their CARD domains were highly divergent from caspase-9 domains of vertebrates and comparative analysis revealed unrelated complex apoptosis networks.

Importantly, analyses from several genomes (cnidarian, nematode, fly, amphioxus, sea urchin, human) clearly identified three well-defined independent clades of paralogous genes

among metazoans, highlighting that Ced-4, Dark and Apaf-1 are not homologous between ecdysozoans and vertebrates²⁵. The divergent evolution history of structural actors of the apoptosome platforms confirm functional analyses carried out so far suggesting that caspase structure and interaction differs among taxa and clades¹⁶.

Despite its fundamental role in the apoptotic cell death pathways and also in non-apoptotic functions^{36–38}, diverse metazoan phyla have independently experienced adapter protein Apaf-1 ortholog losses or independent molecular evolution. Consequently, the modality of apoptosome formation and the subsequent caspases activation are convergent among metazoan, and the similar mechanisms observed are hypothesised to be related to relaxation of functional constraints on molecular Apaf-1-like molecules oligomerisation process.

Finally, in-depth analysis of apoptosome structure argues in favour of distinct evolutionary origins, most likely modulated by functional interactions involving distinct initiator caspases.

The regulation of apoptosis by the Bcl-2 family is convergent among metazoans

Intrinsic apoptosis is ultimately regulated by the Bcl-2 proteins, composed of several Bcl-2 homologous (BH) domains^{39,40}. In mammals, the balance between pro-survival (four BH1-BH4 domains) and pro-apoptotic proteins (Bax/Bak/Bok and BH3-only) of the Bcl-2 controls initiation of intrinsic apoptosis. Conformational changes of the three-dimensional structures and interactions between Bcl-2 actors enables the assembly of pore-like structures controlling MOMP⁴¹.

Multiple sequence alignments of metazoan Bcl-2 family proteins (Supplementary Figure 3, Table S3) (but with over-represented chordates reflective of greater availability of

vertebrate genomes) confirm the distribution of Bcl-2 in Metazoa^{16,24,40,42}. Consistent with functional Bcl-2 classification, proteins clustered into five monophyletic groups (i.e. three ‘pro-apoptotic’ clades: Bok, Bak, Bax and two less supported pro-survival - ‘anti-apoptotic’ groups: Bcl-2/W/XL and a more complex Bcl-B / Mcl-1 / Bfl-1 clade) (Supplementary Figure 3). However, respective relationships among each of these well-supported groups were not clearly resolved, and some divergent or less characterised sequences from molluscs such as *Biomphalaria glabrata* (Bcl-like2, Bcl-like3), from the cnidarian *Hydra vulgaris* (Bcl-like1) or from the urochordate *Ciona intestinalis* (Bcl-like1) were not strictly attributed to a particular class.

Caenorhabditis is deprived of pro-apoptotic Bcl-2 and possesses only the pro-survival Ced-9 (orthologous to vertebrates Bcl-2/w/xl), and two BH-3 only proteins (Egl-1 and Ced-13). Conversely, only two pro-apoptotic Bok-like close paralogs (Debcl and Buffy) were present in *Drosophila* (Figure 2, Supplementary Figure 3), but their functions remains unclear⁴³. In both animals, there is no MOMP and apoptosome assembly does not require cytochrome c binding⁴⁴. Presence of both anti- and pro-apoptotic Bcl-2 in molluscs underlines the divergent particularities observed inside Protostomia. Contrarily, similarities of Bcl-2 family composition are observed within some deuterostomes (mammals and echinoderms).

Finally, if multiple Bcl-2 genes were acquired early in metazoan evolution, and despite conservation of almost homologous genes, key differences accumulate, making initiation mechanisms in intrinsic apoptotic signalling pathways convergent among animals.

Apoptotic mitochondrial pathways are convergent among metazoans

Functional evidences emphasise that caspase-2 members play a critical role in various cell deaths, but have independently involved in a range of non-apoptotic functions, including

cell cycle regulation, DNA repair and tumour suppression^{45–48}. This implication of caspase-2 in a myriad of signalling pathways and interaction with a panel of adaptor molecules demonstrates its functional versatility^{45,46,49–52}. As previously reported and unlike other initiator caspases studied, our phylogenetic analyses corroborate the wide distribution of caspase-2 in bilaterians and suggest its ancestral multifunctionality.

The major structural and functional similarities that led to an erroneous interpretation of the phylogenetic position of Ced-3 and Dronc underlines a probable common evolutionary origin of caspase-2 and -9 genes. We propose here that caspase-9 originates from deuterostome-specific duplication of a caspase-2-like gene, followed by functional specialisation of paralogs. In the case of vertebrates (and probably in Cephalochordates), the two families of paralogs have been preserved. Caspase-2 retains multifunctional activity, and in mammals can interact with PIDDosome platform containing P53, adapter molecules RAIDD, and signalling complex DISC, activating both extrinsic, intrinsic, and DNA damage pathways^{51,53–55}. Conversely, the caspase-9 gene underwent a functional divergence in connection with its specialisation in allosteric interactions with the apoptosome³.

Due to its pivotal role as a mediator of genomic stability through involvement in cell proliferation, oxidative stress, aging and cell death, the molecular divergence of the caspase-2 gene is highly constrained during evolution, probably because destabilisation of any signalling cascade is sufficient to initiate tumorigenesis. Therefore, purifying selection is likely important during caspase-2 evolution, except during the radiation of deuterostomes, when it could have been relaxed due to the duplication and the modification of the functionalisation of caspase duplicates (cf. Duplication-Degeneration-Complementation model)^{56–58}. This model could explain the loss of caspase-9 concomitantly with a caspase-2 duplication in urochordates (Figure 1, Supplementary Figure 2). Similarly, orthologs of caspase-2 have been

lost, or strongly diverged, while caspase-9 orthologs present a *de novo* relative expansion in *Strongylocentrotus purpuratus* (Echinodermata) and *Saccoglossus kowalevskii* (Hemichordata).

Due to generalised gene losses in ecdysozoans and in comparison with other bilaterians, *Caenorhabditis* and *Drosophila* apoptotic pathways are generally considered simpler than those of vertebrates^{9,10}. However, what distinguishes ecdysozoans from lophotrochozoans (molluscs, annelids, and their relatives) and deuterostomes is not only a smaller number of genes but also above all the shape of pathways organised around different paralogous genes. The absence of orthologous relationships among genes results in a very different structural organisation of platforms but also generates important functional divergences (i.e. mechanisms of regulating assembly, CARD-CARD interactions with procaspases)¹⁵.

Consistent with mammalian caspase-2 functions, *Caenorhabditis* Ced-3 has both initiator as well as executioner activities but is activated by the Ced-4 platform in a specific manner^{8,51,59,60}. Unlike the organisation of the mammal apoptosome, the *Caenorhabditis* Ced-4 platform presents neither MOMP nor the release of cytochrome c^{4,10}. Due to its interaction with Dark (Apaf-1 paralog), the only CARD-caspase in *Drosophila* (Dronc) has been wrongly classified as caspase-9^{6,61}. Likewise, involvement of Dronc in various processes such as compensatory cell proliferation, inhibition of cell migration or spermatid differentiation, brings this protein closer to the functionalities of caspase-2 clade^{62,63}.

Like other protostomes, molluscs are deprived of caspase-9 but caspase-2 orthologs have been identified in bivalves and were suspected to function in “a caspase-9-like manner”⁶⁴. Hence, caspase-2 is responsible for apoptotic process during larval metamorphosis in the oyster *Crassostrea gigas*, but more surprisingly, despite the absence of caspase-9 and Apaf-1

(and thus, of a mammalian-like apoptosome), this peculiar pathway is amazingly associated with cytochrome c release^{30,33,65–68}. The complexity of intrinsic apoptosis in molluscs seems to be important, but divergent from what was observed in ecdysozoans or vertebrates, and more specifically shows a putative expansion of initiator and executioner caspases that participate both in immunity, stress responses, and apoptosis (Figure 2)^{33,64,69–72}.

Unexpectedly, mammals present almost the unique case (with probably the cephalochordates) in which both caspase-2 and caspase-9 are conserved and involved in apoptosis. This putative functional redundancy (i.e. recruitment, autoactivation or transactivation, homodimerisation and subsequent interchain proteolytic cleavage) undoubtedly led to the functional specialisation observed for caspase-9. This appears to be fundamentally linked to the mammalian mitochondrial pathway and non-apoptotic activity most often indirectly via caspase-3 activation^{73–76}. Finally, echinoderms seem to uniquely have an intrinsic apoptosis similar to mammals, with a caspase-9, Bcl-2, Apaf-1, and a MOMP with cytochrome c release (Figure 2)^{5,77}.

Although we can envisage a weak parsimonious scenario that showcases a common ancestral apoptotic pathway in deuterostomes (but implying independent secondarily losses in hemichordates, cephalocordates and urochordates), the similarities observed between echinoderms and mammals more probably reflect functional convergences based on independent recruitment of apoptotic actors.

CONCLUSION

The apoptotic networks of *Caenorhabditis* and *Drosophila* do not exemplify ancestral conditions from which mammalian-grade apoptotic complexity emerged but are on the

contrary, and as suggested recently, the result of a derived condition specific to ecdysozoans among animals^{24,69,78,79}.

The core components of intrinsic apoptotic pathways, especially initiator caspases and the apoptosome platform, are not ancestral in metazoans. Our phylogenetic analyses highlight an unexpected evolutionary history: while the bilaterian caspase-2-mediated apoptotic toolkit emerged ancestrally and remains multifunctional, the caspase-9 mediator of the mammalian apoptosome is specific to deuterostomes.

The major functional divergences in mitochondrial apoptotic pathways observed among animals^{8,80–82} mainly originated in the recruitment of paralogous actors from the same multigenic families reflecting the adaptive processes specific to each taxon, which lead ultimately to convergent evolutionary histories. Interestingly, this richness of the apoptotic genetic repertoire was suggested to be links to the persistence of stem cells in adults from different phyla^{83,84}.

Finally, mitochondria-mediated apoptosis, like other programmed cell deaths, has likely evolved before and throughout metazoan diversification to shape developmental processes, immune response, or to adapt cellular environment to environmental constraint.

RESOURCE AVAILABILITY

Lead contact

Further information and requests should be address to the Lead Contact, Gabriel Krasovec (gabriel.krasovec@nuigalway.ie).

Materials availability

Any request should be address to the Lead Contact.

282

283 **Data and code availability**

284 No code was generated during this study. Source data are available upon request.

285

286 **EXPERIMENTAL MODEL AND SUBJECT DETAILS**

287 No experimental model was used during this study.

288

289 **METHOD DETAILS**

290 **Sequences dataset construction**

291 Putative metazoans CARD-caspases were identified by using tBLASTn and BLASTp
 292 searches with human caspases (caspase-1 to 10), Ced-3, and Dronc as query on NCBI, ANISEED
 293 (ascidians), EchinoBase (*Strongylocentrotus purpuratus*), and neurobase.rc.ufl.edu
 294 (*Pleurobrachia bachei*) databases, and followed by reciprocal BLAST. After identification of
 295 CARD-caspases in target species, sequences were added as query to conduct BLAST searches
 296 in close relative (i.e. identified CARD-caspases of *Crassostrea gigas* were used as query to look
 297 for in other molluscs). Sequences with an e-value inferior to 1e-10 was retained. All identified
 298 sequences were analysed with ScanProsite (ExPaSy)⁸⁵ and InterProScan (EMBL-EBI)⁸⁶ to
 299 double check the presence of specific caspases domains. Another sequence was added to
 300 verify its identification proposed as a caspase-2 in literature: *Crassostrea angulata* casp-2⁶⁸.
 301 Caspases family are short proteins (containing the large common P20 and the small P10
 302 domains) with a high number of genes per species that rapidly limits the relevance of the
 303 phylogenetic analyses. To reduce the artefact branching and unreadable topology, the dataset
 304 was built using CARD-caspase gene repertoires of selected species and in order to maximize

phylogenetic diversity across Metazoa. A full list of all caspase sequences is provided in Table S1.

Metazoan Bcl-2 were identified by using tBLASTn and BLASTp searches with human Bcl-2 as query on NCBI, are followed by reciprocal BLAST. All identified sequences were analysed with ScanProsite (ExPaSy) and InterProScan (EMBL-EBI) to double check the presence of BH domains. Because of their too short sequences, BH3-only were not taking in account. A full list of all Bcl-2 sequences is provided in Table S3.

Multiple alignments of protein sequences were generated using the MAFFT software version 7⁸⁷ with default parameters and also Clustal Omega⁸⁸ to verify the congruence of the different alignments. All sequences were then manually checked in BloEdit 7.2 software⁸⁹ to verify the presence of the specific domains previously identified. Gblocks version 0.91b⁹⁰ was used to remove vacancies and blur sites. Final alignments are composed of 230, 235, and 147 amino acids for metazoan CARD-caspases alignment, deuterostomian CARD-caspases alignment, and metazoan Bcl-2 alignment, respectively.

APAF-1 detection

Identification of metazoan Apaf-1 was made by tBLASTn and BLASTp using human APAF-1, nematode Ced-4, and fly Dark as query on NCBI, ANISEED (ascidians), and neurobase.rc.ufl.edu (*Pleurobrachia bachei*) databases, and followed by reciprocal BLAST. E-value threshold was specified to be 0.1 to increase the chance of finding sequences that match. Potential resulting sequences were analyzed with ScanProsite and also InterProScan. A full list of all Apaf-1 sequences is provided in Table S2.

Phylogenetic analysis

Phylogenetic analyses were carried out from the amino-acid alignment by Maximum-Likelihood (ML) method using PhyML 3.1⁹¹, combined ML tree search with 1000 bootstrap replicates, and tree were visualized using Seaview⁹². Best amino-acid evolution model to conduct analysis were determined using MEGA11⁹³ and are WAG and LG model for CARD-caspases alignments and Bcl-2 alignment, respectively.

Bayesian analyses were performed using MrBayes (v3.2.6)⁹⁴ under mixed model. For each analysis, one fourth of the topologies were discarded as burn-in values, while the remaining ones were used to calculate the posterior probability. The run for metazoan CARD-caspases alignment was carried out for 2 000 000 generations with 15 randomly started simultaneous Markov chains (1 cold chain, 14 heated chains) and sampled every 100 generations. The run for deuterostomian CARD-caspases alignment was carried out for 500 000 generations with 5 randomly started simultaneous Markov chains (1 cold chain, 4 heated chains) and sampled every 100 generations. The run for metazoan Bcl-2 alignment was carried out for 5 000 000 generations with 20 randomly started simultaneous Markov chains (1 cold chain, 19 heated chains) and sampled every 100 generations.

ML bootstrap values higher than 50% and Bayesian posterior probabilities are indicated on the Bayesian tree (Figure 1; Supplementary Figures 2, 3)

For the metazoans caspase-CARD phylogeny, outgroup used is the only one caspase with a pro-domain Card of the Porifera *Amphimedon queenslandica* (XP_003383519) (Figure 1). Analyses of caspase-Card were made independently at the deuterostomes scale with four different outgroup to test their effect on the stability of the topology: i) caspase-Card-Y of the annelid *Capitella teleta* (ELT97848.1), ii) caspase-Card 2 of the mollusk *Aplysia californica* (XP_005113266), iii) caspase-Card-X2 of cnidarian *Hydra vulgaris* (NP_001274285.1) iv) caspase-Card Ced-3 of the ecdysozoan *Caenorhabditis elegans* (AAG42045.1) (Supplementary

Figure 2). For the metazoans Bcl-2 phylogenies, outgroup used to test their effect on the stability of the topology are: *i*) Bcl-2 like1 (XP_003383425.1) and Bcl-2 like2 (XP_003387574.1) of Porifera *Amphimedon queenslandica* (Supplementary Figure 3).

ACKNOWLEDGEMENTS

Authors acknowledge Sébastien Darras (Sorbonne Université, Banyuls-sur-mer), Christine Vesque (Sorbonne Université, Paris), Jérôme Gros (Institut Pasteur, Paris), Sabine Hennequin (Sorbonne Université, Paris), Uri Frank (NUIG, Galway), and Helen Horkan (NUIG, Galway) for helpful comments. GK was supported by a Ph.D. fellowship from the French Ministry of Education, Research and Innovation.

AUTHOR CONTRIBUTIONS

JP and EQ managed the project. GK made BLAST, phylogenetic analysis, and figures. GK and EQ wrote the manuscript.

DECLARATION OF INTERESTS

Authors declare no competing interests.

BIBLIOGRAPHY

1. Kerr, J.F., Wyllie, A.H., and Currie, A.R. (1972). Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br. J. Cancer* 26, 239–257.
2. Hengartner, M.O. (2000). The biochemistry of apoptosis. *Nature* 407, 770–776.
3. Galluzzi, L., Vitale, I., Aaronson, S.A., Abrams, J.M., Adam, D., Agostinis, P., Alnemri, E.S., Altucci, L., Amelio, I., Andrews, D.W., et al. (2018). Molecular mechanisms of cell death: recommendations of the Nomenclature Committee on Cell Death 2018. *Cell Death Differ.* 25, 486–541.

- 379 4. Lettre, G., and Hengartner, M.O. (2006). Developmental apoptosis in *C. elegans*: a complex
380 CEDnario. *Nat Rev Mol Cell Biol* 7, 97–108.
- 381 5. Bender, C.E., Fitzgerald, P., Tait, S.W.G., Llambi, F., McStay, G.P., Tupper, D.O., Pellettieri, J.,
382 Sánchez Alvarado, A., Salvesen, G.S., and Green, D.R. (2012). Mitochondrial pathway of apoptosis
383 is ancestral in metazoans. *Proc. Natl. Acad. Sci. U.S.A.* 109, 4904–4909.
- 384 6. Steller, H. (2008). Regulation of apoptosis in *Drosophila*. *Cell Death Differ.* 15, 1132–1138.
- 385 7. Driscoll, M. (1996). Cell death in *C. elegans*: molecular insights into mechanisms conserved
386 between nematodes and mammals. *Brain Pathol* 6, 411–425.
- 387 8. Fuchs, Y., and Steller, H. (2015). Live to die another way: modes of programmed cell death and
388 the signals emanating from dying cells. *Nat. Rev. Mol. Cell Biol.* 16, 329–344.
- 389 9. S, E. (2007). Apoptosis: a review of programmed cell death. *Toxicologic pathology* 35.
390 <https://pubmed.ncbi.nlm.nih.gov/17562483/>.
- 391 10. Meier, P., Finch, A., and Evan, G. (2000). Apoptosis in development. *Nature* 407, 796–801.
- 392 11. Jacobson, M.D., Weil, M., and Raff, M.C. (1997). Programmed cell death in animal development.
393 *Cell* 88, 347–354.
- 394 12. Hipfner, D.R., and Cohen, S.M. (2004). Connecting proliferation and apoptosis in development
395 and disease. *Nat Rev Mol Cell Biol* 5, 805–815.
- 396 13. Vríz, S., Reiter, S., and Galliot, B. (2014). Cell death: a program to regenerate. *Curr. Top. Dev. Biol.*
397 108, 121–151.
- 398 14. Ellis, H.M., and Horvitz, H.R. (1986). Genetic control of programmed cell death in the nematode
399 *C. elegans*. *Cell* 44, 817–829.
- 400 15. Dorstyn, L., Akey, C.W., and Kumar, S. (2018). New insights into apoptosome structure and
401 function. *Cell Death Differ* 25, 1194–1208.
- 402 16. Young, N.D., Harris, T.J., Evangelista, M., Tran, S., Wouters, M.A., Soares da Costa, T.P., Kershaw,
403 N.J., Gasser, R.B., Smith, B.J., Lee, E.F., et al. (2020). Diversity in the intrinsic apoptosis pathway
404 of nematodes. *Commun Biol* 3, 478.
- 405 17. Bell, R.A.V., and Megeney, L.A. (2017). Evolution of caspase-mediated cell death and
406 differentiation: twins separated at birth. *Cell Death Differ* 24, 1359–1368.
- 407 18. Shrestha, S., Tung, J., Grinshpon, R.D., Swartz, P., Hamilton, P.T., Dimos, B., Mydlarz, L., and
408 Clark, A.C. (2020). Caspases from scleractinian coral show unique regulatory features. *J Biol*
409 *Chem* 295, 14578–14591.
- 410 19. Uren, A.G., O'Rourke, K., Aravind, L., Pisabarro, M.T., Seshagiri, S., Koonin, E.V., and Dixit, V.M.
411 (2000). Identification of Paracaspases and Metacaspases: Two Ancient Families of Caspase-like
412 Proteins, One of which Plays a Key Role in MALT Lymphoma. *Molecular Cell* 6, 961–967.
- 413 20. Crawford, E.D., Seaman, J.E., Barber, A.E., David, D.C., Babbitt, P.C., Burlingame, A.L., and Wells,
414 J.A. (2012). Conservation of caspase substrates across metazoans suggests hierarchical

- importance of signaling pathways over specific targets and cleavage site motifs in apoptosis. *Cell Death Differ* 19, 2040–2048.
21. Thornberry, N.A., and Lazebnik, Y. (1998). Caspases: enemies within. *Science* 281, 1312–1316.
22. Terajima, D., Shida, K., Takada, N., Kasuya, A., Rokhsar, D., Satoh, N., Satake, M., and Wang, H.-G. (2003). Identification of candidate genes encoding the core components of the cell death machinery in the *Ciona intestinalis* genome. *Cell Death Differ.* 10, 749–753.
23. Delsuc, F., Brinkmann, H., Chourrout, D., and Philippe, H. (2006). Tunicates and not cephalochordates are the closest living relatives of vertebrates. *Nature* 439, 965–968.
24. Zmasek, C.M., and Godzik, A. (2013). Evolution of the animal apoptosis network. *Cold Spring Harb Perspect Biol* 5, a008649.
25. Zmasek, C.M., Zhang, Q., Ye, Y., and Godzik, A. (2007). Surprising complexity of the ancestral apoptosis network. *Genome Biol.* 8, R226.
26. Zou, H., Henzel, W.J., Liu, X., Lutschg, A., and Wang, X. (1997). Apaf-1, a human protein homologous to *C. elegans* CED-4, participates in cytochrome c-dependent activation of caspase-3. *Cell* 90, 405–413.
27. Zou, H., Li, Y., Liu, X., and Wang, X. (1999). An APAF-1.cytochrome c multimeric complex is a functional apoptosome that activates procaspase-9. *J. Biol. Chem.* 274, 11549–11556.
28. Dorstyn, L., Read, S., Cakouros, D., Huh, J.R., Hay, B.A., and Kumar, S. (2002). The role of cytochrome c in caspase activation in *Drosophila melanogaster* cells. *J Cell Biol* 156, 1089–1098.
29. Shi, Y. (2006). Mechanical aspects of apoptosome assembly. *Curr Opin Cell Biol* 18, 677–684.
30. Li, Y., Zhang, L., Qu, T., Tang, X., Li, L., and Zhang, G. (2017). Conservation and divergence of mitochondrial apoptosis pathway in the Pacific oyster, *Crassostrea gigas*. *Cell Death Dis* 8, e2915.
31. Kiss, T. (2010). Apoptosis and its functional significance in molluscs. *Apoptosis* 15, 313–321.
32. Romero, A., Novoa, B., and Figueras, A. (2015). The complexity of apoptotic cell death in mollusks: An update. *Fish Shellfish Immunol.* 46, 79–87.
33. Estévez-Calvar, N., Romero, A., Figueras, A., and Novoa, B. (2013). Genes of the mitochondrial apoptotic pathway in *Mytilus galloprovincialis*. *PLoS ONE* 8, e61502.
34. Srivastava, M., Simakov, O., Chapman, J., Fahey, B., Gauthier, M.E.A., Mitros, T., Richards, G.S., Conaco, C., Dacre, M., Hellsten, U., et al. (2010). The *Amphimedon queenslandica* genome and the evolution of animal complexity. *Nature* 466, 720–726.
35. Srivastava, M., Begovic, E., Chapman, J., Putnam, N.H., Hellsten, U., Kawashima, T., Kuo, A., Mitros, T., Salamov, A., Carpenter, M.L., et al. (2008). The *Trichoplax* genome and the nature of placozoans. *Nature* 454, 955–960.
36. Zermati, Y., Mouhamad, S., Stergiou, L., Besse, B., Galluzzi, L., Boehrer, S., Pauleau, A.-L., Rosselli, F., D’Amelio, M., Amendola, R., et al. (2007). Nonapoptotic role for Apaf-1 in the DNA damage checkpoint. *Mol Cell* 28, 624–637.

37. Ferraro, E., Pesaresi, M.G., De Zio, D., Cencioni, M.T., Gortat, A., Cozzolino, M., Berghella, L., Salvatore, A.M., Oettinghaus, B., Scorrano, L., et al. (2011). Apaf1 plays a pro-survival role by regulating centrosome morphology and function. *J Cell Sci* 124, 3450–3463.
38. Shakeri, R., Kheirollahi, A., and Davoodi, J. (2017). Apaf-1: Regulation and function in cell death. *Biochimie* 135, 111–125.
39. Lee, E.F., Clarke, O.B., Evangelista, M., Feng, Z., Speed, T.P., Tchoubrieva, E.B., Strasser, A., Kalinna, B.H., Colman, P.M., and Fairlie, W.D. (2011). Discovery and molecular characterization of a Bcl-2-regulated cell death pathway in schistosomes. *Proc. Natl. Acad. Sci. U.S.A.* 108, 6999–7003.
40. Banjara, S., Suraweera, C.D., Hinds, M.G., and Kvansakul, M. (2020). The Bcl-2 Family: Ancient Origins, Conserved Structures, and Divergent Mechanisms. *Biomolecules* 10.
41. Kalkavan, H., and Green, D.R. (2018). MOMP, cell suicide as a BCL-2 family business. *Cell Death Differ* 25, 46–55.
42. Dunn, S.R., Phillips, W.S., Spatafora, J.W., Green, D.R., and Weis, V.M. (2006). Highly conserved caspase and Bcl-2 homologues from the sea anemone *Aiptasia pallida*: lower metazoans as models for the study of apoptosis evolution. *J. Mol. Evol.* 63, 95–107.
43. Clavier, A., Rincheval-Arnold, A., Colin, J., Mignotte, B., and Guénal, I. (2016). Apoptosis in *Drosophila*: which role for mitochondria? *Apoptosis* 21, 239–251.
44. Abdelwahid, E., Yokokura, T., Krieser, R.J., Balasundaram, S., Fowle, W.H., and White, K. (2007). Mitochondrial disruption in *Drosophila* apoptosis. *Dev Cell* 12, 793–806.
45. Krumschnabel, G., Sohm, B., Bock, F., Manzl, C., and Villunger, A. (2009). The enigma of caspase-2: the laymen’s view. *Cell Death & Differentiation* 16, 195–207.
46. Olsson, M., Forsberg, J., and Zhivotovsky, B. (2015). Caspase-2: the reinvented enzyme. *Oncogene* 34, 1877–1882.
47. Lassus, P., Opitz-Araya, X., and Lazebnik, Y. (2002). Requirement for caspase-2 in stress-induced apoptosis before mitochondrial permeabilization. *Science* 297, 1352–1354.
48. Zhivotovsky, B., and Orrenius, S. (2005). Caspase-2 function in response to DNA damage. *Biochem. Biophys. Res. Commun.* 331, 859–867.
49. Krumschnabel, G., Manzl, C., and Villunger, A. (2009). Caspase-2: killer, savior and safeguard--emerging versatile roles for an ill-defined caspase. *Oncogene* 28, 3093–3096.
50. Olsson, M., Vakifahmetoglu, H., Abruzzo, P.M., Höglstrand, K., Grandien, A., and Zhivotovsky, B. (2009). DISC-mediated activation of caspase-2 in DNA damage-induced apoptosis. *Oncogene* 28, 1949–1959.
51. Lavrik, I.N., Golks, A., Baumann, S., and Krammer, P.H. (2006). Caspase-2 is activated at the CD95 death-inducing signaling complex in the course of CD95-induced apoptosis. *Blood* 108, 559–565.
52. Braga, M., Sinha Hikim, A.P., Datta, S., Ferrini, M.G., Brown, D., Kovacheva, E.L., Gonzalez-Cadavid, N.F., and Sinha-Hikim, I. (2008). Involvement of oxidative stress and caspase 2-mediated

488 intrinsic pathway signaling in age-related increase in muscle cell apoptosis in mice. *Apoptosis* 13,
489 822–832.

490 53. Duan, H., and Dixit, V.M. (1997). RAIDD is a new “death” adaptor molecule. *Nature* 385, 86–89.

491 54. Tinel, A., and Tschopp, J. (2004). The PIDDosome, a protein complex implicated in activation of
492 caspase-2 in response to genotoxic stress. *Science* 304, 843–846.

493 55. Haupt, S., Berger, M., Goldberg, Z., and Haupt, Y. (2003). Apoptosis - the p53 network. *J Cell Sci*
494 116, 4077–4085.

495 56. Force, A., Lynch, M., Pickett, F.B., Amores, A., Yan, Y.L., and Postlethwait, J. (1999). Preservation
496 of duplicate genes by complementary, degenerative mutations. *Genetics* 151, 1531–1545.

497 57. Conant, G.C., and Wolfe, K.H. (2008). Turning a hobby into a job: how duplicated genes find new
498 functions. *Nat Rev Genet* 9, 938–950.

499 58. Fares, M.A., Keane, O.M., Toft, C., Carretero-Paulet, L., and Jones, G.W. (2013). The roles of
500 whole-genome and small-scale duplications in the functional specialization of *Saccharomyces*
501 *cerevisiae* genes. *PLoS Genet* 9, e1003176.

502 59. Degterev, A., Boyce, M., and Yuan, J. (2003). A decade of caspases. *Oncogene* 22, 8543–8567.

503 60. Mancini, M., Machamer, C.E., Roy, S., Nicholson, D.W., Thornberry, N.A., Casciola-Rosen, L.A.,
504 and Rosen, A. (2000). Caspase-2 Is Localized at the Golgi Complex and Cleaves Golgin-160 during
505 Apoptosis. *J Cell Biol* 149, 603–612.

506 61. Kumar, S., and Doumanis, J. (2000). The fly caspases. *Cell Death Differ* 7, 1039–1044.

507 62. Ouyang, Y., Petritsch, C., Wen, H., Jan, L., Jan, Y.N., and Lu, B. (2011). Dronc caspase exerts a non-
508 apoptotic function to restrain phospho-Numb-induced ectopic neuroblast formation in
509 *Drosophila*. *Development* 138, 2185–2196.

510 63. Huh, J.R., Guo, M., and Hay, B.A. (2004). Compensatory proliferation induced by cell death in the
511 *Drosophila* wing disc requires activity of the apical cell death caspase Dronc in a nonapoptotic
512 role. *Curr. Biol.* 14, 1262–1266.

513 64. Vogeler, S., Carboni, S., Li, X., and Joyce, A. (2021). Phylogenetic analysis of the caspase family in
514 bivalves: implications for programmed cell death, immune response and development. *BMC*
515 *Genomics* 22, 80.

516 65. Pirger, Z., Rácz, B., and Kiss, T. (2009). Dopamine-induced programmed cell death is associated
517 with cytochrome c release and caspase-3 activation in snail salivary gland cells. *Biol. Cell* 101,
518 105–116.

519 66. Romero, A., Estévez-Calvar, N., Dios, S., Figueras, A., and Novoa, B. (2011). New insights into the
520 apoptotic process in mollusks: characterization of caspase genes in *Mytilus galloprovincialis*.
521 *PLoS ONE* 6, e17003.

522 67. Zhang, L., Li, L., and Zhang, G. (2011). Gene discovery, comparative analysis and expression
523 profile reveal the complexity of the *Crassostrea gigas* apoptosis system. *Dev. Comp. Immunol.*
524 35, 603–610.

68. Yang, B., Li, L., Pu, F., You, W., Huang, H., and Ke, C. (2015). Molecular cloning of two molluscan caspases and gene functional analysis during *Crassostrea angulata* (Fujian oyster) larval metamorphosis. *Mol. Biol. Rep.* 42, 963–975.
69. Plachetzki, D.C., Pankey, M.S., MacManes, M.D., Lesser, M.P., and Walker, C.W. (2020). The Genome of the Softshell Clam *Mya arenaria* and the Evolution of Apoptosis. *Genome Biol Evol* 12, 1681–1693.
70. Piquet, B., Shillito, B., Lallier, F.H., Duperron, S., and Andersen, A.C. (2019). High rates of apoptosis visualized in the symbiont-bearing gills of deep-sea *Bathymodiolus* mussels. *PLoS ONE* 14, e0211499.
71. Sokolova, I.M., Evans, S., and Hughes, F.M. (2004). Cadmium-induced apoptosis in oyster hemocytes involves disturbance of cellular energy balance but no mitochondrial permeability transition. *J. Exp. Biol.* 207, 3369–3380.
72. Sokolova, I.M. (2009). Apoptosis in molluscan immune defense. *Invertebrate Survival Journal* 6, 49–58.
73. An, H.-K., Chung, K.M., Park, H., Hong, J., Gim, J.-E., Choi, H., Lee, Y.W., Choi, J., Mun, J.Y., and Yu, S.-W. (2020). CASP9 (caspase 9) is essential for autophagosome maturation through regulation of mitochondrial homeostasis. *Autophagy* 16, 1598–1617.
74. Madadi, Z., Akbari-Birgani, S., Monfared, P.D., and Mohammadi, S. (2019). The non-apoptotic role of caspase-9 promotes differentiation in leukemic cells. *Biochim Biophys Acta Mol Cell Res* 1866, 118524.
75. Hollville, E., and Deshmukh, M. (2018). Physiological functions of non-apoptotic caspase activity in the nervous system. *Semin Cell Dev Biol* 82, 127–136.
76. Tran, H.T., Fransen, M., Dimitrakopoulou, D., Van Imschoot, G., Willemarck, N., and Vleminckx, K. (2017). Caspase-9 has a nonapoptotic function in *Xenopus* embryonic primitive blood formation. *J Cell Sci* 130, 2371–2381.
77. Tamura, R., Takada, M., Sakaue, M., Yoshida, A., Ohi, S., Hirano, K., Hayakawa, T., Hirohashi, N., Yura, K., and Chiba, K. (2018). Starfish Apaf-1 activates effector caspase-3/9 upon apoptosis of aged eggs. *Sci Rep* 8, 1611.
78. Belyi, V.A., Ak, P., Markert, E., Wang, H., Hu, W., Puzio-Kuter, A., and Levine, A.J. (2010). The origins and evolution of the p53 family of genes. *Cold Spring Harb Perspect Biol* 2, a001198.
79. Lu, W.-J., Amatruda, J.F., and Abrams, J.M. (2009). p53 ancestry: gazing through an evolutionary lens. *Nat Rev Cancer* 9, 758–762.
80. Berthelet, J., and Dubrez, L. (2013). Regulation of Apoptosis by Inhibitors of Apoptosis (IAPs). *Cells* 2, 163–187.
81. Ribeiro Lopes, M., Parisot, N., Callaerts, P., and Clevro, F. (2019). Genetic Diversity of the Apoptotic Pathway in Insects. In *Evolution, Origin of Life, Concepts and Methods*, P. Pontarotti, ed. (Springer International Publishing), pp. 253–285.

82. Ribeiro Lopes, M., Parisot, N., Gaget, K., Huygens, C., Peignier, S., Duport, G., Orlans, J., Charles, H., Baatsen, P., Jousselin, E., et al. (2020). Evolutionary novelty in the apoptotic pathway of aphids. *Proc Natl Acad Sci U S A* 117, 32545–32556.
83. Aubrey, B.J., Janic, A., Chen, Y., Chang, C., Lieschke, E.C., Diepstraten, S.T., Kueh, A.J., Bernardini, J.P., Dewson, G., O'Reilly, L.A., et al. (2018). Mutant TRP53 exerts a target gene-selective dominant-negative effect to drive tumor development. *Genes Dev* 32, 1420–1429.
84. Aubrey, B.J., Kelly, G.L., Janic, A., Herold, M.J., and Strasser, A. (2018). How does p53 induce apoptosis and how does this relate to p53-mediated tumour suppression? *Cell Death Differ* 25, 104–113.
85. Gattiker, A., Gasteiger, E., and Bairoch, A. (2002). ScanProsite: a reference implementation of a PROSITE scanning tool. *Appl Bioinformatics* 1, 107–108.
86. Quevillon, E., Silventoinen, V., Pillai, S., Harte, N., Mulder, N., Apweiler, R., and Lopez, R. (2005). InterProScan: protein domains identifier. *Nucleic Acids Res* 33, W116–120.
87. Katoh, K., and Standley, D.M. (2013). MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* 30, 772–780.
88. Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M., Söding, J., et al. (2011). Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol. Syst. Biol.* 7, 539.
89. HALL, T.A. (1999). BioEdit : a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp. Ser.* 41, 95–98.
90. Castresana, J. (2000). Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol. Biol. Evol.* 17, 540–552.
91. Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O. (2010). New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* 59, 307–321.
92. Gouy, M., Guindon, S., and Gascuel, O. (2010). SeaView version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol* 27, 221–224.
93. Tamura, K., Stecher, G., and Kumar, S. (2021). MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Mol Biol Evol* 38, 3022–3027.
94. Ronquist, F., and Huelsenbeck, J.P. (2003). MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19, 1572–1574.

Figure 1:

Topology of metazoan CARD-caspases phylogeny obtained by Bayesian inference. Maximum likelihood method produces same topology. Three strongly supported monophyletic groups are identified: caspase-9, caspase-2 and a more heterogeneous group named here [Inflammatory Caspases + Caspase-Y]. Together they form a clade strictly corresponding to

bilaterian animals. Caspase-2 is widely distributed among bilaterians [Deutostomia + Ecdysozoa + Lophotrochozoa/Spiralia] reflecting an ancestral origin. Conversely, caspase-9 is strictly restricted to deuterostomian animals. Sequences of non-bilaterians (Cnidaria + Ctenophora + Placozoa) form divergent paraphyletic groups. We used as outgroup the unique CARD-caspase of the Porifera *Amphimedon queenslandica* (XP_003383519).

Figure 2:

Reconstruction of convergent hypothetical intrinsic apoptotic pathways among metazoans according to molecular actors detected and identified in their genomes. Variability of intrinsic apoptotic pathways among animals emerged from convergences and recruitment of distinct actors having independent evolutionary history. Caspase-2 is bilaterian-specific and the initiator of ecdysozoans intrinsic apoptosis. Caspase-9 is restricted to deuterostomes and the specific initiator of mammalian intrinsic apoptosis. Deuterostomes exhibit several losses (i.e. caspase 2 in cephalochordates, caspase 9 in urochordates) or duplication (i.e. caspase 9 in echinoderms), highlighting a putative evolutionary flexibility in apoptotic pathways establishment. Mitochondrial functions diverge among phyla and cytochrome c release thanks to the mitochondrial outer membrane permeabilisation (MOMP) is only specific to mammals

and possibly echinoderms. The convergent evolutionary histories certainly reflect a probable
phylum specific adaptive processes leading to parallel evolution of mitochondrial apoptotic
pathways observed among animals.

Supplementary Figure 1:

Monophyly of each CARD-caspase group is conserved using Bayesian inference and maximum likelihood analyses. Caspase-9, caspase-2 and [Inflammatory Caspases + Caspase-Y] groups remain conserved, confirming the bilaterian-specificity of caspase-2 and deuterostomian-specificity of caspase-9. Relationships among these three clades differ depending on the methodology employed. Divergent sequences of non-bilaterians animals (Cnidaria + Ctenophora + Placozoa) always form a paraphyletic group of caspases.

696

697

698

699

700

701

702

703

704 *californica* caspase-2 (XP_005113266) (**A2**), the *Hydra vulgaris* caspase-X2 (NP_001274285.1)
 705 (**A3**), and the *Caenorhabditis elegans* Ced3 (AAG42045.1) (**A4**).

706
 707
 708
 709
 710
 711
 712
 713
 714
 715
 716
 717
 718
 719
 720
 721
 722
 723
 724
 725
 726
 727
 728
 729
 730

Bayesian posterior probabilities 0 to 1.00

● ML Bootstrap >90

● ML Bootstrap 70-90

○ ML Bootstrap 50-70

Supplementary Figure 3:

Topology of Bcl-2 family phylogeny from Bayesian inference and maximum likelihood at the metazoan scale. Bcl-2 proteins clustered into five monophyletic groups: three 'pro-apoptotic' clades (Bok, Bak, Bax) and two less supported 'anti-apoptotic' groups (Bcl-2/W/XL) and a more complex (Bcl-B/Mcl-1/Bfl-1) clade. Each group includes bilaterians as well as non-bilaterians sequences. Clustering into these five groups is consistent between analyses, while relationship

among them is not well resolved. Numbers given on branches correspond to posterior probabilities. We used respectively as outgroup the *Amphimedon queenslandica* Bcl-like 1 (XP_003383425.1) (**A**), and *Amphimedon queenslandica* Bcl-like 2 (XP_003387574.1) (**B**).

772 Supplemental table 1: List of CARD-caspases used for phylogenetic analysis.

773

	Species	Accession number	Name
Porifera	<i>Amphimedon queenslandica</i>	XP_003383519	Casp. Card
Cnidaria	<i>Acropora digitifera</i>	XP_015766400	Casp. X2
		XP_015768208	Casp. X1
	<i>Exaiptasia pallida</i>	KXJ12672.1	Casp. X1
		KXJ20965.1	Casp. X2
		KXJ12683.1	Casp. X3
	<i>Hydra vulgaris</i>	ACY95435.1	Casp. X1
		ACY95436.1	Casp. X2
Ctenophora	<i>Pleurobrachia bachei</i>	Sb 2658116	Casp. X
Placozoa	<i>Trochoplax adherens</i>	RDD44781.1	Casp. X
Lophotrochozoa	<i>Aplysia californica</i>	XP_005113266	Casp. 2
		XP_012945422.1	Casp. Y
	<i>Biomphalaria glabrata</i>	XP_013082356.1	Casp.
	<i>Capitella teleta</i>	ELU00616.1	Casp. 2
		ELT97848.1	Casp. Y
	<i>Crassostrea angulata</i>	AGN75137.1	Casp. 2
	<i>Crassostrea gigas</i>	XP_011419292	Casp. 2
		XP_011414267	Casp. Y1
		XP_011449817	Casp. Y3
		XP_011432762	Casp. Y2
	<i>Octopus bimaculatus</i>	XP_014783442.1	Casp. 2
		XP_014790087.1	Casp. Y1
		XP_014784582.1	Casp. Y2
Chelicerata	<i>Limulus polyphemus</i>	XP_013788138.1	Casp. 2
		XP_013776997.1	Casp. 2
Insecta	<i>Agrilus planipennis</i>	XP_018324425.1	Casp. 2
	<i>Apis mellifera</i>	XP_016771440.1	Casp. 2
	<i>Drosophila melanogaster</i>	CAB53565.1	Casp. 2
	<i>Habropoda laboriosa</i>	XP_017788772.1	Casp. 2
	<i>Helicoverpa armigera</i>	AEK20835.1	Casp. 2
	<i>Papilio machaon</i>	XP_014362236.1	Casp. 2
	<i>Tribolium castaneum</i>	XP_001813274	Casp. 2
Nematoda	<i>Ascaris suum</i>	ERG86894.1	Casp. 2
	<i>Caenorhabditis elegans</i>	AAG42045.1	Casp. 2
Vertebrata	<i>Homo sapiens</i>	AAH17839.1	Casp. 4
		AAI13407.1	Casp. 5
	<i>Mus musculus</i>	NP_033937.2	Casp. 1
		EDL13489.1	Casp. 2
		AAH56447.1	Casp. 9
	<i>Xenopus laevis</i>	NP_001081223.1	Casp. 1
		NP_001081404.1	Casp. 2
		NP_001079035.1	Casp. 9
Echinodermata	<i>Strongylocentrotus purpuratus</i>	XP_789183.3	Casp. 9A
		XP_011661242.1 1	Casp. 9B

		XP_011661359.1	Casp. 9C
Hemichordata	<i>Saccoglossus kowalevskii</i>	XP_006811879.1	Casp. 9
Cephalochordata	<i>Branchiostoma belcheri</i>	XP_019646757.1	Casp. 2A
		XP_019642903.1	Casp. 2B
		XP_019623612.1	Casp. 9A
		XP_019644208.1	Casp. 9B
	<i>Branchiostoma floridae</i>	EEN68002.1	Casp. 2
Urochordata	<i>Botryllus schlosseri</i>	Boschl.CG.Botznik2013.chrUn.g02816.01.p	Casp. 2A1
		Boschl.CG.Botznik2013.chrUn.g08767.01.p	Casp. 2A2
		Boschl.CG.Botznik2013.chrUn.g09831.01.p	Casp. 2B
	<i>Ciona intestinalis</i>	KH2012:KH.C4.463.v1.A.ND1-1	Casp. 2A
	<i>Ciona savignyi</i>	CISAVI-CG-ENS81-R15-461426-463207-09349-P	Casp. 2A
		CISAVI-CG-ENS81-R54-461275-467018-10343-P	Casp. 2B
	<i>Molgula oculata</i>	Moocul-CG-ELv1_2-S113854-g13164-01-p	Casp. 2A
		Moocul-CG-ELv1_2-S103067-g09526-01-p	Casp. 2B
	<i>Phallusia mammillata</i>	PHMAMM-CG-MTP2014-S310-G07060-01-P	Casp. 2A
		PHMAMM-CG-MTP2014-S92-G02989-01-P	Casp. 2B
	<i>Halocynthia roretzi</i>	HARORE-CG-MTP2014-S35-G03390-01-P	Casp. 2A
		HARORE-CG-MTP2014-S130-G07320-01-P	Casp. 2B

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

Supplemental Table 2: List of Apaf-1 among metazoans. N/A: No APAF-1 has been identified.

Species	Accession number
Porifera	<i>Amphimedon queenslandica</i>
Ctenophora	<i>Pleurobrachia bachei</i>
Placozoa	<i>Trichoplax sp.</i>
Cnidaria	<i>Acropora digitifera</i>
	<i>Exaiptasia pallida</i>
	<i>Hydra vulgaris</i>
Nematoda	<i>Caenorhabditis elegans</i>
Chelicerata	<i>Limulus polyphemus</i>
Insecta	<i>Agrilus planipennis</i>
	<i>Apis mellifera</i>
	<i>Drosophila melanogaster</i>
	<i>Habropoda laboriosa</i>
	<i>Helicoverpa armigera</i>
	<i>Papilio machaon</i>
	<i>Tribolium castaneum</i>
	<i>Aplysia californica</i>
	<i>Biomphalaria glabrata</i>
Lophotrochozoa	<i>Crassostrea angulata</i>
	<i>Crassostrea gigas</i>
	<i>Octopus bimaculatus</i>
	<i>Capitella teleta</i>
Vertebrata	<i>Homo sapiens</i>
	<i>Mus musculus</i>
	<i>Xenopus laevis</i>
Echinodermata	<i>Strongylocentrotus purpuratus</i>
Hemichordata	<i>Saccoglossus kowalevskii</i>
Cephalochordata	<i>Branchiostoma belcheri</i>
	<i>Branchiostoma floridae</i>
Urochordata	<i>Botryllus schlosseri</i>
	<i>Ciona intestinalis</i>
	<i>Ciona savignyi</i>
	<i>Molgula oculata</i>
	<i>Phallusia mammillata</i>
	<i>Halocynthia roretzi</i>

Supplemental table 3: List of Bcl-2 used for phylogenetic analysis.

Species		Accession number	Names
Porifera	<i>Amphimedon queenslandica</i>	XP_003383425.1	Bcl-2 like 1
		XP_003387574.1	Bcl-2 like 2
Cnidaria	<i>Hydra vulgaris</i>	EF104645	Bak 1
		EU035760	Bak 2
		XP_012562061.1	Bax
		EU035764	Bcl-2 like 1
		EF104646	Bcl-2 like 2
		EU035765	Bcl-2 like 3
		EU035763	Bcl-2 like 4
		EU035761	Bok
		EF104647	Mcl-1 1
		EU035762	Mcl-1 2
Nematoda	<i>Caenorhabditis elegans</i>	AAA20080.1	Ced-9
Insecta	<i>Drosophila melanogaster</i>	AAF44120.1	Buffy (Bok-1)
		AAF26289.1	Buffy (Bok-2)
Lophotrocozoa	<i>Biomphalaria glabrata</i>	XP_013085524.1	Bax 1
		XP_013096338_1	Bax 2
		XP_013086802.1	Bax 3
		XP_013070177.1	Bak
		XP_013081872.1	Bcl-2 like 1
		XP_013093137.1	Bcl-2 like 2
		XP_013068612.1	Bcl-2 like 3
		XP_013083436.1	Bcl-2 like 4
		XP_013069706.1	Bok
		XP_013065969.1	Mcl-1
		XP_011424481_1	Bax
		XP_011439700_1	Bak
		XP_011449013_1	Bcl-2 like 1
		ACH42081_1	Bcl-2 like 2
		EKC18663_1	Bcl-2 like 3
		XP_011436990_1	Mcl-1 1
		EKC40007_1	Mcl-1 2
	<i>Crassostrea gigas</i>	XP_011424481_1	Bax
		XP_011439700_1	Bak
		XP_011449013_1	Bcl-2 like 1
		ACH42081_1	Bcl-2 like 2
		EKC18663_1	Bcl-2 like 3
		XP_011436990_1	Mcl-1 1
		EKC40007_1	Mcl-1 2
		XP_011424481_1	Bax
		XP_011439700_1	Bak
		XP_011449013_1	Bcl-2 like 1
Vertebrata	<i>Homo sapiens</i>	AAA74466.1	Bak
		NP_001278357.1	Bax
		API71171.1	Bcl-2
		AAK48715.1	Bcl-B
		AAB09055.1	Bcl-w
		CAA80661.1	Bcl-xL
		NP_004040.1	Bfl-1
		NP_115904.1	Bok
		AAF64255.1	Mcl-1
	<i>Mus musculus</i>	NP_031549.2	Bak
		NP_031553.1	Bax
		AAH95964.1	Bcl-2

		Q9Z0F3.1	Bcl-B
		AAA51039.1	Bcl-xL
		AAB09056.1	Bcl-w
		Q07440.1	Bfl-1
		NP_058058.1	Bok
		NP_032588.1	Mcl-1
	<i>Xenopus laevis</i>	NP_001089587.1	Bak
		AAR84081.1	Bax
		BAH28834.1	Bcl-2
		AAI10791.1	Bcl-xl
		XP_018089640.1	Bcl-w
		NP_001139563.1	Bok
		ACI47310.1	Mcl-1
Urochordata	<i>Ciona intestinalis</i>	KH2012:KH.C4.794.v1.A.SL1-1	Bax
		KH2012:KH.S653.2.v2.A.SL1-1	Bcl-xl
		KH2012:KH.L87.39.v1.A.ND1-1	Bok
	<i>Molgula oculata</i>	Moocul.CG.ELv1_2.S96550.g08147.01.p	Bax
		Moocul.CG.ELv1_2.S112899.g12639.01.p	Bok

798
799
800
801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

KEY RESOURCES TABLE

Software and Algorithms		
ScanProsite	Gattiker et al., 2002 ⁸⁵	https://prosite.expasy.org/scanprosite/
InterProScan	Quevillon et al., 2005 ⁸⁶	https://www.ebi.ac.uk/interpro/search/sequence/
MAFFT 7	Katoh et Standley, 2013 ⁸⁷	https://mafft.cbrc.jp/alignment/server/
Clusta Omega	Sievers et al., 2011 ⁸⁸	https://www.ebi.ac.uk/Tools/msa/clustalo/
BioEdit 7.04	Hall, 1999 ⁸⁹	https://bioedit.software.informer.com/7.2/
Gblocks 0.91b	Castresana, 2000 ⁹⁰	http://molevol.cmima.csic.es/castresana/Gblocks_server.html
PhyML 3.1	Guindon et al., 2005 ⁹¹	http://www.atgc-montpellier.fr/phyml/versions.php
Seaview	Gouy et al., 2010 ⁹²	http://doua.prabi.fr/software/seaview
Mega11	Tamura et al., 2021 ⁹³	https://www.megasoftware.net/
MrBayes 3.1.2	Ronquist et Huelsenbeck, 2003 ⁹⁴	https://nbisweden.github.io/MrBayes/download.html