Continental-scale Hyperspectral tree species classification in the National Ecological Observatory Network

Sergio Marconi¹, Ben. G. Weinstein¹, Sheng Zou³, Stephanie A. Bohlman², Alina Zare³, Aditya Singh⁴, Dylan Stewart³, Ira Harmon⁵, Ashley Steinkraus¹, Ethan P. White¹

¹Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, Florida, USA

²School of Forest, Fisheries and Geomatics Sciences, University of Florida, Gainesville, Florida, USA

³Department of Electrical and Computer Engineering, University of Florida, Gainesville, Florida, USA

⁴Department of Agricultural & Biological Engineering, University of Florida, Gainesville, FL, USA

⁵Department of Computer and Information Sciences and Engineering, University of Florida, Gainesville, FL, USA

Abstract

Advances in remote sensing imagery and computer vision applications unlock the potential for developing algorithms to classify individual trees from remote sensing at unprecedented scales. However, most approaches to date focus on site-specific applications and a small number of taxonomic groups. This limitation makes it hard to evaluate whether these approaches generalize well across broader geographic areas and ecosystems. Leveraging field surveys and hyperspectral remote sensing data from the National Ecological Observatory Network (NEON), we developed a continental extent model for tree species classification that can be applied to the entire network including a wide range of US terrestrial ecosystems. We compared the performance of the generalized approach to models trained at each individual site, evaluating advantages and challenges posed by training species classifiers at the US scale. We evaluated the effect of geography, environmental, and ecological conditions on the accuracy and precision of species predictions.

On average, the general model resulted in good overall classification accuracy (micro-F1 score), with better accuracy than site-specific classifiers (average individual tree level accuracy of 0.77 for the general model and 0.72 for site-specific models). Aggregating species to the genus-level increased accuracy to 0.83. Regions with more species exhibited lower classification accuracy. Trees were more likely to be confused with congeneric and co-occurring species and confusion was highest for trees with structural damage and in complex closed-canopy forests. The model produced accurate estimates of uncertainty, correctly identifying trees where confusion was likely. Using only data from NEON this single integrated classifier can make predictions for 20% of all tree species found in forest ecosystems across the US, suggesting the potential for broad scale general models for species classification from hyperspectral imaging.

1. Introduction

Forest ecosystems play a central role in essential services like assessment of natural resources and biodiversity conservation at national scales (Wiens, 2016; Pecl et al., 2017), but understanding this role at the large range of scales over which ecological processes operate can be challenging. A common strategy to monitor biodiversity and biomass of forests at broad scales is to use field surveys of plots as a sample for landscape level characteristics (USDA Forest Service, 2001, Lawrence et al. 2010). Data collection within survey plots requires extensive effort, limiting even the most extensive national forest inventories to several thousand permanent plots sampled every few years (White et al., 2016), which can be too sparse for investigating the effects of local and landscape ecological properties on forest structure and diversity (Tomppo et al., 2008). Remote sensing can help bridge this gap between local and regional scales by providing individual tree level data at scales beyond what is feasible for traditional plot-level inventories (Anderson, 2018). Models linking remotely sensed imagery to field surveys can identify the location and species identity of individual trees (Henry & Jarvis, 2019), alleviating the challenge of inferring local patterns from sparsely sampled data (Ayrey et al., 2019, Bastin et al., 2019, Kandare et al., 2017) for understanding species distributions and abundances.

Numerous approaches have been developed for pixel- or canopy-scale species-level classification using hyperspectral remote sensing based on exploiting differences in foliar properties and canopy structure (Mayra et al, 2021, Belgiu & Dragut, 2016, Ballanti et al., 2016, Ab Majid et al., 2016). In general, however, most approaches are site- and/or ecosystem-specific (Fassnacht et al., 2016). Site and/or ecosystem-specific techniques have the potential to limit the generality of algorithms and make it difficult to: 1) conduct research at regional to continental scales due to the lack of general models that can be applied across ecosystems; 2) identify rare or uncommon species due to limited data for training models, which often results in

studies focusing on a limited subset of common species; and 3) accurately apply the model beyond the scale of the associated field data. Furthermore, training data from single site studies often lack the full range of variation in spectral characteristics that can occur for each species due to intraspecific variation and variability from the remote sensing acquisition process. Developing species classification approaches that combine data from across broad geographic extents has the potential to address these limitations and increase the utility of remote sensing models in duplicated effort across sites and projects.

Developing species classification models that span geographic areas, forest types and species pools poses a novel set of challenges. First, it requires building a library of coregistered field and remote sensing data that includes data from multiple sites and ecoregions for training and testing algorithms. Second, increasing the geographic extent of species classification risks confusing species with similar spectral properties that do not overlap in their geographic distributions. Third, combining data from multiple sites may introduce variation in spectral reflectance due to differences in phenology (which affect leaf greenness) and environmentally driven intraspecific variation, which affect leaf biochemistry (Sims & Gamon, 2002). Finally, aggregating remote sensing data from multiple flights, sensors, and sites may increase variation in spectral signatures due to complex sources of spatial and temporal variation that are linked, but not limited to, acquisition dates, solar angles, ecosystems types and variation in sensor calibrations (Pax-Lenney et al., 2001). Therefore, while there are many potential benefits to more broad scale models for species classification, it is unclear how they will perform compared to models developed for specific ecosystems.

Here we leverage newly available data from the National Ecological Observatory Network (NEON) to develop a continental level model for tree species classification that can be applied to the entire network and compare its performance to the traditional approach of building individual models for each site. We used NEON remote sensing and field data on individual trees at 30 terrestrial sites from Puerto Rico to Alaska, covering a wide range of ecoregions and

biomes across the United States (US). Several studies have developed species classification models for NEON data, but all these studies focused on individual NEON sites (Scholl et al., 2020, Fricker et al., 2019, Marrs & Ni-Meister, 2019, Marconi et al., 2020), or 2-3 sites in the same region (Rana et al. 20XX, Graves et al. 2021), and a small number of species within each site. Even outside of NEON, species classification efforts from airborne remote sensing focus on site-specific algorithms, making it hard to assess how well these approaches transfer to different regions (but see Rana et al., 2018). We build on this work to develop a general model that can be applied across the entire National Ecological Observatory Network by connecting field-identified tree stems to hyperspectral images then using an ensemble of machine learning algorithms to identify species. This ensemble-based approach allows for leveraging the strengths of different machine learning classifiers and provides effective ways to estimate the uncertainty of predictions. Using this model, we (1) assess whether a general model approach improves performance compared to separate models for each site, (2) determine the importance of reflectance, geography, environmental and ecological conditions on the accuracy and precision of species predictions; (3) evaluate the uncertainty in predictions; and (4) discuss the potential for this general model to be used for ecological applications from monitoring populations of individual species to understanding patterns of biodiversity.

2. Methods

2.1. Field Data

Vegetation structure field data (<https://data.neonscience.org/data-products/DP1.10098.001>) were collected by the NEON terrestrial observatory system (TOS) between 2015 and 2019 (Table S.1). This dataset, sampled from 400 m² plots distributed across the landscape of each NEON site, includes information about individual trees' geolocation and

properties such as species identity, health status, canopy position, crown diameter, and tree height. Vegetation structure plot locations are located randomly across the sites stratified by vegetation type within each site with the aim of capturing landscape level biodiversity and structural features at each site. Each subplot is assigned to an ecosystem type extracted from the National Land Cover Dataset. For this study we used data from 30 of the 41 NEON sites with partial to complete forest cover, encompassing 17 out of 18 ecoclimatic domains in the US (Figure S.1). Data from the other NEON sites could not be used because field data was either missing information about tree stem positions or the remote sensing imagery contained gaps in the hyperspectral or missing information about the sensor angle at the time of data collection. We only included individual stem data that met the following criteria: (a) the stem had a species label assigned to it, (b) it was marked as “alive” and “tree” in the NEON field inventory, and (c) it belonged to a species with more than 5 entries for the entire dataset. We also did not use stems designated in the NEON vegetation structure data as fully shaded, shrubs or sapling, as these stems are most likely not visible in the remote sensing imagery and would therefore be erroneously paired with pixels belonging to species from neighboring overstory crowns. The final dataset used for species classification consisted of 5867 individual trees of 80 species.

2.2. Remote sensing data

For this study we used the hyperspectral L3 data from the NEON Airborne Observatory Platform (NEON, 2021). This data consists of 1km² tiles with 426 channels recording reflectance in 5nm bands from 350 to 2450nm. Reflectance data was already atmospherically corrected using the ATCOR-4 approach (Krause et al., 2011). Pixel resolution is 1m². We applied bidirectional reflectance distribution function (BRDF) correction, topographic correction, and L2 normalization to reduce the effect of peripheral light and non-Lambertian scattering with

the goal of minimizing variation in reflectance ascribable to flight path and airplane position (Marconi et al., 2020). For all tiles ($n = 4500$), we used the same general parameterization to define the BRDF kernel. We also dropped bands in the water absorption regions of the spectra as well as the spectrometer's peripheral bands to reduce the effects of noise and artifacts. Thus, the hyperspectral data was reduced to a total of 350 channels. In the models, we included terrain elevation (1m^2 spatial resolution) along with the hyperspectral data because of elevation's potential information in discriminating species within landscapes (Strahler et al., 1978, Scholl et al., 2020). This data was derived from a LiDAR sensor mounted along with the hyperspectral sensor on the aircraft, which was converted into a 1m spatial resolution raster, and appended to the hyperspectral data as an additional band.

We assigned each individual tree from the filtered field dataset to a square clip of 16 pixels (4m crown diameter), centered around the stem's GPS coordinates. This threshold was selected because it is smaller than more than 95% of individual tree crowns diameter measured from the NEON vegetation structure dataset. We adopted this strategy to reduce the number of mislabeled pixels at the edges of the crown that belong to neighboring trees, especially in dense closed canopies. To remove shaded and non-vegetation pixels from these clips, we removed all pixels with $\text{NDVI} < 0.5$ and low reflectance in the NIR (reflectance at $825\text{nm} < 0.2$). Since stem positions often do not match precisely with the center of the tree crown in the canopy, pixels will sometimes be assigned to the wrong label (hereafter referred to as pixel pollution). To reduce pixel pollution, we filtered out pixels that were much shorter than the maximum height of the crown. These pixels are less likely to belong to the sunlit portion of the target crown or may even measure the reflectance from neighboring shorter tree crowns, or the understory within a gap in the target crown. We filtered out pixels that were $\geq 5\text{m}$ below the top height of the tree as determined by the maximum height of the tree from the LiDAR data in the 16-pixel clip. Finally, we removed stems with field GPS locations that fell within 3 meters of one another where the

stems belonged to different taxa to decrease the chance of confusing closely neighboring, and potentially intermixed tree crowns of different species. After all these steps, the final, filtered dataset used ~50,000 out of 200,000 initial pixels and 7464 out of ~21,000 crowns in the original vegetation structure dataset.

Due to the large number of bands in hyperspectral data it is necessary to reduce the number of features used in classifiers to reduce the number of correlated bands and limit the potential for overfitting (Li et al., 2011). We reduced the 356 hyperspectral bands to 45 spectral features using a hierarchical dimensionality reduction, consisting of clustering bands with similar standardized distributions according to Kullback-Leibler divergence (KLD) (Zare et al., 2019). We chose this approach because it is well suited as a dimensionality reduction tool for data in the form of probability densities like hyperspectral reflectance spectra (Delicado, 2011) and allows for a direct identification of the most informative spectral regions. For this study we used 15 spectral regions, and we used the minimum, mean and maximum of the reflectance in each region as the hyperspectral features for the species classifier, resulting in 45 spectral predictors.

2.3. Site effects

To provide the model with information on site location that could be useful to reduce confusion across species that do not co-occur in a site but are characterized by similar spectral signatures, we included the latitude and longitude of the centroid of each site in the model. This approach incorporates information on the proximity of different sites and can be readily generalized to use outside of NEON. To help control for potential differences resulting from variation in sensor calibration of the specific flight missions, which would be specific to each site, we added a “site identifier” to the remote sensing features in the model. The site identifier was calculated using a Leave-One-Out regression encoding strategy to transform site categories into real positive numbers (https://contrib.scikit-learn.org/category_encoders; Wright & König, 2019). The advantage of this approach over the commonly used one-hot-encoder (ref)

is that it compresses the information into a single feature (rather than one feature per site), which avoids undesired sparsity and potential overfitting due to a large number of encoded classes (30 in this study) (Rodriguez et al., 2018). We used data in the training set to fit the encoder and assigned its average value to each site category. The final model input for the general model was hyperspectral features, elevation, latitude and longitude and site. For the site-specific models, only spectral features and elevation were used.

2.4. Species classification

For both the general and site-specific models, we performed species classification at the pixel level using an ensemble of five classifiers (Figure S.2): (1) a random forest classifier (Belgiu & Dragut, 2016), (2) a k-nearest neighbors classifier (Laaksonen & Oja, 1996), (3) a histogram gradient boosting classifier (Guryanov, 2019), (4) a fully connected multilayer perceptron (Pacífico et al., 2018), and (5) a bagging classifier with support vector machine as base estimators, using tools from the scikit-learn python package (Pedrosa et al., 2011). Details for each classifier can be found in supplementary materials (Supplement 1: classifiers). Ensemble-based approaches generally provide better performance and limit overfitting compared to using one classifier alone (Knauer et al., 2019). We chose the individual classifiers to ensemble because they have been shown to perform well for species classification on NEON data (Marconi et al., 2019). All predictors were normalized for model fitting by subtracting the mean and dividing by the standard deviation (i.e., setting the mean to zero and the standard deviation to 1). Parameters for all models and the ensemble were extracted by performing cross validation.

We used entropy loss to measure the quality of tree-splits for random forests, categorical cross-entropy as the loss function for the histogram-gradient boosting, a radial basis function kernel to allow for a non-linear decision surface for the support vector classifiers, and the

Manhattan distance for calculating the distance between k-nearest neighbors in the KNN classifier. We stacked these five pixel-based models by using the probability vectors produced by each classifier as features for a meta-ensemble elastic net logistic model (Tang et al., 2015, Hui & Hastie, 2005). We chose this approach because logistic classifiers are easily interpretable and use maximum likelihood to obtain estimates of the coefficients, returning as a result confidence scores that match the probability of a label-match and not just the single best predicted classification (Maddala, 1986), which is fundamental for assigning each prediction to a reliable uncertainty score. Pairing predictions to robust estimates of uncertainty is fundamental to increase the utility of remote sensing tree surveys for ecological analysis because it allows for (1) selecting trees and areas that meet or exceed minimum confidence in the derived measures for being used for scientific analyses, and (2) allows for cascading the uncertainty in predictions onto the results of analyses downstream (Dietze, 2017). Training the logistic meta-ensemble on calibrated scores from sub-classifiers offers an advantage over other modern algorithms, whose estimates of uncertainty do not match true probabilities and are not well calibrated to the output of interest (Guo et al., 2017, Mukhati et al. 2020).

One of the main challenges of species classification algorithms is the imbalance between number of individual samples for rare and common species, which can cause models to overfit to highly abundant classes. In our data set, the number of pixels per species ranged from 44-28000 and the number of individual trees per species ranged from 5-1000. We used SMOTETomek technique (Batista et al., 2003) to reduce the effects of species class imbalances in the training set. SMOTETomek consists of a combination of under and oversampling which resulted in roughly 1000 spectral signatures (pixels) per species. First we undersampled pixels from the most abundant species using Tomek links, which removes noisy and borderline pixels (Tomek, 1976). Then, we used a SMOTE oversampling approach (Chawla et al., 2002) to create non-identical synthetic pixels for any species with fewer pixels than the

majority class, thus balancing each class to roughly 1000 pixels each. No over-undersampling was applied to the test data. Because of the stratified design of the train-test split, most species and sites had a number of test trees proportional to their frequency in the original dataset. We also used the same train-test split to repeat the entire analysis once for each NEON site by building and testing site-specific models built using only data from each particular site.

2.5. Evaluation

We evaluated the performance of the models by training the model on 80% of the data and evaluating its performance on the remaining 20%. Since spatial autocorrelation across train and test data can lead to optimistic bias in classification (Millard & Richardson 2015), we placed all individuals within a plot together into either the training or testing data sets. A series of randomizations of the plots were performed to create a 80:20 split of individuals that optimized the number of species in the train and test data sets. For each randomization, we calculated the total number of species in the test set and repeated this random operation until we found the split which maintained the highest number of species from the original data in both train and test set. For the general model, the training data set contained 5867 individuals of 80 species and the test data set 1599 individuals of 75 species. The resulting data represents 56% of the total tree species in the original unfiltered vegetation structure dataset and these species account for an average of 89% of individuals per site (Figure S.3).

Predictions for the species class of each individual in the test set were made using a 4x4 clip centered on the location of the test stem. Model performance was then evaluated using overall accuracy, individual tree level (micro) and average species-level (macro) F1 scores (hereafter referred to as individual-level and species-level accuracy respectively). The F1 score combines precision and recall to provide a general measure of the overall accuracy of the

species classification, allowing for direct comparison between models using a single metric (Chinchor, 1992). For each site, F1 scores for the general model were compared to those produced by equivalent single site models to determine how the general model performed relative to the traditional single site approach.

To understand the performance of the general model in different ecological contexts, we evaluated how performance varied across the United States, how performance correlated with the number of species being predicted at the NEON site, and which components of the model (site effects, elevation, geographic location, and hyperspectral reflectance) were most important for prediction. We identified the relative importance of site, geolocation, elevation and reflectance by performing bootstrap features importance (Breiman, 2001). This approach is based on evaluating how the overall accuracy is affected by each individual feature. At every iteration, one feature is selected and the values are randomly shuffled among the samples, effectively removing the information held in it. The accuracy is recorded with the shuffled feature to determine the loss of performance compared to the unshuffled data. We used the same approach to quantify the relative importance of the 15 spectral regions in which we grouped the hyperspectral data.

We also evaluated the characteristics of trees and forests associated with the most confusion between species (i.e., misclassification) based on forest type (using the National Land Cover Database; Homer et al., 2001) and information from the NEON field data on canopy position, tree status, and growth form from the NEON field data. We also assessed spatial structure in confusion by determining, for every misclassified tree, whether the species it was incorrectly classified as also occurred in the same NEON field plot. Finally, since confusion commonly occurred within genera we also evaluated model performance for predicting genus instead of species.

2.6. Prediction

We generated predictions for individual trees at the landscape scale (~350 km²) by integrating our approach with individual tree detections from previous work (Weinstein et al., 2021). The Weinstein et al. (2021) dataset consists of 100 million individual tree crowns from 37 NEON sites identified using a retinanet neural network object detector and represented by quadrangular polygons (i.e. bounding boxes) roughly representing the surface of the sunlit portion of the crown. For consistency between the approach used for training and testing the model (16-pixel clips), we extracted the pixels from the centroid of each estimated bounding box. First, we extracted a 4x4 square window of pixels around the centroid of each detection. For bounding boxes smaller than 16m², we dropped the pixels falling outside the bounding boxes. Second, we filtered vegetation pixels from the background using the same procedure as applied to the training/test data set. We finally selected all pixels with uncertainty scores > 0.5 to be used to make predictions at the level of individual trees. We assigned each tree to a species class by averaging the probability vectors (i.e., probability that the pixel is assigned to any of the 80 classes) of each pixel in the crown and selecting the species with the highest average probability. We assigned each individual-tree prediction an uncertainty score consisting of the average pixel probability, which ranged from 0-1.

3. Results

The general (cross-site) model yielded more accurate species classifications (larger F1 scores) than site-level models for 17 (species-level F1) and 18 (individual-level F1) of the 30 sites and identical accuracies for 8 (species-level F1) and 10 (individual-level F1) additional sites. There were only two sites that showed better site-level species-level and individual-level F1 scores: Blandy Experimental Farm, Washington (BLAN) and Rocky Mountain National Park,

Colorado (RMNP) (Figure 1, Figure S. 4, Figure S. 5). On average, the general model resulted in increases in the accuracy of individual tree level classification (increases in individual-level F1) from 0.72 to 0.77 and species-average accuracy (increases in species-level F1) from 0.46 to 0.54. Accuracy of the ensemble was higher than its sub-models trained singularly (raw outputs available at <https://doi.org/10.5281/zenodo.5796142>), which is consistent with the general observation that ensemble-based approaches produce more accurate predictions (Healey et al., 2018). Since the general ensemble model proved to be the best performing approach in this study, we focus primarily on it from this point forward.

Our results show a link between classification accuracy and ecological properties such as ecosystem type, tree health, and growth form (Figure 2, Figure S.6). Damaged trees, including broken boles and other types of damage (but not diseased trees), exhibited higher rates of misclassification than healthy crowns (Figure 2), with broken boles exhibiting a >50% misclassification rate. The general model performed best in shrub scrub and evergreen forests and worst in wetlands (~32% misclassification rate), with deciduous forests falling in between. Average classification accuracy was higher in eastern forests compared to western forests (Figure 3a), and was significantly correlated with the number of species within the site (Figure 3b,d). The algorithm generally underperformed in the Prairie Peninsula and Central and Southern Plains ecoregions which are characterized by patches of closed forest at the edges of prairies or farmland (Figure 3a, Figure S.6). These results align with previous work in showing that classification from remote sensing is more challenging for more complex canopies, overlapping crowns, and coexisting species with similar life history and spectral properties (Heinzel & Koch 2016, Bioucas-Dias, 2013).

Roughly 80% of the information used by the algorithm for classifying species was from the hyperspectral reflectance (Figure 4). Important information was present across the entire spectrum, but our results show that each spectral region is characterized by groups of more informative bands. Specifically, the most important band groupings are the blue and green

(0.450 to 0.550 nm) in the visible region, the red-edge in the near infrared (0.62 to 0.85), 1.15 to 1.27 nm in SWIR1 and 1.62 to 1.68 nm in SWIR2. The site's coordinates, which represented the geographic location of sites, explained 11% of total variation and were the second most important variable (Figure 4). Elevation, a proxy of potential local changes in the environment within each site, accounted for another 4%. The site effect, a proxy of other site level ancillary information (e.g. sensor calibration, flight and atmospheric conditions), only accounted for 3% of the total explained variance.

Comparing misclassification among species shows there is greater confusion for rare species, congeners, and species that co-occur within NEON field plots, and that model-estimated uncertainty accurately reflects confidence in the model prediction. All species performing poorly ($F1 < 0.5$) belonged to taxa with low sample sizes (less than 50 trees for training) (Figure S6, Figure S7). In general, most of the confusion was among species co-occurring within plot (74%) and site (93%). A large amount of confusion also occurred among congeneric species (~27% of total misclassifications), mostly within pines, poplars, oaks and maples, which make up 57% of the test dataset (Figure S.9). Oaks, pines and poplars in particular accounted for ~87% of the total within genus confusion, and most misclassifications had confidence scores >0.8 . Aggregating predictions at the genus level improved the overall accuracy by 6% (individual-level F1 accuracy of 83%), confirming that part of the confusion is embedded in physiological similarities across taxonomically related trees. The model produced reliable estimates of uncertainty for all species regardless of the accuracy. Uncertainty scores matched closely with the probability of correct classification ($R^2 = 0.89$, Figure 5).

4. Discussion

Using a single general model that integrated data across a continental scale resulted in more accurate classification of tree species identity from remote sensing data than building separate models for individual forest sites. The more accurate classification occurred despite the continental data set containing many different forest types, structures and species compositions across 30 sites. This suggests that the benefits of increasing the number of samples for less common species and more fully characterizing within-species variance outweighs the costs associated with including species that do not overlap geographically and including components of within-species variance not observed at individual sites (Figure 1). To our knowledge this is the first study to attempt to develop a generalized model across dozens of sites and the success of the general model here suggests that developing generalized algorithms offers a potential step forward in species classification from remote sensing more broadly. Our generalized approach leveraged the information from multiple locations and survey efforts, increasing the number of individuals from rarely sampled or highly variable classes and allowing models to learn more broadly about how to distinguish species in the taxonomic group of interest. In addition to yielding improved predictions, generalized cross-site approaches can potentially generate predictions for a wide range of ecosystems, including those with limited or no training data, allowing other studies to leverage the same shared model and thereby facilitating large-scale ecological research (Weinstein et al. 2021).

By virtue of classification of the most common tree species occurring in the canopy, the results of this model are potentially useful for several ecological applications, such as mapping biomass and modeling carbon, energy and water flux. Our model captured species making up ~80% of the individual trees in the upper canopy, although this varied among sites. Furthermore, given the stratified sampling of the NEON vegetation structure data used to develop the generalized model, the model is likely to capture the major vegetation types and

most common species at each site. Canopy trees, which are visible from optical remote sensing devices, contribute the majority of biomass in forests (Lutz et al., 2012). Because they form the interface between the atmosphere and land surface, the canopy layer also is particularly important for water and energy flux (Paul-Limogens et al., 2017). Because carbon storage, water and energy flux can vary among species (Wright et al., 2006), the ability to map the location and coverage of canopy species is important for assessing these important ecosystem characteristics. Other ecological applications, such as assessing total forest species richness, and quantifying tree regeneration, cannot be addressed using the model because our model could not classify rare canopy species or understory individuals,

One of the main challenges in developing models that generalize well across the continent is overcoming differences across sites in factors including seasonality, background, and sensor calibration (Hesketh & Sanchez-Azofeifa, 2012, Clark et al., 2005, Pu, 2021). To quantify the sensitivity of the algorithm to this ancillary information, we evaluated the relative importance of the site-effect features compared to reflectance, geography and elevation. Our results showed that the relative importance of the site-effect is marginal and accounts for less than 3% of the total information captured (Figure 4). This suggests that the spectral signal from NEON data is comparable across different flights and that flight-specific noise can be minimized using BRDF corrections and vector normalization to limit the impact on the accuracy of generalized algorithms. This is due in part to NEON data being highly standardized and using the same image pre-processing protocol across the entire network (Kampe et al., 2014). NEON remote sensing data is also collected at the peak of vegetation productivity for each site, reducing the confounding effect of different phenological stages for species occurring at multiple sites (Gartner et al., 2016). Expanding large scale surveys outside the NEON network would require integrating information from less standardized sources, raising new challenges related to fusion of sensors that are not cross-calibrated and images collected in different seasons (Brook & Ben-Dor, 2015, Zou et al., 2018). Further investigation is therefore fundamental to evaluating

whether our findings apply to applications that involve integrating multiple sensors, missions, or resolutions.

One of the advantages to broad scale general models is that they allow assessment of how different ecological and environmental conditions influence the quality of the species classification. Understanding variation in model performance across space, forest types, and taxa is fundamental to better understanding where and when these models can be applied and improvement of large-scale surveys from remote sensing. In our analysis, eastern forests showed lower accuracy compared to western ecosystems. We believe this is at least partly because eastern ecosystems are characterized by the higher species diversity of canopy trees as well as crown geometry that makes aligning stems to crowns more difficult compared to western conifer stands (Figure 2, 3, S.6). Higher species diversity in eastern forests (mean species per site ~15) compared to western forests (mean species diversity per site ~4), inherently makes classification tasks more challenging due to larger numbers of classes typically resulting in lower accuracy predictions (Takahashi et al., 2020). Continuous closed canopies also increase the likelihood pixels selected in a window centered on the stem will be from neighboring tree crowns. This is due to the difficulty of obtaining accurate GPS points of stems in closed canopy (Rodriguez-Perez et al., 2007), as well as the increased likelihood of sunlit portions of the crown being displaced from the stem location in continuous broadleaf forests (Strigul et al., 2008). This is a common problem, since field surveys often provide only the geographic coordinates of tree stems and lack information about crown position or size, making it impossible to correctly align crown borders with species labels. For example, pixel mislabeling may be one of the reasons why our classifier was weaker at sites in the Great Plains region (e.g., the NEON sites of Lyndon B. Johnson National Grasslands, CLBJ and University of Kansas Field Station, UKFS), where patches of grasslands alternate with dense forests characterized by multiple oak species forming a complex mosaic of crowns that may have shifted their stem locations. In contrast, conifers in western US forests tend to be

dominated by species characterized by apical dominance (e.g. aspens and firs) with crowns centered directly above the main stem reducing pixel mislabeling and improving classification. Finally, savannas, such as the San Joaquin Experimental Range (SJER), characterized by isolated trees of few species (mostly broadleaves) may be less likely to suffer from confounding effects like crown displacement and stem-crown misalignment, making them less prone to spectral mixing or potential pixel-mislabeling (Heinzel & Koch, 2012). The most challenging ecosystem type in our analysis, wetlands, combines all of these challenges. Species like *Carpinus caroliniana* and *Betula papyrifera*, found often in plots from wetland ecosystems, were among the species with the worst classification accuracy, partly because they are generally smaller trees that can occur in the understory, grow in closed canopies in the overstory (e.g. an average dbh ~16.5cm and average height of ~10m), and often include limited training samples because they are found in riparian ecosystems which make up a small fraction of the landscapes from the NEON sites included in this study (less than 50 individuals per species). Because of these challenges, the accuracy of the species predictions need to be assessed depending on the site and ecosystem types within sites to ensure it is sufficient for the intended ecological application.

Because species predictions from remote sensing are imperfect, it is important that classification models produce robust estimates of uncertainty to allow this uncertainty to be propagated through ecological analyses and considered during decision making. This is particularly important when generating large numbers of predictions at large scales, because this will result in including species located in undersampled areas and challenging ecosystems as well as species that are difficult to classify due to rarity or similarity to other closely related species. Our results confirmed that stacking scores from different classifiers using a logistic regression produces accurate estimation of classification uncertainty (Figure 6).

While our generalized approach resulted in significant improvements over site-level models, it is important to recognize that the accuracy of this approach is still insufficient for

ecological analyses contingent on rare or untrained species. For example, biodiversity patterns are often driven by rare species (Leitao et al., 2016, Mouilllot et al., 2013), which are the most challenging taxa for species classification, especially species so rare that they cannot be included in the model due to data limitations ($n < 5$ individual trees in this study). Extrapolating outside of NEON sites, a goal for general models, would also result in the presence of additional species missing from the field dataset, restricting the range of ecological analyses to species sampled within the footprint of NEON sites. Some of these limitations may be mitigated by classifying trees into higher level taxonomic levels. In this study we observed that misclassified trees were generally limited to species in the same genus and species co-occurring in the same plot (Figure S.9, Figure S.10). Oaks, pines and poplars in particular accounted for ~87% of the total within genus confusion. One possible driver of confusion among oak species is their similar physiological and spectral characteristics (Figure S.11). Some co-occurring oaks species like *Quercus alba* and *Quercus stellata* can also cross-breed and therefore be physiologically very similar (Hardin, 1975), making them particularly hard to distinguish from imagery. For these reasons, most of the cases leading to misclassification resulted from within-genus confusion. This implies that uncertainty can be significantly reduced by aggregating predictions to the genus level, offering a more robust solution for large scale ecological applications that can be successfully addressed by accurately classifying trees at the level of genus, families or plant functional type. For example, earth system models use plant functional types as the taxonomic unit for quantifying carbon dynamics at continental to global scale (Lawrence et al., 2019), large scale fire risk assessment and management can be achieved by using genus level surveys of the most dominant taxa (Ma et al., 2021), and patterns of forest biomass largely depending on which taxa dominate the ecosystem (Cheng et al., 2018). An increase in taxonomic level also reduces issues with applying general models beyond the training data (e.g., outside of NEON sites) because it is much more likely that all genera or families have been sampled in the training data.

Building generalized algorithms provides an approach to overcome the significant field data limitations present in most remote sensing tasks in ecology, by allowing pooling data from ever growing sources of spatially explicit field surveys and high-resolution remote sensing imagery. Our results showed that by integrating field surveys from dozens of NEON sites, it is possible to produce a general model that provides improvements over single-site models for species classification, with good estimates of uncertainty, and the ability to increase accuracy further by aggregating predictions at the genus level. This general approach also unlocks the potential for making predictions outside of NEON sites. The ultimate goal is to develop general models that can be used anywhere in the region of interest (in our case the United States). Using only NEON data, we successfully built a single integrated classifier that includes 20% of all tree species found in forest ecosystems across the US (n=80 out of 396 surveyed by the United States Forest Inventory and Analysis project; appendix F; Woudenberg, et al., 2010). Beyond NEON, more and more openly available field and hyperspectral datasets are being released from aerial (airborne and UAV) and satellite hyperspectral missions worldwide (Cook et al., 2013, Vangi et al., 2021, Claverie et al., 2018). Our results show that instead of training hundreds of separate models for local applications, there is the potential for integrating field and remote sensing collections from multiple locations and sources to build general models with improved accuracy for a broader range of landscapes and geographic locations. Further integration with more field and remote sensing datasets could potentially provide remote sensing based surveys of hundreds of millions of trees, making it possible to investigate the properties of ecosystems from local to continental scales.

5. Conclusions

Remote sensing is facing a revolution in data quality and methods accuracy, making it a good candidate for developing applications to survey species assembly and forest properties at scale. Leveraging data collected from NEON across the US, we demonstrated that building continental scale algorithms for generalized species classification offers several advantages over the more traditional site level applications. Despite being very high for dominant taxa, accuracy in predictions for less represented species can be taunted by limitations in field-to-image misalignment, the number of species and individuals from rarely sampled taxa, making surveys from remote sensing unsuited to date for analyzing patterns in species alpha diversity at scale. Yet, building generalized algorithms is a fundamental cornerstone to overcome these limitations, because it allows for pooling from ever growing sources of geo-explicit field surveys and high resolution remote sensing imagery. Our results showed that by integrating field surveys with NEON airborne data, it is possible already to generate highly accurate predictions at the genus level and overall good estimates of uncertainty for individual trees. This allows for generating surveys of hundreds of millions of individual crowns across the continent, unlocking the potential for investigating large scale ecological applications focusing on the sun-exposed part of the canopy, dominant species, genres or functional types.

Acknowledgements

This work was supported by the Gordon and Betty Moore Foundation's Data-Driven Discovery Initiative through grant GBMF4563 to E. P. White and by the National Science Foundation through grant 1926542 to E. P. White, S. A. Bohlman, A. Zare, D. Z. Wang, and A. Singh; by the NSF Dimension of Biodiversity program grant (DEB-1442280) and USDA/NIFA McIntire-Stennis program (FLA-FOR-005470) to S. A. Bohlman; by the University of Florida

Biodiversity Institute (UFBI) and Informatics Institute (UFII) Graduate Fellowship to Sergio

Marconi. There was no additional external funding received for this study.

All data can be found in the following Zenodo archive:

<https://doi.org/10.5281/zenodo.5796143>. All code for data preprocessing, model training and

testing and analyses can be found in the following GitHub repo:

<https://github.com/MarconiS/Continental-scale-Hyperspectral-tree-species-classification-in-the->

National-Ecological-Observatory-N

References

1. Ab Majid, Ibtisam, Zulkiflee Abd Latif, and Nor Aizam Adnan. "Tree species classification using worldview-3 data." 2016 7th IEEE Control and System Graduate Research Colloquium (ICSGRC). IEEE, 2016.
2. Alina Zare, Susan Meerdink, Yutai Zhou, Caleb Robey, Ron Fick, John Henning, & Paul Gader. (2019, April 12). GatorSense/hsi_toolkit_py: Initial Release (Version v1.0). Zenodo. <http://doi.org/10.5281/zenodo.2638117>
3. Anderson-Teixeira, K.J., Davies, S.J., Bennett, A.C., Gonzalez-Akre, E.B., Muller-Landau, H.C., Joseph Wright, S., Abu Salim, K., Almeyda Zambrano, A.M., Alonso, A., Baltzer, J.L. and Basset, Y., 2015. CTFS-ForestGEO: a worldwide network monitoring forests in an era of global change. *Global change biology*, 21(2), pp.528-549.
4. Anderson, C.B. "Biodiversity monitoring, earth observations and the ecology of scale." *Ecology letters* 21.10 (2018): 1572-1585.
5. Ayrey, E., et al. "Synthesizing Disparate LiDAR and Satellite Datasets through Deep Learning to Generate Wall-to-Wall Regional Forest Inventories." *BioRxiv* (2019): 580514.
6. Ballanti, Laurel, et al. "Tree species classification using hyperspectral imagery: A comparison of two classifiers." *Remote Sensing* 8.6 (2016): 445.

7. Bastin, J.F., et al. "The global tree restoration potential." *Science* 365.6448 (2019): 76-79.
8. Belgiu, Mariana, and Lucian Drăguț. "Random forest in remote sensing: A review of applications and future directions." *ISPRS journal of photogrammetry and remote sensing* 114 (2016): 24-31.
9. Bioucas-Dias, José M., et al. "Hyperspectral remote sensing data analysis and future challenges." *IEEE Geoscience and remote sensing magazine* 1.2 (2013): 6-36.
10. Breiman, Leo. "Random forests." *Machine learning* 45.1 (2001): 5-32.
11. Brook, Anna, and Eyal Ben-Dor. "Supervised vicarious calibration (SVC) of multi-source hyperspectral remote-sensing data." *Remote Sensing* 7.5 (2015): 6196-6223.
12. Chawla, Nitesh V., et al. "SMOTE: synthetic minority over-sampling technique." *Journal of artificial intelligence research* 16 (2002): 321-357.
13. Cheng, Yanxia, et al. "Biomass-dominant species shape the productivity-diversity relationship in two temperate forests." *Annals of Forest Science* 75.4 (2018): 1-9.
14. Chinchor, N. "MUC-4 evaluation metrics in Proc. of the Fourth Message Understanding Conference 22–29." (1992).
15. Claverie, Martin, et al. "The Harmonized Landsat and Sentinel-2 surface reflectance data set." *Remote Sensing of Environment* 219 (2018): 145-161.
16. Cook, Bruce D., et al. "NASA Goddard's LiDAR, hyperspectral and thermal (G-LiHT) airborne imager." *Remote Sensing* 5.8 (2013): 4045-4066.
17. Delicado, Pedro. "Dimensionality reduction when data are density functions." *Computational Statistics & Data Analysis* 55.1 (2011): 401-420.
18. Dietze, Michael C. "Prediction in ecology: A first-principles framework." *Ecological Applications* 27.7 (2017): 2048-2060.
19. Fassnacht, Fabian Ewald, et al. "Review of studies on tree species classification from remotely sensed data." *Remote Sensing of Environment* 186 (2016): 64-87.

20. Fricker, G. A., Ventura, J. D., Wolf, J. A., North, M. P., Davis, F. W., & Franklin, J. (2019). A convolutional neural network classifier identifies tree species in mixed-conifer forest from hyperspectral imagery. *Remote Sensing*, 11(19), 2326.
21. G. Batista, B. Bazzan, M. Monard, "Balancing Training Data for Automated Annotation of Keywords: a Case Study," In WOB, 10-18, 2003.
22. Gärtner, Philipp, Michael Förster, and Birgit Kleinschmit. "The benefit of synthetically generated RapidEye and Landsat 8 data fusion time series for riparian forest disturbance monitoring." *Remote Sensing of Environment* 177 (2016): 237-247.
23. Guo, Chuan, et al. "On calibration of modern neural networks." *International Conference on Machine Learning*. PMLR, 2017.
24. Guryanov, Aleksei. "Histogram-based algorithm for building gradient boosting ensembles of piecewise linear decision trees." *International Conference on Analysis of Images, Social Networks and Texts*. Springer, Cham, 2019.
25. Hardin, James W. "Hybridization and introgression in *Quercus alba*." *Journal of the Arnold Arboretum* 56.3 (1975): 336-363.
26. Healey, Sean P., et al. "Mapping forest change using stacked generalization: An ensemble approach." *Remote Sensing of Environment* 204 (2018): 717-728.
27. Heinzl, Johannes, and Barbara Koch. "Investigating multiple data sources for tree species classification in temperate forest and use for single tree delineation." *International Journal of Applied Earth Observation and Geoinformation* 18 (2012): 101-110.
28. Henrys, Peter A., and Susan G. Jarvis. "Integration of ground survey and remote sensing derived data: Producing robust indicators of habitat extent and condition." *Ecology and evolution* 9.14 (2019): 8104-8112.

29. Hesketh, Michael, and G. Arturo Sánchez-Azofeifa. "The effect of seasonal spectral variation on species classification in the Panamanian tropical forest." *Remote Sensing of Environment* 118 (2012): 73-82.
30. Homer, Collin, et al. "Development of a 2001 national land-cover database for the United States." *Photogrammetric Engineering & Remote Sensing* 70.7 (2004): 829-840.
31. Kampe, T., et al. "NEON imaging spectrometer geolocation processing algorithm theoretical basis document." NEON Doc.# 001290 Rev A (2014).
32. Kandare, K., et al. "Prediction of species-specific volume using different inventory approaches by fusing airborne laser scanning and hyperspectral data." *Remote Sensing* 9.5 (2017): 400.
33. Knauer, Uwe, et al. "Tree species classification based on hybrid ensembles of a convolutional neural network (CNN) and random forest classifiers." *Remote Sensing* 11.23 (2019): 2788.
34. Krause, Keith S., et al. "Early algorithm development efforts for the National Ecological Observatory Network Airborne Observation Platform imaging spectrometer and waveform lidar instruments." *Imaging Spectrometry XVI*. Vol. 8158. International Society for Optics and Photonics, 2011.
35. Laaksonen, Jorma, and Erkki Oja. "Classification with learning k-nearest neighbors." *Proceedings of International Conference on Neural Networks (ICNN'96)*. Vol. 3. IEEE, 1996.
36. Lawrence, David M., et al. "The Community Land Model version 5: Description of new features, benchmarking, and impact of forcing uncertainty." *Journal of Advances in Modeling Earth Systems* 11.12 (2019): 4245-4287.
37. Lawrence, M., et al. "Comparisons of national forest inventories." *National forest inventories*. Springer, Dordrecht, 2010. 19-32.

38. Leitaõ RP, Zuanon J, Ville´ger S, Williams SE, Baraloto C, Fortunel C, Mendonça FP, Mouillot D. 2016 Rare species contribute disproportionately to the functional structure of species assemblages. *Proc. R. Soc. B* 283: 20160084.
<http://dx.doi.org/10.1098/rspb.2016.0084>
39. Li, Wei, et al. "Locality-preserving dimensionality reduction and classification for hyperspectral image analysis." *IEEE Transactions on Geoscience and Remote Sensing* 50.4 (2011): 1185-1198.
40. Loizzo, R., et al. "PRISMA: The Italian hyperspectral mission." *IGARSS 2018-2018 IEEE International Geoscience and Remote Sensing Symposium*. IEEE, 2018.
41. Lutz, James A., et al. "Ecological importance of large-diameter trees in a temperate mixed-conifer forest." *PloS one* 7.5 (2012): e36131.
42. M. L. Clark, D. A. Roberts, and D. B. Clark, "Hyperspectral discrimination of tropical rain forest tree species at leaf to crown scales," *Remote Sensing of Environment*, vol. 96, no. 3–4, pp. 375–398, 2005.
43. Ma, Wu, et al. "Assessing climate change impacts on live fuel moisture and wildfire risk using a hydrodynamic vegetation model." *Biogeosciences* (2021).
44. Maddala, Gangadharrao S. *Limited-dependent and qualitative variables in econometrics*. No. 3. Cambridge university press, 1986.
45. Marconi, Sergio, et al. "Rethinking the fundamental unit of ecological remote sensing: Estimating individual level plant traits at scale." *bioRxiv* (2019): 556472.
46. Mäyrä, Janne, et al. "Tree species classification from airborne hyperspectral and LiDAR data using 3D convolutional neural networks." *Remote Sensing of Environment* 256 (2021): 112322.
47. Mouillot D, Bellwood DR, Baraloto C, Chave J, Galzin R, et al. Rare Species Support Vulnerable Functions in High-Diversity Ecosystems. *PLoS Biology*, 2013 DOI: 10.1371/journal.pbio.1001569

48. Mukhoti, Jishnu, et al. "Calibrating deep neural networks using focal loss." arXiv preprint arXiv:2002.09437 (2020).
49. NEON (National Ecological Observatory Network). Spectrometer orthorectified surface directional reflectance - mosaic, RELEASE-2021 (DP3.30006.001).
<https://doi.org/10.48443/qeae-3x15>. Dataset accessed from <https://data.neonscience.org> on March 7, 2021
50. Pacifico, Luciano DS, Valmir Macario, and Joao FL Oliveira. "Plant classification using artificial neural networks." 2018 International Joint Conference on Neural Networks (IJCNN). IEEE, 2018.
51. Paul-Limoges, Eugénie, et al. "Below-canopy contributions to ecosystem CO₂ fluxes in a temperate mixed forest in Switzerland." *Agricultural and Forest Meteorology* 247 (2017): 582-596.
52. Pax-Lenney, Mary, et al. "Forest mapping with a generalized classifier and Landsat TM data." *Remote Sensing of Environment* 77.3 (2001): 241-250.
53. Pecl, G.T., Araújo, M.B., Bell, J.D., Blanchard, J., Bonebrake, T.C., Chen, I.C., Clark, T.D., Colwell, R.K., Danielsen, F., Evengård, B. and Falconi, L., 2017. Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science*, 355(6332), p.eaai9214.
54. Pedregosa, Fabian, et al. "Scikit-learn: Machine learning in Python." *the Journal of machine Learning research* 12 (2011): 2825-2830.
55. Pu, Ruiliang. "Mapping Tree Species Using Advanced Remote Sensing Technologies: A State-of-the-Art Review and Perspective." *Journal of Remote Sensing* 2021 (2021).
56. Rana, Parvez, et al. "Towards a generalized method for tree species classification using multispectral airborne laser scanning in Ontario, Canada." *IGARSS 2018-2018 IEEE International Geoscience and Remote Sensing Symposium*. IEEE, 2018.

57. Rodríguez-Pérez, José R., M. F. Álvarez, and Enoc Sanz-Ablanedo. "Assessment of low-cost GPS receiver accuracy and precision in forest environments." *Journal of Surveying Engineering* 133.4 (2007): 159-167.
58. Scholl, V. M., Cattau, M. E., Joseph, M. B., & Balch, J. K. (2020). Integrating national ecological observatory network (neon) airborne remote sensing and in-situ data for optimal tree species classification. *Remote Sensing*, 12(9), 1414.
59. Sims, Daniel A., and John A. Gamon. "Relationships between leaf pigment content and spectral reflectance across a wide range of species, leaf structures and developmental stages." *Remote sensing of environment* 81.2-3 (2002): 337-354.
60. Strahler, A. H., Thomas L. Logan, and Nevin A. Bryant. "Improving forest cover classification accuracy from Landsat by incorporating topographic information." (1978).
61. Strahler, Alan H., Thomas L. Logan, and Nevin A. Bryant. "Improving forest cover classification accuracy from Landsat by incorporating topographic information." (1978).
62. Strigul, Nikolay, et al. "Scaling from trees to forests: tractable macroscopic equations for forest dynamics." *Ecological Monographs* 78.4 (2008): 523-545.
63. Takahashi Miyoshi, Gabriela, et al. "Evaluation of hyperspectral multitemporal information to improve tree species identification in the highly diverse atlantic forest." *Remote Sensing* 12.2 (2020): 244.
64. Tang, J., S. Alelyani, and H. Liu. "Data Classification: Algorithms and Applications." *Data Mining and Knowledge Discovery Series*, CRC Press (2015): pp. 498-500.
65. Tavares, P. A., Beltrão, N. E. S., Guimarães, U. S., & Teodoro, A. C. (2019). Integration of sentinel-1 and sentinel-2 for classification and LULC mapping in the urban area of Belém, eastern Brazilian Amazon. *Sensors*, 19(5), 1140.
66. Tomek, Ivan. "Two modifications of CNN." (1976).
67. Tomppo, E., et al. "Combining national forest inventory field plots and remote sensing data for forest databases." *Remote Sensing of Environment* 112.5 (2008): 1982-1999.

68. USDA Forest Service, 2001. Forest Inventory and Analysis National Core Field Guide, Volume I: Field Data Collection Procedures For Phase 2 Plots, Version 1.5. US Department of Agriculture, Forest Service, Washington, DC.
69. Vangi, Elia, et al. "The new hyperspectral satellite PRISMA: Imagery for forest types discrimination." *Sensors* 21.4 (2021): 1182.
70. Weinstein, B.G., Gardner, L., Saccomanno, V., Steinkraus, A., Ortega, A., Brush, K., Yenni, G., McKellar, A.E., Converse, R., Lippitt, C. and Wegmann, A., 2021. A general deep learning model for bird detection in high resolution airborne imagery. *bioRxiv*. <https://doi.org/10.1101/2021.08.05.455311>
71. White, J. C., et al. "Remote sensing technologies for enhancing forest inventories: A review." *Canadian Journal of Remote Sensing* 42.5 (2016): 619-641.
72. Wiens, J.J., 2016. Climate-related local extinctions are already widespread among plant and animal species. *PLoS biology*, 14(12).
73. Woudenberg, S. W., et al. "The Forest Inventory and Analysis Database: Database description and users manual version 4.0 for Phase 2." Gen. Tech. Rep. RMRS-GTR-245. Fort Collins, CO: US Department of Agriculture, Forest Service, Rocky Mountain Research Station. 336 p. 245 (2010).
74. Wright, Marvin N., and Inke R. König. "Splitting on categorical predictors in random forests." *PeerJ* 7 (2019): e6339.
75. Wright, Ian J., et al. "Cross-species patterns in the coordination between leaf and stem traits, and their implications for plant hydraulics." *Physiologia Plantarum* 127.3 (2006): 445-456.
76. Xi, Zhouxin, et al. "See the forest and the trees: Effective machine and deep learning algorithms for wood filtering and tree species classification from terrestrial laser scanning." *ISPRS Journal of Photogrammetry and Remote Sensing* 168 (2020): 1-16.

738 77. Zou, Hui, and Trevor Hastie. "Regularization and variable selection via the elastic net."
739 Journal of the royal statistical society: series B (statistical methodology) 67.2 (2005):
740 301-320.
741
742
743

Figure 1. Performance of generalized vs site-specific classification models for each NEON site. Positive values are sites for which the generalized model performed better than site-level. Negative values are sites for which the generalized model performed worse compared to site-level. Blue bars represent species-level F1 score, yellow bars individual-level F1. Numbers separated by (:) on top of each site name represent the total number of species in the training and test set specific for each site. Numbers separated by (/) in BLAN represent the different number of species in the test set for the site specific and general models.

Figure 2. Fraction of misclassified trees across ecosystem types (blue), growth form (yellow), canopy position (red) and health status (grey). Numbers above the x-axis labels are the number of trees in each category.

Figure 3. Variation in accuracy of the generalized algorithm across the US. (a) map of average individual-level F1 scores for each ecological domain. Dots represent the location of each NEON site. Blue polygons represent the Prairie Peninsula and Central-Southern Plains. (b) Relationship between individual-level F1 and number of species in the training dataset for each site. The blue line is the loess smoother relationship over the 29 sites. (c) Kernel density estimate of the distribution of individual-level F1 scores (averages per site). (d) Relationship between species-level F1 and number of trained species found in site; orange line is the loess smoother relationship over the 29 sites. (e) Kernel density estimate of the distribution of species-level F1 scores (averages per site).

Figure 4: Features importance calculated from the permutation feature importance procedures described in Breiman, 2001, on the meta-ensemble model. (a) Relative contribution of different feature types. reflectance (grey), site coordinates (yellow), elevation (blue) and site effect (red). (b) Relative importance of each Kullback-Leibler group used for dimensionality reduction of reflectance. Blue bars represent the reflectance for the average minimum, mean and maximum band in the specific KL group. Bar width represents the range of bands covered by the specific KL group. Color intensity represents the relative importance of the specific KL group for the classifier (from light blue being of little importance, to dark blue being highly important). Black lines represent the reflectance of a randomly selected pixel to illustrate a typical vegetation reflectance pattern. Reflectance was normalized using L2 normalization.

794

795 Figure 5. Evaluation of model confidence score (the probability of assigning the correct label to
796 a prediction) as a measure of uncertainty. Confidence score was binned into 34 equal-width
797 bins (each bin representing an interval of 0.03). Bin centers were plotted against the fraction of
798 trees in that confidence score bin that were correctly classified. The blue line shows the fitted
799 linear relationship between the confidence score and the proportion of correctly classified trees.
800 The black line is the 1:1 line.

801

802

Figure 6. Example of species classification maps for all individual trees at the Bartlett Experimental Forest (BART) NEON site in New Hampshire. Species in legend include six of the most abundant taxa predicted at the site. Individual crown boundaries were estimated using predictions from Weinstein et al., 2021. The background is gray scale imagery of the site, so the gray areas on the right hand panels are areas without any trees including roads and other open areas.

Supplement 1: parameterization of classifiers and meta ensemble

KNN classifier was trained using 20 neighboring points, with distance weighted by the inverse of their Manhattan distance. The Random Forest classifier was trained using 300 trees with up to 7 features (square root of the total predictors) considered for better split, validated using cross-entropy loss function on out-of-bag samples. The gradient boosting classifier was trained using 1000 maximum iterations, a learning rate of 0.01, max depth of 25 and 0.5 L2 regularization. Loss was calculated using categorical cross-entropy on out-of-bag samples. Multi-layer Perceptron classifier was trained for 1200 max iterations, using relu activation, 1 hidden layer, weight optimization through adam booster with exponential decay rate of 0.9. The Bagging Classifier was trained using 10 support vector machine classifiers as base estimators. We used loose regularization ($C = 1000$), RBF kernel, and 5-fold cross validation to calibrate probability estimates. The meta ensemble was trained using probability vectors produced by each weak classifier. We used a regularized logistic regression (elasticnet), with 0.5 L1 to L2 penalty ratio. We used a saga solver to optimize the loss function.

Supplementary Figures

Figure S1 Geographic distribution of NEON sites included for this study. Colored regions represent ecological regions defined by NEON (<https://www.neonscience.org/field-sites/about-field-sites>). A description of each site and their ecological domain can be found in the Supplementary Table 1.

Figure S. 2: Flowchart of the species classification pipeline developed for this study

849
850 Figure S.3 For each site, fraction of all species (red) and their relative frequency in number of
851 trees (blue) covered by the species included in the analysis, compared to the total number of
852 species (red) and individual trees (blue) included in the NEON vegetation structure dataset.
853 Numbers separated by (/) above each site name represent the total number of species in the
854 original dataset and in the filtered data respectively, specific for each site. Trees in the canopy
855 (blue bars) were determined by canopy position data in the vegetation structure data where
856 trees in the canopy were designated as "Full sun", "Mostly shaded", "Partially shaded", "Open
857 growth", or non classified ("NA")..

858

859

860

861 Figure S.4 Density functions of the difference in $\Delta F1$ scores between the generalized and each
862 single-site algorithm for species-level F1 (yellow) and individual-level F1 (blue). Positive $\Delta F1$
863 values (17 out of 29 sites) represent sites where the generalized algorithm outperformed its site-
864 specific counterpart. Dashed vertical lines represent the average $\Delta F1$ across sites (species-
865 level F1 = 0.09, individual-level F1 = 0.05).

866

867

868 Supplement S.5. Difference in accuracy between the general and site-specific approaches for
869 each species-site combination. Negative values (red) represent taxa whose accuracy is higher
870 in the general approach. Blue values represent taxa whose accuracy is higher in the site-
871 specific approach. White values where accuracy was similar for the general and site-specific
872 approaches. Grey are species that do not occur at the site. Sites are sorted by geographic
873 similarity.

Figure S.6 Example of 400m² plots for 6 sites from western (left panel), central (center panel) and eastern US (right panel). Dots represent field stem data collected from NEON vegetation structure. Different dot colors represent different species. Only stems that have been filtered to include only stems that are likely to be in the canopy. From top left to bottom right sites acronyms are Abby Road (ABBY), Delta Junction (DEJU), Harvard Forest (HARV), San Joaquin Experimental Range (SJER), Blandy Experimental Farm (BLAN), Lenoir Landing (LENO),

Figure S.7. Precision, Recall and F1 for each individual species included in the dataset. Results can be found in the supplement table in a separate supplementary file.

Figure S.8. The relationship between individual species F1 scores and number of individual trees available for training for that species. The blue line shows a fitted relationship using local polynomial regression fitting (loess) and the grey region shows the 95% confidence interval around that relationship.

Figure S.9. Confidence score $P(x)$ for those taxa each species was most confused with. Taxa with a $P(x)$ lower than 0.02 were not included.

907

908

909

910

Figure S.10. Distribution of species across sites.

Figure S.11 Example of spectral signature overlap between often confused congeneric species.

Lines represent the average spectra, shaded areas represent the standard deviation for all pixels extracted for that particular species. The first species in legend is in blue, the second in grey. The X-axis is the band number from brdf corrected hyperspectral image. Couples of species are: (a) *Quercus marilandica* and *Quercus stellata*, (b) *Quercus alba* and *Quercus rubra*, (c) *Pinus palustris* and *Pinus elliottii*, (d) *Abies magnifica* and *Abies lowiana*.

922

Site Name	Ecological Domain	Domain Name	State	Geolocation (Lat-Lon)
Abby Road NEON (ABBY)	D16	Pacific Northwest	Washington	45.76
				-122.33
Bartlett Experimental Forest NEON (BART)	D01	Northeast	New Hampshire	44.06
				-71.29
Blandy Experimental Farm NEON (BLAN)	D02	Mid Atlantic	Virginia	39.03
				-78.04
Caribou-Poker Creeks Research Watershed NEON (BONA)	D19	Taiga	Alaska	65.15
				-147.5
Dead Lake NEON (DELA)	D08	Ozarks Complex	Alabama	32.54
				-87.8
Delta Junction NEON (DEJU)	D19	Taiga	Alaska	63.88
				-145.75
Disney Wilderness Preserve NEON (DSNY)	D03	Southeast	Florida	28.13
				-81.44
Guanica Forest NEON (GUAN)	D04	Atlantic Neotropical	Puerto Rico	17.97
				-66.87
Harvard Forest & Quabbin	D01		Massachusetts	42.54

Watershed NEON (HARV)		Northeast		-72.17
KU Field Station NEON (UKFS)	D06	Prairie Peninsula	Kansas	39.04
				-95.19
Konza Prairie Biological Station NEON (KONZ)	D06	Prairie Peninsula	Kansas	39.1
				-96.56
Lenoir Landing NEON (LENO)	D08	Ozarks Complex	Alabama	31.85
				-88.16
Lyndon B. Johnson National Grassland NEON (CLBJ)	D11	Southern Plains	Texas	33.4
				-97.57
Moab NEON (MOAB)	D13	Southern Rockies / Colorado Plateau	Utah	38.25
				-109.39
Mountain Lake Biological Station NEON (MLBS)	D07	Appalachians / Cumberland Plateau	Virginia	37.38
				-80.52
Niwot Ridge NEON (NIWO)	D13	Southern Rockies / Colorado	Colorado	40.05
				-105.58

		Plateau		
Ordway-Swisher Biological Station NEON (OSBS)	D03	Southeast	Florida	29.69
				-81.99
Rocky Mountains NEON (RMNP)	D10	Central Plains	Colorado	40.28
				-105.55
San Joaquin Experimental Range NEON (SJER)	D17	Pacific Southwest	California	37.11
				-119.73
Smithsonian Conservation Biology Institute NEON (SCBI)	D02	Mid Atlantic	Virginia	38.89
				-78.14
Smithsonian Environmental Research Center NEON (SERC)	D02	Mid Atlantic	Maryland	38.89
				-76.56
Steigerwaldt-Chequamegon NEON (STEI)	D05	Great Lakes	Wisconsin	45.51
				-89.59
Talladega National Forest NEON (TALL)	D08	Ozarks Complex	Alabama	32.95
				-87.39
The Jones Center At Ichauway NEON (JERC)	D03	Southeast	Georgia	31.19
				-84.47
Treehaven NEON (TREE)	D05	Great Lakes	Wisconsin	45.49
				-89.59

University of Notre Dame Environmental Research Center NEON (UNDE)	D05	Great Lakes	Michigan	46.23
				-89.54
Wind River Experimental Forest NEON (WREF)	D16	Pacific Northwest	Washington	45.82
				-121.95
Yellowstone National Park NEON (YELL)	D12	Northern Rockies	Wyoming	44.95
				-110.54

Table S.1 Description of NEON sites and ecological domains used in this study.