New cell biological explanations for kinesin-linked axon degeneration

Yu-Ting Liew^{1*}, André Voelzmann¹, Liliana M. Pinho-Correia¹, Thomas Murphy¹, Haydn Tortoisshell¹, Jill Parkin¹, David M.D. Bailey², Matthias Landgraf², Andreas Prokop^{1#}

1) The University of Manchester, Manchester Academic Health Science Centre, Faculty of Biology, Medicine and Health, School of Biology, Manchester, UK

2) Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ

◆ Please, note that Y.-T.L. is the key contributor to this work as part of her PhD thesis (Liew, 2018). Unfortunately, we have not been able to establish contact for over two years, so that direct consent by this author to publication of this work has not been given. However, consent is assumed from prior discussions we had.

Author for correspondence: Andreas.Prokop@manchester.ac.uk

Running title: Kinesin-linked axonopathies

Key words: *Drosophila*, neurodegeneration, axons, microtubules, axonal transport, mitochondria, ROS

Abstract

Axons are the slender, up to meter-long projections of neurons that form the biological cables wiring our bodies. Most of these delicate structures must survive for an organism's lifetime, meaning up to a century in humans. Axon maintenance requires life-sustaining motor protein-driven transport distributing materials and organelles from the distant cell body. It seems logic that impairing this transport causes systemic deprivation linking to axon degeneration. But the key steps underlying these pathological processes are little understood. To investigate mechanisms triggered by motor protein aberrations, we studied more than 40 loss- and gain-of-function conditions of motor proteins, cargo linkers or further genes involved in related processes of cellular physiology. We used one standardised *Drosophila* primary neuron system and focussed on the organisation of axonal microtubule bundles as an easy to assess readout reflecting axon integrity. We found that bundle disintegration into curled microtubules is caused by the losses of Dynein heavy chain and the Kif1 and Kif5 homologues Unc-104 and Kinesin heavy chain (Khc). Using point mutations of Khc and functional loss of its linker proteins, we studied which of Khc's sub-functions might link to microtubule curling. One cause was emergence of harmful reactive oxygen species through loss of Milton/Miro-mediated mitochondrial transport. In contrast, loss of the Kinesin light chain linker caused microtubule curling through an entirely different mechanism appearing to involve increased mechanical challenge to microtubule bundles through de-inhibition of Khc. The wider implications of our findings for the understanding of axon maintenance and pathology are discussed.

Introduction

Axons are the long and slender processes of neurons which form the biological cables that wire the nervous system and are indispensable for its function. In humans, axons can be up to 2 metres long whilst displaying diameters of only 0.1-15µm (Prokop, 2020). Most of these delicate cellular processes must survive for an organism's lifetime, meaning up to a century in humans. Unsurprisingly, mammals lose about 40% of their axon mass towards high age (Calkins, 2013; Coleman, 2005; Marner et al., 2003). This rate is drastically increased in hereditary forms of axonopathies (Prokop, 2021).

Of particular interest to this article are mutations of microtubule-binding motor proteins that cause axonopathies, of which the OMIM® database (Online Mendelian Inheritance in Man®; Amberger et al., 2015) currently lists DYNACTIN 1 (ALS1, OMIM® reference #105400), DYNEIN HEAVY CHAIN 1 (CMT2A1, #614228), KIF1B (CMT20, #118210), KIF5A (SPG10, #604187; ALS, #617921), KIF1A (SPG30, #610357; HSN2C, #614213), KIF1B (CMT2A1, #118210) and KIF1C (SPAX2, #611302); in the case of KIF1A, links to HSP and ataxias are likely to be added soon (Nicita et al., 2020). The listed motor proteins are actively involved in live-sustaining axonal transport of RNAs, proteins, lipids and organelles (Guedes-Dias and Holzbaur, 2019; Hirokawa et al., 2010). Genetic aberration of such transport is thought to lead to systemic collapse of axonal structure and physiology, hence axonopathy. However, we have little understanding of the concrete mechanisms leading to these pathologies.

To bridge this important knowledge gap, we took a novel approach based on two strategic decisions: Firstly, we used axonal microtubules (MTs) as our main readout. These MTs are arranged into loose bundles that run all along axons to form the essential highways for axonal transport and to provide a source of MTs for axon growth and branching at any life stage (Prokop, 2020). Accordingly, aberrations of MT bundles (presenting as gaps or areas of bundle disorganisation in the form of MT curling) are sensitive indicators of axonal pathology (Prokop, 2021). Mechanisms that help to maintain these MT bundles are starting to emerge (Hahn et al., 2019).

Our second strategic decision was the use of *Drosophila* primary neurons as a cost-effective and fast model system, where the complexity of mechanisms can be addressed with powerful genetics, in ways hard to achieve in vertebrate models (Prokop et al., 2013). For example, in this study alone, we used over 40 different mutations or transgenic constructs - some of them in genetic combinations to address functional redundancies or hierarchies (e.g. Beaven et al., 2015; Gonçalves-Pimentel et al., 2011; Koper et al., 2012). Loss-of-function analyses in *Drosophila* are facilitated by the fact that key factors, such as Kinesin heavy chain (Khc, kinesin-1), Kinesin light chain (Klc) or Milton, are each encoded by a single gene in *Drosophila*, as opposed to three, four or two in mammals, respectively. Furthermore, genetic and pharmacological tools are readily available to manipulate virtually any genes in question - and all these functional approaches can be combined with efficient and well-established readouts for axonal physiology and MTs (Hahn et al., 2016; Prokop et al., 2013; Prokop et al., 2012; Sánchez-Soriano et al., 2010).

Here we demonstrate that the losses of three motor proteins cause MT curling: the Kif5A/B homologue Kinesin heavy chain (Khc), the Kif1A homologue Unc-104, and Dynein heavy chain (Dhc/DYNC1H1). We find that all three are required for axonal transport of mitochondria and synaptic proteins. Focussing on Khc and employing available means to dissect its various sub-functions, we identified two mechanisms linking to MT curling: Firstly, loss of Khc/Milton/Miro-mediated transport causes harmful reactive oxygen species (ROS) likely linking to mitochondrial transport. Secondly, loss of the Kinesin light chain linker appears to cause de-inhibition of Khc as a condition that we find

to cause MT curling. Both mechanisms align with the recently proposed 'dependency cycle of local axon homeostasis' as a conceptual model of axonopathy (Prokop, 2021).

Results

Loss of Khc, Unc-104 or Dhc cause axonal MT curling

To assess whether loss of motor protein function impacts on axonal MT organisation, we tested mutant alleles for axonal transport-related motor proteins: (a) Dynein heavy chain (Dhc) is an obligatory component of the dynein/Dynactin complex and essential for most, if not all, MT-based retrograde transport (Reck-Peterson et al., 2018); (b) Klp64D is an obligatory subunit of heterodimeric kinesin-2 (KIF3 homologue) reported to mediate anterograde axonal transport of acetylcholine-related synaptic enzymes or olfactory receptors (Baqri et al., 2006; Jana et al., 2021; Kulkarni et al., 2017; Ray et al., 1999); (c) the PX-domain-containing type 3 kinesin Klp98A (KIF16B homologue) was shown to mediate autophagosome-lysosome dynamics and endosomal Wingless transport in non-neuronal cells but is also strongly expressed in the nervous system (Mauvezin et al., 2016; Witte et al., 2020; flybase.org: [FBgn0004387](https://flybase.org/FASTA/unc-104/FBgn0004387)); (d) the PH-domain-containing type 3 kinesin Unc-104 (Kif1A homologue) is essential for synaptic transport in *Drosophila* axons (Pack-Chung et al., 2007; Voelzmann et al., 2016); (e) Kinesin heavy chain/Khc is the sole kinesin-1 in *Drosophila* (Kif5A-C homologue) involved in multiple transport functions in *Drosophila* neurons (see details below; e.g. Bowman et al., 2000; Gindhart et al., 2003; Glater et al., 2006; Loiseau et al., 2010; Rosa-Ferreira et al., 2018; Saxton et al., 1991).

To assess potential roles of these motor proteins in MT regulation, we cultured primary neurons obtained from embryos lacking these gene functions (see Methods; Fig.1) and analysed them at 5DIV (days *in vitro*). MT curling phenotypes (where bundles deteriorate into curled, intertwined, crisscrossing MTs; curved arrows and enlarged insets in Fig.1B,F,G) occurred as a moderate phenotype in Dhc-deficient neurons and were prominent in neurons lacking Khc or Unc-104 (Fig.1H).

To assess whether MT curling phenotypes were accompanied by transport defects, we analysed additional sub-cellular markers in *Khc*⁸, *unc-104*¹⁷⁰ and *Dhc64C*⁴⁻¹⁹ homozygous mutant neurons. First, using the pre-synaptic protein Synaptotagmin (Syt) as an indicator of vesicular transport (Voelzmann et al., 2016), we found reduced presynaptic spots within axons of neurons mutant for any one of these three motor proteins, suggesting they all contribute to axonal vesicular transport (Fig.2A-D,O). Second, the axonal number and distribution of mitochondria (visualised with mitoTracker; Klionsky et al., 2012), is significantly reduced in neurons lacking either Khc, Unc-104 or Dhc64C function (Fig.2H-K,P). Third, in *Khc*⁸ mutant neurons, we also assessed the distribution of endoplasmic reticulum (ER) using the genomically tagged *Rtnl1-GFP* allele. In wild-type neurons, ER is distributed evenly along the entirety of the axon (Fig.S1A); loss of Khc does not affect this distribution, but about three quarters of neurons show an abnormal accumulation of Rtnl1::GFP-labelled ER material at their tips (Fig.S1B,C; details in legend).

In conclusion, the three motor proteins that display MT curling are also required for normal axonal transport of synaptic vesicles and mitochondria in *Drosophila* primary neurons. In addition, at least Khc is also required for the normal axonal distribution of ER.

Khc displays strong maternal effects

Of these three motor proteins, we performed more detailed analyses on Khc because many genetic tools are available for the systematic dissection of its various functions (Fig.3B). First, to validate its MT-related phenotypes, we tested additional mutant alleles (*Khc*²⁷ and *Khc*⁸ in homozygosis or over deficiency) as well as RNAi mediated knockdown of Khc (via pan-neuronal *elav-Gal4*). In all cases we found that the MT curling phenotypes were equally present at 5DIV (Fig.S2A,B,D).

However, the MT phenotypes were not evident at earlier stages in *Khc*^{8/Df} mutant neurons, either at 6 hours *in vitro* (HIV) or at 3DIV (Fig.S2B,D), suggesting that phenotypes either accumulate gradually (as observed upon loss of Efa6; Qu et al., 2019) or are masked by perdurance of maternal product (meaning wild-type *Khc* gene product deposited in the egg by the heterozygous mothers; Prokop, 2013).

To distinguish between these two possibilities, we used a pre-culture technique where neurons are kept in centrifuge tubes for 5 days to deplete maternal gene product before plated in culture (Prokop et al., 2012; Sánchez-Soriano et al., 2010). Such pre-cultured neurons displayed prominent MT curling already at 1DIV (Fig.S2C,E), arguing that Khc has a prominent maternal contribution that persists for more than 3 days. Similar observations were made with the *Khc*^{1ts} mutant allele (details in Fig.S2C).

MT sliding functions of Khc do not link to MT curling

A C-terminal MT-binding site enables Khc to cross-link MTs and move them against each other (Fig.3Bi; Andrews et al., 1993; Jolly et al., 2010; Lu et al., 2013; Lu et al., 2015; Winding et al., 2016). We hypothesised that Khc might contribute to MT bundle maintenance by using its MT sliding function, for example by shifting MTs to achieve even distribution along axons. The Khc sliding function is selectively inhibited by the genomically engineered, lethal *Khc*^{mutA} allele that abolishes C-terminal MT binding without interfering with other linkers or autoinhibition of Khc (Fig.3A; Winding et al., 2016).

To test whether Khc-mediated sliding contributes to MT bundle regulation, we cultured *Khc*^{mutA} mutant neurons in different ways: embryo-derived neurons were cultured for 1DIV after 5d pre-culture or for 5DIV without pre-culture, and neurons from larval brains were cultured for 2DIV. In all cases, these neurons failed to show enhanced MT curling (Fig.3Ci), suggesting that the *Khc* mutant phenotype is not caused by the loss of its MT sliding function.

Loss of Milton and Miro causes MT curling phenotypes

Next, we focussed on the transport functions of Khc. For example, Pat1 (Protein interacting with APP tail-1) had been shown to link Khc to non-vesicular transport in *Drosophila* oocytes (Fig.3Bii; Loiseau et al., 2010). It is also strongly expressed in the *Drosophila* nervous system (flybase.org: FBgn0029878) but potential neuronal cargoes are unknown. Pat1 function can be eliminated by the gene-specific small deficiencies *Pat1*^{robin} and *Pat1*^{grive} which represent viable null alleles (Loiseau et al., 2010). When analysing cultures of larval neurons homozygous for either allele, we did not find any obvious enhancement of MT curling (Fig.3Cii).

The linker protein Milton and its binding partner Miro (a small GTPase) link the C-terminus of Khc to organelles including mitochondria (Fig.3A,Biii; Harbauer, 2017; Misgeld and Schwarz, 2017; Sheng,

2017; Smith and Gallo, 2018) and potentially peroxisomes (Castro et al., 2018; Covill-Cooke et al., 2017; Okumoto et al., 2018; Tang, 2018).

Using the loss-of-function mutant alleles *milt*⁹², *Miro*^{Sd32} or *Miro*^{B682}, we first confirmed the functional contributions of Milt and Miro in primary fly neurons. For this, we stained homozygous mutant neurons with mitoTracker and anti-Syt. We found the axonal localisation of Syt to be unaffected whereas mitochondria were strongly reduced in number, thus confirming the expected cargo specificity (Fig.2E-G,L-P). The mitochondrial phenotype was milder for loss of Miro than Milt, as is consistent also with previous findings in fly neurons *in vivo* as well as mouse neurons (Glater et al., 2006; Guo et al., 2005; López-Doménech et al., 2018; Russo et al., 2009; Vagnoni et al., 2016). In *milt*⁹² mutant neurons, mitochondria were virtually absent from axons, restricted mostly to cell bodies and proximal axon segments (Fig.2L). This absence of mitochondria as the major ATP source does not affect synaptic transport because it is self-sufficient through local glycolysis on transported vesicles (Fig.S3A,B; Hinckelmann et al., 2016; Zala et al., 2013).

Having confirmed that the *milt*⁹², *Miro*^{Sd32} or *Miro*^{B682} alleles selectively inhibit mitochondrial transport, we then assessed potential impacts on MT organisation. We found that all three mutant alleles caused significant increases in MT curling (Fig.3Ciii). This finding suggests that loss of Khc-mediated organelle transport triggers MT curling (Fig.3Biii).

Excessive ROS triggers MT disorganisation

Reduced numbers of axonal mitochondria can be expected to impair local homeostasis of calcium, ATP, reactive oxygen species (ROS), and AAA+ protease-mediated protein quality control systems (Glynn, 2017; Misgeld and Schwarz, 2017; Paupe and Prudent, 2018). We started by manipulating the ROS homeostasis through the application of DEM (diethyl maleate). DEM is an effective inhibitor of the anti-oxidant compound glutathione, hence causing the elevation of ROS levels (Fig.4A; Albano et al., 2015; Dasgupta et al., 2012; Pompella et al., 2003). We found that 12 hr-long application of 100 µM DEM before fixation (from 4.5 to 5DIV) induced robust MT curling (Fig.4F). To validate this finding, we then used genetic tools to generate loss- or gain-of-function conditions for ROS-regulating enzymes (yellow highlighted in Fig.4A):

Firstly, we used a null allele of Catalase (*Cat*^{l⁹¹}), an enzyme removing hydrogen-peroxide (H₂O₂; Walker et al., 2018), and null alleles of two members of the Superoxide dismutase family, *Sod1* (*Sod1*ⁿ¹ or *Sod1*ⁿ⁶⁴) and *Sod2* (*Sod1*ⁿ²⁸³), which convert superoxide anions (O₂⁻) into H₂O₂ (Palma et al., 2020). Of these, Catalase is enriched in peroxisomes, copper-zink-dependent *Sod1* (linked to amyotrophic lateral sclerosis/ALS1; #105400; Saccon et al., 2013) is primarily cytoplasmic, and the manganese-dependent *Sod2* enzyme is predominantly mitochondrial (Fig.4A). When assessed in primary *Drosophila* neurons, the functional deficiencies of either *Sod1*, *Sod2* or *Cat* caused MT curling (Fig.4F).

Secondly, we used targeted expression (a) of *Nox* (NADPH oxidase) to enhance O₂⁻ levels, (b) of *Sod1* to reduce O₂⁻ levels and enhance H₂O₂, and (c) of *Duox* (Dual oxidase) to increase H₂O₂ levels (Anh et al., 2011; Bedard and Krause, 2007; Zelko et al., 2002). All these manipulations caused increased MT curling (Fig.4F).

Taken together these results clearly indicate that insults to ROS homeostasis have a strong tendency to trigger MT curling. Upregulation of either O₂⁻ or H₂O₂ seems to cause this effect, although O₂⁻ may elicit its effects through conversion into H₂O₂ (Bedard and Krause, 2007).

Harmful ROS appears to relate to mitochondria and links loss of Khc or Milt to MT curling

To assess whether harmful ROS might be responsible for linking loss of Khc/Milton/Miro to MT curling, we treated mutant neurons with Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid), an α -tocopherol/vitamin E analogue that displays beneficial antioxidant effects in many cell systems by inhibiting fatty acid peroxidation and quenching singlet oxygen and superoxide (Giordano et al., 2020). Neurons mutant for *Khc*⁸ or *milt*⁹² were either pre-cultured for a day in the presence of 100 μ M Trolox, or they were cultured directly for 5 days with Trolox. Under both conditions, application of Trolox strongly suppressed or even abolished the MT curling phenotype of *Khc*⁸ and *milt*⁹² mutant neurons (Fig.5); this indicated harmful ROS to be the main reason for MT bundle disintegration. It might explain why rat neurons depleted of the Khc homologue KIF5C were reported to be more vulnerable to H₂O₂ application (Iworima et al., 2016).

To understand how loss of Khc and Milt might trigger harmful levels of ROS, we first set out to identify the potential source. For example, loss of Catalase causes MT curling (Fig.4A,F), potentially suggesting that peroxisomes are required to keep H₂O₂ levels down. To test this possibility, we blocked peroxisome biogenesis using the *Pex3*² mutation (Faust et al., 2014). Abolishing peroxisomes in this way did not induce any obvious MT curling phenotypes, but the *Pex3*² mutant neurons had shorter axons (Fig.S4) potentially due to lack of peroxisomal lipidogenesis (Wanders et al., 2020). The absence of obvious MT phenotypes seems to contradict MT curling observed upon Catalase deficiency (Fig.4F), but it might be explained by observations that Catalase can localise outside peroxisomes in the cytoplasm (Zhou and Kang, 2000).

We concluded that MT curling observed upon loss of Khc, Milt or Miro is more likely to link to their roles in axonal transport of mitochondria; disturbing mitochondrial dynamics could either generate harmful ROS (via the ETC; Fig.4A) or affect their ability to quench local ROS (via Sod2; Fig.4A).

ROS-absorbing properties rather than disrupted fission/fusion of mitochondria might provide links to MT curling

Loss of Khc/Milt/Miro might cause harmful ROS by affecting fission/fusion processes required to maintain a healthy mitochondrial population (Cagalinec et al., 2013; Liu et al., 2009; Wang et al., 2015). In support of this notion fission/fusion factors are linked to axonopathies; this is the case for the fission factor DNM1L/DYNAMIN-LIKE PROTEIN 1 (Optic atrophy 5; OMIM® #610708), as well as the fusion factors OPA1/OPA1 MITOCHONDRIAL DYNAMIN-LIKE GTPase (Optic atrophy 1; #165500) and MFN/MITOFUSIN (CMT2A2A, CMT2A2B, HMSN6A; #609260, 617087, 601152).

To test whether loss of fission/fusion is a condition that affects MT bundling, we used mutant alleles abolishing the functions of the fly homologues of mammalian DNM1L (*Drp1*^{T26}; Dynamin related protein 1), of mammalian OPA1 (*Opa1*^{s3475}, Optic atrophy 1) and of mammalian MFN (*Marf*^B; Mitochondrial assembly regulatory factor). In axons of wild-type neurons, mitochondria mostly displayed dash-like shapes (Fig.S5A) and occasionally appeared dot-like or formed longer lines (not shown). In contrast, within axons of neurons with impaired fusion (*Opa1*^{s3475} or *Marf*^B) mitochondrial shapes were primarily short and dot-like (Fig.S5C,D), whereas loss of fission (*Drp1*^{T26}) caused string-of-pearl arrangements where a continuous thread of mitochondria ran all along the main axon but was excluded from side branches (Fig.S5B). These findings are consistent with reports for mammalian neurons (Smirnova et al., 2001; Uo et al., 2009; Yu et al., 2011).

When analysed for MT organisation, none of the three fission/fusion-deficient conditions caused curling in axons, neither at 5 DIV nor upon pre-culture (Fig.S5E,F). This might suggest that MT curling upon loss of Khc/Milt/Miro is unlikely to be caused by mitochondrial fission/fusion defects; the

absence of fission/fusion events seems not to affect mitochondria in ways that cause harmful ROS leakage, as is also consistent with views of other authors (Misgeld and Schwarz, 2017). Accordingly, also MFN2-deficient mouse neurons seem not to experience oxidative stress (Baloh et al., 2007). A further argument against the involvement of fission/fusion is based on the observation that mitochondria in Milt-deficient neurons tend to stay in the cell body: it is unlikely that harmful ROS generated in the soma were to reach the distal axon via long-range diffusion, especially when considering the abundance of ROS-buffering systems (Fundu et al., 2019; Kükürt et al., 2021; Oswald et al., 2018).

We prefer therefore the explanation that MT curling in Khc/Milt/Miro-deficient neurons (Fig.2P) might be caused by the absence of mitochondria from critical positions in the axon, thus depleting these areas from Sod2 activity (for details see Discussion).

Klc also causes MT curling but through a mechanism distinct from Milt

Given the comparable strength of MT phenotypes upon loss of Khc, Milt or Miro (Fig.3Ciii) and their shared links to harmful ROS production (Fig.5), mitochondrial transport defects seemed to offer the perfect explanation for why loss of Khc induces MT curling. We expected therefore that loss of the vesicular transport linker Klc would not cause these MT phenotypes.

Surprisingly, we found that also loss of Klc (*Klc*^{8ex94} or *Klc*^{8ex94/Df} mutant neurons at 5 DIV) caused MT curling, and that this phenotype was at least as strong as observed with Khc or Milt deficiency (Fig.3Ciii,iv). We obtained the same results when performing 5d pre-cultures with *Khc*⁸, *Klc*^{8ex94} and *milt*⁹² mutant neurons (to deplete their maternal products; Figs.6A-E), confirming that the three factors generate comparably strong MT curling phenotypes.

To establish whether Klc might work synergistically with Khc and Milt, we applied Trolox to *Klc*^{8ex94} mutant neurons. However, in contrast to *Khc*⁸ and *milt*⁹² mutant neurons, the MT curling in Klc-deficient neurons was not suppressed by Trolox, neither in pre-cultured neurons nor in 5DIV cultures (Fig.5). This clearly demonstrated that Klc works through an independent mechanism.

Klc's links to MT curling do not depend on vesicular cargo transport

We first tested whether Klc's impact on MT regulation might link to vesicular cargo transport, capitalising on reports that Khc-mediated vesicular cargo transport requires a protein complex of a number of factors including Klc and Sunday driver (Syd, the JIP3 homologue; Fig.3Biv; Gindhart et al., 2003; Horiuchi et al., 2005; Koushika, 2008). Accordingly, functional loss of either Klc or Syd was shown to abolish Khc-mediated vesicular transport, with motoraxons in peripheral larval nerves displaying synaptic protein accumulation that were similarly strong upon Klc or Syd deficiency as observed upon loss of Khc (Bowman et al., 2000; Fügler et al., 2012; Gauger and Goldstein, 1993; Gindhart et al., 1998; Hurd and Saxton, 1996; Pilling et al., 2006). Equally in primary culture, we found that the number of Synaptotagmin-positive dots in axons was reduced in *Klc*^{8ex94} and *syd*^{z4} null mutant neurons, and the phenotypes were similarly strong as observed in Khc-deficient neurons (Fig.2B,G,O). These effects seem specific to vesicular cargo transport since mitochondrial numbers in axons of *Klc*^{8ex94} mutant neurons appeared normal (Fig.2N,P).

Our data confirm therefore that Klc, Syd and Khc closely co-operate during vesicular cargo transport in primary *Drosophila* neurons. However, in contrast to severe MT curling in Klc- and Khc-deficient neurons, *syd*^{z4} or *syd*^{z4/Df} mutant neurons at 5DIV failed to show similar phenotypes (Fig.3Civ). This

observation was further supported by experiments where we knocked down GAPDH (glyceraldehyde-3-phosphate dehydrogenase), an enzyme required for glycolysis that is known to fuel the axonal transport of vesicles but not of mitochondria (details in Fig.S3A,B; Zala et al., 2013). When GAPDH was knocked down in primary *Drosophila* neurons, axons displayed a reduction in synaptic dots but not in mitochondrial numbers (Fig.S3C), as is consistent with analyses in larval nerves (Zala et al., 2013). However, GAPDH knock-down did not cause obvious MT curling phenotypes (Fig.S3C), thus mirroring results obtained with Syd deficiency.

We concluded that blocking vesicular axonal transport appears not to be a cause for MT curling, and that loss of Klc is likely to trigger its MT phenotypes through a different mechanism. This view was also supported by genetic interaction studies using trans-heterozygous pairings of *Khc*⁸, *milt*⁹² and *Klc*^{8ex94} (i.e. combining heterozygosity for two genes at a time in the same neurons). Of the three constellations, only *Khc*^{8/+} *milt*^{92/+} trans-heterozygote mutant neurons generated a MT curling phenotype that was significantly enhanced over single heterozygous conditions (Fig.6G), supporting functional links between Khc and Milt but not with Klc.

Taken together, MT curling upon loss of Khc appears to relate to Milt/Miro-mediated mitochondrial transport as explained before, but not to Klc-mediated vesicular transport. Milt, Miro and Khc seem to have comparably strong mutant phenotypes because their loss leads to the same transport defect, whereas phenotypes observed upon loss of Klc (which has binding sites on Khc that overlap with those of Milt; details in Fig.3A) seem not to relate to its function as a transport linker but work through an entirely different mechanism.

Excessive pools of active Khc might explain the Klc-deficient MT phenotype

We hypothesised that MT curling upon loss of Klc may relate to its roles in regulating the activation state of Khc. Thus, Khc pools that are not linked to cargo tend to be auto-inhibited and detached from MTs. This inactivation requires intramolecular loop formation via binding of the N- to the C-terminus, and this also involves the association with Klc (co-regulated through its own auto-inhibition/activation mechanism; Figs.3A,Biv; Bowman et al., 2000; Koushika, 2008; Verhey and Hammond, 2009; Verhey et al., 1998; Wong and Rice, 2010; Yip et al., 2016). In non-neuronal cells, overriding auto-inhibition of the Khc-Klc complex causes MT curling (Paul et al., 2020; Randall et al., 2017).

To test whether Klc-deficient MT curling in neurons might involve excessive pools of active Khc, we first targeted the expression of GFP-tagged constructs of Khc to neurons. We found that full-length Khc::GFP was homogeneously distributed along axons and failed to increase MT curling when analysed at 5DIV (Fig.7), consistent with the idea that extra pools of Khc tend to be inactive and detached from MTs.

We then expressed two non-inactivating Khc derivatives (*Khc*¹⁻⁸¹¹::GFP and *Khc*⁸²⁻⁷¹¹::GFP; top of Fig.3A) which both lack the C-terminal domain needed for auto-inhibition but also for their roles in cargo transport and MT sliding (dark grey and green in Fig.3A). When analysed in neurons at 5DIV, both constructs accumulated at axon tips, as is typical of non-inactivating kinesins (Niwa et al., 2013). Of these, *Khc*¹⁻⁸⁵⁰::GFP caused a very mild MT phenotype, suggesting that extra pools of free-running Khc *per se* cause little harm (Fig.7B,D). In contrast, *Khc*⁸²⁻⁷¹¹::GFP caused severe MT curling (Fig.7C,D), potentially because this truncated form also has a small N-terminal deletion – and short deletions of the N-terminus have been shown to display damaging effects on MTs (Budaitis et al., 2021; details in Fig.3A). In principle, our findings with this dys-regulated construct supported our

previously published hypothesis that the activity of kinesins is harmful to axonal MT bundles and can explain MT curling (Hahn et al., 2019; Prokop, 2021).

However, since all essential C-terminal binding sites were removed in $Khc^{1-850::GFP}$ and $Khc^{82-711::GFP}$ (Fig.3A), our experiments so far only assessed free-running Khc that could not engage in movement of any cargo (Fig.3A,B). We hypothesised that active transport would be expected to generate higher forces than free-running Khc and, hence, be more challenging to MT bundles. We therefore tested the genomically engineered Khc^{E177K} and $Khc^{E177R, R947E}$ mutant alleles (Fig.3A), which have point mutations in the E177 and R947 residues that are known to form a required salt bridge with each other during auto-inactivation (Kaan et al., 2011; Kelliher et al., 2018); these mutant alleles cause lethality and distal accumulations of Khc (Brendza et al., 1999; Kelliher et al., 2018). When analysing Khc^{E177K} and $Khc^{E177R, R947E}$ homozygous mutant neurons at 3DIV, we found that they display robust MT curling (Fig.7D). This clearly indicated that extra pools of actively engaging Khc harm MT bundles, which might therefore explain the *Klc* mutant phenotype (see Discussion).

Discussion

Using MT bundles of *Drosophila* neurons as a powerful approach to dissect motor-related pathomechanisms

Motor proteins involved in axonal transport clearly are key drivers of neuronal survival, yet their links to axonopathies remain poorly understood and speculative (Coleman, 2005; Guo et al., 2020; Kawaguchi, 2013; Sleight et al., 2019). Here, we aimed to unravel concrete mechanisms through which motor protein loss can affect axons.

Our approach was unprecedented in that we performed a systematic genetic study in one standardised neuron system and used MT bundles as key readout. We studied the organisation of axonal MT bundles because they are good indicators of axon integrity (Prokop, 2020) that are easy to quantify and have an intricate interdependent relationship with motor proteins (Prokop et al., 2013). The key phenotype we observed upon motor manipulation is MT curling which appears conserved across species, since depleting Dynein or the kinesin-1 linker JIP3 causes the same kind of curling in axons of mammalian neurons (Ahmad et al., 2006; Rafiq et al., 2020).

The easily accessible and quantifiable MT curling readout allowed us to determine roles of motor proteins and their interactors, or of proteins regulating potential downstream processes - and many of these factors have known links to neurodegeneration or axonopathies. Our approach was facilitated by using a standardised neuronal culture system in which findings could be integrated and were highly accessible to powerful *Drosophila* genetics. The additional advantage of this system is low genetic redundancy of factors involved in axonal transport, with one *Drosophila* gene having on average almost 3 mammalian orthologues (*Khc/Kif5*: 1 paralogue in fly vs. 3 paralogues in mammals; *Miro/RHOT*: 1 vs. 2; *Milt/TRAK*: 1 vs. 2; *Klc*: 1 vs. 4; *Marf/MFN*: 1 vs. 2; *Unc-104/Kif1*: 1 vs. 3). This low redundancy in fly enormously facilitates loss-of-function analyses and combinatorial genetics.

Capitalising on these advantages, our unprecedented strategy enabled us to generate new understanding and conceptual explanations. So far, we found that deficiencies of three motor proteins (*Khc*, *Unc-104*, *Dhc*) cause MT curling. Notably, the homologues of all three factors have OMIM®-listed links to human axonopathies (see Introduction) potentially reflecting evolutionarily conserved mechanisms of axon pathology that might even be shared between these motor protein classes.

For example, we found that phenotypes upon loss of Khc and Unc-104 are very similar with respect to enhanced MT curling and the reduction in axonal numbers of mitochondria and synaptic dots (Figs.1, 2), and both seem involved in mRNA transport (L.M.P.C., unpublished results; Lyons et al., 2009). This functional overlap is in agreement with reports that kinesin-1 and -3 collaborate during transport (Arpağ et al., 2019; Zahavi et al., 2021). Consequently, these two motors might therefore link to axonopathy through comparable mechanisms.

An intricate relationship: kinesins simultaneously harm and care for MT bundles

The challenges of studying axonal transport are ample due to (1) the parallel involvement of different motor protein classes (which might act redundantly; Hirokawa et al., 2010; see previous section), (2) the enormous wealth of their cargoes, (3) the involvement of many different linkers (that might interact promiscuously with different motors; Brady and Morfini, 2017; Drerup et al., 2016; Gindhart, 2006; Hirokawa et al., 2010; Maday et al., 2014), (4) additional roles in slow transport (transient 'hitchhiking' of proteins on transported vesicles; Roy, 2020; Tang et al., 2013), and (5) complications caused by the interdependence of kinesins and dynein/Dynactin (Hancock, 2014; Moughamian et al., 2013; Twelvetrees et al., 2016; potentially explaining the rather counter-intuitive observation that loss of anterograde Khc transport causes distal ER accumulations; details in legend of Fig.S1).

For kinesin-1 alone (Fig.3B), we tested roles of Khc in MT sliding (*Khc^{mutA}*), roles of Khc/Milt/Miro in mitochondrial/peroxisomal transport, of Khc/Klc/Syd/GAPDH in vesicular transport, of Khc/Pat1 in potential non-vesicular transport, and potential roles of Klc and certain Khc domains/residues in Khc auto-inhibition. These extensive studies still left out further known linkers, such as SKIP/SNW1/SKIIP and Arl8 (lysosome transport; Keren-Kaplan and Bonifacino, 2021; Rosa-Ferreira and Munro, 2011; Rosa-Ferreira et al., 2018) or Tropomyosin (mRNA transport; Fig.3A; Dimitrova-Paternoga et al., 2021; Veeranan-Karmegam et al., 2016). Nevertheless, the analyses we performed suggested two distinct mechanisms:

Firstly, Khc/Milt/Miro-mediated transport is required to uphold ROS homeostasis, with harmful ROS being a strong inducer of MT curling (demonstrated by our studies with DEM, Trolox and ROS-regulating enzymes; Figs.4, 5). Secondly, we found that the movement and active transport of Khc along MTs damages axonal bundles, as demonstrated by the expression of Khc deletion constructs and analyses of non-inactivating *Khc* mutant alleles (Fig.7). These latter findings align with published *in vitro* experiments demonstrating kinesin-1-induced MT damage (Andreu-Carbó et al., 2021; Budaitis et al., 2021; Dumont et al., 2015; Triclin et al., 2021; VanDelinder et al., 2016), MT curling observed in kinesin-1-based gliding assays *in vitro* (Hahn et al., 2019; Lam et al., 2016), and the curling observed upon kinesin-1 activation in non-neuronal cells (Paul et al., 2020; Randall et al., 2017). Notably, MT curling is not specifically linked to motor proteins, but is similarly observed upon loss of various MT-binding and -regulating proteins (Hahn et al., 2019) and in a model of chemotherapy-induced peripheral neuropathy (Rozario et al., 2021).

All these causes of MT curling, including the two mechanisms described in this work, can be explained with the previously proposed "local axon homeostasis" model (Hahn et al., 2019) and the subsequently derived "dependency cycle of axon homeostasis" (Prokop, 2021; details in Fig.8). These models propose that kinesins that trail along MT bundles during axonal transport ('2' in Fig.8) pose a mechanical challenge that leads to MT curling ('3'); active machinery of MT-regulating proteins and support through the cortical actin-spectrin sleeve is therefore required to prevent disintegration and maintain these bundles long-term ('4'). However, the machinery that maintains MT bundles is itself dependent on materials and physiology provided by axonal transport ('5'), thus

establishing a cycle of mutual dependency where interruption at any point will have a knock-on effect on all other aspects of axon physiology and function (Prokop, 2021). The mechanisms we described here act in either direction of this cycle (thick read arrows in Fig.8).

Mitochondria regulate ROS homeostasis required for MT bundle maintenance

Our finding that harmful ROS is a key trigger of MT curling aligns with reports that actin as well as MTs are modified or even damaged by ROS (Goldblum et al., 2021; Wilson et al., 2016; Wioland et al., 2021) and that oxidative stress induces axon swellings in models of Parkinson's disease, multiple sclerosis or ALS (Czaniecki et al., 2019; Nikić et al., 2011; Song et al., 2013). Unfortunately, a more generalised statement cannot be made because axonal MTs have rarely been analysed in oxidative stress experiments (De Vos et al., 2007; Debattisti et al., 2017; Fischer et al., 2012; Saccon et al., 2013; Song et al., 2013).

Pinpointing the precise source of harmful ROS upon Khc/Milt/Miro loss is a tedious task when considering (1) the intricate network of ROS regulation (Fig.4A) where manipulations of very different regulators caused comparable phenotypes (Fig.4F), and (2) the spectrum of organelles involved in ROS homeostasis regulation: these involve the finely tuned mitochondria-peroxisome system (Fransen et al., 2017; Pascual-Ahuir et al., 2017), but also the ER which contains oxidases required for protein folding (Hudson et al., 2015). Unfortunately, removing or affecting the ER to assess its involvement is not trivial (O'Sullivan et al., 2012; Yalcin et al., 2017); but our studies of *Pex3* mutant conditions suggested that peroxisomes are unlikely to link to MT curling (Fig.S4). This said, peroxisomes certainly play important further roles in maintaining healthy axons (Wali et al., 2016).

In our view, the most likely organelles involved in MT curling are the mitochondria. We were surprised to find that not the presence of damaged mitochondria leaking harmful ROS seems to trigger MT curling (Fig.S5), but rather the absence of mitochondria. This is best illustrated by *milt* mutant neurons where mitochondria are mostly restricted to cell bodies (Fig.2L,P), yet strong ROS-induced MT curling occurs in axons (Figs.3Ciii, 5, 6E).

As already mentioned in the Results part, we believe that the best model combining all observations is the absence of mitochondria and Sod2 as their ROS scavenger from critical locations in axons. For example, we know from live imaging experiments that MT curling starts at growth cones or branch points (A.V., unpublished data), and both are typical sites where mitochondria localise (Bunge, 1973; Mandal and Drerup, 2019). Failure to quench harmful ROS in these critical locations could therefore promote the initiation of MT curling; this would also explain why drastic mitochondrial depletion upon Milt deficiency triggers similarly strong MT curling as moderate depletion upon loss of Khc or Miro (Figs.2P, 3C): not the number of mitochondria is essential but their adequate localisation, and this aspect is regulated through a Khc/Milt/Miro-dependent mechanism (Misgeld and Schwarz, 2017).

The drastic depletion of axonal mitochondria upon Milt deficiency as compared to the moderate number reductions upon loss of Khc, Unc-104 or Dhc (Fig.2P) might suggest Milt as a 'master linker' for mitochondrial transport in fly neurons. Indeed, Milt is known to link to Khc and Dynein in both flies and mammals (Russo et al., 2009; van Spronsen et al., 2013), whereas there are currently no such reports for Unc-104; its mammalian homologue Kif1 was reported so far to perform mitochondrial transport through KBP (Kif1 binding protein; Campbell et al., 2014; Nangaku et al., 1994; Tanaka et al., 2011; Wozniak et al., 2005).

Khc activation as a further factor leading to MT curling

As discussed above, our data with non-inactivating constructs and mutant alleles of Khc strongly suggested that excess engagement of this motor can trigger MT curling (Fig.7). Since Klc is involved in Khc auto-inhibition, the MT curling phenotype we observe in axons of Klc-deficient neurons might therefore link to this mechanism as a potential cause for axonopathy.

Also Kif1A/Unc-104 undergoes inactivation involving intramolecular loop formation and the KBP linker (Cong et al., 2021; Kevenaar et al., 2016). KBP mutations impair axon growth, are disruptive to axonal MT bundles (Lyons et al., 2008) and cause the devastating neurological disorder Goldberg-Shprintzen syndrome in humans (Chang et al., 2019; Hirst et al., 2017). Similarly, non-inactivating mutations of Kif1A cause spastic paraplegia (Chiba et al., 2019; Gabrych et al., 2019). Also KLC2 has been linked to neuropathy (SPOAN; #609541) and KLC4 mutations have recently been reported to cause excessive axon branching (Haynes et al., 2021). Axonopathy-linked human mutations of Kif5A were mapped exclusively to the motor domain or the very C-terminal end so far, but none were reported in the auto-inactivation domains or the KLC binding site (Nicolas et al., 2018). However, this does not mean that such mutations are not detrimental: mutations affecting auto-inhibition might rather confer lethality (as is the case in *Drosophila*; Brendza et al., 1999; Kelliher et al., 2018) and therefore escape the spectrum of diagnosed diseases.

The modest curling observed upon overexpression of Khc^{1-850::GFP}, versus the strong phenotype with genomically engineered *Khc*^{E177K} and *Khc*^{E177R, R947E} mutant alleles (Fig.3A), suggests that free-running Khc is insufficient to cause a strong phenotype; instead it seems that the C-terminus of Khc has to interact with cargo to generate forces strong enough to affect MT bundles. But what cargoes might be involved?

Some insights might come from *milt*^{92/92} *Klc*^{8ex94/8ex94} double-mutant neurons which show an intriguing pattern: loss of Klc and Milt trigger MT curling through completely different mechanisms (Fig.8), and their phenotypes should therefore be additive. However, double-mutant neurons show the same amount of curling as the single mutants (Figs.5B, 6F). Even more, MT curling in double-mutant neurons is partly cured by Trolox, although Klc-deficiency alone does not respond to Trolox (Fig.5B). The easiest explanation for these findings is that surplus pools of activated Khc triggered by loss of Klc engage in force-generation that depends on mitochondria-derived ATP (unlike vesicular transport; Fig.S3A). Since mitochondria and their ATP are absent from axons of *milt* *Klc* double-mutant neurons, the extra pool of Khc lacks the necessary fuel to contribute to the joint MT curling phenotype. So far, our attempts to pinpoint such force-generating activities of surplus Khc pools have not been successful: they seem not to involve microtubule sliding and Pat1-mediated transport (Fig.3Bi,ii) as suggested by failed suppression of MT curling in *Khc*^{mutA/mutA} *Klc*^{8ex94/8ex94} or *Pat1*^{robin/robin} *Klc*^{8ex94/8ex94} double-mutant neurons (details in Fig.S6).

Conclusion

Using our unconventional strategy (MT curling as key readout for systematic genetic analyses in a standardised *Drosophila* primary neuron system) we were able to develop new concepts for how molecular motor mutations might trigger axonopathies. Given the breadth of genetics versus lack of mechanistic detail of our studies, our results are certainly more suggestive than definite. But they are astonishingly consistent with many reports in the field (as mentioned throughout this work) and align well with the 'dependency cycle of local axon homeostasis' as a model describing the fundamental principle of axon maintenance and pathology (Prokop, 2021). We hope therefore that our findings and ideas will stimulate further studies in which the model and proposed mechanisms

are put to the test, incorporating also other aspects of axon physiology, such as ATP and calcium regulation. Whatever the outcome, such studies will be highly informative and contribute to the battle against a class of diseases that are of enormous socioeconomic burden and personal hardship.

Conflict of Interest

None of the authors has a conflict of interests

Acknowledgements

This work was made possible through support by the Biotechnology and Biological Sciences Research Council (BBSRC) to A.P. (BB/I002448/1, BB/P020151/1, BB/L000717/1, BB/M007553/1) and M.L. (BB/R016666/1), by parents to Y.-T.L., H.T. and T.M. The Fly Facility has been supported by funds from The University of Manchester and the Wellcome Trust (087742/Z/08/Z). We thank colleagues and the Bloomington *Drosophila* Stock Center (NIH P40OD018537) for providing fly stocks as mentioned in Materials and Methods.

Methods

Fly stocks

All human homology statements are based on information listed on flybase.org (Marygold et al., 2016; Millburn et al., 2016), all statements about genetic links to human diseases on information provided by www.omim.org (Online Mendelian Inheritance in Man®; Amberger et al., 2015). The following stocks were used in this study (reference and source provided in brackets; BL indicates Bloomington *Drosophila* Stock Collection): null mutant alleles (unless indicated differently) we used were

- *unc-104*¹⁷⁰ (Pack-Chung et al., 2007; Tom Schwarz)
- *Klp64D*^{K1} (Ray et al., 1999; hypomorphic allele; BL #5578)
- *Klp64D*ⁿ¹²³ (Perez and Steller, 1996; BL #5674)
- *Klp98A*^{Δ47} (Derivery et al., 2015; Marcos Gonzalez-Gaitan)
- *Dhc64C*⁴⁻¹⁹ (Gepner et al., 1996; BL #5274)
- *Khc*⁸ (Saxton et al., 1991; BL #1607)
- *Khc*²⁷ (Saxton et al., 1991; Isabel Palacios)
- *Klc*^{1ts} (Saxton et al., 1991; BL #31994; a temperature-sensitive allele which is homozygous viable at 18°C but usually kept over balancer)
- *Khc*^{mutA} (Winding et al., 2016; Vladimir Gelfand; confirmed by lethality of hetero-allelic *Khc*^{mutA/8} animals)
- *Khc*^{E177K} and *Khc*^{E177K,R947E} (Kelliher et al., 2018; Jill Wildonger)
- *Df(Khc)* (*Df(2R)BSC309*; Cook et al., 2012; BL #23692)
- *milt*⁹² (Cox and Spradling, 2006; Stowers et al., 2002; Tom Schwarz)
- *Df(milt)* (*Df(2L)ED440*, *P{w[+mW.Scer\FRT.hs3]=3'.RS5+3.3}ED440*; Ryder et al., 2004; Kyoto #150498)
- *Miro*^{Sd32} (Guo et al., 2005)
- *Miro*^{B682} (Guo et al., 2005; BL #52003)
- *Df(Miro)* (*Df(3R)Exel6197*; Parks et al., 2004; BL #7676)
- *Klc*^{8ex94} (Gindhart et al., 1998; BL #31997)

- *syd⁴* (Bowman et al., 2000; BL #32016)
- *Df(3L)syd^{A2}* (Bowman et al., 2000; deleting C-terminus; BL #32017)
- *Pat1^{grive}* and *Pat1^{robin}* (Loiseau et al., 2010; Isabel Palacios)
- *Sod1ⁿ¹* (Phillips et al., 1989; BL #24492)
- *Sod1ⁿ⁶⁴* (Phillips et al., 1995; BL #7451)
- *Rtnl1-YFP (PBac{681.P.FSVS-1}Rtnl1CPT1001291*; Cahir O'Kane; O'Sullivan et al., 2012)
- *Drp1^{T26}* (Verstreken et al., 2005; BL #3662)
- *Mar^B* (Sandoval et al., 2014; Hugo Bellen)
- *P{lacW}Opa1^{s3475}* (Spradling et al., 1999; BL #12188)
- *Catⁿ¹* (Mackay and Bewley, 1989; Matthias Landgraf)
- *Sod2ⁿ²⁸³* (Duttaroy et al., 2003; BL#34060)
- *Pex3²* (Faust et al., 2014; BL#64251)

Gal4 driver lines used were the

- *elav-Gal4* (Luo et al., 1994)
- *tubP-Gal4* (Lee and Luo, 1999; Liquan Luo)

UAS lines

- *UAS-Khc^{FL}::GFP* (3rd, unpublished; Isabel Palacios)
- *UAS-Khc⁸²⁻⁷¹¹-GFP* (2nd, BL #9648; constitutively active Khc consisting of base pairs 248-2134 / aa 82-711; fused with EGFP sequence; flybase.org: [FB0198610](https://flybase.org/locus/FB0198610))
- *UAS-Khc1-850-GFP* (Loiseau et al., 2010; Isabel Palacios)
- *UAS-Khc-RNAi* (Lu et al., 2013; Vagnoni et al., 2016; BL #35770)
- *UAS-Sod1* (J. Hu and J.P. Phillips, unpublished)
- *UAS-Duox* (Ha et al., 2005; Matthias Landgraf)
- *UAS-Gapdh-IR* (*Gapdh1^{GD7467}*; Vienna *Drosophila* Resource Centre)

Cloning of *UAS-Nox-YPet*

10xUAS-IVS-Nox::YPet was generated by using the *pJFRC12-10xUAS-IVS-myr-GFP* vector (Addgene 26222; Pfeiffer et al., 2010) as a backbone which was modified by substituting GFP with YPet (Nguyen and Daugherty, 2005) plus an N-terminal flexible linker, amplified from dFlex_YPet_phase0 (Gärtig et al., 2019) using primer ML1 and ML2, and inserted by the Klenow Assembly Method (tinyurl.com/4r99uv8m) into the XbaI/BamHI sites producing Vector 1: *pJFRC12-10xUAS-IVS-myr-linker-YPet*. Nox cDNA was amplified from a DGRC (*Drosophila* Genomics Resource Center) cDNA library clone using primers ML5 and ML6, located in a *pOTB7* vector backbone, and inserted into BamHI/XhoI sites of Vector 1. Constructs were sent to FlyORF for transgenesis, and targeted via PhiC31-mediated site-specific insertion to the *PBac{y⁺-attP-3B}VK00040* landing site (Bloomington line #9755) on the third chromosome (3R, 87B10).

ML1	gacatcatcagaccacgcggatccggctccgccggctccgccgccggctccggcgagttcgtgtcca agggcgag
ML2	gttccttcacaaagatcctctagattactgtacagctcgttcattgccc
ML5	ggagccggcgagccggatccgaagcactccttacgaaaggcaaatccgt
ML6	cttcaggcgggccggctcgagaatcaaaatgaacgcggaccaggagtc

Drosophila primary cell culture

Drosophila primary neuron cultures were performed as published previously (Prokop et al., 2012; Qu et al., 2017). In brief, stage 11 embryos were treated for 1 min with bleach to remove the chorion, sterilized for ~30 s in 70% ethanol, washed in sterile Schneider's/FCS, and eventually homogenized with micro-pestles in 1.5 centrifuge tubes containing 21 embryos per 100µl dispersion medium and left to incubate for 5 min at 37°C. Cells were washed with Schneider's medium (Gibco), spun down for 4 mins at 650g, supernatant was removed and cells re-suspended in 90µl of Schneider's medium containing 20% fetal calf serum (Gibco). 30µl drops were placed on cover slips. Cells were allowed to adhere for ~2hrs either directly on glass or on cover slips coated with a 5 µg/ml solution of concanavalin A, and then grown as a hanging drop culture for hours or days at 26°C as indicated in each experiment.

To abolish maternal rescue of mutants, i.e. masking of the mutant phenotype caused by deposition of normal gene product from the healthy gene copy of the heterozygous mothers in the oocyte (Prokop, 2013), we used a pre-culture strategy (Prokop et al., 2012; Sánchez-Soriano et al., 2010) where cells were kept for 5 days in a tube before they were plated on a coverslip.

Cells were treated with 100 µM Trolox (Sigma; stepwise diluted from a 100mM stock solution in ethanol) or 100 µM DEM prepared in 100% ethanol. For controls (vehicle treatment), equivalent concentrations of vehicle (sterile H₂O or 100% ethanol) were diluted in cell culture medium. All reagents were purchased from Sigma-Aldrich, unless otherwise stated.

For visualisation of mitochondria, cell cultures were incubated with 400nM MitoTracker Red CMXRos (Invitrogen; Klionsky et al., 2012) for 30min at room temperature (RT); stock solutions were prepared in DMSO and diluted in cell culture medium to the final concentration. Following incubation, cultures were then fixed and stained following the procedures below.

Immunohistochemistry

Primary fly neurons were fixed in 4% paraformaldehyde (PFA) in 0.05M phosphate buffer (PB; pH 7–7.2) for 30min at room temperature (RT). Antibody staining and washes were performed with PBT. Staining reagents: anti-tubulin (clone DM1A, mouse, 1:1000, Sigma; alternatively, clone YL1/2, rat, 1:500, Millipore Bioscience Research Reagents); anti-Syt (1:1000; rabbit; Sean Sweeney); anti-GFP (1:500, rabbit, ab290, Abcam); Cy3-conjugated anti-HRP (goat, 1:100, Jackson ImmunoResearch); FITC-, Cy3- or Cy5-conjugated secondary antibodies (1:200; donkey, purified, Jackson Immuno Research); F-actin was stained with Phalloidin conjugated with TRITC/Alexa647, FITC or Atto647N (1:200; Invitrogen and Sigma). Specimens were embedded in ProLong Gold Antifade mounting medium.

Microscopy and data analysis

Standard documentation was performed with AxioCam monochrome digital cameras (Carl Zeiss Ltd.) mounted on BX50WI or BX51 Olympus compound fluorescent microscopes. To determine the degree of MT disorganisation in axons we used the "MT disorganisation index" (MDI) (Qu et al., 2017): the area of disorganisation was measured using the freehand selection tool in Fiji/ImageJ; this value was then divided by axon length (see above) multiplied by 0.5 µm (typical axon diameter, thus approximating the expected area of the axon if it were not disorganised). To quantify the number of synaptic densities in mature neurons in culture, we used ImageJ, first thresholding to select synaptic densities from axons of single isolated cells, followed by particle analysis. For statistical

analyses, Kruskal–Wallis one-way ANOVA with *post hoc* Dunn’s test or Mann–Whitney Rank Sum Tests were used to compare groups. The data used for our analyses will be made available on request from the authors.

Ethical statement

An ethical statement is not required.

References

- Ahmad, F. J., He, Y., Myers, K. A., Hasaka, T. P., Francis, F., Black, M. M., Baas, P. W. (2006). Effects of dynactin disruption and dynein depletion on axonal microtubules. *Traffic* **7**, 524-37 -- <https://doi.org/10.1111/j.1600-0854.2006.00403.x>
- Albano, R., Raddatz, N. J., Hjelmhaug, J., Baker, D. A., Lobner, D. (2015). Regulation of System x(c)(-) by Pharmacological Manipulation of Cellular Thiols. *Oxidative Medicine and Cellular Longevity* **2015**, 269371 -- <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4407525/>
- Alves-Silva, J., Sánchez-Soriano, N., Beaven, R., Klein, M., Parkin, J., Millard, T., Bellen, H., Venken, K. J. T., Ballestrem, C., Kammerer, R. A., Prokop, A. (2012). Spectraplakins promote microtubule-mediated axonal growth by functioning as structural microtubule-associated proteins and EB1-dependent +TIPs (Tip Interacting Proteins). *J. Neurosci* **32**, 9143-58 -- <https://doi.org/10.1523/JNEUROSCI.0416-12.2012>
- Amberger, J. S., Bocchini, C. A., Schiettecatte, F., Scott, A. F., Hamosh, A. (2015). OMIM.org: Online Mendelian Inheritance in Man (OMIM(R)), an online catalog of human genes and genetic disorders. *Nucleic Acids Res* **43**, D789-98 -- <http://www.ncbi.nlm.nih.gov/pubmed/25428349>
- Andreu-Carbó, M., Fernandes, S., Velluz, M.-C., Kruse, K., Aumeier, C. (2021). Motor usage imprints microtubule stability along the shaft. *Dev Cell* -- <https://doi.org/10.1016/j.devcel.2021.11.019>
- Andrews, S. B., Gallant, P. E., Leapman, R. D., Schnapp, B. J., Reese, T. S. (1993). Single kinesin molecules crossbridge microtubules in vitro. *Proc Natl Acad Sci U S A* **90**, 6503-7 -- <https://doi.org/10.1073/pnas.90.14.6503>
- Anh, N. T., Nishitani, M., Harada, S., Yamaguchi, M., Kamei, K. (2011). Essential role of Duox in stabilization of *Drosophila* wing. *J Biol Chem* **286**, 33244-51 -- <https://doi.org/10.1074/jbc.M111.263178>
- Arpağ, G., Norris, S. R., Mousavi, S. I., Soppina, V., Verhey, K. J., Hancock, W. O., Tüzel, E. (2019). Motor dynamics underlying cargo transport by pairs of kinesin-1 and kinesin-3 motors. *Biophys J* **116**, 1115-1126 -- <https://doi.org/10.1016/j.bpj.2019.01.036>
- Baloh, R. H., Schmidt, R. E., Pestronk, A., Milbrandt, J. (2007). Altered axonal mitochondrial transport in the pathogenesis of Charcot-Marie-Tooth disease from mitofusin 2 mutations. *J Neurosci* **27**, 422-30 -- <http://www.ncbi.nlm.nih.gov/pubmed/17215403>
- Baqri, R., Charan, R., Schimmelpfeng, K., Chavan, S., Ray, K. (2006). Kinesin-2 differentially regulates the anterograde axonal transports of acetylcholinesterase and choline acetyltransferase in *Drosophila*. *J Neurobiol* **66**, 378-92 -- <http://www.ncbi.nlm.nih.gov/pubmed/16408306>
- Beaven, R., Dzhindzhev, N. S., Qu, Y., Hahn, I., Dajas-Bailador, F., Ohkura, H., Prokop, A. (2015). *Drosophila* CLIP-190 and mammalian CLIP-170 display reduced microtubule plus end association in the nervous system. *Mol Biol Cell* **26**, 1491-508 -- <https://doi.org/10.1091/mbc.E14-06-1083>

- Bedard, K., Krause, K.-H. (2007). The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Phys Rev* **87**, 245-313 -- <https://doi.org/10.1152/physrev.00044.2005>
- Bowman, A. B., Kamal, A., Ritchings, B. W., Philp, A. V., McGrail, M., Gindhart, J. G., Goldstein, L. S. B. (2000). Kinesin-dependent axonal transport is mediated by the Sunday driver (Syd) protein. *Cell* **103**, 583-94 -- [http://dx.doi.org/10.1016/S0092-8674\(00\)00162-8](http://dx.doi.org/10.1016/S0092-8674(00)00162-8)
- Brady, S. T., Morfini, G. A. (2017). Regulation of motor proteins, axonal transport deficits and adult-onset neurodegenerative diseases. *Neurobiol Dis* **105**, 273-282 -- <https://doi.org/10.1016/j.nbd.2017.04.010>
- Brendza, K. M., Rose, D. J., Gilbert, S. P., Saxton, W. M. (1999). Lethal kinesin mutations reveal amino acids important for ATPase activation and structural coupling. *J Biol Chem* **274**, 31506-14 -- <https://doi.org/10.1074/jbc.274.44.31506>
- Budaitis, B. G., Badiyan, S., Yue, Y., Blasius, T. L., Reinemann, D. N., Lang, M. J., Cianfrocco, M. A., Verhey, K. J. (2021). A kinesin-1 variant reveals motor-induced microtubule damage in cells. *bioRxiv*, 2021.10.19.464974 -- <https://www.biorxiv.org/content/biorxiv/early/2021/10/20/2021.10.19.464974.full>
- Bunge, M. B. (1973). Fine structure of nerve fibers and growth cones of isolated sympathetic neurons in culture. *J Cell Biol* **56**, 713-35 -- <http://doi.org/10.1083/jcb.56.3.713>
- Caglinec, M., Safiulina, D., Liiv, M., Liiv, J., Choubey, V., Wareski, P., Veksler, V., Kaasik, A. (2013). Principles of the mitochondrial fusion and fission cycle in neurons. *J Cell Sci* **126**, 2187-97 -- <http://jcs.biologists.org/content/joces/126/10/2187.full.pdf>
- Calkins, D. J. (2013). Age-Related Changes in the Visual Pathways: Blame It on the AxonAge-Related Changes in the Visual Pathways. *Invest Ophthalmol Vis Sci* **54**, ORSF 37-41 -- <https://dx.doi.org/10.1167/iovs.13-12784>
- Campbell, P. D., Shen, K., Sapio, M. R., Glenn, T. D., Talbot, W. S., Marlow, F. L. (2014). Unique function of kinesin Kif5A in localization of mitochondria in axons. *J Neurosci* **34**, 14717-32 -- <http://www.ncbi.nlm.nih.gov/pubmed/25355224>
- Cao, L., Cantos-Fernandes, S., Gigant, B. (2017). The structural switch of nucleotide-free kinesin. *Scientific Reports* **7**, 42558 -- <https://doi.org/10.1038/srep42558>
- Cao, Z., Lindsay, J. G. (2017). The Peroxiredoxin Family: An Unfolding Story. *Subcell Biochem* **83**, 127-147 -- <https://www.ncbi.nlm.nih.gov/pubmed/28271475>
- Castro, I. G., Richards, D. M., Metz, J., Costello, J. L., Passmore, J. B., Schrader, T. A., Gouveia, A., Ribeiro, D., Schrader, M. (2018). A role for Mitochondrial Rho GTPase 1 (MIRO1) in motility and membrane dynamics of peroxisomes. *Traffic* **19**, 229-242 -- <https://onlinelibrary.wiley.com/doi/abs/10.1111/tra.12549>
- Chang, H. Y., Cheng, H. Y., Tsao, A. N., Liu, C., Tsai, J. W. (2019). Multiple functions of KBP in neural development underlie brain anomalies in Goldberg-Shprintzen syndrome. *Front Mol Neurosci* **12**, 265 -- <https://www.ncbi.nlm.nih.gov/pubmed/31736709>
- Chiba, K., Takahashi, H., Chen, M., Obinata, H., Arai, S., Hashimoto, K., Oda, T., McKenney, R. J., Niwa, S. (2019). Disease-associated mutations hyperactivate KIF1A motility and anterograde axonal transport of synaptic vesicle precursors. *Proc Natl Acad Sci* **116**, 18429-34 -- <https://doi.org/10.1073/pnas.1905690116>
- Coleman, M. (2005). Axon degeneration mechanisms: commonality amid diversity. *Nat Rev Neurosci* **6**, 889-98 -- <https://doi.org/10.1038/nrn1788>
- Cong, D., Ren, J., Zhou, Y., Wang, S., Liang, J., Ding, M., Feng, W. (2021). Motor domain-mediated autoinhibition dictates axonal transport by the kinesin UNC-104/KIF1A. *PLoS Genet* **17**, e1009940 -- <https://doi.org/10.1371/journal.pgen.1009940>

- 743 Cook, R. K., Christensen, S. J., Deal, J. A., Coburn, R. A., Deal, M. E., Gresens, J. M., Kaufman,
744 T. C., Cook, K. R. (2012). The generation of chromosomal deletions to provide extensive
745 coverage and subdivision of the *Drosophila melanogaster* genome. *Genome Biol* **13**, R21 --
746 <http://doi.org/10.1186/gb-2012-13-3-r21>
- 747 Covill-Cooke, C., Lopez-Domenech, G., Birsa, N., Kittler, J. T. (2017). The mitochondrial Rho-
748 GTPase, Miro, is resident at peroxisomes and regulates peroxisomal trafficking and
749 morphology. *bioRxiv* -- <https://www.biorxiv.org/content/biorxiv/early/2017/12/30/241208.full.pdf>
- 750 Cox, R. T., Spradling, A. C. (2006). Milton controls the early acquisition of mitochondria by
751 *Drosophila* oocytes. *Development* **133**, 3371-3377 -- <https://doi.org/10.1242/dev.02514>
- 752 Czanietcki, C., Ryan, T., Stykel, M. G., Drolet, J., Heide, J., Hallam, R., Wood, S., Coackley, C.,
753 Sherriff, K., Bailey, C. D. C., Ryan, S. D. (2019). Axonal pathology in hPSC-based models of
754 Parkinson's disease results from loss of Nrf2 transcriptional activity at the Map1b gene locus.
755 *Proc Natl Acad Sci* **116**, 14280-14289 -- <https://doi.org/10.1073/pnas.1900576116>
- 756 Dasgupta, A., Zheng, J., Bizzozero, O. A. (2012). Protein carbonylation and aggregation precede
757 neuronal apoptosis induced by partial glutathione depletion. *ASN Neuro* **4** --
758 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3322383/>
- 759 De Vos, K. J., Chapman, A. L., Tennant, M. E., Manser, C., Tudor, E. L., Lau, K. F., Brownlees, J.,
760 Ackerley, S., Shaw, P. J., McLoughlin, D. M., Shaw, C. E., Leigh, P. N., Miller, C. C. J.,
761 Grierson, A. J. (2007). Familial amyotrophic lateral sclerosis-linked SOD1 mutants perturb fast
762 axonal transport to reduce axonal mitochondria content. *Hum Mol Genet* **16**, 2720-2728 --
763 <http://www.ncbi.nlm.nih.gov/pubmed/17725983>
- 764 Debattisti, V., Gerencser, A. A., Saotome, M., Das, S., Hajnóczky, G. (2017). ROS control
765 mitochondrial motility through p38 and the motor adaptor Miro/Trak. *Cell Reports* **21**, 1667-1680
766 -- <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5710826/>
- 767 del Castillo, U., Gnazzo, M. M., Sorensen Turpin, C. G., Nguyen, K. C. Q., Semaya, E., Lam, Y.,
768 de Cruz, M. A., Bembenek, J. N., Hall, D. H., Riggs, B., Gelfand, V. I., Skop, A. R. (2019).
769 Conserved role for Ataxin-2 in mediating endoplasmic reticulum dynamics. *Traffic* **20**, 436-447 --
770 <https://doi.org/10.1111/tra.12647>
- 771 Derivery, E., Seum, C., Daeden, A., Loubéry, S., Holtzer, L., Jülicher, F., Gonzalez-Gaitan, M.
772 (2015). Polarized endosome dynamics by spindle asymmetry during asymmetric cell division.
773 *Nature* **528**, 280-285 -- <http://dx.doi.org/10.1038/nature16443>
- 774 Dimitrova-Paternoga, L., Jagtap, P. K. A., Cyrklaff, A., Vaishali, Lapouge, K., Sehr, P., Perez, K.,
775 Heber, S., Löw, C., Hennig, J., Ephrussi, A. (2021). Molecular basis of mRNA transport by a
776 kinesin-1-atypical tropomyosin complex. *Genes Dev* **35**, 976-91 --
777 <http://doi.org/10.1101/gad.348443.121>
- 778 Drerup, C. M., Lusk, S., Nechiporuk, A. (2016). Kif1B interacts with KBP to promote axon
779 elongation by localizing a microtubule regulator to growth cones. *J Neurosci* **36**, 7014-26 --
780 <https://doi.org/10.1523/JNEUROSCI.0054-16.2016>
- 781 Dumont, E., L. P., Do, C., Hess, H. (2015). Molecular wear of microtubules propelled by surface-
782 adhered kinesins. *Nat Nanotechnol* **10**, 166-9 -- <http://dx.doi.org/10.1038/nnano.2014.334>
- 783 Duttaroy, A., Paul, A., Kundu, M., Belton, A. (2003). A Sod2 null mutation confers severely reduced
784 adult life span in *Drosophila*. *Genetics* **165**, 2295-9 --
785 <https://doi.org/10.1093/genetics/165.4.2295>
- 786 Faust, J. E., Manisundaram, A., Ivanova, P. T., Milne, S. B., Summerville, J. B., Brown, H. A.,
787 Wangler, M., Stern, M., McNew, J. A. (2014). Peroxisomes are required for lipid metabolism and
788 muscle function in *Drosophila melanogaster*. *PLoS One* **9**, e100213 --
789 <https://doi.org/10.1371/journal.pone.0100213>

- 790 Fischer, L. R., Li, Y., Asress, S. A., Jones, D. P., Glass, J. D. (2012). Absence of SOD1 leads to
791 oxidative stress in peripheral nerve and causes a progressive distal motor axonopathy. *Exp*
792 *Neurol* **233**, 163-71 -- <http://www.ncbi.nlm.nih.gov/pubmed/21963651>
- 793 Fransen, M., Lismont, C., Walton, P. (2017). The Peroxisome-Mitochondria Connection: How and
794 Why? *Int J Mol Sci* **18**, E1126 -- <https://doi.org/10.3390/ijms18061126>
- 795 Fügler, P., Sreekumar, V., Schule, R., Kern, J. V., Stanchev, D. T., Schneider, C. D., Karle, K. N.,
796 Daub, K. J., Siegert, V. K., Flötenmeyer, M., Schwarz, H., Schöls, L., Rasse, T. M. (2012).
797 Spastic paraplegia mutation N256S in the neuronal microtubule motor KIF5A disrupts axonal
798 transport in a *Drosophila* HSP model. *PLoS Genet* **8**, e1003066 --
799 <http://doi.org/10.1371/journal.pgen.1003066>
- 800 Fundu, T. M., Kapepula, P. M., Esimo, J. M., Remacle, J., Ngombe, N. K. (2019). Subcellular
801 Localization of Glutathione Peroxidase, Change in Glutathione System during Ageing and
802 Effects on Cardiometabolic Risks and Associated Diseases. *Intech Open*, 1-19 --
803 <https://doi.org/10.5772/intechopen.89384>
- 804 Gabrych, D. R., Lau, V. Z., Niwa, S., Silverman, M. A. (2019). Going too far is the same as falling
805 short: kinesin-3 family members in hereditary spastic paraplegia. *Front Cell Neurosci* **13**, 419 --
806 <https://doi.org/10.3389/fncel.2019.00419>
- 807 Gärtig, P.-A., Ostrovsky, A., Manhart, L., Giachello, C. N. G., Kovacevic, T., Lustig, H., Chwalla, B.,
808 Cachero, S., Baines, R. A., Landgraf, M., Evers, J. F. (2019). Motor circuit function is stabilized
809 during postembryonic growth by anterograde trans-synaptic Jelly Belly - Anaplastic Lymphoma
810 Kinase signaling. *bioRxiv*, 841106 -- <https://doi.org/10.1101/841106>
- 811 Gauger, A. K., Goldstein, L. S. (1993). The *Drosophila* kinesin light chain. Primary structure and
812 interaction with kinesin heavy chain. *J Biol Chem* **268**, 13657-66 --
813 [https://doi.org/10.1016/S0021-9258\(19\)38698-3](https://doi.org/10.1016/S0021-9258(19)38698-3)
- 814 Gepner, J., Li, M., Ludmann, S., Kortas, C., Boylan, K., Iyadurai, S. J., McGrail, M., Hays, T. S.
815 (1996). Cytoplasmic dynein function is essential in *Drosophila melanogaster*. *Genetics* **142**,
816 865-78 -- <https://doi.org/10.1093/genetics/142.3.865>
- 817 Gigant, B., Wang, W., Dreier, B., Jiang, Q., Pecqueur, L., Plückthun, A., Wang, C., Knossow, M.
818 (2013). Structure of a kinesin–tubulin complex and implications for kinesin motility. *Nature*
819 *Structural & Molecular Biology* **20**, 1001-1007 -- <https://doi.org/10.1038/nsmb.2624>
- 820 Gindhart, J. G. (2006). Towards an understanding of kinesin-1 dependent transport pathways
821 through the study of protein-protein interactions. *Brief Funct Genomic Proteomic* **5**, 74-86 --
822 <https://doi.org/10.1093/bfqp/ell002>
- 823 Gindhart, J. G., Chen, J., Faulkner, M., Gandhi, R., Doerner, K., Wisniewski, T., Nandlestadt, A.
824 (2003). The kinesin-associated protein UNC-76 is required for axonal transport in the *Drosophila*
825 nervous system. *Mol Biol Cell* **14**, 3356-65 -- <http://www.ncbi.nlm.nih.gov/pubmed/12925768>
- 826 Gindhart, J. G., Jr., Desai, C. J., Beushausen, S., Zinn, K., Goldstein, L. S. (1998). Kinesin light
827 chains are essential for axonal transport in *Drosophila*. *J Cell Biol* **141**, 443-54 --
- 828 Giordano, M. E., Caricato, R., Lionetto, M. G. (2020). Concentration dependence of the antioxidant
829 and prooxidant activity of trolox in HeLa cells: involvement in the induction of apoptotic volume
830 decrease. *Antioxidants* **9**, 1058 -- <https://doi.org/doi:10.3390/antiox9111058>
- 831 Glater, E. E., Megeath, L. J., Stowers, R. S., Schwarz, T. L. (2006). Axonal transport of
832 mitochondria requires Milton to recruit kinesin heavy chain and is light chain independent. *J Cell*
833 *Biol* **173**, 545-57 -- <https://doi.org/10.1083/jcb.200601067>
- 834 Glynn, S. E. (2017). Multifunctional Mitochondrial AAA Proteases. *Front Mol Biosci* **4**, 34 --
835 <http://www.ncbi.nlm.nih.gov/pubmed/28589125>
- 836 Goldblum, R. R., McClellan, M., White, K., Gonzalez, S. J., Thompson, B. R., Vang, H. X., Cohen,
837 H., Higgins, L., Markowski, T. W., Yang, T.-Y., Metzger, J. M., Gardner, M. K. (2021). Oxidative

- 838 stress pathogenically remodels the cardiac myocyte cytoskeleton via structural alterations to the
839 microtubule lattice. *Developmental Cell* **56**, 2252-2266.e6 --
840 <https://doi.org/10.1016/j.devcel.2021.07.004>
- 841 Gonçalves-Pimentel, C., Gombos, R., Mihály, J., Sánchez-Soriano, N., Prokop, A. (2011).
842 Dissecting regulatory networks of filopodia formation in a *Drosophila* growth cone model. *PLoS*
843 *ONE* **6**, e18340 -- <https://doi.org/10.1371/journal.pone.0018340>
- 844 Guedes-Dias, P., Holzbaur, E. L. F. (2019). Axonal transport: Driving synaptic function. *Science*
845 **366**, eaaw9997 -- <https://doi.org/10.1126/science.aaw9997>
- 846 Guo, W., Stoklund Dittlau, K., Van Den Bosch, L. (2020). Axonal transport defects and
847 neurodegeneration: molecular mechanisms and therapeutic implications. *Semin Cell Dev Biol*
848 **99**, 133-150 -- <https://doi.org/10.1016/j.semcdb.2019.07.010>
- 849 Guo, X., Macleod, G. T., Wellington, A., Hu, F., Panchumathi, S., Schoenfield, M., Marin, L.,
850 Charlton, M. P., Atwood, H. L., Zinsmaier, K. E. (2005). The GTPase dMiro is required for
851 axonal transport of mitochondria to *Drosophila* synapses. *Neuron* **47**, 379-93 --
852 <https://doi.org/10.1016/j.neuron.2005.06.027>
- 853 Ha, E. M., Oh, C. T., Bae, Y. S., Lee, W. J. (2005). A direct role for dual oxidase in *Drosophila* gut
854 immunity. *Science* **310**, 847-50 -- <https://doi.org/10.1126/science.1117311>
- 855 Hahn, I., Ronshaugen, M., Sánchez-Soriano, N., Prokop, A. (2016). Functional and genetic
856 analysis of spectraplakins in *Drosophila*. *Methods Enzymol* **569**, 373-405 --
857 <https://doi.org/10.1016/bs.mie.2015.06.022>
- 858 Hahn, I., Voelzmann, A., Liew, Y.-T., Costa-Gomes, B., Prokop, A. (2019). The model of local axon
859 homeostasis - explaining the role and regulation of microtubule bundles in axon maintenance
860 and pathology *Neural Dev* **14**, 10.1186/s13064-019-0134-0 -- [https://doi.org/10.1186/s13064-](https://doi.org/10.1186/s13064-019-0134-0)
861 [019-0134-0](https://doi.org/10.1186/s13064-019-0134-0)
- 862 Hahn, I., Voelzmann, A., Parkin, J., Fuelle, J. B., Slater, P. G., Lowery, L. A., Sanchez-Soriano, N.,
863 Prokop, A. (2021). Tau, XMAP215 and Eb co-operatively regulate microtubule polymerisation
864 and bundle formation in axons. *PLoS Genet* **17**, e1009647 --
865 <https://doi.org/10.1371/journal.pgen.1009647>
- 866 Hancock, W. O. (2014). Bidirectional cargo transport: moving beyond tug of war. *Nat Rev Mol Cell*
867 *Biol* **15**, 615-28 -- <http://doi.org/10.1038/nrm3853>
- 868 Harbauer, A. B. (2017). Mitochondrial health maintenance in axons. *Biochem Soc Trans* **45**, 1045-
869 1052 -- <http://www.ncbi.nlm.nih.gov/pubmed/28778985>
- 870 Haynes, E. M., He, J., Jean-Pierre, M., Eliceiri, K. W., Huiskens, J., Halloran, M. (2021). KLC4
871 shapes axon arbors during development and mediates adult behavior. *bioRxiv*,
872 2021.09.26.461872 -- <https://doi.org/10.1101/2021.09.26.461872>
- 873 Higuchi, H., Bronner, C. E., Park, H.-W., Endow, S. A. (2004). Rapid double 8-nm steps by a
874 kinesin mutant. *EMBO J* **23**, 2993-2999 -- <https://doi.org/10.1038/sj.emboj.7600306>
- 875 Hinckelmann, M. V., Virlogeux, A., Niehage, C., Poujol, C., Choquet, D., Hoflack, B., Zala, D.,
876 Saudou, F. (2016). Self-propelling vesicles define glycolysis as the minimal energy machinery
877 for neuronal transport. *Nat Commun* **7**, 13233 -- <http://doi.org/10.1038/ncomms13233>
- 878 Hirokawa, N., Niwa, S., Tanaka, Y. (2010). Molecular motors in neurons: transport mechanisms
879 and roles in brain function, development, and disease. *Neuron* **68**, 610-638 --
880 <http://www.ncbi.nlm.nih.gov/pubmed/21092854>
- 881 Hirst, C. S., Stamp, L. A., Bergner, A. J., Hao, M. M., Tran, M. X., Morgan, J. M., Dutschmann, M.,
882 Allen, A. M., Paxinos, G., Furlong, T. M., McKeown, S. J., Young, H. M. (2017). Kif1bp loss in
883 mice leads to defects in the peripheral and central nervous system and perinatal death. *Sci Rep*
884 **7**, 16676 -- <https://www.ncbi.nlm.nih.gov/pubmed/29192291>

- 885 Horiuchi, D., Barkus, R. V., Pilling, A. D., Gassman, A., Saxton, W. M. (2005). APLIP1, a kinesin
886 binding JIP-1/JNK scaffold protein, influences the axonal transport of both vesicles and
887 mitochondria in *Drosophila*. *Curr Biol* **15**, 2137-41 -- <https://doi.org/10.1016/j.cub.2005.10.047>
- 888 Hudson, D. A., Gannon, S. A., Thorpe, C. (2015). Oxidative protein folding: From thiol–disulfide
889 exchange reactions to the redox poise of the endoplasmic reticulum. *Free Radic Biol Med* **80**,
890 171-182 -- <https://doi.org/10.1016/j.freeradbiomed.2014.07.037>
- 891 Hunter, B., Allingham, J. S. (2020). These motors were made for walking. *Protein Sci* **29**, 1707-
892 1723 -- <https://doi.org/10.1002/pro.3895>
- 893 Hurd, D. D., Saxton, W. M. (1996). Kinesin mutations cause motor neuron disease phenotypes by
894 disrupting fast axonal transport in *Drosophila*. *Genetics* **144**, 1075-85 --
895 <http://www.ncbi.nlm.nih.gov/pubmed/8913751>
- 896 Iworima, D. G., Pasqualotto, B. A., Rintoul, G. L. (2016). Kif5 regulates mitochondrial movement,
897 morphology, function and neuronal survival. *Mol Cell Neurosci* **72**, 22-33 --
898 <https://doi.org/10.1016/j.mcn.2015.12.014>
- 899 Jana, S. C., Dutta, P., Jain, A., Singh, A., Adusumilli, L., Girotra, M., Kumari, D., Shirolkar, S., Ray,
900 K. (2021). Kinesin-2 transports Orco into the olfactory cilium of *Drosophila melanogaster* at
901 specific developmental stages. *PLOS Genet* **17**, e1009752 --
902 <https://doi.org/10.1371/journal.pgen.1009752>
- 903 Jolly, A. L., Kim, H., Srinivasan, D., Lakonishok, M., Larson, A. G., Gelfand, V. I. (2010). Kinesin-1
904 heavy chain mediates microtubule sliding to drive changes in cell shape. *Proc Natl Acad Sci U S*
905 *A* **107**, 12151-6 -- <https://doi.org/10.1073/pnas.1004736107>
- 906 Kaan, H. Y. K., Hackney, D. D., Kozielski, F. (2011). The structure of the kinesin-1 motor-tail
907 complex reveals the mechanism of autoinhibition. *Science* **333**, 883-5 --
908 <https://www.science.org/doi/abs/10.1126/science.1204824>
- 909 Kawaguchi, K. (2013). Role of kinesin-1 in the pathogenesis of SPG10, a rare form of hereditary
910 spastic paraplegia. *Neuroscientist* **19**, 336-44 -- <http://doi.org/10.1177/1073858412451655>
- 911 Kelliher, M. T., Yue, Y., Ng, A., Kamiyama, D., Huang, B., Verhey, K. J., Wildonger, J. (2018).
912 Autoinhibition of kinesin-1 is essential to the dendrite-specific localization of Golgi outposts. *J*
913 *Cell Biol* **217**, 2531-2547 -- <http://doi.org/10.1083/jcb.201708096>
- 914 Keren-Kaplan, T., Bonifacio, J. S. (2021). ARL8 relieves SKIP autoinhibition to enable coupling of
915 lysosomes to kinesin-1. *Curr Biol* **31**, 540-554.e5 -- <https://doi.org/10.1016/j.cub.2020.10.071>
- 916 Kevenaar, Josta T., Bianchi, S., van Spronsen, M., Olieric, N., Lipka, J., Frias, Cátia P.,
917 Mikhaylova, M., Harterink, M., Keijzer, N., Wulf, Phebe S., Hilbert, M., Kapitein, Lukas C.,
918 de Graaff, E., Ahkmanova, A., Steinmetz, Michel O., Hoogenraad, Casper C. (2016). Kinesin-
919 Binding protein controls microtubule dynamics and cargo trafficking by regulating kinesin motor
920 activity. *Curr Biol* **26**, 849-61 -- <http://dx.doi.org/10.1016/j.cub.2016.01.048>
- 921 Khan, S. J., Abidi, S. N. F., Skinner, A., Tian, Y., Smith-Bolton, R. K. (2017). The *Drosophila* Duox
922 maturation factor is a key component of a positive feedback loop that sustains regeneration
923 signaling. *PLOS Genet* **13**, e1006937 -- <https://doi.org/10.1371/journal.pgen.1006937>
- 924 Klionsky, D. J., Abdalla, F. C., Abeliovich, H., Abraham, R. T., Acevedo-Arozena, A., Adeli, K.,
925 Agholme, L., Agnello, M., Agostinis, P., Aguirre-Ghiso, J. A., Ahn, H. J., Ait-Mohamed, O., Ait-
926 Si-Ali, S., Akematsu, T., Akira, S., Al-Younes, H. M., Al-Zeer, M. A., Albert, M. L., Albin, R. L.,
927 Alegre-Abarrategui, J., Aleo, M. F., Alirezaei, M., Almasan, A., Almonte-Becerril, M., Amano, A.,
928 Amaravadi, R. K., Amarnath, S., Amer, A. O., Andrieu-Abadie, N., Anantharam, V., Ann, D. K.,
929 Anoopkumar-Dukie, S., Aoki, H., Apostolova, N., Arancia, G., Aris, J. P., Asanuma, K., Asare,
930 N. Y. O., Ashida, H., Askanas, V., Askew, D. S., Auberger, P., Baba, M., Backues, S. K.,
931 Baehrecke, E. H., Bahr, B. A., Bai, X.-Y., Bailly, Y., Baiocchi, R., Baldini, G., Balduini, W.,
932 Ballabio, A., Bamber, B. A., Bampton, E. T. W., Juhász, G., Bartholomew, C. R., Bassham, D.
933 C., Bast, R. C., Batoko, H., Bay, B.-H., Beau, I., Béchet, D. M., Begley, T. J., Behl, C.,

- Behrends, C., Bekri, S., Bellaire, B., Bendall, L. J., Benetti, L., Berliocchi, L., Bernardi, H., Bernassola, F., Besteiro, S., Bhatia-Kissova, I., Bi, X., Biard-Piechaczyk, M., Blum, J. S., Boise, L. H., Bonaldo, P., Boone, D. L., Bornhauser, B. C., Bortoluci, K. R., Bossis, I., Bost, F., Bourquin, J.-P., Boya, P., Boyer-Guittaut, M., Bozhkov, P. V., Brady, N. R., Brancolini, C., Brech, A., Brenman, J. E., Brennand, A., Bresnick, E. H., Brest, P., Bridges, D., Bristol, M. L., Brookes, P. S., Brown, E. J., Brumell, J. H., et al. (2012). Guidelines for the use and interpretation of assays for monitoring autophagy. *Autophagy* **8**, 445-544 -- <https://doi.org/10.4161/auto.19496>
- Koper, A., Schenck, A., Prokop, A. (2012). Analysis of adhesion molecules and basement membrane contributions to synaptic adhesion at the *Drosophila* embryonic NMJ. *PLoS One* **7**, e36339 -- <http://doi.org/10.1371/journal.pone.0036339>
- Koushika, S. P. (2008). "JIP"ing along the axon: the complex roles of JIPs in axonal transport. *Bioessays* **30**, 10-4 -- <http://www.ncbi.nlm.nih.gov/pubmed/18081006>
- Kükürt, A., Gelen, V., Başer, Ö. F., Deveci, H. A., Karapehlivan, M. (2021). Thiols: Role in Oxidative Stress-Related Disorders. *Intech Open*, 1-21 -- <https://doi.org/10.5772/intechopen.96682>
- Kulkarni, A., Khan, Y., Ray, K. (2017). Heterotrimeric kinesin-2, together with kinesin-1, steers vesicular acetylcholinesterase movements toward the synapse. *FASEB J* **31**, 965-974 -- <http://doi.org/10.1096/fj.201600759RRR>
- Lam, A. T., VanDelinder, V., Kabir, A. M. R., Hess, H., Bachand, G. D., Kakugo, A. (2016). Cytoskeletal motor-driven active self-assembly in *in vitro* systems. *Soft Matter* **12**, 988-997 -- <http://dx.doi.org/10.1039/C5SM02042E>
- Lee, T., Luo, L. (1999). Mosaic analysis with a repressible neurotechnique cell marker for studies of gene function in neuronal morphogenesis. *Neuron* **22**, 451-61 -- [https://doi.org/10.1016/s0896-6273\(00\)80701-1](https://doi.org/10.1016/s0896-6273(00)80701-1)
- Liew, Y.-T. (2018). Roles of microtubule-binding proteins during axon growth and maintenance. In "Faculty of Biology, Medicine and Health", Vol. PhD, pp. 160. The University of Manchester, Manchester.
- Liu, X., Weaver, D., Shirihi, O., Hajnoczky, G. (2009). Mitochondrial 'kiss-and-run': interplay between mitochondrial motility and fusion-fission dynamics. *EMBO J* **28**, 3074-89 --
- Loiseau, P., Davies, T., Williams, L. S., Mishima, M., Palacios, I. M. (2010). *Drosophila* PAT1 is required for Kinesin-1 to transport cargo and to maximize its motility. *Development* **137**, 2763-2772 -- <https://doi.org/10.1242/dev.048108>
- López-Doménech, G., Covill-Cooke, C., Ivankovic, D., Halff, E. F., Sheehan, D. F., Norkett, R., Birsa, N., Kittler, J. T. (2018). Miro proteins coordinate microtubule- and actin-dependent mitochondrial transport and distribution. *The EMBO Journal* **37**, 321-336 -- <http://doi.org/10.15252/emboj.201696380>
- Lu, W., Fox, P., Lakonishok, M., Davidson, Michael W., Gelfand, Vladimir I. (2013). Initial neurite outgrowth in *Drosophila* neurons is driven by Kinesin-powered microtubule sliding. *Curr Biol* **23**, 1018-1023 -- <http://dx.doi.org/10.1016/j.cub.2013.04.050>
- Lu, W., Lakonishok, M., Gelfand, V. I. (2015). Kinesin-1-powered microtubule sliding initiates axonal regeneration in *Drosophila* cultured neurons. *Mol Biol Cell* -- <http://doi.org/10.1091/mbc.E14-10-1423>
- Luo, L., Liao, Y. J., Jan, L. Y., Jan, Y. N. (1994). Distinct morphogenetic functions of similar small GTPases: *Drosophila* Drac1 is involved in axonal outgrowth and myoblast fusion. *Genes Dev.* **8**, 1787-1802 -- <https://doi.org/10.1101/gad.8.15.1787>
- Lyons, D. A., Naylor, S. G., Mercurio, S., Dominguez, C., Talbot, W. S. (2008). KBP is essential for axonal structure, outgrowth and maintenance in zebrafish, providing insight into the cellular

- 982 basis of Goldberg-Shprintzen syndrome. *Development* **135**, 599-608 --
983 <https://doi.org/10.1242/dev.012377>
- 984 Lyons, D. A., Naylor, S. G., Scholze, A., Talbot, W. S. (2009). Kif1b is essential for mRNA
985 localization in oligodendrocytes and development of myelinated axons. *Nat Genet* **41**, 854-8 --
986 <https://doi.org/10.1038/ng.376>
- 987 Mackay, W. J., Bewley, G. C. (1989). The genetics of catalase in *Drosophila melanogaster*.
988 isolation and characterization of acatalasemic mutants. *Genetics* **122**, 643-52 --
989 <https://doi.org/10.1093/genetics/122.3.643>
- 990 Maday, S., Twelvetrees, Alison E., Moughamian, Armen J., Holzbaur, Erika L. F. (2014). Axonal
991 transport: cargo-specific mechanisms of motility and regulation. *Neuron* **84**, 292-309 --
992 <http://dx.doi.org/10.1016/j.neuron.2014.10.019>
- 993 Mandal, A., Drerup, C. M. (2019). Axonal transport and mitochondrial function in neurons. *Front*
994 *Cell Neurosci* **13**, 373 -- <https://doi.org/10.3389/fncel.2019.00373>
- 995 Marelja, Z., Dambowsky, M., Bolis, M., Georgiou, M. L., Garattini, E., Missirlis, F., Leimkühler, S.
996 (2014). The four aldehyde oxidases of *Drosophila melanogaster* have different gene expression
997 patterns and enzyme substrate specificities. *J Exp Biol* **217**, 2201-11 --
998 <https://doi.org/10.1242/jeb.102129>
- 999 Marner, L., Nyengaard, J. R., Tang, Y., Pakkenberg, B. (2003). Marked loss of myelinated nerve
1000 fibers in the human brain with age. *J Comp Neurol* **462**, 144-52 --
1001 <http://www.ncbi.nlm.nih.gov/pubmed/12794739>
- 1002 Marygold, S. J., Crosby, M. A., Goodman, J. L. (2016). Using FlyBase, a database of *Drosophila*
1003 genes and genomes. *Methods Mol Biol* **1478**, 1-31 --
1004 <http://www.ncbi.nlm.nih.gov/pubmed/27730573>
- 1005 Mauvezin, C., Neisch, A. L., Ayala, C. I., Kim, J., Beltrame, A., Braden, C. R., Gardner, M. K.,
1006 Hays, T. S., Neufeld, T. P. (2016). Coordination of autophagosome–lysosome fusion and
1007 transport by a Klp98A–Rab14 complex in *Drosophila*. *J Cell Sci* **129**, 971-982 --
1008 <https://doi.org/10.1242/jcs.175224>
- 1009 Millburn, G. H., Crosby, M. A., Gramates, L. S., Tweedie, S. (2016). FlyBase portals to human
1010 disease research using *Drosophila* models. *Dis Model Mech* **9**, 245-52 --
1011 <http://www.ncbi.nlm.nih.gov/pubmed/26935103>
- 1012 Misgeld, T., Schwarz, T. L. (2017). Mitostasis in Neurons: Maintaining Mitochondria in an Extended
1013 Cellular Architecture. *Neuron* **96**, 651-666 -- <http://doi.org/10.1016/j.neuron.2017.09.055>
- 1014 Moughamian, A. J., Osborn, G. E., Lazarus, J. E., Maday, S., Holzbaur, E. L. (2013). Ordered
1015 recruitment of dynactin to the microtubule plus-end is required for efficient initiation of retrograde
1016 axonal transport. *J Neurosci* **33**, 13190-203 -- <http://www.ncbi.nlm.nih.gov/pubmed/23926272>
- 1017 Nangaku, M., Sato-Yoshitake, R., Okada, Y., Noda, Y., Takemura, R., Yamazaki, H., Hirokawa, N.
1018 (1994). KIF1B, a novel microtubule plus end-directed monomeric motor protein for transport of
1019 mitochondria. *Cell* **79**, 1209-20 --
- 1020 Nguyen, A. W., Daugherty, P. S. (2005). Evolutionary optimization of fluorescent proteins for
1021 intracellular FRET. *Nat Biotechnol* **23**, 355-360 -- <https://doi.org/10.1038/nbt1066>
- 1022 Nicita, F., Ginevrino, M., Travaglini, L., D'Arrigo, S., Zorzi, G., Borgatti, R., Terrone, G.,
1023 Catteruccia, M., Vasco, G., Brankovic, V., Siliquini, S., Romano, S., Veredice, C., Pedemonte,
1024 M., Armando, M., Lettori, D., Stregapede, F., Bosco, L., Sferra, A., Tessarollo, V., Romaniello,
1025 R., Ristori, G., Bertini, E., Valente, E. M., Zanni, G. (2020). Heterozygous KIF1A variants
1026 underlie a wide spectrum of neurodevelopmental and neurodegenerative disorders. *J Med*
1027 *Genet* -- <https://doi.org/10.1136/jmedgenet-2020-107007>
- 1028 Nicolas, A., Kenna, K. P., Renton, A. E., Ticozzi, N., Faghri, F., Chia, R., Dominov, J. A., Kenna, B.
1029 J., Nalls, M. A., Keagle, P., Rivera, A. M., van Rheenen, W., Murphy, N. A., van Vugt, J. J. F. A.,

- 1030 Geiger, J. T., Van der Spek, R. A., Pliner, H. A., Shankaracharya, Smith, B. N., Marangi, G.,
1031 Topp, S. D., Abramzon, Y., Gkazi, A. S., Eicher, J. D., Kenna, A., Logullo, F. O., Simone, I.,
1032 Logroscino, G., Salvi, F., Bartolomei, I., Borghero, G., Murru, M. R., Costantino, E., Pani, C.,
1033 Puddu, R., Caredda, C., Piras, V., Tranquilli, S., Cuccu, S., Corongiu, D., Melis, M., Milia, A.,
1034 Marrosu, F., Marrosu, M. G., Floris, G., Cannas, A., Capasso, M., Caponnetto, C., Mancardi, G.,
1035 Origone, P., Mandich, P., Conforti, F. L., Cavallaro, S., Mora, G., Marinou, K., Sideri, R., Penco,
1036 S., Mosca, L., Lunetta, C., Pinter, G. L., Corbo, M., Riva, N., Carrera, P., Volanti, P., Mandrioli,
1037 J., Fini, N., Fasano, A., Tremolizzo, L., Arosio, A., Ferrarese, C., Trojsi, F., Tedeschi, G.,
1038 Monsurrò, M. R., Piccirillo, G., Femiano, C., Ticca, A., Ortu, E., La Bella, V., Spataro, R.,
1039 Colletti, T., Sabatelli, M., Zollino, M., Conte, A., Luigetti, M., Lattante, S., Santarelli, M., Petrucci,
1040 A., Pugliatti, M., Pirisi, A., Parish, L. D., Occhineri, P., Giannini, F., Battistini, S., Ricci, C.,
1041 Benigni, M., Cau, T. B., Loi, D., Calvo, A., Moglia, C., Brunetti, M., et al. (2018). Genome-wide
1042 analyses identify KIF5A as a novel ALS gene. *Neuron* **97**, 1268-83.e6 --
1043 <http://dx.doi.org/10.1016/j.neuron.2018.02.027>
- 1044 Nikić, I., Merkler, D., Sorbara, C., Brinkoetter, M., Kreutzfeldt, M., Bareyre, F. M., Brück, W.,
1045 Bishop, D., Misgeld, T., Kerschensteiner, M. (2011). A reversible form of axon damage in
1046 experimental autoimmune encephalomyelitis and multiple sclerosis. *Nat Med* **17**, 495-499 --
1047 <https://doi.org/10.1038/nm.2324>
- 1048 Niwa, S., Takahashi, H., Hirokawa, N. (2013). (beta)-Tubulin mutations that cause severe
1049 neuropathies disrupt axonal transport. *EMBO J* **32**, 1352-1364 --
1050 <http://dx.doi.org/10.1038/emboj.2013.59>
- 1051 O'Sullivan, N. C., Jahn, T. R., Reid, E., O'Kane, C. J. (2012). Reticulon-like-1, the Drosophila
1052 orthologue of the hereditary spastic paraplegia gene reticulon 2, is required for organization of
1053 endoplasmic reticulum and of distal motor axons. *Hum Mol Genet* **21**, 3356-65 --
1054 <http://www.ncbi.nlm.nih.gov/pubmed/22543973>
- 1055 Okumoto, K., Ono, T., Toyama, R., Shimomura, A., Nagata, A., Fujiki, Y. (2018). New splicing
1056 variants of mitochondrial Rho GTPase-1 (Miro1) transport peroxisomes. *J Cell Biol* **217**, 619-33
1057 -- <http://doi.org/10.1083/jcb.201708122>
- 1058 Orr, W. C., Radyuk, S. N., Sohal, R. S. (2013). Involvement of redox state in the aging of
1059 *Drosophila melanogaster*. *Antioxid Redox Signal* **19**, 788-803 --
1060 <https://doi.org/10.1089/ars.2012.5002>
- 1061 Oswald, M. C. W., Garnham, N., Sweeney, S. T., Landgraf, M. (2018). Regulation of neuronal
1062 development and function by ROS. *FEBS Lett* **592**, 679-691 -- <https://doi.org/10.1002/1873-3468.12972>
- 1064 Pack-Chung, E., Kurshan, P. T., Dickman, D. K., Schwarz, T. L. (2007). A *Drosophila* kinesin
1065 required for synaptic bouton formation and synaptic vesicle transport. *Nat Neurosci* **10**, 980-9 --
1066 <http://doi.org/10.1038/nn1936>
- 1067 Palma, F. R., He, C., Danes, J. M., Paviani, V., Coelho, D. R., Gantner, B. N., Bonini, M. G. (2020).
1068 Mitochondrial Superoxide Dismutase: What the Established, the Intriguing, and the Novel
1069 Reveal About a Key Cellular Redox Switch. *Antioxid Redox Signal* **32**, 701-714 --
1070 <https://www.ncbi.nlm.nih.gov/pubmed/31968997>
- 1071 Parks, A. L., Cook, K. R., Belvin, M., Dompe, N. A., Fawcett, R., Huppert, K., Tan, L. R., Winter, C.
1072 G., Bogart, K. P., Deal, J. E., Deal-Herr, M. E., Grant, D., Marcinko, M., Miyazaki, W. Y.,
1073 Robertson, S., Shaw, K. J., Tabios, M., Vysotskaia, V., Zhao, L., Andrade, R. S., Edgar, K. A.,
1074 Howie, E., Killpack, K., Milash, B., Norton, A., Thao, D., Whittaker, K., Winner, M. A., Friedman,
1075 L., Margolis, J., Singer, M. A., Kopczynski, C., Curtis, D., Kaufman, T. C., Plowman, G. D.,
1076 Duyk, G., Francis-Lang, H. L. (2004). Systematic generation of high-resolution deletion
1077 coverage of the *Drosophila melanogaster* genome. *Nature Genetics* **36**, 288-292 --
1078 <https://doi.org/10.1038/ng1312>

- 1079 Pascual-Ahuir, A., Manzanares-Estreder, S., Proft, M. (2017). Pro- and antioxidant functions of the
1080 peroxisome-mitochondria connection and its impact on aging and disease. *Oxid Med Cell*
1081 *Longev* **2017**, 9860841 -- <https://doi.org/10.1155/2017/9860841>
- 1082 Paul, D. M., Mantell, J., Borucu, U., Coombs, J., Surridge, K. J., Squire, J. M., Verkade, P.,
1083 Dodding, M. P. (2020). In situ cryo-electron tomography reveals filamentous actin within the
1084 microtubule lumen. *J Cell Biol* **219** -- <https://doi.org/10.1083/jcb.201911154>
- 1085 Paupe, V., Prudent, J. (2018). New insights into the role of mitochondrial calcium homeostasis in
1086 cell migration. *Biochem Biophys Res Commun* **500**, 75-86 --
1087 <https://doi.org/10.1016/j.bbrc.2017.05.039>
- 1088 Perez, S. E., Steller, H. (1996). Molecular and genetic analyses of lama, an evolutionarily
1089 conserved gene expressed in the precursors of the *Drosophila* first optic ganglion. *Mech Dev*
1090 **59**, 11-27 -- [https://doi.org/10.1016/0925-4773\(96\)00556-4](https://doi.org/10.1016/0925-4773(96)00556-4)
- 1091 Pfeiffer, B. D., Ngo, T.-T. B., Hibbard, K. L., Murphy, C., Jenett, A., Truman, J. W., Rubin, G. M.
1092 (2010). Refinement of tools for targeted gene expression in *Drosophila*. *Genetics* **186**, 735-755 -
1093 - <http://doi.org/10.1534/genetics.110.119917>
- 1094 Phillips, J. P., Campbell, S. D., Michaud, D., Charbonneau, M., Hilliker, A. J. (1989). Null mutation
1095 of copper/zinc superoxide dismutase in *Drosophila* confers hypersensitivity to paraquat and
1096 reduced longevity. *Proc Natl Acad Sci U S A* **86**, 2761-5 --
1097 <https://doi.org/10.1073/pnas.86.8.2761>
- 1098 Phillips, J. P., Tainer, J. A., Getzoff, E. D., Boulianne, G. L., Kirby, K., Hilliker, A. J. (1995).
1099 Subunit-destabilizing mutations in *Drosophila* copper/zinc superoxide dismutase:
1100 neuropathology and a model of dimer dysequilibrium. *Proc Natl Acad Sci U S A* **92**, 8574-78 --
1101 <http://doi.org/10.1073/pnas.92.19.8574>
- 1102 Pilling, A. D., Horiuchi, D., Lively, C. M., Saxton, W. M. (2006). Kinesin-1 and Dynein are the
1103 primary motors for fast transport of mitochondria in *Drosophila* motor axons. *Mol Biol Cell* **17**,
1104 2057-68 -- <https://doi.org/10.1091/mbc.e05-06-0526>
- 1105 Pompella, A., Visvikis, A., Paolicchi, A., Tata, V. D., Casini, A. F. (2003). The changing faces of
1106 glutathione, a cellular protagonist. *Biochemical Pharmacology* **66**, 1499-1503 --
1107 <http://www.sciencedirect.com/science/article/pii/S0006295203005045>
- 1108 Prokop, A. (2013). A rough guide to *Drosophila* mating schemes. *figshare*,
1109 dx.doi.org/10.6084/m9.figshare.106631 -- <http://dx.doi.org/10.6084/m9.figshare.106631>
- 1110 Prokop, A. (2020). Cytoskeletal organization of axons in vertebrates and invertebrates. *J Cell Biol*
1111 **219**, e201912081 -- <https://doi.org/10.1083/jcb.201912081>
- 1112 Prokop, A. (2021). A common theme for axonopathies? The dependency cycle of local axon
1113 homeostasis. *Cytoskeleton* **78**, 52–63 -- <https://doi.org/10.1002/cm.21657>
- 1114 Prokop, A., Beaven, R., Qu, Y., Sánchez-Soriano, N. (2013). Using fly genetics to dissect the
1115 cytoskeletal machinery of neurons during axonal growth and maintenance. *J. Cell Sci.* **126**,
1116 2331-41 -- <http://dx.doi.org/10.1242/jcs.126912>
- 1117 Prokop, A., Küppers-Munther, B., Sánchez-Soriano, N. (2012). Using primary neuron cultures of
1118 *Drosophila* to analyse neuronal circuit formation and function. *The making and un-making of*
1119 *neuronal circuits in Drosophila* **69**, 225-47 -- http://dx.doi.org/10.1007/978-1-61779-830-6_10
- 1120 Qu, Y., Hahn, I., Lees, M., Parkin, J., Voelzmann, A., Dorey, K., Rathbone, A., Friel, C., Allan, V.,
1121 Okenve Ramos, P., Sánchez-Soriano, N., Prokop, A. (2019). Efa6 protects axons and regulates
1122 their growth and branching by inhibiting microtubule polymerisation at the cortex. *eLife* **8**,
1123 e50319 -- <https://doi.org/10.7554/eLife.50319>
- 1124 Qu, Y., Hahn, I., Webb, S. E. D., Pearce, S. P., Prokop, A. (2017). Periodic actin structures in
1125 neuronal axons are required to maintain microtubules. *Mol Biol Cell* **28** 296-308 --
1126 <https://doi.org/10.1091/mbc.e16-10-0727>

- 1127 Rafiq, N. M., Lyons, L. L., Gowrishankar, S., De Camilli, P., Ferguson, S. M. (2020). JIP3 links
1128 lysosome transport to regulation of multiple components of the axonal cytoskeleton. *bioRxiv*,
1129 2020.06.24.169219 -- <https://doi.org/10.1101/2020.06.24.169219>
- 1130 Randall, T. S., Yip, Y. Y., Wallock-Richards, D. J., Pfisterer, K., Sanger, A., Ficek, W., Steiner, R.
1131 A., Beavil, A. J., Parsons, M., Dodding, M. P. (2017). A small-molecule activator of kinesin-1
1132 drives remodeling of the microtubule network. *Proc Natl Acad Sci U S A* **114**, 13738-43 --
1133 <http://doi.org/10.1073/pnas.1715115115>
- 1134 Ray, K., Perez, S. E., Yang, Z., Xu, J., Ritchings, B. W., Steller, H., Goldstein, L. S. B. (1999).
1135 Kinesin-II is required for axonal transport of choline acetyltransferase in *Drosophila*. *J. Cell Biol.*
1136 **147**, 507-517 --
- 1137 Reck-Peterson, S. L., Redwine, W. B., Vale, R. D., Carter, A. P. (2018). The cytoplasmic dynein
1138 transport machinery and its many cargoes. *Nature Reviews Molecular Cell Biology* **19**, 382-398
1139 -- <https://doi.org/10.1038/s41580-018-0004-3>
- 1140 Rosa-Ferreira, C., Munro, S. (2011). Arl8 and SKIP act together to link lysosomes to kinesin-1. *Dev*
1141 *Cell* **21**, 1171-8 -- <https://doi.org/10.1016/j.devcel.2011.10.007>
- 1142 Rosa-Ferreira, C., Sweeney, S. T., Munro, S. (2018). The small G protein Arl8 contributes to
1143 lysosomal function and long-range axonal transport in *Drosophila*. *Biol Open* **7** --
1144 <https://doi.org/10.1242/bio.035964>
- 1145 Roy, S. (2020). Finding order in slow axonal transport. *Curr Opin Neurobiol* **63**, 87-94 --
1146 <https://doi.org/10.1016/j.conb.2020.03.015>
- 1147 Rozario, A. M., Duwé, S., Elliott, C., Hargreaves, R. B., Moseley, G. W., Dedecker, P., Whelan, D.
1148 R., Bell, T. D. M. (2021). Nanoscale characterization of drug-induced microtubule filament
1149 dysfunction using super-resolution microscopy. *BMC Biology* **19**, 260 --
1150 <https://doi.org/10.1186/s12915-021-01164-4>
- 1151 Russo, G. J., Louie, K., Wellington, A., Macleod, G. T., Hu, F., Panchumarthi, S., Zinsmaier, K. E.
1152 (2009). *Drosophila* Miro is required for both anterograde and retrograde axonal mitochondrial
1153 transport. *J Neurosci* **29**, 5443-55 -- <http://www.ncbi.nlm.nih.gov/pubmed/19403812>
- 1154 Ryder, E., Blows, F., Ashburner, M., Bautista-Llacer, R., Coulson, D., Drummond, J., Webster, J.,
1155 Gubb, D., Gunton, N., Johnson, G., O'Kane, C. J., Huen, D., Sharma, P., Asztalos, Z., Baisch,
1156 H., Schulze, J., Kube, M., Kittlaus, K., Reuter, G., Maroy, P., Szidonya, J., Rasmuson-
1157 Lestander, A., Ekstrom, K., Dickson, B., Hugentobler, C., Stocker, H., Hafen, E., Lepesant, J.
1158 A., Pflugfelder, G., Heisenberg, M., Mechler, B., Serras, F., Corominas, M., Schneuwly, S.,
1159 Preat, T., Roote, J., Russell, S. (2004). The DrosDel collection: a set of P-element insertions for
1160 generating custom chromosomal aberrations in *Drosophila melanogaster*. *Genetics* **167**, 797-
1161 813 -- <https://doi.org/10.1534/genetics.104.026658>
- 1162 Sablin, E. P., Jon Kull, F., Cooke, R., Vale, R. D., Fletterick, R. J. (1996). Crystal structure of the
1163 motor domain of the kinesin-related motor ncd. *Nature* **380**, 555-559 --
1164 <https://doi.org/10.1038/380555a0>
- 1165 Saccon, R. A., Bunton-Stasyshyn, R. K. A., Fisher, E. M. C., Fratta, P. (2013). Is SOD1 loss of
1166 function involved in amyotrophic lateral sclerosis? *Brain* **136**, 2342-2358 --
1167 <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3722346/>
- 1168 Sánchez-Soriano, N., Gonçalves-Pimentel, C., Beaven, R., Haessler, U., Ofner, L., Ballestrem, C.,
1169 Prokop, A. (2010). *Drosophila* growth cones: a genetically tractable platform for the analysis of
1170 axonal growth dynamics. *Dev Neurobiol* **70**, 58-71 -- <https://doi.org/10.1002/dneu.20762>
- 1171 Sandoval, H., Yao, C.-K., Chen, K., Jaiswal, M., Donti, T., Lin, Y. Q., Bayat, V., Xiong, B., Zhang,
1172 K., David, G., Charng, W.-L., Yamamoto, S., Duraine, L., Graham, B. H., Bellen, H. J. (2014).
1173 Mitochondrial fusion but not fission regulates larval growth and synaptic development through
1174 steroid hormone production. *Elife* **3**, e03558 -- <https://doi.org/10.7554/eLife.03558>

- 1175 Saxton, W. M., Hicks, J., Goldstein, L. S., Raff, E. C. (1991). Kinesin heavy chain is essential for
1176 viability and neuromuscular functions in *Drosophila*, but mutants show no defects in mitosis. *Cell*
1177 **64**, 1093-102 -- [https://doi.org/10.1016/0092-8674\(91\)90264-y](https://doi.org/10.1016/0092-8674(91)90264-y)
- 1178 Sheng, Z. H. (2017). The interplay of axonal energy homeostasis and mitochondrial trafficking and
1179 anchoring. *Trends Cell Biol* **27**, 403-416 -- <http://www.ncbi.nlm.nih.gov/pubmed/28228333>
- 1180 Sleight, J. N., Rossor, A. M., Fellows, A. D., Tosolini, A. P., Schiavo, G. (2019). Axonal transport
1181 and neurological disease. *Nat Rev Neurol* **15**, 691-703 -- [https://doi.org/10.1038/s41582-019-](https://doi.org/10.1038/s41582-019-0257-2)
1182 [0257-2](https://doi.org/10.1038/s41582-019-0257-2)
- 1183 Smirnova, E., Griparic, L., Shurland, D. L., van der Bliek, A. M. (2001). Dynamin-related protein
1184 Drp1 is required for mitochondrial division in mammalian cells. *Mol Biol Cell* **12**, 2245-56 --
1185 <http://www.ncbi.nlm.nih.gov/pubmed/11514614>
- 1186 Smith, G. A., Lin, T.-H., Sheehan, A. E., Van der Goes van Naters, W., Neukomm, L. J., Graves,
1187 H. K., Bis-Brewer, D. M., Züchner, S., Freeman, M. R. (2019). Glutathione S-transferase
1188 regulates mitochondrial populations in axons through increased glutathione oxidation. *Neuron*
1189 **103**, 52-65.e6 -- <https://doi.org/10.1016/j.neuron.2019.04.017>
- 1190 Smith, G. M., Gallo, G. (2018). The role of mitochondria in axon development and regeneration.
1191 *Dev Neurobiol* **78**, 221-237 -- <http://www.ncbi.nlm.nih.gov/pubmed/29030922>
- 1192 Song, W., Song, Y., Kincaid, B., Bossy, B., Bossy-Wetzel, E. (2013). Mutant *SOD1*^{G93A} triggers
1193 mitochondrial fragmentation in spinal cord motor neurons: neuroprotection by SIRT3 and PGC-
1194 1α. *Neurobiol Dis* **51**, 72-81 -- <https://doi.org/10.1016/j.nbd.2012.07.004>
- 1195 Spradling, A. C., Stern, D., Beaton, A., Rhem, E. J., Lavery, T., Mozden, N., Misra, S., Rubin, G.
1196 M. (1999). The Berkeley *Drosophila* genome project gene disruption project. Single P-element
1197 insertions mutating 25% of vital *Drosophila* genes. *Genetics* **153**, 135-77 --
1198 <https://doi.org/10.1093/genetics/153.1.135>
- 1199 Stapper, Z. A., Jahn, T. R. (2018). Changes in Glutathione Redox Potential Are Linked to
1200 Aβ₄₂-Induced Neurotoxicity. *Cell Reports* **24**, 1696-1703 --
1201 <https://doi.org/10.1016/j.celrep.2018.07.052>
- 1202 Stowers, R. S., Megeath, L. J., Gorska-Andrzejak, J., Meinertzhagen, I. A., Schwarz, T. L. (2002).
1203 Axonal transport of mitochondria to synapses depends on Milton, a novel *Drosophila* protein.
1204 *Neuron* **36**, 1063-77 -- [https://doi.org/10.1016/s0896-6273\(02\)01094-2](https://doi.org/10.1016/s0896-6273(02)01094-2)
- 1205 Tanaka, K., Sugiura, Y., Ichishita, R., Mihara, K., Oka, T. (2011). KLP6: a newly identified kinesin
1206 that regulates the morphology and transport of mitochondria in neuronal cells. *J Cell Sci* **124**,
1207 2457-2465 -- <http://jcs.biologists.org/content/124/14/2457.abstract>
- 1208 Tang, B. (2018). Miro—working beyond mitochondria and microtubules. *Cells* **7**, 18 --
1209 <http://www.mdpi.com/2073-4409/7/3/18>
- 1210 Tang, Y., Scott, D., Das, U., Gitler, D., Ganguly, A., Roy, S. (2013). Fast vesicle transport is
1211 required for the slow axonal transport of synapsin. *J Neurosci* **33**, 15362-75 --
1212 <http://www.ncbi.nlm.nih.gov/pubmed/24068803>
- 1213 Triclin, S., Inoue, D., Gaillard, J., Htet, Z. M., DeSantis, M. E., Portran, D., Derivery, E., Aumeier,
1214 C., Schaedel, L., John, K., Letierrier, C., Reck-Peterson, S. L., Blanchoin, L., Théry, M. (2021).
1215 Self-repair protects microtubules from destruction by molecular motors. *Nat Mater* --
1216 <https://doi.org/10.1038/s41563-020-00905-0>
- 1217 Tsuda, M., Ootaka, R., Ohkura, C., Kishita, Y., Seong, K.-H., Matsuo, T., Aigaki, T. (2010). Loss of
1218 Trx-2 enhances oxidative stress-dependent phenotypes in *Drosophila*. *FEBS Letters* **584**, 3398-
1219 3401 -- <https://doi.org/10.1016/j.febslet.2010.06.034>
- 1220 Twelvetrees, A. E., Pernigo, S., Sanger, A., Guedes-Dias, P., Schiavo, G., Steiner, R. A., Dodding,
1221 M. P., Holzbaur, E. L. (2016). The dynamic localization of cytoplasmic dynein in neurons is
1222 driven by kinesin-1. *Neuron* **90**, 1000-15 -- <http://www.ncbi.nlm.nih.gov/pubmed/27210554>

- 1223 Uo, T., Dworzak, J., Kinoshita, C., Inman, D. M., Kinoshita, Y., Horner, P. J., Morrison, R. S.
1224 (2009). Drp1 levels constitutively regulate mitochondrial dynamics and cell survival in cortical
1225 neurons. *Exp Neurol* **218**, 274-85 -- <http://www.ncbi.nlm.nih.gov/pubmed/19445933>
- 1226 Vagnoni, A., Hoffmann, P. C., Bullock, S. L. (2016). Reducing Lissencephaly-1 levels augments
1227 mitochondrial transport and has a protective effect in adult *Drosophila* neurons. *J Cell Sci* **129**,
1228 178-190 -- <http://doi.org/10.1242/jcs.179184>
- 1229 van Spronsen, M., Mikhaylova, M., Lipka, J., Schlager, Max A., van den Heuvel, Dave J., Kuijpers,
1230 M., Wulf, Phebe S., Keijzer, N., Demmers, J., Kapitein, Lukas C., Jaarsma, D., Gerritsen,
1231 Hans C., Akhmanova, A., Hoogenraad, Casper C. (2013). TRAK/Milton motor-adaptor proteins
1232 steer mitochondrial trafficking to axons and dendrites. *Neuron* **77**, 485-502 --
1233 <http://dx.doi.org/10.1016/j.neuron.2012.11.027>
- 1234 VanDelinder, V., Adams, P. G., Bachand, G. D. (2016). Mechanical splitting of microtubules into
1235 protofilament bundles by surface-bound kinesin-1. *Sci Rep* **6**, 39408 --
1236 <http://doi.org/10.1038/srep39408>
- 1237 Veeranan-Karmegam, R., Boggupalli, D. P., Liu, G., Gonsalvez, G. B. (2016). A new isoform of
1238 *Drosophila* non-muscle Tropomyosin 1 interacts with Kinesin-1 and functions in oskar mRNA
1239 localization. *J Cell Sci* **129**, 4252-64 -- <https://doi.org/10.1242/jcs.194332>
- 1240 Verhey, K. J., Hammond, J. W. (2009). Traffic control: regulation of kinesin motors. *Nat Rev Molec*
1241 *Cell Biol* **10**, 765-77 -- <https://doi.org/10.1038/nrm2782>
- 1242 Verhey, K. J., Lizotte, D. L., Abramson, T., Barenboim, L., Schnapp, B. J., Rapoport, T. A. (1998).
1243 Light chain-dependent regulation of Kinesin's interaction with microtubules. *J Cell Biol* **143**,
1244 1053-66 -- <http://www.ncbi.nlm.nih.gov/pubmed/9817761>
- 1245 Verstreken, P., Ly, C. V., Venken, K. J., Koh, T. W., Zhou, Y., Bellen, H. J. (2005). Synaptic
1246 mitochondria are critical for mobilization of reserve pool vesicles at *Drosophila* neuromuscular
1247 junctions. *Neuron* **47**, 365-78 -- <https://doi.org/10.1016/j.neuron.2005.06.018>
- 1248 Voelzmann, A., Okenve-Ramos, P., Qu, Y., Chojnowska-Monga, M., del Caño-Espinel, M., Prokop,
1249 A., Sánchez-Soriano, N. (2016). Tau and spectraplakins promote synapse formation and
1250 maintenance through Jun kinase and neuronal trafficking. *eLife* **5**, e14694 --
1251 <http://dx.doi.org/10.7554/eLife.14694>
- 1252 Wali, G., Sutharsan, R., Fan, Y., Stewart, R., Tello Velasquez, J., Sue, C. M., Crane, D. I., Mackay-
1253 Sim, A. (2016). Mechanism of impaired microtubule-dependent peroxisome trafficking and
1254 oxidative stress in SPAST-mutated cells from patients with Hereditary Spastic Paraplegia. *Sci*
1255 *Rep* **6**, 27004 -- <https://doi.org/10.1038/srep27004>
- 1256 Walker, C. L., Pomatto, L. C. D., Tripathi, D. N., Davies, K. J. A. (2018). Redox regulation of
1257 homeostasis and proteostasis in peroxisomes. *Physiol Rev* **98**, 89-115 --
1258 <http://doi.org/10.1152/physrev.00033.2016>
- 1259 Wanders, R. J. A., Vaz, F. M., Waterham, H. R., Ferdinandusse, S. (2020). Fatty Acid Oxidation in
1260 Peroxisomes: Enzymology, Metabolic Crosstalk with Other Organelles and Peroxisomal
1261 Disorders. *Adv Exp Med Biol* **1299**, 55-70 -- <https://www.ncbi.nlm.nih.gov/pubmed/33417207>
- 1262 Wang, C., Du, W., Su, Q. P., Zhu, M., Feng, P., Li, Y., Zhou, Y., Mi, N., Zhu, Y., Jiang, D., Zhang,
1263 S., Zhang, Z., Sun, Y., Yu, L. (2015). Dynamic tubulation of mitochondria drives mitochondrial
1264 network formation. *Cell Res* **25**, 1108-1120 -- <http://dx.doi.org/10.1038/cr.2015.89>
- 1265 Wilson, C., Terman, J. R., Gonzalez-Billault, C., Ahmed, G. (2016). Actin filaments - a target for
1266 redox regulation. *Cytoskeleton (Hoboken)* **73**, 577-95 --
1267 <http://www.ncbi.nlm.nih.gov/pubmed/27309342>
- 1268 Winding, M., Kelliher, M. T., Lu, W., Wildonger, J., Gelfand, V. I. (2016). Role of kinesin-1-based
1269 microtubule sliding in *Drosophila* nervous system development. *Proc Natl Acad Sci U S A* **113**,
1270 E4985-94 -- <http://www.ncbi.nlm.nih.gov/pubmed/27512046>

- p>1271 Wioland, H., Frémont, S., Guichard, B., Echard, A., Jégou, A., Romet-Lemonne, G. (2021). Actin
-
- p>1272 filament oxidation by MICAL1 suppresses protections from cofilin-induced disassembly.
- EMBO*
-
- p>1273
- reports*
- 22**
- , e50965 --
- <https://doi.org/10.15252/embr.202050965>
-
- p>1274 Witte, L., Linnemannstöns, K., Schmidt, K., Honemann-Capito, M., Grawe, F., Wodarz, A., Gross,
-
- p>1275 J. C. (2020). The kinesin motor Klp98A mediates apical to basal Wg transport.
- Development*
-
- p>1276
- 147**
-
- <https://doi.org/10.1242/dev.186833>
-
- p>1277 Wong, H.-S., Dighe, P. A., Mezera, V., Monternier, P.-A., Brand, M. D. (2017). Production of
-
- p>1278 superoxide and hydrogen peroxide from specific mitochondrial sites under different bioenergetic
-
- p>1279 conditions.
- J Biol Chem*
-
-
- p>1280
- <http://www.jbc.org/content/early/2017/08/31/jbc.R117.789271.abstract>
-
- p>1281 Wong, Y. L., Rice, S. E. (2010). Kinesin's light chains inhibit the head- and microtubule-binding
-
- p>1282 activity of its tail.
- Proc Natl Acad Sci*
- 107**
- , 11781-6 --
- <https://doi.org/10.1073/pnas.1005854107>
-
- p>1283 Wozniak, M. J., Melzer, M., Dorner, C., Haring, H. U., Lammers, R. (2005). The novel protein KBP
-
- p>1284 regulates mitochondria localization by interaction with a kinesin-like protein.
- BMC Cell Biol*
- 6**
- , 35
-
- p>1285 --
- <http://www.ncbi.nlm.nih.gov/pubmed/16225668>
-
- p>1286 Yalcin, B., Zhao, L., Stofanko, M., O'Sullivan, N. C., Kang, Z. H., Roost, A., Thomas, M. R.,
-
- p>1287 Zaessinger, S., Blard, O., Patto, A. L., Sohail, A., Baena, V., Terasaki, M., O'Kane, C. J. (2017).
-
- p>1288 Modeling of axonal endoplasmic reticulum network by spastic paraplegia proteins.
- Elife*
- 6**
- ,
-
- p>1289 e23882 --
- <https://doi.org/10.7554/eLife.23882>
-
- p>1290 Yip, Y. Y., Pernigo, S., Sanger, A., Xu, M., Parsons, M., Steiner, R. A., Dodding, M. P. (2016). The
-
- p>1291 light chains of kinesin-1 are autoinhibited.
- Proc Natl Acad Sci U S A*
- 113**
- , 2418-23 --
-
- p>1292
- <https://www.ncbi.nlm.nih.gov/pubmed/26884162>
-
- p>1293 Yu, W., Sun, Y., Guo, S., Lu, B. (2011). The PINK1/Parkin pathway regulates mitochondrial
-
- p>1294 dynamics and function in mammalian hippocampal and dopaminergic neurons.
- Hum Mol Genet*
-
- p>1295
- 20**
- , 3227-40 --
- <http://www.ncbi.nlm.nih.gov/pubmed/21613270>
-
- p>1296 Zahavi, E. E., Hummel, J. J. A., Han, Y., Bar, C., Stucchi, R., Altelaar, M., Hoogenraad, C. C.
-
- p>1297 (2021). Combined kinesin-1 and kinesin-3 activity drives axonal trafficking of TrkB receptors in
-
- p>1298 Rab6 carriers.
- Dev Cell*
-
-
- p>1299
- <https://www.sciencedirect.com/science/article/pii/S153458072100068X>
-
- p>1300 Zala, D., Hinckelmann, M.-V., Yu, H., da Cunha, L., Menezes, M., Liot, G., Cordelières, Fabrice P.,
-
- p>1301 Marco, S., Saudou, F. (2013). Vesicular glycolysis provides on-board energy for fast axonal
-
- p>1302 transport.
- Cell*
- 152**
- , 479-491 --
- <http://dx.doi.org/10.1016/j.cell.2012.12.029>
-
- p>1303 Zelko, I. N., Mariani, T. J., Folz, R. J. (2002). Superoxide dismutase multigene family: a
-
- p>1304 comparison of the CuZn-SOD (SOD1), Mn-SOD (SOD2), and EC-SOD (SOD3) gene structures,
-
- p>1305 evolution, and expression.
- Free Radic Biol Med*
- 33**
- , 337-49 --
- [https://doi.org/10.1016/s0891-](https://doi.org/10.1016/s0891-5849(02)00905-x)
-
- p>1306
- [5849\(02\)00905-x](https://doi.org/10.1016/s0891-5849(02)00905-x)
-
- p>1307 Zhou, Z., Kang, Y. J. (2000). Cellular and subcellular localization of catalase in the heart of
-
- p>1308 transgenic mice.
- J Histochem Cytochem*
- 48**
- , 585-94 --
-
- p>1309
- <https://doi.org/10.1177/002215540004800502>
-
- p>1310 Zorov, D. B., Juhaszova, M., Sollott, S. J. (2014). Mitochondrial reactive oxygen species (ROS)
-
- p>1311 and ROS-induced ROS release.
- Phys Rev*
- 94**
- , 909-950 --
-
- p>1312
- <https://journals.physiology.org/doi/abs/10.1152/physrev.00026.2013>
-
- p>1313
-
- p>1314

Figures

Fig.1 Deficiencies of three motor proteins cause MT curling. **A-G)** Examples of neurons of different genotype (indicated top right) and stained for tubulin at 5DIV; asterisks indicate cell bodies, arrow heads axon tips, curved arrows areas of MT curling, white rectangles shown as twofold magnified, yellow emboxed insets; scale bar in A represents 20µm in all images. **H)** Quantification of MT curling phenotypes measured as MT disorganisation index (MDI) and normalised to wild-type controls (red stippled line); mean ± SEM is indicated in blue, numbers of analysed neurons in orange, results of Mann Whitney rank sum tests are shown in grey/black.

Fig.2 Impacts of motor protein and linker mutations on numbers of axonal mitochondria and synaptic spots. **A-N)** Examples of neurons of different genotype (indicated top right) and stained at 5DIV for tubulin (tub, magenta) and either Synaptotagmin (Syt, green in A-G) or with mitoTracker (green in H-N); scale bar in A represents 20µm in all images. **O,P)** Quantification of axonal numbers of Syt-positive spots (O) or mitochondria (P), all normalised to wild-type controls (red stippled line); medians are indicated in blue, numbers of analysed neurons in orange, results of Mann Whitney rank sum tests are shown in grey/black.

Fig.S1 Endoplasmic reticulum accumulates at axon tips upon loss of Khc. **A,B)** Primary neurons at 5 DIV carrying the genomically tagged *Rtln1-YFP* allele labelling endoplasmic reticulum (ER; del Castillo et al., 2019; O'Sullivan et al., 2012), either in wild-type (wt; A) or *Khc*⁸ mutant background (B); inset with blue outline in A displays the green channel of the neuron (reduced to 50% in size) to illustrate the continuous nature of *Rtln1::GFP*-labelled ER throughout its neurites; the yellow emboxed area in B is shown as twofold increased inset of the green channel to illustrate the netlike organisation of ER visible in axonal swellings. Asterisks indicate cell bodies and arrow heads axon tips (note that there are two neurons in B), white/orange chevrons point at strong/weak axonal tip accumulations of ER. Accumulations might indicate an imbalance of antero- and retrograde organelle movement potentially caused by loss of Khc-dependent Dynein transport to axon tips (Moughamian et al., 2013; Twelvetrees et al., 2016) expected to reduce the retrograde drift of ER. The scale bar in A represents 20 µm in A and B. **C)** Quantification of axonal tip accumulation of ER: numbers of neurons analysed are shown in orange, numbers in bars the rounded percentages of neurons with no/weak/strong accumulations, the number above bars show the P value of the X² test.

Fig.3 Assessing contributions of Khc subfunctions to MT regulation. **A)** Schematic representation of *Drosophila* Khc drawn to scale. Domains are colour-coded and start/end residues are indicated by numbers: motor domain (red; according to Sablin et al., 1996), coiled-coil domains required for homo- and/or heterodimerisation (green; as predicted by Ncoils in ensembl.org), the C-terminal ATP-independent MT-binding motif (blue; according to Winding et al., 2016), and the C-terminal auto-inactivation domain (dark grey; according to Kaan et al., 2011); grey lines above the protein scheme indicate the three expression constructs used in this study; below the protein scheme further details are shown: the sequence of the C-terminal MT-binding domain (*mutA* mutations indicated in orange; Winding et al., 2016), the sequence of the auto-inactivation domain (indicating the IAK motif and R947E mutation; Kelliher et al., 2018), the binding areas (darker green coiled-coils) of Klc (according to Veeranan-Karmegam et al., 2016), Mlt (known to overlap with Klc; Glater et al., 2006; Verhey et al., 1998) and Tropomyosin 1 (Dimitrova-Paternoga et al., 2021), and the two-fold enlarged motor

domain. The secondary structure of the motor domain is indicated below (α helices in black, β sheets in blue, loops/L in red); this map was generated by matching the resolved structure of Khc (UniProt code: P17210, PDB id 2y65) with descriptions of the kinesin consensus (Sablin et al., 1996); regions/motifs that bind ADP/ATP (nt, orange; according to Cao et al., 2017; Gigant et al., 2013; Sablin et al., 1996) and/or MTs (dark red; according to Hunter and Allingham, 2020) are also indicated below; N1-4 in the motor domain indicate highly conserved motifs (according to Sablin et al., 1996); abbreviations above the motor domain indicate the locations of the cover strand (CS; according to Budaitis et al., 2021), P-loops (PL) and switch domains I and II (SI, SII; according to Cao et al., 2017; Gigant et al., 2013; Sablin et al., 1996). The N-terminal deletion of the above *Khc*(82-711) construct is shown in pink: it does not affect MT-binding sites, but it removes the cover strand (known to affect kinesin's MT affinity and processivity; Budaitis et al., 2021) and the first P-loop (with potential impact on the ATP/ADP cycle); it might also affect the behaviour of the second P-loop which was shown to accelerate Khc movement when harbouring the T94S mutation (Cao et al., 2017; Higuchi et al., 2004). **B**) Schematic representation of some sub-functions of Khc (details and abbreviations in main text; red and stippled black lines indicate processive transport; for further sub-functions see Discussion): via a C-terminal MT-binding domain Khc can slide MTs (i), associating with Pat1 (and potentially Klc) it is expected to transport non-vesicular cargoes including mRNA (ii), with Milt and Miro organelle transport (iii), and with a protein complex containing Klc and Syd vesicular transport (iv); in the absence of such associations Khc is auto-inhibited and detaches from MTs assisted by Klc (v); to interfere with these subfunctions in this study, different genes were genetically removed (orange crosses) or specific *Khc* mutant alleles used (italic orange text). **C**) Quantified effects on MT curling caused by specific mutations affecting Khc sub-functions (numbers in grey circles indicate which function in A is affected): MT curling is quantified as MT disorganisation index (MDI) normalised to wild-type controls (red stippled line); bars at bottom indicate type of culture ('5 DIV', embryonic neurons 5 days *in vitro*; 'L3 1/2 DIV', late larval neurons 1/2 days *in vitro*; '5d pre 1 DIV', embryonic neurons pre-cultured for 5 days and cultured for 1 day); mean $\pm$ SEM is indicated in blue, numbers of analysed neurons in orange, results of Mann Whitney rank sum tests are shown in grey/black.

Fig.S2 Validation of Khc's MT phenotype and demonstration of maternal contribution. **A-C**) Quantification of MT curling phenotypes measured as MT disorganisation index (MDI) and normalised to wild-type controls (red stippled line); mean $\pm$ SEM is indicated in blue, numbers of analysed neurons in orange, results of Mann Whitney rank sum tests are shown in grey/black; A) shows data for *Khc*²⁷ in homozygosis (27/27) or over deficiency (27/Df), for Khc knock-down (*elav>Khc-IR*) and wild-type (wt) and driver line (*elav*) controls; B) shows data for *Khc*⁸ over deficiency (8/Df) and wild-type controls at different culture times (HIV, hours *in vitro*; DIV, days *in vitro*); C) shows data for *Klc*^{B/Df} and *Klc*^{1ts} at 1DIV following 5d pre-culture; note that *Klc*^{1ts} is a temperature-sensitive allele (see methods) and was pre-cultured at 26°C and cultured at 29°C. **D,E**) Examples of neurons at different times in culture (D; relating to data in B) and after pre-culture (E; relating to C); asterisks indicate cell bodies, arrow heads axon tips, curved arrows areas of MT curling; scale bar in D represents 20 μ m in D and E.

Fig.4 ROS enhancing manipulations cause MT curling phenotypes. **A**) Scheme illustrating the complexity of ROS-regulating systems in *Drosophila*; ROS-generating factors (bold green): two cytoplasmic NADPH oxidases (Nox/NADPH Oxidase, Duox/Dual oxidase with its essential maturation factor Mol/Moladietz; Khan et al., 2017); enzymes of the mitochondrial EMT/electron transport chain (Wong et al., 2017; Zorov et al., 2014); peroxisomal ACOX1/acyl-CoA oxidase 1

(Walker et al., 2018); Xanthine/aldehyde oxidases (Rosy, AOX1, AOX2, AOX3, AOX4; all jointly silenced by loss of Mal/Maroon-like sulfurtransferase; Marelja et al., 2014); ROS removal mechanisms (red): superoxide dismutases turn superoxide ($O_2^{\bullet-}$) into H_2O_2 (cytoplasmic CuZn-dependent Sod1, mitochondrial Mn-dependent Sod2, extracellular Sod3); H_2O_2 is scavenged by peroxisomal Cat/Catalase and neuronal peroxiredoxins (Jafrac1, Prx5; Cao and Lindsay, 2017; Orr et al., 2013; Smith et al., 2019; Stapper and Jahn, 2018) and the GSH transferase Gzfz (GST-containing FLYWCH zinc-finger protein; Smith et al., 2019; Stapper and Jahn, 2018); the latter three depend on the redox cycle of the Glu-Cys-Gly tripeptide GSH/Glutathione, synthesised by glutathione synthetases (Gss1, Gss2) and Gclc/Glutamate-cysteine ligase (Smith et al., 2019; Stapper and Jahn, 2018) and regenerated via Thioredoxins (primarily Trx-2 in neurons; Orr et al., 2013; Tsuda et al., 2010) and Thioredoxin reductases (primarily TrxR-1 in neurons; Orr et al., 2013; Smith et al., 2019); pharmacological agents (black italics): DEM/diethyl maleate blocks the GSH system (Pompella et al., 2003); agents/factors used in our study are highlighted in yellow. **B-E**) Examples of neurons, either wild-type (wt) expressing Sod1 or Duox (driven by *elav-Gal4*) or homozygous for *Ca²⁺*, all cultured for 1DIV and stained for actin (act, magenta) and tubulin (tub; green); asterisks indicate cell bodies, arrow heads axon tips, curved arrows areas of MT curling; yellow emboxed areas are shown as 1.5-fold enlarged insets (green channel only); scale bar in B represents 20µm in B-E. **F**) Quantification of MT curling phenotypes measured as MT disorganisation index (MDI) and normalised to wild-type controls (red stippled line); bars at bottom indicate type of culture ('1/3/5 DIV/HIV', embryonic neurons 1/3/5 days/hours *in vitro*); mean ± SEM is indicated in blue, numbers of analysed neurons in orange, results of Mann Whitney rank sum tests are shown in grey/black.

Fig.S4 MT bundle and axon length phenotypes of *pex3²* mutant neurons. Quantification of phenotypes of wild-type (wt) and *Pex³* homozygous mutant neurons: **A**) MT curling phenotypes measured as MT disorganisation index (MDI); **B**) axon length; both measured are normalised to wild-type controls (red stippled line); mean ± SEM is indicated in blue, numbers of analysed neurons in orange, results of Mann Whitney rank sum tests are shown in grey/black.

Fig.5 Ameliorating effects of Trolox on mutant MT curling phenotypes. Quantification of MT curling phenotypes measured as MT disorganisation index (MDI) and normalised to wild-type controls (red stippled line); neurons of different genotype (indicated below) were cultured for 1day after preculture (A) of for 5 days (B) in the presence of vehicle (blue) or 100µm Trolox (green; molecule depicted in A); mean ± SEM is indicated in blue/green, numbers of analysed neurons in orange, results of Mann Whitney rank sum tests are shown in grey/black.

Fig.S5 Impaired fission/fusion of mitochondria does not affect MT bundles. **A-D**) Neurons at 5 days *in vitro* (DIV) and stained with anti-tubulin (tub, magenta) and mitoTracker (green); they are wild-type (wt) or homozygous mutant for the mitochondrial fission factor Drp or the mitochondrial fusion factors Marf or Opa, as indicated; asterisks indicate cell bodies, arrow heads axon tips, yellow emboxed areas are shown as 2-fold enlarged insets (green channel only), and the scale bar in A represents 20µm in A-D; note that mitochondria tend to appear as dashed lines in controls (A), as a continuous string of pearls excluded from side branches upon loss of fission (B), and as sparse dots upon loss of fusion (C,D). **E**) Quantification of MT curling phenotypes from experiments shown in A-D, measured as MT disorganisation index (MDI) and normalised to wild-type controls (red stippled line). **F**) Similar experiments with the same mutations using 5 day pre-culture and culture for 1 day. In E

and F, mean $\pm$ SEM is indicated in blue, numbers of analysed neurons in orange, results of Mann Whitney rank sum tests are shown in grey/black.

Fig.6 Genetic studies of functional links between Khc, Milt and Klc. **A-D)** Examples of neurons, either wild-type (wt) or homozygous for *Khc*, *milt* or *Klc* null mutant alleles, cultured for 1DIV following 5d pre-culture (to deplete maternal product) and stained against tubulin (tub); asterisks indicate cell bodies, arrow heads axon tips, curved arrows areas of MT curling; scale bar in D represents 20 μ m in A-D. **E-G)** Quantification of MT curling phenotypes measured as MT disorganisation index (MDI) and normalised to wild-type controls (red stippled line) shown for precultured neurons (E; as in A-D), single/double-homozygous mutant neurons (F) and upon genetic interaction (G; single heterozygous and trans-heterozygous); mean $\pm$ SEM is indicated in blue, numbers of analysed neurons in orange, results of Mann Whitney rank sum tests are shown in grey/black.

Fig.S3 Phenotypes upon *Gapdh1* knock-down in primary neurons at 5 DIV. **A)** Illustration of the NADH- and ATP-generating steps of glycolysis; names of proteins are shown in bold, other molecules in italics: GAPDH (glyceraldehyde-3-phosphate dehydrogenase), PGK (phosphoglycerate kinase), GAP (glyceraldehyde-3-phosphate), 1,3 BPG (1,3-biphosphoglycerate), 3-PGA (3-phosphoglycerate). **B)** GAPDH and PGK are present on transported vesicles together with other factors relevant for glycolysis (Hinckelmann et al., 2016; Zala et al., 2013) providing ATP to drive kinesin-mediated processive transport (red and stippled black lines). **C)** In the absence of *Gapdh1*, the transport of synaptic vesicles but not mitochondria is impaired (assessed via anti-Syt and mitoTracker staining; see Fig.2), as is consistent with *in vivo* observations in *Drosophila* (larval motor nerves; Zala et al., 2013). **D)** Absence of GAPDH does not cause MT curling. Quantification of MT curling phenotypes in C and D is measured as MT disorganisation index (MDI) and normalised to wild-type controls (red stippled line); median in C and mean $\pm$ SEM in D are indicated in blue, numbers of analysed neurons in orange, results of Mann Whitney rank sum tests are shown in grey/black.

Fig.7 Impacts of activated Khc on MT curling. **A-C)** Examples of neurons at 5DIV expressing different Khc constructs (indicated bottom left; compare Fig.3A) and stained for tubulin (tub, magenta) and GFP (green), also shown as greyscale single channel images on the right; asterisks indicate cell bodies, arrow heads axon tips, curved arrows areas of MT curling; scale bar in A represents 20 μ m in all images. **D)** Quantification of MT curling phenotypes measured as MT disorganisation index (MDI) and normalised to wild-type controls (red stippled line); genotypes are shown below, also indicating the culture period (5DIV, 3DIV); mean $\pm$ SEM is indicated in blue, numbers of analysed neurons in orange, results of Mann Whitney rank sum tests are shown in grey/black.

Fig.8 Mapping findings on the dependency cycle of local axon homeostasis. The numbered green arrows and red T-bar make up the previously published 'dependency cycle of local axon homeostasis' (Prokop, 2021): 1) axonal transport provides materials, components and organelles required for axon function; 2) this transport requires MT bundles as the essential highways; 3) however, this live-sustaining transport damages MT bundles; 4) the axonal cortex and MT binding proteins (MTBPs) support and maintain MT bundles (emboxed names in orange and grey at bottom left list factors that were shown in the *Drosophila* neuron culture system to be involved in bundle-maintaining cortical and MT regulation; Alves-Silva et al., 2012; Hahn et al., 2021; Qu et al., 2019;

Qu et al., 2017); 5) bundle maintenance requires transport-dependent components and physiology, thus closing the circle. The original model of 'local axon homeostasis' comprised arrows 1-4 (Hahn et al., 2019). Khc contributes to the MT bundle damage, and this is enhanced by non-inactivating mutations (vibrating red arrow and *Khc* alleles top left). Loss of function of Khc, Milt, Miro, Unc-104 and Dhc contribute to mitochondrial transport (large green arrow, top right). Mitochondria harbour Sod2 that can quench harmful ROS (green T-bar); 8 independent pharmacological and genetic manipulation of ROS regulation (bottom right) demonstrated that dysregulation of ROS causes MT curling (dashed red arrow). Examples of mammalian factors that can be mapped onto this cycle are explained in the Discussion and previous reviews (Hahn et al., 2019; Prokop, 2021).

Fig.S6 *Pat1* and *Khc*^{mutA} mutations fail to suppress the *Klc*-deficient MT curling phenotype. Quantification of MT curling phenotypes measured as MT disorganisation index (MDI) and normalised to wild-type controls (red stippled line); genotypes are shown below, also indicating the culture period (2DIV, 5DIV); mean ± SEM is indicated in blue, numbers of analysed neurons in orange, results of Mann Whitney rank sum tests are shown in grey/black. The fact that *Khc*^{mutA} and *Pat1*^{robin} fail to suppress *Klc*^{8ex94}-induced MT curling suggests that potential surplus pools of non-inactivated Khc do not engage in MT sliding or Pat1-mediated transport to cause MT curling.

Fig. 1 Liew et al.

Fig. 2 Liew et al.

Fig. 4 Liew et al.

Fig. 5 Liew et al.

Fig. 6 Liew et al.

Fig. 7 Liew et al.

Fig. S1 Liew et al.

Fig. S2 Liew et al.

Fig. S3 Liew et al.

A **B**

Fig. S4 Liew et al.

Fig. S5 Liew et al.

Double-mutant constellations with *Klc*^{8ex94}

Fig. S6 Liew et al.