Spatiotemporal transcriptomics reveals pathogenesis of viral myocarditis

Authors: Madhav Mantri¹, Meleana M. Hinchman², David W. McKellar¹, Michael F. Z. Wang¹, Shaun T. Cross^{1,2,3}, John S. L. Parker^{2,3*}, Iwijn De Vlaminc^{1,3*}

Affiliations: ¹Nancy E. and Peter C. Meinig School of Biomedical Engineering, Cornell University, Ithaca, New York, ²Baker Institute for Animal Health, College of Veterinary Medicine, Cornell University, Ithaca, New York, ³Cornell Institute for Host-Microbe Interactions and Disease, Cornell University, Ithaca, New York.

*To whom correspondence should be addressed: id93@cornell.edu and jsp7@cornell.edu

ABSTRACT

A significant fraction of sudden death in children and young adults is due to viral myocarditis, an inflammatory disease of the heart. Here we used integrated single-cell and spatial transcriptomics to study the temporal, spatial, and cellular heterogeneity of host-virus interactions associated with reovirus-induced myocarditis in neonatal murine hearts. We further assayed the intestine, the primary site of reovirus infection to establish the cellular targets of the virus and the chronology of molecular events that ultimately lead to myocarditis. Our data give insight into the cell-type specificity of innate immune responses, and into the transcriptional states of inflamed cardiac cells that recruit circulating immune cells, including cytotoxic T cells which induce pyroptosis. Analyses of spatially restricted gene expression in myocarditic regions and the border zone identified immune-mediated cell-type specific injury and stress responses. Overall, we observe a complex network of cellular phenotypes and cell-cell interactions associated with viral myocarditis.

INTRODUCTION

Viral infection is the most common cause of myocarditis^{1,2}. The resulting inflammatory cardiomyopathy can lead to arrhythmias, dilated cardiomyopathy, and death^{1,3,4}. In humans, viral myocarditis is challenging to study because of the low sensitivity of available diagnostic testing, the acute onset of the disease, the focal nature of the disease, and the extreme heterogeneity of immune-virus interactions in complex cardiac tissues⁴⁻⁶. In mice, mammalian orthoreovirus offers a flexible model system⁷. After oral inoculation, the Type 1 Lang (T1L) reovirus strain initially infects the gastrointestinal tract. Within days the infection then spreads to secondary sites in the body, including the heart, leading to myocarditis in up to 50% of infections⁷⁻⁹. Yet, even in this mouse model, the molecular pathogenesis of viral myocarditis is difficult to study because of the complex network of cardiac and immune cell types involved and the cellular, spatial, and temporal heterogeneity of the disease^{2,10}. Consequently, neither the cell types that are responsible for the innate immune response, nor the cell types that are infected *in vivo* have been identified. Similarly, the responses of infected and uninfected bystander cells within the heart have not been characterized. In addition, the protective versus damaging effects of adaptive immune responses have not been quantified. Experiments in mice with severe combined immunodeficiency (SCID) indicate that adaptive immune responses are not required for myocardial injury and heart failure^{7,11}, but these observations do not exclude the possibility that immune-cell-mediated injury is important in immunocompetent mice. Unbiased characterization of all cellular phenotypes as a function of time and location within infected cardiac tissues is needed to address these knowledge gaps.

Here we used integrated single-cell and spatially resolved RNA-sequencing (RNA-seq) to study the cellular and spatial heterogeneity of myocarditic processes in the hearts of reovirus-infected neonatal mice at multiple time points after infection. We also applied these technologies to study the innate response to reovirus infection in the intestine. In addition, we performed time-series single-cell RNA-seq (scRNA-seq) of cardiac tissues of mice infected with a reovirus point mutant that does not cause myocarditis. To establish viral tropism, we implemented molecular enrichment of non-polyadenylated viral transcripts that were otherwise poorly represented in the transcriptomes. Our measurements give insight into the cell-type specificity of innate immune responses, into the tropism of the virus in the intestine and the heart, and into the transcriptional states of cell types involved in the production of inflammatory cytokines and the recruitment of circulating immune cells. Analyses of spatially restricted gene expression in myocarditic regions and the border zone around those regions identified injury and stress responses in different cell types, including cardiomyocytes. Overall, our data identify spatially restricted cellular interactions and cell-type specific host responses during reovirus-induced myocarditis.

61

62 RESULTS

63 Single-cell and spatial transcriptomics of reovirus T1L-infected neonatal mice hearts

64 To elucidate the pathogenesis of reovirus-induced myocarditis, we analyzed heart tissues
 65 collected from neonatal mice infected orally with either the T1L strain of reovirus or a mock
 66 control (**Methods, Fig. 1A**). We generated scRNA-seq data for 31,684 cells from infected
 67 hearts and mock controls at 4, 7, and 10 days post-infection (dpi), and 8,243 spatial
 68 transcriptomes for four tissue sections from infected hearts and mock controls at 4 and 7 dpi
 69 from the same litter (10x Chromium and 10x Visium, **Methods, Supp Fig. 1A-1B and Fig. 1B-
 70 1C**). The single-cell transcriptomes represented 18 distinct cell types, including
 71 cardiomyocytes, endocardial cells, cardiac fibroblasts, endothelial cells, mural cells,
 72 macrophages, neutrophils, NK cells, dendritic cells, T cells, and B cells (**Methods, Fig 1B,
 73 Supp Fig 1C-E**). Clustering of the spatial transcriptomic data revealed distinct transcriptional
 74 programs for myocarditic regions and the border zone surrounding the myocarditic regions in
 75 the 7 dpi reovirus-infected heart that corresponded to areas of tissue damage identified by
 76 H&E staining (**Fig. 1C**). The combination of scRNA-seq and spatial transcriptomics allowed us
 77 to resolve and visualize cell types and gene expression in a spatial context (**Methods, Supp
 78 Fig 1F**). Because the virus first infects the gastrointestinal tract before it spreads to other body
 79 sites including the heart, we also performed scRNA-seq and spatial transcriptomics on ileum.
 80 We obtained 7,695 single-cell transcriptomes and 8,027 spatial spot transcriptomes for ileum
 81 from mock and infected samples at 1 and 4 dpi (**Fig. 1D, Supp Fig. 2A-C**).

82 To faithfully identify reovirus transcripts in the ileum and heart, which are not polyadenylated,
 83 we performed hybridization-based enrichment of viral fragments captured in the scRNA-seq
 84 libraries (**Methods**). In the ileum, we captured a total of 13,100 unique viral transcripts, with
 85 viral load decreasing from 1 dpi to 4 dpi (**Supp Fig. 2D-2E**). At 1 dpi, entero-endocrine cells
 86 had the highest fraction of infected cells followed by enterocytes and goblet cells, all of which
 87 are present in the gut epithelium. Lymphatic endothelial cells were infected at 4 dpi, suggesting
 88 that the virus reaches the bloodstream via lymphatic drainage to allow transmission of the virus
 89 to secondary sites in the body, including the heart, as shown before¹² (**Supp Fig. 2E**). We
 90 captured 2,762 unique viral transcripts from 392 cells in the T1L-infected hearts. The viral load
 91 first increased from 4 dpi to 7 dpi and then decreased from 7 dpi to 10 dpi, consistent with viral
 92 titer assays performed on whole hearts^{9,13} (**Fig. 1E, Supp Fig. 2F-2G**). Endocardial and
 93 endothelial cells were the most frequently infected cell types at 4 dpi, suggesting that
 94 endocardial cells lining the ventricular lumen and endothelial cells lining the cardiac
 95 vasculature are among the first cells to be infected in the heart (**Fig. 1E**). We detected viral
 96 transcripts in neutrophils, dendritic cells, and T cells in the 7 dpi heart (**Fig. 1E**). This
 97 observation suggests that antigen-presenting cells and immune cells may contribute to the
 98 spread of infection to other organs in the body. The role of infected dendritic cells in bringing
 99 more reovirus to the cardiac tissue during systemic infection has been discussed previously⁸.

Figure 1: Single-cell and spatial transcriptomics of cardiac and ileum tissue of reovirus-infected neonatal mice. **A)** Experiment and analysis workflow. **B)** UMAP plot of 31,684 single-cell transcriptomes from mock-infected and reovirus-infected hearts at 4, 7, and 10 dpi, clustered by gene expression and colored by cell type (left). UMAP plots showing the gaussian kernel density of cells across samples for the heart scRNA-seq data (right). **C)** 8,243 spatial transcriptomes of cardiac tissue sections from mock-infected and reovirus-infected mice at 4 and 7 dpi. Haematoxylin and Eosin (H&E) stained image of reovirus-infected myocarditic tissue section used for spatial transcriptomics at 7 dpi (in box). **D)** UMAP plot of 7,695 single-cell transcriptomes from mock-infected and reovirus-infected ileum at day 1 and 4 dpi, clustered and colored by cell type (left). UMAP plots showing the gaussian kernel density of cells across samples for the ileum scRNA-seq data (right). **E)** Dot plot showing the percent of cells with non-zero viral transcripts and the mean viral transcript counts (UMIs) in ileal and cardiac cell types.

Endothelial cells are primed with basal interferon response and play an important role in initiating host innate immune response

To detect early transcriptional differences in the cardiac tissue after infection, we performed Differential Gene Expression Analysis (DGEA, mock vs infected hearts at 4 dpi, **Methods**). This analysis revealed a significant upregulation of 226 genes in the infected heart (two-sided Wilcoxon test, log fold-change > 1.0 and p-value < 0.01), including genes related to the interferon- β pathway, interferon signaling, and innate immune responses (**Supp Fig 3A-3B, Fig. 2A**).

To quantify and compare the overall magnitude of early infection responses across different cell types, we computed a gene module score (infection response score, IR, module of 226 genes selected above). Comparison of the IR of different cell types in absence of infection revealed a small, but higher IR in endothelial cells as compared to other cardiac cell types (**Fig. 2B**). In response to infection, an increase in IR was observed for all cardiac cell types, but the greatest increase in IR was observed for endothelial cells (**Fig. 2B**). These data suggest that endothelial cells lining the cardiac vasculature are important initiators of the host defense to viral infection. Comparison of IR scores using the spatial transcriptomic data shows increased IR scores in the infected hearts at 4 and 7 dpi with the highest scores found in myocarditic regions (**Fig. 2C**). Given our observation that endothelial cells within the heart had the highest IR score in absence of infection, we asked if this observation was unique to heart tissue or was a more general phenomenon. To this end, we used the Tabula Muris scRNA-seq mouse atlas¹⁴ and estimated the IR of ~16,000 cells of five major cell types (epithelial cells, fibroblasts, endothelial cells, smooth muscle cells, and mesenchymal cells) across 10 different organs and tissues. This analysis revealed that endothelial cells consistently had the highest IR score across all tissues in mice (**Fig. 2D**). These results indicate that endothelial cells lining the vasculature have a higher basal expression of innate response genes within most tissues, which may prime these cells to respond to viral dissemination within the blood and lymphatics.

To investigate the cell-type-specific IR in the ileum, the primary site of reovirus infection, we performed DGEA on reovirus-infected at 1 dpi compared to mock-infected ileal cells and found a significant upregulation of 438 genes (two-sided Wilcoxon test, log fold-change > 1.0 and p-value < 0.01), related to the interferon-beta pathway, interferon signaling, and innate immune responses (**Supp Fig. 3C-3D**). We computed an IR score using this module of 438 genes and observed higher basal IR scores in enterocytes and entero-endocrine cells as compared to other ileal cell types (**Fig. 2E**). Enterocytes further showed the highest increase in IR score after infection, followed by entero-endocrine, endothelial, and lymphatic cells (**Fig. 2E**). Comparison of IR scores for spatial transcriptomic data further supported our analysis of the scRNA-seq data, showing increased IR scores in the infected ileum at 1 and 4 dpi with the highest scores evident within intestinal mucosa and villi (**Fig. 2F**). The intestinal epithelial cells must tolerate commensal microorganisms present in the lumen of the gut and yet still be responsive to invasive pathogens. Our data suggest that to achieve this, enterocytes and entero-endocrine cells in the gut epithelium are primed with a basal interferon response and play an important part in mounting innate immune responses in the early stages of viral infection.

Figure 2: Endothelial cells have the highest basal interferon response and the highest increase in innate response upon reovirus infection **A)** Heatmap showing the expression of the 25 most upregulated genes in the reovirus-infected heart as compared to mock at 4 dpi. **B)** Infection response score for cardiac cell types in scRNA-seq data across mock-infected and reovirus-infected hearts at three distinct stages. The infection response score represents the gene module score for a panel of 226 genes that are significantly upregulated (two-sided Wilcoxon test, log fold-change > 1.0 and p-value < 0.01) in the reovirus-infected heart as compared to the mock-infected heart at 4 dpi. **C)** Infection response score (defined above) across spots in spatial transcriptomics data. **D)** Infection response score calculated for five common cell types across 13 tissues from the tabula-muris mouse atlas data. **E)** Infection response score for ileal cell types in scRNA-seq data across mock-infected and reovirus-infected ileum at two distinct stages. The infection response score represents the gene module score for a panel of 438 genes significantly upregulated (two-sided Wilcoxon test, log fold-change > 1.0 and p-value < 0.01) in the reovirus-infected ileum at 1 dpi as compared to the mock-infected ileum. **F)** Infection response score for spatial transcriptomics data from mock-infected and reovirus-infected ileum at two distinct stages.

Cytotoxic T cells recruited by inflamed endothelial cells induce pyroptosis

To explore the heterogeneity of endothelial cell phenotypes in more detail, we reclustered all 9,786 cardiac endothelial cells in the scRNA-seq data. We observed four distinct phenotypes: **i)** uninfamed venous endothelial cells expressing *Nr2f2* and *Ap1nr* mainly derived from the mock controls, **ii)** arterial endothelial cells expressing *Gja4*, *Gja5*, and *Cxcl12* derived from both mock and infected cardiac hearts, **iii)** inflamed endothelial cells derived from infected hearts at 4 and 10 dpi, and **iv)** inflamed endothelial cells from the heart at 7 dpi, with both inflamed endothelial cell clusters expressing *Isg15*, *ligp1*, and *Ly6a* (**Fig. 3A-B**). DGEA across endothelial subclusters revealed that the inflamed 7 dpi endothelial cells overexpressed chemokines *Cxcl9* and *Cxcl10*, which are generally involved in immunoregulatory and

inflammatory processes, but more specifically in the recruitment of T cells (**Fig. 3B-3C, Supp Fig. 3E**). In line with this observation, T cells in the 7 dpi hearts expressed the *Cxcr3* receptor (see below). The *Cxcl9*-high inflamed endothelial cells furthermore expressed high levels of cell adhesion marker genes *Vcam1* and *Icam1*, which help immune cells in the blood to attach to endothelial cells¹⁵ (**Fig. 3C, Supp Fig. 3E, 3I**). The endothelial cells also overexpressed MHC class 1 (*H2-D1* and *H2-K1*) and MHC class 2 (*Cd74*) molecules, suggesting their involvement in antigen presentation to adaptive immune cells (**Fig. 3B-C, Supp Fig. 3E**). Endothelial cells have been shown to be involved in antigen presentation and shaping the cellular immune response in infectious myocarditis¹⁵. Gene ontology (GO) term enrichment analysis identified pathways further supporting the *Cxcl9*-high endothelial cells' involvement in leukocyte cell-cell adhesion, T cell activation, regulation of interleukin-8 production, and response to cytokines, interferon-gamma, interleukin-1, and tumor necrosis factors (**Supp Fig. 3F**).

The observation that endothelial cells are involved in the recruitment of T cells prompted us to explore the heterogeneity of T cells in the infected hearts in more detail. To this end, we reclustered 2,205 T cell single-cell transcriptomes, leading to four subclusters representing three T cell subtypes, *i*) *Cd8+* cytotoxic T cells, *ii*) *Cd4+* helper T cells, and *iii*) naive T cells (**Fig 3E-F**). Both the cytotoxic and helper T cells identified within infected hearts expressed *Cxcr3* receptor, interferon-gamma (*Ifng*), and the chemokines *Ccl3*, *Ccl4*, *Ccl5*, *S100A4*, and *S100A6*, suggesting their involvement in neutrophil recruitment and activation (**Fig. 3G, Supp Fig. 3G**). The *Cxcr3* receptor binds selectively to the chemokines *Cxcl9* and *Cxcl10*, promoting chemotaxis (**Fig. 3G**). Cytotoxic T cells represented the majority of infiltrating T cells and expressed *Prf1*, *Gzma*, *Gzmb*, and *Gzmk*, coding for lytic molecules associated with the granzyme-dependent exocytosis pathway¹⁶ (**Fig. 3F-G, Supp Fig. 3G, 3K**). These cells also expressed tumor necrosis factor superfamily genes *Fasl* and *Tradd*, which are involved in the Fas-induced cell death pathway. *Fasl* binds to *Fas* on the surface of target cells and mediates programmed cell death signaling and NF- κ B activation (**Fig. 3G**). The *Fasl*-*Fas* apoptosis pathway is important in regulating T cells, in promoting tolerance to self-antigens, and is a mechanism by which cytotoxic T cells kill target cells¹⁶. GO term enrichment analysis identified pathways involved in neutrophil activation and degranulation, processing and presentation of exogenous peptide antigen, interleukin-1-mediated signaling pathway, tumor necrosis factor-mediated signaling, NIK/NF-kappaB signaling, cellular response to lectin, and apoptotic processes (**Supp Fig. 3H**).

The downstream gene markers for cell death-associated pathways *Pycard*, *Acer2*, *Zbp1*, and Caspases *Casp1*, *Casp4*, and *Casp12* were enriched in the *Cxcl9*-high endothelial cells. This raises the possibility that cytotoxic lymphocytes are responsible for inflamed endothelial cell death (**Fig. 3B-3C, Supp Fig. 3I**). GO term enrichment of endothelial cells confirmed an upregulation of cell death pathways including activation of cysteine-type endopeptidase activity involved in the apoptotic process, positive regulation of extrinsic apoptotic signaling pathway, and pyroptosis pathway (**Supp Fig. 3F**). We assessed the spatial transcriptomic data to

validate direct interactions between *Cxcl9*-high inflamed endothelial cells and T cells and found that they were indeed spatially co-localized in the myocarditic regions and the border zone (**Supp Fig. 1F**). We calculated gene module scores for genes associated with ontology terms enriched in *Cxcl9*-high endothelial and cytotoxic T cells for spatial transcriptomics data and found these pathways to be enriched in the myocarditic regions (**Fig. 3D, 3H, and Supp Fig. 3I-3L**). Collectively, these results suggest that endothelial cells lining the cardiac vasculature act as a blood-heart barrier and play an important role in the recruitment and activation of the host adaptive immune system. These cells may be the target of both direct viral damage and immune-mediated damage during reovirus-induced myocarditis. Damage to the microvasculature within the heart may then cause loss of blood supply and be a factor in the subsequent death of cardiomyocytes independent of direct viral replication.

Figure 3 Cytotoxic T cells recruited by inflamed endothelial cells induce pyroptosis in myocarditic tissue
A) UMAP plot of 9,786 single-cell endothelial cell transcriptomes from mock-infected and reovirus-infected hearts at 4, 7, and 10 dpi colored by endothelial cell (EC) subtype clusters (phenotypes) (top) and condition (bottom). **B)** Heatmap showing top-five differentially expressed genes (two-sided Wilcoxon test, log fold-change > 1.0 and p-value < 0.01) for endothelial cell subtypes. **C)** UMAP plot showing the expression of genes upregulated in *Cxcl9*-high endothelial cells. **D)** Spatial transcriptomic maps of cardiac tissue from reovirus infected hearts at 7 dpi showing gene module scores calculated for four GO terms enriched in *Cxcl9*-high endothelial cells. **E)** UMAP plot of 2,205 single-cell T cell (TC) transcriptomes from mock-infected and reovirus-infected hearts at 4, 7, and 10 dpi colored by T cell subtype clusters (top) and condition (bottom). **F)** Heatmap showing top-five differentially expressed genes (two-sided Wilcoxon test, log fold-change > 1.0 and p-value < 0.01) for T cell subtypes. **G)** UMAP plot showing the expression of genes upregulated in cytotoxic T cells from myocarditic heart at 7 dpi. **H)** Spatial transcriptomics maps of cardiac tissue from reovirus infected hearts at 7 dpi showing gene module scores calculated for four GO terms enriched in cytotoxic T cells.

Spatially restricted gene expression in myocarditic tissue

The spatially restricted nature of myocarditis motivated us to explore the spatial heterogeneity of gene expression in reovirus-infected hearts. Our initial clustering of the spatial transcriptomic data revealed distinct transcriptional programs for myocarditic regions, the tissue bordering these myocarditic regions, and the rest of the ventricular tissue (**Fig. 1C, 4A**). Differential spatial gene expression analysis for these regions revealed upregulation in the myocarditic regions of cell-type markers for infiltrating immune cells, (*Cd8a* and *Gzma* for T cells, *Nkg7* for NK cells, *S100a8* for neutrophils), markers of inflammation (*Cd52* and *Lyc62*, **Supp Fig. 4A**), and chemokines and cytokines (*Ccl5*, *Ccl2*, *Cxcl9*, and *Cxcl10*). Analysis of the corresponding scRNA-seq data showed that *Ccl5* is expressed by dendritic cells, *Ccl2* by fibroblasts, and *Cxcl9* and *Cxcl10* by endothelial cells. The receptor for *Ccl2*, *Ccr2*, is expressed in macrophages, indicating that fibroblasts use the *Ccl2-Ccr2* axis for macrophage recruitment during myocardial inflammation, as described recently^{17,18} (**Supp Fig. 4A-B**). Collectively these analyses show that chemokine-producing endothelial and cytokine-producing fibroblast cells recruit immune cells to the myocarditic tissue.

Closer inspection of the myocarditic regions and border zones showed an upregulation of additional genes of interest, including *Timp1*, *AW112010*, *Clu*, *Ankrd1*, *Gm4841*, and *Ctss* (**Fig. 4B**). *Timp1* is mainly expressed by inflamed fibroblasts in the scRNA-seq data (**Supp Fig. 4D**). *Timp1* is a natural inhibitor of the matrix metalloproteinases (MMPs), a group of peptidases involved in the degradation of the extracellular matrix. Upregulation of *Timp1* in patients with deteriorating heart failure was reported previously¹⁹. *AW112010* is expressed by inflamed endothelial cells and fibroblasts in the scRNA-seq data and was previously found to code for an interferon-induced small secreted protein with a crucial role in the innate immune response to infection and inflammation²⁰. *Clu* is expressed in a subset of inflamed cells from all cardiac cell types in our data and was previously shown to be upregulated during severe myocarditis²¹ (**Supp Fig. 4D**). *Ctss*, expressed mainly in monocytes, encodes a protease used for degradation of antigenic proteins to peptides for presentation on MHC class II molecules. Increased formation of immunoproteasomes in susceptible mice has been shown to affect the generation of antigenic peptides and subsequent T cell activity in viral myocarditis^{22,23}. GO term analysis of genes upregulated in the border zone revealed enrichment of terms related to the response to tumor necrosis factor, response to interleukin-1, and NIK/ NF-κB signaling (**Supp Fig. 4E**).

To further understand the effect of immune cell infiltration on the cell type composition around the myocarditic regions, we assessed cell type proportions as a function of distance from myocarditic regions in the tissue. We quantified the cell type proportions in myocarditic regions, the border zones, and the rest of the ventricular tissue, and found that the fraction of *Cxcl9*-high endothelial cells, *Ccl2*+ fibroblasts, T cells, dendritic cells, and NK cells was increased in the myocarditic regions, and the fraction of cardiomyocytes was reduced in myocarditic regions (**Fig. 4C, Supp Fig. 1F**).

283

284 To understand the phenotype of *Ccl2*⁺ fibroblasts enriched in myocarditic region and border
285 zone, we reclustered 9,192 fibroblast cells from the scRNA-seq dataset and identified a distinct
286 cluster of inflamed *Ccl2*⁺ fibroblasts from the infected heart at 7 dpi (**Supp. Fig. 4G-4H**). The
287 *Ccl2*⁺ fibroblast cells expressed high levels of MHC class 1 (*H2-D1* and *H2-K1*), adhesion
288 marker genes *Vcam1* and *Icam1*, and other genes like *Serpina3g*, *C3*, and *Ms4a4d* (**Supp Fig.**
289 **4I**). Moreover, these fibroblast cells also expressed *Casp1* and *Casp4*, suggesting pyroptosis
290 activity in response to cytotoxic T cells (**Supp Fig. 4I**).

291 To investigate the effect of inflammation on cardiomyocytes in myocarditic hearts, we
292 reclustered 502 cardiomyocyte cells from the scRNA-seq dataset and identified three distinct
293 phenotypes: **i**) ventricular myocytes expressing *Myl2*, *Myl3*, and *Mb* derived from mock and
294 infected hearts at 4 and 10 dpi, **ii**) atrial myocytes expressing markers *Myl4*, *Myl7*, and *Nppa*
295 derived from mock and infected hearts at 4 and 10 dpi, and **iii**) inflamed myocytes from the
296 infected heart at 7 dpi expressing innate immunity genes *Isg15*, *Igtp*, and *Iigp1* (**Fig. 4D-E**).
297 Inflamed myocytes from the infected heart at 7 dpi had a distinct phenotype as compared to
298 the myocytes from hearts at 4 and 10 dpi, which clustered with myocyte cells from mock-
299 infected hearts (**Fig. 4E**). To find transcriptional signatures for myocytes present in the border
300 zone, we selected genes that were both enriched in cardiomyocyte cells in the scRNA-seq
301 data and upregulated in the border zone. This analysis revealed that cardiomyocytes in the
302 border zone expressed *Gm4841*, *Gm12185*, *Mt1*, *Mt2*, *Ankrd1*, and *Nppb* (**Fig. 4F, Supp Fig.**
303 **4F**). *Gm4841* and *Gm12185* are interferon-inducible genes produced in response to interferon-
304 gamma. *Mt1* and *Mt2* genes modulate inflammation and support remodeling in ischemic
305 cardiomyopathy in mice²⁴. Upregulation of *Ankrd1*, a myocyte survival factor, occurs during
306 late-stage heart disease in patients with idiopathic dilated cardiomyopathy²⁵. A recent study
307 shows that cardiomyocytes expressing *Ankrd1* are localized in the border zone on day 1 post-
308 myocardial infarction²⁶. Together, our analysis reveals that tissue injury is localized to
309 myocarditic regions with remodeling programs being active in the border zone and
310 demonstrates the importance of spatially resolved molecular measurements to study viral
311 myocarditis.

Figure 4 Myocarditic regions and the border zone have distinct transcriptomic profiles and cell type specific signatures. **A)** Spatial transcriptomics map of cardiac tissue section from reovirus-infected mice at 7 dpi colored by spot clusters representing transcriptionally distinct tissue regions. **B)** Spatial transcriptomics maps of cardiac tissue sections from reovirus-infected mice at 7 dpi showing the expression of differentially expressed genes of interest in the myocarditic and the border zone. **C)** Changes in average predicted cell-type proportions across the infected ventricle, for cell types enriched in the myocarditic region and the border zone. **D)** UMAP plot of 502 single-cell cardiomyocyte cell transcriptomes from mock-infected and reovirus-infected hearts at 4, 7, and 10 dpi colored by myocyte cell subtype (phenotypes) (left) and condition (right). **E)** Heatmap showing top-five differentially expressed genes (two-sided Wilcoxon test, log fold-change > 1.0 and p-value < 0.01) for cardiomyocyte cell subtypes. **F)** Venn Diagram showing myocyte-specific genes upregulated in the border zone around the myocarditic regions (left). UMAP plot showing the expression of myocyte-specific genes which are upregulated in the border zone of myocarditic regions (right).

Reduced adaptive immune cell infiltration associated with reovirus K287T mutant

We recently reported a reovirus mutant T1L S4-K287T (K287T) which has a point mutation in the S4 gene encoding outer capsid protein sigma-3 ($\sigma 3$), a double-stranded (ds) RNA-binding multifunctional protein that promotes viral protein synthesis and facilitates viral entry and assembly⁹. K287T successfully infects the heart but produces lower viral titers relative to reovirus wild-type (WT) and does not cause myocarditis⁹. To validate our findings, we performed additional scRNA-seq for K287T infected hearts at 4, 7, and 10 dpi. We generated a total of 16,771 single-cell transcriptomes and integrated the data with the data from the wild-type (WT) virus. K287T-infected cells clustered with WT-infected cells for almost all cell types (Fig. 5A, Supp Fig. 4J). We performed viral transcript enrichment and compared the mean viral transcripts in WT- and mutant-infected cells. We found similar levels of viral transcripts for WT and K287T viruses at 4 dpi but a 60-fold lower viral load for K287T at 7 dpi, consistent with viral titer assays⁹ (Supp Fig. 4K-4L). We then compared early host responses among cardiac cell types due to K287T and WT infection. K287T induced a similar level of innate immune responses as WT reovirus with endothelial cells showing the highest increase in cardiac IR score (as defined before) at 4 dpi (Fig. 5B). We analyzed the cell type composition of Cxcl9-high endothelial cells and immune cells between K287T- and WT-infected hearts and found

that the 7 dpi K287T-infected heart at 7 dpi had significantly fewer *Cxcl9*-high endothelial cells and immune cells infiltrating the heart compared to WT-infected heart. These differences are in line with the reduced levels of inflammation and tissue injury related to the K287T mutant (**Fig. 5C**). Our results show that cardiac endothelial cells mount a potent and robust innate immune response even when infected with the K287T mutant virus with reduced replication in most cell types. Clearance of the virus from the infected cells by 7 dpi and a weaker adaptive host response then leads to the non-myocarditic phenotype of mutant K287T-infected hearts.

Figure 5 A robust innate immune response but reduced adaptive immune cell infiltration explains the non-myocarditic phenotype on infection with reovirus K287T mutant. A) UMAP plot of 48,455 single-cell cell transcriptomes from mock-infected, reovirus-wildtype (WT) infected, and reovirus mutant (K287T) infected hearts at 4, 7, and 10 dpi colored by cell-type clusters. **B)** Infection response score for cardiac cell types in scRNA-seq data across mock-infected, reovirus-WT infected, and reovirus-K287T infected hearts on 4 dpi. The infection response score represents the gene module score for a panel of 226 genes that are significantly upregulated in the reovirus-WT infected sample as compared to the mock-infected sample at 4 dpi. **C)** Changes in cell-type proportions with time for cell types detected in the myocarditic regions. Panels show the changes in cell-type proportions across mock-infected, reovirus-WT infected, and reovirus-K287T infected cells.

DISCUSSION

Viral myocarditis has been recognized as a cause of heart failure for more than 50 years, but it is still a challenging disease to study, diagnose, and treat²⁷. Here, we used integrated spatial and single-cell RNA-seq to dissect the temporal, spatial, and cellular heterogeneity of reovirus-induced acute myocarditis in a neonatal mouse model. We assayed ileum and heart tissues at multiple time points after infection. We investigated the cell types that are infected, and the cellular and spatial heterogeneity of innate and adaptive immune responses. We generated a total of thirteen scRNA-seq and eight spatial transcriptomics datasets, spanning two organs, four time points, and three infection conditions. Our data provide detailed insight into the chronology of molecular events that lead to reovirus-induced myocarditis. After oral inoculation, reovirus T1L infects entero-endocrine and enterocyte cells in the gut epithelium within 1 dpi. These cells mount a potent innate immune response to inhibit viral replication. The virus then infects the gut lymphatic cells within 4 dpi and through lymphatic drainage, the

virus is transmitted via the bloodstream to secondary sites in the body, including the heart. Around 4 dpi, the virus infects the endothelial cells lining the cardiac vasculature. Endothelial cells mount a potent innate immune response in the heart. In symptomatic cases, inflamed endothelial cells secrete chemokines to recruit circulating immune cells, including cytotoxic T cells. Tissue-infiltrating cytotoxic T cells then induce pyroptosis in the myocarditic tissue. Overall, our experiments reveal a dynamic and spatially heterogeneous network of cellular phenotypes and cell-cell interactions associated with reovirus-induced myocarditis.

Integrated high-throughput scRNA-seq and spatial transcriptomics was recently used to study heart development^{28,29} and heart disease^{26,30}, but these methods have not been used to study viral myocarditis prior to our work. Bulk RNA-seq has been used previously to profile transcriptomic signatures of infection, inflammation, and tissue injury associated with viral myocarditis^{9,31,32}. Yet, these ensemble-level approaches do not capture the cellular and spatial heterogeneity of host response to infection. scRNA-seq has recently been used to study Coxsackievirus B3 (CVB3)-induced myocarditis in a mouse model³³. Lasrado et al. report inflammatory phenotypes of myeloid cells, the role of fibroblasts in remodeling and inflammation, and the role of cytotoxic T-cells in CVB3-induced myocarditis. However, the cardiac cell types that are targeted by the virus, the cell type heterogeneity in basal interferon response and innate immune response, and the spatial restriction of transcriptional programs were not explored in this study.

Previous studies have claimed that the direct cytopathic effect of viral replication on cardiac cells is the main cause of cardiac damage during reovirus-induced myocarditis^{7,34}. Notably, Sherry et al. found that reovirus infection can induce myocarditis in immunodeficient mice lacking B and/or T cells, suggesting that reovirus-induced myocarditis does not strictly require adaptive immunity^{7,11}. However, these previous experiments do not rule out the possibility that the host adaptive immune response can augment or delimit the nature and amount of host damage in immune-competent mouse, as is suggested by our work. Holm et al. and Stewart et al. have studied the protective role of innate immune responses in reovirus-induced myocarditis^{13,35}. However, the temporal, spatial, and cell type heterogeneity of basal type-I IFN and innate immune responses to infection had not been characterized before our work. Miyamoto et al. and Stewart et al. compared basal levels of type-I IFN between cardiac myocytes and fibroblasts *in vitro* but these studies did not include all the cell types that make up complex cardiac tissues^{36,37}.

Spatiotemporal characterization of viral myocarditis is crucial to understanding the viral and host factors that are important for disease pathology. This knowledge may ultimately lead to novel diagnostic approaches and better treatments. Several viruses that frequently infect humans can cause myocarditis, including Adenovirus, enteroviruses, Epstein-Barr virus, human Herpesvirus 6, parvovirus B19, and SARS-CoV2. The approaches that we have implemented here can be used in future studies to investigate how the induction, pathophysiology, and course of myocarditis induced by these viruses differs. We hope that the

data and analysis routines that we make available here will be a valuable resource for such future studies.

METHODS

Reovirus infections of neonatal C57BL/6J mice.

Confirmed pregnant female C57BL/6J mice were ordered from Jackson Laboratories to be delivered at embryonic stage E14.5. Litters weighing 3 gram/ pup were gavaged using intramedic tubing (Becton Dickinson 427401) per os with 50 μ l with 10^7 PFU reovirus type 1 lang (T1L): wildtype or K287T mutant in 1x phosphate buffered saline (PBS) containing green food color (McCormick) via a 1ml tuberculin slip tip syringe (BD 309659) and 30G x 1/2 needle (BD 305106). Litters treated with 1x PBS containing green food color alone on the same day were used as mock controls for the respective infection groups. The mock-infected and reovirus-infected mice pups were weighed daily until the time points used in the study (days 1, 4, 7, and 10). All animal work was conducted ethically, conforming to the U.S. Public Health Service policy, and was approved by the Institutional Animal Care and Use Committee at Cornell University (IACUC Number 2019-0129).

Sample preparation for single-cell transcriptomics of cardiac tissue.

We sacrificed mock-infected and reovirus-infected C57BL/6J mice on day 4, day 7, and day 10 post-infection and collected cardiac tissues for single cell transcriptomics. Hearts from respective stages were isolated aseptically, washed with ice-cold Hank's Balanced Salt Solution, HBSS (with calcium and magnesium chloride; Gibco 14025-134), and minced into 1-2mm pieces. Cardiac tissue pieces were then digested in tissue dissociation media with 200U/mL collagenase type II (Gibco 17100-015), 1 mg/ml dispase (Sigma D4693), and 3mM calcium chloride in HBSS for four cycles of 10 minutes under mild agitation at 37°C in 1.5 ml eppendorf tubes. After every 10-minute cycle, cell suspension was collected, added to ice-cold 1x PBS with 0.04% bovine serum albumin (BSA; Sigma A3803) and new dissociation media was added to the tubes. At the end of the digestion, the cells were passed through a 70 μ m filter and centrifuged into a pellet. To remove most blood contaminants, samples were resuspended in an ammonium-chloride-potassium (ACK) lysis buffer (Lonza #10-548E) for 3-5 minutes and centrifuged. Samples were then washed again in PBS with 0.04% BSA and then resuspended at 1×10^6 cells per ml. Cells from each sample were stained with Trypan Blue and cell viability was calculated on an automated cell counter (Countess II) before loading the cells on 10x Chromium. We used these cell viabilities to adjust the number of cells loaded on 10x Chromium to get the desired number of transcriptomes from viable cells for each sample (5000 cells per sample).

Sample preparation for single-cell transcriptomics of intestinal tissue.

We sacrificed mock-infected and reovirus-infected C57BL/6J mice on days 1 and 4 post-infection and collected intestinal ileum tissue for single cell transcriptomics. Intestinal ileum

from respective stages were isolated aseptically, washed with ice-cold Hank's Balanced Salt Solution, HBSS (without calcium and magnesium chloride; Gibco 14175-095) to remove contamination. The ileum tissue was then opened longitudinally, washed again with HBSS, and minced into 1-2mm pieces. To isolate the epithelial layer of cells, ileum tissue pieces were incubated in HBSS with 10mM Ethylenediaminetetraacetic acid (EDTA, Invitrogen 15575-038) and 1mM Dithiothreitol, (DTT, Sigma 43816-10ML) for two cycles of 10 minutes under mild agitation at 37°C. After every 10-minute cycle, cell suspension containing the intestinal epithelial cells was collected, added to ice-cold 1x PBS with 0.04% bovine serum albumin (BSA; Sigma A3803). The undigested pieces of lamina propria were then washed thoroughly with PBS (with calcium and magnesium chloride; Gibco 14080-055) to get rid of all EDTA. These pieces were then transferred to fresh tubes and incubated in 200U/ml Collagenase type I (Gibco 17100-017) and 3mM calcium chloride in PBS for three cycles of 10 minutes under mild agitation at 37°C in 1.5 ml eppendorf tubes. After every 10-minute cycle, cell suspension containing the lamina propria cells was collected, added to ice-cold phosphate buffered saline, PBS with 0.04% BSA in separate tubes. At the end of the digestion, the cells were passed through a 40µm filter and washed twice in PBS with 0.04% BSA and then resuspended at 1×10^6 cells per ml. Cells from intestinal epithelium and lamina propria for each sample were stained with Trypan Blue and cell viability was calculated on automated cell counters (Countess II). Cell counts adjusted with viability were then pooled as 40% epithelial cells and 60% lamina propria to adjust the number of cells loaded on 10x Chromium and to get the desired number of transcriptomes from viable cells for each sample (5000 cells per sample).

Single-cell RNA sequencing library preparation.

5000-6000 viable cells per sample (for heart and ileum tissues) were targeted on the Chromium platform (10x Genomics) using one lane per sample per time point. Single-cell libraries were built using the Chromium Next GEM Single Cell 3' Library Construction V3 Kit (10x Genomics) and were then sequenced on an Illumina NextSeq 500 using 75 cycle high output kits (Index 1 = 8, Read 1 = 28, and Read 2 = 55) for all samples. Sequencing data were aligned to a combined mouse and reovirus reference genome (described below) using the Cell Ranger 6.0.0 pipeline (10x Genomics).

Sample preparation for Visium spatial transcriptomics.

Whole hearts and intestinal ileum were isolated using aseptic techniques and placed in ice cold sterile Hank's Balanced Salt Solution, HBSS (without calcium and magnesium chloride; Gibco 14175-095). Blood and other contamination were carefully removed by perfusing the tissues with fresh HBSS. Fresh tissues were immediately embedded in Optimal Cutting Compound (OCT) media (SAKURA 25608-930) and frozen in a liquid-nitrogen-cooled isopentane (EMD Millipore, MX0760) bath for spatial transcriptomics experiments. The tissue blocks were cut into 10µm sections using Thermo Scientific CryoStar NX50 cryostat and mounted on Visium Gene Expression slides (10x Genomics), which were pre-cooled to -20°C and used for the Visium Spatial Gene Expression experiment.

Visium spatial transcriptomics library preparation.

We used the Visium Spatial Gene Expression (10x Genomics) platform for the spatial transcriptomics experiments. Tissue sections from fresh-frozen hearts (mock-infected and reovirus-infected at day 4 and day 7 post infection) and ileum (mock infected and reovirus infected at day 1 and day 4 post infection) were mounted with one section per capture area on individual Visium Gene Expression slides. These sections are then fixed in pre-chilled methanol for 30 minutes and then hematoxylin and eosin (H&E) stained and imaged, which is later used by the 10x Genomics Space Ranger (version 1.0.0) software to detect the spots which are covered by the tissue. The optimal permeabilization time for 10 μ m thick sections was found to be 18 minutes for the heart and 12 minutes for the ileum using the 10x Genomics Visium Tissue Optimization kit. Spatially tagged cDNA libraries were built using the 10x Genomics Visium Spatial Gene Expression 3' Library Construction V1 Kit. H&E-stained heart tissue sections were imaged using Zeiss PALM MicroBeam laser capture microdissection system at 20x objective and the images were stitched and processed using Fiji ImageJ software. cDNA libraries were sequenced on an Illumina NextSeq 500/550 using 150 cycle high output kits (Read 1 = 28, Read 2 = 120, Index 1 = 10, and Index 2 = 10) for ileum and on an Illumina NextSeq 2K (P2 flow cell) using the 100-cycle kit (Read 1 = 28, Read 2 = 96, Index 1 = 10, and Index 2 = 10). Fiducial frames around the capture area on the Visium slide were aligned manually and spots covering the tissue were selected using Loop Browser 4.0.0 software (10x Genomics). Sequencing data was then aligned to a combined mouse and reovirus reference genome (described below) using the Space Ranger 1.0.0 (10x Genomics) pipeline to derive a feature spot-barcode expression matrix. Visium slide number V19B23-046 was used for spatial transcriptomics experiment on mice hearts (mock-infected 4 dpi: capture area D1, reovirus-infected 4 dpi: capture area B1, mock-infected 7 dpi: capture area C1, and reovirus-infected 7 dpi: capture area A1). Visium slide number V19B23-045 was used for spatial transcriptomics experiment on mice ileum tissue (mock-infected 1 dpi: capture area D1, reovirus-infected 1 dpi: capture area B1, mock-infected 4 dpi: capture area C1, and reovirus-infected 4 dpi: capture area A1).

Hybridization-based enrichment of viral fragments

We performed a hybridization-based enrichment of viral fragments on a part of scRNA-seq libraries using xGen NGS target enrichment kit (IDT; 1080577). In this approach, a panel of 5'-biotinylated oligonucleotides is used for capture and pulldown of target molecules of interest, which are then PCR amplified and sequenced. We designed a panel of 202 biotinylated probes tiled across the entire reovirus T1L genome to selectively sequence viral molecules from the scRNA-seq libraries (**Supp Table 1**). 300ng of fragmented and indexed scRNA-seq libraries from reovirus-WT infected hearts, reovirus-mutant infected hearts, and reovirus-infected ileum were pooled in three separate reactions for xGen hybridization capture. Two rounds of hybridization capture using the xGen enrichment protocol were performed for every reaction to enrich viral transcripts. Amplification was performed for a total of 18 PCR cycles after the first round of capture. 50% of the amplified product was used for the second round of hybridization

capture and amplification was performed for a total of 5 PCR cycles after the second round of enrichment. Post-enrichment products were pooled and sequenced on Illumina Mini-seq for ileum libraries and NextSeq 500 for heart libraries.

Reference genome and annotation

Mus musculus genome and gene annotations (assembly: GRCm38) were downloaded from the Ensembl genome browser, and reovirus strain type-1 Lang genome and gene annotations were downloaded and compiled from the NCBI browser. We have shared reovirus genome sequence and annotation files on figshare with the identifier <https://doi.org/10.6084/m9.figshare.c.5726372>. Genomes were processed using the Cell Ranger v-3.0.0 (10x Genomics) pipeline's mkref command.

Single-cell RNAseq data processing and visualization

Cells with fewer than 200 unique genes or more than 20 percent of transcripts aligning to mitochondrial genes were removed. The single-cell transcriptomes were log-transformed and normalized using the Scanpy package version-1.8.1³⁸. We used Scanpy to choose the highly variable genes with min_disp=0.5 and max_mean=3 thresholds. We then performed mean centering and scaling while regressing out total UMI counts, percent mitochondrial transcripts, S score, and G2M score, followed by principal component analysis (PCA) to reduce the dimensions of the data to the top 20 principal components (PCs). Uniform Manifold Approximation and Projection (UMAP) and the Nearest Neighbor (NN) graph were initialized in this PCA space using the first 20 PCs. The cells were then clustered using the Leiden method with multiple values of clustering resolution to get fine (resolution=0.5) and broad (resolution=0.3) celltype clusters. Cell-type-specific canonical gene markers along with differentially expressed genes (wilcoxon method) for each cluster were used to assign cell type labels. Normalized gene expression was visualized on DotPlots, UMAP plots, and Violin plots across cell type groups. A few cell type clusters representing cell states of the same cell type were grouped into broad cell type groups using cell type marker genes and then used for downstream analysis. Differential gene expression analysis (DGEA) was performed using the rank_gene_groups function in Scanpy with the Wilcoxon statistical method. All gene module scores were calculated using the score_genes function in scanpy.

Reclustering and analysis of endothelial cells, T cells, fibroblasts, and cardiomyocytes

Normalized gene expression for a specific cell type group was extracted from the combined scRNA-seq dataset. We used Scanpy to reselect the highly variable genes within that cell type group with min_disp=0.5 and max_mean=3 thresholds. We then performed mean centering and scaling while regressing out total UMI counts, percent mitochondrial transcripts, S score, and G2M score, followed by principal component analysis (PCA) to reduce the dimensions of the data to the top 20 principal components (PCs). Uniform Manifold Approximation and Projection (UMAP) and the Nearest Neighbor (NN) graph were initialized in this PCA space

using the first 20 PCs. The cells were then reclustered using the Leiden method (resolution=0.5 for endothelial cells, resolution=0.3 for T cells, resolution=0.2 for fibroblasts, and resolution=0.3 for cardiomyocytes) to get cell type subclusters. Differentially expressed genes (wilcoxon method) for each subcluster were then used to assign cell subtype labels. Subclusters representing doublets and expressing markers of multiple cell types were then removed from the analysis. Normalized gene expression for differentially expressed genes and genes of interest was visualized on DotPlots and UMAP plots across celltype subgroups. Differential gene expression analysis (DGEA) was performed using the rank_gene_groups function in Scanpy with the Wilcoxon statistical method. All gene module scores were calculated using the score_genes function in Scanpy.

Spatial transcriptomics data processing, integration, analysis, and visualization

Spatial transcriptomics data from barcoded spatial spots from four heart sections were log-normalized using the Scanpy package (1.8.1). Scanpy package was then used to select highly variable genes for spatial transcriptomics data with min_disp=0.5 and max_mean=3 thresholds. We then performed mean centering and scaling while regressing out total UMI counts, percent mitochondrial UMIs, S score, and G2M score, followed by PCA on the spot gene expression matrix, and reduced the dimensions of the data to the top 20 principal components. UMAP and the NN graph were initialized in this PCA space. The spots were then clustered using the Leiden method with multiple values of clustering resolution. The method returned spot clusters representing different tissue regions, which were then visualized on H&E images as spatial transcriptomics maps for individual samples to assign anatomical regions. Normalized gene expression was visualized on spatial transcriptomics maps for all tissue sections. Spot clusters representing the same tissue regions were grouped into broad anatomical region groups using marker genes and then used for downstream analysis. Stereoscope deconvolution method³⁹ implemented in the scvi-tools⁴⁰ package was used for integration of spatial transcriptomics data with time-matched scRNA-seq data and cell type prediction values for spatial transcriptomics spots were estimated for the infected heart at 7 dpi. DGEA for anatomical regions was performed using the rank_gene_groups function in Scanpy with the Wilcoxon statistical method.

Viral transcript sequencing data processing, filtering, and visualization

Enriched viral transcript data were aligned to a combined mouse and reovirus type-1 Lang genome for all infected samples. Viral unique molecule (UMI) counts were taken from the combined expression matrices and added as metadata in the host gene expression data. Viral UMI counts in empty droplets, droplets with low-quality cells (< 200 host UMI counts), droplets with viable cells (>=200 host UMI counts) were sorted by viral UMI and visualized on a histogram to filter out the cell-free ambient viral RNA enriched in the hybridization protocol. Using the distribution of viral UMI counts in empty droplets, thresholds of two viral UMIs and ten viral UMIs were used to identify infected cells in the heart and ileum respectively (**Supp Fig**

2D). Viral transcripts in the infected cells were then visualized on a DotPlot to determine viral tropism in tissues.

Gene Ontology term enrichment analysis for scRNA-seq and spatial transcriptomics

Gene Ontology (GO) term enrichment analysis was performed on differentially expressed genes using gseapy wrapper package⁴¹. Differentially expressed genes (two-sided Wilcoxon test, log fold-change threshold = 2.0, p-value < 10⁻⁴ for scRNA-seq cells, and log fold-change threshold = 0.5, p-value < 10⁻² for spatial transcriptomics spots) were selected and used for GO term enrichment analysis using GO_Biological_Processes_2021 gene sets in enrichr command⁴². The enriched GO terms of interest were selected and visualized on a BarPlot. The genes associated with GO terms of interest were used to calculate module scores using score_genes command in Scanpy.

ACKNOWLEDGEMENTS

We would like to thank Peter Schweitzer and the Cornell Genomics Center for help with single-cell and spatial sequencing assays, Cornell Bioinformatics facility for assistance with bioinformatics, and Dr. Danica M. Sutherland from the lab of Dr. Terence S. Dermody at University of Pittsburgh for assistance with animal experiments. We also thank the members of the Parker and De Vlaminc labs for many valuable discussions. This work was supported by R21AI144557 (to J.S.P. and I.D.V.), and DP2AI138242 (to I.D.V.). S.T.C. was supported by the National Institutes of Health and National Institute of Allergy and Infectious Diseases Award T32AI145821.

AUTHOR CONTRIBUTIONS

M.M., J.S.P., and I.D.V. designed the study. M.M., M.M.H., and S.T.C. performed the animal experiments. M.M. and M.M.H. performed the scRNA-seq and spatial transcriptomics experiments. M.M., D.W.M., and M.F.Z.W. analyzed the data. M.M., J.S.P., and I.D.V. wrote the manuscript. All authors provided feedback and comments.

DATA AVAILABILITY

The authors declare that all sequencing data supporting the findings of this study have been deposited in NCBI's Gene Expression Omnibus (GEO)⁴³ with GEO series accession number [GSE189636](https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE189636). Raw and processed H&E-stained tissue images and tissue-spot alignment files matched to spatial transcriptomics datasets have been made publicly available on figshare with identifier <https://doi.org/10.6084/m9.figshare.c.5726372>⁴⁴. Scripts to reproduce the analysis presented in this study have been deposited on GitHub (https://github.com/madhavmantri/reovirus_induced_myocarditis).

CONFLICTS

636 The authors declare no conflicts.

637

638 REFERENCES

- 639 1. Pollack, A., Kontorovich, A. R., Fuster, V. & Dec, G. W. Viral myocarditis—diagnosis, treatment
640 options, and current controversies. *Nat. Rev. Cardiol.* 2015 1211 **12**, 670–680 (2015).
- 641 2. Rose, N. R. Viral Myocarditis. *Curr. Opin. Rheumatol.* **28**, 383 (2016).
- 642 3. Yajima, T. & Knowlton, K. U. Viral myocarditis from the perspective of the virus. *Circulation* **119**,
643 2615–2624 (2009).
- 644 4. Tschöpe, C. *et al.* Myocarditis and inflammatory cardiomyopathy: current evidence and future
645 directions. *Nat. Rev. Cardiol.* **18**, 169–193 (2021).
- 646 5. Woodruff, J. F. Viral myocarditis. A review. *Am. J. Pathol.* **101**, 425 (1980).
- 647 6. Lasrado, N. & Reddy, J. An overview of the immune mechanisms of viral myocarditis. *Rev. Med.*
648 *Viol.* **30**, 1–14 (2020).
- 649 7. Sherry, B., Schoen, F. J., Wenske, E. & Fields, B. N. Derivation and characterization of an
650 efficiently myocarditic reovirus variant. *J. Virol.* **63**, 4840–4849 (1989).
- 651 8. Boehme, K. W., Lai, C. M. & Dermody, T. S. Mechanisms of reovirus bloodstream dissemination.
652 *Adv. Virus Res.* **87**, 1–35 (2013).
- 653 9. Guo, Y. *et al.* The multi-functional reovirus $\sigma 3$ protein is a virulence factor that suppresses stress
654 granule formation and is associated with myocardial injury. *PLOS Pathog.* **17**, e1009494 (2021).
- 655 10. Swirski, F. K. & Nahrendorf, M. Cardioimmunology: the immune system in cardiac homeostasis
656 and disease. *Nat. Rev. Immunol.* 2018 1812 **18**, 733–744 (2018).
- 657 11. Sherry, B., Li, X. Y., Tyler, K. L., Cullen, J. M. & Virgin, H. W. Lymphocytes protect against and
658 are not required for reovirus-induced myocarditis. *J. Virol.* **67**, 6119–6124 (1993).
- 659 12. Phillips, M. B., Dina Zita, M., Howells, M. A., Weinkopff, T. & Boehme, K. W. Lymphatic Type 1
660 Interferon Responses Are Critical for Control of Systemic Reovirus Dissemination. *J. Virol.* **95**,
661 (2021).
- 662 13. Holm, G. H. *et al.* Interferon Regulatory Factor 3 Attenuates Reovirus Myocarditis and
663 Contributes to Viral Clearance. *J. Virol.* (2010). doi:10.1128/jvi.01742-09
- 664 14. Schaum, N. *et al.* Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris. *Nat.*
665 *2018 5627727* **562**, 367–372 (2018).
- 666 15. Woudstra, L., Juffermans, L. J. M., van Rossum, A. C., Niessen, H. W. M. & Krijnen, P. A. J.
667 Infectious myocarditis: the role of the cardiac vasculature. *Heart Fail. Rev.* **23**, 583–595 (2018).
- 668 16. Chávez-Galán, L., Arenas-Del Angel, M. C., Zenteno, E., Chávez, R. & Lascurain, R. Cell death
669 mechanisms induced by cytotoxic lymphocytes. *Cell. Mol. Immunol.* **6**, 15–25 (2009).

- 670 17. Leuschner, F. *et al.* Silencing of CCR2 in myocarditis. *Eur. Heart J.* **36**, 1478–1488 (2015).
- 671 18. Miteva, K. *et al.* Mesenchymal Stromal Cells Modulate Monocytes Trafficking in Coxsackievirus
672 B3-Induced Myocarditis. *Stem Cells Transl. Med.* **6**, 1249–1261 (2017).
- 673 19. Barton, P. J. R. *et al.* Increased expression of extracellular matrix regulators TIMP1 and MMP1 in
674 deteriorating heart failure. *J. Heart Lung Transplant.* **22**, 738–744 (2003).
- 675 20. Jackson, R. *et al.* The translation of non-canonical open reading frames controls mucosal
676 immunity. *Nature* **564**, 434–438 (2018).
- 677 21. Swertfeger, D. K., Witte, D. P., Stuart, W. D., Rockman, H. A. & Harmony, J. A. K. Apolipoprotein
678 J/clusterin induction in myocarditis: A localized response gene to myocardial injury. *Am. J.*
679 *Pathol.* **148**, 1971–1983 (1996).
- 680 22. Szalay, G. *et al.* Ongoing Coxsackievirus Myocarditis Is Associated with Increased Formation
681 and Activity of Myocardial Immunoproteasomes. *Am. J. Pathol.* **168**, 1542–1552 (2006).
- 682 23. Van Der Borght, K. *et al.* Myocarditis elicits dendritic cell and monocyte infiltration in the heart
683 and self-antigen presentation by conventional type 2 dendritic cells. *Front. Immunol.* **9**, 2714
684 (2018).
- 685 24. Duerr, G. D. *et al.* Metallothioneins 1 and 2 Modulate Inflammation and Support Remodeling in
686 Ischemic Cardiomyopathy in Mice. *Mediators Inflamm.* **2016**, (2016).
- 687 25. Bogomolovas, J. *et al.* Induction of Ankrd1 in dilated cardiomyopathy correlates with the heart
688 failure progression. *Biomed Res. Int.* **2015**, (2015).
- 689 26. Yamada, S. *et al.* Spatiotemporal single-cell analysis reveals critical roles of mechano-sensing
690 genes at the border zone in remodeling after myocardial infarction. doi:10.21203/rs.3.rs-
691 620498/v1
- 692 27. Gupta, S., Markham, D. W., Drazner, M. H. & Mammen, P. P. A. Fulminant myocarditis. *Nat.*
693 *Clin. Pract. Cardiovasc. Med.* **2008 511 5**, 693–706 (2008).
- 694 28. Mantri, M. & Scuderi, G. J. Spatiotemporal single-cell RNA sequencing of developing chicken
695 hearts identifies interplay between cellular differentiation and morphogenesis.
696 *Github:chicken_heart* (2020). doi:<https://doi.org/10.5281/zenodo.4517120>
- 697 29. Asp, M. *et al.* A Spatiotemporal Organ-Wide Gene Expression and Cell Atlas of the Developing
698 Human Heart. *Cell* **179**, 1647-1660.e19 (2019).
- 699 30. Kuppe, C. *et al.* Spatial multi-omic map of human myocardial infarction. *bioRxiv*
700 2020.12.08.411686 (2020). doi:10.1101/2020.12.08.411686
- 701 31. Lindner, D. *et al.* Association of Cardiac Infection With SARS-CoV-2 in Confirmed COVID-19
702 Autopsy Cases. *JAMA Cardiol.* **5**, 1281–1285 (2020).
- 703 32. Bräuninger, H. *et al.* Cardiac SARS-CoV-2 infection is associated with pro-inflammatory
704 transcriptomic alterations within the heart. *Cardiovasc. Res.* (2021). doi:10.1093/CVR/CVAB322
- 705 33. Lasrado, N., Borchering, N., Arumugam, R., Starr, T. K. & Reddy, J. Dissecting the Cellular
706 Landscape and Transcriptome Network in Viral Myocarditis by Single-Cell RNA Sequencing.

707 *bioRxiv* 2021.05.16.444367 (2021). doi:10.1101/2021.05.16.444367

708 34. Baty, C. J. & Sherry, B. Cytopathogenic effect in cardiac myocytes but not in cardiac fibroblasts
709 is correlated with reovirus-induced acute myocarditis. *J. Virol.* **67**, 6295–6298 (1993).

710 35. Stewart, M. J., Blum, M. A. & Sherry, B. PKR's protective role in viral myocarditis. *Virology* **314**,
711 92–100 (2003).

712 36. Miyamoto, S. D. *et al.* Cardiac Cell-specific Apoptotic and Cytokine Responses to Reovirus
713 Infection: Determinants of Myocarditic Phenotype. *J. Card. Fail.* **15**, 529–539 (2009).

714 37. Stewart, M. J., Smoak, K., Blum, M. A. & Sherry, B. Basal and Reovirus-Induced Beta Interferon
715 (IFN- β) and IFN- β -Stimulated Gene Expression Are Cell Type Specific in the Cardiac Protective
716 Response. *J. Virol.* **79**, 2979–2987 (2005).

717 38. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: Large-scale single-cell gene expression data
718 analysis. *Genome Biol.* **19**, 1–5 (2018).

719 39. Andersson, A. *et al.* Single-cell and spatial transcriptomics enables probabilistic inference of cell
720 type topography. *Commun. Biol.* 2020 31 **3**, 1–8 (2020).

721 40. Gayoso, A. *et al.* scvi-tools: a library for deep probabilistic analysis of single-cell omics data.
722 *bioRxiv* 2021.04.28.441833 (2021). doi:10.1101/2021.04.28.441833

723 41. Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for
724 interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci.* **102**, 15545–15550 (2005).

725 42. Xie, Z. *et al.* Gene Set Knowledge Discovery with Enrichr. *Curr. Protoc.* **1**, e90 (2021).

726 43. Edgar, R. Gene Expression Omnibus: NCBI gene expression and hybridization array data
727 repository. *Nucleic Acids Res.* (2002). doi:10.1093/nar/30.1.207

728 44. Mantri, M. Reovirus-induced-myocarditis. (2021).
729 doi:<https://doi.org/10.6084/m9.figshare.c.5726372>

730