Metal transporter SLC39A14/ZIP14 modulates regulation between the gut microbiome and host metabolism

Trista L. Thorn^{1#}, Samuel B. Mitchell^{1#}, Yongeun Kim¹, Min-Ting Lee¹, Janine M. C. Comrie¹, Elizabeth L. Johnson¹, Tolunay B. Aydemir^{1*}

Division of Nutritional Sciences, Cornell University, Ithaca, 14853 NY

[#]Equally contributed

^{*}Corresponding author

Key Words: Zinc, zinc transporter, gut microbiome, metabolomics, intestine

Abstract

Zinc (Zn) plays a critical role in maintaining intestinal homeostasis by regulating intestinal epithelial cells, host immune cells, and gut microbiome community composition. Deletion of metal transporter *Slc39a14/Zip14* causes spontaneous intestinal permeability with low-grade chronic inflammation, mild hyperinsulinemia, and greater body fat with insulin resistance in adipose, suggesting a role for ZIP14-mediated intestinal metal transport in regulating both intestinal homeostasis and systemic metabolism. Here, we showed the function of ZIP14-mediated Zn transport in the gut microbiome composition and how ZIP14-linked changes to gut microbiome community composition are correlated with changes in host metabolism. Deletion of *Zip14* generated Zn-deficient epithelial cells and luminal content in the entire intestinal tract; reduced bacterial diversity and *Saccharomyces cerevisiae* (*S. cerevisiae*) overgrowth; altered host metabolome; and shifted host energy metabolism toward glucose utilization. This work provides evidence for the regulation of gut microbiome composition, host metabolome, and energy metabolism by metal transporter ZIP14.

Significance

Intestinal permeability, gut dysbiosis, and Zn dyshomeostasis are emerging signatures of inflammatory bowel diseases and metabolic disorders such as type-2-diabetes and obesity. Zn deficiency is a common clinical finding among these diseases. Zn is essential for the regulation of the intestinal epithelial cells, host immune cells, and the gut microbiome. Transporter-mediated mobilization of Zn plays a critical role in maintaining intestinal homeostasis by facilitating the targeted tissue/cell-specific function. However, studies are lacking in linking transporter-mediated Zn mobilization, gut microbiome, host's intestinal health, and metabolism. Using the systems-level

approach, this study revealed novel findings that deletion of *Slc39a14/Zip14* resulted in altered intestinal Zn homeostasis, gut microbiome composition, host metabolome and energy metabolism.

Introduction

The essential trace metal Zn is the second most abundant metal in the body, with diverse and indispensable catalytic, structural, and regulatory functions¹. Zn transporters maintain Zn homeostasis by facilitating the cellular and subcellular distribution of Zn to and within tissues. Two families of metal transporters, SLC39A/ZIP1-14 (Zrt-Irt-like proteins) and SLC30A/ZnT1-10 (Zn transporter), have opposing functions. ZIP family transporters are responsible for the transport of metals into the cytoplasm from the extracellular space and within the lumen of intracellular compartments. ZnT family transporters transport metals from the cytoplasm to either extracellular space or intracellular compartments. Intestinal Zn transport is central to both intestinal and systemic Zn homeostasis. Zn is absorbed mainly by the function of apically localized Zn transporter, ZIP4²⁻⁴. On the basolateral side, ZnT1 and, ZIP5 and ZIP14 are proposed to transport Zn from the enterocyte into the circulation and from circulation to enterocytes, respectively^{5,6}. Metallothioneins and the ZnT2, ZnT5A, ZnT6, ZnT7, ZIP7, and ZIP14 transporters provide intracellular Zn movement⁶⁻⁹.

Emerging evidence suggests that Zn transporter-mediated mobilization of Zn plays a critical role in maintaining intestinal homeostasis by facilitating the targeted tissue/cell-specific function. Zn is essential for the regulation of the intestinal epithelial cells, host immune cells, and intestinal commensal bacteria¹⁰. Mice lacking ZIP4 in the intestinal epithelial cells showed loss of Zn in Paneth cells with impaired proliferation and epithelial architecture¹¹. Deleting ZIP7 in the intestinal epithelial cells in mice caused defects in stem cell survival, the proliferation of crypts, and resolving endoplasmic reticulum stress¹². ZnT2 is expressed on the secretory granules in Paneth cells, promotes the accumulation of zinc, and regulates the secretion of antimicrobial molecules¹³. Genome-wide association studies revealed an association between a variant of Zn transporter ZIP8 and type of inflammatory bowel disease (IBD), Crohn's disease (CD) with altered gut microflora¹⁴. Intestinal permeability and inflammation are signatures of IBD. In a humanized mouse model of the *ZIP8* variant, enhanced disease susceptibility was reported in chemically induced colitis^{15,16}.

We have previously shown that deletion of *Zip14*, the phylogenetically closest relative of *Zip8*¹⁷, induced spontaneous intestinal permeability with low-grade chronic inflammation (metabolic

endotoxemia), enlarged islets with mild compensatory hyperinsulinemia, and greater body fat with insulin resistance in adipose tissue^{6,18–20}. Besides Zn, ZIP14 also transports manganese and iron²¹. We have previously shown that intestinal manganese elimination was impaired in *Zip14* KO mice^{22,23}. Here we present our novel findings on the function of ZIP14-mediated Zn transport in intestinal tissue, microbial profile, and how ZIP14-linked changes in gut microbiome and metabolites modulate host metabolism.

Results

Ablation of *Zip14* alters gastrointestinal Zn homeostasis. We first examine the effect of the *Zip14* deletion on Zn homeostasis in the entire intestinal tract using microwave plasma atomic emission spectrometry (MP-AES). In the *Zip14* KO mice, Zn concentrations were lower in the proximal and distal intestine and the colon tissue (Fig. 1A). The total intestinal content is composed of both unabsorbed dietary Zn and Zn from endogenous sources, including transporter-mediated transepithelial flux of Zn in the serosal to the mucosal direction. We collected total intestinal content and separated it into two phases by centrifugation. Luminal content (supernatant) represents ionic and protein-bound Zn, and digesta (pellet) represents the unavailable Zn for host absorption²⁴. Both luminal content and digesta had decreased concentrations of Zn in the *Zip14* KO (Fig. 1B). These data collectively suggested that in the absence of basolaterally localized ZIP14, Zn transport was impaired in serosal to mucosal direction, resulting in a chronic Zn deficiency in the entire GI tract's tissue, luminal content, and digesta. Zn absorption occurs throughout the intestinal tract as a function of Zn transporters. Since the highest Zn absorption occurs in the proximal intestine, we measured the abundance of known intestinal Zn transporters from isolated enterocytes. We observed more pronounced changes in the female mice. Compared to WT, the abundance of apical Zn importer ZIP4 and basolateral Zn exporter ZnT1 in *Zip14* KO mice was increased and decreased, respectively (Fig. 1C), supporting the Zn-deficient conditions in the enterocytes of *Zip14* KO mice. However, these changes were not sufficient to compensate for the deficiency caused by the absence of ZIP14.

Ablation of *Zip14* modulates gut microbiome, and ZIP14-shaped microbiome can influence intestinal permeability and metabolic phenotypes in the host. Intestinal barrier function and consequent metabolic alterations are regulated by dynamic and reciprocal interactions among the intestinal microbiota, epithelium, and immune system²⁵. Dietary metals are important factors influencing the intestinal microbiome²⁶, as evidenced by the observations that metal availability is crucial for the outcome of host-pathogen interactions. However, much less is known about the effects of dietary metals on commensal bacteria. A limited number of studies have demonstrated

that both Zn deficiency and excess can influence the composition of the gut microbiome. However, it is largely unknown whether Zn transporters regulate the gut microbiome through modulating luminal Zn levels. Due to impaired Zn transport (serosal to mucosal), *Zip14* deletion results in lower Zn levels in the gastrointestinal lumen than those in WT mice (Fig. 1B). We hypothesized that a low Zn environment would affect the composition of the gut microbiome. To test this hypothesis, we determined microbiome community composition using 16S rRNA gene sequencing from fecal pellets of *Zip14* KO versus WT mice. We found significantly lower microbiome α -diversity in WB-KO mice when compared to WT mice (Fig. 2A). This suggests that the presence of certain taxa may be reliant on intestinal luminal concentrations of Zn. To better understand which taxa might be susceptible to Zn concentrations in the lumen of the intestine, we looked for differential abundances of taxa between the microbial communities of *Zip14* KO and WT mice (Fig. 2B). We found a decreased abundance of Bacteroidaceae and Clostridiaceae families and increased abundance of Ruminococcaceae families (Fig. 2C), all of which have been observed in IBD patients as well ^{27,28}. To better understand how ZIP14-mediated changes in microbiome community composition may function to alter host phenotypes, we transplanted microbiome-depleted mice with fecal contents from *Zip14* KO and WT mice (Fig. 3A). Following the fecal transplant, we have conducted a permeability assay using FITC-Dextran gavage. We found significantly higher levels of FITC-Dextran in the serum of mice that received the fecal slurry from *Zip14* KO mice (Fig. 3B), indicating that the ZIP14-shaped microbiome contributed to the progression of intestinal permeability. Furthermore, the mice that received the fecal slurry from *Zip14* KO were mildly hypoglycemic (Fig. 3C) with significantly greater % body fat (Fig. 3D) and lower % lean mass (Fig. 3E), recapitulating the metabolic phenotype of *Zip14* KO mice ²⁰. Collectively these data indicated that the ZIP14-shaped microbiome could influence metabolic phenotypes in the host.

***Saccaramicea cerevisiae* (*S. cerevisiae*) overgrowth may contribute to intestinal permeability in *Zip14* KO mice.** The gut microbiota produces a broad range of metabolic products that impact host biology. To identify the host and microbial-originated metabolites in WT and KO mice serum, we conducted untargeted metabolomics using Gas chromatography (GC) coupled to time-of-flight mass spectrometry (TOF-MS). Principal component analysis (PCA) of overall serum samples showed that metabolite abundance in *Zip14* KO mice significantly differed from WT mice (Fig. 4A), indicating that the *Zip14* KO group could be distinguished from WT according to their different metabolic profiles. Metabolites with a fold change of >1.5 and p-value < 0.05 were considered significantly different (Fig. 4B). The serum concentration of 11 identifiable metabolites varied significantly between WT and WB-KO (Fig. 4B, C). Serum adenosine,

serotonin, β -sitosterol, creatinine, N-acetyl-D-tryptophan, and saccharic acid were significantly greater, and only linoleic acid was significantly lower in the serum of *Zip14* KO mice (Fig. 4C).

The upregulated metabolite, N-acetyl-D-tryptophan, in the serum of *Zip14* KO mice could be unique to *S. cerevisiae* in the gut due to the production of a unique enzyme acting on D-amino acids, D-amino acid N-acetyltransferase^{29,30}. We hypothesized that the Zn deficient gut in the *Zip14* KO mice with lower bacterial α -diversity could provide the opportunity for yeast overgrowth. Our qPCR results revealed a greater amount of *S. cerevisiae* DNA in the feces of WB-KO mice when compared to their controls (Fig. 4D). To investigate the direct effect of the gut microbiome on the metabolic phenotype of *Zip14* KO mice, we conventionalized antibiotics (abx)-treated WT mice with fecal slurry from either WT (WTCv) or *Zip14* KO (KOCv) mice. We measured *S. cerevisiae* DNA following conventionalization. The KOCv mice had a greater amount of *S. cerevisiae*; however, it did not reach the statistical significance ($p < 0.07$) (Fig. 4D). To understand the effect of *S. cerevisiae* on the increased intestinal permeability phenotype of WB-KO mice, the mice's drinking water was supplemented with the antifungal drug fluconazole for ten days. We found a significant reduction in plasma FITC-Dextran levels when WB-KO mice were treated with fluconazole, suggesting that *S. cerevisiae* overgrowth along with reduced bacterial diversity in the WB-KO contributed to the development of intestinal permeability.

Host energy metabolism is shifted towards continuous utilization of glucose as an energy source in *Zip14* KO mice. To assist in the biological interpretation of our metabolomics data, we conducted pathway and enrichment analyses using MetaboAnalyst³¹ to explore whether significantly differing metabolites belong to a common pathway. There was an enrichment of significantly differing compounds in several metabolic pathways (Fig. 5A). Among them, the Tricarboxylic Acid Cycle (TCA) cycle pathway had the most significant impact score. TCA cycle intermediate metabolites, citric acid, α -ketoglutaric acid, fumaric acid, and malic acid were significantly upregulated in the serum of *Zip14* KO mice. Of note, α -ketoglutarate, fumaric acid, and malic acid are also responsible for the "alanine, aspartate, and glutamine metabolism"³². Disease signature analysis was conducted using metabolite sets reported in human blood. In agreement with the presence of intestinal permeability and low-grade chronic inflammation in *Zip14* KO mice, the top signatures were septic shock and diabetes (Fig. 5B).

Despite the increased intestinal permeability, low-grade chronic inflammation, and insulin resistance in adipose, *Zip14* KO mice do not develop hyperglycemia. In addition, our metabolomics data revealed the most significant impact score was on TCA cycle (Fig. 5A), collectively suggesting that *Zip14* deletion may cause a shift in host energy metabolism. To

quantify these observed effects on whole-body metabolism, we conducted metabolic phenotyping by using a Comprehensive Laboratory Animal Monitoring System (CLAMS). In the light cycle, oxygen consumption levels (Fig. 6A), carbon dioxide production (Fig. 6B), and respiratory exchange ratio (Fig. 6C) were greater in the *Zip14* KO mice than WT. These data suggested that glucose was preferentially used as an energy source in the *Zip14* KO mice. This may partially explain the prevention of hyperglycemia phenotype. Next, we fed mice with HFD for 16 weeks to force fatty acids (consumption of excess). At the end of the study, fasting blood glucose levels were elevated in HFD fed WT mice, but there was no increase in the HFD-fed *Zip14* KO mice, even though both WT and KO had an equal increase in percent body fat in the HFD fed group (data not shown).

Discussion

In this study, we showed that metal transporter *Slc39a14/Zip14* regulates intestinal Zn homeostasis and gut microbiome composition, resulting in altered host metabolism. Deletion of the host gene encoding metal transporter, *Slc39a14/Zip14*, generated Zn-deficient epithelial and luminal content in the entire intestinal tract; reduced microbial diversity; altered host metabolome; and; shifted host energy metabolism toward glucose utilization (Fig. 7). To our knowledge, this is the first investigation that links the function of a mammalian metal transporter to gut microbiome composition, host serum metabolites, and energy metabolism.

Our previous studies showed localization of ZIP14 both to the basolateral side of the enterocyte and endosomes⁶. In the *Zip14* KO mice, increased endosomal Zn concentrations suggested that Zn was entrapped in these compartments and generated Zn-deficient enterocytes. Here we showed that deletion of *Zip14* caused a deficiency in the intestinal tissue and the luminal content (Fig. 1A, B). These mice consumed a standard chow diet, and there was no difference in food consumption (data not shown). Therefore, low intestinal and luminal Zn suggested that Zn transport in serosal to mucosal direction was impaired in addition to endosomal entrapment in the absence of basolaterally localized ZIP14. ZIP4 is an apically localized Zn transporter and is responsible for the absorption of Zn. It is known to be upregulated during Zn deficiency to increase Zn absorption³³. ZnT1 on the basolateral side of the intestine was proposed to facilitate Zn transport from enterocytes to blood⁸. ZnT1 is also regulated by Zn status⁵. It was shown that during Zn deficiency, ZnT1 was downregulated to retain Zn in the cells. Upregulation of ZIP4 and downregulation of ZnT1 in isolated enterocytes from *Zip14* KO mice further supported the Zn-deficient environment.

ZIP14, besides Zn, can transport manganese and iron ²¹. We have previously shown that Zn and manganese homeostasis were altered in *Zip14* KO mice at a steady-state ^{18,20,23,34}, while ZIP14-related alterations in iron transport were largely observed in iron overload ³⁵. Our current (Fig. 1) and previous results confirmed that both Zn and manganese transport in serosal to mucosal direction was impaired when *Zip14* was deleted ^{23,34}. Since Zn, iron, and manganese are essential for microbial growth, the changes that we observe in gut microbiome composition could be caused by either Zn or Mn deficient intestine of *Zip14* KO mice. Manganese deficiency in humans is extremely rare ^{36,37}. However, a rare genetic mutation in metal transporter, *ZIP8*, causes manganese deficiency with enhanced susceptibility to dextran sulfate sodium-induced injury ¹⁵. Contrary to manganese, Zn deficiency is recognized by WHO as a major global risk factor for disease mortality and morbidity ³⁸. Importantly, Zn deficiency induces diarrhea and increases the risk of gastrointestinal diseases ^{39,40}. Zn deficiency is found in IBD patients, and Zn supplementation studies have shown improved disease outcomes ¹⁰. However, human studies linking Zn, Zn transporters, gut microbiome composition, and intestinal health are lacking. In animal models, both Zn excess and deficiency were shown to modulate the microbiome profile. High dietary Zn in mice reduced the diversity of the gut microbiome community⁴¹, while in piglets, it caused either no change or increased microbiota diversity of ileal digesta and decreased microbiota diversity of the colonic digesta based on the dose and duration of Zn supplementation or the growth phase of the animals ⁴². A less diverse cecal microbiome composition was detected in gallus gallus in response to chronic dietary Zn deficiency ⁴³, supporting our finding of lower bacterial α -diversity in the Zn-deficient intestine of *Zip14* KO mice.

The gut microbiome consists of bacteria, viruses, protozoa, and fungi in the gastrointestinal tract. The gut microbiota produces a broad range of metabolic products that impact host biology by acting on cells within the gastrointestinal tract or entering circulation and affecting distal sites within the body ^{44,45}. Among the upregulated metabolites in the serum of *Zip14* KO mice (Fig. 3), N-acetyl-D-tryptophan, serotonin, and creatinine levels were shown to be different between germ-free and conventional mice ⁴⁶, suggesting microbial origin/contribution. Enteric bacteria or yeast can generate tryptophan metabolites such as serotonin and N-acetyl-tryptophan ^{47,48}. While either host or microbiome could make serotonin, N-acetyl-D-tryptophan could be unique to *Saccharomyces cerevisiae* (*S. Cerevisiae*) in the gut due to the production of a unique enzyme acting on D-amino acids, D-amino acid N-acetyltransferase ^{29,30}. Lower bacterial diversity in the gut of *Zip14* KO mice could provide an opportunity for *S. cerevisiae* overgrowth. Of note, yeast cells survive during Zn deficiency due to multiple homeostatic and adaptive strategies ⁴⁹. Increased intestinal permeability is observed both in Zn deficiency and inflammatory bowel

disease (IBD) ¹⁰. Furthermore, Zn deficiency ⁵⁰ and *S. cerevisiae* overgrowth ⁵¹ have been shown in IBD. Interestingly, serotonin was also shown to be upregulated in Crohn's disease ⁵². These are, collectively, in agreement with the phenotype of *Zip14* KO mice with intestinal permeability and low-grade chronic inflammation.

Here we found a shift in energy metabolism towards continuous glucose utilization as an energy source with significantly greater TCA cycle intermediate metabolites in the serum of *Zip14* KO mice. We have previously shown that in the absence of *Zip14*, hepatic glucose uptake was increased along with greater glycogen stores ⁵³ suggesting a greater amount of glucose available for energy production. The liver may have two ways of responding to this: increasing inefficient metabolism (Fig. 5-increased TCA cycle metabolites) and exporting these calories in the form of fat for deposition in peripheral tissues ²⁰. Energy metabolism is also affected by inflammation. During chronic inflammation, adaptive immune cells, specifically T lymphocytes, go through metabolic reprogramming in favor of oxidative phenotype ⁵⁴, in agreement with low-grade chronic inflammation and increased TCA cycle intermediates in *Zip14* KO mice. We have previously shown mitochondrial localization of ZIP14 in hepatocytes ¹⁸. In addition to metabolic reprogramming in adaptive immune cells of *Zip14* KO mice, the shift in energy metabolism could also be due to ZIP14-mediated regulation of mitochondrial metal concentration and the function of corresponding metalloenzymes. It is important to acknowledge that further studies with tissue/cell type-specific conditional *Zip14* KO mice along with dietary interventions for Zn and manganese are warranted to test these possible roles for ZIP14.

Methods

Animals and treatments

Animals. Heterozygous (*Zip14*^{+/-}) mice are a generous gift of Dr. Robert J. Cousins. Heterozygous *Zip14*^{+/-} on the mixed background of C57BL6 and 129S5 mice were bred to obtain homozygous whole-body (WB) *Zip14* KO (*Zip14*^{-/-}) and WT (*Zip14*^{+/+}) mice ⁵⁵. The mice were maintained using standard rodent husbandry and received a commercial, irradiated chow diet (Harlan Teklad 7912; Indianapolis, IN, with 60 mg Zn/kg as ZnO) and tap water. Age-matched mice from both sexes were used as young adults (8–16 wk of age). Euthanasia was through exsanguination by cardiac puncture under isoflurane anesthesia. Protocols were approved by the Cornell University Institutional Animal Care and Use Committees. **Metabolic phenotyping.** Metabolic parameters of the mice were measured by the Comprehensive Laboratory Animal Monitoring System (CLAMS) by using Promethion (Sable Systems). Mice were transferred to the CLAMS metabolic cages and

allowed to acclimate for 2-3 days with free access to food and water. Following the acclimation, measurements of VO₂ and VCO₂ gas exchanges and food and water consumption were collected for five days. The data were analyzed using Expedata software system (v1.9.27), Macro interpreter (v2.40), and then CalR software (v.1.2). **Body composition.** Measurements were conducted in awake animals by time-domain nuclear magnetic resonance using the Minispec LF65 Body Composition Mice Analyzer. **Fasting blood glucose.** Following overnight fasting, blood glucose levels were measured by tail bleeding using the OneTouch® Ultra Blood glucose monitoring system. **Antibiotics treatment.** Antibiotics (Abx) (Ampicillin 1 g/L, neomycin 1 g/L, gentamycin 500 mg/L; penicillin 100U/L, vancomycin hydrochloride 1g/L, metronidazole 500mg/L) were given in the drinking water, ad libitum as described previously^{56,57} for three weeks. Abx water was replaced with control water 4 hours before fecal transplant. **Anti-fungal treatment.** Drinking water was supplemented with fluconazole (0.5 mg/mL, Sigma) for ten days^{58,59}. **Fecal transplant.** Fecal pellets were collected from co-caged WT and WB KO mice to prepare the fecal slurry. 0.2 g of the fecal pellet was homogenized in 1mL of phosphate-buffered saline (PBS) using bullet blender (Next Advance). Following centrifuging at 800 x g for 3 min and filtering by 30 µm cell strainer, antibiotics-treated mice were inoculated for three consecutive days with 0.2 ml of the fecal slurry^{57,60}. Following conventionalization, metabolic phenotypes were assessed weekly by measuring fasting blood glucose and body composition by NMR. At the end of the three weeks, blood and tissue were collected for further biochemical analysis. **Permeability Assay.** Following morning fasting (removal of food and water for 4 hours), mice were given FITC-dextran (60 mg/100 g FITC-dextran 4) (Sigma-MW 3000-5000) via oral gavage. Blood was collected into EDTA tubes by cardiac puncture to obtain plasma at two hours post-gavage. FITC-dextran levels in plasma were measured fluorometrically at 485 nm emission/528 nm excitation.

Analytical methods

Metal Assays. The entire excised small intestine was cut into two sections as proximal (the first 11cm) and distal (rest) and perfused with a metal chelating buffer (10 mM EDTA, 10 mM HEPES, and 0.9% NaCl₂) (equal volume/section). Similarly, the colon was excised and perfused with the above buffer. The contents of the intestine and colon were centrifuged at 3000xg to obtain the lumen (supernatant) and digesta (pellet). Aliquots of tissue, lumen, digesta, and the fecal pellets were digested at 95°C overnight in HNO₃ to measure metal concentrations using Microwave Plasma-Atomic Emission Spectrometry (MP-AES). Normalization was to tissue weight or volume (intestinal contents).

Gut microbiome and serum metabolite profiling

Fresh fecal pellets were collected from mice that were co-caged for one month prior. **DNA extraction and sequencing.** DNA samples were extracted from the mice fecal samples using DNeasy PowerSoil Kit (QIAGEN, Inc.) according to the manufacturer's instruction. Region V4 of the 16S rRNA gene was amplified using the universal primers 515F and GoLay-barcoded 806R and the PCR program as previously described⁶¹. Samples were amplified in duplicate with the following thermocycler protocol: hold at 94°C for 3 min; 30 cycles of 94°C for 45 s, 50°C for 1 min, 72°C for 1.5 min; and hold at 72°C for 10 min, and the duplicate final amplified products were pooled. Amplicons were further purified using Mag-Bind® RxnPure Plus (Omega Bio-tek, Inc.) and quantified with NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific), and 150 ng of amplicons from each sample were pooled and paired-end sequenced (2 x 250bp) on an Illumina MiSeq instrument. **16S rRNA gene sequencing analysis.** Reads were demultiplexed using QIIME v1.9.1 (split_libraries_fastq.py) before denoising and processing with Divisive Amplicon Denoising Algorithm 2 (DADA2)⁶². Taxonomy was assigned by mapping with the Greengenes 16S rRNA Gene Database⁶³. The QIIME output data was then imported to RStudio (Version 1.0.136) with the Bioconductor package phyloseq⁶⁴ to normalize and plot the input data. All statistical analyses were done in R using the vegan⁶⁵ and DESeq2⁶⁶. A Benjamini–Hochberg FDR of 0.01 was used as the cutoff for statistical significance for heatmap plotting unless otherwise noted. **Saccharomyces cerevisiae detection.** For isolating fungal DNA, an equal amount of colon digesta from each mouse were suspended in 50 mM Tris buffer (pH 7.5) supplemented with one mM EDTA, 0.2% b-mercaptoethanol and 1000 units/ml of lyticase (Sigma), incubated at 37°C for 30 min to disrupt fungal cells as described^{58,59} prior to processing through the QIAamp Fast DNA Stool Mini Kit (Qiagen). The quantitative PCR (qPCR) reactions with an equal amount of DNA were set up using power track SYBR green master mix and run in Roche Lightcycler 480-II RT-PCR system. The *S. cerevisiae*-specific primer sequences were forward 5-AGGAGTGC GGTTCTTTG-3 and reverse 5-TACTTACCGAGGCAAGCTACA-3⁶⁷. **Serum untargeted metabolomics profiling.** Gas chromatography (GC) coupled to time-of-flight mass spectrometry (TOF-MS) was performed using Gerstel CIS4 –with dual MPS Injector/ Agilent 6890 GC-Pegasus III TOF MS by UC Davis West Coast Metabolomics Center. Statistical and pathway analyses were performed using MetaboAnalyst 5.0³¹. Data were normalized to the median to account for variance between samples and generalized logarithm transformed to account for differences in metabolite abundance. Metabolites with a fold change >1.5 and p < 0.05 were considered to be significantly different for pathway analysis. Pathway analysis was conducted using KEGG human metabolic pathways. Disease signature analysis was conducted using metabolite sets reported in human blood.

Isolation of Enterocytes and Western blot analysis

Luminal contents of the proximal intestine were flashed by PBS prior to evert them using bamboo sticks with pointed ends. Everted intestines were incubated in PBS with 1.5 mM EDTA solution for 20 min and released enterocytes were collected by centrifugation at 4°C at 500 x g for 5 min. Following two washes, the pellets were resuspended in lysis buffer (Tris-HCl, 137mM NaCl, 10% glycerol, 1% Triton X-100, 2 mM EDTA) with protease and phosphatase inhibitors (Thermo Scientific) added along with the PMSF (Sigma-Aldrich). During enterocyte separation, all the buffers were supplemented with protease inhibitors (AGScientific). Solubilized proteins were separated by 10% SDS-PAGE. Visualization was done by chemiluminescence (SuperSignal, Thermo Fisher) and digital imaging (Protein Simple). Rabbit anti-mouse ZIP14 antibody was custom made by Genscript. Antibodies for ZIP4, ZnT1, and GAPDH were obtained from Proteintech (20625), Novusbio (BP-94196), and Cell Signaling (14C10), respectively.

Statistical Analyses

Data are presented as means ± SE. Significance was assessed by Student's t-test for single comparisons. Statistical significance was set at $P < 0.05$. Analyses were performed using GraphPad Prism. The statistical criteria for metabolomics analysis and 16S rRNA gene sequencing analysis were included in the methods.

References

1. Present Knowledge in Nutrition - 11th Edition. Available at: <https://www.elsevier.com/books/present-knowledge-in-nutrition/marriott/978-0-323-66162-1>. (Accessed: 23rd June 2021)
2. King, J. C., Shames, D. M. & Woodhouse, L. R. Zinc homeostasis in humans. *J. Nutr.* **130**, 1360S–6S (2000).
3. Cousins, R. J. Theoretical and practical aspects of zinc uptake and absorption. *Adv. Exp. Med. Biol.* **249**, 3–12 (1989).
4. Hoadley, J. E., Leinart, A. S. & Cousins, R. J. Kinetic analysis of zinc uptake and serosal transfer by vascularly perfused rat intestine. *Am. J. Physiol. Liver Physiol.* **252**, G825–G831 (1987).
5. McMahon, R. J. & Cousins, R. J. Regulation of the zinc transporter ZnT-1 by dietary zinc. *Proc. Natl. Acad. Sci.* **95**, 4841–4846 (1998).
6. Guthrie, G. J. et al. Influence of ZIP14 (slc39A14) on intestinal zinc processing and barrier function. *Am. J. Physiol. Gastrointest. Liver Physiol.* **308**, G171-8 (2015).

7. Dufner-Beattie, J., Langmade, S. J., Wang, F., Eide, D. & Andrews, G. K. Structure, function, and regulation of a subfamily of mouse zinc transporter genes. *J. Biol. Chem.* **278**, 50142–50 (2003).
8. Liuzzi, J. P., Bobo, J. A., Cui, L., McMahon, R. J. & Cousins, R. J. Zinc transporters 1, 2 and 4 are differentially expressed and localized in rats during pregnancy and lactation. *J. Nutr.* **133**, 342–51 (2003).
9. Dufner-Beattie, J., Kuo, Y.-M., Gitschier, J. & Andrews, G. K. The adaptive response to dietary zinc in mice involves the differential cellular localization and zinc regulation of the zinc transporters ZIP4 and ZIP5. *J. Biol. Chem.* **279**, 49082–90 (2004).
10. Ohashi, W. & Fukada, T. Contribution of zinc and zinc transporters in the pathogenesis of inflammatory bowel diseases. *Journal of Immunology Research* **2019**, (2019).
11. Geiser, J., Venken, K. J. T., de Lisle, R. C. & Andrews, G. K. A mouse model of acrodermatitis enteropathica: Loss of intestine zinc transporter ZIP4 (Slc39a4) disrupts the stem cell niche and intestine integrity. *PLoS Genet.* **8**, (2012).
12. Ohashi, W. *et al.* Zinc Transporter SLC39A7/ZIP7 Promotes Intestinal Epithelial Self-Renewal by Resolving ER Stress. *PLOS Genet.* **12**, e1006349 (2016).
13. Podany, A. B., Wright, J., Lamendella, R., Soybel, D. I. & Kelleher, S. L. ZnT2-Mediated Zinc Import Into Paneth Cell Granules Is Necessary for Coordinated Secretion and Paneth Cell Function in Mice. *CMGH* **2**, 369–383 (2016).
14. Li, D. *et al.* A Pleiotropic Missense Variant in SLC39A8 Is Associated With Crohn's Disease and Human Gut Microbiome Composition. *Gastroenterology* **151**, 724–732 (2016).
15. Sunuwar, L. *et al.* Pleiotropic ZIP8 A391T implicates abnormal manganese homeostasis in complex human disease. *JCI Insight* **5**, (2020).
16. Nakata, T. *et al.* A missense variant in SLC39A8 confers risk for Crohn's disease by disrupting manganese homeostasis and intestinal barrier integrity. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 28930–28938 (2020).
17. Girijashanker, K. *et al.* Slc39a14 gene encodes ZIP14, a metal/bicarbonate symporter: similarities to the ZIP8 transporter. *Mol. Pharmacol.* **73**, 1413–23 (2008).
18. Aydemir, T. B., Troche, C., Kim, M.-H. & Cousins, R. J. Hepatic ZIP14-mediated Zinc Transport Contributes to Endosomal Insulin Receptor Trafficking and Glucose Metabolism. *J. Biol. Chem.* **291**, 23939–23951 (2016).
19. Troche, C., Aydemir, T. B. & Cousins, R. J. Zinc transporter Slc39a14 regulates inflammatory signaling associated with hypertrophic adiposity. *Am. J. Physiol. Endocrinol.*

- 394 *Metab.* **310**, aipendo.00421.2015 (2015).
- 395 20. Aydemir, T. B. *et al.* Zinc Transporter ZIP14 Functions in Hepatic Zinc, Iron and Glucose
396 Homeostasis during the Innate Immune Response (Endotoxemia). *PLoS One* **7**, e48679
397 (2012).
- 398 21. Aydemir, T. B. & Cousins, R. J. The Multiple Faces of the Metal Transporter ZIP14
399 (SLC39A14). *J. Nutr.* **148**, 174–184 (2018).
- 400 22. Aydemir, T. B. *et al.* Metal Transporter *Zip14* (*Slc39a14*) Deletion in Mice Increases
401 Manganese Deposition and Produces Neurotoxic Signatures and Diminished Motor
402 Activity. *J. Neurosci.* **37**, 5996–6006 (2017).
- 403 23. Aydemir, T. B. *et al.* Intestine-specific deletion of metal transporter *Zip14* (Slc39a14)
404 causes brain manganese overload and locomotor defects of manganism. *Am. J. Physiol.*
405 *- Gastrointest. Liver Physiol.* **318**, G673–G681 (2020).
- 406 24. Starke, I. C., Pieper, R., Neumann, K., Urgan Zentek, J. & Vahjen, W. The impact of high
407 dietary zinc oxide on the development of the intestinal microbiota in weaned piglets.
408 doi:10.1111/1574-6941.12233
- 409 25. Festi, D. *et al.* Gut microbiota and metabolic syndrome. *World J. Gastroenterol.* **20**,
410 16079 (2014).
- 411 26. Lopez, C. A. & Skaar, E. P. Cell Host & Microbe The Impact of Dietary Transition
412 Metals on Host-Bacterial Interactions. *Cell Host Microbe* **23**, 737–748 (2018).
- 413 27. KL, G., BP, A. & EMM, Q. The microbiome and inflammatory bowel disease. *J. Allergy*
414 *Clin. Immunol.* **145**, 16–27 (2020).
- 415 28. Morgan, X. C. *et al.* Dysfunction of the intestinal microbiome in inflammatory bowel
416 disease and treatment. *Genome Biol.* 2012 139 **13**, 1–18 (2012).
- 417 29. Yow, G. Y., Uo, T., Yoshimura, T. & Esaki, N. D-Amino acid-N-acetyltransferase of
418 *Saccharomyces cerevisiae*: A close homologue of histone acetyltransferase Hpa2p acting
419 exclusively on free D-amino acids. *Arch. Microbiol.* **182**, 396–403 (2004).
- 420 30. Sampath, V. *et al.* Biochemical characterization of Hpa2 and Hpa3, two small closely
421 related acetyltransferases from *saccharomyces cerevisiae*. *J. Biol. Chem.* **288**, 21506–
422 21513 (2013).
- 423 31. Pang, Z. *et al.* MetaboAnalyst 5.0: narrowing the gap between raw spectra and functional
424 insights. *Nucleic Acids Res.* (2021). doi:10.1093/nar/gkab382
- 425 32. Wang, M. *et al.* Metabonomic strategy for the detection of metabolic effects of probiotics
426 combined with prebiotic supplementation in weaned rats†. *RSC Adv.* **8**, 5042–5057
427 (2018).

33. Küry, S. *et al.* Identification of SLC39A4, a gene involved in acrodermatitis enteropathica. *Nat. Genet.* **31**, 239–40 (2002).
34. Aydemir, T. B. *et al.* Metal transporter Zip14 (Slc39a14) deletion in mice increases manganese deposition and produces neurotoxic signatures and diminished motor activity. *J. Neurosci.* **37**, (2017).
35. Jenkitkasemwong, S. *et al.* SLC39A14 Is Required for the Development of Hepatocellular Iron Overload in Murine Models of Hereditary Hemochromatosis. *Cell Metab.* **22**, 138–150 (2015).
36. Erdman, J. W., MacDonald, I. A. & Zeisel, S. H. *Present Knowledge in Nutrition: Tenth Edition. Present Knowledge in Nutrition: Tenth Edition* (Wiley-Blackwell, 2012).
doi:10.1002/9781119946045
37. Ross, A. C., Caballero, B. H., Cousins, R. J., Tucker, K. L. & Ziegler, T. R. *Modern nutrition in health and disease.* (Wolters Kluwer Health/Lippincott Williams & Wilkins, 2014).
38. WHO. The World health report: 2002: Reducing the risks, promoting healthy life. *Therapie* **52**, 569–571 (2002).
39. Skrovanek, S. Zinc and gastrointestinal disease. *World J. Gastrointest. Pathophysiol.* **5**, 496–513 (2014).
40. Ross, A. Catharine, Cousins, Robert J. and Caballero, B. *et. al. Modern Nutrition in Health and Disease - Wolters Kluwer Health, Lippincott Williams & Wilkins - Literati by Credo.* (2014).
41. Zackular, J. P. *et al.* Dietary zinc alters the microbiota and decreases resistance to *Clostridium difficile* infection. *Nat. Med.* **22**, 1330–1334 (2016).
42. Yu, T. *et al.* Dietary high zinc oxide modulates the microbiome of ileum and colon in weaned piglets. *Front. Microbiol.* **8**, 825 (2017).
43. Reed, S. *et al.* Chronic zinc deficiency alters chick gut microbiota composition and function. *Nutrients* **7**, 9768–9784 (2015).
44. Donia, M. S. & Fischbach, M. A. Small Molecules from the Human Microbiota.
doi:10.1126/science.1254766
45. Krautkramer, K. A., Fan, J. & Bäckhed, F. Gut microbial metabolites as multi-kingdom intermediates. *Nat. Rev. Microbiol.* **19**, (2021).
46. Wikoff, W. R. *et al.* Metabolomics analysis reveals large effects of gut microflora on mammalian blood metabolites. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 3698–3703 (2009).
47. Roager, H. M. & Licht, T. R. Microbial tryptophan catabolites in health and disease.

Nature Communications **9**, 1–10 (2018).

48. Agus, A., Planchais, J. & Sokol, H. Gut Microbiota Regulation of Tryptophan Metabolism in Health and Disease. *Cell Host and Microbe* **23**, 716–724 (2018).
49. Eide, D. J. Homeostatic and adaptive responses to zinc deficiency in *saccharomyces cerevisiae*. *J. Biol. Chem.* **284**, 18565–18568 (2009).
50. Siva, S., Rubin, D. T., Gulotta, G., Wroblewski, K. & Pekow, J. Zinc deficiency is associated with poor clinical outcomes in patients with inflammatory bowel disease. *Inflamm. Bowel Dis.* **23**, 152–157 (2017).
51. Chiaro, T. R. *et al.* A member of the gut mycobiota modulates host purine metabolism exacerbating colitis in mice. *Sci. Transl. Med.* **9**, (2017).
52. Manocha, M. & Khan, W. I. Serotonin and GI disorders: An update on clinical and experimental studies. *Clinical and Translational Gastroenterology* **3**, (2012).
53. Aydemir, T. B., Troche, C., Kim, M.-H. & Cousins, R. J. Hepatic ZIP14-mediated zinc transport contributes to endosomal insulin receptor trafficking and glucose metabolism. *J. Biol. Chem.* **291**, (2016).
54. Moro-García, M. A., Mayo, J. C., Sainz, R. M. & Alonso-Arias, R. Influence of Inflammation in the Process of T Lymphocyte Differentiation: Proliferative, Metabolic, and Oxidative Changes. *Front. Immunol.* **9**, 339 (2018).
55. Aydemir, T. B., Sitren, H. S. & Cousins, R. J. The zinc transporter Zip14 influences c-met phosphorylation and hepatocyte proliferation during liver regeneration in mice. *Gastroenterology* **142**, (2012).
56. Ramakrishnan, S. K. *et al.* Intestinal non-canonical NFκB signaling shapes the local and systemic immune response. *Nat. Commun.* **10**, (2019).
57. Das, N. K. *et al.* Microbial Metabolite Signaling Is Required for Systemic Iron Homeostasis. *Cell Metab.* **31**, 115-130.e6 (2020).
58. Iliev, I. D. *et al.* Interactions between commensal fungi and the C-type lectin receptor dectin-1 influence colitis. *Science (80-.).* **336**, 1314–1317 (2012).
59. Jiang, T. T. *et al.* Commensal Fungi Recapitulate the Protective Benefits of Intestinal Bacteria. *Cell Host Microbe* **22**, 809-816.e4 (2017).
60. Johnson, E. L. *et al.* Sphingolipids produced by gut bacteria enter host metabolic pathways impacting ceramide levels. *Nat. Commun.* **11**, (2020).
61. Caporaso, J. G. *et al.* Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 4516–4522 (2011).
62. Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon

data. *Nat. Methods* 2016 137 **13**, 581–583 (2016).

63. McDonald, D. *et al.* An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME J.* 2012 63 **6**, 610–618 (2011).

64. McMurdie, P. J. & Holmes, S. phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. *PLoS One* **8**, e61217 (2013).

65. Oksanen, J. *et al.* Package ‘vegan’ Title Community Ecology Package Version 2.5-7. (2020).

66. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).

67. Chang, H. W. *et al.* Quantitative real time PCR assays for the enumeration of *Saccharomyces cerevisiae* and the *Saccharomyces sensu stricto* complex in human feces. *J. Microbiol. Methods* **71**, 191–201 (2007).

68. Guthrie, G. J. G. J. G. J. *et al.* Influence of ZIP14 (slc39A14) on intestinal zinc processing and barrier function. *Am. J. Physiol. - Gastrointest. Liver Physiol.* **308**, G171–G178 (2015).

FIGURES.

Figure 1. Altered gastrointestinal Zn homeostasis in *Zip14* KO mice. Zn concentrations in gastrointestinal tissues (A) and the luminal contents (B) were measured by MP-AES. Values are means $\pm$ SE; n = 16-19 (both female and male mice were included). Student's t-test for WT vs. WB-KO comparison. C) Abundance of Zn transporters, ZIP14, ZIP4, and ZnT1, in isolated enterocytes was shown by western blot analysis.

Figure 2. Decreased gut microbial diversity in the WB-Zip14 KO mice. A, B) WB-Zip14 KO displayed decreased α -diversity, as shown by 16S rRNA gene sequencing. Fold change of >2 and p-value < 0.05 were considered significantly different. C) The most significant changes at the family and genus levels (WB-KO/WT) (n=7-8). The details of the analysis are provided in the methods section.

Figure 3. The ZIP14-shaped microbiome can influence intestinal permeability and metabolic phenotypes in the host. A) Following three weeks of antibiotics treatment, mice have received a fecal transplant from either WT (WTCv) or WB-KO (KOCv) mice. B) Plasma FITC-Dextran concentrations were measured fluorometrically. B) Fasting blood glucose was measured by tail bleeding using the OneTouch® Ultra Blood glucose monitoring system. Percent body fat (C) and percent lean mass (D) were assessed by nuclear magnetic resonance. Values are means $\pm$ SE; $n = 5-7$ (both female and male mice were included). Student's t-test.

Figure 4. Serum untargeted metabolomics profiling of *Zip14* KO and WT mice. Serum metabolites were measured Gas chromatography (GC) coupled to time-of-flight mass spectrometry (TOF-MS) (n=6 per group). A) The PCA score plot of serum samples from WT and WB-KO mice. B) Volcano plot. C) The serum metabolites with the known identity were statistically different. D) *S. Cerevisiae* DNA was quantified by qPCR in colonic digesta from either WT and WB-KO or WT mice that received fecal transplant either from WT (WTCv) or KO (KOCv) mice. Values are means $\pm$ SE; n = 3. Student's t-test. E) Following ten days of antifungal (fluconazole) treatment, plasma FITC-Dextran concentrations were measured fluorometrically. Values are means $\pm$ SE; n = 6-9. Student's t-test.

Figure 5.. Pathway and enrichment analyses using MetaboAnalyst. For the significantly different metabolites (details in method section), pathway and enrichment analyses were conducted using MetaboAnalyst. A) Pathway analysis was conducted using KEGG human metabolic pathways. B) Disease signature analysis was conducted using metabolite sets reported in human blood.

Figure 6. Altered energy metabolism in the WB-KO mice. Following 2-3 days of acclimation, metabolic parameters of the mice were measured by the Comprehensive Laboratory Animal Monitoring System (CLAMS). Measurements of VO₂ (A) and VCO₂ (B) gas exchanges were collected for five days. C) The respiratory exchange rate (RER) was obtained based on these values. Values are means $\pm$ SE; n= 4 (both female and male mice were included). Student t-test.

Figure 7. Model for a mechanism of intestinal permeability in *Zip14* KO mice. Deletion of the host gene encoding metal transporter, *Slc39a14/Zip14*, generates Zn-deficient epithelial and luminal content in the intestine. Reduced Zn concentrations in the enterocytes were shown to dysregulate tight junction proteins (TJP), subsequently increasing intestinal permeability⁶⁸. Reduced Zn concentrations in the lumen lower bacterial diversity and give the opportunity to *S. cerevisiae* overgrowth, further contributing to intestinal permeability. Consequently, altered host and microbial metabolites, inflammation, and shift in energy metabolism are found in the host.