Draft version 2021/12/23, submitted for peer review

Pharmacological blockade of muscle afferents and perception of effort: a systematic review with meta-analysis

Maxime Bergevin^{1,2}, James Steele³, Marie Payen de la Garanderie^{1,2}, Camille Feral-Basin^{1,2}, Samuele M. Marcora⁴, Pierre Rainville^{2,5}, Jeffrey G. Caron¹⁻⁶ and Benjamin Pageaux^{1,2}

¹ École de kinésiologie et des sciences de l'activité physique (EKSAP), Faculté de médecine, Montreal, Canada

² Centre de recherche de l'Institut universitaire de gériatrie de Montréal (CRIUGM), Montreal, Canada

³ School of Sport, Health and Social Sciences, Southampton, United Kingdom

⁴ Department of Biomedical and Neuromotor Sciences (DiBiNeM), University of Bologna, Bologna, Italy

⁵ Département de stomatologie, Faculté de médecine dentaire, Université de Montréal, Montreal, Canada

⁶ Centre de recherche interdisciplinaire en réadaptation du Montréal métropolitain, Montreal, Canada

Corresponding author :

Benjamin Pageaux

École de kinésiologie et des sciences de l'activité physique (EKSAP)

Centre d'éducation physique et des sports (CEPSUM)

Faculté de médecine, Université de Montréal

2100, boul. Édouard-Montpetit

Montréal, QC H3T 1J4

Canada

+ 1 514-343-6111

benjamin.pageaux@umontreal.ca

29
 30 **ORCID :**
 31 Maxime Bergevin: [https://orcid.org/ 0000-0002-0343-8900](https://orcid.org/0000-0002-0343-8900)
 32 James Steele : <https://orcid.org/0000-0002-8003-0757>
 33 Camille Féral-Basin : [https://orcid.org/ 0000-0003-0045-8268](https://orcid.org/0000-0003-0045-8268)
 34 Samuele M. Marcora : <https://orcid.org/0000-0002-1570-7936>
 35 Pierre Rainville: <https://orcid.org/0000-0001-9801-757X>
 36 Benjamin Pageaux : <https://orcid.org/0000-0001-9302-5183>

37 **Statement and Declarations**

38 **Funding statement.** MB is supported by the Canadian Institutes of Health Research through the Canada
 39 Graduate Scholarships – Master’s Frederick Banting and Charles Best grant, the “Formation de maîtrise”
 40 scholarship from the Fond de recherche en santé du Québec and an MSc scholarship from the Centre de
 41 recherche de l’Institut universitaire de gériatrie de Montréal (CRIUGM). BP’s research is supported by
 42 the Natural Sciences and Engineering Research Council of Canada—Discovery Grants Program.

43 **Competing interests.** The authors have no conflict of interest to disclose in respect to this manuscript.

44 **Availability of data and material.** All data and materials are available within the manuscript or in
 45 supplemental materials.

46 **Code availability.** Not applicable

47

48

49 **Keywords**

50 Perceived exertion, sense of effort, motor drive, motor command, neural feedback, central command,
 51 peripheral feedback, opioid, spinal anesthesia, epidural anesthesia, impaired afference, exercise,
 52 endurance

53

Abstract

The perception of effort (PE) provides information on task difficulty and influences physical exercise regulation and human behavior. This perception differs from other-exercise related perceptions such as pain. There is no consensus on the role of group III-IV muscle afferents as a signal processed by the brain to generate PE. The aim of this meta-analysis was to investigate the effect of pharmacologically blocking muscle afferents on the PE. Six databases were searched to identify studies measuring the ratings of perceived effort (RPE) during physical exercise, with and without pharmacological blockade of muscle afferents. Articles were coded based on the operational measurement used to distinguish studies in which PE was assessed specifically (*effort dissociated*) or as a composite experience including other exercise-related perceptions (*effort not dissociated*). Articles that did not provide enough information for coding were assigned to the *unclear* group. The *effort dissociated* group ($n=6$) demonstrated a slight RPE increase with reduced muscle afferents feedback (standard mean change raw (SMCR), 0.39; 95%CI, 0.13 to 0.64). The group *effort not dissociated* ($n=2$) did not reveal conclusive results (SMCR, -0.29; 95%CI, -2.39 to 1.8). The group *unclear* ($n=8$) revealed a slight RPE decrease with reduced muscle afferents feedback (SMCR, -0.27; 95%CI, -0.50 to -0.04). The heterogeneity in results between groups reveals that the inclusion of other perceptions than effort in its rating influences the RPE scores reported by the participants. The absence of decreased RPE in the *effort dissociated* group suggests that muscle afferents feedback is not a sensory signal generating PE.

250 words

Key points (2-3 sentences summarizing, in non-technical language, the key findings/implications of the manuscript)

To date, there is no consensus on the neurophysiological signal processed by the brain to generate the perception of effort. Following a systematic search in six databases, this meta-analysis reveals that reducing afferent feedback from the working muscles via epidural anesthesia does not reduce perception of effort. This systematic review emphasizes the role of the corollary discharge associated with the motor command as the signal processed by the brain to generate the perception of effort.

1. Introduction

During physical exercise, the perception of effort provides information on the intensity and difficulty of the task being performed. The perception of effort is involved in the regulation of human behavior and influences how the nervous system selects a given movement amongst a myriad of possibilities [1, 2]. The perception of effort is exacerbated in the presence of fatigue [3] and various pathologies such as chronic fatigue syndrome [4, 5], stroke [6] and cancer [7]. This perception is used to prescribe and monitor exercise in both rehabilitation programs [8, 9] and athletic training [10-12]. Despite the growing interest in this perception, to date, researchers have failed to reach a consensus on the signal(s) processed by the brain leading to its generation [13-15].

One popular model amongst exercise physiologists, referred to as the afferent feedback model, suggests that the feedback originating from the peripheral organs active during physical exercise (i.e., skeletal muscles, heart, lungs) is processed by the central nervous system to generate the perception of effort [16, 17]. Notably, authors suggested that feedback from group III-IV muscle afferents plays an important role in the generation of the perception of effort [16, 18]. The ratings of perception of effort intensity would then be predicted to increase with higher discharge rates of the muscle afferents accompanying intense exercise [19, 20]. In contrast, a popular model amongst neuroscientists, cardiovascular physiologists and physiologists interested in kinesthesia is the corollary discharge model. This model proposes that the perception of effort is generated by the processing of a copy of the central motor command, named the corollary discharge [21-24]. In this model, an increase in the magnitude of the central motor command should result in an increase in the perception of effort intensity [21, 25]. It is important to note that this model does not bar peripheral contributions to the regulation of central commands during voluntary movement, but states that central processing of afferent feedback does not generate the perception of effort and that effort could be perceived in the absence of afferent feedback [13, 14, 26, 27]. For example, any peripheral or central mechanism able to alter the muscle force production capacity [21, 28] or the corticospinal excitability – the ease with which the central motor command is relayed to the motoneurons [29] – may modulate the perception of effort by increasing or decreasing the magnitude of the central motor command needed to sustain a given level of performance [30].

A powerful technique to gain insights into the neurophysiology of the perception of effort and test the existing models is by pharmacologically blocking muscle afferents while monitoring the perception of effort [e.g., 26, 31, 32, 33]. Since the afferent feedback model considers group III-IV muscle afferents as the signal processed by the brain to generate the perception of effort, pharmacologically blocking muscle afferents should decrease the ratings of perceived effort. On the other hand, observing stable or increased

ratings of perceived effort in the presence of reduced muscle afferents feedback would support a centrally generated perception of effort. Intrathecal and epidural injection of anesthetics or analgesics, such as lidocaine or fentanyl, has traditionally been used to investigate the role of group III-IV muscle afferents and the motor command in cardio-respiratory responses to exercises in both healthy [e.g., 32] and symptomatic participants [e.g., 34] as well as in human performance during endurance exercises [35]. In these studies, participants performed an exercise protocol, usually cycling or isolated knee exercises, with and without intact feedback from group III-IV muscle afferents. While the primary variables of interest were the cardio-respiratory responses to the tasks, these studies often measured the perception of effort as a secondary or tertiary variable. Interestingly, there are conflicting results from these studies. In the presence of pharmacological blockade of muscle afferents, several authors observed a decrease [e.g., 36] in the ratings of perceived effort while others observed no difference [e.g., 33] or an increase [e.g., 37] when compared to a sham or control intervention. This heterogeneity is also found in patients with cardiovascular diseases [e.g., 34, 38]. To the best of our knowledge, only one published article has narratively reviewed the use of pharmacological blockade to explore the neurophysiological mechanisms underlying the perception of effort [26], and a systematic approach has yet to be conducted.

The conflicting findings on the effects of pharmacologically blocking muscle afferents on the perception of effort may be explained by differences in its operational definitions, leading to inconsistencies in the instructions provided to the participants on how to quantify the perception of effort. In his seminal work, Borg defined the perception of effort as how *heavy* and *laborious* a physical task is [39, 40]. However, he also mentioned that this perception results from the integration of various peripheral factors, including the organs of circulation and respiration, the muscles, the skin, and the joints. Since this first definition and associated description provided by Borg, two lines of research have investigated the perception of effort in exercise sciences (see figure 1). A first line of research considers effort as a construct encompassing a mix of exercise-related perceptions [17, 41]. This approach results from the original description proposed by Borg, and later authors supplemented this definition with the notions of fatigue and discomfort: *the subjective intensity of effort, strain, discomfort, and/or fatigue that is experienced during physical exercise* [17]. However, experimental data demonstrated that the perception of effort can be dissociated from other exercise-related perceptions, such as pain [42, 43], discomfort [44-46], muscle tension [47, 48] and fatigue [49, 50]. In line with the aforementioned evidence, a second line of research considers effort as a construct dissociated from other perceptions. This approach follows Borg's original definition conceptualizing the perception of effort as one's appreciation of the difficulty of a task (how *hard* it is). For instance, Preston and Wegner [51] described the perception of effort as the *feeling of difficulty and labor experienced during exertion*. In 2010, Marcora proposed that the perception of effort is the *conscious sensation of how hard*,

heavy and strenuous a physical task is [52]. More recently, Steele defined the perception of effort as the *perception of current task demands relative to the perception of capacity to meet those demands* [15]. In light of the disparate definitions proposed in the literature, it appears crucial to consider the definition used to investigate the perception of effort when interpreting such data. However, to the best of our knowledge, such consideration has not been made in the literature when discussing the signal(s) generating perception of effort.

In this context, the aim of this systematic review with meta-analysis was to explore the impact of pharmacologically blocking muscle afferents on the perception of effort during physical tasks. To explore whether the inclusion or not of other exercise-related perceptions influence the quantification of perceived effort, a qualitative approach was used to group included studies by their definition prior to the statistical analyses.

****Please insert figure 1****

2. Methods

The present review was conducted and is reported as per the Preferred Reporting Items for Systematic Reviews and Meta-analyses statement (PRISMA; [53]). All methods were pre-specified in a protocol registered on PROSPERO (CRD401913921) prior to the screening process and any deviations from the pre-registered methods are noted throughout.

2.1 Search

The following electronic databases were searched to identify studies: MEDLINE, EMBASE, CINAHL, SPORTDiscuss, Web of Science and PsycINFO. To ensure the inclusion of recently published articles, the search was conducted on three separate occasions (April 2019 and March 2020 and October 2021). A search strategy was developed for MEDLINE and adapted for each database. The first of the two concepts included was the perception of effort and included the following terms: *perception of effort*, *sense of effort*, *perceived exertion*, *central motor command*, *central motor drive*. Because the perception of effort has been used by some in the neuroscience literature as an index of the central motor command during exercise, the latter has been included in this concept. The second concept related to the pharmacological blockade and included the following terms: *epidural anesthesia*, *spinal anesthesia*, *neural blockade*, *nerve block*, *sensory dysesthesia*, *fentanyl*, *lidocaine*, *bupivacaine*, *muscle afferent*, *neural feedback*, *afferent feedback*, and *group III-IV*. It is noteworthy that, despite using the same search strategy in all three instances, PsycINFO returned fewer articles each time the database was scanned (65 against 30 against 27 articles) and that 47 of the 65 original articles were not found in the subsequent searches. Additionally, no limitation to the publication date was set during the search. The complete search strategy for every scanned database is available in the supplementary material S1.

2.2 Article inclusion

Eligibility criteria were defined accordingly with the PICOS model. Articles qualified for inclusion if they met the following criteria: 1) *population*: intervention was done on human participants, 2) *intervention*: consisted of a blockade of spinal afferents by epidurally or intrathecally injecting a local anesthetic or analgesics, 3) *comparators*: intervention was compared against a control or placebo condition, 4) *outcome*: the perception of effort was a primary, secondary, or tertiary outcome, and 5) *study design*: repeated-measure designs. Additionally, only articles published in peer-reviewed journals and written in English were retained. We also opted not to include self-paced protocols to allow comparison between studies. This decision is based on the necessity to compare the intensity of the perception of effort when the task demand (e.g., power output) is matched between conditions [15]. The study selection

process was done separately by MB and MPDLG, using the online platform Covidence (<https://www.covidence.org/home>). Conciliation of study selection was done after screening at the title/abstract level and after the full-text screening. Disagreements were settled through discussion and, if necessary, through consultation with BP and JS intervention.

2.3 Risks of bias

Risks of bias were appraised with a modified version [54] of the Effective Public Health Practice Project (EPHPP) Quality Assessment Tool for Quantitative Studies [55]. By not applying the selection bias component, this version was adapted to accommodate sport sciences studies where self-referral is common [54]. Further adaptations were made to reflect the need of the present review following the Cochrane Collaboration's guidelines [56]. The EPHPP does not provide explicit instructions on the assessment of cross-over trials, which was a type of design in the included studies. Appropriately randomized cross-over trials were considered equivalent to randomized controlled trials whereas cross-over trials performed in a counter-balanced order or with inappropriate randomization were considered equivalent to controlled clinical trials. Studies testing participants before and after opioid injection in a pre-post fashion were considered cohort studies. Risks of bias may be different across several outcomes and thus warranted confounders to be appraised specifically for the perception of effort. This is important because authors often consider the perception of effort when interpreting their results [e.g., 57] or use their results to draw conclusions on the regulation of the perception of effort [e.g., 58]. The risk of unblinding in the included studies is high, owing to the side effects the epidural anesthesia may have [e.g., pruritus, dysesthesia; 59]. As such, studies were considered blinded only if they detailed adequate blinding methods, such as the use of a sham injection. Because correctly guessing the interventions may influence the outcomes [60], the reviewers noted whether the participants were asked to guess the order of the interventions when applicable. However, this had no impact on the labelling of the "blinding" component of the EPHPP. The *withdrawals and drop-outs* component was not applied because none was reported in the included studies. This is probably because the included studies were brief, usually spanning over 3 to 5 visits. Finally, all the included studies were either cross-over trial or cohort studies, in which the risk of carry-over is critical. This component was considered when making a judgment for the intervention integrity. As such, the reviewer noted whether the risks of carry-over were minimized by allowing sufficient time in-between experimental visits for the effects of both protocol and the anesthesia to dissipate (wash-out period). Alternatively, risks of carry-over were considered minimal if baseline values between conditions were similar.

2.4 Coding process

Included articles were classified into three distinct groups: 1) effort dissociated from other exercise-related perceptions (*effort dissociated* group), 2) effort including other exercise-related perceptions (*effort not-dissociated* group) and 3) not enough information (*unclear* group). A first round attempted to code articles by the reported definition of the perception of effort or the instructions provided to their participants. However, this method was insufficient as none of the studies explicitly reported either of those. Following this, a finer decision-making process was developed by JC, CFB, MB, MPDLG and BP (see figure 2a). During a second round, the reviewers noted whether other exercise-related perceptions(s) were measured (coded as *effort dissociated*) and whether the perception of effort was used interchangeably with other exercise-related perception(s) (coded as *effort not dissociated*). If this information was not available, studies using the perception of effort as an index of the central motor command were coded as *effort dissociated*. Studies that could not be coded on that basis were coded as *unclear*. Articles were separately coded by MB and MPDLG. Decisions were then conciliated, and disagreements settled through discussion with BP. Notably, this process was done separately to the data extraction for meta-analysis (described below). The analyst (JS) was not provided with the final coding until after extraction was complete.

****Please insert figure 2****

2.5 Data extraction

Data extraction tables were prepared to map: (a) author and year of publication; (b) the test/exercise conditions used; (c) the method and scale used to capture ratings of perception of effort; (d) the means and standard errors or standard deviations for ratings of perception of effort including both control and intervention conditions; and (e) sample sizes. Of note, we included data for all perception of effort outcomes reported in studies and for all time points for which they were measured; thus, if a study reported multiple perception of effort measurements were captured, these were all included but appropriately coded as either being taken during submaximal tasks or maximal tasks (*i.e.*, at task failure). Some studies only reported data graphically or did not report outcomes in a manner conducive to extraction for our analysis, or despite investigating spinal blockade and capturing ratings of perception of effort did not report these outcomes as they were secondary in those studies. In these cases, authors were contacted for sharing either appropriate summary statistics or raw data to facilitate inclusion in our analysis. Three authors were

contacted of which one shared the required summary statistics whereas others were either unable or unwilling to share their data. For the studies reporting only graphical data, WebPlotDigitizer (v4.3, Ankit Rohatgi; <https://apps.automeris.io/wpd/>) was used to extract data for inclusion in our analysis.

2.6 Synthesis and analysis

The meta-analysis was performed using the ‘metafor’ [61] package in R (v 4.0.2; R Core Team, <https://www.r-project.org/>). All analysis codes are available on the Open Science Framework website (<https://osf.io/cy5n4/>). As all studies relied on within-participant design, the standardized mean change using the raw values (SMCR) as described by Becker [62] was calculated between the conditions where the pooled standard deviation from both conditions was used as the denominator for standardization [63]:

$$\text{Equation 1 : } SMCR = c(n - 1) \frac{\mu_{con} - \mu_{int}}{SD_{con+int}}$$

Where μ_{con} and μ_{int} are the means for control and intervention conditions respectively, c is a bias correction factor [64], and $SD_{(con+int)}$ is the pooled standard deviation calculated as:

$$\text{Equation 2 : } SD_{con+int} = \sqrt{\frac{(n_{con}-1)SD_{con}^2 + (n_{int}-1)SD_{int}^2}{n_{con}+n_{int}-2}}$$

As some studies reported zero variances (*e.g.*, some showing ceiling effects), a small constant (six sigma *i.e.*, 3×10^{-7}) was added to all studies prior to calculating effect sizes. The magnitude of standardized effect sizes was interpreted with reference to Cohen [65] thresholds: *trivial* (<0.2), *small* (0.2 to <0.5), *moderate* (0.5 to <0.8), and *large* (>0.8). Standardized effects were calculated in such a manner that negative effect size values indicated the presence of an intervention effect (*i.e.*, a drop in rating of perception of effort), whereas a positive effect size values indicated an effect favoring the control conditions.

As a departure from the pre-registered analysis due to the nested structure of the effect sizes calculated from the studies included (*i.e.*, effects nested within groups nested within studies), multilevel mixed effects meta-analyses reflecting these nested random effects in the model were performed. Cluster robust point estimates and precision of those estimates using 95% compatibility (confidence) intervals (CIs) were produced, weighted by inverse sampling variance to account for the within- and between-study variance (tau-squared). Restricted maximal likelihood estimation was used in all models. A main model was produced including all effects sizes. Several exploratory sub-group model analyses were also conducted and detailed in the pre-registration with slight departure here because of the coding process for methods

to capture ratings of perception of effort. In addition to a model for all studies, we also produced models for only studies where ratings of perception of effort were captured as *effort dissociated*, *effort not dissociated*, and *unclear* separately. Further, in all models we additionally sub-grouped for tasks as ‘maximal’ (*i.e.*, ratings of perception of effort captured at the point of task failure or during a similar maximal task *e.g.*, an MVC), and ‘submaximal’ (*i.e.*, ratings of perception of effort recorded during exercise prior to the point of task failure, or during tasks not necessarily requiring participants to reach task failure *e.g.*, a fixed duration task of absolute demands). This was to separate potential ceiling effects of maximal task conditions. Note, in the preregistration we initially stated we would calculate a pooled effect for all submaximal values reported in any given study. However, to increase the number of effect sizes for analysis, and thus statistical power, we opted to calculate each separately and thus use a multilevel mixed effects meta-analysis model to account for this. For each sub-group analysis multilevel models were produced except for the ‘*effort not dissociated*’ ‘maximal’ conditions where only one effect size was available.

In contrast to our pre-registration, and in light of the heterogeneity and poor reporting of methods to capture ratings of perception of effort, dichotomizing the existence of an effect for the main results was avoided. Therefore, traditional null hypothesis significance testing, which has been extensively critiqued [66, 67], was not employed. Instead, the implications of all results compatible with these data, from the lower limit to the upper limit of the interval estimates, was considered with the greatest interpretive emphasis placed on the point estimate.

Risk of small study bias was examined visually through contour-enhanced funnel plots. Q and I^2 statistics were also produced and reported [68]. A significant Q statistic is typically considered indicative of effects likely not being drawn from a common population. I^2 values indicate the degree of heterogeneity in the effects: 0-40% were not important, 30-60% moderate heterogeneity, 50-90% substantial heterogeneity, and 75-100% considerable heterogeneity [56]. For within participant effects pre-post correlations for measures were rarely reported; thus, a range of values for correlation coefficients ($r = 0.5$, 0.7 , and 0.9) and explored the sensitivity of results to each of these was assumed. As overall findings were relatively insensitive to this range, the results for $r = 0.7$ are reported here. Results for inclusion of the other assumed correlation coefficients are reported in the supplementary material available on OSF ($r = 0.5$, <https://osf.io/gqby6/>; $r = 0.9$, <https://osf.io/qbe2n/>). [graphical display of study heterogeneity; 69]

315

316 **3. Results**

317 The search across all databases returned 902 articles. After the removal of 319 duplicates, 583 original
 318 articles remained and were screened at the title/abstract level, leaving 80 articles. A total of 18 articles
 319 were retained. The screening process is shown in figure 3. None of the included articles defined the
 320 perception of effort nor provided instructions for rating it. Therefore, included studies could not be coded
 321 on that basis and instead, other cues have been used (see figure 2a). Based upon this, among the 18 articles,
 322 only 6 could be classified in the *effort dissociated* group, while 4 were placed in the *effort not dissociated*
 323 group and 8 in the *unclear* group. The pooled number of participants in the studies included for qualitative
 324 analysis was 163 across 21 groups within the 18 studies, and with sample sizes ranging from 5 to 16
 325 participants (median = 8). However, data was not available for quantitative synthesis in 2 studies, both
 326 belonging to the *effort not-dissociated* group thus reducing the sample size on which quantitative synthesis
 327 was based to $n = 164$. Full details of all studies included in quantitative synthesis can be seen in the data
 328 extraction table (<https://osf.io/ku3w7/>).

329 ****Please insert figure 3 here****

330

331 **Risks of bias**

332 None of the studies were labelled “strong”, whereas 5 were “moderate” and 13 “weak”. Most studies
 333 used a randomized or counter-balanced within-subject design, but 7 studies were classified as cohort
 334 studies (pre-test, post-test). Almost half ($n = 7$) of the studies were deemed to have strongly controlled for
 335 known confounders of the perception of effort, but 6 of them were labelled “moderate”, with the remaining
 336 5 being considered “weak”. None of the studies described how the assessor was blinded except for one
 337 [70]. In contrast, 5 studies blinded their participants with a placebo injection. The remaining 13 studies
 338 compared the epidural anaesthesia with a “no-intervention” control condition and the participants could
 339 therefore not be considered blind. Furthermore, none of the included studies reported asking the
 340 participants to guess the order of the intervention after completion of the protocols. All studies used either
 341 the Borg’s scale (RPE 6-20) or the CR10 scale, which are known psychophysical scales in the context of
 342 physical exercise. However, several ($n = 9$) articles reported to have used a modified version of the CR10
 343 scale without providing any information on the modifications performed. The validity and reliability of
 344 these modifications were therefore unclear. The table 1 shows the risk of bias within studies for every
 345 included articles. An audit trail is available in the supplementary material 2.

****Please insert table 1 here****

Meta-Analysis

The quantitative analysis could be performed on only 16 studies, the remaining two providing insufficient information. The main model including all combined effects sizes ($k = 49$ across 16 clusters [median = 2, range = 1 to 8 effects per cluster]) revealed a negative trivial point estimate with precision ranging from negative small to positive trivial effects for the interval estimate (-0.05 [95% CIs = -0.28 to 0.18]), yet with moderate to substantial heterogeneity ($Q_{(48)} = 127.22$, $p < 0.0001$, $I^2_{\text{between_study}} = 77\%$, $I^2_{\text{between_group}} = 0\%$, $I^2_{\text{within_group}} = 0\%$). When considering only submaximal conditions the model ($k = 39$ across 13 clusters [median = 2, range = 1 to 8 effects per cluster]) revealed a point estimate close to zero with precision ranging from negative small to positive small effects for the interval estimate (0.05 [95% CIs = -0.32 to 0.42]), with substantial heterogeneity ($Q_{(38)} = 103.00$, $p < 0.0001$, $I^2_{\text{between_study}} = 77\%$, $I^2_{\text{between_group}} = 0\%$, $I^2_{\text{within_group}} = 0\%$). Considering only maximal conditions the model ($k = 10$ across 9 clusters [median = 1, range = 1 to 2 effects per cluster]) revealed a negative small point estimate with precision ranging from negative moderate to positive trivial effects for the interval estimate (-0.17 [95% CIs = -0.47 to 0.13]), with moderate to substantial heterogeneity ($Q_{(9)} = 22.92$, $p = 0.0064$, $I^2_{\text{between_study}} = 0\%$, $I^2_{\text{between_group}} = 30\%$, $I^2_{\text{within_group}} = 30\%$). Figure 4 presents the funnel plot for all studies with color coding by method for capturing rating of perception of effort. Figures 5 and 6 present all effect sizes for the submaximal models and the overall model estimates for all coding and combined.

****Please insert figure 4 here****

Ratings of perception of effort as ‘effort dissociated’. The subgroup model where the method of rating of perception of effort was coded as *effort* ($k = 19$ across 6 clusters [median = 2.5, range = 1 to 8 effects per cluster]) revealed an overall small point estimate with precision ranging from positive trivial to positive moderate effects for the interval estimate (0.39 [95% CIs = 0.13 to 0.64]), with relative homogeneity ($Q_{(18)} = 36.96$, $p = 0.0053$, $I^2_{\text{between_study}} = 77\%$, $I^2_{\text{between_group}} = 0\%$, $I^2_{\text{within_group}} = 49\%$). When considering only submaximal conditions the model ($k = 16$ across 6 clusters [median = 2, range = 1 to 7 effects per cluster]) revealed a small point estimate with precision ranging from positive trivial to positive large effects for the interval estimate (0.54 [95% CIs = 0.07 to 1.0]), with moderate to substantial heterogeneity ($Q_{(15)} = 31.44$, $p = 0.0077$, $I^2_{\text{between_study}} = 51\%$, $I^2_{\text{between_group}} = 0\%$, $I^2_{\text{within_group}} = 8\%$). When considering only maximal conditions the model ($k = 3$ across 3 clusters [1 effect per cluster]) revealed a negative trivial

point estimate with precision ranging from negative small to positive small effects for the interval estimate (-0.05 [95% CIs = -0.24 to 0.15]), with relative homogeneity ($Q_{(2)} = 0.12$, $p = 0.9412$, $I^2_{\text{between_study}} = 0\%$, $I^2_{\text{between_group}} = 0\%$, $I^2_{\text{within_group}} = 0\%$).

****Please insert figure 5 here****

Ratings of perception of effort as 'effort not dissociated'. The subgroup model where the method of rating of perception of effort capture was coded as *effort not dissociated* ($k = 8$ across 2 clusters [4 effects per cluster]) revealed an overall negative small point estimate with poor precision ranging from negative large to positive large effects for the interval estimate (-0.29 [95% CIs = -2.39 to 1.8]), with relative homogeneity ($Q_{(7)} = 9.62$, $p = 0.2113$, $I^2_{\text{between_study}} = 7\%$, $I^2_{\text{between_group}} = 7\%$, $I^2_{\text{within_group}} = 14\%$). When considering only submaximal conditions the model ($k = 7$ across 2 clusters [median = 3.5, range = 3 to 4 effects per cluster]) revealed a negative small point estimate poor precision ranging from negative large to positive large effects for the interval estimate (-0.32 [95% CIs = -3.07 to 2.42]), with moderate heterogeneity ($Q_{(6)} = 9.51$, $p = 0.1467$, $I^2_{\text{between_study}} = 12\%$, $I^2_{\text{between_group}} = 12\%$, $I^2_{\text{within_group}} = 20\%$). When considering only maximal conditions only a single effect met these conditions which revealed a negative small point estimate with precision ranging from negative moderate to positive small effects for the interval estimate (-0.22 [95% CIs = -0.76 to 0.33]).

Ratings of perception of effort as 'unclear'. The subgroup model where the method of rating of perception of effort capture was coded as *unclear* ($k = 22$ across 8 clusters [median = 2, range = 1 to 8 effects per cluster]) revealed an overall negative small point estimate with precision ranging from negative moderate to negative trivial effects for the interval estimate (-0.27 [95% CIs = -0.50 to -0.04]), with moderate heterogeneity ($Q_{(21)} = 42.14$, $p = 0.004$, $I^2_{\text{between_study}} = 6\%$, $I^2_{\text{between_group}} = 0\%$, $I^2_{\text{within_group}} = 45\%$). When considering only submaximal conditions the model ($k = 16$ across 5 clusters [median = 2, range = 1 to 8 effects per cluster]) revealed a negative small point estimate with precision ranging from negative large to positive trivial effects for the interval estimate (-0.37 [95% CIs = -0.88 to 0.14]), with moderate to substantial heterogeneity ($Q_{(15)} = 19.27$, $p = 0.2020$, $I^2_{\text{between_study}} = 59\%$, $I^2_{\text{between_group}} = 0\%$, $I^2_{\text{within_group}} = 0\%$). When considering only maximal conditions the model ($k = 6$ across 5 clusters [median = 1, range = 1 to 2 effects per cluster]) revealed a negative small point estimate with poor precision ranging from negative large to positive moderate effects for the interval estimate (-0.25 [95% CIs = -0.91 to 0.41]), with

407 considerable heterogeneity ($Q_{(5)} = 22.47$, $p = 0.004$, $I^2_{\text{between_study}} = 0\%$, $I^2_{\text{between_group}} = 39\%$, $I^2_{\text{within_group}} =$
 408 39%).

409 ****Please insert figure 6 here****

410

411

4. Discussion

This systematic review with meta-analysis was conducted to investigate the effect of pharmacologically blocking muscle afferents on the perception of effort during physical exercise. We also sought to explore the differences in reported perception of effort according to the approach used to investigate this construct. Eighteen articles were coded as *effort dissociated* ($n = 6$), *effort not dissociated* ($n = 4$) or *unclear* ($n = 8$) according to whether the authors included other exercise-related perceptions in the investigation of the perception of effort. Considering all subgroups combined, a trivial negative point estimate for ratings of perceived effort reduction following pharmacological blockade of muscle afferents was observed, with interval estimate ranging from negative small to positive trivial effects. The magnitude of the interval estimate crossing zero as well as the trivial negative point of estimate (0.05) suggests that regardless of the theoretical approach used to investigate the perception of effort, pharmacologically blocking muscle afferents feedback does not reduce the perception of effort during physical exercise.

The subgroup analysis also revealed a clear influence of the theoretical approach used to investigate the perception of effort (i.e., as a construct dissociated or not from other exercise-related perceptions). To the best of our knowledge, this systematic review is the first to highlight an influence of the theoretical approach used to investigate this perception, and strongly suggests that future experimental studies should carefully report the instructions provided to the participants for rating perception of effort.

Effort dissociated subgroup. Only considering the *effort dissociated* subgroup, pooling 6 studies yielded a small positive point estimate with a positive confidence interval suggesting that impairing group III-IV muscle afferents increased ratings of perceived effort. Participants working at the same absolute demands (i.e., same external workload in both conditions) reported higher perception of effort with epidural anesthesia [32, 33, 37]. However, when working at similar relative intensities (i.e., taking into consideration any muscle strength reduction following injection of local anesthetic [36]), the same participants reported similar perception of effort. Likewise, ratings of perceived effort were also similar at a given oxygen uptake during graded exercises [32, 37]. The *effort dissociated* subgroup also showed relatively low heterogeneity, though for submaximal tasks the majority came from between study factors. Differences across studies could be explained by differences in experimental designs [69], as results differed with the calibration of exercise demands according to absolute or relative workloads. It is also important to note that the increased perception of effort in these studies is likely due to the use of lidocaine and/or bupivacaine to block III-IV muscle afferent feedback. Indeed, contrary to fentanyl which acts more specifically on sensory transmission [71], lidocaine and bupivacaine is known to affect sodium and potassium channels [72] and reduce force production capacity [73] thereby requiring the participants to

increase their motor command to maintain the same absolute workload. According to the corollary discharge model, this increased motor command increases the magnitude of the associated corollary discharge, which in turn increases the perception of effort [21]. This subgroup analysis reveals that when effort is investigated as dissociated from other exercise-related perceptions, pharmacological blockade of muscle afferents does not reduce perception of effort. Therefore, as perception of effort is not reduced, muscle afferent feedback cannot be considered as a sensory signal processed by the brain to generate the perception of effort. This result reinforces the potential of using the perception of effort intensity as a psychophysiological index of the motor command [21, 25, 74-76], as traditionally performed in the neuroscience, cardiovascular physiology and kinesthesia literatures [21-25].

Effort not-dissociated subgroup. Only considering the *effort not dissociated* subgroup, pooling 2 studies yielded an overall small negative point estimate. This negative point estimate was associated with an important imprecision based upon the confidence interval range likely due to the low number of studies and small cluster sample correction for robust estimates. Both studies observed lower ratings of perceived effort in epidurally anaesthetized participants [31, 70]. Interestingly, Amann et al. [31] observed lower ratings of perceived effort only at higher cycling power output (*i.e.*, 80% peak power output, 325 ± 19 W). Participants reported similar ratings of perceived effort at 50, 100 and 150 W. Likewise, chronic obstructive pulmonary disease patients performing a constant-load cycling protocol at 80% peak power output tended to report lower perceived effort intensity at iso-time points [70]. These authors found similar ratings of perceived effort at exhaustion. According to the Oxford Dictionary, discomfort can be defined as a slight pain and something that makes a person feel physically uncomfortable. Because of its relation to the concept of pain, the inclusion of discomfort in the definition of the perception of effort may bias the ratings of perceived effort whenever there is a change in the perception of pain [13, 77]. Although there seems to exist wide interindividual variability in pain threshold during cycle ergometry, muscle pain is known to increase with increased exercise demands during this task [78]. Attending to these perceptions when measuring the perception of effort may attenuate the perceptual differences between conditions during lower cycling demands where discomfort is already low and thus, any difference with epidural anesthesia would likely be minimal. Conversely, when working at 80% peak power output, participants reported substantially lower ratings of perceived effort with epidural anesthesia, probably because they felt less discomfort and/or pain than they normally would have. Pain and associated unpleasant sensations are transmitted through group III-IV fibres [79], and thus are attenuated with epidural anesthesia. Amann et al., [31] and Gagnon et al. [70] both observed lower ratings of perceived effort, suggesting that the reduction in muscle pain and associated discomfort may have biased the ratings of perceived effort when other exercise-related perceptions were included in the definition of effort. A study employing a self-paced

protocol that could not be added to this meta-analysis found similar results [36]. Participants performed a 5-km cycling time trial with and without lidocaine with a mean power output similar to that of Amann et al. [31]. The authors however found a decrease in ratings of perceived effort of nearly 2 unit-points on the CR10 scale (CTRL: 8.4 ± 0.4 SEM; Lidocaine: 6.8 ± 0.4 SEM). Given the high-power output, the fact that feedback from group III-IV is known to be the signal processed by the brain to generate muscle pain [79], and that the authors reported investigating “limb discomfort”, the lower values likely reflect a decreased pain and discomfort when cycling with lidocaine. As such, the inclusion of discomfort in the definition of effort would likely result in a decrease in reported ratings of perceived effort. Interestingly, a similar protocol, with the only difference of using fentanyl instead of lidocaine, observed an increase in ratings of perceived effort at the completion of the 5 km time-trial [58]. However, the authors also observed an excessive development of fatigue resulting in an increase in central motor drive, known to exacerbate the perception of effort [3, 21, 25, 28]. Moreover, when an effect of the epidural anesthesia is detected (e.g., 13% reduction in “limb discomfort”[31]), the magnitude of its effect is small. Even when the theoretical approach encompasses several exercise-related perceptions, the contribution of group III-IV muscle afferents appears to be limited as previously suggested [77].

Unclear subgroup. Only considering the *unclear* subgroup, pooling 8 articles yielded an overall small negative point estimate with a negative confidence interval. Among those articles, 2 observed lower ratings of perceived effort at task failure when exercising with impaired muscle afferents with similar integrated forces between conditions [80, 81]. Interestingly, 2 other studies observed similar ratings of perceived effort at task failure. Amann et al. [58] found a 27% decrease in perception of effort intensity with epidural anesthesia at the 3-min mark (average time, placebo: 8.7 ± 0.3 , fentanyl: 6.8 ± 0.3), but not at exhaustion in trained athletes cycling at 80% of their peak power output previously measured. Similarly, Sidhu et al. [82] observed a decrease in perception of effort intensity with epidural anesthesia only at 25% of endurance time during a similar exercise protocol. It must be noted that a 1-unit difference on the CR10 was consistently maintained throughout the protocol (25% endurance time [ET], 5.9 ± 0.4 vs 4.9 ± 0.4 ; 50% ET, 8.3 ± 0.3 vs 7.4 ± 0.5 ; 70% ET, 9.4 ± 0.2 , 8.5 ± 0.3), until exhaustion where values were nearly identical (100% ET, $9.9 \pm .01$ vs 10.0 ± 0). It is indicative of a tendency for ratings of perceived effort to be lower in the epidural anesthesia condition, albeit not reaching statistical significance. Similar values at the end of the endurance time would also be consistent with previous studies suggesting that the perception of effort attain near maximal values at exhaustion [e.g., 83]. Three other studies using fentanyl in hypertensive and heart failure patients also did not find different ratings of perceived effort compared to a sham or control condition [38, 84, 85]. Exercise demands, determined with cycling power output, was extremely low in one of the studies (40 W; [84]). Because participants reported similar ratings of perceived

effort when also considering discomfort at lower intensities [31] and that fentanyl does not lead to loss in muscle strength [73], it is not possible to determine which was the cause of the lack of differences. However, when exercising at higher intensities (*e.g.*, 65% - 80% W_{peak}), healthy controls and heart failure participants reported similar ratings of perceived effort during both conditions [38]. Furthermore, healthy and heart failure participants did not differ in ratings of perceived effort despite large differences in external workload [38], further suggesting that it is the relative and not the absolute workload determining the perception of effort.

Although the involvement of the central motor drive in the perception of effort seems widely accepted [21, 24, 36], there is still confusion about the role of group III-IV muscle afferents as a signal processed by the brain to generate the perception of effort [13, 27]. In fact, physiologists have still not reached a consensus after more than 150 years of debate [86-89]. Central projections of group III-IV muscle afferents to several spinal and supra-spinal sites, including sensory cortices, anatomically support the afferent feedback model [90, 91]. This model also finds experimental evidence from studies involving epidural anesthesia [31, 70]. However, as mentioned above, the inclusion of other exercise-related perceptions likely biased the ratings of perceived effort measured. Furthermore, if these muscle afferents constitute a centrally processed signal generating the perception of effort, stimulation of group III-IV muscle afferents would generate a sense of effort even in the absence of central motor drive. However, injections of physiological concentrations of metabolites known to stimulate those muscle afferents do not generate perception of effort at rest [92]. This manipulation however elicits sensations related to discomfort (*e.g.*, itch, tingling) and pain. It appears that the presence of the motor command, and therefore the voluntary engagement of the participant in the task, is crucial for experiencing the perception of effort.

There is extensive support that the neurocognitive processing of the corollary discharge generates the perception of effort. First, significant correlations between the ratings of perceived effort and the amplitude of movement-related cortical potential (MRCP), an index of the motor command, have previously been observed [21, 25, 74, 93, 94]. For example, a reduction in force production capacity is associated with an increased MRCP amplitude and an increased perception of effort intensity to maintain the same absolute force [21]. When caffeine is ingested, perception of effort intensity to maintain the same absolute force is reduced in association with a decreased MRCP amplitude [25]. This positive effect of caffeine on the perception of effort is most likely due to the increased excitability of the corticospinal pathway induced by its ingestion [95-97], leading to a lower motor command required to activate the working muscles as revealed by the decreased MRCP amplitude. Second, support in favor of the role of the corollary discharge as an internal signal generating the perception of effort can be found in various neuroscience or

psychophysiological studies. For example, Zenon et al [98] demonstrated that disrupting the supplementary motor area via continuous theta burst transcranial magnetic stimulation increases perception of effort. Other studies demonstrated a close relationship between perception of effort and physiological variables known to be strongly influenced by the motor command, such as the respiratory frequency [99] or the electromyographic signal [76].

Strengths and limitations

One strength of this systematic review is that our search was not restrained by publication date, despite the articles spanning three decades. Rather, the shift from muscle weakness-inducing lidocaine and bupivacaine to the highly selective μ -opioid receptor agonist fentanyl demonstrated the importance of the magnitude of the central motor drive in generating the perception of effort. In the presence of muscle weakness (induced by some opioids binding to spinal motoneurons), exercisers must increase the magnitude of their motor command to maintain the same absolute level of performance [30]. This was observed in older studies via the higher ratings of perceived effort reported when participants were working at similar external workload [e.g., 37, 100]. Another strength of this meta-analysis is that we opted to pool the data per time-point, instead of averaging the ratings of perceived effort within studies. This allows us to avoid any potential ceiling effects from maximal conditions, where ratings of perceived effort are expected to attain near maximal values. Moreover, having each effect reported in each study increased the number of effect sizes and thus the statistical power and the precision of this meta-analysis.

One limitation of this systematic review is that none of the included studies explicitly provided the definition of the perception of effort, or the instructions given to the participants to report their ratings of perceived effort. Therefore, to overcome this limitation, a unique coding was created to be able to classify the included articles based on cues leading to the assumption or not of the inclusion of other exercise-related perceptions in the definition of effort. While some may argue that such approach opens the doors for interpretation, we would like to emphasize that the coding process was clear and objective, and could be reproduced by other research groups, as presented in figure 2a. We are therefore confident that our coding process was successful at separating studies according to the definition of the perception of effort used and the associated approach used by the researcher (dissociated perception or not). Importantly, this limitation strongly emphasizes the need for better reporting of the definition of the perception of effort and associated instructions provided to the participants in the methods or in supplementary materials. Moreover, we had far fewer studies in the *effort not dissociated* group than the other 2 groups, decreasing the precision of the meta-analysis for this group.

Conclusion

Our results indicate that the group III-IV muscle afferents does not contribute as a signal processed by the brain to generate the perception of effort. Therefore, during voluntary movements, the perception of effort appears to be generated by the brain processing of the corollary discharge associated with the motor command. This meta-analysis also underscores the importance to provide clear and standardized instructions to the participants to avoid the confounding effect of other exercise-related perceptions in the ratings of perceived effort. We therefore recommend, similarly to others [13, 27, 101], that researchers and clinicians instruct their participants to rate their perception of effort specifically by excluding other exercise-related perception. Such approach is crucial for researchers investigating the perception of effort as a psychophysiological marker of the magnitude of the motor command [e.g., 74, 76] and using muscle pain as a psychophysiological marker of feedback from group III-IV muscle afferents [e.g., 102, 103]. Investigating effort as a unique and dissociated perception is also crucial to better understand how perception of effort interacts with other exercise-related perceptions, such as pain, and influence performance and the regulation of human behavior. It could unravel underlying mechanisms generating and regulating the perception of effort, and lead to the development of unique multidisciplinary interventions aimed at decreasing perception of effort to improve the adherence to an exercise training program [e.g., 104].

591 **Acknowledgements**

592 The authors warmly thank the librarian Marc-Olivier Croteau for his precious assistance.

593

594

595 **5. Bibliography**

- 596 1. Gaveau J, Grospretre S, Berret B, Angelaki DE, Papaxanthis C. A cross-species neural
597 integration of gravity for motor optimization. *Science Advances*. 2021;7(15):eabf7800.
- 598 2. Izawa J, Rane T, Donchin O, Shadmehr R. Motor Adaptation as a Process of Reoptimization.
599 *The Journal of Neuroscience*. 2008;28(11):2883-91.
- 600 3. Pageaux B, Lepers R. Fatigue induced by physical and mental exertion increases perception of
601 effort and impairs subsequent endurance performance. *Frontiers in physiology*. 2016 2016;7:587.
- 602 4. Cook DB, O'Connor PJ, Lange G, Steffener J. Functional neuroimaging correlates of mental
603 fatigue induced by cognition among chronic fatigue syndrome patients and controls. *NeuroImage*. 2007
604 2007/05/15/;36(1):108-22.
- 605 5. Barhorst EE, Andrae WE, Rayne TJ, Falvo MJ, Cook DB, Lindheimer JB. Elevated Perceived
606 Exertion in People with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome and Fibromyalgia: A
607 Meta-analysis. *Medicine & Science in Sports & Exercise*. 2020;52(12):2615-27.
- 608 6. Kuppaswamy A, Clark EV, Turner IF, Rothwell JC, Ward NS. Post-stroke fatigue: a deficit in
609 corticomotor excitability? *Brain*. 2014;138(1):136-48.
- 610 7. Fernandez C, Firdous S, Jehangir W, Behm B, Mehta Z, Berger A, et al. Cancer-Related Fatigue:
611 Perception of Effort or Task Failure? *American Journal of Hospice and Palliative Medicine®*.
612 2020;37(1):34-40.
- 613 8. Piña IL, Apstein CS, Balady GJ, Belardinelli R, Chaitman BR, Duscha BD, et al. Exercise and
614 heart failure: a statement from the American Heart Association Committee on exercise, rehabilitation,
615 and prevention. *Circulation*. 2003 2003;107(8):1210-25.
- 616 9. AACVPR. Guidelines for cardiac rehabilitation and secondary prevention programs: Human
617 Kinetics; 2013.
- 618 10. Eston R, Parfitt G. Perceived exertion, heart rate and other non-invasive methods for exercise
619 testing and intensity control. *Kinanthropometry and Exercise Physiology: Volume One: Anthropometry*.
620 2018 2018:464.
- 621 11. Impellizzeri FM, Rampinini E, Coutts AJ, Sassi A, Marcora SM. Use of RPE-based training load
622 in soccer. *Medicine & Science in sports & exercise*. 2004 2004;36(6):1042-7.
- 623 12. ACSM. ACSM's exercise testing and prescription: Lippincott williams & wilkins; 2017.
- 624 13. Pageaux B. Perception of effort in exercise science: definition, measurement and perspectives.
625 *European Journal of Sport Science*. 2016 2016;16(8):885-94.
- 626 14. Smirmaul BdPC. Sense of effort and other unpleasant sensations during exercise: clarifying
627 concepts and mechanisms. *Br J Sports Med*. 2012 2012;46(5):308-11.
- 628 15. Steele J. What is (perception of) effort? Objective and subjective effort during task performance.
629 *PsyArXiv*. 2020 2020.
- 630 16. St Gibson AC, Lambert EV, Rauch LHG, Tucker R, Baden DA, Foster C, et al. The role of
631 information processing between the brain and peripheral physiological systems in pacing and perception
632 of effort. *Sports medicine*. 2006 2006;36(8):705-22.
- 633 17. Noble BJ, Robertson RJ. The Borg scale: development, administration and experimental use.
634 *Perceived exertion: Champaign: Human Kinetics*; 1996. p. 59-89.
- 635 18. Tucker R. The anticipatory regulation of performance: the physiological basis for pacing
636 strategies and the development of a perception-based model for exercise performance. *British journal of*
637 *sports medicine*. 2009 2009;43(6):392-400.

19. Amann M, Venturelli M, Ives SJ, McDaniel J, Layec G, Rossman MJ, et al. Peripheral fatigue limits endurance exercise via a sensory feedback-mediated reduction in spinal motoneuronal output. *Journal of applied physiology*. 2013 2013;115(3):355-64.
20. Keller JL, Housh TJ, Hill EC, Smith CM, Schmidt RJ, Johnson GO. Neuromuscular responses of recreationally active women during a sustained, submaximal isometric leg extension muscle action at a constant perception of effort. *European journal of applied physiology*. 2018 2018;118(12):2499-508.
21. de Morree HM, Klein C, Marcora SM. Perception of effort reflects central motor command during movement execution. *Psychophysiology*. 2012 2012;49(9):1242-53.
22. Enoka RM, Stuart DG. Neurobiology of muscle fatigue. *Journal of applied physiology*. 1992 1992;72(5):1631-48.
23. Taylor JL. Proprioception. 2009 2009.
24. Winchester PK, Williamson JW, Mitchell JH. Cardiovascular responses to static exercise in patients with Brown-Sequard syndrome. *The Journal of physiology*. 2000 2000;527(Pt 1):193.
25. de Morree HM, Klein C, Marcora SM. Cortical substrates of the effects of caffeine and time-on-task on perception of effort. *Journal of Applied Physiology*. 2014 2014;117(12):1514-23.
26. Pageaux B, Gaveau J. Studies using pharmacological blockade of muscle afferents provide new insights into the neurophysiology of perceived exertion. *The Journal of physiology*. 2016 2016;594(18):5049.
27. Marcora SM. Perception of effort during exercise is independent of afferent feedback from skeletal muscles, heart, and lungs. *Journal of applied physiology*. 2009 2009;106(6):2060-2.
28. Marcora SM, Bosio A, De Morree HM. Locomotor muscle fatigue increases cardiorespiratory responses and reduces performance during intense cycling exercise independently from metabolic stress. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 2008 2008;294(3):R874-R83.
29. Weavil JC, Amann M. Corticospinal excitability during fatiguing whole body exercise. *Progress in brain research*. 2018 2018;240:219-46.
30. Gandevia SC. Spinal and supraspinal factors in human muscle fatigue. *Physiological reviews*. 2001 2001.
31. Amann M, Blain GM, Proctor LT, Sebranek JJ, Pegelow DF, Dempsey JA. Group III and IV muscle afferents contribute to ventilatory and cardiovascular response to rhythmic exercise in humans. *Journal of Applied Physiology*. 2010 2010;109(4):966-76.
32. Fernandes A, Galbo H, Kjaer M, Mitchell JH, Secher NH, Thomas SN. Cardiovascular and ventilatory responses to dynamic exercise during epidural anaesthesia in man. *The Journal of physiology*. 1990 1990;420(1):281-93.
33. Smith SA, Querry RG, Fadel PJ, Gallagher KM, Strømstad M, Ide K, et al. Partial blockade of skeletal muscle somatosensory afferents attenuates baroreflex resetting during exercise in humans. *The Journal of physiology*. 2003 2003;551(3):1013-21.
34. Amann M, Venturelli M, Ives SJ, Morgan DE, Gmelch B, Witman MA, et al. Group III/IV muscle afferents impair limb blood in patients with chronic heart failure. *International journal of cardiology*. 2014;174(2):368-75.
35. Amann M, Wan H-Y, Thurston TS, Georgescu VP, Weavil JC. On the influence of group III/IV muscle afferent feedback on endurance exercise performance. *Exercise and Sport Sciences Reviews*. 2020 2020;48(4):209-16.
36. Amann M, Proctor LT, Sebranek JJ, Eldridge MW, Pegelow DF, Dempsey JA. Somatosensory feedback from the limbs exerts inhibitory influences on central neural drive during whole body endurance exercise. *Journal of applied physiology*. 2008 2008;105(6):1714-24.
37. Kjaer M, Secher NH, Bach FW, Sheikh S, Galbo H. Hormonal and metabolic responses to exercise in humans: effect of sensory nervous blockade. *American Journal of Physiology-Endocrinology And Metabolism*. 1989 1989;257(1):E95-E101.

38. Olson TP, Joyner MJ, Eisenach JH, Curry TB, Johnson BD. Influence of locomotor muscle afferent inhibition on the ventilatory response to exercise in heart failure. *Experimental physiology*. 2014 2014;99(2):414-26.
39. Borg GAV. Physical performance and perceived exertion. 1962 1962.
40. Borg GAV. Borg's perceived exertion and pain scales: Human kinetics; 1998.
41. Gamberale F. Perception of effort in manual materials handling. *Scandinavian journal of work, environment & health*. 1990:59-66.
42. Hamilton AL, Killian KJ, Summers E, Jones NL. Quantification of intensity of sensations during muscular work by normal subjects. *Journal of Applied Physiology*. 1996 1996;81(3):1156-61.
43. O'Connor PJ, Cook DB. Moderate-intensity muscle pain can be produced and sustained during cycle ergometry. *Medicine & Science in Sports & Exercise*. 2001 2001;33(6):1046-51.
44. Christian RJ, Bishop D, Girard O, Billaut F. The role of sense of effort on self-selected cycling power output. *Frontiers in physiology*. 2014 2014;5:115.
45. Steele J, Fisher J, McKinnon S, McKinnon P. Differentiation between perceived effort and discomfort during resistance training in older adults: reliability of trainee ratings of effort and discomfort, and reliability and validity of trainer ratings of trainee effort. *Journal of Trainology*. 2016 2016;6(1):1-8.
46. Stuart C, Steele J, Gentil P, Giessing J, Fisher JP. Fatigue and perceptual responses of heavier- and lighter-load isolated lumbar extension resistance exercise in males and females. *PeerJ*. 2018 2018;6:e4523.
47. Jones LA. The senses of effort and force during fatiguing contractions. 1995 1995: Springer; 1995. p. 305-13.
48. McCloskey DI, Ebeling P, Goodwin GM. Estimation of weights and tensions and apparent involvement of a "sense of effort". *Experimental neurology*. 1974 1974;42(1):220-32.
49. Marcora S. Psychobiology of fatigue during endurance exercise. *Endurance performance in sport*: Routledge; 2019. p. 15-34.
50. Richter M, Gendolla GH, Wright RA. Three decades of research on motivational intensity theory: What we have learned about effort and what we still don't know. *Advances in motivation science*: Elsevier; 2016. p. 149-86.
51. Preston J, Wegner DM. Elbow grease: When action feels like work. *Social cognition and social neuroscience Oxford handbook of human action*. 2009 2009:569-86.
52. Marcora SM. Perception: Effort of. *Encyclopedia of perception*. Goldstein, E. B. ed. Thousand Oak, CA: Sage; 2010. p. 380-3.
53. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *Journal of clinical epidemiology*. 2009 2009;62(10):e1-e34.
54. McCormick A, Meijen C, Marcora SM. Psychological determinants of whole-body endurance performance. *Sports medicine*. 2015 2015;45(7):997-1015.
55. Thomas BH, Ciliska D, Dobbins M, Micucci S. A process for systematically reviewing the literature: providing the research evidence for public health nursing interventions. *Worldviews on Evidence-Based Nursing*. 2004 2004;1(3):176-84.
56. Higgins, Green S. *Cochrane handbook for systematic reviews of interventions*. Version 5.1. 0 [updated March 2011]. The Cochrane Collaboration. www.cochrane-handbook.org. 2011 2011.
57. Kjær M, Hanel B, Worm L, Perko G, Lewis SF, Sahlin K, et al. Cardiovascular and neuroendocrine responses to exercise in hypoxia during impaired neural feedback from muscle. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 1999 1999;277(1):R76-R85.

58. Amann M, Blain GM, Proctor LT, Sebranek JJ, Pegelow DF, Dempsey JA. Implications of group III and IV muscle afferents for high-intensity endurance exercise performance in humans. *The Journal of physiology*. 2011 2011;589(21):5299-309.
59. Chaney MA. Side effects of intrathecal and epidural opioids. *Canadian Journal of Anaesthesia*. 1995 1995;42(10):891-903.
60. Schnoll RA, Epstein L, Audrain J, Niaura R, Hawk L, Shields PG, et al. Can the blind see? Participant guess about treatment arm assignment may influence outcome in a clinical trial of bupropion for smoking cessation. *Journal of substance abuse treatment*. 2008 2008;34(2):234-41.
61. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *Journal of statistical software*. 2010 2010;36(3):1-48.
62. Becker BJ. Synthesizing standardized mean-change measures. *British Journal of Mathematical and Statistical Psychology*. 1988 1988;41(2):257-78.
63. Morris SB. Estimating effect sizes from pretest-posttest-control group designs. *Organizational research methods*. 2008 2008;11(2):364-86.
64. Morris SB. Distribution of the standardized mean change effect size for meta-analysis on repeated measures. *British Journal of Mathematical and Statistical Psychology*. 2000 2000;53(1):17-29.
65. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. Hillsdale: Erlbaum. Conner, BE (1988). *The Box in the Barn*. Columbus: Highlights for ...; 1988.
66. Amrhein V, Greenland S, McShane B. Scientists rise up against statistical significance. *Nature*. 2019 2019;305-7.
67. McShane BB, Gal D, Gelman A, Robert C, Tackett JL. Abandon statistical significance. *The American Statistician*. 2019 2019;73(sup1):235-45.
68. Higgins, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *Bmj*. 2003 2003;327(7414):557-60.
69. Olkin I, Dahabreh IJ, Trikalinos TA. GOSH—a graphical display of study heterogeneity. *Research Synthesis Methods*. 2012 2012;3(3):214-23.
70. Gagnon P, Bussires JS, Ribeiro F, Gagnon SL, Saey D, Gagn N, et al. Influences of spinal anesthesia on exercise tolerance in patients with chronic obstructive pulmonary disease. *American journal of respiratory and critical care medicine*. 2012 2012;186(7):606-15.
71. Kanai A, Osawa S, Suzuki A, Ozawa A, Okamoto H, Hoka S. Regression of sensory and motor blockade, and analgesia during continuous epidural infusion of ropivacaine and fentanyl in comparison with other local anesthetics. *Pain Medicine*. 2007;8(7):546-53.
72. Olschewski A, Hempelmann G, Vogel W, Safronov BV. Blockade of Na⁺ and K⁺ currents by local anesthetics in the dorsal horn neurons of the spinal cord. *The Journal of the American Society of Anesthesiologists*. 1998;88(1):172-9.
73. Amann M, Proctor LT, Sebranek JJ, Pegelow DF, Dempsey JA. Opioid-mediated muscle afferents inhibit central motor drive and limit peripheral muscle fatigue development in humans. *The Journal of Physiology*. 2009 2009;587(1):271-83.
74. Jacquet T, Lepers R, Poulin-Charronnat B, Bard P, Pfister P, Pageaux B. Mental fatigue induced by prolonged motor imagery increases perception of effort and the activity of motor areas. *Neuropsychologia*. 2021 2021;150:107701.
75. Siemionow V, Yue GH, Ranganathan VK, Liu JZ, Sahgal V. Relationship between motor activity-related cortical potential and voluntary muscle activation. *Experimental Brain Research*. 2000;133(3):303-11.
76. Kozlowski B, Pageaux B, Hubbard EF, Peters BS, Millar PJ, Power GA. Perception of effort during an isometric contraction is influenced by prior muscle lengthening or shortening. *European Journal of Applied Physiology*. 2021:1-12.
77. Marcora SM. Role of feedback from Group III and IV muscle afferents in perception of effort, muscle pain, and discomfort. *Journal of Applied Physiology*. 2011;110(5):1499-.

78. Cook DB, O'Connor PJ, Eubanks SA, Smith JC, Lee M. Naturally occurring muscle pain during exercise: assessment and experimental evidence. *Medicine and science in sports and exercise*. 1997;29(8):999-1012.
79. O'Connor PJ, Cook DB. 5 Exercise and Pain: The Neurobiology, Measurement, and Laboratory Study of Pain in Relation to Exercise in Humans. *Exercise and sport sciences reviews*. 1999 1999;27(1):119-66.
80. Broxterman RM, Layec G, Hureau TJ, Morgan DE, Bledsoe AD, Jessop JE, et al. Bioenergetics and ATP synthesis during exercise: role of group III/IV muscle afferents. *Medicine and science in sports and exercise*. 2017 2017;49(12):2404.
81. Broxterman RM, Hureau TJ, Layec G, Morgan DE, Bledsoe AD, Jessop JE, et al. Influence of group III/IV muscle afferents on small muscle mass exercise performance: a bioenergetics perspective. *The Journal of physiology*. 2018 2018;596(12):2301-14.
82. Sidhu SK, Weavil JC, Venturelli M, Garten RS, Rossman MJ, Richardson RS, et al. Spinal μ -opioid receptor-sensitive lower limb muscle afferents determine corticospinal responsiveness and promote central fatigue in upper limb muscle. *The Journal of physiology*. 2014 2014;592(22):5011-24.
83. Staiano W, Bosio A, de Morree HM, Rampinini E, Marcora S. The cardinal exercise stopper: Muscle fatigue, muscle pain or perception of effort? *Progress in brain research: Elsevier*; 2018. p. 175-200.
84. Barbosa TC, Vianna LC, Fernandes IA, Prodel E, Rocha HNM, Garcia VP, et al. Intrathecal fentanyl abolishes the exaggerated blood pressure response to cycling in hypertensive men. *The Journal of physiology*. 2016 2016;594(3):715-25.
85. Sidhu SK, Weavil JC, Rossman MJ, Jessop JE, Bledsoe AD, Buys MJ, et al. Exercise pressor reflex contributes to the cardiovascular abnormalities characterizing: hypertensive humans during exercise. *Hypertension*. 2019 2019;74(6):1468-75.
86. Müller J. *Elements of physiology*. London, England: Lea and Blanchard; 1842.
87. Bain A. *The senses and the intellect*: Longman, Green, Longman, Roberts, and Green; 1864.
88. Bastian HC. *The brain as an organ of mind*: Appleton; 1880.
89. James W. *The feeling of effort*. Anniversary Memoirs of the Boston Society of Natural History. Boston, MA: The Society of Natural History; 1880. p. 3-32.
90. Craig AD. How do you feel? Interoception: the sense of the physiological condition of the body. *Nature reviews neuroscience*. 2002 2002;3(8):655-66.
91. Craig AD. Interoception: the sense of the physiological condition of the body. *Current opinion in neurobiology*. 2003 2003;13(4):500-5.
92. Pollak KA, Swenson JD, Vanhaitsma TA, Huguen RW, Jo D, Light KC, et al. Exogenously applied muscle metabolites synergistically evoke sensations of muscle fatigue and pain in human subjects. *Experimental physiology*. 2014 2014;99(2):368-80.
93. Berchicci M, Menotti F, Macaluso A, Di Russo F. The neurophysiology of central and peripheral fatigue during sub-maximal lower limb isometric contractions. *Frontiers in human neuroscience*. 2013 2013;7:135.
94. Guo F, Sun Y-J, Zhang R-H. Perceived exertion during muscle fatigue as reflected in movement-related cortical potentials: an event-related potential study. *Neuroreport*. 2017 2017;28(3):115-22.
95. Cerqueira V, De Mendonça A, Minez A, Dias AR, De Carvalho M. Does caffeine modify corticomotor excitability? *Neurophysiologie Clinique/Clinical Neurophysiology*. 2006 2006;36(4):219-26.
96. Kalmar JM. The influence of caffeine on voluntary muscle activation. *Medicine and science in sports and exercise*. 2005;37(12):2113-9.
97. Tarnopolsky MA. Effect of caffeine on the neuromuscular system—potential as an ergogenic aid. *Applied physiology, nutrition, and metabolism*. 2008;33(6):1284-9.

98. Zénon A, Sidibé M, Olivier E. Disrupting the supplementary motor area makes physical effort appear less effortful. *Journal of Neuroscience*. 2015;35(23):8737-44.
99. Nicolò A, Marcora SM, Sacchetti M. Respiratory frequency is strongly associated with perceived exertion during time trials of different duration. *Journal of sports sciences*. 2016;34(13):1199-206.
100. Friedman DB, Brennum J, Sztuk F, Hansen OB, Clifford PS, Bach FW, et al. The effect of epidural anaesthesia with 1% lidocaine on the pressor response to dynamic exercise in man. 1993 1993;470(1):681-91.
101. Halperin I, Emanuel A. Rating of perceived effort: methodological concerns and future directions. *Sports Medicine*. 2020 2020;50(4):679-87.
102. Angius L, Pageaux B, Crisafulli A, Hopker J, Marcora SM. Ischemic preconditioning of the muscle reduces the metaboreflex response of the knee extensors. *European Journal of Applied Physiology*. 2021:1-15.
103. Pageaux B, Angius L, Hopker JG, Lepers R, Marcora SM. Central alterations of neuromuscular function and feedback from group III-IV muscle afferents following exhaustive high-intensity one-leg dynamic exercise. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 2015;308(12):R1008-R20.
104. Marcora SM. Can doping be a good thing? Using psychoactive drugs to facilitate physical activity behaviour: Springer; 2016.

Fig. 1: Overview of the two lines of research investigating the perception of effort in exercise sciences and the primary definitions used in the literature. In blue, the perception of effort does not include other exercise-related perceptions and is investigated as a construct dissociated from other perceptions. This approach follows Borg’s original definition conceptualizing the perception of effort as one’s appreciation of the difficulty of a task (how *hard* it is). In green, the perception of effort includes other exercise-related perceptions and is investigated as a construct encompassing a mix of exercise-related perceptions. This approach results from the original description proposed by Borg.

860

861

862 **Figure 2:** Panel A illustrates the coding process used for the classification of the articles included in the
 863 systematic review. Panel B illustrates the changes overtime in the use of the perception of effort
 864 construct in the included studies.

Figure 3: Flow diagram of systematic review inclusion/exclusion adapted from the Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

Figure 4: Contour enhanced funnel plot for all effects (*i.e.*, rating of perception of effort as *effort dissociated*, *effort not-dissociated*, *unclear* color coded; see legend).

A. Submaximal Exercise Tasks - *effort dissociated* group

B. Submaximal Exercise Tasks - *effort not dissociated* group

C. Submaximal Exercise Tasks - *unclear* group

D. Submaximal Exercise Tasks - Overall model estimates for all codings, and combined

Figure 5: Forest plot for the effect of an epidural anesthesia on the perception of at submaximal exercise demands (comparison epidural vs. placebo or no intervention). Panel A illustrates the effect sizes for the *effort dissociated* group. Panel B illustrates the effect sizes for the *effort not dissociated* group. Panel C illustrates effect sizes for the *unclear* group. Panel D illustrates the overall effect sizes for all coding and combined. Standardized mean different with 95% confidence intervals are shown.

A. Maximal Exercise Tasks - *effort dissociated* group

B. Maximal Exercise Tasks - *effort not dissociated* group

C. Maximal Exercise Tasks - *unclear* group

D. Maximal Exercise Tasks - Overall model estimates for all codings, and combined

Figure 6: Forest plot for the effect of an epidural anesthesia on the perception of effort at maximal demands (comparison epidural vs. placebo or no intervention). Panel A illustrates the effect sizes for the *effort dissociated* group. Panel B illustrates the effect sizes for the *effort not dissociated* group. Panel C illustrates effect sizes for the *unclear* group. Panel D illustrates the overall effect sizes for all coding and combined. Standardized mean different with 95% confidence intervals are shown.

Table 1. Risks of bias within included studies

	Design	Confounders	Blinding		Data collection methods			Intervention integrity		Overall label	
	Label	Label	Assessor	Subjects	Label	Valid	Reliable	Label	Consistency	Contamination	Label
<i>Ammann et al., 2010</i>	S	S	?	Y	M	?	?	W	Y	Y	M
<i>Amann et al., 2011</i>	S	S	?	Y	M	?	?	W	Y	Y	M
<i>Amann et al., 2014</i>	M	M	N	N	W	?	?	W	Y	Y	W
<i>Barbosa et al., 2014</i>	M	S	N	N	W	?	?	W	N	Y	W
<i>Blain et al., 2016</i>	S	S	N	N	W	?	?	W	N	Y	W
<i>Broxterman et al., 2017</i>	S	S	N	N	W	?	?	W	N	Y	W
<i>Broxterman et al., 2018</i>	S	S	N	N	W	?	?	W	N	Y	W
<i>Fernandes et al., 1990</i>	S	W	N	N	W	Y	Y	S	Y	Y	W
<i>Friedman et al., 1993</i>	M	W	N	N	W	Y	Y	S	Y	Y	W
<i>Gagnon et al., 2012</i>	S	M	Y	Y	S	N	N	W	N	Y	M
<i>Kjaer et al., 1989</i>	S	M	N	N	W	Y	Y	S	Y	Y	M
<i>Kjaer et al., 1999</i>	M	M	N	N	W	Y	Y	S	Y	Y	W
<i>Mitchell et al., 1989</i>	M	W	N	N	W	Y	Y	S	Y	Y	W
<i>Olson et al., 2014</i>	S	S	?	Y	M	N	N	W	Y	Y	M
<i>Sidhu et al., 2014</i>	M	W	N	N	W	N	N	W	N	Y	W
<i>Sidhu et al., 2019</i>	S	M	N	N	W	?	?	W	N	Y	W
<i>Smith et al., 2003</i>	M	M	N	N	W	Y	Y	S	Y	Y	W
<i>Smith et al., 2020</i>	S	W	N	Y	M	?	?	W	Y	Y	W

W: weak; M: moderate; S: strong; N: no; Y: yes; ?: unclear

"the feeling of how heavy, strenuous and laborious exercise is" Borg

"The particular feeling of energy that is being exerted... accompanied by a sense of strain and labour, that intensifies the harder a person tries" Preston & Wegner

"the conscious sensation of how hard, heavy, and strenuous a physical task is" Marcora

Effort dissociated

Effort not dissociated

1962

1996

2009

2010

"the sensation from the organs of circulation and respiration, from the muscles, the skin, the joints and force" Borg

"the subjective intensity of effort, strain, discomfort, and/or fatigue that is experienced during exercise" Noble and Robertson

A.

B.

Records identified through database
searching and other sources
(n = 902)

MEDLINE	200
EMBASE	455
SPORTDiscuss	10
CINHAL	13
Web of Science	147
PsycINFO	77

Duplicates removed
(n = 319)

Records after duplicates removed
(n = 583)

Records screened for
relevance in title and
abstract
(n = 583)

Record excluded
(n = 503)

Full-text articles screened
for eligibility
(n = 80)

Full-text articles excluded, with reasons
(n = 62)

- Wrong outcomes (n = 30)
- Wrong type of publication (n = 20)
- Wrong intervention (n = 9)
- Wrong study design (n = 1)
- Self-paced exercise (n = 2)

Qualitative analyses : 18
Quantitative analyses : 16

• Data not available for
quantitative analyses (n = 2)

Funnel plot of all studies (combined *effort*, *gestalt*, and *unclear* RPE scores)

A. Submaximal Exercise Tasks - *effort dissociated* group

B. Submaximal Exercise Tasks - *effort not dissociated* group

C. Submaximal Exercise Tasks - *unclear* group

D. Submaximal Exercise Tasks - Overall model estimates for all codings, and combined

A. Maximal Exercise Tasks - *effort dissociated* group

B. Maximal Exercise Tasks - *effort not dissociated* group

C. Maximal Exercise Tasks - *unclear* group

D. Maximal Exercise Tasks - Overall model estimates for all codings, and combined