The influence of non-neural factors on BOLD signal magnitude

Jan W. Kurzawski¹, Omer Faruk Gulban^{2,3}, Keith Jamison⁴, Jonathan Winawer^{1*}, Kendrick Kay^{5*}

¹Department of Psychology, New York University, New York, NY, 10003, USA

²Cognitive Neuroscience Department, Faculty of Psychology and Neuroscience, Maastricht University, 62229, The Netherlands

³Brain Innovation, Maastricht, 62229, The Netherlands

⁴Department of Radiology, Weill Cornell Medicine, New York, NY, 10021, USA

⁵Center for Magnetic Resonance Research, Department of Radiology, University of Minnesota, Minneapolis, MN, 55455, USA

*Co-senior author

Corresponding Author: Jan W. Kurzawski, Department of Psychology, 6 Washington Place, New York University, NY, 10003 (jan.kurzawski@nyu.edu)

Keywords: BOLD signal, fMRI, vasculature, horizontal meridian, vertical meridian, cerebral sinus.

Number of pages: 25

Number of figures: 10

Number of words for: abstract | introduction | discussion: 238/250 | 648/650 | 1427/1500

Conflict of interest: The authors declare no competing financial interests.

Acknowledgments: This research was funded by the US National Eye Institute R01-EY027964, R01-MH111417 and R01-EY027401 to JW. Collection of the NSD dataset was supported by NSF CRCNS grants IIS-1822683 (to KK) and NSF IIS-1822929.

Abstract

To what extent is the size of the blood-oxygen-level-dependent (BOLD) response influenced by factors other than neural activity? In a re-analysis of three neuroimaging datasets, we find large systematic inhomogeneities in the BOLD response magnitude in primary visual cortex (V1): stimulus-evoked BOLD responses, expressed in units of percent signal change, are up to 50% larger along the representation of the horizontal meridian than the vertical meridian. To assess whether this surprising effect can be interpreted as differences in local neural activity, we quantified several factors that potentially contribute to the size of the BOLD response. We find strong relationships between BOLD response magnitude and cortical thickness, cortical curvature, and the presence of large veins. These relationships are consistently found across subjects and suggest that variation in BOLD response magnitudes across cortical locations reflects, in part, differences in anatomy and vascularization. To compensate for these factors, we implement a regression-based correction method and show that after correction, BOLD responses become more homogeneous across V1. The correction reduces the horizontal/vertical difference by about half, indicating that some of the difference is likely not due to neural activity differences. Additionally, we find that while the cerebral sinuses overlap with the vertical meridian representation in V1, they do not explain the observed horizontal/vertical difference. We conclude that interpretation of variation in BOLD response magnitude across cortical locations should consider the influence of the potential confounding factors of cortical thickness, curvature, and vascularization.

Significance statement

The magnitude of the BOLD signal is often used as a surrogate of neural activity, but the exact factors that contribute to its strength have not been studied on a voxel-wise level. Here, we examined several anatomical and measurement-related factors to assess their relationship with BOLD magnitude. We find that BOLD magnitude correlates with cortical anatomy and macrovasculature. To remove the contribution of these factors, we propose a simple, data-driven model that can be used in any functional magnetic resonance imaging (fMRI) experiment. After accounting for the confounding factors, BOLD magnitude becomes more spatially homogenous. Our correction method improves the ability to make more accurate inferences about local neural activity from fMRI data.

Introduction

The blood-oxygen-level-dependent (BOLD) signal measured by fMRI is an important tool for non-invasive study of the human nervous system. However, the neural mechanisms underlying BOLD remain an active area of investigation (Herman et al., 2017). One clear conclusion is that the BOLD signal is strongly influenced by neural activity (Arthurs et al., 2000; Heeger et al., 2000; Attwell and Iadecola, 2002; Heeger and Ress, 2002; Logothetis, 2002; Lee et al., 2010; Siero et al., 2014). For a given location in the brain, and within a constrained paradigm (e.g., viewing different images and measuring the response that they elicit in visual cortex), the BOLD signal magnitude appears to be lawfully related to basic

measures of neural activity. For example, as stimulus contrast increases, neural firing rates and BOLD magnitude increase in proportion (Heeger et al., 2000). Similarly, increase in coherence of stimulus motion boosts BOLD magnitude and firing rates in V5/MT (Britten et al., 1993; Rees et al., 2000). When comparing different experimental paradigms or different brain locations, however, it is less clear how to interpret differences in the magnitude of the BOLD signal. For example, seeing a stimulus and expecting a stimulus can both elicit robust BOLD signals in V1, but the underlying neural activity is very different in the two paradigms (Sirotin and Das, 2009; Herman et al., 2017). It is also the case that similar BOLD signal magnitudes in two locations may be linked to very different underlying neural activity. These two limitations are reviewed by (Logothetis, 2008).

There are several reasons to believe that BOLD signal magnitudes, even within a fixed experimental paradigm, are influenced by factors that are not directly related to neural activity. The BOLD response, quantified as percent signal change above baseline, can be especially high in voxels containing large veins (Menon et al., 1993; Kim et al., 1994; Hoogenraad et al., 1999; Kay et al., 2019) or unusually low, delayed, and/or displaced in voxels near cerebral sinuses (Winawer et al., 2010; Jamison et al., 2017). The choice of MRI sequence, field strength (van der Zwaag et al., 2009), and sequence parameters like echo time (Gorno-Tempini et al., 2002) can also affect BOLD signal magnitude, and these effects may vary across the brain (Herman et al., 2017). Indeed, it has been reported that BOLD may vary across the cortex up to 40% simply due to different orientation of vasculature relative to the direction of the static magnetic field (Gagnon et al., 2015; Gagnon et al., 2016). Furthermore, recent high-resolution fMRI studies have shown that BOLD signal magnitude clearly depends on cortical depth. It is highest in the superficial layers which are positioned near large pial veins and decreases with cortical depth (Polimeni et al., 2010; Koopmans et al., 2011; Zimmermann et al., 2011; Yu et al., 2014; Fracasso et al., 2016a; Fracasso et al., 2016b; Dumoulin, 2017; Dumoulin et al., 2018; Kay et al., 2019; Self et al., 2019; van Dijk et al., 2020).

In this paper, we study variations in BOLD signal magnitude within a fixed paradigm, focusing our efforts on primary visual cortex (V1). We believe that by focusing on a single brain region in well-controlled visual paradigms, we are in the best position to derive sound interpretations of differences in BOLD signal magnitudes across the cortex. In three distinct datasets, we demonstrate large differences between the meridional locations: the BOLD magnitude in V1 is up to 50% higher along the representation of the horizontal meridian than along the representation of the vertical meridian. We then investigate the potential basis of these inhomogeneities by analyzing factors that are in principle distinct from neural activity. We find that several of these factors are systematically related to observed variation in BOLD magnitudes across V1. To remove their influence, we propose a simple correction method and show that the correction increases BOLD signal homogeneity across V1, reducing the horizontal-vertical difference by about half.

Methods

Datasets

We used three publicly available visual fMRI datasets: the Human Connectome Project (HCP) 7T Retinotopy Dataset (Benson et al., 2018), the Natural Scenes Dataset (NSD) (Allen et al., 2021), and the Temporal Decomposition Method (TDM) Dataset (Kay et al., 2020). All data were acquired on 7T MR scanners using gradient-echo pulse sequences (technical details provided in Supplementary Table 1). The datasets varied in stimulus properties and experimental design. HCP stimuli consisted of rings, wedges, and bars in a retinotopic mapping experiment; NSD stimuli consisted of natural scene images; and TDM stimuli consisted of high-contrast rings presented at different eccentricities. Experimental details are shown in **Figure 1**. The analyses performed in this paper start with pre-processed data from each dataset (details below).

Temporal Decomposition Method Dataset (TDM)

- 5 subjects
- $0.8 \times 0.8 \times 0.8 \text{ mm}^3$
- Rings varying in eccentricity
- 9 runs per subject
- $TR = 2.2 \text{ s}$
- $3.5 \text{ s ON} / 0.5 \text{ s OFF}$
- Maximum eccentricity = 5.5 degrees

Natural Scenes Dataset (NSD)

- 8 subjects
- $1.8 \times 1.8 \times 1.8 \text{ mm}^3$
- Natural scene images
- 360-480 runs per subject
- $TR = 1.6 \text{ s}$
- $3 \text{ s ON} / 1 \text{ s OFF}$
- Maximum eccentricity = 4.2 degrees

Human Connectome Project (HCP)

- 181 subjects
- $1.6 \times 1.6 \times 1.6 \text{ mm}^3$
- Textures viewed through spatial apertures
- 6 runs per subject
- $TR = 1 \text{ s}$
- Continuous design
- Maximum eccentricity = 8 degrees

Figure 1 – Datasets used in this study. Stimulus images for each of the datasets are shown. For TDM, stimuli consisted of 6 rings varying in eccentricity. For NSD, stimuli consisted of natural scene images. For HCP, the experiment consisted of several retinotopic mapping runs that included expanding and contracting rings, rotating wedges, and moving bars filled with a colorful object-based texture. Additional acquisition details are provided in Supplementary Table 1.

Extracting BOLD magnitude

From each dataset, we extracted a measure of BOLD signal magnitude at each cortical surface vertex. For TDM, we started with the pre-processed fMRI time-series data provided with the dataset and analyzed the data with a GLM. Specifically, we convolved a canonical HRF with stimulus onsets to create a regressor for each experimental condition, and then

used these regressors with GLMdenoise (Kay et al., 2013b) to estimate a beta weight for each condition. We computed the maximum beta weight across all conditions for each voxel as the measure of BOLD signal magnitude. These results are defined at six different cortical depths (equidistant from 10% to 90% of the cortical depth) in each subject's native surface space. (Depth assignment was achieved by a spatial interpolation of each fMRI volume at the locations of the six depth-dependent cortical surfaces; see Kay et al. (2020) for details.) For NSD, we took the 'meanbeta' values (1 mm data preparation, beta version 2) provided with the dataset; these values indicate the average BOLD percent signal change observed across all stimulus trials and all scan sessions. We then mapped these values to the 3 cortical depth surfaces provided in NSD (positioned at 25%, 50%, and 75% of the cortical depth), and then averaged across depth to produce one value per surface vertex in the subjects' native surface space. The HCP dataset was previously analyzed (Benson et al., 2018) with a population receptive field (pRF) model (Dumoulin and Wandell, 2008) implemented in analyzePRF (Kay et al., 2013a). The model includes a gain parameter that describes the amplitude of the BOLD response of a given voxel (or vertex) to the object-based texture (covering the entire pRF) for a single repetition time ($TR = 1$ s). We quantified BOLD in terms of percent signal change (%BOLD) by dividing the gain parameter by mean signal intensity and multiplying by 100. The results are prepared in FreeSurfer's fsaverage space.

Visual field mapping

We used retinotopic mapping to divide the primary visual cortex into a set of regions. For HCP, we used polar angle and eccentricity estimates available from the data release. For the TDM and NSD datasets, we mapped Benson's polar angle and eccentricity atlas using neuropythy software (Benson and Winawer, 2018). We use the following convention for all 3 datasets: the upper vertical meridian corresponds to 0 deg, the horizontal meridian corresponds to 90 deg, and the lower vertical meridian corresponds to 180 deg. Note that the polar angle estimates are rescaled for the correlation and linear regression analysis (see next section). We used Benson's definition of the extent of visual areas V1, V2, and V3 for all 3 datasets (Benson et al., 2014).

Quantification of non-neural factors

For each dataset, we quantified a number of factors that might be related to variation in the magnitude of the BOLD signal across cortical locations. These factors include the following:

1. *Cortical curvature* was obtained from FreeSurfer outputs (Dale et al., 1999; Fischl and Dale, 2000), and refers to the geometry of the folding pattern of the cortical surface. Negative values correspond to gyri while positive values correspond to sulci. Curvature is quantified as $1/r$, where r is the radius of an inscribed circle measured in mm.
2. *Cortical thickness* was also obtained from FreeSurfer outputs. It is measured in mm and corresponds to the distance between the outermost (close to cerebrospinal fluid) and innermost (close to white matter) boundaries of gray matter.
3. *Angle with respect to B_0* was calculated by considering the angle (θ) between the pial surface normal and the direction of the B_0 static magnetic field as estimated

from NIFTI header information. Angle was quantified in degrees and was normalized as $\text{abs}(\theta - 90)$ such that a final value of 0 deg indicates that the cortical surface is parallel to the magnetic field and a final value of 90 deg indicates that the cortical surface is perpendicular to the magnetic field.

4. *RF coil bias* was taken to be the result of fitting a 3D polynomial to the mean signal intensity in the fMRI data. The values are in raw scanner units. This method has been previously used (Kay et al., 2019).
5. *Mean bias-corrected EPI* was calculated as the mean signal intensity in the fMRI data divided by the RF coil bias (Kay et al., 2019). The units range from approximately 0 to 2, and indicate percentages (e.g., 0.5 means 50% of the strength of typical signal intensities).
6. *Polar angle (θ)* was obtained from Benson's atlas (Benson et al., 2014) and normalized as $\text{abs}(\theta - 90)$. After this normalization, 0 deg corresponds to the horizontal meridian, while 90 deg corresponds to the upper and lower vertical meridians.
7. *Depth* was estimated (in the TDM dataset) by generating 6 cortical surfaces equally spaced between 10% and 90% of the distance from the pial surface to the boundary between gray and white matter. These surfaces are numbered from 1–6, where 1 is outermost and 6 is innermost.

Definition of regions of interest

Using the visual field mapping results, we defined regions of interest (ROIs) corresponding to the representation of the horizontal and vertical meridians within V1. The ROIs were defined by limiting the eccentricity to the maximum stimulus eccentricity used in each dataset and limiting the angle to a specific range (e.g., to create a V1 ROI for the upper vertical meridian with a width of 20 deg, we created a mask where polar angle estimates were higher than 0 and lower than 20).

Modelling variation in BOLD signal magnitude

To account for non-neural contribution to %BOLD, we used a multiple regression model. The modeled data (Y) consisted of the %BOLD value observed at each surface vertex in visual areas V1–V3. Although this study focuses on BOLD homogeneity in V1, we include %BOLD in V1–V3. This is because we are attempting to establish relationships that might generalize across different cortical regions. Furthermore, if we were to include only vertices in V1, we would be at high risk of removing genuine neural activity differences (e.g. those that may exist between the horizontal and vertical meridians) that correlate with the non-neural factors.

The variables used to model the data included cortical thickness, cortical curvature, and mean bias-corrected EPI intensity. (Only these three factors showed evidence of being substantially related to BOLD magnitude; see Results.) The variables were standardized (z -scored) and, together with a constant term, were included as predictors in the design matrix (X). Ordinary least-squares estimates for beta weights were obtained in the following linear model:

$$Y = X\beta + n \quad (\text{Eq. 1})$$

where Y is the %BOLD magnitude at each vertex, X is the 4-column design matrix, β is a set of beta weights (4 per vertex), and n is a set of residuals.

Major cortical sulci

In several figures we show outlines of major cortical sulci. These include the calcarine sulcus (CALC), parieto-occipital sulcus (POS), intraparietal sulcus (IPS), occipitotemporal sulcus (OTS), and superior temporal sulcus (STS). These sulci were manually labelled on the fsaverage surface and then mapped to each individual's native surface.

Atlas of cerebral sinuses

The cerebral sinuses are the largest draining veins in the brain and are known to cause artifacts in fMRI data (Winawer et al., 2010). To aid in identifying the location of the sinuses, we first created an atlas of the sinuses using the T1- and T2-weighted volumes from the HCP1200 dataset (Van Essen et al., 2012). Specifically, we performed the following steps:

- 1) We obtained T1-weighted and T2-weighted volumes from the 494 subjects from the HCP1200 dataset for which twin pairings are available. We used the versions of these volumes that are already nonlinearly transformed to MNI space.
- 2) Because the sinuses have high intensity (bright) in T1-weighted volumes and low intensity (dark) in T2-weighted volumes, we divided T1 intensity values by T2 intensity values to accentuate the locations of the sinuses. We refer to the resulting ratios as the sinus map.
- 3) The sinus map from each subject was intensity normalized, trimmed for outliers, and averaged across subjects to produce a group-average sinus map (**Figure 2A**).
- 4) We manually defined four control points on the group-average sinus map that mark the geometrical termination of the four sinus segments, namely, the superior sagittal sinus, the straight sinus, and the left and right transverse sinus. Using the fast-marching method (Sethian, 1996), we traced paths that follow the sinus between the control points and used the branching of these paths to define the sinus segments (**Figure 2B**).

The primary outputs of this atlas creation procedure are 1-mm volumes in MNI space that represent likely locations of sinus voxels as well as their segment labels (**Figure 2C**). These resources may be useful for other studies and are freely available at <https://osf.io/2nc4x/>. In the current study, we used the sinus atlas to identify cortical locations that may be affected by sinus artifacts. Specifically, we prepared a binary version of the sinus atlas in the native volumetric space of a given subject and then blurred this binary volume with a Gaussian kernel with full-width-half-maximum of 5 mm. This enables identification of cortical gray matter positioned at varying distances from the sinuses.

Figure 2 – Constructing an atlas of the cerebral sinuses. This figure illustrates our procedure for creating a sinus atlas using T1- and T2-weighted volumes prepared in MNI space. A) Procedure for creating a group-averaged sinus map. B) Procedure for finding the branches of the sinuses. C) The final representation of the sinus atlas. Different colors show different branches of the sinuses. The atlas is freely available at <https://osf.io/2nc4x/>.

Data and code availability

The datasets that are used in this paper are freely available online: NSD (<http://naturalscenesdataset.org>), HCP (<https://osf.io/bw9ec/>), and TDM (<https://osf.io/j2wsc/>). Code that reproduces the main figures in this paper is available at <https://github.com/jk619/meridianbias/>. Associated data files as well as the sinus atlas are available at <https://osf.io/2nc4x/>.

Results

Stronger BOLD responses along the V1 horizontal meridian

We examined BOLD response magnitudes in three freely available datasets: the Natural Scenes Dataset (NSD; Allen et al., 2021), the data used for the Temporal Decomposition Method (TDM; Kay et al., 2020), and the Human Connectome Project 7T Retinotopy Dataset (HCP; Benson et al., 2018). Each dataset contains BOLD responses to different types of visual stimulation (see Methods). We defined one region of interest (ROI) for the horizontal meridian (HM) and one for the vertical meridian (VM) (**Figure 3A–B**). These ROIs represent a wedge-shaped region in the visual field centered at the horizontal meridian with a width of 40 deg (horizontal) and two wedges abutting the vertical meridian each with a width of 20 deg (vertical).

In each of the three datasets, we compared BOLD magnitudes expressed in percent signal change (%BOLD) observed for the VM with BOLD magnitudes for the HM (**Figure 3C–E**). In each dataset, we find higher %BOLD in the HM ROIs compared to the VM ROIs. We summarize this difference with an asymmetry ratio: $(\text{HM} - \text{VM}) / \text{mean}(\text{HM}, \text{VM})$. All datasets

show strong asymmetry, with an asymmetry ratio of ~30%. Positive values for the asymmetry ratio indicate greater response for the horizontal meridian.

Figure 3 – BOLD magnitude is higher at the horizontal meridian in V1. **A)** Polar angle map of group-average HCP subject (999999) with V1 boundary outlined in dotted black lines. **B)** Horizontal and vertical regions of interest (ROIs) are indicated in gray and magenta, respectively. White text indicates major brain sulci (see Methods). **C-E)** Mean BOLD magnitude for horizontal and vertical ROIs in the three datasets. Error bars indicate standard error across subjects.

One possibility is that the horizontal and vertical V1 BOLD responses are in fact similar, but the vertical ROIs appear to have lower signal due to mixing with signal from V2. V2 and V1 border along the vertical meridian representation, and blurring might occur either in acquisition or in pre-processing and analysis. To further our understanding of the V1 response asymmetries, we re-computed asymmetry ratios using smaller wedges at many locations (**Figure 4A**). While the asymmetry is strongest at the cardinal meridians, some horizontal/vertical asymmetry is found at least 30 deg away from the meridians in all three datasets (**Figure 4B**). This argues against the explanation that the asymmetry is caused by spillover from V2.

Figure 4 – BOLD asymmetries generalize to off-cardinal locations. To further understand V1 BOLD asymmetry, we manipulated the location of the wedge ROIs in 5-deg increments. **A)** The upper row shows the visual field location of wedge ROIs and the lower row shows the corresponding cortical locations (flattened left hemisphere). For clarity, we show only every other set of ROIs. **B)** Asymmetry ratio as a function of angular distance from the cardinal meridians. Error bars indicate standard error across subjects.

HM/VM asymmetry persists at inner cortical depths

The BOLD signal is strongly influenced by properties of the brain's vasculature. Uneven venous contributions across the brain can cause variation in BOLD magnitude (Menon et al., 1993; Kim et al., 1994; Hoogenraad et al., 1999; Kay et al., 2019). One possibility is that the meridional asymmetries we observe arise from non-uniformities in the vascular network. To investigate this possibility, we took advantage of the sub-millimeter resolution of the TDM dataset and examined HM/VM asymmetry as a function of cortical depth. Because macroscopic venous effects are larger in the superficial cortex due to large pial veins (Duvernoy et al., 1981; Turner, 2002; Polimeni et al., 2010; Kay et al., 2019), by sampling BOLD responses from deeper cortical depths, we minimize contributions from pial veins. We find that the HM/VM asymmetry is larger at the superficial depths, suggesting that part of the asymmetry may be due to differential properties in macroscopic vasculature (**Figure 5**). This depth effect is systematic: every subject shows higher asymmetry at the superficial depth than the middle depth. Nonetheless, there remains a substantial horizontal/vertical asymmetry at all cortical depths (**Figure 5**), suggesting that macroscopic vessels near the pial surface are not the entire explanation. At the innermost cortical depth sampled, which is

least influenced by pial vessels, the HM/VM asymmetry is 26% (average across subjects) and is positive in each of the 5 subjects. The middle depths appear to have the least asymmetry. This could be due to a difference in neural responses at intermediate depths, which generally correspond to input-related cortical layers.

Figure 5 – BOLD asymmetry in V1 persists at inner cortical depths. We exploit the high-resolution TDM dataset to discriminate V1 BOLD responses across cortical depth and estimate response asymmetries as a function of depth (asymmetry is calculated in the same way as in **Figure 3**). The presence of asymmetry at the innermost cortical depth suggests that response asymmetries exist even with minimal contribution of large pial veins.

Assessing and modeling non-neural contributions to BOLD signal magnitude

In addition to vascular effects, other factors unrelated to neural activity evoked by the experimental manipulation may influence variation in %BOLD across the cortical surface. These additional factors are often neglected in fMRI analysis pipelines. Although some of the factors are known to vary across the cortex, their influence on the BOLD signal is poorly understood. Here, we attempt to understand how these factors may induce BOLD magnitude variations. To the best of our knowledge, we are unaware of any previous study that has examined this issue in detail, especially at the level of individual voxels (or vertices) within individual subjects.

We first identified a list of possible confounding factors based on consideration of basic anatomical properties of the brain and the nature of fMRI measurement. These factors are cortical curvature, cortical thickness, radiofrequency (RF) coil bias, mean bias-corrected EPI signal intensity, and angle between the cortical surface and the direction of the static magnetic field B0. Each of these factors can be interpreted as spatial maps, with a value at each vertex on the cortical surface mesh. The five maps can be obtained from standard

anatomical scans (T1-weighted) or from the fMRI measurements themselves without additional MRI experiments (see Methods for details). Example surface visualizations of these maps together with %BOLD are shown in **Figure 6**. We hypothesize that inhomogeneities in some of these maps might explain some of the observed inhomogeneity in %BOLD across V1.

Figure 6 – Anatomical and acquisition factors across cortex. Each sphere shows data mapped on the left hemisphere for subject S1 in the NSD dataset. Below each surface map is a histogram of the plotted values. White outlines indicate major cortical sulci. %BOLD represents the average response to the natural scene stimuli used in the NSD dataset. Some of the spatial variability in %BOLD might be due to variability in the depicted non-neural factors.

To understand the potential relationships amongst these five identified factors and %BOLD, we first performed voxel-wise correlation analyses. For these analyses, we used the TDM dataset, as its high spatial resolution facilitates the identification of vascular effects (Kay et al., 2019). We examined data from V1–V3 where neural activity magnitudes can be expected to be relatively homogeneous given the simple contrast patterns used. In **Figure 7A**, we show pairwise correlations across these five quantities, as well as retinotopic polar angle preference (rescaled between 0 and 90; see Methods) and cortical depth. We find that %BOLD correlates substantially with four factors: curvature ($r = 0.26$), thickness ($r = -0.17$), mean bias-corrected EPI intensity ($r = -0.25$), and depth ($r = -0.27$). We do not find a strong correlation between %BOLD and polar angle. Although results from **Figure 3C–E**, **Figure 4C** and **Figure 5** suggest a strong negative correlation, the previous analysis included data only from V1. Here we analyze vertices from V1–V3 where this relationship becomes weaker ($r = -0.05$). Overall, we can summarize as follows: %BOLD extracted from V1–V3 tends to be higher at locations that correspond to sulci, in thinner parts of the cortex, in voxels with lower mean bias-corrected EPI intensities, and at more superficial cortical depths. To assess the generality of these findings, we also examined the NSD dataset where substantially more cortex (beyond V1–V3) is activated by the natural scene images. Across the nsdgeneral ROI (drawn to cover visually responsive posterior cortex), we again find a correlation between %BOLD and curvature ($r = 0.12$; $p < 0.01$, one-sample t -test across subjects) and between %BOLD and thickness ($r = -0.25$; $p < 0.01$). Correlation with mean bias-corrected EPI is small ($r = -0.01$, likely due to the lower resolution of the NSD data (1.8-mm) which tends to obscure the macrovasculature.

Examination of correlations amongst factors (excluding %BOLD) yields additional insights. The strongest correlation that we find is between curvature and thickness ($r = -0.28$), indicating that sulci tend to be thin. Curvature is correlated with mean bias-corrected EPI ($r = 0.16$) and with polar angle ($r = -0.19$), and thickness is correlated with polar angle ($r = 0.15$). Our interpretation of these effects is that venous effects tend to be stronger in gyri (consistent with previous findings in Kay et al., 2019), and that the correlations related to polar angle simply reflect the tendency for horizontal meridian representations to fall on sulci (e.g. the calcarine sulcus). Overall, these complex relationships suggest that making sense of non-neural influences on %BOLD requires a broad perspective that considers multiple factors.

Figure 7 – Modeling variations in BOLD signal magnitude. A) Correlation (Pearson’s r) between neural and non-neural factors and %BOLD extracted from V1–V3 from the TDM dataset. Main plot shows results from data concatenated across all subjects, while inset plots show results from individual-subject data. P-values indicate significance of one sample t -test across subjects; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. **B)** Regression model for %BOLD. Based on the results of panel A, we selected curvature, thickness, and mean bias-corrected EPI as the main non-neural factors that confound %BOLD. These three factors were then used in a multiple linear regression model to predict %BOLD (top). The amount of variance explained by the model is shown in the inset.

Correcting BOLD signal magnitude for non-neural factors

We now explore whether we can develop a statistical model to compensate for the influence of non-neural factors on %BOLD. We operate under the assumption that any observed correlation between the factors and %BOLD is incidental and does not reflect genuine neural activity variation. For our model, we do not use cortical depth as a factor because it is not practically accessible in standard-resolution fMRI measurements and, more importantly, because aggressively removing its contribution would clearly remove true (linear) variations in depth-dependent activity. Our model is a multiple regression model (**Figure 7B, top**) that uses the main factors of curvature, thickness, and mean bias-corrected EPI intensity as continuous variables and attempts to determine a weighted sum of these factors that optimally accounts for variations in %BOLD across cortical locations (see Methods for details).

Fitting the model, we find a strong positive contribution of curvature and negative contributions of cortical thickness and EPI intensity (**Figure 7B, bottom**), consistent with the earlier voxel-wise correlation analyses. Estimated beta weights are fairly consistent across subjects, and the model on average across subjects explains 19% of the variance in %BOLD.

A multiple regression model using all 5 factors resulted in only minimally larger explained variance, 20% vs. 19%, consistent with the earlier correlation analyses indicating that RF coil bias and B0 angle bear little or no relationship with %BOLD.

To better understand the relationship between the identified non-neural factors and %BOLD, we construct a 2D histogram relating the model fit (BOLD prediction based on non-neural factors obtained by multiplying the design matrix and estimated beta weights) and %BOLD (**Figure 8A**). This reveals a clear nonlinear relationship. To accommodate this nonlinearity, we fit a nonlinear function relating the linear model fit and %BOLD (blue line in **Figure 8A**). Finally, we remove the contribution of non-neural factors by dividing %BOLD observed at each cortical location by the fit of the nonlinear model. The interpretation is that the influence of non-neural factors on %BOLD might impose a type of ‘gain’ field on fMRI responses observed in a given experiment. Note that our method of rescaling BOLD magnitudes does not change the pattern of responses across different experimental conditions within a voxel. For example, if the response to condition A is 25% higher than the response to condition B, this will continue to be the case after rescaling.

Figure 8 – Correction of V1 BOLD inhomogeneity. A) Removal of non-neural factors. First, linear combinations of non-neural factors are used to predict %BOLD within V1–V3 using the TDM dataset. The model is fit on data concatenated from all 5 TDM subjects. The model is augmented with a nonlinear power-law function (blue line), which is controlled by a gain parameter ($p(1)$), an exponent parameter ($p(2)$), and a constant term ($p(3)$). **B)** Each voxel’s BOLD responses are divided by the model fit, yielding the corrected %BOLD. **C)** BOLD signal magnitude within V1 before and after the correction (TDM dataset, subject S3, most superficial cortical depth). Asterisk indicates the fovea and dashed lines indicate the boundary between V1 and V2. After correction,

some vertices with very high BOLD are eliminated (see black arrows). Within each plot, the color range extends from 0 to the 99th percentile of the observed values.

The result of the proposed correction procedure is shown in **Figure 8B**. We see that after the correction procedure, the distribution of BOLD response becomes flatter, indicating the efficacy of the procedure. Increased homogeneity of BOLD magnitude is also visible on the cortical surface (**Figure 8C**). As a further check of the validity of our procedure, we quantified the homogeneity of BOLD magnitudes across V1 using the coefficient of determination (CoV; ratio of standard deviation to mean). We find that the CoV decreases from 88% before correction to 64% after correction (**Figure 9C**). We also summarize the effect of the correction by showing the correlations between %BOLD and non-neural factors before and after the correction (**Figure 9D**). Correlations after the correction are substantially reduced, though the correlation between %BOLD and depth remains strong.

We now return to the experimental effect that motivated this study, namely, BOLD response asymmetries across the horizontal and vertical meridians in V1. We quantify the asymmetry before (**Figure 9A**) and after (**Figure 9B**) the correction. We find that before accounting for the non-neural factors, the asymmetry is 41.7%, but after the correction, the asymmetry is reduced to 23.1%. We thus suggest that some of the observed differences in BOLD response magnitudes are due to non-neural factors, but the remaining differences may reflect genuine differences in local neural activity across these subregions of V1.

Figure 9 – The effect of BOLD inhomogeneity correction. **A)** Dependence of %BOLD on polar angle in V1 before correction (TDM dataset, results summarized across subjects and depths using local linear regression). Shaded area indicates the width of the wedge used to calculate the asymmetry ratio reported in the plot title.

B) Same as panel A, except showing results after correction. **C)** Coefficient of variation of %BOLD across V1 before and after correction (error bars indicate standard error across subjects). **D)** Voxel-wise correlation between the various factors and %BOLD before and after correction.

Cerebral sinuses do not explain horizontal/vertical asymmetry

The cerebral sinuses are the largest veins that drain blood away from the brain and are composed of three main branches (superior sagittal sinus, straight sinus, and transverse sinus) (**Figure 10A**). Due to their very large size, the cerebral sinuses introduce substantial magnetic field inhomogeneities that may result in BOLD signal dropout and/or distortion (Winawer et al., 2010; Jamison et al., 2017). One possibility is that sinus-related artifacts may be related to the observation of horizontal/vertical asymmetry. Specifically, it might be that the cerebral sinuses tend to be located near the vertical meridians, cause loss of BOLD signal, and therefore produce horizontal/vertical asymmetry. Modelling the effects of the sinus is difficult. First, both veins and sinuses result in low EPI intensity values, making them difficult to distinguish using conventional EPI data. Second, they have different effects on the BOLD signal. BOLD responses tend to be higher near small and moderately sized veins but tend to be lower (or non-existent) near the sinuses. Third, mapping the exact cortical location of the sinuses is challenging.

To assess the impact of the sinuses on our measurements, we constructed an MNI atlas of likely sinus locations (see Methods). We then mapped this atlas to the native space of individual subjects and identified surface vertices that are likely to be near the sinuses. Consistent with previous studies (Winawer et al., 2010), we find that several branches of the sinuses tend to overlap with primary visual cortex (**Figure 10B**) and more specifically with the representation of the vertical meridian (**Figure 10C–D**; results for all subjects can be found in **Supplementary Figure 1**). To test whether presumed signal loss from the sinuses might be responsible for horizontal-vertical asymmetry, we removed vertices close to the sinuses and re-computed horizontal-vertical asymmetry. Across different inclusion levels (removing voxels close to or further away from the sinuses), we find very little effect on the asymmetry (**Figure 10E–F**). Our interpretation is that the sinuses tend to co-localize with the VM, but this correspondence is not perfect (and is variable across subjects), and there are sufficient remaining VM vertices for which to compute horizontal-vertical asymmetry. We therefore conclude that horizontal-vertical asymmetry is not a simple consequence of BOLD signal loss induced by the sinuses.

Figure 10 – Assessing the effect of the cerebral sinuses on V1 response asymmetry. **A)** Sinus atlas in MNI space visualized against a 3D glass brain. There are three main branches (marked by different colors) with a confluence located close to the occipital pole. **B)** Sinus atlas mapped to the cortical surface of subject S5 from the TDM dataset (cut-off = 0.8). The underlying data are mean bias-corrected EPI intensity values. **C)** Same as panel B except the underlying data are polar angle estimates from the Benson atlas (thresholded at a stimulus eccentricity of 5.5 deg). **D)** Stacked histogram showing polar angle values from all subjects inside the V1 ROI (cut-off = 0.8). White bins indicate locations outside of the sinus atlas; colored bins indicate locations within the sinus atlas. **E)** Spatial extent of sinus influence. After blurring a binary version of the sinus atlas with a 5-mm FWHM Gaussian, we use different cut-off values to select different spatial extents of the influence of the sinuses. **F)** Corrected %BOLD obtained from the TDM dataset as a function of polar angle. Here we use different cut-off values to exclude sinus-related vertices and plot the resulting corrected %BOLD (local regression estimate; same as Figure 9A–B). For each cut-off value we calculate the asymmetry ratio and report the value in the text inset. Shaded area indicates the width of the wedge used to calculate the asymmetry.

Discussion

In this paper, we used three publicly available datasets to assess the degree of homogeneity of BOLD signal magnitude in primary visual cortex. We found that stimulus-evoked BOLD responses, expressed as percent signal change, are up to 50% stronger along the horizontal meridian than the vertical meridian. To investigate whether these magnitude differences can be attributed to differences in local neural activity, we systematically evaluated the potential contribution of several non-neural factors to the observed effect. We found that BOLD signal magnitude correlates with cortical curvature, cortical thickness, and macrovasculature (as indexed by bias-corrected EPI intensities). Using a regression-based correction procedure, we were able to increase the homogeneity of BOLD signal magnitude and found that the meridional differences were reduced by half.

Spatial variations in BOLD magnitude

This study tackles the issue of the neural basis of variation in BOLD signal magnitude. Specifically, we address variation in BOLD across cortical locations for a fixed experimental manipulation, as opposed to variation in BOLD across experimental manipulations for a fixed cortical location. The latter has been heavily studied (Heeger et al., 2000; Logothetis et al., 2001; Heeger and Ress, 2002; Logothetis and Wandell, 2004; Mishra et al., 2021), whereas the former has not yet been systematically studied to the best of our knowledge. If there are indeed non-neural factors that influence BOLD signal variation, taking this into account is critical when interpreting differences in fMRI responses across brain regions. We acknowledge that a challenge in understanding the neural basis of the BOLD signal is that directly comparable ground-truth measurements of neural activity are typically not available. In our analyses, we relied on the working assumption that the experimental paradigms of the three datasets (combined with suitable averaging and analysis procedures) are expected to generate relatively homogeneous patterns of neural activity in early visual cortex. Of course, this may not be exactly the case.

We note that some analyses of fMRI data are insensitive to spatial variation of BOLD magnitude. For example, population receptive field models incorporate an explicit gain parameter that is distinct from other parameters (such as receptive field position) in the model. As such, such approaches can discount the gain observed at a given voxel. However, in general, conventional approaches to fMRI analysis in which the size of the BOLD response is used for inference (including subtraction-based analyses) are susceptible to the influence of non-neural factors on BOLD signal magnitude.

Non-neural factors that affect BOLD magnitude

Cortical anatomy. We find that cortical curvature and cortical thickness correlate with BOLD signal magnitude (see **Figure 7A**). It is known that many anatomical properties vary with thickness and with curvature (Jiang et al., 2021): (i) total neuron count is higher in gyri than it is in sulci (Hilgetag and Barbas, 2005), (ii) gyri tend to be thicker than sulci (Welker, 1990; Hilgetag and Barbas, 2005), (iii) venous effects (resulting in higher BOLD signal amplitude) are more prominent in gyri than they are in sulci (Kay et al., 2019); and (iv) there may even be intrinsic causal relationships between curvature and thickness during anatomical development (Hilgetag and Barbas, 2005).

However, the exact anatomical and biophysical mechanisms that might link curvature and thickness to BOLD signal magnitudes and whether they reflect purely non-neural sources of variance or might be linked to variations in neural activity is largely unknown, to our knowledge. This is an important issue for future research. Here, we operate under the working assumption that correlations between the BOLD signal and cortical curvature or thickness reflect incidental factors unrelated to local neural activity. We therefore assume that a correction which removes their influence from the BOLD signal is desirable.

Cerebral sinuses. One possible explanation of the horizontal/vertical asymmetry pertains to the severe magnetic field inhomogeneities introduced by the cerebral sinuses. Consistent with Winawer et al. (2010; their figure 10), we find that the transverse sinus is close to the upper vertical meridian of V1 (ventral bank of calcarine sulcus) and that vertices near the sinus tend to be darker (lower mean EPI). Although the sinus is known to corrupt the fMRI signal (Winawer et al, 2010), we found that removing voxels near the sinus did not eliminate the horizontal/vertical asymmetry. The atlas we create in this paper helps identify cortical regions likely to be influenced by the cerebral sinuses. Note that the location of the sinuses is fairly consistent across subjects in volume-based MNI space but is quite variable across subjects when projecting to individual subjects' native surfaces (see **Supplementary Figure 1**). We expect this atlas will prove valuable for studying sinus-related regions in future studies.

Orientation of pial veins. It has been reported that regions where the cortical surface is oriented perpendicular to the main magnetic field produce lower BOLD signal than regions where the surface is oriented parallel (Gagnon et al., 2015; Fracasso et al., 2018). The proposed explanation is that this effect is caused by the orientation of pial veins, which lie parallel to the cortical surface. Our analyses did not replicate this result and indicated little relationship between BOLD magnitude and B0 angle (see **Figure 7A**). One possible explanation could be related to the pre-processing pipeline for the TDM dataset, in which fMRI signals were sampled in the gray matter at 10% or more of the cortical thickness away from the pial surface. This may have dampened effects related to pial veins, though a detailed examination would be necessary to resolve these discrepancies.

RF coil effects. Due to cortical folding, gyri tend to be closer to the RF coil than sulci. Locations that are further from the coil might have lower mean signal intensities and therefore lower SNR (Srirangarajan et al., 2021), but this should not affect BOLD *magnitudes* expressed in terms of percent signal change. We are not aware of any mechanism that would alter the percent signal change in brain locations that are further away from the RF coil (e.g., lower BOLD responses in sulci compared to gyri). Indeed, we did not find any relationship between RF coil bias and BOLD magnitude (see **Figure 7A**).

Correction for the impact of non-neural factors

Our results show that voxel-wise %BOLD is likely contaminated by several non-neural factors. To account for these factors, we developed a regression-based correction method. The goal of this method was to introduce a simple, data-driven approach that can be applied irrespectively of the specific experiment or brain region that is under consideration. The underlying premise of the method is that by removing the contribution of non-neural factors, the resulting measures would constitute a better representation of the underlying neural activity. After application of the method, we found that %BOLD becomes more homogenous and correlations between %BOLD and non-neural factors become significantly reduced. Thus, our results indicate that some variation in %BOLD that might be interpreted as change in neural activity likely reflects the variation of non-neural factors. We believe the results presented in this paper constitute a first step towards developing a cogent strategy for compensating for non-neural biases in BOLD signal magnitudes.

There are other methods that can be used to suppress the contribution of non-neural factors to BOLD signal magnitudes. By identifying early and late components of evoked hemodynamic responses, a temporal decomposition method can be used to estimate BOLD response components more closely linked to the microvasculature, which presumably more closely reflect local neural activity (Kay et al., 2020). Another analysis method focuses on BOLD fluctuations where estimates of slow oscillations (< 0.1 Hz) are used to suppress vascular-related effects (Kazan et al., 2016). Similarly, some methods use the amplitude of fluctuations in resting-state data to rescale the BOLD signal (Di et al., 2013; Guidi et al., 2020). Finally, acquisition methods, such as spin-echo pulse sequences, can be used to suppress unwanted venous effects. Note that all these methods concern effects of the macrovasculature, but systematic biases in BOLD signal magnitudes may in theory persist even if BOLD responses were restricted to the microvasculature. Further investigation is necessary to resolve these possibilities.

More generally, our findings here are part of a broader effort to make meaningful inferences about the underlying biology from MRI measurements. While the pixel intensity of most MR images is expressed in arbitrary units, there has been some progress in removing the influence of unwanted factors to quantify specific properties of the measured tissue. For example, using anatomical imaging, one can estimate the macromolecular tissue volume within a voxel (Mezer et al., 2013), a quantity that has clinical relevance for neurodegenerative diseases. Our study is a step towards linking BOLD amplitude more directly to neural activity.

Author Contributions

J.W.K, K.K. and J.W. conceived and designed this study, K.K. acquired the data. J.W.K and K.J. performed data analyses, with K.K. and J.W. providing guidance. J.W.K., J.W, O.F.G and K.K. wrote and edited the manuscript.

Dataset	TDM	NSD	HCP
Field strength	7T	7T	7T
TR	2200 ms	1600 ms	1000 ms
TE	22.4 ms	22.0 ms	22.2 ms
Flip angle	80	62	45
Number of slices	84	84	85
Matrix size	200 × 162	120 × 120	130 × 130
Field of view	160 mm × 129.6 mm	216 mm × 216 mm	208 mm × 208 mm
Nominal spatial resolution	0.8 mm	1.8 mm	1.6 mm
Multiband factor	2	3	5
iPAT factor	3	2	2
Partial Fourier	6/8	7/8	7/8

Supplementary Table 1 – Details on the fMRI pulse sequence used in each of the datasets. Each column describes different dataset.

Supplementary Figure 1 – The location of the cerebral sinuses relative to primary visual cortex. Polar angle map (see legend) derived from Benson's atlas (Benson et al., 2014) is shown on a spherical cortical surface for each subject and hemisphere in the TDM dataset. Light gray and dark gray indicate gyri and sulci, respectively. The location of V1 (obtained from the atlas) is outlined in black. The estimated location of the sinuses is outlined in white. In some subjects, the sinuses are positioned near the dorsal and ventral boundaries of V1 (corresponding to the vertical meridian), but in other subjects, the sinuses do not align well with these boundaries.

- Allen EJ, St-Yves G, Wu Y, Breedlove JL, Prince JS, Dowdle LT, Nau M, Caron B, Pestilli F, Charest I, Hutchinson JB, Naselaris T, Kay K (2021) A massive 7T fMRI dataset to bridge cognitive neuroscience and artificial intelligence. *Nature Neuroscience*.
- Arthurs OJ, Williams EJ, Carpenter TA, Pickard JD, Boniface SJ (2000) Linear coupling between functional magnetic resonance imaging and evoked potential amplitude in human somatosensory cortex. *Neuroscience* 101:803-806.
- Attwell D, Iadecola C (2002) The neural basis of functional brain imaging signals. *Trends Neurosci* 25:621-625.
- Benson NC, Winawer J (2018) Bayesian analysis of retinotopic maps. *Elife* 7.
- Benson NC, Butt OH, Brainard DH, Aguirre GK (2014) Correction of distortion in flattened representations of the cortical surface allows prediction of V1-V3 functional organization from anatomy. *PLoS Comput Biol* 10:e1003538.
- Benson NC, Jamison KW, Arcaro MJ, Vu AT, Glasser MF, Coalson TS, Van Essen DC, Yacoub E, Ugurbil K, Winawer J, Kay K (2018) The Human Connectome Project 7 Tesla retinotopy dataset: Description and population receptive field analysis. *J Vis* 18:23.
- Britten KH, Shadlen MN, Newsome WT, Movshon JA (1993) Responses of neurons in macaque MT to stochastic motion signals. *Vis Neurosci* 10:1157-1169.
- Dale AM, Fischl B, Sereno MI (1999) Cortical Surface-Based Analysis: I. Segmentation and Surface Reconstruction. *NeuroImage* 9:179-194.
- Di X, Kannurpatti SS, Rypma B, Biswal BB (2013) Calibrating BOLD fMRI activations with neurovascular and anatomical constraints. *Cereb Cortex* 23:255-263.
- Dumoulin SO (2017) Layers of Neuroscience. *Neuron* 96:1205-1206.
- Dumoulin SO, Wandell BA (2008) Population receptive field estimates in human visual cortex. *Neuroimage* 39:647-660.
- Dumoulin SO, Fracasso A, van der Zwaag W, Siero JCW, Petridou N (2018) Ultra-high field MRI: Advancing systems neuroscience towards mesoscopic human brain function. *Neuroimage* 168:345-357.
- Duvernoy HM, Delon S, Vannson JL (1981) Cortical blood vessels of the human brain. *Brain Res Bull* 7:519-579.
- Fischl B, Dale AM (2000) Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proc Natl Acad Sci U S A* 97:11050-11055.
- Fracasso A, Petridou N, Dumoulin SO (2016a) Systematic variation of population receptive field properties across cortical depth in human visual cortex. *Neuroimage* 139:427-438.
- Fracasso A, Luijten PR, Dumoulin SO, Petridou N (2018) Laminar imaging of positive and negative BOLD in human visual cortex at 7T. *Neuroimage* 164:100-111.
- Fracasso A, van Veluw SJ, Visser F, Luijten PR, Spliet W, Zwanenburg JJM, Dumoulin SO, Petridou N (2016b) Lines of Baillarger in vivo and ex vivo: Myelin contrast across lamina at 7T MRI and histology. *Neuroimage* 133:163-175.
- Gagnon L, Sakadzic S, Lesage F, Pouliot P, Dale AM, Devor A, Buxton RB, Boas DA (2016) Validation and optimization of hypercapnic-calibrated fMRI from oxygen-sensitive two-photon microscopy. *Philos Trans R Soc Lond B Biol Sci* 371.
- Gagnon L, Sakadzic S, Lesage F, Musacchia JJ, Lefebvre J, Fang Q, Yucel MA, Evans KC, Mandeville ET, Cohen-Adad J, Polimeni JR, Yaseen MA, Lo EH, Greve DN, Buxton RB, Dale AM, Devor A, Boas DA (2015) Quantifying the microvascular origin of BOLD-fMRI from first principles with two-photon microscopy and an oxygen-sensitive nanoprobe. *J Neurosci* 35:3663-3675.
- Gorno-Tempini ML, Hutton C, Josephs O, Deichmann R, Price C, Turner R (2002) Echo time dependence of BOLD contrast and susceptibility artifacts. *Neuroimage* 15:136-142.
- Guidi M, Huber L, Lampe L, Merola A, Ihle K, Moller HE (2020) Cortical laminar resting-state signal fluctuations scale with the hypercapnic blood oxygenation level-dependent response. *Hum Brain Mapp* 41:2014-2027.

- Heeger DJ, Ress D (2002) What does fMRI tell us about neuronal activity? *Nature Reviews Neuroscience* 3:142-151.
- Heeger DJ, Huk AC, Geisler WS, Albrecht DG (2000) Spikes versus BOLD: what does neuroimaging tell us about neuronal activity? *Nat Neurosci* 3:631-633.
- Herman MC, Cardoso MMB, Lima B, Sirotin YB, Das A (2017) Simultaneously estimating the task-related and stimulus-evoked components of hemodynamic imaging measurements. *Neurophotonics* 4:031223.
- Hilgetag CC, Barbas H (2005) Developmental mechanics of the primate cerebral cortex. *Anat Embryol (Berl)* 210:411-417.
- Hoogenraad FG, Hofman MB, Pouwels PJ, Reichenbach JR, Rombouts SA, Haacke EM (1999) Sub-millimeter fMRI at 1.5 Tesla: correlation of high resolution with low resolution measurements. *J Magn Reson Imaging* 9:475-482.
- Jamison K, Vizioli L, Zhang R, Tao J, Winawer J, Kay K (2017) A tool for automatic identification of cerebral sinuses and corresponding artifacts in fMRI. *Journal of Vision* 17:295-295.
- Jiang X, Zhang T, Zhang S, Kendrick KM, Liu T (2021) Fundamental functional differences between gyri and sulci: implications for brain function, cognition, and behavior. *Psychoradiology* 1:23-41.
- Kay K, Jamison KW, Zhang RY, Ugurbil K (2020) A temporal decomposition method for identifying venous effects in task-based fMRI. *Nat Methods* 17:1033-1039.
- Kay K, Jamison KW, Vizioli L, Zhang R, Margalit E, Ugurbil K (2019) A critical assessment of data quality and venous effects in sub-millimeter fMRI. *Neuroimage* 189:847-869.
- Kay KN, Winawer J, Mezer A, Wandell BA (2013a) Compressive spatial summation in human visual cortex. *J Neurophysiol* 110:481-494.
- Kay KN, Rokem A, Winawer J, Dougherty RF, Wandell BA (2013b) GLMdenoise: a fast, automated technique for denoising task-based fMRI data. *Front Neurosci* 7:247.
- Kazan SM, Mohammadi S, Callaghan MF, Flandin G, Huber L, Leech R, Kennerley A, Windischberger C, Weiskopf N (2016) Vascular autorescaling of fMRI (VasA fMRI) improves sensitivity of population studies: A pilot study. *Neuroimage* 124:794-805.
- Kim SG, Hendrich K, Hu X, Merkle H, Ugurbil K (1994) Potential pitfalls of functional MRI using conventional gradient-recalled echo techniques. *NMR Biomed* 7:69-74.
- Koopmans PJ, Barth M, Orzada S, Norris DG (2011) Multi-echo fMRI of the cortical laminae in humans at 7 T. *Neuroimage* 56:1276-1285.
- Lee JH, Durand R, Gradinaru V, Zhang F, Goshen I, Kim DS, Fenno LE, Ramakrishnan C, Deisseroth K (2010) Global and local fMRI signals driven by neurons defined optogenetically by type and wiring. *Nature* 465:788-792.
- Logothetis NK (2002) The neural basis of the blood-oxygen-level-dependent functional magnetic resonance imaging signal. *Philos Trans R Soc Lond B Biol Sci* 357:1003-1037.
- Logothetis NK (2008) What we can do and what we cannot do with fMRI. *Nature* 453:869-878.
- Logothetis NK, Wandell BA (2004) Interpreting the BOLD signal. *Annu Rev Physiol* 66:735-769.
- Logothetis NK, Pauls J, Augath M, Trinath T, Oeltermann A (2001) Neurophysiological investigation of the basis of the fMRI signal. *Nature* 412:150-157.
- Menon RS, Ogawa S, Tank DW, Ugurbil K (1993) Tesla gradient recalled echo characteristics of photic stimulation-induced signal changes in the human primary visual cortex. *Magn Reson Med* 30:380-386.
- Mezer A, Yeatman JD, Stikov N, Kay KN, Cho N-J, Dougherty RF, Perry ML, Parvizi J, Hua LH, Butts-Pauly K, Wandell BA (2013) Quantifying the local tissue volume and composition in individual brains with magnetic resonance imaging. *Nat Med* 19:1667-1672.

- Mishra A, Hall CN, Howarth C, Freeman RD (2021) Key relationships between non-invasive functional neuroimaging and the underlying neuronal activity. *Philosophical Transactions of the Royal Society B: Biological Sciences* 376:20190622.
- Polimeni JR, Fischl B, Greve DN, Wald LL (2010) Laminar analysis of 7T BOLD using an imposed spatial activation pattern in human V1. *Neuroimage* 52:1334-1346.
- Rees G, Friston K, Koch C (2000) A direct quantitative relationship between the functional properties of human and macaque V5. *Nature Neuroscience* 3:716-723.
- Self MW, van Kerkoerle T, Goebel R, Roelfsema PR (2019) Benchmarking laminar fMRI: Neuronal spiking and synaptic activity during top-down and bottom-up processing in the different layers of cortex. *Neuroimage* 197:806-817.
- Sethian JA (1996) A fast marching level set method for monotonically advancing fronts. *Proceedings of the National Academy of Sciences* 93:1591.
- Siero JC, Hermes D, Hoogduin H, Luijten PR, Ramsey NF, Petridou N (2014) BOLD matches neuronal activity at the mm scale: a combined 7T fMRI and ECoG study in human sensorimotor cortex. *Neuroimage* 101:177-184.
- Sirotnin YB, Das A (2009) Anticipatory haemodynamic signals in sensory cortex not predicted by local neuronal activity. *Nature* 457:475-479.
- Srirangarajan T, Mortazavi L, Bortolini T, Moll J, Knutson B (2021) Multi-band FMRI compromises detection of mesolimbic reward responses. *NeuroImage* 244:118617.
- Turner R (2002) How much cortex can a vein drain? Downstream dilution of activation-related cerebral blood oxygenation changes. *Neuroimage* 16:1062-1067.
- van der Zwaag W, Francis S, Head K, Peters A, Gowland P, Morris P, Bowtell R (2009) fMRI at 1.5, 3 and 7 T: characterising BOLD signal changes. *Neuroimage* 47:1425-1434.
- van Dijk JA, Fracasso A, Petridou N, Dumoulin SO (2020) Linear systems analysis for laminar fMRI: Evaluating BOLD amplitude scaling for luminance contrast manipulations. *Sci Rep* 10:5462.
- Van Essen DC et al. (2012) The Human Connectome Project: a data acquisition perspective. *Neuroimage* 62:2222-2231.
- Welker W (1990) Why Does Cerebral Cortex Fissure and Fold? In: *Cerebral Cortex: Comparative Structure and Evolution of Cerebral Cortex, Part II* (Jones EG, Peters A, eds), pp 3-136. Boston, MA: Springer US.
- Winawer J, Horiguchi H, Sayres RA, Amano K, Wandell BA (2010) Mapping hV4 and ventral occipital cortex: the venous eclipse. *J Vis* 10:1.
- Yu X, Qian C, Chen DY, Dodd SJ, Koretsky AP (2014) Deciphering laminar-specific neural inputs with line-scanning fMRI. *Nat Methods* 11:55-58.
- Zimmermann J, Goebel R, De Martino F, van de Moortele PF, Feinberg D, Adriany G, Chaimow D, Shmuel A, Ugurbil K, Yacoub E (2011) Mapping the organization of axis of motion selective features in human area MT using high-field fMRI. *PLoS One* 6:e28716.