Effects of land use and weather on the presence and abundance of mosquito-borne disease vectors in a urban and agricultural landscape in Eastern Ontario, Canada

Miarisoa Rindra Rakotoarinia¹, F. Guillaume Blanchet^{2,3,4}, Dominique Gravel², David Lapen⁵, Patrick A. Leighton¹, Nicholas H. Ogden^{1,6}, Antoinette Ludwig^{1,6}

¹ Groupe de recherche en épidémiologie des zoonoses et santé publique (GREZOSP), Faculté de médecine vétérinaire, Université de Montréal, Montréal, Canada

² Département de biologie, Université de Sherbrooke, Sherbrooke, Canada

³ Département de mathématique, Université de Sherbrooke, Sherbrooke, Canada

⁴ Département des sciences de la santé communautaire, Université de Sherbrooke, Sherbrooke, Canada

⁵ Eastern Cereal and Oilseed Research Center, Agriculture and Agrifood Canada, Ottawa, Canada

⁶ Public Health Sciences Division, National Microbiology Laboratory, Public Health Agency of Canada, St-Hyacinthe, Canada

*Corresponding author

E-mail : mr.rakotoarinia.randriamialy@umontreal.ca

Abstract

Weather and land use can significantly impact mosquito abundance and presence, and by consequence, mosquito-borne disease (MBD) dynamics. Knowledge of vector ecology and mosquito species response to these drivers will help us better predict risk from MBD. In this study, we evaluated and compared the independent and combined effects of weather and land use on mosquito species occurrence and abundance in Eastern Ontario, Canada. Data on occurrence and abundance (245,591 individuals) of 30 mosquito species were obtained from mosquito capture at 85 field sites in 2017 and 2018. Environmental variables were extracted from weather and land use datasets in a 1-km buffer around trapping sites. The relative importance of weather and land use on mosquito abundance (for common species) or occurrence (for all species) was evaluated using multivariate hierarchical statistical models. Models incorporating both weather and land use performed better than models that include weather only for approximately half of species (59% for occurrence model and 50% for abundance model). Mosquito occurrence was mainly associated with temperature whereas abundance was associated with precipitation and temperature combined. Land use was more often associated with abundance than occurrence. For most species, occurrence and abundance were positively associated with forest cover but for some there was a negative association. Occurrence and abundance of some species (47% for occurrence model and 88% for abundance model) were positively associated with wetlands, but negatively associated with urban (*Culiseta melanura* and *Anopheles walkeri*) and agriculture (*An. quadrimaculatus*, *Cs. minnesotae* and *An. walkeri*) environments. This study provides predictive relationships between weather, land use and mosquito occurrence and abundance for a wide range of species including those that are currently uncommon, yet known as arboviruses vectors.

Elucidation of these relationships has the potential to contribute to better prediction of MBD risk, and thus more efficiently targeted prevention and control measures.

Keywords: land use, weather, mosquito-borne disease, mosquito vectors, occurrence, abundance, Ontario

Introduction

Mosquitos have been considered as the most dangerous creatures worldwide [1]. With their widespread occurrence and ability to transmit diseases, they are responsible for hundreds of thousands of deaths each year. Over the past 40 years, in Canada, mosquito populations have evolved in terms of composition, abundance and geographic distribution. There were 74 mosquito species identified as occurring in Canada in an exhaustive list compiled in the 1980s [2]. Recently, six new endemic species have been added to the list: *Ochlerotatus ventrovittis*; *Oc. japonicus*; *Culex salinarius*; *Cx. erraticus*; *Anopheles perplexens*, and *An. crucians* [3, 4]. Furthermore, new exotic invasive species from the United States have been repeatedly detected in southern Ontario. *Aedes aegypti* were detected in 2016 and 2017 in southern Ontario. These are thought to be repeated introductions, with no evidence of over-winter survival, however the more cold-tolerant *Ae. albopictus* is now thought to have become established [5].

Arboviruses endemic to Canada include West Nile Virus (WNV), Eastern Equine Encephalitis virus (EEEV) and Californian serogroup viruses (CSV) belonging to the Bunyaviridae, including Jamestown Canyon virus (JCV) and Snowshoe Hare virus (SSHV).

Three mosquito species are mainly responsible for the transmission of WNV in eastern Canada: *Aedes vexans* and the two species of the *Culex pipiens* and *Cx. restuans*, which are often enumerated together due to difficulty in distinguishing them [6]. *Culiseta melanura* is the main vector for EEEV [7, 8]. CSV have several potential vectors including *Ochlerotatus* and *Culiseta* spp. [9-21]. Other mosquito species present in Canada may also be involved in the transmission of arboviruses as suggested by laboratory studies [22-24]. Therefore, monitoring communities of mosquito species, rather than single species, may be more effective for understanding the ecology of arboviruses and predicting risk from them.

Many studies have highlighted the importance of weather and climate as determinants of the spatio-temporal distribution and abundance of mosquitoes [25-28]. Climate change [29], may be driving current expansions in the geographical range of mosquitoes, particularly northern expansion of ranges [30-32]. Weather and climate impact the occurrence and abundance of mosquitos in multiple ways [33]. Increasing temperature can lead to faster inter-stadial development and shorter life cycles. However, the association between weather and mosquito occurrence as well as abundance is not straightforward. High temperatures above 35 °C kills most mosquitoes species [34, 35] but higher average temperatures could also increase the winter survival rate of eggs and promote earlier hatching, and prolong the season of mosquito activity [36]. Increased rainfall may increase reproduction and abundance by increasing suitable larval habitats. Conversely, excessive rainfall can also decrease abundance via egg destruction and leaching of larvae [37].

Weather and climate are not the sole factors influencing species distributions, abundance and diversity. Land use change related to agriculture and urban expansion, and increased forest loss is the largest driver of landscape modification worldwide [38, 39]. Several studies have

linked land cover and land use change to mosquito community changes [40-49]. Urban greening, as a result of urban conservation initiatives, will likely enhance resources for arthropods including mosquitoes [50, 51]. Increased crop irrigation in agricultural environments and associated drainage networks may increase larval habitat for several mosquito species. Habitat structure determines mosquito breeding and resting sites, and nutrition of the different mosquito developmental stages [49, 50, 52-54]. Thus, attempting to understand how land use activities, as well as climate and weather, influence composition and abundance of mosquitos is essential to understand the ecology of mosquito species.

Most studies on mosquitos [25, 55-58] or MBDs [59-63] focus either on weather and/or climate and only occasionally account for land use [40, 64, 65]. Despite evidence that integrated study of weather and/or climate and land use effects can help us better understand mosquito occurrence, abundance and range expansion [66, 67], few studies have done so [68-71]. Furthermore, studies tend to focus on one or a few species and ignore the rarer ones even though the community of vectors as a whole may be important in MBD dynamics [72]. In this study, we aim first to assess the combined impacts of land use and weather variables as determinants of mosquito occurrence and abundance, and to develop a more community-wide approach to assess the risk represented by mosquito species endemic to regions where MBDs have an important impact on public health. Individual differences in species traits were included in this study as they can influence the species response to the explanatory variables. As a first cut analysis, we used readily available remote sensing and meteorological data. Much finer analyzes may follow using micro-climatic and micro-land use data that should improve models accuracy.

Materials and methods

Study region

The study sites are located in the greater Ottawa area and the adjacent South Nation river catchment (SN) in eastern Ontario, Canada (Fig 1). The city of Ottawa covers 2,778.13 km² with ~1 million inhabitants, while the SN river basin covers an area of ~3900 km² with an estimated population of approximately 115,000. The SN study area is predominately agricultural land, with patchy forested areas and some suburban locations [73]. The climate is semi-continental with a warm, humid summer and a very cold winter [74].

Fig 1. Study region and sampling site locations (Public Health Ontario in Ottawa (29 sites) and the South Nation River watershed (56 sub-sites)).

Mosquito variables, sampling design and mosquito identification

Mosquito data came from two sources. Prospective field study was undertaken at 56 SN field sites in 2017 and 2018. Data were also obtained from the ongoing West Nile virus surveillance program of the Ottawa Health Unit during 2017 and 2018, in which mosquitoes were captured at 29 sites. At the SN sites, a stratified sampling design was used. Traps were set in locations with (1) suburban and, (2) agricultural land uses class, that each contained seven sites. Each site regrouped four sub-sites according to the presence of water and woodlot features: (1) an area with no water and no wooded area, (2) an area with water but no woods, (3) an area

with water that is partially wooded and (4) an area with water that is densely wooded. The number of traps employed by a public health unit in Ontario at a given location varied significantly from year to year. To ensure comparability between the two datasets, we kept only the traps that were in operation during a minimum of 15 weeks. For both datasets, mosquitoes were collected on each site one night every other week from May to end of September (end of August for 2017 in the SN sites), using CO₂ baited Centers for Disease Control and Prevention (CDC) miniature light-traps. Trapped mosquitoes were identified and counted by species. *Cx. pipiens* and *Cx. restuans* were lumped into a single group *Cx. pipiens-restuans* due to their morphological similarity

Weather variables

Temperature (minimum and maximum) and precipitation during the study period were extracted from a gridded weather data : Daymet (Daily Surface Weather And Climatological Summaries) (<https://daymet.ornl.gov/>) [75]. These data are available at a daily time step across a 1km x 1km spatial resolution. Weather data were associated with mosquito trapping locations by averaging weather conditions in a one-kilometer radius around each trapping site. Given that larval development to adult stage can occur between 5 to 90 days [76, 77], these daily variables were averaged over different time period: 5, 30 and 90 days before the capture. With the weather during the day of capture, a total of 12 weather variables were used as explanatory in the models (S3 Table).

Land use variables

A buffer zone with a radius of 1 km was created around each trapping site, to approximate the average flight range of the 27 studied species [78, 79]. The percent coverage of land use classes was extracted from annual crop inventory data for 2017 and 2018 [78], within each buffer. Crop inventory data were reclassified into seven land use variables (S4 Table).

Mosquito traits

In this study, we also studied how traits influenced the species response to the environment [80]. The traits, included as a binary variable, were five important behavioral characteristics related to mosquito life cycle and mosquito habitat: 1) number of generations per year, 2) stage of the life cycle that overwinters, 3) type of substrate where the eggs are laid, 4) resistance of the egg to desiccation and 5) typical larval habitat (S6 Table).

Statistical methods

Where possible we used either presence-absence or abundance of each species detected as response variables. However, for species with low abundance (<10 individuals), only presence-absence models were used. The temporal unit was a period of two weeks (S1 Table), species abundance was averaged biweekly for each trapping site (S7 Fig). The temporal unit as well as the sample identifier were used as random effects to account for repeated sampling from the same site.

Spatial autocorrelation was taken into account using the distance-based Moran's eigenvector maps (db-MEM) [81]. This method is a spectral decomposition of the spatial

coordinates of the sampling locations and allows to account for the proximity of certain sampling locations. The db-MEMs that significantly characterize spatial autocorrelation were included as covariates in the model. Construction and test of db-MEM were performed using the adespatial R package [81].

Relationships between mosquito occurrence or abundance and environmental variables were explored using a Hierarchical Modeling of species Community (HMSC) analysis [80]. Mosquito abundance data were explored using over-dispersed Poisson models, whereas for mosquito presence/absence data probit models were used. HMSC model parameters are estimated using a Gibbs sampler. For the probit model, the MCMC chain was run for 300,000 and 320,000 iterations, for weather-only and weather-and-land-use models respectively, half of which were burn-in iterations. For the over-dispersed Poisson model, weather-only and weather-and-land-use models were run for 200,000 and 220,000 iterations respectively, half of which were burn-in iterations. Convergence was assessed using the Geweke's convergence diagnostics [82]. Convergence was assumed to be reached for a parameter if it had a Geweke diagnostic value lower than two. Trace and density plots for all parameters were also checked visually to ensure convergence was reached.

Adjusted Efron's pseudo- R^2 was computed to compare the occurrence and abundance models. This metric is a relative measure of model fit, obtained by calculating the proportion of explained variance over the total variance in the data [83] adjusted for the number of parameters in the model. We also assessed the predictive performance of occurrence-based models by using the area under the receiver operating characteristic curve (AUC) and the root-mean-squared error (RMSE). AUC evaluates the ability of the model to discriminate accurately occurrence and absence. We used the RMSE and the correlation between predicted and observed abundance for

the abundance models. All multivariate models were carried out with the “HMSC” package [84] through the R statistical language version 4.0.2 [85].

Models with and without land use were compared by looking at the values of their respective adjusted pseudo- R^2 . The explained variation was partitioned into components related to fixed effects (weather, land use, spatial component) and random effects (two weeks’ period and sample identification) for each species, following the procedure proposed by Ovaskainen et al. (2017) [80].

Results

Mosquito sampling

Thirty species belonging to seven genera (*Aedes* (*Ae*), *Anopheles* (*An*), *Coquillettidia* (*Cq*), *Culiseta* (*Cs*), *Culex* (*Cx*), *Ochlerotatus* (*Oc*), *Psorophora* (*Ps*)) were detected with a total of 245,591 adult female mosquitos captured. The biweekly average abundance of each species varied over the two years of collection (Fig 2). The total abundance of all combined species was higher in 2017 (163,211) than in 2018 (81,628).

Fig 2. Seasonal changes in biweekly averaged abundance for all species over 2017 and 2018 collection period in Ottawa and the South Nation river sites. The “other species” are rare species with average abundance less than 1. They include *An. barberi*, *An. walkeri*, *Cs. melanura*, *Cs. minnesotae*, *Cs. morsitatus*, *Cx. salinarius*, *Cx. territans*, *Oc. abserratus*, *Oc.*

cantator, *Oc. communis*, *Oc. dorsalis*, *Oc. excrucians*, *Oc. fitchii*, *Oc. intrudens*, *Oc. japonicus*,
Oc. provocans, *Oc. punctator*, *Oc. sticticus*, *Oc. triseriatus*, *Ps. ciliata* and *Ps. ferox*.

Added value of land use as a predictor of mosquito occurrence and abundance

Of the thirty species observed, only 27 were included in statistical analysis due to lack of information on the traits of three species (*An. barberi*, *Oc. sticticus* and *Cx. salinarius*). All models for all species converged. Density plots are provided in the supporting information (S17-20 Fig). The community-level explained variation, illustrated by the mean-values of the species-specific adjusted Efron's pseudo- R^2 , was computed for each model ranging from 0.21 to 0.28 (Table 1). The values of the community-level adjusted Efron's pseudo- R^2 suggest that including land-use in the model in addition to weather can modestly improve overall predictive power. However, because the community-level explained variation is the average R^2 across all species, it hides the significant variation in independent species (from 0.053 to 0.988 for the occurrence model and from 0.040 to 0.725 for the abundance) (Fig 3). For some species, the adjusted Efron's pseudo- R^2 was negative, indicating that these models are less likely than sampling a random variable from a probability distribution, so results from these species are not interpreted.

Table 1. Community-level explained variance of the four models

	Community-level adjusted pseudo- R^2 of weather-only model	Community-level adjusted pseudo- R^2 of weather-and-land-use model
Occurrence (27 species)	0.26	0.28
Abundance (12 species)	0.21	0.22

When variation was partitioned according to each species, there was an increase of the adjusted Efron's pseudo- R^2 values in 59% of species from the occurrence model and 50% of species from the abundance model (Fig 3 and 4). This suggests that adding the land use component does not necessarily improve the quality of the models and this is true for both occurrence and abundance. The community-level adjusted pseudo- R^2 corresponding to land use was low (0.013 and 0.015 for occurrence and abundance, respectively) compared to that of the weather variable (0.069 and 0.057 for occurrence and abundance respectively). In all cases, weather information was most crucial for predicting mosquito occurrence and abundance data in the region.

Fig 3. Variation partitioning according to species among weather, land use, spatial fixed effect and the temporal and spatial unit random effects from the occurrence-model.

Community-level adjusted Efron's pseudo- R^2 of weather-only model (0.24) was compared to that of weather-and-land-use model (0.28). Each bar corresponds to one species.

Fig 4. Variation partitioning according to species among weather, land use, spatial fixed effect and the temporal and spatial unit random effects from the abundance-model.

Community-level adjusted Efron's pseudo- R^2 of weather-only model (0.21) was compared to that of weather-and-land-use model (0.22). Each bar corresponds to one species.

The results also highlight that other components of the models may play important roles in explaining the response variables. For example, approximately half of the whole variation explained was attributed to the sample random effect (10%) suggesting that the sample have relatively strong effects on patterns of species occurrence and abundance while the spatial fixed

effect variance ranged between 3.66 and 8.37%. However, the community-level adjusted Efron's pseudo- R^2 corresponding to the temporal unit ("Biweek") remained very small for all species (<1%). The comparison between the occurrence-based model and the abundance-based model revealed that modeling occurrence explains more variation in the data (Community-level adjusted Efron's pseudo- $R^2 = 0.28$) than the models for abundance (Community-level adjusted Efron's pseudo- $R^2 = 0.22$) except for two species (*Oc. trivittatus* and *An. walkeri*). Finally, the proportion of variance related to each group of covariates and random variables was relatively similar for the same species whether in the occurrence model or in the abundance model.

Weather predictors of mosquito occurrence and abundance

As our primary goal was to evaluate the added value of including land use when modeling mosquito occurrence and abundance, we only present the results obtained from weather-and-land-use models. However, results of the weather-only models are presented in the supporting information (S12 and 13 Fig). Furthermore, since the community-level adjusted Efron's pseudo- R^2 does not reflect precisely the variation in the coefficient of determination specific to each species, from here, we focus only on species-level results.

Model results indicated that various weather variables influence occurrence and abundance of a mosquito species (Table 2, S8 and 9 Fig) with weather variables averaged over one to three months prior to mosquito capture being those most important.

Table 2. Comparison of the significant and relatively strong effects of weather on mosquito occurrence and abundance of the 12 commonest species. + and - signs represent a positive

(average of the marginal distribution of the model parameter above zero) or a negative
(average of the marginal distribution of the model parameter below zero) association
between the variables and mosquito occurrence or abundance. Orange cells show
relationships between weather variables and the response variables (occurrence /
abundance) that are most precise (narrow distribution) and important. Grey cells
represent species that showed either a negative value of the adjusted Efron's pseudo-R² or
a deterioration of model goodness of fit once land use was added.

	Genus	Species	Adjusted pseudo-R ²	Weather adjusted pseudo-R ²	Weather day of capture			Weather averaged 5d before capture			Weather averaged 30d before capture			Weather averaged 90d before capture		
					Precipitation	Min. temperature	Max. temperature	Precipitation	Min. temperature	Max. temperature	Precipitation	Min. temperature	Max. temperature	Precipitation	Min. temperature	Max. temperature
OCCURRENCE MODEL	<i>Aedes</i>	<i>Ae. cinereus</i>	-0.02	0.00					-	+	-		-	+		
		<i>Ae. vexans</i>	0.68	0.27					-	+	-	+	-	-	+	-
	<i>Anopheles</i>	<i>An. punctipennis</i>	0.99	0.10		-	+	+				-			+	+
		<i>An. quadrimaculatus</i>	0.41	0.09		-		+					+	+		
		<i>An. walkeri</i>	0.16	0.03						+	+					
	<i>Coquillettidia</i>	<i>Cq. perturbans</i>	0.31	0.09							+	-	+		+	-
	<i>Culiseta</i>	<i>Cs. melanura</i>	0.15	0.05		+	-	+	-	-	+	-	+		-	
		<i>Cs. minnesotae</i>	0.23	0.05	-	-		+				-	+	+		-
		<i>Cs. morsitans</i>	0.07	0.02					-					-	+	-
	<i>Culex</i>	<i>Cx. pipiens-restuans</i>	0.55	0.11												
		<i>Cx. territans</i>	-0.02	0.00					-	-			+			
	<i>Ochlerotatus</i>	<i>Oc. abserratus</i>	0.20	0.05	-		-	+				-	+	+	-	-
		<i>Oc. canadensis</i>	0.25	0.04					-			-		-	+	-
		<i>Oc. cantator</i>	0.14	0.02		+	-					+		+	-	
		<i>Oc. communis</i>	0.05	0.01		+	-			-	+	-	+		+	-
		<i>Oc. dorsalis</i>	0.07	0.01			-			-	+		+	+		
		<i>Oc. excrucians</i>	0.26	0.03				-					+	+	+	-

		<i>Oc. fitchii</i>	0.09	0.02	-	+			-	-	+			-	+	-
		<i>Oc. intrudens</i>	0.12	0.02		+	-			-		-	+		+	-
		<i>Oc. japonicus</i>	0.61	0.04	-	+	-				-	+	-	+		+
		<i>Oc. provocans</i>	0.35	0.21				+	-	-	-	-	+		+	-
		<i>Oc. triseriatus</i>	0.42	0.05						+	-	+	-			
		<i>Oc. trivittatus</i>	0.41	0.15	+					+			-	-	+	-
		<i>Oc. stimulans</i>	0.25	0.03								-	+		+	-
		<i>Oc. punctor</i>	0.63	0.35		-	+	+		-			+		-	-
	<i>Psorophora</i>	<i>Ps. ciliata</i>	-0.02	0.00	-	+						+		+		
		<i>Ps. ferox</i>	0.19	0.04		+	-	+	-	-	-					

ABUNDANCE MODEL	<i>Aedes</i>	<i>Ae. cinereus</i>	0.07	0.02	-				-	+	-	-		+		+
		<i>Ae. vexans</i>	-0.06	-0.02				-	-	+	-	-	-	+	-	+
	<i>Anopheles</i>	<i>An. punctipennis</i>	0.21	0.01					-						+	
		<i>An. quadrimaculatus</i>	0.04	0.03				+	-	+	-		+	+		+
		<i>An. walkeri</i>	0.19	0.07					-	+	+		+	+	-	+
	<i>Coquillettia</i>	<i>Cq. perturbans</i>	0.72	0.30	-					+		-	+		-	
	<i>Culex</i>	<i>Cx. pipiens-restuans</i>	0.11	0.02		-	+		-	+		-	+	+	+	
	<i>Ochlerotatus</i>	<i>Oc. canadensis</i>	0.51	0.07					-	+	-	-	-		+	
		<i>Oc. provocans</i>	0.33	0.06				+	-		-	-	+	+	-	-
		<i>Oc. trivittatus</i>	0.18	0.03				-		+	-	+	-	+	+	
		<i>Oc. japonicus</i>	0.15	0.04		+	-								+	
		<i>Oc. stimulans</i>	0.24	0.06		+							-	+	+	-

Precipitation - Occurrence and abundance displayed generally positive responses to precipitation. Precipitation had a greater effect when it was averaged over a large number of days prior to sampling relative to a smaller number of days, especially for the abundance models. Total precipitation during the day of capture did not have a major effect. The occurrence of only *Oc. provocans*, and the abundance of only *An. quadrimaculatus* were positively associated with precipitation averaged over 5 days before capture (Prp₅). Precipitation averaged over 30 days had a positive relationship with *An. walkeri* and a negative relationship with *Oc. canadensis*. Precipitation averaged over 90 days was positively associated with the abundance of *An. quadrimaculatus*, *An. walkeri*, *Cx. pipiens-restuans* and *Oc. stimulans*, and the occurrence of *Cs.*

minnesotae and *Oc. excrucians*. The occurrence of *Oc. fitchii* was negatively associated with the precipitation averaged over 90 days.

Minimum temperature – The effect of the minimum temperature differed according to the time lag before capture. The minimum temperature averaged over 90 days before capture (T_{min90}) was a significant determinant for occurrence in 42% of species. The minimum temperature averaged over 30 days (T_{min30}) was positively associated with the occurrence of *Ae. vexans* and *Oc. cantator* and negatively associated with the occurrence of *Cq. perturbans* and *Oc. stimulans*. In the abundance model, the effect of T_{min30} was negative for 38% of species. The minimum temperature averaged over 5 days (T_{min5}) was a determinant of abundance for only two species (*An. quadrimaculatus* and *Oc. canadensis*) but with a negative effect. Finally, the minimum temperature of the day of capture was positively associated with the occurrence of *Cs. melanura* and *Oc. japonicus*.

Maximum temperature - The maximum temperature was significantly associated with mosquito occurrence and abundance. In the occurrence model, the maximum temperature averaged over 30 days (T_{max30}) was positively associated with nine of 19 species, but negatively associated with *Ae. Vexans*. In addition, T_{max30} was positively associated with the abundance of *Cq. perturbans* and *Oc. stimulans* and negatively associated with the abundance of *Oc. trivittatus*. In the abundance model, the maximum temperature averaged over 5 days (T_{max5}) stood out as the variable most frequently associated with mosquito abundance with a positive association with five of eight species. The maximum temperature of the day of capture was negatively associated with the occurrence of *Cs. melanura*.

Land use predictors of mosquito occurrence and abundance

Land use variables had generally weaker associations with mosquito occurrence and abundance than weather variables as shown by the low adjusted Efron's pseudo- R^2 values for the land use component. We nevertheless decided to retain the results of the species for which an improvement in the model fit was observed.

Overall, the land use effects were consistent between occurrence and abundance models. The major land use drivers common to all mosquito species were wetland, forest and bare ground (Table 3). There was a positive relationship between wetland coverage and mosquito occurrence (for 9 of 19 species) and abundance (for 7 of 8 species). Coverage of the forest land use class had a positive relationship with mosquito occurrence (for 7 of 19 species) and abundance (for 3 of 8 species) with the exception of *Cx pipiens-restuans* in the occurrence and abundance model and *An. quadrimaculatus*, *An. walkeri* in the abundance model. Coverage with bare ground was negatively associated with most species (12 of 19 species from the occurrence model and 4 of 8 species for the abundance model).

Other land use classes (water, agriculture, urban and shrubland) were infrequently associated with mosquito occurrence and abundance. Coverage with water was positively associated with occurrence and abundance of *Anopheles* species, and negatively associated with occurrence of *Oc. canadensis*. There was a negative relationship between coverage with the agriculture land class and occurrence of *An. quadrimaculatus* and *Cs. minnesotae*, and the abundance of *An. quadrimaculatus* and *An. walkeri*. Coverage with the urban land class had a negative relationship with the occurrence of *Cs. melanura* and the abundance of *An. walkeri*. The shrubland land class was negatively associated with the occurrence of four species (21%) and the

abundance of three species (38%) but was positively associated with abundance of *An. quadrimaculatus*.

Table 3. Comparison of the significant and relatively strong effects of land use on mosquito occurrence and abundance of the 12 most common species. + and - signs represent a positive (average of the marginal distribution of the model parameter above zero) or a negative (average of the marginal distribution of the model parameter below zero) association between the variables and the mosquito occurrence or abundance. Orange cells show the relationships between the weather variables and the response variables (occurrence / abundance) that are more precise and important. Grey cells represent species that showed either a negative value of the adjusted Efron's pseudo-R² or a deterioration of model goodness of fit once land use was added.

	Genus	Species	Adjusted pseudo-R ²	Land use adjusted pseudo-R ²	Land use						
					Water	Wetland	Agriculture	Urban	Forest	Shrubland	Bare
OCCURRENCE MODEL	<i>Aedes</i>	<i>Ae. cinereus</i>	-0.02	-0.0017		+	-		+	+	-
		<i>Ae. vexans</i>	0.68	0.0143	-	+	+			+	-
	<i>Anopheles</i>	<i>An. punctipennis</i>	0.99	0.0118	+	+	-		-		+
		<i>An. quadrimaculatus</i>	0.41	0.0521	+		-	-	-	+	+
		<i>An. walkeri</i>	0.16	0.0383	+	+	-	-	-	+	-
	<i>Coquillettidia</i>	<i>Cq. perturbans</i>	0.31	0.0175		+	-		+	-	-
	<i>Culiseta</i>	<i>Cs. melanura</i>	0.15	0.0131	+	+	-	-		-	-
		<i>Cs. minnesotae</i>	0.23	0.0153	-	-	-	-	+	-	
		<i>Cs. morsitans</i>	0.07	0.0076	-	+			+	-	-
	<i>Culex</i>	<i>Cx. pipiens-restuans</i>	0.55	0.0464		+	-	+	-		+
		<i>Cx. territans</i>	-0.02	-0.0008			-	-	-		+
	<i>Ochlerotatus</i>	<i>Oc. abserratus</i>	0.20	0.0097	-		-	-	+	-	-
		<i>Oc. canadensis</i>	0.25	0.0379	-	+		+	+	-	-
		<i>Oc. cantator</i>	0.14	0.0053		+	-	-			-
		<i>Oc. communis</i>	0.05	0.0015	-	+		-	+	-	-
		<i>Oc. dorsalis</i>	0.07	0.0017		+	-	-			-
		<i>Oc. excrucians</i>	0.26	0.0059		-			+	-	-
		<i>Oc. fitchii</i>	0.09	0.0051	-	+	-	-	+	-	-

		<i>Oc. intrudens</i>	0.12	0.0027	-				+	-	-
		<i>Oc. japonicus</i>	0.61	0.0145	-	+		-	-		-
		<i>Oc. provocans</i>	0.35	0.0108	-	-	+		+	-	-
		<i>Oc. triseriatus</i>	0.42	0.0073	-	+				-	-
		<i>Oc. trivittatus</i>	0.41	0.0151	-	+	+	-	+	+	-
		<i>Oc. stimulans</i>	0.25	0.0042		+	+	+	+	-	-
		<i>Oc. punctator</i>	0.63	0.0113	-	-	-		+	-	
	<i>Psorophora</i>	<i>Ps. ciliata</i>	-0.02	-0.0007	-	+	-	-	+	-	-
		<i>Ps. ferox</i>	0.19	0.0074	-	+	-	-	+	-	-

ABUNDANCE MODEL	<i>Aedes</i>	<i>Ae. cinereus</i>	0.07	0.0074	-	+	-		+		-
		<i>Ae. vexans</i>	-0.06	-0.0013	-	+		-	-		-
	<i>Anopheles</i>	<i>An. punctipennis</i>	0.21	0.0107		+	-	-	-		+
		<i>An. quadrimaculatus</i>	0.04	0.0010	+		-	-	-	+	+
		<i>An. walkeri</i>	0.19	0.0045	+	+	-	-	-	+	-
	<i>Coquilletidia</i>	<i>Cq. perturbans</i>	0.72	0.0113	-	+	-	-	+	-	
	<i>Culex</i>	<i>Cx. pipiens-restuans</i>	0.11	0.0222		+			-	+	+
	<i>Ochlerotatus</i>	<i>Oc. canadensis</i>	0.51	0.0108	-	+	-	+	+	-	-
		<i>Oc. provocans</i>	0.33	0.0463	-	+	+		+	-	-
		<i>Oc. trivittatus</i>	0.18	0.0462	-	+	+	+	+	-	-
		<i>Oc. japonicus</i>	0.15	0.0083	-	+	-	-	-	-	-
		<i>Oc. stimulans</i>	0.24	0.0105		+	+	+	+	-	-

Discussion

This study confirms the central role of weather in the ecology of mosquito in Eastern Canada, while highlighting the importance of land use in determining the occurrence and abundance of certain species. By jointly analyzing data from multiple species using the HMSC approach, we were able to gain unique insight into ecological drivers of species that are less abundant and therefore infrequently studied, yet are known vectors for several important mosquito-borne diseases.

The importance of cumulative weather effects

Our results clearly demonstrate the importance of the accumulation over several days, or weeks to months of favorable climatic conditions for the development of mosquitoes. Mosquitoes are ectothermic organisms, each developmental stage may be sensitive to weather conditions, and there is a species-specific range of temperature conditions that permit completion of their life cycle and, if we are in the optimum of the range, is the shortest possible [86]. However, for univoltine species, i.e. having only one generation per year, high temperature for one month may speed up the life cycle which leads to an earlier emergence of the adults. As adult lifespan can last few weeks at most [77], the cohorts that appeared after one month are no longer present after 90 days. Indeed, we observed that, together with the positive effect of a higher temperature averaged over 30 days before capture, we often saw a negative association when a longer period is considered, i.e. 90 days.

Weather variables that affect mosquito occurrence are not necessarily the same that affect abundance. The importance of precipitation, especially accumulated over several weeks before mosquito capture, appears only in our abundance models. This cumulative effect of precipitation has to be considered alongside the time period between egg laying and adult emergence. The multiplication of mosquitoes depends closely on the availability of breeding sites favored by rainfall as well as the presence of water in these same sites, whether natural or artificial. If the water evaporates quickly, this will give less time for larvae to develop and limit the number of emerging adults [87].

Lastly, warm temperatures on the day of capture had a limited impact on mosquito abundance, compared to the cumulative effects of temperatures over several days. However, for

some species, optimal temperatures for adult daily activity apparently remained relatively important, and not in the expected direction. Two species, *Oc. japonicus* and *Cs. melanura*, were positively associated with a warm minimal temperature on the day of the capture. The scarce information from the existing literature indicates that *Cs. melanura* is nocturnal [88] and therefore, adults may be more active during the night when it is generally cooler (T_{min_0}) than during the day (T_{max_0}), and making warm nights potentially favorable to *Cs. melanura*. Explanations for *Ae. japonicus* are less obvious. This species is known to be cold tolerant [89], and may be more active during cool days (intermediate temperatures).

An apparent limitation of our study was to use the weather as a proxy of mosquito seasonality. However, mosquitoes don't respond instantaneously to weather but accumulate past effects of this latter to develop cohorts over the season. Using the average weather over previous time interval partly solves this issue but our approach doesn't represent a seasonal model per se. Our model will still predict high abundance of mosquitos in early or late summer where weather conditions are exceptionally favorable, no matter if the population had enough time to develop. Thus, our study highlights the need of a seasonal model for community, currently non-existent, that will be constructed on population dynamics to reflect this cumulative build up of the population over the season.

Land use influence on mosquito

Overall, the adjusted pseudo- R^2 values related to land use were found to be relatively low. In part, a relatively low effect of land use categories could be due to the spatial resolution of land use data being 30m, which may be too coarse to adequately represent variation in mosquito

habitat; a finer land use data resolution (~ 1m) could lead to high adjusted pseudo-R² values [46]. The sample random effect and / or the spatial fixed effects included in the models often showed a relatively high proportion of variance, which could be further evidence that the full impact of land use on mosquito occurrence was not captured in this study. Indeed, our spatial descriptors (db-MEM) likely captured relevant land use structure that were not accounted for in our study such as biogeographic barriers [81].

Wetland appeared to be an important driver of mosquito occurrence or abundance for many species. This is not surprising since wetlands and water constitute a fundamental habitat for the mosquito life cycle, in particular for *An. quadrimaculatus*, *An. walkeri*, *Cq. perturbans*, *Oc. canadensis*, *Oc. cantator*, *Oc. japonicus*, *Oc. trivittatus* and *Ps. ferox* [90, 91], among which are known potential vectors of EEEV and CSV (*Cq. perturbans*, *Oc. canadensis*, *Cs. melanura*).

Occurrence of ‘forest’ habitat was also a strong determinant of occurrence and abundance of several mosquito species. Wooded areas generally serve as refuge for adult mosquitoes, especially for the CSV vectors *Oc. excrucians*, *Oc. communis* and *Oc. stimulans*. These results are consistent with the limited information in the literature which describes, for example, that *Oc. excrucians* can be found in a wide range of habitats according to Wood et al. [90] while Darsie et al [91] mention that these species larvae tend to prefer wooded spring pools and semi-permanent wetlands with abundant vegetation. The ‘forest’ class was negatively associated with *Cx. pipiens-restuans* (presence) and *Anopheles* (abundance), which is consistent with several studies [58, 63, 92]. *Cx. pipiens-restuans* are also known to be urban species although a positive association with ‘urban’ was not detected in our study. Indeed, urban landscapes are very heterogeneous and could themselves be classified into several categories according to the presence of woodlands and aquatic environments. A more detailed characterization of the urban

micro-habitat may help better define the relationships between this land use class (urban) and mosquitoes. The negative association between ‘forest’ and the abundance of *An. walkeri* could be explained by the fact that the adults of this species generally stay near their larval aquatic habitats, in contrast to other species of *Anopheles* (*An. quadrimaculatus* and *An. punctipennis*) which leave their breeding sites during the day to seek shelters [91].

The ‘bare ground’ class was negatively associated with mosquito occurrence and abundance, as could be expected as the lack of an aquatic environment and vegetation would prevent completion of the mosquito life cycle.

Intensive agricultural practices, as are common in the study region, could negatively influence some species, including *Anopheles* species and one *Culiseta* species. *Culiseta minnesotae* is naturally adapted to specific marshy environments including sedge and cattail marshes which may be found at the edge of some agricultural fields [91, 93]. However, cattails are considered as invasive plants and are subject to considerable efforts to eliminate them by most farmers, reducing habitat for this species. *Anopheles* species generally prefer pristine freshwater environments such as peat bogs and unpolluted freshwater swamps [90, 91], while wetlands in agricultural land are often contaminated with chemical agents (fertilizers, pesticides, etc.) [94, 95], which may explain the negative association between these species and the agriculture land use class.

The negative relationship between the shrubland class and some mosquito species (*Cq. perturbans*, *Oc. canadensis*, *Oc. communis*, *Oc. intrudens* and *Oc. stimulans*) may be explained by the lack of wetland in this type of habitat, dominated by shrubs, including grasses, herbs, and geophytes.

In summary, we found evidence that mosquito occurrence and abundance are related to land use variables. We detected some common patterns among mosquito species with regard to their response to natural land use variables (water and forest) but more specific response to other land use variables such as agricultural land and shrubland.

Conclusion

Efforts to connect mosquito distribution and abundance to environmental variables are generally limited to the most regionally relevant species in terms of public health, most being focused on mosquito abundance. However, it is important to determine how multiple species are influenced by these variables since models based on a single or small number of species do not provide a portrait of the whole community including rare disease vectors. This study provides baseline multi-species associations where species respond differently to a given set of environmental conditions. Our study also demonstrates the relevance of including land use in predicting mosquito occurrence and abundance. If temperature and precipitation are primary drivers in mosquito development, appropriate habitat is also required for many species. Our findings suggest that the impact of land use and weather together on mosquito occurrence and abundance is species-specific and may have diverging impacts on whether a species can survive (occurrence) or thrive (abundance). Future research should develop our understanding of mosquito community preferences in terms of habitat and climatic conditions. Several analytical approaches are now available to work at the multi-species level and study the ecological connections between species. One of the main outcome of such study would be to confirm if human-altered environments favor the multiplication of a few species of mosquitoes that are

flexible habitat generalists, to while excluding a diversity of more specialized mosquitoes which are becoming increasingly rare. We expect that this imbalance in terms of biodiversity would favor species that are efficient vectors of human disease. Further information about mosquito community ecology could allow vector management strategies to be more targeted with respect to anthropogenic pressures including deforestation, urbanization and agriculture.

Acknowledgement

We are grateful to all AAFC field teams, Curtis Russel and folks from Public Health Ontario for providing Ontario surveillance data, Julie Légaré for mosquito identification and Christopher Fernandez Prada for lab facilities. We thank the Groupe de Recherche sur l'Épidémiologie des Zoonoses et Santé Publique (GREZOSP) and the Public Health Agency of Canada (PHAC) which funded the study and supported its execution.

Author Contributions

Conceptualization: Miarisoa Rindra Rakotoarinia, Antoinette Ludwig, Patrick Leighton, Nicholas H. Ogden, David Lapen, Guillaume Blanchet, Dominique Gravel.

Formal Analysis: Miarisoa Rindra Rakotoarinia, Guillaume Blanchet, Antoinette Ludwig.

Funding Acquisition: Antoinette Ludwig, David Lapen.

489 **Methodology:** Miarisoa Rindra Rakotoarinia, Antoinette Ludwig, Patrick Leighton, Nicholas H.
490 Ogden, David Lapen, Guillaume Blanchet, Dominique Gravel.

491 **Resources:** Antoinette Ludwig, Patrick Leighton, Nicholas H. Ogden, David Lapen, Guillaume
492 Blanchet, Dominique Gravel.

493 **Software:** Guillaume Blanchet, Dominique Gravel.

494 **Supervision:** Antoinette Ludwig, Patrick Leighton, Nicholas H. Ogden, David Lapen,
495 Guillaume Blanchet, Dominique Gravel.

496 **Visualization:** Miarisoa Rindra Rakotoarinia, Guillaume Blanchet, Antoinette Ludwig.

497 **Writing – Original Draft Preparation:** Miarisoa Rindra Rakotoarinia, Antoinette Ludwig.

498 **Writing – Review & Editing:** Miarisoa Rindra Rakotoarinia, Antoinette Ludwig, Patrick
499 Leighton, Nicholas H. Ogden, David Lapen, Guillaume Blanchet, Dominique Gravel.

500

501

502 **References**

503 1. Organisation Mondiale de la Santé O. Maladies Tropicales Négligées. 2020 [updated
504 2020-09-29]. Available from:

505 https://www.who.int/neglected_diseases/vector_ecology/mosquito-borne-diseases/fr/.

506 2. Wood D, Dang P, Ellis R. The insects and arachnids of Canada. Part 6. The mosquitoes
507 of Canada. Diptera: Culicidae: Canadian Government Publishing Centre.; 1979.

3. Darsie RF, Jr. Revision of Darsie and Ward (1981) to include *Ochlerotatus japonicus* Theobald and a checklist of species referred to the genus *Ochlerotatus* in the Nearctic Region. J Am Mosq Control Assoc. 2002;18(4):237-40. PubMed PMID: 12542180.
4. Giordano BV, Gasparotto A, Hunter FF. A Checklist of the 67 Mosquito Species of Ontario, Canada. Journal of the American Mosquito Control Association. 2015;31(1):101-3. doi: 10.2987/14-6456R.1.
5. Giordano B, Gasparotto A, Liang P, Nelder M, Russell C, Hunter F. Discovery of an *Aedes* (*Stegomyia*) *albopictus* population and first records of *Aedes* (*Stegomyia*) *aegypti* in Canada. Medical and veterinary entomology. 2020;34(1):10-6.
6. Kramer LD, Styer LM, Ebel GD. A global perspective on the epidemiology of West Nile virus. Annu Rev Entomol. 2008;53:61-81.
7. Armstrong PM, Andreadis TG. Eastern equine encephalitis virus in mosquitoes and their role as bridge vectors. Emerging infectious diseases. 2010;16(12):1869.
8. Molaei G, Andreadis TG, Armstrong PM, Anderson JF, Vossbrinck CR. Host feeding patterns of *Culex* mosquitoes and West Nile virus transmission, northeastern United States. Emerging infectious diseases. 2006;12(3):468.
9. Andreadis TG, Anderson JF, Armstrong PM, Main AJ. Isolations of Jamestown Canyon virus (Bunyaviridae: Orthobunyavirus) from field-collected mosquitoes (Diptera: Culicidae) in Connecticut, USA: a ten-year analysis, 1997-2006. Vector Borne Zoonotic Dis. 2008;8(2):175-88. doi: 10.1089/vbz.2007.0169. PubMed PMID: 18386967.
10. LeDuc JW. Review Article: The Ecology Of California Group Viruses. Journal of Medical Entomology. 1979;16(1):1.

- 530 11. Andreadis TG, Capotosto PM, Shope RE, Tirrell SJ. Mosquito and arbovirus surveillance
531 in Connecticut, 1991-1992. J Am Mosq Control Assoc. 1994;10(4):556-64. PubMed PMID:
532 7707064.
- 533 12. Goff G, Whitney H, Drebot MA. Roles of Host Species, Geographic Separation, and
534 Isolation in the Seroprevalence of Jamestown Canyon and Snowshoe Hare Viruses in
535 Newfoundland. Applied and Environmental Microbiology. 2012;78(18):6734-40. doi:
536 10.1128/AEM.01351-12. PubMed PMID: PMC3426688.
- 537 13. Harris MC, Yang F, Jackson DM, Dotseth EJ, Paulson SL, Hawley DM. La Crosse Virus
538 Field Detection and Vector Competence of Culex Mosquitoes. Am J Trop Med Hyg.
539 2015;93(3):461-7. Epub 2015/07/16. doi: 10.4269/ajtmh.14-0128. PubMed PMID: 26175029;
540 PubMed Central PMCID: PMCPMC4559680.
- 541 14. Defoliart GR, Watts DM, Grimstad PR. Changing patterns in mosquito-borne
542 arboviruses. J Am Mosq Control Assoc. 1986;2(4):437-55. Epub 1986/12/01. PubMed PMID:
543 2906983.
- 544 15. DeFoliart GR, Anslow RO, Hanson RP, Morris CD, Papadopoulos O, Sather GE.
545 Isolation of Jamestown Canyon serotype of California encephalitis virus from naturally infected
546 Aedes mosquitoes and tabanids. Am J Trop Med Hyg. 1969;18(3):440-7. PubMed PMID:
547 5768778.
- 548 16. Parkin WE, Hammon WM, Sather GE. Review of current epidemiological literature on
549 viruses of the California arbovirus group. Am J Trop Med Hyg. 1972;21(6):964-78. Epub
550 1972/11/01. PubMed PMID: 4564448.

- 551 17. Heard PB, Zhang MB, Grimstad PR. Isolation of Jamestown Canyon virus (California
552 serogroup) from Aedes mosquitoes in an enzootic focus in Michigan. J Am Mosq Control Assoc.
553 1990;6(3):461-8. PubMed PMID: 2230774.
- 554 18. Mokry J, Artsob H, Butler R. Studies on California serogroup virus activity in
555 Newfoundland, Canada, 1980-83. Mosq News. 1984;44:310-4.
- 556 19. Boromisa RD, Grimstad PR. Virus-vector-host relationships of Aedes stimulans and
557 Jamestown Canyon virus in a northern Indiana enzootic focus. Am J Trop Med Hyg.
558 1986;35(6):1285-95. PubMed PMID: 2878626.
- 559 20. Iversen J, Hanson RP, Papadopoulos O, Morris CV, DeFoliart GR. Isolation of viruses of
560 the California encephalitis virus group from boreal Aedes mosquitoes. Am J Trop Med Hyg.
561 1969;18(5):735-42. PubMed PMID: 5810801.
- 562 21. Boromisa RD, Grayson MA. Oral transmission of Jamestown Canyon virus by Aedes
563 provocans mosquitoes from northeastern New York. J Am Mosq Control Assoc. 1991;7(1):42-7.
564 Epub 1991/03/01. PubMed PMID: 1675258.
- 565 22. Kramer LD, Reisen WK, Chiles RE. Vector competence of Aedes dorsalis (Diptera:
566 Culicidae) from Morro Bay, California, for western equine encephalomyelitis virus. J Med
567 Entomol. 1998;35(6):1020-4. PubMed PMID: 9835696.
- 568 23. Turell MJ, O'Guinn ML, Dohm DJ, Jones JW. Vector competence of North American
569 mosquitoes (Diptera: Culicidae) for West Nile virus. J Med Entomol. 2001a;38(2):130-4. doi:
570 10.1603/0022-2585-38.2.130. PubMed PMID: 11296813.
- 571 24. Vaidyanathan R, Edman JD, Cooper LA, Scott TW. Vector competence of mosquitoes
572 (Diptera:Culicidae) from Massachusetts for a sympatric isolate of eastern equine
573 encephalomyelitis virus. J Med Entomol. 1997;34(3):346-52. PubMed PMID: 9151501.

- 574 25. Hongoh V, Berrang-Ford L, Scott M, Lindsay L. Expanding geographical distribution of
575 the mosquito, *Culex pipiens*, in Canada under climate change. *Applied geography*. 2012;33:53-
576 62.
- 577 26. Ogden NH, Milka R, Caminade C, Gachon P. Recent and projected future climatic
578 suitability of North America for the Asian tiger mosquito *Aedes albopictus*. *Parasites & vectors*.
579 2014;7(1):532.
- 580 27. Ogden N, Gachon P. Changements climatiques et maladies infectieuses: À quoi pouvons-
581 nous nous attendre. *Relevé des maladies transmissibles au Canada*. 2019;45(4):83-8.
- 582 28. Ogden NH, St-Onge L, Barker IK, Brazeau S, Bigras-Poulin M, Charron DF, et al. Risk
583 maps for range expansion of the Lyme disease vector, *Ixodes scapularis*, in Canada now and with
584 climate change. *International Journal of Health Geographics*. 2008;7(1):1-15.
- 585 29. Solon S, Qin M, Manning Z. Contribution of working group I to the fourth assessment
586 report of the intergovernmental panel on climate change, 2007. Cambridge University Press,
587 Cambridge; 2007.
- 588 30. Hertig E. Distribution of *Anopheles* vectors and potential malaria transmission stability in
589 Europe and the Mediterranean area under future climate change. *Parasites & vectors*.
590 2019;12(1):18.
- 591 31. Kraemer MU, Reiner RC, Brady OJ, Messina JP, Gilbert M, Pigott DM, et al. Past and
592 future spread of the arbovirus vectors *Aedes aegypti* and *Aedes albopictus*. *Nature microbiology*.
593 2019;4(5):854-63.
- 594 32. Armstrong PM, Andreadis TG, Shepard JJ, Thomas MC. Northern range expansion of the
595 Asian tiger mosquito (*Aedes albopictus*): Analysis of mosquito data from Connecticut, USA.
596 *PLoS neglected tropical diseases*. 2017;11(5):e0005623.

33. Ogden NH, Lindsay LR. Effects of climate and climate change on vectors and vector-borne diseases: ticks are different. *Trends in parasitology*. 2016;32(8):646-56.
34. Buxton PA. The effect of climatic conditions upon populations of insects. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 1933;26(4):325-56.
35. Heinrich B. *Insect thermoregulation*. 1981.
36. Eisen L, Monaghan AJ, Lozano-Fuentes S, Steinhoff DF, Hayden MH, Bieringer PE. The impact of temperature on the bionomics of *Aedes (Stegomyia) aegypti*, with special reference to the cool geographic range margins. *Journal of medical entomology*. 2014;51(3):496-516.
37. Clements AN. *The biology of mosquitoes: development, nutrition and reproduction*: Chapman & Hall London; 1992.
38. Curtis PG, Slay CM, Harris NL, Tyukavina A, Hansen MC. Classifying drivers of global forest loss. *Science*. 2018;361(6407):1108-11.
39. DeFries RS, Rudel T, Uriarte M, Hansen M. Deforestation driven by urban population growth and agricultural trade in the twenty-first century. *Nature Geoscience*. 2010;3(3):178-81.
40. Rosà R, Marini G, Bolzoni L, Neteler M, Metz M, Delucchi L, et al. Early warning of West Nile virus mosquito vector: climate and land use models successfully explain phenology and abundance of *Culex pipiens* mosquitoes in north-western Italy. *Parasites & vectors*. 2014;7(1):269.
41. Zahouli JB, Koudou BG, Müller P, Malone D, Tano Y, Utzinger J. Effect of land-use changes on the abundance, distribution, and host-seeking behavior of *Aedes* arbovirus vectors in oil palm-dominated landscapes, southeastern Côte d'Ivoire. *PloS one*. 2017;12(12):e0189082.

42. Marí RB, Jiménez-Peydró R. Differences in mosquito (Diptera: Culicidae) biodiversity across varying climates and land-use categories in Eastern Spain. *Entomologica Fennica*. 2011;22(3):190–8—8.
43. de Valdez MRW. Mosquito species distribution across urban, suburban, and semi-rural residences in San Antonio, Texas. *Journal of Vector Ecology*. 2017;42(1):184-8.
44. Steiger DBM, Ritchie SA, Laurance SG. Mosquito communities and disease risk influenced by land use change and seasonality in the Australian tropics. *Parasites & vectors*. 2016;9(1):387.
45. Asigau S, Parker PG. The influence of ecological factors on mosquito abundance and occurrence in Galápagos. *Journal of Vector Ecology*. 2018;43(1):125-37.
46. Landau KI, van Leeuwen WJ. Fine scale spatial urban land cover factors associated with adult mosquito abundance and risk in Tucson, Arizona. *Journal of Vector Ecology*. 2012;37(2):407-18.
47. Steiger DBM, Ritchie SA, Laurance SG. Land use influences mosquito communities and disease risk on remote tropical islands: a case study using a novel sampling technique. *The American journal of tropical medicine and hygiene*. 2016;94(2):314-21.
48. Chaves LF, Hamer GL, Walker ED, Brown WM, Ruiz MO, Kitron UD. Climatic variability and landscape heterogeneity impact urban mosquito diversity and vector abundance and infection. *Ecosphere*. 2011;2(6):1-21.
49. Roiz D, Ruiz S, Soriguer R, Figuerola J. Landscape effects on the presence, abundance and diversity of mosquitoes in Mediterranean wetlands. *PLoS One*. 2015;10(6):e0128112.

50. Yang L, Turo KJ, Riley CB, Inocente EA, Tian J, Hoekstra NC, et al. Can urban greening increase vector abundance in cities? The impact of mowing, local vegetation, and landscape composition on adult mosquito populations. *Urban Ecosystems*. 2019;22(5):827-39.
51. Wilke AB, Chase C, Vasquez C, Carvajal A, Medina J, Petrie WD, et al. Urbanization creates diverse aquatic habitats for immature mosquitoes in urban areas. *Scientific reports*. 2019;9(1):1-11.
52. Stanton M, Rejmanek M, Galen C. Changes in vegetation and soil fertility along a predictable snowmelt gradient in the Mosquito Range, Colorado, USA. *Arctic and Alpine Research*. 1994;26(4):364-74.
53. Thullen JS, Sartoris JJ, Walton WE. Effects of vegetation management in constructed wetland treatment cells on water quality and mosquito production. *Ecological Engineering*. 2002;18(4):441-57.
54. Gardner AM, Anderson TK, Hamer GL, Johnson DE, Varela KE, Walker ED, et al. Terrestrial vegetation and aquatic chemistry influence larval mosquito abundance in catch basins, Chicago, USA. *Parasites & vectors*. 2013;6(1):9.
55. Ripoche M, Campagna C, Ludwig A, Ogden NH, Leighton PA. Short-term Forecasting of Daily Abundance of West Nile Virus Vectors *Culex pipiens-restuans* (Diptera: Culicidae) and *Aedes vexans* Based on Weather Conditions in Southern Québec (Canada). *Journal of medical entomology*. 2019;56(3):859-72.
56. Wang J, Ogden NH, Zhu H. The impact of weather conditions on *Culex pipiens* and *Culex restuans* (Diptera: Culicidae) abundance: a case study in Peel region. *Journal of medical entomology*. 2011;48(2):468-75.

661 57. Lebl K, Brugger K, Rubel F. Predicting Culex pipiens/restuans population dynamics by
662 interval lagged weather data. Parasites & vectors. 2013;6(1):129.

663 58. Diuk-Wasser MA, Brown HE, Andreadis TG, Fish D. Modeling the spatial distribution of
664 mosquito vectors for West Nile virus in Connecticut, USA. Vector-Borne & Zoonotic Diseases.
665 2006;6(3):283-95.

666 59. Chen CC, Jenkins E, Epp T, Waldner C, Curry PS, Soos C. Climate change and West
667 Nile virus in a highly endemic region of North America. Int J Environ Res Public Health.
668 2013;10(7):3052-71. doi: 10.3390/ijerph10073052. PubMed PMID: 23880729; PubMed Central
669 PMCID: PMC3734476.

670 60. Bartlow AW, Manore C, Xu C, Kaufeld KA, Del Valle S, Ziemann A, et al. Forecasting
671 Zoonotic Infectious Disease Response to Climate Change: Mosquito Vectors and a Changing
672 Environment. Veterinary sciences. 2019;6(2):40.

673 61. Giordano BV, Kaur S, Hunter FF. West Nile virus in Ontario, Canada: A twelve-year
674 analysis of human case prevalence, mosquito surveillance, and climate data. PLoS One.
675 2017;12(8):e0183568.

676 62. Trawinski P, Mackay D. Meteorologically conditioned time-series predictions of West
677 Nile virus vector mosquitoes. Vector-Borne and Zoonotic Diseases. 2008;8(4):505-22.

678 63. Trawinski PR, Mackay DS. Identification of environmental covariates of West Nile virus
679 vector mosquito population abundance. Vector-Borne and Zoonotic Diseases. 2010;10(5):515-
680 26.

681 64. Ferraguti M, Martinez-de La Puente J, Figuerola J. Ecological Effects on the Dynamics
682 of West Nile Virus and Avian Plasmodium: The Importance of Mosquito Communities and

683 Landscape. Viruses-Basel. 2021;13(7). doi: 10.3390/v13071208. PubMed PMID:
684 WOS:000676895200001.

685 65. Reisen WK. Landscape epidemiology of vector-borne diseases. Annual review of
686 entomology. 2010;55:461-83.

687 66. Madzokere ET, Hallgren W, Sahin O, Webster JA, Webb CE, Mackey B, et al.
688 Integrating statistical and mechanistic approaches with biotic and environmental variables
689 improves model predictions of the impact of climate and land-use changes on future mosquito-
690 vector abundance, diversity and distributions in Australia. Parasites & Vectors. 2020;13(1):1-13.

691 67. Franklins LH, Jones KE, Redding DW, Abubakar I. The effect of global change on
692 mosquito-borne disease. The Lancet Infectious Diseases. 2019;19(9):e302-e12.

693 68. Chuang T-W, Hildreth MB, Vanroekel DL, Wimberly MC. Weather and land cover
694 influences on mosquito populations in Sioux Falls, South Dakota. Journal of medical
695 entomology. 2011;48(3):669-79.

696 69. Hunt SK, Galatowitsch ML, McIntosh AR. Interactive effects of land use, temperature,
697 and predators determine native and invasive mosquito distributions. Freshwater Biology.
698 2017;62(9):1564-77.

699 70. Kache PA, Eastwood G, Collins-Palmer K, Katz M, Falco RC, Bajwa WI, et al.
700 Environmental Determinants of Aedes albopictus Abundance at a Northern Limit of Its Range in
701 the United States. The American Journal of Tropical Medicine and Hygiene. 2020;102(2):436-
702 47.

703 71. McClure KM, Lawrence C, Kilpatrick AM. Land use and larval habitat increase Aedes
704 albopictus (Diptera: Culicidae) and Culex quinquefasciatus (Diptera: Culicidae) abundance in
705 lowland Hawaii. Journal of medical entomology. 2018;55(6):1509-16.

72. Becker N, Petric D, Zgomba M, Boase C, Dahl C, Lane J, et al. Mosquitoes and their control Kluwer Academic. Plenum Publishers, New York; 2003.
73. Butt S, Ramprasad P, Fenech A. Changes in the landscape of southern Ontario Canada since 1750: impacts of European colonization. Integrated Mapping Assessment. 2005:83-92.
74. Baldwin DJ, Desloges JR, Band LE. Physical geography of Ontario. Ecology of a managed terrestrial landscape: patterns and processes of forest landscapes in Ontario. 2000:12-29.
75. Thornton P, Thornton M, Mayer B, Wei Y, Devarakonda R, Vose R, et al. Daymet: Daily Surface Weather Data on a 1-km Grid for North America, Version 3. ORNL DAAC, USA. 2017.
76. Roiz D, Ruiz S, Soriguer R, Figuerola J. Climatic effects on mosquito abundance in Mediterranean wetlands. Parasites & vectors. 2014;7(1):333.
77. Illinois Department of Public Health. Prevention and control - Mosquitoes and disease 2017 [updated March 29, 2017]. Available from: <https://www.cdc.gov/dengue/resources/factsheets/mosquitolifecyclefinal.pdf>.
78. Lowe AM, Forest-Bérard K, Trudel R, Lo E, Gamache P, Tandonnet M, et al. Mosquitoes Know No Borders: Surveillance of Potential Introduction of Aedes Species in Southern Québec, Canada. Pathogens. 2021;10(8). Epub 2021/08/29. doi: 10.3390/pathogens10080998. PubMed PMID: 34451462; PubMed Central PMCID: PMC8400959.
79. Verdonschot PF, Besse-Lototskaya AA. Flight distance of mosquitoes (Culicidae): a metadata analysis to support the management of barrier zones around rewetted and newly constructed wetlands. Limnologica. 2014;45:69-79.

80. Ovaskainen O, Tikhonov G, Norberg A, Guillaume Blanchet F, Duan L, Dunson D, et al.
How to make more out of community data? A conceptual framework and its implementation as
models and software. *Ecology Letters*. 2017;20(5):561-76.
81. Dray S, Legendre P, Peres-Neto PR. Spatial modelling: a comprehensive framework for
principal coordinate analysis of neighbour matrices (PCNM). *Ecological modelling*. 2006;196(3-
4):483-93.
82. Geweke J. Evaluating the accuracy of sampling-based approaches to the calculation of
posterior moments: Federal Reserve Bank of Minneapolis, Research Department Minneapolis,
MN; 1991.
83. Efron B. Regression and ANOVA with zero-one data: Measures of residual variation.
Journal of the American Statistical Association. 1978;73(361):113-21.
84. Blanchet F, Tikhonov G, Norberg A. HMSC: Hierarchical modelling of species
community. R package version 2.0-10. 2017.
85. Team RC. R: A language and environment for statistical computing. Vienna, Austria;
2020.
86. CDC. Mosquito life cycle 2021 [October 13, 2021]. Available from:
<https://www.cdc.gov/dengue/resources/factsheets/mosquitolifecyclefinal.pdf>.
87. Balestrino F, Benedict M, Gilles J. A new larval tray and rack system for improved
mosquito mass rearing. *Journal of Medical Entomology*. 2012;49(3):595-605.
88. Morris C. Phenology of trophic and gonobiologic states in *Culiseta morsitans* and
Culiseta melanura (Diptera: Culicidae). *Journal of medical entomology*. 1984;21(1):38-51.
89. Kaufman MG, Fonseca DM. Invasion biology of *Aedes japonicus japonicus* (Diptera:
Culicidae). *Annual review of entomology*. 2014;59:31-49.

90. Wood DMD, P.T.; R.A. Ellis. . The insects and arachnids of Canada, part 6, the mosquitoes of Canada (Diptera: Culicidae. Biosystematics Research Institute, Ottawa, Ontario1979.
91. Darsie R, Hutchinson M. The mosquitoes of Pennsylvania: Identification of adult females and fourth instar larvae, geographical distribution, biology and public health importance. Technical Bulletin. 2009;1.
92. Brown H, Diuk-Wasser M, Andreadis T, Fish D. Remotely-sensed vegetation indices identify mosquito clusters of West Nile virus vectors in an urban landscape in the northeastern United States. Vector-Borne and Zoonotic Diseases. 2008;8(2):197-206.
93. Crans WJ. A classification system for mosquito life cycles: life cycle types for mosquitoes of the northeastern United States. Journal of Vector Ecology. 2004;29:1-10.
94. Winteringham FPW. Environment and chemicals in agriculture. 1985.
95. Plimmer J. Chemicals for agriculture. IAEA Bulletin. 1984;26(2):13-6.

Supporting information

S1 File.

(PDF)

Fig 1

Fig 2

Fig 3

Fig 4