1 **Nuclear Progesterone Receptor Mediated Linkage of Blood Coagulation** 2 **and Ovulation**

3

4 Jing Huang ^a, Chao Sun ^a, Dong Teng Liu^a, Nan Nan Zhao ^a, Jordan A. Shavit ^b, Yong Zhu ^{a, c, *},
5 Shi Xi Chen ^{a, d, *}

6 ^a State Key Laboratory of Marine Environmental Science, College of Ocean and Earth
7 Sciences, Xiamen University, Xiamen 361102, China

8 ^b Departments of Pediatrics and Human Genetics, University of Michigan, Ann Arbor, MI
9 48109, USA

10 ^c Department of Biology, East Carolina University, 1000 E 5th Street, Greenville, NC
11 27858, USA

12 ^d State-Province Joint Engineering Laboratory of Marine Bioproducts and Technology,
13 Xiamen University, Xiamen 361102, China

14

15 **Short title:** Pgr regulate coagulation factor for ovulation

16

17 **Keywords:** Pgr, progesterone, coagulation factor, ovulation

18

19

20

21

22

23

*** Corresponding authors and request for reprints:**

Dr. Shi Xi Chen

College of Ocean and Earth Sciences, Xiamen University, Xiamen 361102, China.

Phone: +86-13515969358, E-mail address: chenshixi@xmu.edu.cn.

Dr. Yong Zhu

Department of Biology, East Carolina University, 1000 E 5th Street, Greenville, NC

27858, USA. Phone: 252-328-6504, E-mail address: zhuy@ecu.edu.

Financial Support:

This work was supported by Fundamental Research Funds for the Central Universities (No. 20720200114), NSFC 31672628, and NSFC 41976092 to SXC, NIH GM100461 to YZ, and NIH R35 HL150784 to JAS.

Author Contributions:

JH performed most experiments, analyzed the data, and wrote the draft with the assistance of CS, DTL and NNZ. JAS generated the *f5* zebrafish knockout, provided protocols and experimental suggestions, discussed experimental results, and revised the manuscript. YZ supervised the project, conducted the fertility experiment, and revised the manuscript. SXC conceived and supervised the project and revised the manuscript.

Disclosure summary: The authors have nothing to disclose.

Abstract

Ovulation is a dramatic remodeling process including rupture of blood capillaries and clotting, but coagulation is not thought to directly regulate this process. Herein, we report remarkable increases of coagulation factors V (*f5*, ~3145-fold) and tissue factor (*f3a*, ~120-fold) in zebrafish ovarian follicle cells during ovulation. This increase was mediated through the nuclear progesterin receptor (Pgr), which is essential for ovulation in zebrafish. This increase was totally abolished in ovarian follicular cells from *pgr*^{-/-} mutants. In addition, promoter activities of *f5* and *f3a* were significantly enhanced by progesterin (DHP) via Pgr. Similar regulation of human *F5* promoter activity was induced via human PGRB, suggesting a conserved mechanism. Site-directed mutagenesis of the zebrafish *f5* promoter further demonstrated a direct regulation of coagulation factors via progesterin response elements. Moreover, a stark increase of erythrocytes occurred in capillaries meshed in wildtype preovulatory follicles, but was absent in *pgr*^{-/-} mutants. Interestingly, the anticoagulant heparin inhibited ovulation *in vitro*. Furthermore, reduced fecundity was observed in *f5*^{+/-} female zebrafish. Taken together, our study provides plausible evidence for steroid regulation of coagulation factors, and a new hypothesis for blood clotting triggered ovulation in vertebrates.

1. Introduction

Ovulation is a tissue remodeling process involving the rupture of follicular layers, basal membrane, and many blood capillaries intertwined in the follicular cells. This dramatic remodeling process share several features with inflammatory responses (1). Blood coagulation was suggested to be part of the innate defense in response to inflammation during ovulation (2). Formation of blood clots at apical vessels was found shortly before follicle rupture in several animal species including sheep, rabbits, and rodents (1,3-5). Cessation of blood flow likely due to blood coagulation was also observed in apical vessels of ovarian follicles during the late ovulatory phase in women (6). In addition, an increase of activity and expression of thrombin (a protease essential for fibrin formation) and its receptors were also reported in luteinized granulosa cells and follicular fluid in human (7,8). Increased fibrinogen secretion in bovine granulosa cells was thought to be related to the proteolytic activity needed for follicle rupture. Circulating F3 (also known as tissue factor) was elevated in women with polycystic ovary syndrome that is characterized by chronic oligo-ovulation or anovulation (9). Although these reports imply possible association between blood coagulation and the ovulation process, the mechanisms for this role in ovulation are still unclear.

Progestin and its nuclear progesterone receptor (PGR) are well-established initiators for vertebrate ovulation (10-15). Recently, we identified coagulation factors *f3a* and *f5* as potential downstream targets of Pgr in a transcriptomic analysis of preovulatory follicles from zebrafish (16). Interestingly, changes in these coagulation factors during ovulation appear to be conserved among human, mice, and zebrafish (16-18). Possible involvements of progestin or PGR in the regulation of coagulation have been hinted from

several studies using progestin contraceptives. F3 was upregulated in endometriosis and at endometrial bleeding sites in women using long-term progestin-only contraception (19). Orally administered progesterone reduced spontaneous miscarriages and improved fertility among women who had a history of recurrent miscarriages due to abnormal bleeding in early pregnancy (20). Increased risk of venous thromboembolic events was associated with combined oral contraceptives, particularly those containing the progestin desogestrel (21,22). A recent genetic study indicated that the present-day carriers of the Neandertal variant of PGR had less bleeding during early pregnancy because of high levels of PGR expression (23). However, evidence of progestin induced or PGR mediated regulation of coagulation factors, especially in ovarian follicles during ovulation, is lacking.

In this study, we examined the expression, hormonal regulation, and potential function of coagulation factors (*f5* & *f3*) in preovulatory follicles of zebrafish. The *f3* gene is duplicated in zebrafish (*f3a* and *f3b*) (24). We first demonstrated the expression and localization of *f5*, *f3a*, and *f3b* in preovulatory follicular cells. We found that progestin (DHP, 17 α , 20 β -dihydroxy-4-pregnen-3-one, a native ovulation-inducing ligand in zebrafish) via Pgr regulated *f5* and *f3a* expression in preovulatory follicles. We also found that follicular cells with strong Pgr expression were adjacent to blood capillaries, and inhibition of blood coagulation reduced ovulation. Finally, fertility and ovulation were reduced in *f5*^{+/-} zebrafish mutants *in vivo*. Our study provides credible evidence that coagulation factors (*f5* & *f3a*) in preovulatory follicular cells are regulated by progestin (DHP) via Pgr, which are well-known upstream initiators for ovulation. Our

results also suggest a previously unknown role of these coagulation factors in ovulation and female fertility.

2. Materials and Methods

2.1 Animals & Reagents

Zebrafish used are offspring of Tübingen strain unless stated otherwise. Detailed information for knockouts, transgenic lines, key chemicals, and reagents are listed in Supplemental Table 1 & Table 2 (25). Fish were fed three times daily with a commercial food (Otohime B2; Marubeni Nisshin Feed, Tokyo, Japan), supplemented with newly hatched brine shrimp, and were maintained in recirculating system (ESEN, Beijing, China) at 28 ± 0.5 °C under a controlled photoperiod (lights on at 08:00, off at 22:00). The daily ovulation cycle in mature adult zebrafish is established according to an established protocol (26). Briefly, each mature adult female and male zebrafish were paired in a spawning tank. These fish typically start spawning soon after lights on in the morning. Around noon, fish water was replaced, and fish were fed at least three times daily with the commercial food and supplemented with newly hatched brine shrimp. Around one hour prior to lights off, fish water was replaced again, a mesh inner tank was inserted into each spawning tank to prevent cannibalism typical for zebrafish adults. After one week, most fish form a daily ovulation cycle. When applicable, siblings from heterozygous in-cross were used for comparison. All experimental protocols were approved by the Institutional Animal Care and Use Committee (IACUC) at Xiamen University.

2.2 Collection of fully-grown follicles (stage □) during a daily spawning cycle

Fully-grown immature follicles (stage IVa, >650 μm, with visible germinal vesicles, i.e., GV, appears opaque) or mature follicles (stage IVb, GV disappears, appears transparent) were collected at four representative timepoints during a daily spawning cycle (Fig. 1A): 12:00 (IVa follicles); 20:00 (IVa follicles); 05:00 (the onset of oocyte maturation, IVa follicles); and 06:40 (completion of oocyte maturation but prior to ovulation, IVb follicles). To be simple and accurate in our description, we will use “IVa” to indicate fully-grown immature stage IVa follicles; “IVb” to indicate mature but yet ovulated follicles (prior to ovulation) throughout this manuscript.

To sacrifice fish humanely, the spinal cord and blood supply behind the gill cover were cut off swiftly using sharp scissors following a lethal dose of anesthetization (MS-222, 300 mg/L buffered solutions). Ovaries were immediately placed in 60 % L-15 media containing 15 mM HEPES at pH 7.2. Individual follicles were obtained according to an established procedure (26,27). At each timepoint, a total of 25 □a or IVb follicles (containing both germ cell and surrounding follicular cells) from 4 females were pooled as one sample. In addition, the follicular cells (~100 follicles) or their enclosed oocytes (~20 follicles) were collected separately from IVb follicles according to an established method (28). These samples were immediately homogenized in RNazol solution (Molecular Research Center, Cincinnati, USA) and stored at a -80 °C freezer until further processing.

151

152 **2.3 Hormone treatments**

153 According to established follicle classification system (26), follicles at the
154 different stages were collected at 14:00 from several females as described above. Intact
155 follicles with no damage were transferred into a 24-well culture plate containing 60 % L-
156 15 medium. These follicles were incubated at 28 °C with various doses of hormones or
157 various exposure times. Several progestin concentrations (DHP, 1-1,000 nM) with a 2 hrs
158 incubation or several exposure times (0.5-6 hrs) with 100 nM DHP were examined.
159 Based on the result of dose- and time-dependent experiments, follicles at different stages
160 were incubated with DHP (100 nM) for 2 hrs. Besides, stage IVa follicles were incubated
161 with hCG (50 IU/mL) (27) or DHP (100 nM) alone or in combination for 2 hrs. Stage IVa
162 follicles collected from *pgr*^{-/-} were incubated with DHP (100 nM) for 2 hrs. After
163 incubation, minimal 25 follicles with no damage were collected and homogenized in
164 RNazol, stored in a -80 °C freezer until RNA purification. Minimal three replicates were
165 collected at each timepoint.

166 **2.4 Real-time quantitative PCR (qPCR)**

167 RevertAid cDNA synthesis kit (Thermo Scientific, Waltham, USA) was used for
168 reverse transcription (500 ng total RNA/sample). All qPCRs were carried out in a 20 µL
169 reaction on the qTOWER 2.2 real-time PCR system (Analytik Jena AG, Jena, Germany)
170 using the PowerUp SYBR green master mix (Applied Biosystems, USA). Each PCR
171 primer of a pair was designed to target different exons of the gene (Supplemental Table

3) (25). Expression of the target genes was determined as relative values using the comparative C_i method (29) with the *eflα* gene serving as an internal control.

2.5 Whole-mount *in situ* hybridization (WISH)

A DNA fragment corresponding to partial *f5* or *f3a* cDNAs was amplified from ovarian cDNAs by PCR using the gene-specific primers (see Supplemental Table 3 for detail) (25) and cloned into the pGEM-T Easy vector (Promega, USA). Correct sequence and orientations of inserts were validated by Sanger sequencing. Antisense or sense digoxigenin-labeled cRNA probes for *f3a* or *f5* were synthesized by PCR using the plasmid cDNA templates, SP6 or T7 RNA polymerase depending on the insert orientation. IVa or IVb follicles were collected at 20:00 and 06:40 (immediately prior to ovulation), respectively, and fixed in 4 % PFA overnight at 4 °C. Following washing with PBST, the follicles were treated with proteinase K (5 µg/mL) for 2 min at 37 °C, then washed and subsequently hybridized with digoxigenin-labeled *f5* or *f3a* sense or antisense probes at 65 °C overnight using a previously established protocol (30). Following WISH, follicles were photographed using an M165FC dissecting microscope and DFC 550 digital camera (Leica, Wetzlar, Germany).

2.6 Dual-luciferase reporter assay

The Pgr expression and reporter assay plasmid constructs were listed in Supplemental Table 4 (25). Site-directed mutagenesis of four potential Pgr response elements (PRE, i.e., Pgr binding sites) in zebrafish *f5* proximal promoter sequences (Fig. 4E, Supplemental Table 6) (25) was conducted using a Q5 site-directed mutagenesis kit

(NEB, Ipswich, MA, USA) according to the manufacturer's protocol. Human embryonic kidney cells (HEK 293T) were used for the transactivation assays (31,32). Briefly, cells were maintained at 37 °C with 5 % CO₂ in phenol red-free Dulbecco's modified eagle's medium (Hyclone, USA) containing penicillin/streptomycin (Meilunbio, Dalian, China) and supplemented with 10 % fetal bovine serum (FBS, Gibco, Brazil). Lipofectamine 3000 (Life Technologies) was used for transient transfection according to manufactures' protocols. Cells were transiently transfected with a firefly luciferase reporter plasmid (3 µg/vector/60mm petri dish), the pRL-TK vector (150 ng/60mm petri dish) containing the Renilla luciferase reporter gene (as an internal control for transfection efficiency), and a Pgr expression vector or a control vector with no insert (1.5 µg/vector/60mm petri dish). After an overnight incubation at 37 °C, the medium was replaced with luciferase assay medium (DMEM without phenol red, supplemented with 10 % charcoal-stripped FBS) containing various concentrations of progestins (100 pM-1 µM DHP for zebrafish Pgr, P4 or human PGRB, respectively) or vehicle. Following additional 24 hrs of incubation at 37 °C, the cells were washed with PBS and harvested to determine firefly and Renilla luciferase activities according to the Dual-Luciferase reporter assay system (Promega, USA) on a luminometer (GloMax 20/20, Promega, USA). Firefly luciferase activities were normalized using Renilla luciferase data. After normalization for transfection efficiency, induction factors were calculated as the ratios of the average luciferase value of the steroid-treated samples versus vehicle-treated samples.

2.7 Blood capillaries in stage □ follicles

IVa immature follicles were obtained from paired female fish (*Tg(pgr:eGFP/fli1:DsRed)*). In this line, the blood capillaries are labeled with DsRed, while the follicular cells are labeled with eGFP. The nucleus of follicles was co-stained with Hoechst 33342 in a 29-mm glass-bottom dish (Cellvis, Hangzhou, China). Images were captured using an SP8 confocal laser scanning microscope (Leica, Germany). IVb mature follicles were obtained from ovaries undergoing ovulation from paired female fish (*Tg(fli1:eGFP)*) at 7:00. A series of images during ovulation was recorded using a Leica digital camera and fluorescent dissecting microscope described previously.

2.8 Erythrocyte numbers inside the capillaries on the surface of preovulatory follicles.

Erythrocytes inside the capillaries on the surface of ovarian follicles were stained with *o*-dianisidine according to an established protocol (33). Ovaries with IVb follicles were collected at 6:40 or 7:00 from siblings of *pgr*^{+/-} in-cross, producing both wildtype (*pgr*^{+/+}) and *pgr*^{-/-}. As sampling moves closely toward 7:00, IVb follicles will be further in the ovulation process in *pgr*^{+/+} females. In contrast, IVb follicles in *pgr*^{-/-} females will never initiate ovulation (13,15), which serves as an excellent control. After collection, the whole ovaries were immediately stained for 15 min at room temperature in the dark in 0.6 mg/ml *o*-dianisidine, 0.01 M sodium acetate (pH 4.5), 0.65 % H₂O₂, and 40 % (vol/vol) ethanol. Stained ovaries were washed 3 times for 2 min each with PBST. At least 3 ovary samples were collected at each timepoint for each genotype. Minimal 10 IVb follicles were randomly chosen from each sample for image capture using DM 2500 LED microscope and DFC 7000T digital camera (Leica), then the number of erythrocytes

under the field of view were counted by a reviewer who was blinded from sampling conditions and genotypes.

2.9 *In vitro* ovulation assay

Ovaries with IVb mature follicles from *Tg(fli1:eGFP)* zebrafish were collected at 06:40, 20 min before ovulation, and were placed in 60 % L-15 medium. Each ovary was immediately cut into several small pieces and then transferred into a 12-well plate with medium containing 2 mM EDTA, 100 µg/mL heparin, or 50 µg/mL warfarin, to examine their inhibitory effect on ovulation. The dose of each drug was chosen based on previous reports (34,35). Follicles were immediately dispersed by pipetting using a glass pipette (~1 mm in diameter). At least 50 IVb preovulatory follicles per well were used in each experiment. The follicles were cultured at 28 °C in a 2 mL culture medium for 2 hrs. Ovulation was continuously monitored until 09:00 under a fluorescent dissecting microscope (Leica M165FC), and the numbers of oocytes that had successfully ovulated were recorded. Minimal replicates for each treatment were eight.

2.10 Consecutive spawning test

The spawning of zebrafish was optimized under enhanced feeding conditions as described previously (26). Multiple pairs and spawning tanks were set up. Ten to twenty mature females (three months old) of *f5^{+/-}* or wildtype (*f5^{+/+}*) female siblings were used. Each female was individually paired and housed with a fertile wildtype male. Around 21:00 (1 hrs prior to lights off), water in spawning tanks was replaced, and a mesh inner tank was inserted into each spawning tank to prevent the eating of eggs by the adults. The

number of fertilized eggs was recorded for each pair of fish around noon of the next day. Then, water would be replaced, and fish would be fed at least three times a day. The $f5^{+/-}$ and wildtype ($f5^{+/+}$) females were continuously monitored for at least 3 weeks. Data from the last two weeks were used for analyses.

2.11 Statistical analyses

Data analyses were performed using GraphPad Prism version 8. Depending on the experimental setup, either independent (unpaired two-tailed) Student's t-test or one-way ANOVA (one-tailed) followed by the Dunnett's post hoc test was used to assess statistical differences in comparison to the control group. Each experiment (e.g. qPCR, reporter assay, WISH, drug exposure, reporter assay, erythrocytes count, *in vitro* ovulation, fertility test) was repeated at least once.

3. Results

3.1 Dramatic increase of $f5$ and $f3a$ transcripts in follicular cells of preovulatory follicles during a natural spawning cycle *in vivo*

Expression of $f5$ and $f3a$ mRNA was exclusively restricted to the follicular cells of mature follicles (IVb) and was not found in the germ cells (oocytes) (Fig. 1B). In contrast, $f3b$ mRNA was expressed in both follicular cells and oocytes (Fig. 1B). The expression of $f5$, $f3a$, and $f3b$ in mature follicles (IVb), just prior to ovulation, collected at 06:40 were ~3,145-fold, ~120-fold, and ~2-fold higher than those in immature but fully-

grown follicles (IVa, collected at 12:00, Fig. 1C). Since the increase in *f3b* expression was modest, we focused the rest of our efforts on *f5* and *f3a*. Results from whole mount *in situ* hybridization (WISH) showed that both *f5* and *f3a* were nearly undetectable in immature follicles (IVa, Fig. 1D, E). In contrast, strong signals of both *f5* and *f3a* transcripts were detected in the follicular cells of the mature follicles (IVb) (Fig. 1D, E), while no signal was detectable using sense probes (data not shown).

3.2 Progesterone upregulates *f5* and *f3a* transcripts through Pgr

3.2a Transcripts of *f5* and *f3a* were primarily upregulated by progesterone in preovulatory follicles

The progesterone (DHP) is a well-established native ligand for Pgr and an upstream hormonal inducer for ovulation in zebrafish (13,31,36). As predicted, the increase of *f5* and *f3a* transcripts were dose- and time-dependent in fully grown immature follicles (IVa) exposed to external progesterone (Fig. 2A-D).

Progesterone increased *f5* expression in vitellogenesis follicles (III, ~7 fold) and fully-grown immature follicles (IVa, ~337 fold) (Fig. 3A). In comparison, only fully-grown immature follicles (IVa) responded to progesterone with increase in *f3a* transcripts, but not the early developing follicles (stage I-III, Fig. 3B). LH is the primary endocrine factor for inducing ovulation in all animals including zebrafish (37). As expected, increased *f5* expression (~4 fold) was observed in IVa immature follicles exposed to hCG (50 IU/mL, Fig. 3C), but not for *f3a* (Fig. 3D). In comparison, progesterone (DHP) had much

stronger stimulatory effects on the expression of *f5* (~318 fold) and *f3a* (~10 fold) than hCG (Fig. 3C, D).

Significantly reduced *f5* and *f3a* expression was found in mature preovulatory follicles (IVb) collected from *pgr*^{-/-} fish compared to those in wildtype siblings (Fig. 3E). Importantly, DHP-induced *f5* and *f3a* expression were blocked in fully-grown immature follicles (IVa) from *pgr*^{-/-} fish (Fig. 3F).

3.2b Progesterin modulates promoter activity of *f5* and *f3a* via Pgr

As expected (31), progesterin induced dose-dependent, Pgr-mediated activation of the MMTV promoters (Fig. 4A). In the presence of Pgr, progesterin significantly increased the *f5* and *f3a* promoter activities, and this was dose-dependent (Fig. 4B, C). No change of promoter activity was observed when pcDNA3.1(+)-Pgr was replaced by a control vector (pcDNA3.1(+), Fig. 4B, C). The increase for *f5* was much bigger than that of the *f3a*. In the presence of human PGRB, P4 also significantly increased human *F5* promoter activity in a dose-dependent manner. No change in promoter activity was observed when pCMV-HA-hPGRB was replaced by control (pCMV-HA, Fig. 4D). Moreover, site-directed mutagenesis of four predicted progesterone response elements (PRE) significantly decreased zebrafish *f5* promoter activity (Fig. 4E, Supplemental Table 5) (25). A PRE (5'-AGGACATTTT-3') localizing at -1438 to -1425 of zebrafish *f5* appears to be the primary mediator of Pgr regulation (Fig. 4E).

3.3 Coagulation factor contribution to ovulation

3.3a Follicular cells with strong Pgr expression localize adjacent to blood vessels

Intriguingly, we found almost all the follicular cells with strong Pgr expression were located adjacent to capillary vessel networks on the surface of the IVa follicles in a dual transgenic zebrafish line *Tg(pgr:eGFP/fli1:DsRed)* (Fig. 5).

3.3b Rupture of blood vessels during ovulation

The ovulation process started with the appearance of a small hole in follicular cells and a ruptured blood vessel (indicated by a red arrow in Fig. 6A, supplemental video clip A) (25) with a concomitant appearance of the ovulating oocyte on the surface of the ovaries. When the hole of the broken follicular cell layer reached a certain size, the oocyte was quickly released from the follicular layer (Fig. 6B-E, supplemental video clip A) (25).

3.3c Increased numbers of erythrocytes appear in the lumen of blood vessels of mature preovulatory follicles as ovulation proceeds

The numbers of erythrocytes in capillaries of mature preovulatory follicles (IVb) increased as they proceeded to follicular rupture, from ~06:40, (Fig. 7A, D, C) to ~07:00 (Fig. 7B, E, G). This effect was absent in the same mature follicles (IVb) collected from anovulatory *pgr*^{-/-} female siblings around 07:00 (Fig. 7C, F, G).

3.3d Heparin inhibits ovulation *in vitro*

The intact IVb follicles or ovulated follicles from *Tg(fli1:eGFP)* were easily recognizable by the interconnecting blood vessels with strong GFP expression and dramatic changes in size (Fig. 8A-D). As expected, the ovulation rate decreased to 41 %

when follicles were exposed to EDTA (2 mM), an inhibitor of metalloproteases that are required for ovulation (35). Intriguingly, heparin (100 µg/mL), an anticoagulant, significantly reduced the rate of ovulation to 66% (Fig. 8E). Another anticoagulant, warfarin (50 µg/mL), had a reduced but not significant adverse effect on ovulation rate (83 %) (Fig. 8E).

3.3e Reduced fertility in $f5^{+/-}$ zebrafish females *in vivo*

Although $f5$ homozygous mutants live to early adulthood, most die by 4-6 months of age and do not breed well (38). Therefore, we examined female fertility in $f5^{+/-}$ mutants in a consecutive spawning test. We observed significantly reduced numbers of embryos per spawning and total embryos over two weeks in comparison to wildtype sibling females. (Fig. 9A, B). The spawning frequency of $f5^{+/-}$ female fish also significantly decreased compared to that of wildtype siblings (Fig. 9C). One $f5^{+/-}$ female only spawned once in two weeks (Fig. 9C). The spawning intervals of $f5^{+/-}$ females were significantly longer in comparison to that of wildtype siblings (Fig. 9D).

4. Discussion

We have provided comprehensive evidence demonstrating that the expression of coagulation factors $f5$ and $f3a$ in preovulatory follicles is controlled by progestin and its cognate receptor Pgr. Increased expression of these coagulation factors is likely critical for the regulation of ovulation and female fertility in addition to their well-established roles in hemostasis. Our conclusions are based on the following evidence: 1) Coagulation factors, $f5$ and $f3a$, are expressed in the follicular cells and increase dramatically prior to

ovulation during natural spawning cycles in zebrafish; 2) Expression of these cofactors in preovulatory follicles is controlled by progesterin via Pgr, which are essential and upstream regulators for ovulation; 3) Follicular cells with strong Pgr expression are adjacent to blood capillaries; 4). As follicular cells near rupture, a significant increase in the numbers of erythrocytes in blood vessels suggests blood clotting which is correlated with a dramatic increase of *f5* and *f3a* in preovulatory follicular cells; 5) Inhibition of coagulation leads to reduced ovulation; 6) Low fertility is found in *f5*^{+/-} zebrafish female mutants.

A major finding of the present study is the regulation of *f5* and *f3a* by progesterin via Pgr in the preovulatory follicles during ovulation (Fig.10). F3 is expressed in a wide array of tissues (39). It has been demonstrated that physiological concentrations of P4 could increase both F3 mRNA and protein in breast cancer cell lines ZR-75 and T47D (40). Increased F3 expression was correlated with an increase of procoagulant activity, which was hypothesized to be responsible for the increased risk of thrombosis in breast cancer patients (41). Studies of the uterus have also shown that RU486, an antagonist of PGR, not only blocks but also reverses progesterin-enhanced expression of F3 mRNA and protein in stromal cells (42). In line with these studies, PGR binding sites were identified in the promoter region of the human F3 gene (Supplemental Table 5) (25,43). The increase of *f3a* expression in the preovulatory follicles is primarily controlled by Pgr (current study). A difference in the weak activity of *f3a* promoters observed in a human immortal cell line (HEK293T) *in vitro* in comparison to progesterin-induced strong *f3a*

promoter activities found in zebrafish follicles both *in vivo* and *in vitro* may be due to missing core *cis*-regulatory elements in the cloned promoter sequence of zebrafish *f3a* (Supplemental Table 5) (25). Nevertheless, our results demonstrate that like *f5*, *f3a* expression is also controlled by progestin via Pgr (Supplemental Table 5, Fig.10) (25).

Until now, no information was available for progestin regulation of F5. Estrogen and progesterone receptors regulate *F5* expression in mouse bone marrow/bone (44,45). The present study demonstrates that a new extrahepatic site of *f5* expression is regulated by progestin through Pgr in zebrafish. Moreover, progestin enhances the promoter activities of *f5* in zebrafish and F5 in humans via PGR. Several putative PREs were predicted in the promoter regions of human, mouse, and zebrafish *F5* (Supplemental Table 6) (25). Our site-direct mutation analyses of these PREs in the zebrafish *f5* promoter suggests direct regulation of progestin/Pgr via binding to and activating these a PREs. One PRE site (5'- TGACTCA-3') in the *f5* promoter is also conserved between zebrafish, mice and humans (Supplemental Table 6) (25). Whether P4 and/or PGR regulate *F5* expression in human ovaries *in vivo* will be an interesting topic for further investigation.

The functions of coagulation factors in hemostasis are well established (46-48). However, the role of coagulation factors during ovulation is significantly less studied. Formation of blood clots at apical blood vessels was found shortly before follicular rupture in several mammalian species (1,3-6). In addition, the permeability of blood vessels increased 3-fold prior to ovulation in rats (49), which likely facilitates material exchange between the follicles and blood vessels. The increase of material exchange

might be important for anticipating local but avoiding systemic activation of blood clotting. Our observation of ruptured blood vessels on the surface of ovulating follicles further supports this hypothesis. In addition, we also found a significant increase of erythrocytes in capillaries on the surface of ovulating follicles as they progress to rupture/ovulation, suggestive of blood clotting. Lack of functional F5 or F8 caused various bleeding symptoms, including recurrent and serious ovulation-related blood loss in women (50). Occurrence of blood clotting in advance of follicle rupture could preemptively stop excessive bleeding caused by ovulation-induced capillary rupture. The granulosa cells appear to be the primary cells for synthesizing F5 and F3a, as these account for most that express Pgr (this study & (10,36)). Intriguingly, we also found that strong Pgr expressing theca cells were localized near capillaries on the follicles. Whether these theca cells are key switchers of F3a-initiated hemostasis in the capillaries around ovarian follicles will be an interesting topic to examine in the future.

The dramatic induction of *f5* and *f3a* expression in preovulatory follicles via Pgr, an essential mediator for activation of various ovulation-related genes, suggests that locally produced F5 and F3a may be required for ovulation. Activation of multiple metalloproteases within the ovulating follicles at the right times is a critical step for sequential digestion of extracellular matrix proteins of the follicular cells that are required for ovulation (1,37). Their activation and substrate binding are precisely regulated by the timing of prodomain cleavage of the enzyme and removal of endogenous tissue inhibitors of metalloproteases (Timp) (51). Protease inhibitors (Timp and other inhibitors) are typically present at high concentrations in the serum to prevent metalloproteases from

428 acting accidentally (52). Vasoconstriction of blood vessels within the thecal layer at the
 429 apex of the follicles was suggested to be critical for the removal of protease inhibition
 430 from the circulation (53). We hypothesize that coagulation that occurs shortly before
 431 ovulation in the follicles may have a similar function. In addition to the removal of
 432 protease inhibitors from serum, the cleavage of propeptide from latent form of
 433 metalloproteinase is required for activation and thus ovulation (Fig.10). An F3-dependent
 434 pathway of thrombin generation was also shown in bovine, equine, and human follicular
 435 fluid (7,54,55). Granulosa cells were shown to express and secrete fibrinogen, which is
 436 catalyzed by thrombin to become fibrin (56). Fibrin formation could stimulate activities
 437 of plasmin/plasminogen activators (plat and plau) signal, which is well-established for
 438 plasminogen (plg) activation (57). Furthermore, the function of Plat, Plau, and Plg in
 439 metalloproteinase activations during ovulation is also recognized (58,59). Therefore, we
 440 hypothesize that a dramatic increase of F3a and F5 expression in the follicular cells may
 441 facilitate the formation of fibrin, followed by the activation of Plat, Plau, and Plg, which
 442 further activate metalloproteinases. Taken together, as blood coagulation occurs, blood
 443 flow decreases, then stops completely, effectively terminating inhibitory effects of
 444 protease inhibitors from the serum, triggering metalloprotease activations to start the
 445 degradation of the follicular extracellular matrix for preparing the release of a mature
 446 oocyte (Fig.10). A patient diagnosed with severe F5 deficiency had to undergo bilateral
 447 ovariectomies because of the presence of hemorrhagic anovulatory follicles and life-
 448 threatening hemorrhage in the pouch of Douglas (60). Circulating F3 has been shown to
 449 be elevated in women with polycystic ovary syndrome, the most common form of
 450 anovulatory infertility (9). Consistent with findings from medaka (35), the rate of

ovulation was suppressed by EDTA treatment in the current study likely via inhibition of metalloproteases, or possibly inhibition of coagulation. EDTA is a known anticoagulant, acting through chelation of calcium, as many coagulation factors depend on this ion for function. Indeed, the known anticoagulant heparin also suppressed ovulation. The anticoagulant warfarin did not, but this was unsurprising since its mechanism of action usually requires days of treatment. Another finding in the present study is that reduced fecundity and impaired ovulation were observed in $f5^{+/-}$ female zebrafish *in vivo*. Together, these results argue for conserved functions of coagulation factors in the regulation of ovulation in vertebrate ovaries. Long-term efforts in development of new instruments and new techniques/methods, with suitable *in vivo* imaging and examining the association of blood flow and timing of metalloproteases activation, are required for further elucidation of underlying mechanisms and testing our hypothetical model (Fig.10).

In summary, our results demonstrate that $f5$ and $f3a$ expression are dramatically induced by progesterin via a Pgr-dependent mechanism in preovulatory follicles (Fig. 10). Moreover, these coagulation factors may regulate ovulation and female fertility in addition to their roles in hemostasis. Future studies on the regulation and functions of coagulation factors in other organisms including mammalian follicles or ovarian fluid during ovulation will likely lead to developments of diagnostic and therapeutic modalities for patients with gynecological problems associated with mutations and/or dysregulation of coagulation.

Acknowledgments

We want to thank Dr. Mingyu Li (Xiamen University, China) and Dr. Yu Xue (Minnan Normal University, China) for providing zebrafish transgenic lines (*Tg(fli1:eGFP)*, *Tg(fli1:DsRed)*), Dr. Shuang-Bo Kong (Xiamen University, Xiamen, China) for sharing human PGRB expression vector (pCMV-HA-hPGRB). This work was supported by Fundamental Research Funds for the Central Universities (No. 20720200114), NSFC 31672628, and NSFC 41976092 to SXC, NIH GM100461 to YZ, and NIH R35 HL150784 to JAS.

Data Availability Statement(s): Some or all data generated or analyzed during this study are included in this published article or in the data repositories listed in References 25.

References

1. Duffy DM, Ko C, Jo M, Brannstrom M, Curry TE. Ovulation: Parallels With Inflammatory Processes. *Endocr Rev.* 2019;40(2):369-416.
2. Esmon CT. The interactions between inflammation and coagulation. *British Journal of Haematology.* 2005;131(4):417-430.
3. Kanzaki H, Okamura H, Okuda Y, Takenaka A, Morimoto K, Nishimura T. Scanning electron-microscopic study of rabbit ovarian follicle microvasculature using resin injection corrosion casts. *Journal of Anatomy.* 1982;134(4):697-704.
4. Cavender JL, Murdoch WJ. Morphological-studies of the microcirculatory system of periovulatory ovine follicles. *Biol Reprod.* 1988;39(4):989-997.
5. Zackrisson U, Lofman CO, Janson PO, Wallin A, Mikuni M, Brannstrom M. Alterations of follicular microcirculation and apex structure during ovulation in the rat. *Eur J Obstet Gyn R B.* 2011;157(2):169-174.
6. Brannstrom M, Zackrisson U, Hagstrom HG, Josefsson B, Hellberg P, Granberg S, Collins WP, Bourne T. Preovulatory changes of blood flow in different regions of the human follicle. *Fertil Steril.* 1998;69(3):435-442.
7. Gentry PA, Plante L, Schroeder MOB, LaMarre J, Young JE, Dodds WG. Human ovarian follicular fluid has functional systems for the generation and modulation of thrombin. *Fertil Steril.* 2000;73(4):848-854.
8. Kim YS, Kim MS, Lee SH, Choi BC, Lim JM, Cha KY, Baek KH. Proteomic analysis of recurrent spontaneous abortion: Identification of an inadequately expressed set of proteins in human follicular fluid. *Proteomics.* 2006;6(11):3445-3454.
9. Gonzalez F, Kirwan JP, Rote NS, Minium J. Elevated circulating levels of tissue factor in polycystic ovary syndrome. *Clin Appl Thromb-Hem.* 2013;19(1):66-72.
10. Natraj U, Richards JS. Hormonal-regulation, localization, and functional-activity of the progesterone-receptor in granulosa-cells of rat preovulatory follicles. *Endocrinology.* 1993;133(2):761-769.
11. Lydon JP, Demayo FJ, Funk CR, Mani SK, Hughes AR, Montgomery CA, Shyamala G, Conneely OM, Omalley BW. Mice Lacking Progesterone-Receptor Exhibit Pleiotropic Reproductive Abnormalities. *Gene Dev.* 1995;9(18):2266-2278.
12. Hagiwara A, Ogiwara K, Katsu Y, Takahashi T. Luteinizing hormone-induced expression of ptger4b, a Prostaglandin E-2 receptor indispensable for ovulation of the medaka *Oryzias latipes*, is regulated by a genomic mechanism involving nuclear progesterin receptor. *Biol Reprod.* 2014;90(6):1-14.
13. Zhu Y, Liu DT, Shaner ZC, Chen SX, Hong WS, Stellwag EJ. Nuclear progesterin receptor (Pgr) knockouts in zebrafish demonstrate role for Pgr in ovulation but not in rapid non-genomic steroid mediated meiosis resumption. *Front Endocrinol.* 2015;6:37.
14. Kubota K, Cui W, Dhakal P, Wolfe MW, Rumi MAK, Vivian JL, Roby KF, Soares MJ. Rethinking progesterone regulation of female reproductive cyclicity. *P Natl Acad Sci USA.* 2016;113(15):4212-4217.
15. Tang HP, Liu Y, Li JZ, Yin YK, Li GF, Chen Y, Li SS, Zhang Y, Lin HR, Liu XC, Cheng CHK. Gene knockout of nuclear progesterone receptor provides insights into the regulation of ovulation by LH signaling in zebrafish. *Sci Rep.* 2016;6:28545.
16. Liu DT, Brewer MS, Chen SX, Hong WS, Zhu Y. Transcriptomic signatures for ovulation in vertebrates. *Gen Comp Endocr.* 2017;247:74-86.
17. Wissing ML, Kristensen SG, Andersen CY, Mikkelsen AL, Hst T, Borup R, Grondahl ML. Identification of new ovulation-related genes in humans by comparing the transcriptome of granulosa cells before and after ovulation triggering in the same controlled ovarian stimulation cycle. *Hum Reprod.* 2014;29(5):997-1010.
18. Carletti MZ, Christenson LK. Rapid effects of LH on gene expression in the mural granulosa cells of mouse periovulatory follicles. *Reproduction.* 2009;137(5):843-855.
19. Runic R, Schatz F, Wan L, Demopoulos R, Krikun G, Lockwood CJ. Effects of norplant on

538 endometrial tissue factor expression and blood vessel structure. *J Clin Endocr Metab.*
539 2000;85(10):3853-3859.

540 20. Haas DM, Hathaway TJ, Ramsey PS. Progestogen for preventing miscarriage in women with
541 recurrent miscarriage of unclear etiology. *Cochrane Db Syst Rev.* 2019(11):1-50.

542 21. Jick H, Jick SS, Gurewich V, Myers MW, Vasilakis C. Risk of Idiopathic Cardiovascular Death
543 and Nonfatal Venous Thromboembolism in Women Using Oral-Contraceptives with Differing
544 Progestagen Components. *Lancet.* 1995;346(8990):1589-1593.

545 22. Farley TMM, Meirik O, Chang CL, Marmot MG, Poulter NR. Effect of different progestogens in
546 low estrogen oral contraceptives on venous thromboembolic disease. *Lancet.*
547 1995;346(8990):1582-1588.

548 23. Zeberg H, Kelso J, Paabo S. The neandertal progesterone receptor. *Mol Biol Evol.*
549 2020;37(9):2655-2660.

550 24. Stein C, Caccamo M, Laird G, Leptin M. Conservation and divergence of gene families
551 encoding components of innate immune response systems in zebrafish. *Genome Biol.*
552 2007;8(11):1-23.

553 25. Huang J, Sun C, Liu DT, Zhao NN, Shavit JA, Zhu Y, Chen SX. Data from: Nuclear Progesterin
554 Receptor Mediated Linkage of Blood Coagulation and Ovulation. *Figureshare.* 2021:
555 <https://doi.org/10.6084/m9.figshare.17427326>.

556 26. Liu DT, Carter NJ, Wu XJ, Hong WS, Chen SX, Zhu Y. Progesterin and nuclear progesterin receptor
557 are essential for upregulation of metalloproteinase in zebrafish preovulatory follicles. *Front*
558 *Endocrinol.* 2018;9:517.

559 27. Liu DT, Hong WS, Chen SX, Zhu Y. Upregulation of adamts9 by gonadotropin in preovulatory
560 follicles of zebrafish. *Mol Cell Endocrinol.* 2020;499:110608.

561 28. Hanna RN, Zhu Y. Controls of meiotic signaling by membrane or nuclear progesterin receptor in
562 zebrafish follicle-enclosed oocytes. *Molecular and Cellular Endocrinology.* 2011;337(1-2):80-
563 88.

564 29. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C-T method. *Nat*
565 *Protoc.* 2008;3(6):1101-1108.

566 30. Hauptmann G, Gerster T. Multicolor whole-mount in situ hybridization. *Methods Mol Biol.*
567 2000;137:139-148.

568 31. Chen SX, Bogerd J, Garcia-Lopez A, de Jonge H, de Waal PP, Hong WS, Schulz RW. Molecular
569 Cloning and Functional Characterization of a Zebrafish Nuclear Progesterone Receptor. *Biol*
570 *Reprod.* 2010;82(1):171-181.

571 32. Zhang YT, Hong WS, Liu DT, Qiu HT, Zhu Y, Chen SX. Involvement of membrane progesterin
572 receptor Beta (mPRbeta/Paqr8) in sex pheromone progesterin-induced expression of
573 luteinizing hormone in the pituitary of male chinese black sleeper (*Bostrychus Sinensis*).
574 *Front Endocrinol.* 2018;9:397.

575 33. Detrich HW, Kieran MW, Chan FY, Barone LM, Yee K, Rundstadler JA, Pratt S, Ransom D, Zon LI.
576 Intraembryonic hematopoietic-cell migration during vertebrate development. *P Natl Acad Sci*
577 *USA.* 1995;92(23):10713-10717.

578 34. Hanumanthaiah R, Thankavel B, Day K, Gregory M, Jagadeeswaran P. Developmental
579 expression of vitamin K-dependent gamma-carboxylase activity in zebrafish embryos: effect
580 of warfarin. *Blood Cells Mol Dis.* 2001;27(6):992-999.

581 35. Ogiwara K, Takano N, Shinohara M, Murakami M, Takahashi T. Gelatinase A and membrane-
582 type matrix metalloproteinases 1 and 2 are responsible for follicle rupture during ovulation
583 in the medaka. *Proc Natl Acad Sci U S A.* 2005;102(24):8442-8447.

584 36. Hanna RN, Daly SCJ, Pang YF, Anglade I, Kah O, Thomas P, Zhu Y. Characterization and
585 expression of the nuclear progesterin receptor in zebrafish gonads and brain. *Biol Reprod.*
586 2010;82(1):112-122.

587 37. Takahashi T, Hagiwara A, Ogiwara K. Follicle rupture during ovulation with an emphasis on
588 recent progress in fish models. *Reproduction.* 2019;157(1):R1-R13.

589 38. Weyand AC, Grzegorski SJ, Rost MS, Lavik KI, Ferguson AC, Menegatti M, Richter CE, Asselta R,
590 Duga S, Peyvandi F, Shavit JA. Analysis of factor V in zebrafish demonstrates minimal levels

needed for early hemostasis. *Blood Adv.* 2019;3(11):1670-1680.

39. Osterud B, Bjorklid E. Sources of tissue factor. *Semin Thromb Hemost.* 2006;32(1):11-23.

40. Kato S, Pinto M, Carvajal A, Espinoza N, Monso C, Sadarangani A, Villalon M, Brosens JJ, White JO, Richer JK, Horwitz KB, Owen GI. Progesterone increases tissue factor gene expression, procoagulant activity, and invasion in the breast cancer cell line ZR-75-1. *J Clin Endocr Metab.* 2005;90(2):1181-1188.

41. Richer JK, Jacobsen BM, Manning NG, Abel MG, Wolf DM, Horwitz KB. Differential gene regulation by the two progesterone receptor isoforms in human breast cancer cells. *J Biol Chem.* 2002;277(7):5209-5218.

42. Lockwood CJ, Krikun G, Papp C, Aigner S, Nemerson Y, Schatz F. Biological mechanisms underlying ru-486 clinical effects - inhibition of endometrial stromal cell tissue factor content. *J Clin Endocr Metab.* 1994;79(3):786-790.

43. Bravo ML, Pinto MP, Gonzalez I, Oliva B, Kato S, Cuello MA, Lange CA, Owen GI. Progesterone regulation of tissue factor depends on MEK1/2 activation and requires the proline-rich site on progesterone receptor. *Endocrine.* 2015;48(1):309-320.

44. Moverare S, Skrtic S, Lindberg MK, Dahlman-Wright K, Ohlsson C. Estrogen increases coagulation factor V mRNA levels via both estrogen receptor-alpha and -beta in murine bone marrow/bone. *Eur J Endocrinol.* 2004;151(2):259-263.

45. Cleuren AC, Van der Linden IK, De Visser YP, Wagenaar GT, Reitsma PH, Van Vlijmen BJ. 17alpha-Ethinylestradiol rapidly alters transcript levels of murine coagulation genes via estrogen receptor alpha. *J Thromb Haemost.* 2010;8(8):1838-1846.

46. Spronk HM, Govers-Riemslog JW, ten Cate H. The blood coagulation system as a molecular machine. *BioEssays : news and reviews in molecular, cellular and developmental biology.* 2003;25(12):1220-1228.

47. Esmon CT. The impact of the inflammatory response on coagulation. *Thromb Res.* 2004;114(5-6):321-327.

48. Hathcock JJ. Flow effects on coagulation and thrombosis. *Arterioscler Thromb Vasc Biol.* 2006;26(8):1729-1737.

49. Mitsube K, Brannstrom M, Haraldsson B. Modulation of microvascular permeability in the preovulatory rat ovary by an ovulatory gonadotropin stimulus. *Fertil Steril.* 2013;99(3):903-909.

50. Spiliopoulos D, Kadir RA. Congenital factor V and VIII deficiency in women: a systematic review of literature and report of two new cases. *Blood Coagul Fibrin.* 2016;27(3):237-241.

51. Zhu Y. Metalloproteases in gonad formation and ovulation. *Gen Comp Endocr.* 2021;314:113924.

52. Travis J, Salvesen GS. Human plasma proteinase inhibitors. *Annu Rev Biochem.* 1983;52:655-709.

53. Migone FF, Cowan RG, Williams RM, Gorse KJ, Zipfel WR, Quirk SM. In vivo imaging reveals an essential role of vasoconstriction in rupture of the ovarian follicle at ovulation. *Proc Natl Acad Sci U S A.* 2016;113(8):2294-2299.

54. Yamada M, Gentry PA. The hemostatic profile of equine ovarian follicular fluid. *Thrombosis Research.* 1995;77(1):45-54.

55. Roach LE, Petrik JJ, Plante L, LaMarre J, Gentry PA. Thrombin generation and presence of thrombin receptor in ovarian follicles. *Biol Reprod.* 2002;66(5):1350-1358.

56. Parrott JA, Whaley PD, Skinner MK. Extrahepatic expression of fibrinogen by granulosa cells: potential role in ovulation. *Endocrinology.* 1993;133(4):1645-1649.

57. Medved L, Nieuwenhuizen W. Molecular mechanisms of initiation of fibrinolysis by fibrin. *Thromb Haemost.* 2003;89(3):409-419.

58. Beers WH, Strickland S, Reich E. Ovarian plasminogen activator: Relationship to ovulation and hormonal regulation. *Cell.* 1975;6(3):387-394.

59. Ogiwara K, Minagawa K, Takano N, Kageyama T, Takahashi T. Apparent involvement of plasmin in early-stage follicle rupture during ovulation in medaka. *Biol Reprod.* 2012;86(4):113.

- 644 60. Montefusco MC, Duga S, Asselta R, Malcovati M, Peyvandi F, Santagostino E, Mannucci PM,
645 Tenchini ML. Clinical and molecular characterization of 6 patients affected by severe
646 deficiency of coagulation factor V: broadening of the mutational spectrum of factor V gene
647 and in vitro analysis of the newly identified missense mutations. *Blood*. 2003;102(9):3210-
648 3216.
649

Figure

Figure 1. Dramatic increase of *f5* and *f3a* transcripts in follicular cells of preovulatory follicles during a daily spawning cycle *in vivo*.

(A) Schematic illustration of daily spawning cycle in zebrafish synchronized with lights on. Oocyte maturation in zebrafish typically occurs around 06:40, while ovulation occurs around 07:00.

(B) Cell-specific expression of *f5*, *f3a*, and *f3b* in mature follicles examined by RT-PCR. Transcripts of *f5* and *f3a* were expressed in the follicular cells but not in the oocytes, while the transcript of *f3b* was expressed in both follicular cells and oocytes of mature preovulatory follicles. MK: molecular makers; I: intact follicles including both surrounding follicular cells and germ cell; O: denuded oocyte (without follicular cells); F: follicular cells (with no germ cell); N: no template control.

(C) Dramatic increase of *f5*, *f3a*, and *f3b* transcripts in preovulatory follicles during a daily spawning cycle *in vivo*. Fully-grown ovarian follicles (stage IVa or IVb) were collected from wildtype mature female zebrafish at four representative timepoints during a daily spawning cycle (see Fig. 1A). Expression of *f5*, *f3a*, and *f3b* mRNA were determined by qPCR and normalized to the internal housekeeping gene (*eflα*).

Data are expressed as the mean $\pm$ SEM (N = 4) relative to the respective transcript levels of *f5* in IVa follicles collected at 12:00. P-values were calculated by one-tailed one-way ANOVA followed by Dunnett's test against respective controls (samples collected at 12:00).

(D-G) Dramatic increase of *f5* and *f3a* transcripts in the follicular cells of mature follicles. Fully-grown but immature (stage IVa) and mature follicles but prior to ovulation (stage IVb) were collected at 20:00 and 06:40, respectively. Strong signals of *f5* and *f3a* transcripts were only observed in mature follicles (IVb) by whole-mount *in situ* hybridization (WISH), while no signals were observed in the immature follicles (IVa, D, E). The positive ISH signals were localized in the follicular cells (F, G). Scale bar, 250 μ m (D, E) and 50 μ m (F, G).

Figure 2. Transient increase of *f5* and *f3a* expression in preovulatory follicles in response to stimulation of progestin *in vitro*.

(A, C) Effects of various doses of progestin (DHP) (2 hours exposure) on *f5* and *f3a* expression in fully grown immature preovulatory follicles (stage IVa) *in vitro*. (B, D) Effects of DHP (100 nM) on *f5* and *f3a* expression in preovulatory follicles (IVa) at various timepoints *in vitro*. P-values were calculated by one-tailed one-way ANOVA followed by Dunnett's test against respective controls (0 timepoint or vehicle control).

Figure 3. Increase of *f5* and *f3a* transcripts in late-stage follicles is mediated by *Pgr*.

Follicles were incubated with progestin (DHP, 100nM), or LH (hCG, 50 IU/ml) alone, or in combination for 2 hrs at 28 °C *in vitro* before qPCR analysis. (A) An increase of *f5* transcript was observed in late-stage (III and IVa) follicles from wildtype (*pgr*^{+/+}) zebrafish, but not in early-stage follicles (I or II) from wildtype zebrafish, in response to progestin stimulation. (B) An increase of *f3a* expression was observed only in fully-grown immature follicles (IVa) from wildtype (*pgr*^{+/+}) zebrafish, but not in developing follicles (I-III) from wildtype fish, in response to progestin stimulation. (C) Synergistic increase of *f5* mRNA by hCG and DHP in fully grown immature follicles (IVa) from wildtype zebrafish. (D) Progestin but not hCG stimulated the expression of *f3a* mRNA in full-grown immature follicles (IVa) from wildtype zebrafish. (E) Both *f3a* and *f5* transcripts in preovulatory follicles (stage IVb) of *pgr*^{-/-} were significantly lower than that of *pgr*^{+/+} (WT). (F) Stimulatory effect of progestin (DHP) for *f3a* and *f5* expression was abolished in the stage IVa follicles of *pgr*^{-/-} mutants. The expression of *f5* and *f3a* transcripts were determined by qPCR and normalized to an internal control (*ef1a*). Data are expressed as the mean ± SEM (N = 3) relative to the respective transcript levels of target gene measured in wildtype, or vehicle treatment group. P-values were calculated by two-tailed Students' t-test.

Figure 4. Progestin enhances promoter activity of *f5* and *f3a* via Pgr.

HEK293T cells were transiently co-transfected with a firefly luciferase reporter vector containing putative Pgr binding elements (3 μ g/vector/60mm petri dish), a pRL-TK vector (150 ng/60mm petri dish) containing the Renilla luciferase reporter gene (as a control for transfection efficiency), and a Pgr

expression vector (1.5 μ g/vector/60mm petri dish) or an empty control expression vector (1.5 μ g/vector/60mm petri dish). In the presence of zebrafish Pgr (zPgr), progestin significantly increased MMTV (panel **A**), *zf5* (panel **B**), and *zf3a* (panel **C**) promoter activities. This increase was dose-dependent. In the presence of human PGRB (hPGRB), P4 also significantly increased human *hF5* promoter activity, and the increase was dose-dependent (Panel **D**). Site-directed mutagenesis of four likely progestin receptor response elements (PRE) decreased the *zf5* promoter activity (Panel **E**). HEK293T cells were incubated for 24 hrs with increasing concentrations of progestins (DHP or P4: 100 pM to 1 μ M). Extracts of the treated cells were assayed for luciferase activity. Values are shown relative to the luciferase activity of the vehicle treatment group. Data are expressed as mean $\pm$ SEM (N = 3). P-values were calculated by one-tailed one-way ANOVA followed by Dunnett's test against respective controls (vehicle controls, zero dose).

Figure 5. Follicular cells with strong Pgr expression are adjacent to blood capillaries.

Images of a representative stage IVa follicle from a double transgenic zebrafish line *Tg(pgr:eGFP/fli1:DsRed)* are shown. (A) Nucleus staining of follicular cells by Hoechst 33342. (B) Pgr promoter drives GFP expression in follicular cells. (C) Red fluorescence-labeled vascular endothelial cells on the follicles. (D) Merged images to show that follicular cells with strong GFP expression driven by the Pgr promoters are adjacent to blood capillaries. Red arrowheads indicate follicular cells with strong GFP expression driven by the Pgr promoters. Scale bars: 100 μ m.

Figure 6. Rupture of blood capillaries and follicular cells during ovulation.

Ovaries undergoing ovulation were collected from *Tg(fli1:eGFP)* females at 07:00. (A-F), a series of images were taken when a mature (stage IVb) ovarian follicle started ovulation. Interconnected blood vessels labeled with GFP were observed easily on the surface of ovarian follicles. A red arrow indicates a ruptured blood vessel on the surface of an ovulating follicle. Red arrowheads and red dotted lines indicate a hole left in the remaining follicular cells after ovulation. The white dotted line represents the remaining

730 follicular cells. Scale bars: 250 μ m. Oc, oocyte. (Please also see supplemental video clip A for additional
731 details) (25).

Figure 7. Increase of erythrocytes in capillaries on the surface of mature follicles during progression but prior to rupture.

(A, D, B, E) Erythrocytes within the surface capillaries of mature follicles (IVb) from wildtype ($pgr^{+/+}$) female zebrafish collected at 6:40 or 7:00 were visualized using *o*-dianisidine staining as follicles progressing towards rupture and ovulation. (C, F) Erythrocytes were also visualized in similar mature follicles that were collected at 7:00 from $pgr^{-/-}$ females, in which ovulation never occurs (13). (G) At least 30 mature follicles were randomly chosen from at least 3 different females of each condition, and the number of erythrocytes in capillaries were counted by a blinded observer. Data are expressed as mean $\pm$ SEM ($N \geq 30$). P-values were calculated by two-tailed Student's t-test. Scale bars: 500 μ m (A-C) or 100 μ m (D-F).

Figure 8. Heparin inhibits ovulation *in vitro*.

Ovulation rate was determined using mature follicles (IVb) isolated from transgenic female zebrafish (*Tg(fli1:eGFP)*). The intact IVb follicles were easily recognizable by the interconnecting blood vessels with strong GFP expression (A, C). After ovulation, oocytes were easily recognizable by the absence of interconnecting GFP-positive vessels and expansion of the vitelline envelope (chorion) (B, D). Follicular layers separated from ovulated oocytes could also be easily distinguished due to interconnecting blood vessels and a dramatic size reduction (B, D). (A, C) Mature follicles (IVb) were collected from *Tg(fli1:eGFP)* female zebrafish. (B, D) Representative images of ovulated mature follicles were taken 2 hrs after incubation at 28°C. Arrows indicate the remaining follicular layer following ovulation. Arrowheads indicate expanded vitelline envelopes. Oc, oocyte. (E) mature follicles were incubated in a medium containing EDTA (2 mM), heparin (100 µg/mL), warfarin (50 µg/mL), or vehicle control, and rates of ovulation were determined 2 hrs following incubation. Data are expressed as the mean ± SEM (N ≥ 7). P-values were calculated by two-tailed Student's t-test. Scale bars: 1 mm (A, B) or 250 µm (C, D).

Figure 9. Reduced fecundity in $f5^{+/-}$ female zebrafish.

(A) Reduced numbers of live embryos per spawning for each $f5^{+/-}$ female in comparison to that of wildtype ($f5^{+/+}$) females. (B) Reduced total numbers of embryos for each $f5^{+/-}$ female fish in comparison to wildtype ($f5^{+/+}$) females. (C) Reduced spawning times in $f5^{+/-}$ females in comparison to that of wildtype ($f5^{+/+}$) females. (D) Increased spawning intervals in $f5^{+/-}$ females in comparison to wildtype ($f5^{+/+}$) females. Data are expressed as the mean $\pm$ SEM ($N \geq 14$). P-values were calculated by two-tailed Student's t-test.

Figure 10. A proposed model for nuclear progestin receptor (Pgr) mediated upregulation of *f5* and *f3a* in blood coagulation that facilitates ovulation in zebrafish.

Luteinizing hormone (LH) induces ovulation via luteinizing hormone receptor (Lhcgr) in preovulatory follicles, namely in stage IV follicles in zebrafish. LH signaling is mainly mediated by progestin (DHP, i.e., 17 α ,20 β -dihydroxy-progesterone, a native ovulation inducing steroid in zebrafish) and its receptor (Pgr) in the production and activation of metalloproteases. The illustration on the left side is showing the upregulation of metalloproteases in preovulatory follicles, to be more specific, during the transition from stage IVa to IVb follicles, which is an important preparation step for ovulation (16,26,27). However, these proteases are produced in a latent form, i.e., inactive form of metalloproteases (iMPs, see Yong Zhu, 2021 for detail), and inhibited by tissue inhibitors of metalloproteases (TIMP) presented at high levels in and delivered by blood capillaries to the follicles. The illustration on the right side is showing a dramatic increase of coagulation factors *f5* and *f3a* shortly before ovulation, specifically in the late stage of IVb follicles, which are also mainly upregulated by progestin via Pgr. These ovarian follicles derived, and locally produced coagulation factors (F5 and F3a) are imported into inside blood vessels messed in the follicles and activated via yet to be defined mechanisms. Once F5 and F3a are activated, these coagulation factors accelerate blood coagulation, which in turn stops blood flow and therefore effectively terminates TIMP inhibition of metalloproteases from the bloodstream. In addition, the production of fibrin (not

782 shown) due to the formation of blood clots stimulates activities of plasmin/plasminogen activators, which
783 are well-known inducers for metalloprotease activation. The combined force of these two pathways triggers
784 rapid activation of metalloproteases, breakdown of extracellular matrix and basal membrane, rupture of
785 follicular cell layers, the final release of a mature oocyte.