Title: Reactive oxygen species signalling is involved in alkamide-induced alterations in root development.

Running title: Alkamides and ROS signalling

Tonatiu Campos-García ^{a, b}, Jorge Molina-Torres ^b, and Kirk Overmyer ^{a, #}

^a Organismal and Evolutionary Biology Research Program, Faculty of Biological and Environmental Sciences, and Viikki Plant Science Centre. University of Helsinki, P.O. Box 65 (Viikinkaari 1), FI-00014 Helsinki, Finland.

^b Departamento de Biotecnología y Bioquímica, Unidad Irapuato, Cinvestav, Irapuato, Guanajuato, México

[#]Corresponding Author: Kirk Overmyer; University of Helsinki, P.O Box 65 (Viikinkaari 1), FI-00014 Helsinki, Finland; +(358) 44 337-7528; kirk.overmyer@helsinki.fi.

Date of submission: 23 December, 2021

Number of tables and figures: 7 (6 figures and 1 table)

Word count: Total: 5212 (Introduction, 1020; Results, 1560; Discussion, 2386; Conclusions, 174; Acknowledgements 72; Materials and Methods, 971).

Highlight

Reactive oxygen species (ROS) are involved in alkamide-induced altered root development. Heterotrimeric G-protein complex, extracellular acidification, and ROS sourced from peroxidases and NADPH-oxidases are involved in these processes.

Abstract

Alkamides are alpha unsaturated *N*-acylamides structurally related to *N*-acyl ethanolamides (NAEs) and *N*-acyl-L-homoserine lactones (AHLs). Studies have shown that alkamides induce prominent changes in root architecture, a significant metabolic readjustment, and transcriptional reprogramming. Some alkamide responses have been associated with redox signalling; however, this involvement and ROS sources have not been fully described. We utilized a genetic approach to address ROS signalling in alkamide-induced processes and found that in *Arabidopsis*, treatment with the alkamide affinin (50μM) increased the *in-situ* accumulation of H₂O₂ in lateral root emergence sites and reduced H₂O₂ accumulation in primary root meristems implying that altered root growth was dependent on endogenous H₂O₂. Results show that ROS sourced from PRX34, RBOHC and RBOHD were involved in promotion of lateral root emergence by alkamides. RBOHC was required for affinin-induced enhanced root hair expansion. Furthermore, affinin-induced changes in lateral root emergence, but not root hair length, were dependent on a change in extracellular pH. Finally, reverse genetic experiments suggest heterotrimeric G-proteins were involved in plant response to alkamides; nevertheless, further studies with additional higher order G-protein mutants will be required to resolve this question. These results support that alkamides recruit specific ROS signaling programs to mediate alterations in root architecture.

Keywords: affinin, alkamides, development, lateral roots, ROS, root, root growth, reverse genetics, pH

Introduction

In plants and animals some metabolites interact with messengers or receptors to modulate diverse metabolic pathways and signal transduction cascades. Plant chemical effectors from different sources can trigger physiological and morphological responses. Alkamides are low molecular weight α -unsaturated acyl-amide plant metabolites distributed in some plant species. These metabolites are known to have a wide range of biological activities in bacteria, fungi, plants, and mammals (Molina-Torres *et al.* 1996; Gertsch, 2008; Prachayasittukul *et al.* 2013). These include antimicrobial and other activities that suggest these compounds may function in defence against competing plants, microbes, and herbivorous pests (Molina-Torres *et al.* 2004). Affinin (*N*-isobutyl-2*E*,6*Z*,8*E*-decatrienamide) is the most abundant alkamide found in the ethanolic root extracts from chilcuague [*Heliopsis longipes* (A. Gray) S. F. Blake], an endemic species from “Sierra Gorda”, México. This molecule has been shown to have bacteriostatic and fungistatic effects (Molina-Torres *et al.* 1996) and to alter plant root growth and development (Ramírez-Chávez *et al.* 2004). More detailed plant developmental studies have found that alkamides also alter signalling in plants, modulating both developmental and stress response pathways, functioning as biochemical elicitors (López-Bucio *et al.* 2007; Méndez-Bravo *et al.* 2011). Most organisms contain amide lipids composed of one or two amines linked to a fatty acid through an amide bond in their inner and outer membranes. *N*-acylethanolamides (NAEs), which include some endocannabinoids, are an example of amine lipids structurally related to alkamides that have important biological signalling functions in plants (Blancaflor *et al.* 2003; Coulon *et al.* 2012; Blancaflor *et al.* 2014) and in mammals (Kunos *et al.* 2000; Wilson and Nicoll, 2002). Faure *et al.* (2014) found that alkamides could alter the NAE metabolic pathway by modulating fatty acid amide hydrolase (FAAH) activity. However, they also concluded that metabolic pathways other than FAAH are also involved in the metabolism of alkamides in plants, indicating the need for further studies in this area. Alkamides also exhibit structural similarities to *N*-acyl-homoserine lactones (AHLs), another amino lipid compound used by many bacteria as quorum-sensing signals to coordinate their collective behaviour. Accumulating evidence indicates that AHL are also perceived by plants, which respond by altering cell immune responses, host defence, stress responses, energy and metabolic

activities, transcriptional regulation, protein processing, cytoskeletal activities, root development and plant hormone responses (Mathesius et al. 2003; Kravchenko et al. 2006).

Affinin isolated from *Heliopsis longipes* roots displays a dramatic effect on *Arabidopsis thaliana* (*Arabidopsis*) root system architecture by altering primary root growth, lateral root emergence, and increasing root hair elongation in a dose dependent manner (Ramírez-Chávez et al. 2004). Méndez-Bravo et al. (2010) found that *N*-isobutyl decanamide and the interacting signals JA and nitric oxide (NO) act downstream independently of auxin-responsive gene expression to promote lateral root formation and emergence, providing compelling evidence that NO is an intermediate in alkamide signalling mediating the root system architecture alterations observed in *Arabidopsis*. Transcriptomic profiling of *Arabidopsis* has shown that exogenous application of *N*-isobutyl decanamide triggers profound physiological changes with activation of developmental, defence, and stress related genes (Mendez-Bravo et al. 2011). Transcripts of several JA-related genes such as *PDFs*, *VSP2*, *JAZ10* and *JAZ8* accumulated to higher levels. Transcripts of at least 70 genes belonging to the functional group “oxygen and radical detoxification”, also exhibited enhanced accumulation by *N*-isobutyl decanamide treatment. They found that alkamides could modulate some - defence responses associated with necrotrophic pathogens through JA-dependent and MPK6-regulated signalling pathways. Additionally, decanamide induces ROS accumulation in leaf tissue (Mendez-Bravo et al. 2011). These results suggest that general defence-associated responses elicited by *N*-isobutyl decanamide and affinin appear to be related to both hormone and ROS signalling pathways.

Recent studies revealed that ROS act as essential signalling molecules in plants and are required for several basic biological processes including cellular proliferation and differentiation, immunity, and stress responses (Mittler, 2017). ROS signalling specificity is dependent on the species produced and their subcellular location. It is well known that ROS, such as superoxide anion ($O_2^{\cdot-}$), hydrogen peroxide (H_2O_2), hydroxyl radical ($\cdot OH$), and singlet oxygen (1O_2), play important signalling roles in plants as key regulators of growth, development, response to biotic and environmental stimuli, plant metabolism, and programmed cell death (Jaspers and Kangasjärvi, 2010; Suzuki et al., 2011; del Río, 2015; Overmyer et al., 2003). In plants, H_2O_2 is the product of catalytic reactions from different

enzymes such as the peroxisomal flavin-containing enzymes glycolate oxidase and the acyl-CoA oxidase, which are involved in the photorespiratory and fatty acid β -oxidation pathways, respectively. H_2O_2 is produced in the chloroplast under normal and stress conditions, as well as from the activities of peroxidases, and in some species, oxalate oxidase (Torres, 2010; Baxter et al., 2014). It is now well established that a major source of the $O_2^{\cdot-}$ is the plasma membrane-localized NADPH-oxidases (del Río, 2015). $O_2^{\cdot-}$ is a charged molecule under most physiological conditions and can not passively transfer across a membrane. However, $O_2^{\cdot-}$ can dismutate into H_2O_2 , either enzymatically by superoxide dismutase (SOD), peroxidases or spontaneously, especially at low pH. H_2O_2 can readily cross membranes passively or via aquaporins. Additionally, $O_2^{\cdot-}$ can mediate the formation of membrane soluble lipid peroxides (Miller *et al.*, 2010; Mittler *et al.*, 2011). In many biological systems (as animals and plants) ROS signalling is mediated by a highly regulated process of ROS accumulation in specific cellular compartments. The NADPH-oxidases, termed *NOX* in animals and RESPIRATORY BURST OXIDASE HOMOLOGs (RBOHs) in plants, are plasma membrane localized enzymes that produce ROS into the apoplast. This family of enzymes are highly regulated via calcium and various phosphorylation/dephosphorylation events (Ogasawara *et al.* 2008; Mittler, 2017). In *Arabidopsis* the ROS produced by the NADPH-oxidases RBOHC, RBOHD, and RBOHF, as well as the class III peroxidases, have been shown to be involved in the lateral root emergence and root hair growth (Ogasawara *et al.* 2008; Li *et al.* 2015; Orman-Ligeza *et al.* 2016; Manzano *et al.* 2014, Fernández-Marcos *et al.* 2017). Nevertheless, the source of alkamide-induced ROS production and the signalling pathways recruited downstream that lead to modifications in root growth, development, and plant metabolism remain unknown. Here we utilize a genetic approach to address the role of ROS signalling in alkamide-induced processes.

Materials and Methods

Plant material and growth conditions.

Arabidopsis thaliana (*Arabidopsis*) accession Columbia-0 (Col-0) was used for all experiments unless otherwise indicated. For reverse genetic experiments, the following mutants were used; the NADPH-oxidases, *respiratory burst oxidase homologC* (*rbohC*),

rbohD, and *rbohF*; class III peroxidase, *peroxidase34* (*prx34*); heterotrimeric G-protein subunits, *g-protein alpha subunit1-2* (*gpa1-2*), *arabidopsis gtp-binding protein beta1-2* (*agb1-2*), *arabidopsis g-protein gamma subunit2-1* (*agg2-1*), and the *regulator of g-protein signaling1-2* (*rgs1-2*). Mutant seeds were obtained from NASC (www.arabidopsis.info). Seeds were surface sterilized with 70% (v/v) ethanol plus 2% triton-X 100 for 5 min, then rinsed in absolute ethanol for 1 min, and dried in a laminar hood over sterile filter paper sheets. Seeds were germinated and grown on agar plates (9 g L⁻¹) containing 0.3x MS medium and sucrose (11 g L⁻¹). Seeds were stratified 72 hrs in darkness at 4°C and then placed in controlled environment growth chambers (Weiss FITOTRON SCG120; www.fitotron.co.uk) vertically to allow root growth along the agar surface and unimpeded hypocotyl growth. Plants were grown under a long-day photoperiod (16 h light, 8 h darkness), 25°C/18°C day/night temperature and a light intensity of 60-100 $\mu\text{M m}^{-2} \text{s}^{-1}$. After germination for four days, seedlings were transferred to plates containing solvent control or affinin treatments (10 or 50 μM). For the pharmacological assay, one-week old seedlings were transferred to a solid medium supplemented individually or in various combinations of affinin (7 or 50 μM), diphenyleneiododinium (DPI; 0.3 μM), and solvent control for seven days. In MES buffer experiments, four-day-old seedlings were transferred to medium supplemented with MES buffer (0.5 g L⁻¹); after the addition of all the reagents the pH was adjusted to 5.7 with NaOH and then autoclaved. After autoclaving, treatments were prepared adding the affinin (10 or 50 μM) to the medium. After seven days the primary root length, number of emerged lateral roots and root hair length were evaluated.

Affinin isolation and purification.

Affinin extraction was performed as previously reported by Ramírez-Chavez *et al.* (2004). After extraction, the concentrated oil was weighted, 2 g were purified by column chromatography as previously reported by Ramírez-Chávez *et al.* (2004); subsequently, the purified fraction was analysed by GC/EIMS to confirm its purity and concentration.

ROS staining.

H₂O₂ accumulation was monitored *in situ* with 3,3'-diaminobenzidine tetrachloride (DAB; Sigma-Aldrich; www.sigmaaldrich.com) based on the method of (Thordal-Christensen *et al.*,

1997). Six-day-old seedlings were immersed in DAB (1 mg ml⁻¹) in PBS buffer (10 mM; pH 7.0) plus 0.05% (v/v) tween-20 and placed overnight at room temperature in the dark. O₂^{•-} was detected based on nitroblue tetrazolium (NBT)-reducing activity. Seedlings were covered in an NBT solution (0.5mg ml⁻¹) in 0.1M PBS buffer (pH 7.4) plus 0.05% triton-X 100 and incubated in the light for 10 minutes. DAB and NBT reactions were stopped by replacing staining solution with fixative / clearing solution of ethanol / acetic acid / glycerol (3:1:1) for 24 hours, mounted on glass slides with a coverslip and visualized with a Leica 2500 microscope (www.leica-microsystems.com).

Root growth analysis and microscopy.

Root growth measurements were performed on images taken from the plates using the SmartRoot plugin on ImageJ (www.imagej.net; <https://smartroot.github.io/>). For the analysis of emerged lateral roots, root samples were visualized with a Leica MZ10F stereo microscope equipped with a model DFC490 digital camera attachment (Leica; www.leica-microsystems.com) and lateral roots protruding beyond the epidermal tissue were scored as emerged. Primary root zone organisation of seedlings with or without 50µM affinin treatment were visualised with 5 mg/ml propidium iodide stain and observed in a Carl Zeiss LSM800 confocal laser microscope (<https://www.zeiss.com/>). Measurements of distance from primary root tip to first root hair bulge, cell length from the elongation/differentiation zone (EZ/DZ) and apical meristem length (AM) were made with ImageJ software (www.imagej.net). DAB staining images were taken with a Leica 2500 microscope with differential interference contrast (DIC) optics (<https://www.leica-microsystems.com/>). For each treatment, at least ten plants were analysed. Representative plants for each treatment were photographed using the Nomarski optics on a Leica 2500 microscope. Quantification of DAB staining intensity was done by ImageJ-based quantification described by Béziat (2017). Prior to quantification, the colour mode of light micrographs was converted from RGB to HSB by using ImageJ software and the saturation channel in the HSB stack was selected. Then the intensity of DAB staining was measured in the saturation channel from a same size area in all pictures. An increase of saturation depicts more “pure” colour, while a decrease denotes a more “washed-out” signal (Béziat *et al.*, 2017).

Bromocresol purple pH assay.

To test pH changes in the roots, seven-day-old *Arabidopsis* seedlings Col-0 were transferred to a solid medium (as described before) with different affinin treatments. At 24 hours, seedlings were transferred from treatments to a glass slides covered with 1ml of the pH indicator bromocresol purple (www.sigmaaldrich.com; 0.04 g L⁻¹) in agarose (0.7%) prepared with distilled water plus CaSO₄ (0.2mM) (Zandonadi *et al.*, 2010). The pH was adjusted to 7.5 with NaOH. After 30 min, glass slides with the seedlings were photographed with a CAMAG TLC Visualizer (www.camag.com). Photographs were analysed with ImageJ software as follows. First, images were converted to an 8-bit format and calibrated with a set of density standards (pH scale bar), a region of interest (ROI) was selected from each pH scale point and the mean grey value of each of those points was recorded and then fitted to a “curve fitting method (linear function)” from the popup menu as described in the ImageJ user manual (<https://imagej.nih.gov>).

Statistical analysis.

Data from mutant sets were analysed with two-way ANOVA ($n = 10$) and a Tukey’s posthoc test using the Minitab 14 software (www.minitab.com). Different letters are used to indicate significance groups (means that differed significantly $P \leq 0.05$). For single genotype experiments, data were analysed with one-way ANOVA ($n = 10$) and Tukey’s posthoc test ($P \leq 0.05$) using InfoStat software (www.infostat.com.ar).

Results

Root phenotype of *Arabidopsis Col-0* in response to affinin.

To establish the principal affinin-induced phenotypes in seedlings of WT Col-0 *Arabidopsis* under our conditions, root architecture was monitored in response to two affinin concentrations (10 μ M and 50 μ M), in aseptic cultures on 0.3x MS solid medium. After seven days of growth, the 10 μ M treatment resulted in an increased number of emerged lateral roots without a significant decrease on the primary root length (Fig. 1A-C). The 50 μ M treatment

reduced growth monitored as primary root length and resulted in a significantly increased number of emerged lateral roots and root hair length (Fig. 1A-D). These results show that affinin treatments influenced *Arabidopsis* root system architecture, consistent with the previously reported phenotypes induced by affinin and decanamide (Ramírez-Chávez *et al.*, 2004; López-Bucio *et al.*, 2007; Méndez-Bravo *et al.*, 2010; Méndez-Bravo *et al.*, 2011).

Affinin alters H₂O₂ accumulation in root meristems and lateral root emergence sites.

Arabidopsis transcriptional signatures (Méndez-Bravo *et al.* 2011) suggest that *N*-isobutyl decanamide modulates ROS signalling, which is involved in root development (Passardi *et al.* 2006; Manzano *et al.* 2014), prompting us to assay for affinin-induced ROS responses in *Arabidopsis* roots. DAB staining was utilized to visualize affinin-induced changes in H₂O₂ accumulation *in situ*. Staining with DAB forms a brown-reddish precipitate in the presence of H₂O₂ and peroxidase activity. In the root meristematic zone, the deposition of DAB stain was reduced by treatment with 50 µM affinin, but no change in DAB staining was seen in the elongation or maturation zones (Fig. 2A). O₂^{•-} accumulation detected by NBT staining remained unchanged (Fig. 2B). DAB staining was significantly higher in the lateral root primordia emergence sites of roots treated with 50 µM affinin but reduced in the apical meristem (Fig. 2A-F). These results suggest that H₂O₂ is involved in affinin-induced changes in lateral root emergence and root apical meristem development leading to an increase in emerged lateral roots and a reduction of primary root length in response to the affinin treatments. These results demonstrate a correlation between sites of H₂O₂ accumulation and developmental responses to affinin, prompting further exploration of the role of ROS in these processes.

Affinin-induced *Arabidopsis* lateral rooting is dependent of endogenous H₂O₂.

To test ROS involvement in affinin-induced changes in root architecture, a pharmacological approach was first employed. The effects of affinin on root development were tested in the presence of diphenyleneiododum (DPI), a NADPH oxidase inhibitor. Plants grown in medium containing DPI plus affinin did not show the characteristic affinin induced phenotypes (Fig. 2E). Specifically, seedlings grown on DPI did not increase the number of lateral roots in the presence of affinin; the number of lateral roots in these seedlings remained

the same as it was prior to being transferred. Affinin at 7 μ M increased the primary root length a response reported by Ramírez-Chávez *et al.* (2004), in contrast primary root growth was totally inhibited by DPI (Fig. 2H-I). These results suggest that endogenous ROS is required for affinin-induced lateral root growth and for primary root development, prompting us to use reverse genetics to define the sources of ROS that are involved in mediating these root architecture changes.

PRX34-mediated H₂O₂ production in affinin-induced responses.

The cell wall associated class III peroxidases (PRXs) have been implicated in lateral root formation and regulation of root tip growth (Passardi *et al.* 2006; Manzano *et al.* 2014). The *prx34* mutant (Bindschedler *et al.*, 2006) was assayed for affinin-induced root structure alterations and associated DAB stain accumulation. The *prx34* mutant exhibited enhanced inhibition of primary root growth in response to high affinin concentration (Fig. 3A), which was associated with a drop in DAB staining intensity in the meristematic zone (Fig. 3C, D), suggesting that modulation of ROS signalling by PRX34 is implicated in the regulation of meristematic processes by affinin. The number of emerged lateral roots in *prx34* was not increased by affinin treatment (Fig. 3B), while the enhanced DAB staining seen in affinin treated wild type Col-0 plants was not observed in the lateral root emergence sites of *prx34* roots (Fig. 3C, D). These results support that PRX34 was required for alkamide-induced signalling leading to lateral root emergence and suggest that the ROS produced or modulated by PRX34 is involved in this process.

NADPH-oxidase mediated ROS production in affinin-induced developmental changes.

We next sought to further define the sources of affinin-induced ROS. To explore the contribution of ROS signalling from the RBOH NADPH oxidases in responses to affinin, the root system architecture of wild type plants and the *rbohC*, *rbohD*, and *rbohF* single mutants were compared following growth in the presence of different affinin concentrations. In medium lacking affinin, all *rboh* mutants exhibited reduced primary root growth as compared to wild type (Fig. 4A) and reduced DAB stain deposition was apparent in the root tips of *rbohC* and *rbohD*, but not *rbohF* under control conditions (Fig. 4C). Primary roots treated with affinin in all mutants showed behaviour similar to WT, however, the *rbohC*, *rbohD*, and

rbohF mutants had enhanced growth inhibition at 50µM affinin (Fig. 4A). DAB staining of the apical meristems (Fig. 4C) showed that affinin at 50 µM reduced the H₂O₂ accumulation in WT, *rbohC* and *rbohD* but no decrease was observed in *rbohF* (Fig. 4C,E). These results suggest that primary root growth inhibition caused by affinin could be mediated via attenuation of ROS accumulation sourced from RBOHC and RBOHD, but not RBOHF. The *rbohC* and *rbohD* mutants lacked a significant affinin-induced increase in emerged lateral root number (Fig. 4B). DAB staining at lateral roots emergence sites was not enhanced in both mutants (*rbohC* and *rbohD*) treated with 50 µM affinin while in *rbohF* staining was higher in response to 50 µM affinin (Fig. 4D, F) suggesting that RBOHC and RBOHD could be involved in promotion of lateral root emergence via alkamide-induced ROS accumulation.

ROS produced by RBOHC is required for cell expansion in root hair growth and characteristically the *rbohC* mutant has greatly reduced root hair length (Foreman *et al.* 2003). Intriguingly, root hair length was unchanged upon affinin-treatment of the *rbohC* mutant indicating that ROS produced by this NADPH-oxidase is necessary for affinin-induced enhanced root hair expansion (Fig. 4G, H). The *rbohD* and *rbohF* mutants had significantly longer root hairs under affinin-treatment (Fig. 4G, H) suggesting that these ROS sources had a minor role as negative regulators of this process. These results highlight the importance of RBOHC-mediated ROS production for enhanced root hair expansion.

G-protein signalling in the *Arabidopsis* affinin response.

To test if the heterotrimeric (ht) G-protein subunits, G_α, G_β and G_γ, are involved in *Arabidopsis* response to affinin, affinin-induced root phenotypes of *gpa1-2*, *agb1-2*, and *agg2-1* mutant plants were examined. In all genotypes primary root length was reduced by 50 µM affinin treatments and in *agb1-2* this effect was slightly but significantly enhanced suggesting AGB2 may act as a negative regulator of this process (Fig. 5A). The *agb1-2* and *agg2-1* mutants exhibited and increased number of lateral roots under control conditions, indicating that they negatively regulate lateral root formation (Fig. 5B). Interestingly, *agb1-2* exhibited a decrease in the number of lateral roots under affinin treatment compared to control (0µM affinin), while *agg2-2* remained equal in affinin treatment and control, and *gpa1-2* had a slight but not significant increase, suggesting that all these subunits could be

required for enhanced lateral root emergence in response to affinin (Fig.5B). In plants G_{α} proteins tend toward their active GTP bound state (Urano and Jones, 2014) and the seven transmembrane domains (7TMD) containing RGS1 protein negatively regulates GPA1-mediated signalling through its GTPase accelerating protein (GAP) activity (Liang *et al.* 2018). The *rgs1-2* mutant was also tested and had a reduced primary root length under control conditions (Fig. 5C) indicating RGS2 is required for full primary root growth. Affinin treatment reduced primary root length to a similar extent in both Col-0 WT and *rgs1-2* (Fig. 5C) suggesting RGS1 is not involved in affinin-induced response on primary root growth. Lateral root emergence was the same in WT and *rgs1-2* under control conditions, but the affinin-induced increase was enhanced in *rgs1-2*, suggesting RGS1 negatively regulates this response (Fig. 5D). Together, these data implicate multiple htG-protein subunits in the regulation of the affinin response.

Extracellular pH and affinin-induced responses.

To test if affinin causes changes in apoplastic pH, an affinin pulse for 24 hours (see Materials and Methods) was applied to seven-day-old *Arabidopsis* roots and changes in the pH in surrounding growth medium was monitored with the indicator bromocresol purple. The results show that affinin treatment induced extracellular acidification in the root elongation zone (Fig. 6A, B). To test the potential functional consequence of these extracellular pH changes, the extracellular pH was stabilized using a medium containing 2-(*N*-morpholino) ethanesulfonic acid (MES) buffer (pH 5.7), which has an active buffering capacity in the relevant range of pH 5.5 - 6.7. In buffered medium, affinin treatment exhibited typical inhibition of primary root growth; however, the number of emerged lateral roots did not increase in response to affinin (Fig. 6C-E). Further, in line with previous studies (Kagenishi *et al.* 2016), root hair growth was inhibited in control plants on buffered medium without affinin; however, root hair length still increased upon affinin treatment on MES buffered medium (Fig. 6F, G). We conclude that the affinin-induced changes in lateral root emergence, but not root hair length, were dependent on a change in extracellular pH.

Discussion

The role of ROS in alkamide-induced processes.

Decanamide-induced transcriptional reprogramming previously observed in *Arabidopsis* (Mendez-Bravo et al. 2011) suggests ROS signalling may mediate alkamide-induced processes. The current study utilizes a genetic approach with the previously established model system (Ramírez-Chávez et al. 2004; Méndez-Bravo et al. 2010) of alkamide-induced alterations to *Arabidopsis* root system architecture to demonstrate that these developmental changes are associated with changes in ROS accumulation, as monitored using NBT to visualize $O_2^{\bullet -}$ production and DAB for H_2O_2 (Figs. 2-4). Additionally, this study builds upon the previous observation of affinin-induced H_2O_2 accumulation in *Arabidopsis* leaves (Méndez-Bravo et al. 2011) by expanding findings to root tissues undergoing developmental changes. We found that 50 μ M affinin induced the accumulation of H_2O_2 in the peripheral cells surrounding the lateral root primordia (Fig. 2C, D), which coincided with an increased number of emerged lateral roots (Fig. 1A, B). These data are consistent with the known signalling role of H_2O_2 in lateral root development as previously reported (Passardi et al. 2006; Manzano et al. 2014; Tsukagoshi et al. 2016; Orman-Ligeza et al. 2016) and suggest that H_2O_2 is involved in root response to alkamides by acting as a signalling intermediate. Further, these results are consistent with the ability of ROS to increase lateral root numbers due to the activation of lateral root pre-branch sites and lateral root primordia. Specifically, ROS are deposited in the apoplast of these cells during lateral root emergence (Orman-Ligeza et al. 2016).

Moreover, our results indicate that affinin attenuated growth of the primary root in a dose dependent manner (Fig. 1A, C), which was accompanied by attenuated DAB staining intensity specifically in the root meristematic region (Fig. 2A, C, D). Root growth is controlled by both the rate of cell division in the meristematic zone and the degree of cell expansion in the elongation zone (Beemster and Baskin, 1998; Tsukagoshi et al. 2010). Increased root meristem size is correlated with an acceleration of root growth and is a result of increased rates of cell division and the delay of cell expansion (Ubeda-Tomas et al. 2009; Tsukagoshi et al. 2010). It has been demonstrated that H_2O_2 and the $O_2^{\bullet -}$ both have distinct accumulation zones and distinct roles in the growing *Arabidopsis* root (Dunand et al. 2007). Our results demonstrate that affinin altered the accumulation of H_2O_2 , but not $O_2^{\bullet -}$, in the meristem (Fig. 2A, B). Affinin treatment reduced the cell size of the root elongation-differentiation zone (Fig. 1E-M) without altering H_2O_2 and $O_2^{\bullet -}$ accumulation in this root

zone (Fig. 2A, B) indicating that affinin can modulate a defined developmental program. This profile of changes is consistent with the action of peroxidases, which are known to modulate primary root growth (Passardi *et al.* 2006). These results agree with the findings that affinin alter meristematic root growth and the expression of *CycB1* gene (Ramírez-Chávez *et al.*, 2004).

We also present evidence supporting the involvement of extracellular acidification in response to affinin. Ion fluxes are a common event that often works in concert with ROS signalling in stress and developmental responses. It is known that changes in apoplastic pH are likely to modulate the activity of several regulatory elements such as cell wall proteins, such as expansins (Cosgrove *et al.* 2002) and pectin methylesterases (Sherrier and Vandenbasch, 1994, Micheli, 2001); plasma membrane proteins, such as pH-sensitive potassium channels (Ilan *et al.* 1996, Hartje *et al.* 2001); and is functionally related with the regulation of root hair growth (Monshausen *et al.* 2007).

Beyond correlating ROS accumulation with affinin-induced changes in root system architecture, pharmacological treatments and mutant plants with compromised ROS pathways (Figs. 3-5) were utilized to reveal a requirement for these ROS-signalling pathways for a subset of affinin-induced developmental responses. Based on this pharmacological and genetic evidence, we conclude that ROS-signalling was involved in affinin-induced responses. This work also defined the sources of various affinin-responses, as further discussed below.

The production of ROS in the apoplast depends on several classes of enzymes, most notably, NADPH-oxidases and class III peroxidases. Several NADPH-oxidases are known ROS sources that mediate cell expansion and determines root system architecture (Torres, 2010). These are known in *Arabidopsis* as RESPIRATORY BURST OXIDASE HOMOLOGS (RBOHs) and act specifically in lateral root emergence, and root hair cell expansion (Foreman *et al.* 2003; Orman-Ligeza *et al.* 2016), while class III peroxidases, such as *AtPrx34*, have been associated with an active role in root cell elongation (Sagi and Fluhr, 2006; Passardi *et al.*, 2006).

To test if these ROS sources are involved in affinin-response, several available *Arabidopsis* mutants were utilized, including *prx34*, which is defective in a class III peroxidase, and the NADPH-oxidase mutants, *rbohC*, *rbohD*, and *rbohF*. Testing the effect of affinin in the *prx34* mutant, the ROS staining pattern of lateral root primordia and the number of emerged lateral roots were indistinguishable from wild type Col-0 plants (Fig. 3C, D). It has been demonstrated that *PRX* activities are important for lateral root development, especially during lateral root emergence (Manzano *et al.*, 2014). The effects of affinin on root system architecture has been previously shown to be independent or downstream of auxin signalling (Méndez-Bravo *et al.*, 2010). Accordingly, lateral root emergence due to *PRX* activity also occurs independent of auxin signalling (Manzano *et al.* 2014). The accumulation of DAB staining in root tips was reduced by affinin 50 μ M in *prx34* as in WT. It has been reported that peroxidases are involved in the regulation of primary root growth (Tsukagoshi *et al.* 2010) and *PRX34* was previously shown to be required as a component of the oxidative burst in response to some pathogens (Bindschedler *et al.*, 2006). Our results strongly suggest that *PRX34* is not required for affinin-induced inhibition of primary root growth, suggesting other *PRXs* may regulate this process. The reduced accumulation of DAB staining in root meristems upon 50 μ M affinin treatment suggests that ROS homeostasis is somehow disrupted due to affinin treatment leading to an alteration in plant cell cycle. This result is supported by the fact that affinin at high concentrations reduced the expression of genes associated with the mitotic cycle, such as *CycB1* (Ramírez-Chávez *et al.*, 2004). On this topic, it has been found that cellular ROS signalling oscillations are rapidly transmitted through MAPK pathways inducing MAP activation and affects microtubules dynamics and organization (Livanos *et al.*, 2012).

Using the *respiratory burst oxidase homologues (RBOH)* mutants (*rbohC*, *rbohD* and *rbohF*) these ROS signalling genes were tested for involvement in alkamide induced signalling. This revealed that primary root growth inhibition by affinin was independent of all the loci represented by these mutant lines (Fig.4A), however H_2O_2 accumulation as monitored by DAB staining in the apical meristem was reduced in the *rbohC* and *rbohD* mutants while unchanged in *rbohF* (Fig. 4C, E). Interestingly the *rbohD* mutant exhibited a significant inhibition of primary root length in response to affinin compared to the other two mutants and wild type. Taken together, the primary root length and DAB staining results do not

support that the ROS produced by *rbohC* and *rbohF* are involved in affinin-induced response in meristematic activity. However, the impact of affinin on root length in *rbohD* supports the hypothesis that ROS homeostasis could be involved in mediating affinin response in meristematic cells. On the other hand, *rbohC* and *rbohD* failed to increase the number of emerged lateral roots in response to affinin (Fig. 4B) and the DAB staining intensity did not show differences in lateral root primordia between the treatments of these mutants (Fig. 4D, F). These results are consistent with Orman-Ligeza *et al.* (2016), who found that *RBOH*-mediated ROS production facilitates lateral root outgrowth by promoting cell wall remodelling of overlying parental tissues. Indeed, the diverse transcription patterns suggest that *RBOHs* function in broad aspects of growth and physiological response (Sagi and Fluhr, 2006). In contrast, *rbohF* response in terms of emerged lateral roots was similar to the wild type of response induced by affinin and the DAB staining showed similar results (Fig. 4D, F).

The effect of affinin on root hair growth has been demonstrated (Ramírez-Chávez, *et al.* 2004) and corroborated (Fig. 4G). Interestingly, *rbohC* was the only mutant that did not respond to affinin-induced enhanced root hair elongation (Fig. 4G), this result not only demonstrates the importance of ROS in affinin induced signalling but also the specificity of ROS produced by *RBOHC* on root hair growth. The *rbohD* and *rbohF* mutants exhibited an increase in root hair length higher than the wild type seedlings (Fig.4G), which indicates that these ROS sources could be negative regulators of root hair growth.

In ROS metabolomic studies, the “oxidative stress” signature includes accumulation of several compounds implicated in ascorbate and glutathione synthesis and degradation pathways, and phytohormones such salicylic acid and jasmonic acid (Noctor *et al.*, 2016). Additionally, this signature also includes several compounds whose connections to antioxidant metabolism and redox homeostasis are not as obvious. For example, the accumulation of branched chain amino acids induced by ABA treatment, a response that seems to occur reproducibly during redox signalling (Ghassemian *et al.*, 2008; Noctor *et al.*, 2015). Recently in tomato seedlings it was found that, affinin induced a dose-dependent metabolomic reprogramming that lead to enhanced accumulation of amino acids, organic acids, sugar alcohols, phenolics, and fatty acids (Campos-García and Molina-Torres, 2021),

all metabolites that have been associated with marker metabolites in the oxidative stress response.

The Alkamide-induced developmental program.

The details concerning how alkamides affect plant growth remain poorly characterized. Exogenous application of affinin to plants has multiple effects on several plant processes, some of which are similar to responses triggered by some well-known stress related signals. Comparison of affinin-response to these other better characterized signalling pathways may offer insight into affinin-induced signalling.

Biotic and abiotic stresses induce the so-called stress-induced morphogenic response (SIMR), which shares some similarities with the alkamide response. NO is an intermediate in both SIMR (Potters et al., 2009) and alkamide signalling mediating alteration in root system architecture in *Arabidopsis* (Méndez-Bravo *et al.* 2010). Altered root branching, inhibition of cell elongation, as well as changes in apoplastic pH, ROS, and redox signalling, are all common to alkamide-response and SIMR (Potters et al., 2007, Potters et al., 2009; Tongnetti et al., 2012) It is possible that alkamide-induced stress results in SIMR. However, auxin features prominently in the regulation of SIMR and it was demonstrated that alkamides act via an auxin-independent signalling pathway (Ramírez-Chávez *et al.* 2004). Nonetheless, the possibility that some alkamide-induced responses are related to SIMR may be worth further consideration.

Although auxins are considered the major plant growth-regulating hormones underlying root system architecture adjustment, the discovery of novel signal molecules such as *N*-acyl amides, *N*-acyl ethanolamides (NAE) and *N*-acyl homoserine lactones (AHLs) has shed light on the intricate signalling networks that trigger root system architecture modifications and physiological responses (Ramírez-Chávez *et al.*, 2004; Méndez-Bravo *et al.*, 2010; Coulon *et al.*, 2012; Schikora *et al.*, 2016). It has been demonstrated that NAEs and AHLs, compounds that are structurally related to alkamides, have a wide range of effects in plant development (Table 1), some of which overlap with affinin-induced responses. Under *in vitro* conditions, affinin shows a trend towards increasing FAAH's capacity to metabolize NAE 12:0, suggesting that affinin may have some effects on plants acting via this modulation of

NAE metabolism. The structural similarity of affinin to NAEs, suggests that affinin might directly influence FAAH activity in plants, but this will also require further investigation (Faure *et al.*, 2015). Recently, Aziz and Chapman (2020) proposed the hypothesis that FAAH proteins hydrolyse a broader range of lipophilic substrates than previously recognized, including affinin, and consequently play a pivotal role in *N*-acyl amide-mediated plant–microbe interactions, a function beyond the established role for FAAH in seedling development in *Arabidopsis*.

In mammals, polyunsaturated NAEs bind to specific 7TMD G-protein coupled receptors (GPCRs) and activate the htG-proteins that regulate multiple downstream signalling pathways (Abadji *et al.*, 1999; Bosier *et al.*, 2010). In plants, htG-protein signalling has been implicated in many processes such as plant growth and development. Plant htG-protein signalling is fundamentally different from mammalian systems; with no canonical 7TMD GPCRs, the presence of novel EXTRA-LARGE G-PROTEINs (XLG1, XLG2, and XLG3), and a single G α subunit that has a low intrinsic GTPase activity and is self-activating in the absence of the GEF activity of GPCRs, thus tends toward a constitutively activated state (Urano *et al.*, 2016). Plant G-protein signalling has been implicated in AHL-signalling. GPA1 was required for AHL-mediated elongation of *Arabidopsis* roots (Liu *et al.*, 2012). However, we found that inhibition of primary root length by affinin was independent of GPA1.

In *Arabidopsis* NAE 18:3 induces stress responses, autophagy, senescence, chlorophyll catabolic genes, represses chlorophyll biosynthesis genes, and these responses require an intact htG protein complex (Yan *et al.*, 2020). Specifically, this study demonstrated the NAE 18:3 response required AGB1, XLGs (using the *xlgl xlg2 xlg3* triple mutant) and AGGs (using the *agg1 agg2 agg3* triple mutant; Yan *et al.*, 2020). The NAE 18:3 response was enhanced in the *gpa1* and *rgs1* mutants. This work also demonstrates that, due to functional redundancy in these gene families, higher order mutants are required to see clear results. Our results also showed complex profile of phenotypes with htG-protein mutants: GPA1, AGG1, and AGB1 were required for, and *rgs1* conferred hypersensitivity to affinin-induced increased lateral root emergence. For affinin-induced primary root elongation, GPA1, AGG1, and RGS1 were not required, but the *agbl* mutant conferred a slight hypersensitivity. Taken

together, these results suggest the similarities between affinin and NAEs warrant further exploration. Especially, further studies with additional higher order G-protein mutants will be required to resolve this question.

Interestingly, the *agbl-2* and *agg2-1* mutants exhibit an increased number of lateral roots under control conditions, and it is known that $G_{\beta\gamma}$ -dimer restrains lateral root formation (Chen *et al.*, 2006). Moreover, it is reported that the *agbl* mutant has a more expanded root architecture presumably due to an increased cell proliferation and lateral root formation (Urano *et al.*, 2016). Our results show that in the *agbl-2* mutant, primary root length and lateral root number were reduced in response to affinin, which is a contrasting response compared to the mutant phenotype. Taken together these results suggest the htG-protein complex mediating plant response to affinin, modulating cell proliferation activity and lateral root emergence. Further studies will be required to clarify the mechanism involved.

Conclusions

Our results provide clear evidence that ROS are molecular intermediates involved in lateral roots emergence and root hair growth in response to alkamides. This is supported by the known roles of ROS as mediators and activators of structural changes in roots. In addition, we demonstrate that alkamides induced root extracellular pH changes, which had an effect on lateral roots emergence. These results provide us with evidence that alkamide-induced modification of root architecture depends on modifications in extracellular pH and ROS homeostasis in the lateral root emergence zone. Further, we also provided evidence that heterotrimeric G proteins can be mediating affinin-induced signalling. This is supported by evidence that some G protein subunits may be activating ROS synthesis, inducing softening of cell walls and facilitating lateral roots emergence. While, on the other hand, ROS formed specifically by RBOHC induce Ca^{+2} channels hyperpolarization, allowing Ca entry into the cell and activating root hairs elongation. Nevertheless, further studies with additional higher order G-protein mutants will be required to resolve how htG-proteins are involved in alkamide-signaling.

Acknowledgements

We thank Tuomas Puukko, Airi Lamminmäki, and Leena Grönholm, for excellent technical support, Eveliina Karjalainen for assistance with mutant seed production, Dr. Huitzimengari Campos-García for help with the statistical analysis, Dr. Vincent-Cervantes Bueno for his help with the confocal microscope, and M.Sc. Enrique Ramírez-Chávez for his support in affinin purification and GC-MS analysis. The members of the Plant Stress Metagroup are acknowledged for the helpful comments and discussions during this project.

Author Contributions

Project conception and planning, TCG, KO, JM-T; experiment design, TCG, KO; experimental work, TCG; writing the manuscript, TCG, KO; editing and approval of manuscript, TCG, KO, JM-T.

Conflicts of Interest

The authors have no conflict of interest to declare. All co-authors have seen and agree with the contents of the manuscript and there is no financial interest to report. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the result.

Funding

This work was supported by the following grants: the Finnish National Agency for Education, Finnish Government Scholarship Pool (decision no. KM-18-10772) and Consejo Nacional de Ciencia y Tecnología (CONACYT) grant (426142) to TCG and the Academy of Finland Center of Excellence in the Molecular Biology of Primary Producers 2014-2019 (decisions #271832 and 307335).

Data availability

569 The data that support the findings of this study are openly available at Campos-García *et al.*
570 (2021): Alkamides and ROS signalling. figshare. Dataset.
571 <https://doi.org/10.6084/m9.figshare.17303615.v1>.

572

References

- Abadji, V., Lucas-Lenard, J. M., Chin, C. N., Kendall, D. A.** 1999. Involvement of the carboxyl terminus of the third intracellular loop of the cannabinoid CB1 receptor in constitutive activation of Gs. *Journal of Neurochemistry*, 72(5), 2032-2038.
- Austin-Brown, S. L., & Chapman, K. D.** 2002. Inhibition of Phospholipase D α by N-Acylethanolamines. *Plant Physiology*, 129(4), 1892-1898.
- Beemster GT, Baskin TI.** 1998. Analysis of cell division and elongation underlying the developmental acceleration of root growth in *Arabidopsis thaliana*. *Plant Physiology* 116,1515-1526.
- Béziat, C., Kleine-Vehn, J., & Feraru, E.** 2017. Histochemical staining of β -Glucuronidase and its spatial quantification. *Methods Molecular Biology* 1497, 73-80.
- Bielski BH, Allen AO.** 1977. Mechanism of the disproportionation of superoxide radicals. *The Journal of Physical Chemistry* 81(11), 1048–1050.
- Bindschedler LV, Dewdney J, Blee KA et al.** 2006. Peroxidase-dependent apoplastic oxidative burst in *Arabidopsis* required for pathogen resistance. *The Plant Journal* 47(6), 851-863.
- Blancaflor EB, Hou G, Chapman KD.** 2003. Elevated levels of N-lauroylethanolamine, an endogenous constituent of desiccated seeds, disrupt normal root development in *Arabidopsis thaliana* seedlings. *Planta* 217, 206–217.
- Blancaflor EB, Kilaru A, Keereetawee J, Khan BR, Faure L, Chapman KD** 2014. N-Acylethanolamines: lipid metabolites with functions in plant growth and development. *The Plant Journal* 79(4), 568-583.
- Bosier, B., Muccioli, G. G., Hermans, E., Lambert, D. M.** 2010. Functionally selective cannabinoid receptor signalling: therapeutic implications and opportunities. *Biochemical Pharmacology* 80(1), 1-12.

Chang Y, Guo J, Gao Y, Chen JG. 2007. Modulation of root cell division by the heterotrimeric G-proteins in Arabidopsis. *Dynamic Cell Biology* 1(1), 72-77.

Chen, J. G., Gao, Y., Jones, A. M. 2006. Differential roles of Arabidopsis heterotrimeric G-protein subunits in modulating cell division in roots. *Plant Physiology* 141(3), 887-897.

Cosgrove DJ, Li LC, Cho HT, Hoffmann-Benning S, Moore RC, Blecker D. 2002. The growing world of expansins. *Plant and Cell Physiology* 43(12), 1436–1444.

Coulon D, Faure L, Salmon M, Wattelet V, Bessoule JJ. 2012. N-Acylethanolamines and related compounds: aspects of metabolism and functions. *Plant Science* 184, 129-140.

Del Río, L. A. 2015. ROS and RNS in plant physiology: an overview. *Journal of Experimental Botany*, 66(10), 2827-2837.

Dunand, C., Crèvecoeur, M., Penel, C. 2007. Distribution of superoxide and hydrogen peroxide in Arabidopsis root and their influence on root development: possible interaction with peroxidases. *New Phytologist*, 174(2), 332-341.

Faure, L., Cavazos, R., Khan, B. R., Petros, R. A., Koulen, P., Blancaflor, E. B., Chapman, K. D. 2015. Effects of synthetic alkamides on Arabidopsis fatty acid amide hydrolase activity and plant development. *Phytochemistry*, 110, 58-71.

Fernández-Marcos, M., Desvoyes, B., Manzano, C., Liberman, L. M., Benfey, P. N., Del Pozo, J. C., Gutierrez, C. 2017. Control of *Arabidopsis* lateral root primordium boundaries by MYB 36. *New Phytologist*, 213(1), 105-112.

Foreman, J., Demidchik, V., Bothwell, J. H., et al 2003. Reactive oxygen species produced by NADPH oxidase regulate plant cell growth. *Nature*, 422(6930), 442-446.

Ghassemian, M., Lutes, J., Chang, H. S., Lange, I., Chen, W., Zhu, T., Wang, X., Lange, B. M. (2008). Absciscic acid-induced modulation of metabolic and redox control pathways in *Arabidopsis thaliana*. *Phytochemistry*, 69(17), 2899-2911.

Gertsch, J. 2008. Immunomodulatory lipids in plants: plant fatty acid amides and the human endocannabinoid system. *Planta Medica*, 74(06), 638-650.

Hartje, S., Zimmermann, S., Klonus, D., & Mueller-Roeber, B. 2000. Functional characterisation of LKT1, a K⁺ uptake channel from tomato root hairs, and comparison with the closely related potato inwardly rectifying K⁺ channel SKT1 after expression in *Xenopus* oocytes. *Planta*, 210(5), 723-731.

Hayyan, M., Hashim, M. A., & Al Nashef, I. M. 2016. Superoxide ion: generation and chemical implications. *Chemical Reviews*, 116(5), 3029-3085.

Ilan, N., Schwartz, A., & Moran, N. 1996. External protons enhance the activity of the hyperpolarization-activated K channels in guard cell protoplasts of *Vicia faba*. *The Journal of Membrane Biology*, 154(2), 169-181.

Jaspers, P., Kangasjärvi, J. 2010. Reactive oxygen species in abiotic stress signaling. *Physiologia Plantarum*, 138(4), 405-413.

Kagenishi, T., Yokawa, K., Baluška, F. 2016. MES buffer affects *Arabidopsis* root apex zonation and root growth by suppressing superoxide generation in root apex. *Frontiers in Plant Science*, 7, 79.

Kravchenko, V. V., Kaufmann, G. F., Mathison, J. C., et al. 2006. N-(3-oxo-acyl) homoserine lactones signal cell activation through a mechanism distinct from the canonical pathogen-associated molecular pattern recognition receptor pathways. *Journal of Biological Chemistry*, 281(39), 28822-28830.

Kunos, G., Járαι, Z., Bátkai, S., Goparaju, S. K., Ishac, E. J., Liu, J., Wang, L., Wagner, J. A. 2000. Endocannabinoids as cardiovascular modulators. *Chemistry and Physics of Lipids*, 108(1-2), 159-168.

Li, N., Sun, L., Zhang, L., Song, Y., Hu, P., Li, C., Hao, F. S. 2015. AtrbohD and AtrbohF negatively regulate lateral root development by changing the localized accumulation of superoxide in primary roots of *Arabidopsis*. *Planta*, 241(3), 591-602.

Liang, X., Ma, M., Zhou, Z., et al. 2018. Ligand-triggered de-repression of *Arabidopsis* heterotrimeric G proteins coupled to immune receptor kinases. *Cell Research*, 28(5), 529-543.

Liu, F., Bian, Z., Jia, Z., Zhao, Q., Song, S. 2012. The GCR1 and GPA1 participate in promotion of *Arabidopsis* primary root elongation induced by N-acyl-homoserine lactones, the bacterial quorum-sensing signals. *Molecular Plant-Microbe Interactions*, 25(5), 677-683.

Livanos, P., Apostolakos, P., Galatis, B. 2012. Plant Cell Division: ROS homeostasis is required. *Plant Signaling & Behavior*, 7(7), 771-778.

López-Bucio, J., Millán-Godínez, M., Méndez-Bravo, A., Morquecho-Contreras, A., Ramírez-Chávez, E., Molina-Torres, J., Pérez-Torres, A., Higuchi, M., Kakimoto T., Herrera-Estrella, L. 2007. Cytokinin receptors are involved in alkamide regulation of root and shoot development in *Arabidopsis*. *Plant Physiology*, 145(4), 1703-1713.

Marklund, S. 1976. Spectrophotometric study of spontaneous disproportionation of superoxide anion radical and sensitive direct assay for superoxide dismutase. *Journal of Biological Chemistry*, 251(23), 7504-7507.

Manzano, C., Pallero-Baena, M., Casimiro, I., De Rybel, B., Orman-Ligeza, B., Van Isterdael, G., Beeckman, T., Draye, X., Del Pozo, J. C. 2014. The emerging role of reactive oxygen species signaling during lateral root development. *Plant Physiology*, 165(3), 1105-1119.

Mathesius, U., Mulders, S., Gao, M., Teplitski, M., Caetano-Anollés, G., Rolfe, B. G., Bauer, W. D. 2003. Extensive and specific responses of a eukaryote to bacterial quorum-sensing signals. *Proceedings of the National Academy of Sciences*, 100(3), 1444-1449.

Méndez-Bravo, A., Calderón-Vázquez, C., Ibarra-Laclette, E., Raya-González, J., Ramírez-Chávez, E., Molina-Torres, J., Guevara-García, A.A., López-Bucio, J., Herrera-Estrella, L. 2011. Alkamides activate jasmonic acid biosynthesis and signaling pathways and confer resistance to *Botrytis cinerea* in *Arabidopsis thaliana*. *PloS One*, 6(11), e27251.

Méndez-Bravo, A., Raya-González, J., Herrera-Estrella, L., López-Bucio, J. 2010. Nitric oxide is involved in alkamide-induced lateral root development in *Arabidopsis*. *Plant and Cell Physiology*, 51(10), 1612-1626.

Micheli, F. 2001. Pectin methylesterases: cell wall enzymes with important roles in plant physiology. *Trends in Plant Science*, 6(9), 414-419.

Miller, E. W., Dickinson, B. C., Chang, C. J. 2010. Aquaporin-3 mediates hydrogen peroxide uptake to regulate downstream intracellular signaling. *Proceedings of the National Academy of Sciences*, 107(36), 15681-15686.

Miller, A. F. 2012. Superoxide dismutases: ancient enzymes and new insights. *FEBS letters*, 586(5), 585-595.

Mittler, R., Vanderauwera, S., Suzuki, N., Miller, G. A. D., Tognetti, V. B., Vandepoele, K., Gollery, M., Shulaev, V., Van Breusegem, F. 2011. ROS signaling: the new wave? *Trends in Plant Science*, 16(6), 300-309.

Mittler, R. 2017. ROS are good. *Trends in Plant Science*, 22(1), 11-19.

Molinatorres, J., Salgado-Garciglia, R., Ramirez-Chavez, E., Del Rio, R. E. 1996. Purely olefinic alkamides in *Heliopsis longipes* and *Acmella* (*Spilanthes*) *oppositifolia*. *Biochemical Systematics and Ecology*, 24(1), 43-47.

Molina-Torres, J., Salazar-Cabrera, C. J., Armenta-Salinas, C., Ramírez-Chávez, E. 2004. Fungistatic and bacteriostatic activities of alkamides from *Heliopsis longipes* roots: affinin and reduced amides. *Journal of Agricultural and Food Chemistry*, 52(15), 4700-4704.

Monshausen, G. B., Bibikova, T. N., Weisenseel, M. H., Gilroy, S. 2009. Ca²⁺ regulates reactive oxygen species production and pH during mechanosensing in *Arabidopsis* roots. *The Plant Cell*, 21(8), 2341-2356.

Ogasawara, Y., Kaya, H., Hiraoka, G., et al. 2008. Synergistic activation of the *Arabidopsis* NADPH oxidase AtrbohD by Ca²⁺ and phosphorylation. *Journal of Biological Chemistry*, 283(14), 8885-8892.

Orman-Ligeza, B., Parizot, B., De Rycke, R., Fernandez, A., Himschoot, E., Van Breusegem, F., Bennett, M.J., Périlleux, C., Beeckman T., Draye, X. 2016. RBOH-mediated ROS production facilitates lateral root emergence in *Arabidopsis*. *Development*, 143(18), 3328-3339.

Ortíz-Castro R, Martínez-Trujillo M, López-Bucio J. 2008. N-acyl-L-homoserine lactones: a class of bacterial quorum-sensing signals alter post-embryonic root development in *Arabidopsis thaliana*. *Plant, Cell & Environment*, 31(10), 1497-1509.

Overmyer, K., Brosché, M., & Kangasjärvi, J. 2003. Reactive oxygen species and hormonal control of cell death. *Trends in Plant Science*, 8(7), 335-342.

Passardi, F., Tognolli, M., De Meyer, M., Penel, C., Dunand, C. 2006. Two cell wall associated peroxidases from *Arabidopsis* influence root elongation. *Planta*, 223(5), 965-974.

Potters, G., Pasternak, T. P., Guisez, Y., Jansen, M. A. (2009). Different stresses, similar morphogenic responses: integrating a plethora of pathways. *Plant, Cell & Environment*, 32(2), 158-169.

Prachayasittikul, V., Prachayasittikul, S., Ruchirawat, S., & Prachayasittikul, V. (2013). High therapeutic potential of *Spilanthes acmella*: a review. *EXCLI journal*, 12, 291.

Qi, J., Wang, J., Gong, Z., Zhou, J. M. 2017. Apoplastic ROS signaling in plant immunity. *Current Opinion in Plant Biology*, 38, 92-100.

Ramírez-Chávez, E., López-Bucio, J., Herrera-Estrella, L., Molina-Torres, J. 2004. Alkamides isolated from plants promote growth and alter root development in *Arabidopsis*. *Plant Physiology*, 134(3), 1058-1068.

Rios, M. Y., Olivo, H. F. 2014. Natural and synthetic alkamides: applications in pain therapy. In: Rahman, E, ed. *Studies in Natural Products Chemistry*. Elsevier. Vol. 43, 79-121.

Sagi, M., Fluhr, R. 2006. Production of reactive oxygen species by plant NADPH oxidases. *Plant Physiology*, 141(2), 336-340.

Schenk, S. T., Stein, E., Kogel, K. H., Schikora, A. 2012. *Arabidopsis* growth and defense are modulated by bacterial quorum sensing molecules. *Plant Signaling & Behavior*, 7(2), 178-181.

Schneider, C. A., Rasband, W. S., Eliceiri, K. W. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*, 9(7), 671-675.

Schuhegger, R., Ihring, A., Gantner, S., et al. 2006. Induction of systemic resistance in tomato by N-acyl-L-homoserine lactone-producing rhizosphere bacteria. *Plant, Cell & Environment*, 29(5), 909-918.

Sherrier, D. J., VandenBosch, K. A. 1994. Secretion of cell wall polysaccharides in *Vicia* root hairs. *The Plant Journal*, 5(2), 185-195.

Song, S., Jia, Z., Xu, J., Zhang, Z., Bian, Z. 2011. N-butyryl-homoserine lactone, a bacterial quorum-sensing signaling molecule, induces intracellular calcium elevation in *Arabidopsis* root cells. *Biochemical and Biophysical Research Communications*, 414(2), 355-360.

Suzuki, N., Miller, G., Morales, J., Shulaev, V., Torres, M. A., Mittler, R. 2011. Respiratory burst oxidases: the engines of ROS signaling. *Current Opinion in Plant Biology*, 14(6), 691-699.

Thordal-Christensen, H., Zhang, Z., Wei, Y., Collinge, D. B. 1997. Subcellular localization of H₂O₂ in plants. H₂O₂ accumulation in papillae and hypersensitive response during the barley—powdery mildew interaction. *The Plant Journal*, 11(6), 1187-1194.

Tognetti, V. B., Mühlenbock, P. E. R., Van Breusegem, F. 2012. Stress homeostasis—the redox and auxin perspective. *Plant, Cell & Environment*, 35(2), 321-333.

Torres, M. A., Dangl, J. L., Jones, J. D. 2002. *Arabidopsis* gp91phox homologues AtrbohD and AtrbohF are required for accumulation of reactive oxygen intermediates in the plant defense response. *Proceedings of the National Academy of Sciences*, 99(1), 517-522.

Torres, M. A. 2010. ROS in biotic interactions. *Physiologia Plantarum*, 138(4), 414-429.

Tripathy, S., Venables, B. J., & Chapman, K. D. 1999. N-Acylethanolamines in signal transduction of elicitor perception. Attenuation of alkalization response and activation of defense gene expression. *Plant Physiology*, 121(4), 1299-1308.

Tripathy, S., Kleppinger-Sparace, K., Dixon, R. A., Chapman, K. D. 2003. N-acylethanolamine signaling in tobacco is mediated by a membrane-associated, high-affinity binding protein. *Plant Physiology*, 131(4), 1781-1791.

Tsukagoshi, H., Busch, W., Benfey, P. N. 2010. Transcriptional regulation of ROS controls transition from proliferation to differentiation in the root. *Cell*, 143(4), 606-616.

Tsukagoshi, H. 2016. Control of root growth and development by reactive oxygen species. *Current Opinion in Plant Biology*, 29, 57-63.

Ubeda-Tomás, S., Federici, F., Casimiro, I., Beemster, G. T., Bhalerao, R., Swarup, R., Doerner, P., Haseloff J., Bennett, M. J. 2009. Gibberellin signaling in the endodermis controls *Arabidopsis* root meristem size. *Current Biology*, 19(14), 1194-1199.

Urano, D., & Jones, A. M. (2014). Heterotrimeric G protein–coupled signaling in plants. *Annual Review of Plant Biology*, 65, 365-384.

von Rad, U., Klein, I., Dobrev, P. I., Kottova, J., Zazimalova, E., Fekete, A., Hartmann, A., Schmitt-Kopplin, P., Durner, J. 2008. Response of *Arabidopsis thaliana* to N-hexanoyl-DL-homoserine-lactone, a bacterial quorum sensing molecule produced in the rhizosphere. *Planta*, 229(1), 73-85.

Wang, Y. S., Shrestha, R., Kilaru, A., Wiant, W., Venables, B. J., Chapman, K. D., Blancaflor, E. B. 2006. Manipulation of *Arabidopsis* fatty acid amide hydrolase expression modifies plant growth and sensitivity to N-acyl ethanolamines. *Proceedings of the National Academy of Sciences*, 103(32), 12197-12202.

Wilson, R. I., Nicoll, R. A. 2002. Endocannabinoid signaling in the brain. *Science*, 296(5568), 678-682.

Yan, C., Cannon, A. E., Watkins, J., Keereetaweep, J., Khan, B. R., Jones, A. M., B Blancaflor, E., Azad, R.K., Chapman, K. D. 2020. Seedling chloroplast responses induced by *N*-linolenylethanolamine require intact G-protein complexes. *Plant Physiology*, 184(1), 459-477.

Zhang, Y., Guo, W. M., Chen, S. M., Han, L., Li, Z. M. 2007. The role of *N*-lauroylethanolamine in the regulation of senescence of cut carnations (*Dianthus caryophyllus*). *Journal of Plant Physiology*, 164(8), 993-1001.

Zandonadi, D. B., Santos, M. P., Dobbss, L. B., Olivares, F. L., Canellas, L. P., Binzel, M. L., Okorokova-Façanha, A.L., Façanha, A. R. 2010. Nitric oxide mediates humic acids-induced root development and plasma membrane H⁺-ATPase activation. *Planta*, 231(5), 1025-1036.

Zhang, W., Jeon, B. W., Assmann, S. M. 2011. Heterotrimeric G-protein regulation of ROS signalling and calcium currents in *Arabidopsis* guard cells. *Journal of Experimental Botany*, 62(7), 2371-2379.

Zhao, J., Davis, L. C., Verpoorte, R. 2005. Elicitor signal transduction leading to production of plant secondary metabolites. *Biotechnology Advances*, 23(4), 283-333.

Zhao, Q., Zhang, C., Jia, Z., Huang, Y., Li, H., Song, S. 2015. Involvement of calmodulin in regulation of primary root elongation by *N*-3-oxo-hexanoyl homoserine lactone in *Arabidopsis thaliana*. *Frontiers in Plant Science*, 5, 807.

Zhao, Y., Zhang, Y., Liu, F., Wang, R., Huang, L., & Shen, W. 2019. Hydrogen peroxide is involved in methane-induced tomato lateral root formation. *Plant Cell Reports*, 38(3), 377-389.

Table 1. *N*-acyl amides in plant development

Type	Chain length: insaturations	Activity	Reference
NAE	12:00	Flower senescence: Increase activity of SOD, Cat, APX and GR.	Zhang <i>et al.</i> , 2007
NAE	12:00	Reduction of: primary root growth, secondary roots, root hair formation, apex swelling, cell invaginations of the plasma membrane, increase levels of vesicles at the cell periphery, improper cell walls near the meristematic region, disorganize endomembrane system and alter vesicular trafficking. Increase the expression of ABRE genes	Blancaflor <i>et al.</i> , 2003
NAE	12:0 to 18:3	Inhibition of PLD α activity	Austin-Brown and Chapman, 2002
NAE	12:0, 18:3	Reduction of root elongation rate.	Wang <i>et al.</i> , 2010; Blancaflor <i>et al.</i> , 2003
NAE	12:0, 14:0, 18:0, 18:1, 18:3, 20:4	Plant defense: Inhibit alkalization of the extracellular medium, modulate the ion flux in the plasma membrane and Induce PAL expression.	Tripathy <i>et al.</i> , 1999; Tripathy <i>et al.</i> , 2003
NAE	12:0, 18:2	Interacts with the ABA signaling pathway to elicit secondary dormancy	Blancaflor <i>et al.</i> , 2014; Wang <i>et al.</i> , 2006
AHL	C4-HSL	Intracellular Ca ²⁺ elevation	Song <i>et al.</i> , 2011
AHL	C4-HSL, C6-HSL, C8-HSL	Increase primary root growth and plant biomass. Resistance against necrotrophic pathogens. Increase SA-levels and defense-gene regulation.	Liu <i>et al.</i> , 2012; Schenk <i>et al.</i> , 2012; von Rad <i>et al.</i> , 2008; Schuëgger <i>et al.</i> , 2006
AHL	C6-HSL	Alteration in auxin/cytokinin level and herbivore susceptibility	von Rad <i>et al.</i> , 2008; Schenk <i>et al.</i> , 2012;
AHL	C10-HSL, C12-HSL	Inhibition of primary root growth.	Zhao <i>et al.</i> , 2015

Figure legends

Figure. 1. Effect of affinin on root architecture in *Arabidopsis* Col-0 accession. (A) Plants grown on affinin treatments (Bars=1cm); (B) primary root length ($n=10$); (C) number of emerged lateral roots ($n=10$) and (D) root hair length ($n=10$). (E, H) confocal images of the primary root from tip to first root hair bulge (Bars=0.1mm), (F, I) confocal images of cells from the elongation-differentiation zone (Bars=0.05mm); (G, J) confocal images of apical meristem size. White arrows indicate root hair bulge and grey arrows shows the transition zone (Bars=0.05mm). (K) Shows the mean distance from QC to the first root hair bulge ($n=5$), (L) is the mean value of cell length in the elongation/differentiation zone ($n=3$) and (M) shows the mean value of the apical meristem length ($n=5$). Data was analysed with one-way ANOVA followed by a Tukey test using the InfoStat software (www.infostat.com.ar). Different lower-case letters are used to indicate means that differ significantly ($P \leq 0.05$). Micrographs were adjusted with the same settings; Colour saturation: 0%, Brightness: 40% and Contrast: 40%

Figure.2. Affinin induced root-developmental changes are mediated by ROS. (A) Detection of endogenous H_2O_2 accumulation with 3, 3'-diaminobenzidine (DAB) staining and (B) nitroblue tetrazolium (NBT) staining, visualizing superoxide ($O_2^{\cdot-}$) accumulation of primary roots treated with affinin (Bars=300 μ m). (C) Visualization of *in situ* accumulation of endogenous H_2O_2 using DAB staining in root apical meristem (Bars=100 μ m) and lateral root emerging sites (Bars=50 μ m) of Col-0 wild type plants. Data shown in (D) represent the mean $\pm$ SE of H_2O_2 staining intensity measured in (C). (E) Affinin-induced *Arabidopsis* (Col-0) lateral rooting is sensitive to synthetic inhibitor of flavoenzymes, diphenyleneiodonium (DPI), used here to target NAD(P)H oxidases (Bars=1cm). Data shown in (F) represent the mean $\pm$ SE of emerged lateral root number measured in (E). Data in (D) was analysed with one-way ANOVA ($n=10$) and a Tukey test using the InfoStat software (www.infostat.com.ar). Data in (F) was analysed with two-way ANOVA ($n=10$) and a Tukey test using Minitab software (<https://www.minitab.com/>). Different lower-case letters are used to indicate means that differ significantly ($P \leq 0.05$). Micrographs were adjusted with the same settings; Colour saturation: 0%, Brightness: 0% and Contrast: 20%.

Figure 3. Effect of affinin on root system architecture in the class III peroxidase mutant *Arabidopsis peroxidase34* (*prx34*). (A) Represents the mean $\pm$ SE of primary root length ($n=10$) while (B) is the number of emerged lateral roots ($n=10$) of plants grown in different affinin treatments. (C) Visualization of *in situ* accumulation of endogenous H_2O_2 using 3, 3'-diaminobenzidine (DAB) staining in root apical meristem (Bars=100 μ m) and lateral root emerging sites (Bars=50 μ m) of *prx34* mutant plants. Data shown in (D) represent the mean $\pm$ SE of H_2O_2 staining intensity measured in (C). Data was analysed with two-way ANOVA ($n=10$) and a Tukey test using Minitab software (<https://www.minitab.com/>). Different lower-case letters are used to indicate means that differ significantly ($P \leq 0.05$). Micrographs were adjusted with the same settings; Colour saturation: 0%, Brightness: 0% and Contrast: 20%.

Figure 4. RESPIRATORY BURST OXIDASE HOMOLOG (RBOH) mediated ROS production is involved in root developmental changes induced by affinin. Data shown in (A) represent the mean $\pm$ SE of primary root length ($n=10$) while data shown in (B) represent the mean $\pm$ SE of emerged lateral roots ($n=10$). (C) DAB staining of root tips and lateral root primordia surrounding area (D). Data in (E) represent the mean $\pm$ SE of H_2O_2 staining intensity in root tips ($n=10$) and (F) in lateral root primordia surrounding area ($n=10$). Photographs in (G) shown the representative morphology of root hairs, while data shown in (H) represent the mean $\pm$ SE of root hair length measured in (G) ($n=10$). Data was analysed with two-way ANOVA ($n=10$) and a Tukey test using Minitab software (<https://www.minitab.com/>). Different lower-case letters are used to indicate means that differ significantly ($P \leq 0.05$). Micrographs were adjusted with the same settings; Colour saturation: 0%, Brightness: 0% and Contrast: 20%.

Figure 5. Effect of affinin on root growth of *Arabidopsis* heterotrimeric G-protein subunit mutants. Data shown in (A) represent the mean $\pm$ SE of primary root length ($n=10$) and (B) the emerged lateral roots number ($n=10$) from heterotrimeric G-protein mutants while (C) represent the mean $\pm$ SE of primary root length ($n=10$) and (D) the emerged lateral roots number ($n=10$) from the regulator of G-protein mutant. Data was analysed with two-way ANOVA ($n=10$) and a Tukey test using Minitab software (<https://www.minitab.com/>). Different lower-case letters are used to indicate means that differ significantly ($P \leq 0.05$).

Figure 6. Involvement of extracellular pH change in affinin-induced developmental responses. (A) Rhizosphere acidification in the maturation zone of *Arabidopsis* roots in response to different affinin treatments (Bars=1cm); the scale on the left is a pH reference scale in the agarose without plants. Data shown in (B) represent the mean $\pm$ SE of medium pH measured in a region of interest (ROI) near the roots of seedlings ($n=7$). (C) Photographs of full seedlings (Bars=1cm) and (F) root hairs (Bars=0.5mm) of plants treated with or without affinin grown in a medium containing or lacking MES buffer (0.5g L^{-1}). Data shown represent the mean $\pm$ SE of primary root length (D), emerged lateral roots (E) and root hair length (G) ($n=10$). Data was analysed with two-way ANOVA and a Tukey test using Minitab software (<https://www.minitab.com/>). Different lower-case letters are used to indicate means that differ significantly ($P \leq 0.05$). Micrographs were adjusted with the same settings; Colour saturation: 0%, Brightness: 0% and Contrast: 20%.

Figures

Figure 1

Fig. 1. Effect of affinin on root architecture in *Arabidopsis* Col-0 accession. (A) Plants grown on affinin treatments (Bars=1cm); (B) primary root length ($n=10$); (C) number of emerged lateral roots ($n=10$) and (D) root hair length ($n=10$). (E, H) confocal images of the primary root from tip to first root hair bulge (Bars=0.1mm), (F, I) confocal images of cells from the elongation-differentiation zone (Bars=0.05mm); (G, J) confocal images of apical meristem size. White arrows indicate root hair bulge and grey arrows shows the transition zone (Bars=0.05mm). (K) Shows the mean distance from QC to the first root hair bulge ($n=5$), (L) is the mean value of cell length in the elongation/differentiation zone ($n=3$) and (M) shows the mean value of the apical meristem length ($n=5$). Data was analysed with one-way ANOVA followed by a Tukey test using the InfoStat software (www.infostat.com.ar) Different lower-case letters are used to indicate means that differ significantly ($P \leq 0.05$). Micrographs were adjusted with the same settings; Colour saturation: 0%, Brightness: 40% and Contrast: 40%

Figure 2

Fig. 2. Affinin induced root-developmental changes are mediated by ROS. (A) Detection of endogenous H_2O_2 accumulation with 3, 3'-diaminobenzidine (DAB) staining and (B) nitroblue tetrazolium (NBT) staining, visualizing superoxide ($O_2^{\bullet -}$) accumulation of primary roots treated with affinin (Bars=300 μm). (C) Visualization of *in situ* accumulation of endogenous H_2O_2 using DAB staining in root apical meristem (Bars=100 μm) and lateral root emerging sites (Bars=50 μm) of Col-0 wild type plants. Data shown in (D) represent the mean $\pm$ SE of H_2O_2 staining intensity measured in (C). (E) Affinin-induced *Arabidopsis* (Col-0) lateral rooting is sensitive to synthetic inhibitor of flavoenzymes, diphenyleneiodonium (DPI), used here to target NAD(P)H oxidases (Bars=1cm). Data shown in (F) represent the mean $\pm$ SE of emerged lateral root number measured in (E). Data in (D) was analysed with one-way ANOVA ($n=10$) and a Tukey test using the InfoStat software (www.infostat.com.ar). Data in (F) was analysed with two-way ANOVA ($n=10$) and a Tukey test using Minitab software (<https://www.minitab.com/>). Different lower-case letters are used to indicate means that differ significantly ($P \leq 0.05$). Micrographs were adjusted with the same settings; Colour saturation: 0%, Brightness: 0% and Contrast: 20%.

Figure 3

Fig. 3. Effect of affinin on root system architecture in *Arabidopsis peroxidase34* (*prx34*) class III peroxidase mutant. (A) represents the mean $\pm$ SE of primary root length ($n=10$) while (B) is the number of emerged lateral roots ($n=10$) of plants grown in different affinin treatments. (C) Visualization of *in situ* accumulation of endogenous H_2O_2 using 3, 3'-diaminobenzidine (DAB) staining in root apical meristem (Bars=100μm) and lateral root emerging sites (Bars=50μm) of *prx34* mutant plants. Data shown in (D) represent the mean $\pm$ SE of H_2O_2 staining intensity measured in (C). Data was analysed with two-way ANOVA ($n=10$) and a Tukey test using Minitab software (<https://www.minitab.com/>). Different lower-case letters are used to indicate means that differ significantly ($P \leq 0.05$). Micrographs were adjusted with the same settings; Colour saturation: 0%, Brightness: 0% and Contrast: 20%.

Figure 4

Fig. 4. RESPIRATORY BURST OXIDASE HOMOLOG (RBOH) mediated ROS production is involved in root developmental changes induced by affinin. Data shown in (A) represent the mean $\pm$ SE of primary root length ($n=10$) while data shown in (B) represent the mean $\pm$ SE of emerged lateral roots ($n=10$). (C) DAB staining of root tips and lateral root

primordia surrounding area (D). Data in (E) represent the mean $\pm$ SE of H₂O₂ staining intensity in root tips ($n=10$) and (F) in lateral root primordia surrounding area ($n=10$). Photographs in (G) shown the representative morphology of root hairs, while data shown in (H) represent the mean $\pm$ SE of root hair length measured in (G) ($n=10$). Data was analysed with two-way ANOVA ($n=10$) and a Tukey test using Minitab software (<https://www.minitab.com/>). Different lower-case letters are used to indicate means that differ significantly ($P \leq 0.05$). Micrographs were adjusted with the same settings; Colour saturation: 0%, Brightness: 0% and Contrast: 20%.

Figure 5

Figure 5. Effect of affinin on root growth of *Arabidopsis* heterotrimeric G-protein subunits mutants. Data shown in (A) represent the mean $\pm$ SE of primary root length ($n=10$) and (B) the emerged lateral roots number ($n=10$) from heterotrimeric G-protein mutants while (C) represent the mean $\pm$ SE of primary root length ($n=10$) and (D) the emerged lateral roots number ($n=10$) from the regulator of G-protein mutant. Data was analysed with two-way ANOVA ($n=10$) and a Tukey test using Minitab software (<https://www.minitab.com/>). Different lower-case letters are used to indicate means that differ significantly ($P \leq 0.05$).

Figure 6

Figure 6. Involvement of extracellular pH change in affinin-induced developmental response. (A) Rhizosphere acidification in the maturation zone of *Arabidopsis* roots in response to different affinin treatments (Bars=1cm); the scale on the left is a pH reference scale in the agarose without plants. Data shown in (B) represent the mean $\pm$ SE of medium pH measured in a region of interest (ROI) near the roots of seedlings ($n=7$). (C) Photographs of full seedlings (Bars=1cm) and (F) root hairs (Bars=0.5mm) of plants treated with or without affinin grown in a medium containing or lacking MES buffer (0.5 g L⁻¹). Data shown represent the mean $\pm$ SE of primary root length (D), emerged lateral roots (E) and root hair length (G) ($n=10$). Data was analysed with two-way ANOVA and a Tukey test using Minitab software (<https://www.minitab.com/>). Different lower-case letters are used to indicate means that differ significantly ($P \leq 0.05$). Micrographs were adjusted with the same settings; Colour saturation: 0%, Brightness: 0% and Contrast: 20%.