Revealing the secrets beneath grapevine and *Plasmopara viticola* early communication: a picture of host and pathogen proteomes

Running title: Grapevine and *P. viticola* proteomes communication

Joana Figueiredo^{1,*}, Rita B. Santos¹, Leonor Guerra-Guimarães^{2,3}, Céline C. Leclercq⁴, Jenny Renaut⁴, Lisete Sousa⁵, Andreia Figueiredo¹

1 Grapevine Pathogen Systems Lab, BioISI – Biosystems & Integrative Sciences Institute, Plant Biology Department, Faculty of Sciences, University of Lisbon, 1749-016 Lisboa, Portugal

2 Centro de Investigação das Ferrugens do Cafeeiro, Instituto Superior de Agronomia, Universidade de Lisboa, 2784-505 Oeiras, Portugal

3 Linking Landscape, Environment, Agriculture and Food, Instituto Superior de Agronomia, Universidade de Lisboa, 1349-017 Lisboa, Portugal

4 Luxembourg Institute of Science and Technology, L-4362 Esch-sur-Alzette, Luxembourg

5 Departamento de Estatística e Investigação Operacional, Centro de Estatística e Aplicações, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

* Correspondence: jffigueiredo@fc.ul.pt

Abstract:

Plant apoplast is the first hub of plant-pathogen communication where pathogen effectors are recognized by plant defensive proteins and cell receptors and several signal transduction pathways are activated. As a result of this first contact, the host triggers a defence response that involves the modulation of several extra and intracellular proteins. In grapevine-pathogen interactions, little is known about the communication between cells and apoplast. Also, the role of apoplastic proteins in response to pathogens still remains a blackbox. In this study we focused on the first 6 hours after *Plasmopara viticola* inoculation to evaluate grapevine proteome modulation in the apoplastic fluid (APF) and whole leaf tissue. *Plasmopara viticola* proteome was also assessed enabling a deeper understanding of plant and pathogen communication. Our results showed that oomycete recognition, plant cell wall modifications, ROS signalling and disruption of oomycete structures are triggered in Regent after *P. viticola* inoculation. Our results highlight a strict relation between the apoplastic pathways modulated and the proteins identified in the whole leaf proteome. On the other hand, *P. viticola* proteins related to growth/morphogenesis and virulence mechanisms were the most predominant. This pioneer study highlights the early dynamics of extra and intracellular communication in grapevine defence activation that leads to the successful establishment of an incompatible interaction.

Keywords (3-10): *Vitis vinifera*, apoplastic fluid, cell compartments, secretomes, oomycete, proteomics

1. Introduction

As sessile organisms, plants have developed mechanisms to rapidly adapt to environmental changes and pathogen attack. To overcome pathogen challenges, plants must quickly recognize the invaders and mount a successful defence strategy. This chess game between the plant and pathogen is illustrated by the “zig zag model”, coined by Jones and Dangl in 2006¹. In this model, the plants first recognize the pathogen-associated molecular patterns (PAMPs) through pattern recognition receptors (PRR) in the apoplast, leading to a PAMP-triggered immunity (PTI). The apoplast is extremely relevant for plant defence since is where plant and pathogen first meet and where recognition begins. Pathogen recognition culminates in the activation of plant defence responses including the induction of defence genes, production of reactive oxygen species (ROS) and deposition of callose. In a second phase, effector-triggered susceptibility or ETS, the pathogen overcomes the plant first response by deploying effectors that increase pathogen virulence, like Crinklers and RxLR effectors¹. In an incompatible interaction, the plant recognizes the pathogen effectors through R-proteins. The interaction between plant R-proteins and pathogen effectors results in an effector-triggered immunity (ETI), that ultimately results in a hypersensitive cell death response (HR) at the pathogen entry site¹. This interaction implies a tight communication between host and pathogen with the traffic of plant proteins and pathogen effector proteins between the apoplast and the intercellular space. While still misgraded, the study of plant apoplast is of extreme importance in plant-pathogen interactions so to identify proteins with a key role in plant defence strategies and better understand their interaction with pathogen molecules. Apoplast proteome was characterized for few plant models, constitutively

or under abiotic/biotic stress, as for example, grapevine^{2,3}, poplar⁴, tobacco⁵, cowpea⁶, rice⁷, coffee^{8,9} and Arabidopsis¹⁰. However, few studies focus on uncovering apoplast proteome modulation considering plant-pathogen interactions and even less when considering woody crop plants, such as grapevine and obligatory biotrophic oomycetes, as the downy mildew etiological agent, *Plasmopara viticola*.

Grapevine (*Vitis vinifera* L.), is one of the major crops grown in temperate climates, however is highly susceptible to downy mildew, caused by *P. viticola* ((Berk. and Curt.) Berl. & de Toni)¹¹. In Europe, *P. viticola* infection leads to heavy crop losses and disease management for downy mildew relies on the massive use of pesticides in susceptible varieties in each growing season. This practice is against the demands of the European Union guidelines for pesticide reduction and sustainable viticulture (Directive 2009/128/EC), so the search for more plant- and environment-friendly solutions is imperative.

In a modern viticulture context, the development of grapevine crossing lines, in breeding programs, is a very well established and accepted strategy to fight against the excessive use of pesticides. These crossing lines are the result of the introgression of pathogen resistant genes, present in Asian and American *Vitis* species, with genes related to the good quality of grapes for wine production, present in susceptible grapevine cultivars. The result is a cultivar that present desired characteristics for wine producers at the same time that resists more to pathogen attack. 'Regent' is a successful example of breeding for resistance and harbours RPV3.1 resistance to *P. viticola* loci¹². Several studies have been performed in 'Regent'-*P. viticola* interaction with the aim to better understand the molecular mechanisms and the key molecules that are responsible to the well-known tolerance that this crossing line has against *P. viticola*¹³⁻¹⁷.

In a climate change scenario, viticulture will face new emerging diseases as well as several outbreaks of the established diseases, such as downy mildew. Thus, a comprehensive knowledge on the grapevine strategies to overcome pathogens, mainly in cultivars with some resistance level, as well as the evolution of pathogen infection mechanisms is paramount to tackle this challenge. Thus, in the present study, we have focused on the early communication between grapevine and *P. viticola* and assessed, for the first time, grapevine apoplast proteome modulation and *P. viticola* proteome and secretomes. We have focused on the first 6 hours post inoculation (hpi), as the events occurring in this time-point were previously shown to be crucial for the outcome of the interaction. We have also highlighted extra and intra-cellular communication pathways by comparing the proteome modulation in the apoplast and in the whole leaf. Up to our knowledge this is the first time where grapevine apoplast and whole leaf proteome communication is revealed during host-pathogen interaction and also the first *P. viticola* proteome sequencing. Our results elucidate the interaction between grapevine and *P. viticola* proteins taking place in the apoplast and how the plant and pathogen proteomes evolve at the first stages of infection.

100

101

2. Results

2.1 Early 'Regent' APF proteome modulation under *P. viticola* infection

The impact of *P. viticola* infection in the modulation of APF proteome in 'Regent' leaves was analysed at 6 hours post inoculation (hpi). By a principal components analysis (PCA), a clear distinction between the proteome of inoculated and mock-inoculated (control) samples was obtained (Fig.1). The distribution of the biological replicates within the PCA scores plot indicates the absence of unwanted variation in the dataset, increasing the confidence in the reproducibility of the differential accumulation analysis.

Fig.1. Principal component analysis of the differential protein profiles in *V. vinifera* cv. 'Regent' at 6 hours post-inoculation with *P. viticola*. The plot shows principal component 1 (PC1) on X axis and principal component 2 (PC2) on Y axis, together they explain 64.7% of protein abundance variability. Ri: 'Regent' inoculated samples; Rm: 'Regent' mock-inoculated samples.

When comparing APF inoculated samples with mock-inoculated samples, a hundred and eighteen proteins were differentially accumulated (DAPs; 74 up accumulated and 44 down accumulated). These proteins are mainly related to stress response, signal transduction, cell wall metabolism, transport and protein metabolism (Fig.2). At 6hpi, *P. viticola* infection leads to an increase in the presence of plant stress response proteins, like glucan endo-1,3- β -glucosidases, disease resistance proteins RPV1-like and RUN1-like [associated to Resistance to *P. viticola* (RPV) loci and Resistance to *Uncinula necator* (RUN) loci], receptor-like protein kinase FERONIA and GDSL esterases/lipases. An up accumulation of LRR kinases related to signal transduction occurs. ROS-related proteins, like peroxidase 4-like and xanthine dehydrogenase 1-like isoform X1, were also detected in 'Regent' APF after oomycete challenged.

Fig.2. Biological process annotation of the differentially accumulated 'Regent' APF proteins, 6 hours post inoculation with *P. viticola*. Dark grey bars: percentage of down accumulated proteins; Light grey bars: percentage of up accumulated proteins.

Several proteins involved in 'Regent' defence mechanism were modulated at 6hpi (Table 1). Indeed, apoplastic proteins related to oomycete perception, that may lead to the activation of several defence signalling pathways, were found to be accumulated (Table 1). Apoplastic proteins associated with the remodelling of plant cell wall were also identified as well as proteins associated with auxin signalling and its regulation in response to *P. viticola* infection (Table 1). Moreover, several proteins associated to ROS production and signalling were modulated at this early time-point of infection (Table 1). Lastly, plant proteins involved in the disruption of oomycete structures were also identified (Table 1).

Table 1. ‘Regent’ APF proteins up accumulated at 6hpi with *P. viticola*, involved in key cellular pathways.

NCBI accession	NCBI description	Biological process	log2(FC)
Modulation of plant physical barriers			
XP_002281842.1	ABC transporter G family member 32 (ABCG32)	stress response	19,00
XP_002263127.1	fatty acyl-CoA reductase 3-like (FAR3)	lipid metabolism	18,83
NP_001268091.1	pectinesterase/pectinesterase inhibitor PPE8B-like (PPE8B)	cell wall modification	15,93
XP_002277293.4	pectinesterase	cell wall metabolism	-1,80
Modulation of plant plasma membrane proteins in response to infection			
XP_019071787.1	disease resistance protein RPV1-like isoform X1 (RPV1)	stress response	19,12
XP_019073586.1	disease resistance protein RUN1-like isoform X1 (RUN1)	stress response	17,99
XP_010644327.1	disease resistance protein At4g27190-like	stress response	4,69
XP_002280315.3	probable disease resistance protein At1g61300	stress response	18,99
XP_019077695.1	probable disease resistance protein At1g61300	stress response	4,47
XP_010654733.1	probable disease resistance protein At5g63020	stress response	21,94
XP_010645387.1	TMV resistance protein N	stress response	21,70
XP_019078946.1	TMV resistance protein N	stress response	17,22
XP_019075299.1	leucine-rich repeat receptor protein kinase MSP1-like	signal transduction	16,43
XP_002267269.1	probable leucine-rich repeat receptor-like protein kinase At1g35710	signal transduction	16,39
XP_002282474.2	serine-threonine protein kinase, plant-type, putative	signal transduction	18,49
XP_010660578.1	receptor-like protein kinase FERONIA (FERONIA)	stress response	18,38
XP_010664467.1	ATPase 9, plasma membrane-type	transport	19,39
XP_002279498.1	putative calcium-transporting ATPase 13, plasma membrane-type (Ca ²⁺ -ATPase)	transport	15,64
XP_002283826.1	protein unc-13 homolog	stomatal regulation	18,55
XP_010647591.2	ankyrin repeat-containing protein ITN1-like isoform X6 (ITN1)	stress response	16,63
Activation of auxin signalling			
XP_002274153.1	protein kinase PINOID	hormone signalling	22,45
XP_002277611.1	protein NDL2	hormone signalling	16,72
XP_002265864.1	ubiquitin-NEDD8-like protein RUB2	protein metabolism	16,62
Regulation of ROS during plant defence response			
XP_002269918.1	peroxidase 4-like	redox	21,05
XP_002274392.1	purple acid phosphatase (PAP)	energetic metabolism	20,55
XP_002285473.1	xanthine dehydrogenase 1-like isoform X1 (XDH1)	redox	16,51
Disruption of oomycete structures			
XP_010664681.1	glucan endo-1,3-beta-D-glucosidase	stress response	19,85
XP_002283647.1	glucan endo-1,3-beta-glucosidase	stress response	15,09
XP_002268991.2	GDSL esterase/lipase (GELP)	stress response	15,72
XP_002276525.1	GDSL esterase/lipase At4g01130 (GELP)	stress response	6,00

2.2 'Regent' whole leaf proteome modulation at 6h post *P. viticola* infection

Whole leaf proteome of 'Regent' during *P. viticola* infection time-course was obtained in 2020¹³. Raw data deposited on the Pride database of the 6hpi was re-analysed following the same pipeline that was implemented for the APF proteome analysis. Hundred and fifty-two DAPs were identified. Of those, 69 proteins were up accumulated and 83 were down accumulated. These proteins were mainly related to stress response, energy and secondary metabolisms and translation (Fig.3). At 6hpi, *P. viticola* infection induces a modulation in the abundance of several stress-related proteins, like heat shock proteins, cysteine proteinases and glucanases. In addition, a great number of photosynthesis-related proteins are down accumulated in 'Regent' leaves in response to infection. Translation and signal transduction-related proteins, like ribosomal proteins and serine/threonine protein kinase (RSTK), respectively, were up accumulated after *P. viticola* infection.

Fig.3. Biological process annotation of the differentially accumulated 'Regent' whole leaf proteins, 6 hours post inoculation with *P. viticola*. Dark grey bars: percentage of down accumulated proteins; Light grey bars: percentage of up accumulated proteins.

The functional annotation of several of the proteins identified in whole leaf are closely related with the pathways that were found to be modulated in APF proteins (Table 2). Indeed, we have identified whole leaf proteins involved in the modulation of plant physical barriers and activation of plant defence signalling through plasma membrane receptors, regulation of ROS levels, and

disruption of oomycete structures (Table 2). Also, proteins associated with calcium signalling, and intracellular trafficking vesicles were modulated in the whole leaf context (Table 2).

Table 2. Regent' whole proteins up accumulated at 6hpi with *P. viticola*, involved in key cellular pathways.

NCBI accession	NCBI description	Biological process	log ₂ (FC)
Modulation of plant physical barriers and plasma membrane receptors in response to infection			
CAN83165.1	pectinesterase inhibitor 9 (PMEI9)	cell wall metabolism	2,39
CBI32865.3	alpha-L-arabinofuranosidase 1 (ASD1)	cell wall metabolism	-18,97
XP_002267434.3	serine/threonine protein kinase (RSTK)	signal transduction	6,90
XP_002265462.3	serine/threonine-protein kinase pakA (RSTK)	signal transduction	17,31
Regulation of ROS during plant defence response			
CBI31928.3	peroxisome biogenesis protein 19-2-like (PEX19-2)	protein metabolism	19,28
CBI32544.3	protein TIC 62, chloroplastic isoform X1	transport	18,74
CAN66554.1	succinate dehydrogenase assembly factor 4, mitochondrial (SDHAF4)	energetic metabolism	17,63
XP_002283860.1	15.7 kDa heat shock protein, peroxisomal	stress response	20,89
CAN67665.1	17.3 kDa class II heat shock protein-like	stress response	6,61
XP_003634522.1	peroxidase 12	redox	-21,36
XP_002285652.2	peroxidase A2-like	redox	-8,08
XP_010647098.1	polyphenol oxidase, chloroplastic-like	redox	-20,80
CBM39273.1	18.2 kDa class I heat shock protein	stress response	-20,66
CBI30632.3	28 kDa heat- and acid-stable phosphoprotein	stress response	-19,75
CBM39216.1	class I heat shock protein	stress response	-21,56
CBI23075.3	small heat shock protein, chloroplastic	stress response	-21,87
Disruption of oomycete structures			
CBI32343.3	endo-1,3;1,4-beta-D-glucanase-like	stress response	-21,33
CBI26171.3	endo-1,3;1,4-beta-D-glucanase-like isoform X3	stress response	-19,29
CAN71820.1	glucan endo-1,3-beta-glucosidase	stress response	-20,49
CBI36040.3	profilin 1	cell regulation	20,96
Modulation of calcium signalling			
CBI15387.3	calcium sensing receptor, chloroplastic	stomatal regulation	-19,74
CBI24493.3	CDGSH iron-sulfur domain-containing protein NEET	secondary metabolism	-7,88
Increase of protein trafficking			
CBI28256.3	putative clathrin assembly protein At5g35200	stress response	21,36

2.3 *P. viticola* proteome during infection establishment

We have sequenced for the first time the *P. viticola* proteome, obtained from grapevine leaves apoplast at 6hpi. Sixty proteins were identified being mainly involved in two biological processes: growth/morphogenesis (e.g. cell division cycle 5 and β -glucan synthesis-associated SKN1) and virulence (RxLR proteins and serine protease trypsin's). Proteins involved in signalling processes like agc kinase (ACG), serine threonine kinase and small GTP-binding Rab28 were also identified (Table 3).

Table 3. *P. viticola* proteins, identified in ‘Regent’ apoplast after 6hpi, involved in virulence and growth/morphogenesis mechanisms.

Protein name	Protein code (INRA Database)
Proteins involved in virulence mechanisms	
Coproporphyrinogen III oxidase (CPOX)	PVIT_0003600.T1
RxLR-like protein (RxLR)	PVIT_0014146.T1
RxLR-like protein (RxLR)	PVIT_0014142.T1
Serine protease trypsin	PVIT_0011817.T1
Serine protease trypsin	PVIT_0011837.T1
Serine threonine kinase	PVIT_0018302.T1
Tetratricopeptide repeat 26	PVIT_0013696.T1
TKL kinase (TKL)	PVIT_0016228.T1
Proteins involved in growth/morphogenesis mechanisms	
Beta-glucan synthesis-associated SKN1 (SKN1)	PVIT_0022780.T1
CAMKK kinase (CaMK)	PVIT_0013015.T1
Cell division cycle 5 (CDC5)	PVIT_0005546.T1
CMGC CDK kinase (CDK)	PVIT_0011642.T1
CMGC MAPK kinase (MAPK)	PVIT_0009065.T1
FAD synthase-like	PVIT_0001050.T1
Serine threonine kinase	PVIT_0018302.T1
Proteins involved in both mechanisms	
AGC kinase (ACG)	PVIT_0009891.T1
Calpain-like protease	PVIT_0019872.T1
Small GTP-binding Rab28	PVIT_0005551.T1

3. Discussion

3.1 *P. viticola* leads to a broad modulation of ‘Regent’ APF and whole leaf proteomes

During grapevine-*P. viticola* interaction, the apoplast compartment is the first hub where plant and pathogen secretomes meet. Several proteins are crucial for the outcome of the interaction, both from the host or pathogen sides. In the apoplast, processes involving pathogen recognition through membrane receptors that activate signal transduction pathways for expression of host defence-associated genes or proteins that directly communicate with pathogen molecules inhibiting infection progress are essential. Considering the whole leaf tissue, trafficking of several proteins to respond to the plant defence requirements must be activated as well as processes that lead to a broad activation of defence-related mechanisms. Thus, communication between the apoplast and the host intracellular organelles is essential for a concerted and quick defence response against the pathogen. Moreover, during the interaction, *P. viticola* develops its infection structures, namely hyphae culminating in plant cell invasion and development of haustorium for feeding. In the first hours of interaction, host and pathogen communications are expected to be very dynamic and to define the outcome of the interaction.

3.1.1 The dual battle at the gate: host strengthens its physical barriers while the pathogen triggers plant cell wall degradation

The cuticle is a barrier coating the outer surface of epidermal cells of organs of the aerial parts of the plants. It protects against water loss and various abiotic and biotic stresses¹⁸. In 'Regent' APF, two cuticle-related proteins were found to be up accumulated after *P. viticola* infection, the ABC transporter G family member 32 (ABCG32) and the fatty acyl-CoA reductase 3-like protein (FAR3), (Table 1; Fig.4). These ABC transporters have been frequently shown to be involved in pathogen response, surface lipid deposition and transport of the phytohormones auxin and abscisic acid^{19,20}. In Arabidopsis, the ABCG32 was reported to be involved in cuticle formation, most likely by exporting cutin precursors from the epidermal cell²¹. The fatty acyl-CoA reductase 3-like protein is involved in cuticular wax biosynthesis²². In incompatible grapevine-*P. viticola* interaction, such as the one that occurs in 'Regent', a higher abundance of these proteins leads to the hypothesis that the host activates processes to promote the strengthening the cuticular barrier in order restrain pathogen penetration (Fig.4).

Also, plants have developed a system for sensing pathogens and monitoring the cell wall integrity, upon which they activate defence responses that lead to a dynamic cell wall remodelling required to prevent pathogen progression. Plant cell wall-associated proteins were found to be differentially modulated in 'Regent' APF in response to *P. viticola*. A down accumulation of a pectinesterase, a protein involved in plant cell wall degradation as well as an up accumulation of pectinesterase/pectinesterase inhibitor PPE8B-like protein (PPE8B), (Table 1; Fig.4). This protein is involved in plant cell wall reconstruction and, in cotton, genes encoding this type of protein are specifically up regulated in plant resistant variety upon *Aspergillus tubingensis* infection²³.

In the whole leaf proteome of 'Regent', the abundance of pectinesterase inhibitor 9 (PMEI9) increased, a protein involved in resistance to pathogens²⁴, and a decrease in the accumulation of alpha-L-arabinofuranosidase 1, a protein involved in cell wall degradation²⁵, was detected (Table 2; Fig.4). The whole leaf and apoplast proteomes are modulated as a defence strategy to prevent cell wall degradation, maintaining its integrity and thus inhibiting the entry of the oomycete. On the other hand, proteins involved in plant cell wall degradation were found in *P. viticola* proteome (Table 3; Fig.4), namely: coproporphyrinogen III oxidase, which is a peroxidase with the ability to degrade lignin, one of the components of plant cell walls²⁶; trypsins, a serine proteases family, also identified in the secretomes of several fungus²⁷, and linked to pathogenicity against plant hosts²⁸. In *P. viticola*, the two identified trypsins were predicted to be apoplastic effectors and so it is expected a direct interaction with plant molecules for cell wall degradation.

During plant-pathogen interactions, plant cell wall is a dynamic structure that functions as a barrier that pathogens need to breach to colonize the plant tissue. Biotrophic pathogens, like *P. viticola*, require a localized and controlled degradation of the cell wall to keep the host cells alive during feeding. The regulation of the abundance of these cell wall-related proteins in the apoplast and whole leaf of 'Regent' suggests an adaptation of the grapevine proteome to prevent cell wall disruption while *P. viticola* secretes proteins that degrade de cell wall to invade the plant cell.

Fig.4. Triggering of host cell wall degradation by *P. viticola* through CPOX and trypsin proteins secretion while in the host several proteins associated to the strengthening of the physical barriers (cuticle – eg ABCG32, FAR3; cell wall - eg PME19, PPE8B) are positively modulated. At the host also a negative modulation of proteins involved in cell wall degradation is promoted, moreover, proteins as 1,3-β-glucanases are translocated to the APF for pathogen cell wall degradation. Proteins represented in red are positively modulated, proteins represented in blue are negatively modulated (compared to mock-inoculated control). Pathogen proteins are represented in grey. 1,3-β-glucanase – includes endo-1,3;1,4-beta-D-glucanase-like, endo-1,3;1,4-beta-D-glucanase-like isoform X3 and glucan endo-1,3-beta-glucosidase; ABCG32 - ABC transporter G family member 32; ASD1 - alpha-L-arabinofuranosidase 1; CDC5 - cell division cycle 5; CPOX - coproporphyrinogen III oxidase; CW - cell wall; GELP - GDSL esterase/lipase; FAR3 - fatty acyl-CoA reductase 3-like protein; PME19 - pectinesterase inhibitor 9; PPE8B - pectinesterase/pectinesterase inhibitor PPE8B-like; SKN1 - beta-glucan synthesis-associated SKN1.

3.1.2 *P. viticola* recognition and signalling events are established as soon as 6hpi

The first layer of plant defence relies on the recognition of conserved microbe-associated molecular patterns (MAMPs) by the so-called pattern recognition receptors (PRRs). PRRs are generally plasma membrane receptors which are often coupled to intracellular kinase domains²⁹. The second layer of plant immunity depends on the ability of the plant to recognize the pathogen effectors, like RxLRs, by disease resistance proteins (R) and trigger a robust resistance response³⁰. Several states such as oxidative burst, cell wall strengthening, induction of defence gene expression, and rapid cell death at the site of infection (hypersensitive response) occur in downstream cellular events leading to the establishment of an incompatible interaction³¹.

These local hypersensitive responses can trigger long-lasting systemic responses (systemic acquired resistance (SAR)) that prime the plant for resistance against a broad spectrum of pathogens^{32,33}.

We have identified several virulence-related proteins that are secreted by *P. viticola* or that are present in oomycete infection structures (Table 3; Fig.5). RxLR effectors were detected in *P. viticola* proteome as soon as 6hpi and were predicted to be secreted to the apoplast. These proteins are key players in virulence for downy mildew species³⁴ since they are known to defeat plant immune responses through many routes, which include reprogramming host gene expression, altering RNA metabolism, and binding to host proteins involved in signalling³⁵. On the host side, eight R-proteins were up-accumulated in the APF after *P. viticola* infection (Table 1; Fig.5), including RPY1-like isoform X1 and RUN1-like isoform X1, which confer resistance to multiple downy and powdery mildews, respectively, by promoting cell death^{36–38}. Also, an up accumulation of several RSTK was observed both in APF (Table 1) and in whole leaf proteome (Table 2). These receptors are involved in a wide array of processes ranging from developmental regulation to disease resistance, including activation of signal transduction for plant defence response initiation^{39,40}. One of the identified receptors in APF is the FERONIA (Fig.5). FERONIA is a plant recognition receptor kinase which plays a significant role in plant immune system. In *Catharanthus roseus*, FERONIA acts as a sensor of cell wall integrity challenged by the host-pathogen interaction and further triggers downstream immune responses in the host cell. In response to changes in the cell wall, FERONIA induces the release of ROS (reviewed in⁴¹). Furthermore, in spinach response to infection with the oomycete *Peronospora effusa*, an up-regulation of the genes encoding for the FERONIA and a LRR receptor-like RSTK was observed⁴². These results highlight the importance of these receptors in the plant cell membrane and in the first line of defence in plant strategies to stop oomycete proliferation.

Moreover, an up accumulation of a H⁺-ATPase (ATPase 9) was also detected in 'Regent' APF at 6hpi. Plasma membrane H⁺-ATPases maintain low cytoplasmic concentrations of H⁺ and are dynamically regulated by biotic and abiotic events, being involved in signalling mechanisms during plant-pathogen interaction (reviewed in⁴³). In response to internal and/or external cues, the trafficking of these plant pumps, to and from the plasma membrane, is also highly regulated. Indeed, we have identified an up accumulation of a protein involved in H⁺-ATPases translocation for the plasma membrane, the protein unc-13 homolog (Table 1). In Arabidopsis, a protein unc-13 homolog (PATROL1) controls the translocation of a H⁺-ATPase to the plasma membrane during stomata opening⁴⁴. PATROL1 resides in the endosome and moves to and from the plasma membrane in response to environmental stimuli. When the stomata open, intracellular vesicles incorporate the plasma membrane and PATROL1 may be carried with them to tether the H⁺-ATPase into the plasma membrane⁴⁴. Modifications of the H⁺-ATPases concentration in the plasma membrane act as an alternative mean to control stomata opening⁴⁵ and in the case of grapevine-*P. viticola* interaction, the stomata are the entry sites of the oomycete. Despite the role of these ATPases in grapevine remains to be elucidated, an up accumulation of ATPase 9 and protein unc-13 homolog could suggest its involvement in stomata opening regulation during 'Regent'-*P. viticola* interaction.

During 'Regent' response to *P. viticola* infection, the up accumulation of another transmembrane protein (TMP), ankyrin repeat-containing protein ITN1-like isoform X6 (ITN1),

was also detected (Table 1; Fig.5). Plant TMPs are essential for normal cellular homeostasis, nutrient exchange, and responses to environmental cues⁴⁶. ITN1 has been implicated in diverse cellular processes such as signal transduction and, in Arabidopsis, this protein was proposed to be related with abscisic acid signalling pathway and promotion of ROS production⁴⁷.

Fig.5. First communication events between host and pathogen: pathogen effectors are recognized by host R proteins leading to a broader remodulation of proteins related to ROS production, signal transduction and immune response establishment. Also, *P. viticola* is able to perceive host molecules activating signalling transduction pathways. Proteins represented in red are positively modulated, proteins represented in blue are negatively modulated (compared to mock-inoculated control). Pathogen proteins are represented in grey. AGC - AGC kinase; Ca²⁺-ATPase - putative calcium-transporting ATPase 13; CaMK - CAMKK kinase; CDK - CMGC CDK kinase; CW - cell wall; FERONIA - receptor-like protein kinase FERONIA; ITN1 - ankyrin repeat-containing protein ITN1-like isoform X6; MAPK - CMGC MAPK kinase; PAP - purple acid phosphatase; PEX19-2 - peroxisome biogenesis protein 19-2-like; RSTK - serine/threonine protein kinase; ROS - reactive oxygen species; RxLR - RxLR-like proteins; R-proteins - includes RPV1-like isoform X1 and RUN1-like isoform X1 and RSTKs; SDHAF4 - succinate dehydrogenase assembly factor 4; TIC62 - protein TIC 62; TKL - TKL kinase; XDH1 - xanthine dehydrogenase 1.

3.1.3 Auxin signalling plays an important role in apoplast

Auxin is a plant growth hormone that plays a role in many aspects of growth and development, and its signalling have been also highly associated to plant stress responses^{48,49}. Several auxin

signalling-related proteins were up accumulated in APF after infection (Table 1): PINOID (a RSTK involved in the regulation of auxin signalling, being a positive regulator of cellular auxin efflux⁴⁸); N-MYC DOWNREGULATED-LIKE2 (NDL2) protein (involved in auxin signalling through the positive regulation of auxin transporter protein 1 (AUX1) a carrier protein responsible for proton-driven auxin influx⁵⁰); and NEDD8 (neural precursor cell expressed developmentally down-regulated 8), a post-translational modifiers of cullin proteins⁵¹. Cullin1 is one of the proteins that forms the “SKP1 cullin F-box” (SCF) ubiquitin ligase complex, that is involved in auxin signalling pathway⁵². Auxin homeostasis could be perturbed by stress-induced changes that affects the auxin efflux carriers and that modify the apoplastic pH disturbing the auxin uptake and distribution^{53,54}. Indeed, in plant-pathogen interactions, auxin was shown to play a dual role both in regulating plant defence and in bacterial virulence⁵⁵. It was shown that some pathogens may promote host auxin accumulation which leads to the suppression of salicylic acid (SA)-mediated host defences. On the other side, host auxin accumulation also modulated expression of particular pathogen virulence genes⁵⁵. Despite the fact that none of the *P. viticola* effectors that we have found to be present in the APF was previously related to host auxin metabolism modulation, we may hypothesize that the modulation of auxin-related host proteins detected may reflect pathogen induced reprogramming. In fact, *P. viticola* being a biotroph, modulation of auxin metabolism leading to a depletion of SA-host defenses could be an infection strategy.

3.1.4 Apoplastic ROS signalling in grapevine immunity

ROS play an important role in pathogen resistance by directly strengthening host cell walls via cross-linking of glycoproteins, promoting lipid peroxidation and activation of ROS-signalling networks^{56–58}. We have previously shown that ROS accumulation occurs in ‘Regent’ at 6hpi with *P. viticola*^{13,59}. In the APF proteome, proteins involved in ROS production, like peroxidase 4-like and purple acid phosphatase (PAP), are up-accumulated in response to the infection (Table 1; Fig.5), which is in accordance with our previous results. In Arabidopsis, PAP5 is induced in the early stages (6hpi) of *Pseudomonas syringae* infection and is involved in ROS generation⁶⁰. Apoplastic ROS are very important in plant development and responses to stress conditions, being involved in the activation of signal transduction from extracellular spaces to the cell interior and may directly eliminate invading pathogens (reviewed in^{61–63}). Small amounts of ROS lead to expression of stress-responsible genes as a plant resistance mechanism. However, high levels of ROS during a long period of time could culminate in damage of plant molecules and consequently in cell death. A strict regulation of ROS levels is thus important to induce a resistance response by the plant without promoting cell injury. In ‘Regent’ APF after infection with *P. viticola*, an up accumulation of xanthine dehydrogenase 1 (XDH1)-like isoform X1 was identified (Table 1; Fig.5). In Arabidopsis, this protein has dual and opposing roles in the ROS metabolism, contributing to H₂O₂ production in epidermal cells to fight pathogen haustoria and producing uric acid to scavenge chloroplast H₂O₂ in mesophyll cells to minimize oxidative damage⁶⁴. In grapevine, the role of this protein was not been elucidated yet. However, in our results, an up accumulation of this ROS-related protein in ‘Regent’ APF after infection was detected, suggesting a possible involvement of XDH1 in plant defence mechanism through ROS metabolism regulation.

In whole leaf proteome, an up accumulation of ROS-related proteins such as peroxisome biogenesis protein 19-2-like (PEX19-2) ^{13,65,66}, protein TIC 62 (TIC62) ⁶⁷ and succinate dehydrogenase assembly factor 4 (SDHAF4) ⁶⁸ was observed at the same time that proteins like peroxidases and polyphenol oxidase ⁶⁹ were less accumulated after infection (Table 2; Fig.5). Also, several heat shock proteins with different abundance levels in infected leaves, when compared to non-infected leaves, were identified (Table 2; ⁷⁰). As we already mentioned, the regulation of ROS metabolism during plant-pathogen interaction presents a complex dynamic. ROS are generated to overcome the infection but at the same time a tight regulation of high ROS levels is needed to protect the plant from oxidative stress (reviewed in ⁶¹⁻⁶³). This regulation is clearly evident in 'Regent' whole leaf and APF through the presence of several ROS-related proteins at different abundance levels, highlighting the importance of ROS in the 'Regent' defence mechanism against *P. viticola* as previously reported ^{13,59}.

3.1.5 Calcium related signalling is prevalent in the APF at 6hpi

In grapevine-*P. viticola* interaction, the increase in calcium leaf concentration in response to infection was already reported ⁷¹. In 'Regent' APF proteome we have found an up accumulation of a putative calcium-transporting ATPase 13 (Ca²⁺-ATPase), (Table 1; Fig.5). Ca²⁺-ATPases play critical roles in sensing calcium fluctuations and relaying downstream signals by activating definitive targets, thus modulating corresponding metabolic pathways (reviewed in ⁷²). We have also detected an up accumulation of PINOID (Table 1). In Arabidopsis, PINOID activity is regulated by several calcium binding proteins, like a calmodulin-like protein, and the binding of these proteins to PINOID is enhanced by calcium ⁷³.

In *P. viticola* proteome, we have detected a Ca²⁺/calmodulin-dependent protein kinase (CaMK) that responds to high levels of calcium ⁷⁴, (Table 3; Fig.5). In pathogens, CaMKs are involved in several pathogenicity-related cellular mechanism. For example, in *C. albicans*, CaMKs functions in cell wall integrity and cellular redox regulation ⁷⁵; in *Neurospora*, a genus of Ascomycete fungi, CaMKs are related to growth and development of the pathogens ⁷⁶; and in *M. oryzae*, conidial germination and appressorial formation were delayed and virulence was attenuated in mutants of a CaMK ⁷⁷. However, in *P. viticola*, up to our knowledge, there is no information about the role of these type of kinases in oomycete development and/or pathogenicity.

At the same time, in 'Regent' whole leaf proteome, calcium-related proteins, like calcium sensing receptor ⁷⁸ and CDGSH iron-sulfur domain-containing protein NEET ⁷⁹, are less accumulated after infection (Table 2).

These results suggest that, at such an early stage of the infection such as 6hpi, this calcium-associated response and regulation, as consequence of high levels of calcium in the infection site, is only taking place in the apoplast, which is the first contact point between plant and pathogen. As such, this regulation of the abundance of proteins associated with calcium metabolism in the leaf tissue is not evident.

3.1.6 Activation of enzymes to disrupt oomycete structures

One of the plant defence mechanisms during pathogen infection is the secretion of proteins involved in the degradation of pathogen structures to inhibit its growth and thus stop its proliferation. Oomycete cell walls consist mainly of β -1,3-glucans, β -1,6-glucans and cellulose rather than chitin, essential constituent of fungal cell walls⁸⁰.

In grapevine APF after *P. viticola* infection an up accumulation of two glucan endo-1,3-beta-D-glucosidases and two GDSL esterase/lipases (GELPs) was observed (Table 1; Fig.4). The first are involved in the degradation of the polysaccharides of the pathogen cell wall and the GELPs possess lipase and antimicrobial activities that directly disrupt pathogen spore integrity⁸¹. In contrast, in whole leaf proteome, proteins like endo-1,3;1,4-beta-D-glucanase-like, endo-1,3;1,4-beta-D-glucanase-like isoform X3 and glucan endo-1,3-beta-glucosidase, are less abundant after infection when compared to the non-infected leaves (Table 2; Fig.4). These results suggest that a demobilization of these proteins from the inside of the cell to the APF might be occurring in response to *P. viticola* infection. The accumulation of these proteins in 'Regent' APF leads to a disruption of oomycete structures as defence mechanism to inhibit the infection progress.

Moreover, a protein with antifungal activity, profilin 1, was found to be accumulated in 'Regent' whole leaf after *P. viticola* infection (Table 2). In Arabidopsis, this protein showed significant intracellular accumulation and cell-binding affinity for fungal cells, being capable to penetrate the fungal cell wall and membrane and act as inhibitor of fungal growth through ROS generation⁸².

3.1.7 The importance of protein trafficking during grapevine-*P. viticola* interaction

During plant-pathogen interaction, protein trafficking is very important for plant cells to quickly respond to pathogen infection. This trafficking occurs through the secretory and endocytic pathways that involves a complex set of proteins associated to vesicle formation, transport, docking, and fusion with the respective target membrane (reviewed in⁸³). Clathrin-mediated endocytosis is the best-known mechanism of endocytosis in plants and involves the generation of small vesicles surrounded by a coat of clathrin and other associated proteins. During interaction of 'Regent' with the oomycete *P. viticola*, we observed an accumulation of clathrin assembly protein in the whole leaf proteome at the first hours of infection (Table 2). Indeed, the increase of proteins involved in the generation of trafficking vesicles, like clathrin assembly protein, reinforces our hypothesis of a relocation of specific proteins within the cell and to the APF as a plant defence mechanism. We have already mentioned the presence of the protein unc-13 homolog in the APF, which is responsible for the translocation of H⁺-ATPases to the plasma membrane⁴⁴ and we have also proposed the translocation of glucanases from the inside of the cell to the APF in response to the infection. These results highlight the massive molecular reprogramming that occurs when a plant is exposed to an environmental stress like pathogen infection.

Even for the pathogen, the molecular trafficking that occurs within the cells is very important for its growth and pathogenicity. In *P. viticola*, we have identified the small GTP-binding Rab28

(Ras homologue from brain), (Table 3). Rab proteins, which constitute the largest family of monomeric GTPases, are small proteins involved in many biological processes (reviewed in ⁸⁴). Members of this family participate in cell regulation, growth, morphogenesis, cell division, and virulence. Also, they are known as master regulators of intracellular bidirectional vesicle transport and, consequently, they localize in ER, vesicles, and multivesicular bodies, as well as in early and late endosomes ⁸⁵. Rabs have been implicated in regulating vesicle motility through interaction with both microtubules and actin filaments of the cytoskeleton ⁸⁶. In fungi, Rab participates in the secretion of metabolites and lytic enzymes and, in *Fusarium graminearum*, Rab GTPases are essential for membrane trafficking-dependent growth and pathogenicity ⁸⁷. However, in *P. viticola* the specific function of Rab proteins was not yet elucidated.

3.1.8 Host and pathogen proteases as hubs during the interaction

Proteases are enzymes that catalyse the breakdown of proteins into smaller polypeptides or single amino acids and play important roles in numerous biochemical processes. Pathogens produce a variety of proteases to degrade host tissue or to disrupt or modify host defence to create suitable conditions for successful colonization (reviewed in ^{88,89}). In *P. viticola*, a calpain-like protease was detected (Table 3) and a possible role in pathogenicity is suggested as previous studies in *M. oryzae* point out that calpains play multiple roles in conidiation, sexual reproduction, cell wall integrity and pathogenicity ⁹⁰. In *Saccharomyces cerevisiae* a calpain is also required for alkaline adaptation and sporulation ⁹¹. Moreover, two trypsins and serine proteases involved in pathogenicity ²⁸ were identified in *P. viticola* proteome during grapevine infection (Table 3).

Host subtilisin-like protease SBT5.3, was found more accumulated in the APF after infection. This accumulation is in accordance with the previously published studies on subtilisin-like proteases in grapevine-*P. viticola* interaction. A 10-fold change in SBT5.3 gene expression was observed also at 6hpi together with the increase of expression of several subtilase genes in 'Regent' ¹⁵. Previously, it has also been hypothesized that subtilases play a crucial role in the establishment of the incompatible interaction between 'Regent' and *P. viticola*. In Arabidopsis, the subtilase SBT3.3 was shown to accumulate in the extracellular matrix after infection and to initiate a downstream immune signalling process ⁹².

Our results highlight that both host and pathogen proteases are essential on the first contact hours and might play an important role in pathogen recognition and in the overcome of host defences.

3.1.9 *P. viticola* proteome at 6hpi reflects actively regulated processes leading to infection development

In *P. viticola*, several kinases are putatively involved in the infection mechanism were also found in the APF at 6hpi (Table 3). We have detected, in the *P. viticola* proteome, a AGC kinase (cAMP-dependent, cGMP-dependent and protein kinase C), (Table 3; Fig.5). This protein family embraces a collection of Ser/Thr kinases that mediate a large number of cellular processes, such as cell growth, response to environmental stresses, and host immunity ⁹³⁻⁹⁶. A study using kinase

inhibitors showed that kinase C protein plays a key role in the signal transduction mechanisms during maintenance of motility of the zoospores, essential during their migration to the stomata⁹⁷. This evidence suggested that AGC kinases might be important pathogenic factors during pathogen infection, leading us to hypothesize that the *P. viticola* AGC kinase might be also relevant for the pathogenicity of this oomycete.

We have also detected a CDK (cyclin-dependent kinase) in *P. viticola* proteome that might be participating in pathogen infection mechanisms through cell polarization for the germ tube formation (Table 3; Fig.5). Indeed, several CDK members have been reported to be pathogenicity-related. In phytopathogenic fungus *U. maydis*, a CDK that is essential for growth and maintenance of cell polarity in this pathogen was identified. *cdk5ts* mutants showed to be drastically less virulent, probably because of the involvement of this protein in the induction of the polar growth required for the infection process⁹⁸. In *P. viticola*, cell polarity guides the emergence and the growth of the germ tube during the infection mechanism, namely the penetration into the stomatal cavity^{99,100}.

A MAPK (Mitogen-Activated Protein Kinase) has also been identified in the *P. viticola* proteome during the interaction with 'Regent' leaves (Table 3; Fig.5). MAPK cascades are very important in numerous cellular mechanisms in pathogens, regulating infection-related morphogenesis, cell wall remodelling, and high osmolarity stress response (reviewed in^{101,102}). For example, in *Peronophythora litchii*, the oomycete pathogen causing litchi downy blight disease, a mutation in PIMAPK10, a MAPKP, led to a reduced mycelial growth rate, less sporulation and weakened pathogenicity, indicating an important function of MAPK signal pathway in oomycete pathogenicity¹⁰³. In *Phytophthora sojae*, another oomycete, the authors showed that the MAPK PsSAK1 controls zoospore development and that it is necessary for pathogenicity¹⁰⁴. In *M. oryzae*, initiation of appressorium formation is controlled by the cyclic AMP-protein kinase A pathway and the appressorium development and invasive growth is regulated by the Pmk1 MAPK pathway^{105,106}.

Tyrosine kinase-like (TKL) was also identified in *P. viticola* proteome after 6 hours of infection (Table 3; Fig.5). The TKL family is present in most eukaryotes and participates in many biological processes, however, information about its role in oomycetes is still scarce. In *P. guiyangense*, TKLs were up-regulated at early infection stages and silencing of TKLs led to reduced mycelia growth, zoospore production and alteration of stress responses. Also, silencing of TKLs resulted in a reduced virulence of *P. guiyangense*¹⁰⁷, suggesting a key role of these kinases in pathogen infection strategies. In *P. infestans* kinome prediction 139 TKLs were identified, however their function is still unknown¹⁰⁸.

Tetratricopeptide repeat (TPR)- and SEL1-containing proteins were also identified in *P. viticola* proteome (Table 3). These have been reported to be directly related to virulence-associated functions in bacterial pathogens^{109–111}. In *Francisella tularensis*, a bacterial pathogen, a TPR-containing protein is a membrane-associated protein that is required for intracellular replication of the microbe, *in vivo* virulence, and heat stress tolerance¹¹². In *P. infestans*, TPR was predicted as one of the most common domains within proteins¹¹³. However, up to our knowledge, the role of TPR-containing proteins in oomycete/fungus pathogenicity is still unknown. Regarding SEL1-containing proteins, in *Candida albicans*, a human fungal pathogen, these proteins are capable of shaping host immune response and the severity of fungal systemic infection and were

suggested as a novel fungus-derived pattern-associated molecular pattern (PAMP)¹¹⁴. However, up to our knowledge, there is no information about pathogen Sel1-containing proteins in plant-oomycete interaction.

Growth-related proteins were also detected in the *P. viticola* proteome sequencing (Table 3; Fig.4). These types of proteins are essential for the development of the pathogen structures during infection and besides they are not directly related to pathogenicity, their presence is an indicator of the pathogen growth and thus the progression of the infection in the host. In *P. viticola*, we have identified the SKN1 protein that is required for synthesis of the β -glucans, the major components of oomycete cell walls¹¹⁵. We have also identified the cell division cycle 5 (Cdc5) protein, a highly conserved nucleic acid binding protein among eukaryotes that plays critical roles in development, however, in pathogens this protein is still poorly characterized¹¹⁶.

4. Conclusions

Plants have developed several defence mechanisms to rapidly respond to pathogen attack. A fast recognition of the pathogen structures and activation of a defence response is primordial for the establishment of the incompatible interaction. Apoplast dynamics becomes an essential part of the battle between plants and pathogens. Here, we reveal for the first time the communication between grapevine extracellular and intracellular spaces at the first hours of interaction with *P. viticola* as well as the pathogen strategies to overcome grapevine defence. Our results highlight several defence mechanisms that are first activated and modulated in grapevine apoplast during infection, namely leading to plant cell wall plasticity to prevent disruption and disturbance of oomycete structures. Several proteins were also identified in the whole leaf proteome of 'Regent' that are closely related with the mechanisms involved in the apoplast modulation during infection, highlighting a tight communication between the APF and cell interior. Moreover, we have shown that *P. viticola* proteome is enriched in virulence-related proteins as a strategy to defeat the plant defence response and in growth-related proteins to develop the infection structures of the *P. viticola*.

The analysis of the plant and oomycete proteins involved in the first hours of the interaction between grapevine and *P. viticola* revealed that both sides are modulating their offense and defence strategies very early on. This reinforces the hypothesis that host-pathogen cross-talk in the first hours of interaction is highly dynamic and processes such as ETI, ETS, PTI and PTS may occur simultaneously.

5. Materials and Methods

5.1 Plant material and inoculation experiments

The tolerant *V. vinifera* cv. 'Regent' (VIVC number 4572) was used in this study. Wood cuttings from 'Regent' were obtained at the Portuguese Grapevine Germplasm Bank⁽¹¹⁷⁾; INIAV – Dois Portos, Portugal) and grown in 2.5 L pots in universal substrate under controlled conditions in a climate chamber at natural day/night rhythm, relative humidity 60% and a photosynthetic photon flux density of 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$.

For plant inoculation, downy mildew symptomatic leaves were harvested at the Portuguese Grapevine Germplasm Bank, sprayed with water and incubated overnight at 22°C in the dark to enhance sporulation. Next day, *P. viticola* sporangia were collected and their vitality was checked by microscopy. Inocula was propagated in the laboratory using the susceptible Müller-Thurgau.

For the experimental assay, 'Regent' abaxial leaf surface was sprayed with an inoculum solution containing 3.5×10^{-5} sporangia/mL. Mock-inoculations with water were also made and used as control. After inoculation, plants were kept in a greenhouse under high humidity conditions and 25°C. The third to fifth fully expanded leaves beneath the shoot apex were harvest at 6 hours post-inoculation for apoplastic fluid extraction.

5.2 Apoplastic fluid extraction

Apoplastic fluid and total soluble protein extraction was performed as described in ³. Briefly, 5 volumes of 0.1 M ammonium acetate in methanol were used to precipitate the proteins overnight at -20°C. Samples were centrifuged at 4000 *g*, during 30 min at -10°C. Recovered pellets were washed (once with 0.1 M ammonium acetate in 100% methanol, twice with 80% (v/v) acetone and twice with 70% (v/v) ethanol), dried and resuspended in 0.03 M Tris-HCl buffer (pH 8.8) solution containing, 7 M urea, 2 M thiourea, 4% (w/v) CHAPS ³. For cytoplasmic content contamination control, malate dehydrogenase assay was used according to ³.

5.3 MS-Based protein identification

Separation and MS-based identification of proteins was performed as described in ³. For identification of *V. vinifera* and *P. viticola* proteins, the genome assembly of *Vitis*12X database (GCF-000003745.3; 41 208 sequences, July 2009) and *Plasmopara viticola* genome database (INRA-PV221 isolate; 15 960 sequences, April 2018), respectively, were used via Mascot Daemon (v.2.6.0. Matrix Science), imported to Progenesis QIP and matched to peptide spectra. The Mascot research parameters were: a peptide tolerance of 20 ppm, a fragment mass tolerance of 0.3 Da, carbamidomethylation of cysteine as fixed modification and oxidation of methionine, N-terminal protein acetylation and tryptophan to kynurenine as variable modifications. Only the proteins identified with a significance MASCOT-calculated threshold P-value < 0.05, at least two significant peptides per proteins and one unique peptide per proteins were accepted.

For identification of *P. viticola* proteins present in apoplast of 'Regent' leaves inoculated with the oomycete, only the sequenced proteins that fulfilled the following two criteria at the same time were considered: be present in at least 2 of the biological replicates in the inoculated samples; and present in only 1 biological replicate or totally absent in control samples. Functional information about *P. viticola* proteins was obtained from *P. viticola* genome database. Protein secretion and effector function were predicted using TargetP 2.0 (<https://services.healthtech.dtu.dk/service.php?TargetP-2.0>; ¹¹⁸) and EffectorP (<http://effectorp.csiro.au/>; ¹¹⁹), respectively.

5.4 Whole leaf proteome data

For 'Regent' whole leaf proteome analysis, 6hpi with *P.viticola*, an already published dataset deposited on the ProteomeXchange Consortium via the PRIDE partner repository with the identifier PXD021613 was used¹³.

5.5 Statistical analysis of APF and whole leaf proteomes

Principal Component Analysis (PCA) of 'Regent' APF proteome at 6hpi with *P. viticola* was performed using the program MetaboAnalyst 5.0 (<http://www.metaboanalyst.ca/>,¹²⁰).

For the statistical analysis and consequent identification of differentially accumulated proteins, both whole leaf and APF proteomes of 'Regent' leaves inoculated with *P. viticola* (6hpi) were matched to the respective control samples, allowing a comparative analysis between datasets. This was done by applying Rank Products (RP), a powerful statistical test designed for identifying differentially expressed genes in microarrays experiments¹²¹, nevertheless, it also provides a simple and straightforward tool to determine the significance of observed changes in other omics data. RP procedure makes weak assumptions about the data and provides a strong performance with very small data sets, allowing for the control of the false discovery rate (FDR) and familywise error rate (probability of a type I error) in the multiple testing situation.

5.5 Subcellular location prediction of APF DAPs

For the APF DAPs from grapevine, a subcellular localization prediction was performed using SignalP 5.0, TargetP 2.0 and SecretomeP 2.0 servers (<http://www.cbs.dtu.dk/services/>,^{122–124}), ApoplastP (<http://apoplastp.csiro.au/>,¹²⁵), BUSCA (<http://busca.biocomp.unibo.it/>,¹²⁶), PredSL (<http://aias.biol.uoa.gr/PredSL/>,¹²⁷), Mercator (<https://www.plabipd.de/portal/mercator-sequence-annotation>,¹²⁸) and Blast2GO (version 5.2.5, <https://www.blast2go.com/>,¹²⁹). The default parameters were used for all the programs.

Based on the subcellular localization prediction, the APF proteins were grouped in 4 different classes according to the following criteria (as described in³): (1) proteins with a predicted signal peptide (SP) by SignalP (Class I); (2) proteins predicted to be secreted through classical secretory pathways but, by other software than SignalP 5.0 (Class II); (3) proteins predicted to be secreted by unconventional secretory pathways (USP) based on SecretomeP (Class III), and proteins with no predicted secretion (Class IV). Only the proteins belonging to the Class I, II and III were considered for further functional analysis.

5.6 APF and whole leaf proteome data functional analysis

Functional annotation based on Gene Ontology annotation (Biological Process) using Blast2GO software was performed.

Data Availability: The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD030508 and 10.6019/PXD030508.

Acknowledgments: Portuguese Foundation for Science and Technology (FCT, Portugal) funded Joana Figueiredo fellowship (SFRH/BD/137066/2018). FCT funded the Research Units and project: BioISI (UIDB/04046/2020 and UIDP/04046/2020), LEAF (UID/AGR/04129/2019) and CEAUL (UIDB/00006/2020), and the project PTDC/BIA-BQM/28539/2017. We thank Dr. Ana Rita Cavaco, Dr. Marisa Maia and Dr. Maria do Céu Silva for the support in the inoculation assay and APF extraction protocol.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Author Contributions: J.F. and A.F. conceived the study; J.F., R.B.S, L.G.G. and A.F. were responsible for the plant material and performed the APF extraction method; J.F. prepared the samples for nanoLS-MS/MS analysis; C.C.L. and J.R. performed the proteome profiling by nanoLC-MS/MS; L.S. performed the statistical analysis; J.F. and A.F. analysed the data and wrote the manuscript. All authors have read and agreed to the published version of the manuscript.

References

- 1 Jones JDG, Dangl JL. The plant immune system. *Nature* 2006; **444**: 323–329.
- 2 Delaunoy B, Colby T, Belloy N *et al.* Large-scale proteomic analysis of the grapevine leaf apoplastic fluid reveals mainly stress-related proteins and cell wall modifying enzymes. *BMC Plant Biology* 2013; **13**: 24.
- 3 Figueiredo J, Cavaco AR, Guerra-Guimarães L *et al.* An apoplastic fluid extraction method for the characterization of grapevine leaves proteome and metabolome from a single sample. *Physiologia Plantarum* 2021; **171**: 343–357.
- 4 Pechanova O, Hsu C-Y, Adams JP *et al.* Apoplast proteome reveals that extracellular matrix contributes to multistress response in poplar. *BMC Genomics* 2010; **11**: 674.
- 5 Dani V, Simon WJ, Duranti M, Croy RRD. Changes in the tobacco leaf apoplast proteome in response to salt stress. *PROTEOMICS* 2005; **5**: 737–745.
- 6 Fecht-Christoffers MM, Braun H-P, Lemaitre-Guillier C, VanDorsselaer A, Horst WJ. Effect of manganese toxicity on the proteome of the leaf apoplast in cowpea. *Plant Physiology* 2003; **133**: 1935–1946.
- 7 Zhou L, Bokhari SA, Dong C-J, Liu J-Y. Comparative proteomics analysis of the root apoplasts of rice seedlings in response to hydrogen peroxide. *PLOS ONE* 2011; **6**: e16723.

- 694 8 Guerra-Guimarães L, Tenente R, Pinheiro C *et al.* Proteomic analysis of apoplastic fluid of
695 *Coffea arabica* leaves highlights novel biomarkers for resistance against *Hemileia vastatrix*.
696 *Frontiers in Plant Science* 2015; **6**. doi:10.3389/fpls.2015.00478.
- 697 9 Guerra-Guimarães L, Vieira A, Chaves I *et al.* Effect of greenhouse conditions on the leaf
698 apoplastic proteome of *Coffea arabica* plants. *Journal of Proteomics* 2014; **104**: 128–139.
- 699 10 Floerl S, Majcherczyk A, Possienke M *et al.* *Verticillium longisporum* infection affects the
700 leaf apoplastic proteome, metabolome, and cell wall properties in *Arabidopsis thaliana*.
701 *PLOS ONE* 2012; **7**: e31435.
- 702 11 Yu Y, Zhang Y, Yin L, Lu J. The mode of host resistance to *Plasmopara viticola* infection of
703 grapevines. *Phytopathology* 2012; **102**: 1094–1101.
- 704 12 Welter LJ, Göktürk-Baydar N, Akkurt M *et al.* Genetic mapping and localization of
705 quantitative trait loci affecting fungal disease resistance and leaf morphology in grapevine
706 (*Vitis vinifera* L). *Mol Breeding* 2007; **20**: 359–374.
- 707 13 B. Santos R, Nascimento R, V. Coelho A, Figueiredo A. Grapevine–downy Mildew
708 rendezvous: proteome analysis of the first hours of an incompatible interaction. *Plants*
709 2020; **9**: 1498.
- 710 14 Guerreiro A, Figueiredo J, Sousa Silva M, Figueiredo A. Linking jasmonic acid to grapevine
711 resistance against the biotrophic oomycete *Plasmopara viticola*. *Front Plant Sci* 2016; **7**:
712 565.
- 713 15 Figueiredo J, Costa GJ, Maia M *et al.* Revisiting *Vitis vinifera* subtilase gene family: a
714 possible role in grapevine resistance against *Plasmopara viticola*. *Front Plant Sci* 2016; **7**:
715 1783.
- 716 16 Cavaco AR, Laureano G, Cunha J, Eiras-Dias J, Matos AR, Figueiredo A. Fatty acid
717 modulation and desaturase gene expression are differentially triggered in grapevine
718 incompatible interaction with biotrophs and necrotrophs. *Plant Physiology and*
719 *Biochemistry* 2021; **163**: 230–238.
- 720 17 Laureano G, Figueiredo J, Cavaco AR *et al.* The interplay between membrane lipids and
721 phospholipase A family members in grapevine resistance against *Plasmopara viticola*. *Sci*
722 *Rep* 2018; **8**: 14538.
- 723 18 Serrano M, Coluccia F, Torres M, L’Haridon F, Métraux J-P. The cuticle and plant defense
724 to pathogens. *Front Plant Sci* 2014; **5**: 274.
- 725 19 Kang J, Park J, Choi H *et al.* Plant ABC transporters. *Arabidopsis Book* 2011; **9**: e0153.
- 726 20 Borghi L, Kang J, de Brito Francisco R. Filling the gap: functional clustering of ABC proteins
727 for the investigation of hormonal transport *in planta*. *Frontiers in Plant Science* 2019; **10**:
728 422.
- 729 21 Bessire M, Borel S, Fabre G *et al.* A Member of the PLEIOTROPIC DRUG RESISTANCE family
730 of ATP binding cassette transporters is required for the formation of a functional cuticle in
731 *Arabidopsis*. *The Plant Cell* 2011; **23**: 1958–1970.

- 732 22 Wang Y, Sun Y, You Q *et al.* Three fatty acyl-coenzyme A reductases, BdFAR1, BdFAR2 and
733 BdFAR3, are involved in cuticular wax primary alcohol biosynthesis in *Brachypodium*
734 *distachyon*. *Plant and Cell Physiology* 2018; **59**: 527–543.
- 735 23 Khizar M, Shi J, Haroon U *et al.* RNA-Seq based exploration of differentially expressed
736 genes (DEGs) in cotton (*Gossypium hirsutum* L.) upon infection with *Aspergillus*
737 *tubingensis*. *Research Square* 2021. doi:10.21203/rs.3.rs-770815/v1.
- 738 24 Liu N, Sun Y, Pei Y *et al.* A pectin methylesterase inhibitor enhances resistance to
739 *Verticillium* wilt. *Plant Physiol* 2018; **176**: 2202–2220.
- 740 25 Chávez Montes RA, Ranocha P, Martinez Y *et al.* Cell wall modifications in Arabidopsis
741 plants with altered alpha-L-arabinofuranosidase activity. *Plant Physiol* 2008; **147**: 63–77.
- 742 26 Franken ACW, Werner ER, Haas H *et al.* The role of coproporphyrinogen III oxidase and
743 ferrochelatase genes in heme biosynthesis and regulation in *Aspergillus niger*. *Appl*
744 *Microbiol Biotechnol* 2013; **97**: 9773–9785.
- 745 27 Leger RJS, Joshi L, Roberts DWY 1997. Adaptation of proteases and carbohydrases of
746 saprophytic, phytopathogenic and entomopathogenic fungi to the requirements of their
747 ecological niches. *Microbiology*; **143**: 1983–1992.
- 748 28 Dubovenko AG, Dunaevsky YE, Belozersky MA, Oppert B, Lord JC, Elpidina EN. Trypsin-like
749 proteins of the fungi as possible markers of pathogenicity. *Fungal Biology* 2010; **114**: 151–
750 159.
- 751 29 Monaghan J, Zipfel C. Plant pattern recognition receptor complexes at the plasma
752 membrane. *Current Opinion in Plant Biology* 2012; **15**: 349–357.
- 753 30 Hammond-Kosack KE, Jones JD. Resistance gene-dependent plant defense responses.
754 *Plant Cell* 1996; **8**: 1773–1791.
- 755 31 Morel J-B, Dangl JL. The hypersensitive response and the induction of cell death in plants.
756 *Cell Death Differ* 1997; **4**: 671–683.
- 757 32 Dong X. Genetic dissection of systemic acquired resistance. *Current Opinion in Plant*
758 *Biology* 2001; **4**: 309–314.
- 759 33 Métraux J-P. Systemic acquired resistance and salicylic acid: current state of knowledge.
760 *European Journal of Plant Pathology* 2001; **107**: 13–18.
- 761 34 Anderson RG, Deb D, Fedkenheuer K, McDowell JM. Recent progress in RxLR effector
762 research. *MPMI* 2015; **28**: 1063–1072.
- 763 35 Wang Y, Wang Y. *Phytophthora sojae* effectors orchestrate warfare with host immunity.
764 *Curr Opin Microbiol* 2018; **46**: 7–13.
- 765 36 Feechan A, Anderson C, Torregrosa L *et al.* Genetic dissection of a TIR-NB-LRR locus from
766 the wild North American grapevine species *Muscadinia rotundifolia* identifies paralogous
767 genes conferring resistance to major fungal and oomycete pathogens in cultivated
768 grapevine. *The Plant Journal* 2013; **76**: 661–674.

- 769 37 Williams S, Yin L, Foley G *et al.* Structure and function of the TIR domain from the grape
770 NLR protein RPV1. *Frontiers in Plant Science* 2016; **7**: 1850.
- 771 38 Horsefield S, Burdett H, Zhang X *et al.* NAD⁺ cleavage activity by animal and plant TIR
772 domains in cell death pathways. *Science* 2019; **365**: 793–799.
- 773 39 Afzal AJ, Wood AJ, Lightfoot DA. Plant receptor-like serine threonine kinases: roles in
774 signaling and plant defense. *Mol Plant Microbe Interact* 2008; **21**: 507–517.
- 775 40 Romeis T. Protein kinases in the plant defence response. *Current Opinion in Plant Biology*
776 2001; **4**: 407–414.
- 777 41 Ji D, Chen T, Zhang Z, Li B, Tian S. Versatile roles of the receptor-like kinase FERONIA in
778 plant growth, development and host-pathogen interaction. *Int J Mol Sci* 2020; **21**: 7881.
- 779 42 Kandel SL, Hulse-Kemp AM, Stoffel K *et al.* Transcriptional analyses of differential cultivars
780 during resistant and susceptible interactions with *Peronospora effusa*, the causal agent of
781 spinach downy mildew. *Sci Rep* 2020; **10**: 6719.
- 782 43 Fuglsang AT, Palmgren M. Proton and calcium pumping P-type ATPases and their
783 regulation of plant responses to the environment. *Plant Physiology* 2021.
784 doi:10.1093/plphys/kiab330.
- 785 44 Hashimoto-Sugimoto M, Higaki T, Yaeno T *et al.* A Munc13-like protein in Arabidopsis
786 mediates H⁺-ATPase translocation that is essential for stomatal responses. *Nat Commun*
787 2013; **4**: 2215.
- 788 45 Gaxiola RA, Palmgren MG, Schumacher K. Plant proton pumps. *FEBS Letters* 2007; **581**:
789 2204–2214.
- 790 46 Lee HC, Carroll A, Crossett B, Connolly A, Batarseh A, Djordjevic MA. Improving the
791 identification and coverage of plant transmembrane proteins in Medicago using bottom–
792 up proteomics. *Frontiers in Plant Science* 2020; **11**: 1925.
- 793 47 Sakamoto H, Matsuda O, Iba K. ITN1, a novel gene encoding an ankyrin-repeat protein that
794 affects the ABA-mediated production of reactive oxygen species and is involved in salt-
795 stress tolerance in *Arabidopsis thaliana*. *The Plant Journal* 2008; **56**: 411–422.
- 796 48 Saini K, AbdElgawad H, Markakis MN *et al.* Perturbation of auxin homeostasis and signaling
797 by PINOID overexpression induces stress responses in Arabidopsis. *Front Plant Sci* 2017; **8**:
798 1308.
- 799 49 Kazan K, Manners JM. Linking development to defense: auxin in plant–pathogen
800 interactions. *Trends in Plant Science* 2009; **14**: 373–382.
- 801 50 Mudgil Y, Uhrig JF, Zhou J, Temple B, Jiang K, Jones AM. Arabidopsis N-MYC
802 DOWNREGULATED-LIKE1, a positive regulator of auxin transport in a G protein–mediated
803 pathway. *Plant Cell* 2009; **21**: 3591–3609.
- 804 51 Parry G, Estelle M. Regulation of cullin-based ubiquitin ligases by the Nedd8/RUB
805 ubiquitin-like proteins. *Semin Cell Dev Biol* 2004; **15**: 221–229.

806 52 Ren C, Pan J, Peng W *et al.* Point mutations in Arabidopsis Cullin1 reveal its essential role
807 in jasmonate response. *Plant J* 2005; **42**: 514–524.

808 53 Potters G, Pasternak TP, Guisez Y, Palme KJ, Jansen MAK. Stress-induced morphogenic
809 responses: growing out of trouble? *Trends in Plant Science* 2007; **12**: 98–105.

810 54 Potters G, Pasternak TP, Guisez Y, Jansen M a. K. Different stresses, similar morphogenic
811 responses: integrating a plethora of pathways. *Plant, Cell & Environment* 2009; **32**: 158–
812 169.

813 55 Djami-Tchatchou AT, Harrison GA, Harper CP *et al.* Dual role of auxin in regulating plant
814 defense and bacterial virulence gene expression during *Pseudomonas syringae* PtoDC3000
815 pathogenesis. *MPMI* 2020; **33**: 1059–1071.

816 56 Apel K, Hirt H. Reactive oxygen species: metabolism, oxidative stress, and signal
817 transduction. *Annual Review of Plant Biology* 2004; **55**: 373–399.

818 57 Baxter A, Mittler R, Suzuki N. ROS as key players in plant stress signalling. *Journal of*
819 *Experimental Botany* 2014; **65**: 1229–1240.

820 58 Waszczak C, Carmody M, Kangasjärvi J. Reactive oxygen species in plant signaling. *Annual*
821 *Review of Plant Biology* 2018; **69**: 209–236.

822 59 Figueiredo A, Martins J, Sebastiana M *et al.* Specific adjustments in grapevine leaf
823 proteome discriminating resistant and susceptible grapevine genotypes to *Plasmopara*
824 *viticola*. *Journal of Proteomics* 2017; **152**: 48–57.

825 60 Ravichandran S, Stone S, Benkel B, Zhang J, Berrue F, Prithiviraj B. Optimal level of purple
826 acid phosphatase 5 is required for maintaining complete resistance to *Pseudomonas*
827 *syringae*. *Frontiers in Plant Science* 2015; **6**: 568.

828 61 Podgórska A, Burian M, Szal B. Extra-cellular but extra-ordinarily important for cells:
829 apoplastic reactive oxygen species metabolism. *Frontiers in Plant Science* 2017; **8**: 1353.

830 62 Shapiguzov A, Vainonen J, Wrzaczek M, Kangasjärvi J. ROS-talk – how the apoplast, the
831 chloroplast, and the nucleus get the message through. *Frontiers in Plant Science* 2012; **3**:
832 292.

833 63 Qi J, Wang J, Gong Z, Zhou J-M. Apoplastic ROS signaling in plant immunity. *Current Opinion*
834 *in Plant Biology* 2017; **38**: 92–100.

835 64 Ma X, Wang W, Bittner F *et al.* Dual and opposing roles of xanthine dehydrogenase in
836 defense-associated reactive oxygen species metabolism in Arabidopsis. *Plant Cell* 2016;
837 **28**: 1108–1126.

838 65 Hadden DA, Phillipson BA, Johnston KA *et al.* Arabidopsis PEX19 is a dimeric protein that
839 binds the peroxin PEX10. *Mol Membr Biol* 2006; **23**: 325–336.

840 66 Su T, Li W, Wang P, Ma C. Dynamics of peroxisome homeostasis and its role in stress
841 response and signaling in plants. *Frontiers in Plant Science* 2019; **10**: 705.

842 67 Küchler M, Decker S, Hörmann F, Soll J, Heins L. Protein import into chloroplasts involves
843 redox-regulated proteins. *EMBO J* 2002; **21**: 6136–6145.

- 844 68 Huang S, Braun H-P, Gawryluk RMR, Millar AH. Mitochondrial complex II of plants: subunit
845 composition, assembly, and function in respiration and signaling. *The Plant Journal* 2019;
846 **98**: 405–417.
- 847 69 Zhang J, Sun X. Recent advances in polyphenol oxidase-mediated plant stress responses.
848 *Phytochemistry* 2021; **181**: 112588.
- 849 70 Scarpeci TE, Zanor MI, Valle EM. Investigating the role of plant heat shock proteins during
850 oxidative stress. *Plant Signal Behav* 2008; **3**: 856–857.
- 851 71 Cesco S, Tolotti A, Nadalini S *et al.* *Plasmopara viticola* infection affects mineral elements
852 allocation and distribution in *Vitis vinifera* leaves. *Sci Rep* 2020; **10**: 18759.
- 853 72 Huda KMdK, Banu MstSA, Tuteja R, Tuteja N. Global calcium transducer P-type Ca²⁺-
854 ATPases open new avenues for agriculture by regulating stress signalling. *Journal of*
855 *Experimental Botany* 2013; **64**: 3099–3109.
- 856 73 Rademacher E, Offringa R. Evolutionary adaptations of plant AGC kinases: from light
857 signaling to cell polarity regulation. *Frontiers in Plant Science* 2012; **3**: 250.
- 858 74 Chang J-Y, Nakahata Y, Hayano Y, Yasuda R. Mechanisms of Ca²⁺/calmodulin-dependent
859 kinase II activation in single dendritic spines. *Nat Commun* 2019; **10**: 2784.
- 860 75 Ding X, Yu Q, Zhang B *et al.* The type II Ca²⁺/calmodulin-dependent protein kinases are
861 involved in the regulation of cell wall integrity and oxidative stress response in *Candida*
862 *albicans*. *Biochemical and Biophysical Research Communications* 2014; **446**: 1073–1078.
- 863 76 Yang Y, Cheng P, Zhi G, Liu Y. Identification of a calcium/calmodulin-dependent protein
864 kinase that phosphorylates the *Neurospora* circadian clock protein FREQUENCY. *Journal of*
865 *Biological Chemistry* 2001; **276**: 41064–41072.
- 866 77 Liu X-H, Lu J-P, Dong B, Gu Y, Lin F-C. Disruption of MoCMK1, encoding a putative
867 calcium/calmodulin-dependent kinase, in *Magnaporthe oryzae*. *Microbiological Research*
868 2010; **165**: 402–410.
- 869 78 Wang W-H, Zheng H-L. Mechanisms for calcium sensing receptor-regulated stomatal
870 closure in response to the extracellular calcium signal. *Plant Signaling & Behavior* 2012; **7**:
871 289–291.
- 872 79 Nechushtai R, Karim O, Zuo K *et al.* The balancing act of NEET proteins: iron, ROS, calcium
873 and metabolism. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 2020;
874 **1867**: 118805.
- 875 80 Mélida H, Sandoval-Sierra JV, Diéguez-Urbeondo J, Bulone V. Analyses of extracellular
876 carbohydrates in oomycetes unveil the existence of three different cell wall types.
877 *Eukaryotic Cell* 2013; **12**: 194–203.
- 878 81 Oh IS, Park AR, Bae MS *et al.* Secretome analysis reveals an Arabidopsis lipase involved in
879 defense against *Alternaria brassicicola*. *The Plant Cell* 2005; **17**: 2832–2847.
- 880 82 Park S-C, Kim IR, Kim J-Y *et al.* Molecular mechanism of *Arabidopsis thaliana* profilins as
881 antifungal proteins. *Biochim Biophys Acta Gen Subj* 2018; **1862**: 2545–2554.

882 83 Nathalie Leborgne-Castel, Bouhidel K. Plasma membrane protein trafficking in plant–
883 microbe interactions: a plant cell point of view. *Front Plant Sci* 2014; **5**: 735.

884 84 Dautt-Castro M, Rosendo-Vargas M, Casas-Flores S. The small GTPases in fungal signaling
885 conservation and function. *Cells* 2021; **10**: 1039.

886 85 Higuchi Y. Membrane traffic related to endosome dynamics and protein secretion in
887 filamentous fungi. *Bioscience, Biotechnology, and Biochemistry* 2021; **85**: 1038–1045.

888 86 Zerial M, McBride H. Rab proteins as membrane organizers. *Nat Rev Mol Cell Biol* 2001; **2**:
889 107–117.

890 87 Zheng H, Zheng W, Wu C *et al.* Rab GTPases are essential for membrane trafficking-
891 dependent growth and pathogenicity in *Fusarium graminearum*. *Environ Microbiol* 2015;
892 **17**: 4580–4599.

893 88 Marshall NC, Finlay BB, Overall CM. Sharpening host defenses during infection: proteases
894 cut to the chase. *Molecular & Cellular Proteomics* 2017; **16**: S161–S171.

895 89 Figaj D, Ambroziak P, Przepiora T, Skorko-Glonek J. The role of proteases in the virulence
896 of plant pathogenic bacteria. *International Journal of Molecular Sciences* 2019; **20**: 672.

897 90 Liu X-H, Ning G-A, Huang L-Y *et al.* Calpains are involved in asexual and sexual
898 development, cell wall integrity and pathogenicity of the rice blast fungus. *Sci Rep* 2016; **6**:
899 31204.

900 91 Futai E, Maeda T, Sorimachi H, Kitamoto K, Ishiura S, Suzuki K. The protease activity of a
901 calpain-like cysteine protease in *Saccharomyces cerevisiae* is required for alkaline
902 adaptation and sporulation. *Mol Gen Genet* 1999; **260**: 559–568.

903 92 Ramírez V, López A, Mauch-Mani B, Gil MJ, Vera P. An extracellular subtilase switch for
904 immune priming in *Arabidopsis*. *PLoS Pathogens* 2013; **9**: e1003445.

905 93 Alves de Castro P, dos Reis TF, Dolan SK *et al.* The *Aspergillus fumigatus* SchASCH9 kinase
906 modulates SakAHOG1 MAP kinase activity and it is essential for virulence. *Molecular*
907 *Microbiology* 2016; **102**: 642–671.

908 94 Fang W, Pava-ripoll M, Wang S, St. Leger R. Protein kinase A regulates production of
909 virulence determinants by the entomopathogenic fungus, *Metarhizium anisopliae*. *Fungal*
910 *Genetics and Biology* 2009; **46**: 277–285.

911 95 Wang C, Shen D, Wang J *et al.* An AGC kinase, PgAGC1 regulates virulence in the
912 entomopathogenic oomycete *Pythium guiyangense*. *Fungal Biology* 2019; **123**: 87–93.

913 96 Berndt P, Lanver D, Kahmann R. The AGC Ser/Thr kinase Aga1 is essential for appressorium
914 formation and maintenance of the actin cytoskeleton in the smut fungus *Ustilago maydis*.
915 *Molecular Microbiology* 2010; **78**: 1484–1499.

916 97 Islam MT, von Tiedemann A, Laatsch H. Protein kinase C is likely to be involved in
917 zoosporogenesis and maintenance of flagellar motility in the peronosporomycete
918 zoospores. *Mol Plant Microbe Interact* 2011; **24**: 938–947.

919 98 Castillo-Lluya S, Alvarez-Tabarés I, Weber I, Steinberg G, Pérez-Martín J. Sustained cell
920 polarity and virulence in the phytopathogenic fungus *Ustilago maydis* depends on an
921 essential cyclin-dependent kinase from the Cdk5/Pho85 family. *Journal of Cell Science*
922 2007; **120**: 1584–1595.

923 99 Kiefer B, Riemann M, Büche C, Kassemeyer H-H, Nick P. The host guides morphogenesis
924 and stomatal targeting in the grapevine pathogen *Plasmopara viticola*. *Planta* 2002; **215**:
925 387–393.

926 100 Riemann M, Büche C, Kassemeyer H-H, Nick P. Cytoskeletal responses during early
927 development of the downy mildew of grapevine (*Plasmopara viticola*). *Protoplasma* 2002;
928 **219**: 13–22.

929 101 Hamel L-P, Nicole M-C, Duplessis S, Ellis BE. Mitogen-activated protein kinase signaling in
930 plant-interacting fungi: distinct messages from conserved messengers. *The Plant Cell* 2012;
931 **24**: 1327–1351.

932 102 Zhao X, Mehrabi R, Xu J-R. Mitogen-activated protein kinase pathways and fungal
933 pathogenesis. *Eukaryotic Cell* 2007; **6**: 1701–1714.

934 103 Jiang L, Situ J, Deng YZ *et al.* PIMAPK10, a mitogen-activated protein kinase (MAPK) in
935 *Peronophythora litchii*, is required for mycelial growth, sporulation, laccase activity, and
936 plant infection. *Front Microbiol* 2018; **9**: 426.

937 104 Li A, Wang Y, Tao K *et al.* PsSAK1, a stress-activated MAP kinase of *Phytophthora sojae*, is
938 required for zoospore viability and infection of soybean. *Mol Plant Microbe Interact* 2010;
939 **23**: 1022–1031.

940 105 Zhang H, Xue C, Kong L, Li G, Xu J-R. A Pmk1-interacting gene is involved in appressorium
941 differentiation and plant infection in *Magnaporthe oryzae*. *Eukaryot Cell* 2011; **10**: 1062–
942 1070.

943 106 Yue X, Que Y, Deng S *et al.* The cyclin dependent kinase subunit Cks1 is required for
944 infection-associated development of the rice blast fungus *Magnaporthe oryzae*. *Environ*
945 *Microbiol* 2017; **19**: 3959–3981.

946 107 Shen D, Dong Y, Wei Y *et al.* Genome-wide and functional analyses of tyrosine kinase-like
947 family genes reveal potential roles in development and virulence in mosquito pathogen
948 *Pythium guiyangense*. *Fungal Genet Biol* 2019; **130**: 11–18.

949 108 Judelson HS, Ah-Fong AM. The kinome of *Phytophthora infestans* reveals oomycete-
950 specific innovations and links to other taxonomic groups. *BMC Genomics* 2010; **11**: 700.

951 109 Allan RK, Ratajczak T. Versatile TPR domains accommodate different modes of target
952 protein recognition and function. *Cell Stress Chaperones* 2011; **16**: 353–367.

953 110 Mittl PRE, Schneider-Brachert W. Sel1-like repeat proteins in signal transduction. *Cell*
954 *Signal* 2007; **19**: 20–31.

955 111 Cervený L, Strasková A, Danková V *et al.* Tetratricopeptide repeat motifs in the world of
956 bacterial pathogens: role in virulence mechanisms. *Infect Immun* 2013; **81**: 629–635.

957 112 Dankova V, Balonova L, Straskova A *et al.* Characterization of tetratricopeptide repeat-like
958 proteins in *Francisella tularensis* and identification of a novel locus required for virulence.
959 *Infect Immun* 2014; **82**: 5035–5048.

960 113 Randall TA, Dwyer RA, Huitema E *et al.* Large-scale gene discovery in the oomycete
961 *Phytophthora infestans* reveals likely components of phytopathogenicity shared with true
962 fungi. *MPMI* 2005; **18**: 229–243.

963 114 Wang W, Deng Z, Wu H *et al.* A small secreted protein triggers a TLR2/4-dependent
964 inflammatory response during invasive *Candida albicans* infection. *Nat Commun* 2019; **10**:
965 1015.

966 115 Roemer T, Delaney S, Bussey H. SKN1 and KRE6 define a pair of functional homologs
967 encoding putative membrane proteins involved in beta-glucan synthesis. *Mol Cell Biol*
968 1993; **13**: 4039–4048.

969 116 Wang C, Li M, Li G *et al.* Two distinct nucleic acid binding surfaces of Cdc5 regulate
970 development. *Biochem J* 2019; **476**: 3355–3368.

971 117 Veloso MM, Almandanim MC, Baleiras-Couto M *et al.* Microsatellite database of grapevine
972 (*Vitis vinifera* L.) cultivars used for wine production in Portugal. *Ciência Téc Vitiv* 2010; **25**:
973 53–61.

974 118 Almagro Armenteros JJ, Salvatore M, Emanuelsson O *et al.* Detecting sequence signals in
975 targeting peptides using deep learning. *Life Sci Alliance* 2019; **2**: e201900429.

976 119 Sperschneider J, Dodds P. EffectorP 3.0: prediction of apoplastic and cytoplasmic effectors
977 in fungi and oomycetes. *MPMI* 2021. doi:10.1094/MPMI-08-21-0201-R.

978 120 Pang Z, Chong J, Zhou G *et al.* MetaboAnalyst 5.0: narrowing the gap between raw spectra
979 and functional insights. *Nucleic Acids Research* 2021; **49**: W388–W396.

980 121 Breitling R, Armengaud P, Amtmann A, Herzyk P. Rank products: a simple, yet powerful,
981 new method to detect differentially regulated genes in replicated microarray experiments.
982 *FEBS Lett* 2004; **573**: 83–92.

983 122 Bendtsen JD, Jensen LJ, Blom N, Von Heijne G, Brunak S. Feature-based prediction of non-
984 classical and leaderless protein secretion. *Protein Eng Des Sel* 2004; **17**: 349–356.

985 123 Armenteros JJA, Tsirigos KD, Sønderby CK *et al.* SignalP 5.0 improves signal peptide
986 predictions using deep neural networks. *Nature Biotechnology* 2019; **37**: 420.

987 124 Emanuelsson O, Brunak S, Heijne G von, Nielsen H. Locating proteins in the cell using
988 TargetP, SignalP and related tools. *Nature Protocols* 2007; **2**: 953.

989 125 Sperschneider J, Dodds PN, Singh KB, Taylor JM. ApoplastP: prediction of effectors and
990 plant proteins in the apoplast using machine learning. *New Phytologist* 2018; **217**: 1764–
991 1778.

992 126 Savojardo C, Martelli PL, Fariselli P, Profiti G, Casadio R. BUSCA: an integrative web server
993 to predict subcellular localization of proteins. *Nucleic Acids Res* 2018; **46**: W459–W466.

994 127 Petsalaki EI, Bagos PG, Litou ZI, Hamodrakas SJ. PredSL: a tool for the N-terminal sequence-
995 based prediction of protein subcellular localization. *Genomics, Proteomics &*
996 *Bioinformatics* 2006; **4**: 48–55.

997 128 Lohse M, Nagel A, Herter T *et al.* Mercator: a fast and simple web server for genome scale
998 functional annotation of plant sequence data. *Plant Cell Environ* 2014; **37**: 1250–1258.

999 129 Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M. Blast2GO: a universal tool
1000 for annotation, visualization and analysis in functional genomics research. *Bioinformatics*
1001 2005; **21**: 3674–3676.

1002

1003