1 An integrative process-based model for biomass 2 and yield estimation of hardneck garlic (*Allium* 3 *sativum*)

4 Kyungdahm Yun
University of Washington

5 Minji Shin
Research Institute of Climate Change and Agriculture

6 Kyung Hwan Moon
Research Institute of Climate Change and Agriculture

7 Soo-Hyung Kim
University of Washington

8 **Abstract**

9 We introduce an integrative process-driven crop model for garlic (*Allium*
10 *sativum*). Building on our previous model that simulated key phenological,
11 morphological, and physiological features of a garlic plant, the new garlic
12 model provides comprehensive and integrative estimations of biomass ac-
13 cumulation and yield formation under diverse environmental conditions.
14 This model also showcases an application of Cropbox to develop a com-
15 prehensive crop model. Cropbox is a crop modeling framework featuring
16 declarative modeling language and unified simulation interface for building
17 and improving crop models. Using Cropbox, we first evaluated the model
18 performance against three datasets with an emphasis on biomass and yield
19 measured under different environmental conditions and growing seasons.
20 We then applied the model to simulate optimal planting dates under future
21 climate conditions for assessing climate adaptation strategies between two
22 contrasting locations in South Korea: the current growing region (Gosan)
23 and an unfavorable cold winter region (Chuncheon, Gangwon-do). The
24 model simulated the growth and development of a southern-type cultivar

(Namdo, Jeju-do) reasonably well. Under RCP (Representative Concentration Pathway) scenarios, an overall delay in optimal planting date from a week to a month and a slight increase in potential yield were expected in Gosan. Expansion of growing region to northern area including Chuncheon was expected due to mild winter temperatures in the future and may allow Namdo cultivar production in more regions. The predicted optimal planting date in the new region was similar to the current growing region that favors early fall planting. Our new integrative garlic model provides mechanistic, process-driven crop responses to environmental cues and can be useful for assessing climate impacts and identifying crop specific climate adaptation strategies for the future.

1 Introduction

Garlic (*Allium sativum*) is a historically important horticultural crop in many countries with global production reaching 30.7 million tons in 2019 after 40 % increase of production in the last decade (FAO, 2020). Physiology of garlic has been extensively studied with an emphasis on characteristics as a bulbous crop where a clove of bulb is planted for growth and a newly grown bulb is harvested for storage and the next round of planting (Takagi, 1989; Kamenetsky, 2007). Some knowledge has been transferred to building crop models specifically targeted for simulating garlic growth and estimating yield at harvest. An early attempt for building a whole-plant garlic model was based on radiation-use efficiency (RUE) to obtain the total amount of carbon assimilates (Rizzalli et al., 2002). CropSyst, which is also a crop model based on RUE, was parameterized for garlic and used for simulating crop rotation between garlic and wheat (Giménez et al., 2016). Other studies focused on a certain aspect of garlic growth and development. The crop coefficient (K_c) needed for calculating evapotranspiration with Penman-Monteith equation was specifically determined for garlic (Villalobos et al., 2004). Photosynthesis and transpiration responses to various environmental conditions were obtained for building a leaf-level gas-exchange model for garlic (Kim et al., 2013). Photosynthetic responses under elevated CO₂ and nitrogen fertilization were further investigated for building a robust model for future climate conditions (Nackley et al., 2016). With a coupled gas-exchange model parameterized for garlic, a process-based model for simulating leaf development and growth of hardneck garlic was developed (Hsiao et al., 2019). Phenology of leaf initiation and appearance was individually tracked

by taking account of multiple cues including thermal time accumulation, bulb storage effect, and photoperiod. Individual leaf elongation was translated and aggregated into leaf area expansion at canopy level which was then divided into two layers of sunlit and shaded leaves for accounting assimilated carbon based on the coupled gas-exchange model. However, carbon partitioning into plant organs such as bulb was not validated and thus not used for yield estimation. Carbon partitioning is a crucial step in yield estimation modeling for horticultural crops in a sense that the final yield is a result of biomass partitioned into a certain organ, such as bulb, to be harvested (Marcelis et al., 1998). The early garlic model used a set of multiple partitioning coefficients dynamically varying with developmental stages of the plant (Rizzalli et al., 2002).

Yield estimation for garlic cultivars grown in each region has been crucial to manage growing practices and maintain a stable supply in the market (Põldma et al., 2005; Lee et al., 2011; Abdalla et al., 2011; Portela et al., 2012). Regression models based on weather data or satellite images were often used for yield estimation at a large scale with a minimum set of historical input data (Choi and Baek, 2016; Gómez et al., 2021). Although worldwide garlic production has been steadily increased in the recent years, it is unclear if yield will be stable in the current growing regions under future climate conditions. For estimating yield under previously unobserved conditions, it is critical to develop a physiological understanding of how garlic would respond to these environmental cues (Nackley et al., 2016). Regional climate changes may lead to changes in current farming practices and shifts in growing regions to maintain or maximize the yield. Crop model is an important tool for supporting such decisions by enabling simulations of plant growth under diverse climate adaptation strategies (Rosenzweig et al., 2014; Holzkämper et al., 2015; Corbeels et al., 2018).

In this study, our primary objective was to build an integrative process-driven garlic model suitable for estimating harvestable biomass as scape and bulb yield under diverse environments including future climate conditions. A new model was developed based on an existing process-based garlic model with an original emphasis on phenology and extended with a focus on biomass accumulation and partitioning (Hsiao et al., 2019). The model was also improved to better reflect physiological responses to temperature by taking account storage conditions of seed garlic bulbs and cold stress response in terms of leaf-level growth and canopy-level mortality. We used Cropbox modeling framework for reimplementing to take advantage of its declarative modeling language and unified interface for coordinating a large batch of simulations with minimum

97 configurations (Yun et al., 2020). For model testing, a new parameter set
 98 was specifically calibrated for Namdo cultivar (*Allium sativum* ‘Namdo’) and
 99 validated with multiple datasets with biomass measurements. For demonstrating
 100 yield estimation capability of the model, a climate adaptation strategy was
 101 assessed by model simulations for the same cultivar grown in two locations
 102 of South Korea. Namdo is a cultivar originally adapted to warm climate in
 103 southern region of Korea (Kim et al., 2009). However, the boundary of growth
 104 region between northern and southern types of garlic has moved northward in
 105 the past decades due to warming climate (Heo et al., 2006). Optimal planting
 106 dates for achieving maximum yield were discovered through model simulation
 107 under current and future climate conditions in the two locations where one is an
 108 already established region for growing southern-type garlic and the other has a
 109 potential to become a new establishment in the future.

110 2 Materials and Methods

111 2.1 Garlic Model

112 The garlic growth model was extended from a process-based model for leaf
 113 development and growth in hardneck garlic (Hsiao et al., 2019). The original
 114 model was capable of simulating leaf area expansion at an individual leaf level and
 115 estimating carbon assimilated in a canopy calculated by coupled gas-exchange,
 116 but assessing biomass allocated into a particular organ, *i.e.* bulb, was not
 117 a primary subject of the model at the time. For realistic yield prediction
 118 under future scenarios, the model should be able to be reliable under diverse
 119 environmental conditions (IPCC, 2014).

120 Model changes made for the experiments reported in this paper include
 121 improved biomass allocation, a dynamically adjusted phyllochron, and cold stress
 122 response (Figure 1). For the sake of technical convenience, model code originally
 123 written in C++ was reimplemented with Cropbox modeling framework using
 124 Julia programming language (Cropbox.jl, 2021). The Cropbox framework allowed
 125 a streamlined model development from model description in a concise declarative
 126 form, iterative parameter adjustments within a notebook environment, and batch
 127 simulation of large sets of parameters and production of figures included in this
 128 paper. The model source code is also available in the public repository (Garlic.jl,
 129 2021).

Figure 1: Diagram of key processes updated in the model presented in this paper. Leaf appearance is driven by accumulation of leaf tip appearance rate (LTAR) which is modified according to dynamic phyllochron (*). Cold stress (†) is implemented in two folds; individual leaf-level growth driven by leaf elongation rate (LER) and plant population-level response accounted by plant density (PD). Carbon assimilated through coupled gas exchange forms a carbon pool which determines the amount of carbon partitioned into individual organs. The bulb carbon constitutes a final yield of garlic. For a complete structure of the model, refer to Figure 1 of Hsiao et al. (2019).

2.1.1 Biomass Allocation

The total amount of carbon assimilation was calculated by a C_3 photosynthesis model coupled with a stomatal conductance model and energy balance model as described in the previous paper (Hsiao et al., 2019). Note that overall model structure including the gas-exchange module was reorganized for taking advantage of domain-specific language provided by Cropbox modeling framework (Yun

et al., 2020). The assimilated carbon accumulates and forms a carbon pool (g) ready to be distributed to plant organs. A potential allocation rate of available carbon (g d^{-1}) is driven by carbon supply rate from the pool (g d^{-1}) excluding maintenance respiration (g d^{-1}). After taking account of carbohydrate synthesis efficiency (Y_g), an actual allocation rate (g d^{-1}) is determined and split into a set of allocation rates for structural organs according to a partitioning table (Figure 2).

The partitioning table is a 2-dimensional array where each row represents a developmental stage and column represents a destination. Developmental stages span from seed, vegetative, bulb growth before scape appearance, bulb growth after scape appearance, bulb growth after scape removal, and death. Partitioning destinations include root, leaf, sheath, scape, and bulb. For each time step of simulation, an actual allocation rate weighted by a partitioning coefficient found in the table was used for biomass accumulation of each destination organ.

Figure 2: Partitioning coefficients for plant organs dynamically adjusted according to developmental stage. V: vegetative stage; R1: reproductive stage between scape initiation and appearance; R2: reproductive stage between scape appearance and removal; R3: reproductive stage after scape removal. R3 is not used if scape is not removed during simulation.

150 2.1.2 Dynamic Phyllochron

151 An interval between leaf appearance, phyllochron, is not necessarily static
 152 throughout plant growth, but can dynamically change depending on the growth
 153 condition. However, maximum leaf tip appearance rate ($\text{LTAR}_{\max}$; d^{-1}) in the
 154 original garlic model was determined by thermal time based on storage duration
 155 between harvest and planting as well as storage temperature for which seed
 156 garlic has been kept during this period. Once initialized, $\text{LTAR}_{\max}$ would have
 157 remained the same until the end of simulation. This assumption was often held
 158 when storage duration was close to an average duration where curve fitting was
 159 originally done for. However, sometimes the leaf tip appearance rate may have
 160 stayed too high at the end of growing season when storage duration was longer
 161 than usual. The opposite would happen when the storage duration was too short.
 162 In other words, scape appearance in the reproductive stage, which is driven
 163 by the same mechanism relying on phyllochron that assumes a scape appears
 164 after three phyllochrons since the onset of reproductive stage, could become
 165 too sensitive to the initial value of $\text{LTAR}_{\max}$. To alleviate this issue, $\text{LTAR}_{\max}$
 166 was adjusted to converge towards a half of the asymptote of maximum leaf tip
 167 appearance rate parameter ($\text{LTAR}_{\max,a}$; d^{-1}). The initial value $\text{LTAR}_{\max,s}$ is
 168 the same as maximal rate of leaf tip appearance rate modified by storage duration
 169 SD defined in the original model Hsiao et al. (2019). SD_m is the storage duration
 170 that results into maximal leaf tip appearance rate and α controls steepness of
 171 the sigmoidal storage function. The same value of parameters were adopted
 172 from the original model (Table 1). After each leaf appearance, $\text{LTAR}_{\max}$ would
 173 linearly increase or decrease depending on the initial rate and the slope towards
 174 the asymptote. Convergence is done when leaf rank k reaches the generic leaf
 175 number N_g which was set to 10 by default.

$$r_0 = \text{LTAR}_{\max,s} = \frac{\text{LTAR}_{\max,a}}{1 + e^{-\alpha(\text{SD} - \text{SD}_m)}} \quad (1)$$

$$r_1 = \frac{\text{LTAR}_{\max,a}}{2} \quad (2)$$

$$\text{LTAR}_{\max} = r_0 + (r_1 - r_0) \frac{k}{N_g} \quad (3)$$

Table 1: Parameters for dynamic phyllochron

Symbol	Value	Units	Description
$\text{LTAR}_{\max,a}$	0.4421	d^{-1}	Maximal leaf tip appearance asymptote
ST	8	$^{\circ}\text{C}$	Storage temperature
SD_m	117.7523	d	Storage days lead to maximal leaf tip appearance rate
α	0.0256	d^{-1}	Steepness of sigmoidal storage function
N_g	10	-	Generic leaf number

2.1.3 Cold Stress

Two types of cold stress response commonly found with garlic plants grown in the field during winter season were added to the model: cold injury and cold damage. Cold injury represents impeded leaf growth under below normal air temperatures. Actual leaf elongation rate (LER ; cm d^{-1}) is scaled down from potential leaf elongation rate (LER_p ; d^{-1}) by cold injury effect (E) which is a function of the number of cold days (D) the plant has experienced. The number of cold days (D) increases when the air temperature (T) is below a critical temperature for cold injury ($T_{c,i}$; $^{\circ}\text{C}$). Once the temperature rises above the threshold $T_{c,i}$, the cumulative days D decreases until it resets to zero. It assumes that the effect of cold injury is prolonged for a certain period time while recovering under normal temperature. The longer a plant is exposed to the cold temperature, the more difficult to recover from the injury. We assumed critical temperature where cold injury begins to appear was at 0°C and had related factors (a , b) fitted with a dataset obtained from a growth chamber experiment (Table 2).

Namdo (ND) cultivar were planted and grown in pots under room temperature until the third leaf emerged. Plants were then placed in a growth chamber subject to constant temperature of 0°C , -5°C , -10°C and -15°C , respectively. Each treatment had three replicates and leaf area of the third leaf blade from survived plants was measured for eight days. Then cold injury effect (E) was fitted to the relative portion of survived plants after a given time.

$$d = \begin{cases} 1, & T < T_{c,i} \\ -1, & T \geq T_{c,i} \end{cases} \quad (4)$$

$$D = \sum \max(d, 0) \quad (5)$$

$$E = 1 - \frac{\log(a(T - T_{c,i}) + b)}{e^{\frac{1}{b}}} \quad (6)$$

$$\text{LER} = E \cdot \text{LER}_p \quad (7)$$

198 Cold damage represents death of the plant under more extreme temperature,
 199 resulting into a reduced plant density (PD). The current plant density (PD)
 200 starts from an initial planting density (PD_0) and then may decrease over time as
 201 the air temperature (T) drops below a critical temperature for cold damage ($T_{c,d}$).
 202 Unlike cold injury where leaf growth only slows down under low temperature
 203 and eventually recovers once temperature rises back up, cold damage leads
 204 to permanent wilting from which plants no longer recover. Mortality due to
 205 cold damage (M) is a logistic curve representing a relative portion of plants
 206 survived at the end of the cold damage treatments. Cold damage shape factor
 207 (s) determines how quickly the plants will die off under such extreme conditions.
 208 Due to lack of experimental datasets available for parameter fitting, we instead
 209 looked up episodes of cold damage and corresponding temperatures reported in
 210 newspaper during the past years to determine parameter values.

$$M = \frac{e^{-s(T-T_{c,d})}}{1 + e^{-s(T-T_{c,d})}} \quad (8)$$

$$S = 1 - M \quad (9)$$

$$\text{PD} = S \cdot \text{PD}_0 \quad (10)$$

Table 2: Parameters for cold stress response

Symbol	Value	Units	Description
$T_{c,i}$	0	°C	Critical temperature for cold injury
a	-0.1	-	Cold injury factor 1
b	1.6	-	Cold injury factor 2
$T_{c,d}$	-15	°C	Critical temperature for cold damage
s	0.9	-	Cold damage shape factor

2.2 Parameter Calibration

A parameter set for Namdo (ND) cultivar was mostly based on existing parameter sets calibrated for Korean Mountain (KM) and Shantung Purple (SP). Only a few parameters were modified to reflect how ND grows in general compared to the other cultivars (Table 3). Minimum length of longest leaf (LM_{min}) was increased to 100 cm which was the largest value found in our datasets. Potential maximum elongation rate (LER_{max}) was accordingly adjusted to 5.56 cm d^{-1} assuming full leaf expansion takes 18 days under optimal condition. Initial leaves at harvest (ILN) was set to 6 as found in a dissected seed garlic clove. Stay green (SG) was set to 1.5 d which was a value calibrated for SP cultivar. Storage temperature (ST) was assumed constant 8°C during entire storage period. Seed bulb was assumed harvested on June 30th and storage duration (SD) was calculated accordingly. Note that it was our intention to select parameters that allow overestimation of biomass and leaf area for tracking potential growth while calibrating parameters to keep timing of phenology as close as possible to the observation.

We validated our new parameter set for ND cultivar using three datasets. The first dataset (D1) was collected from an experiment plot located at Research Institute of Climate Change and Agriculture (RICCA), Jeju, South Korea. ND cultivar was planted on October 8th, 2014 and harvested on June 19th, 2015. Growth and development measurements, such as leaf count, leaf area, biomass for each part, were recorded from mid-vegetative stage until harvest. This dataset was used for evaluating overall response of the ND parameter set we obtained above. The second dataset (D2) was collected from a temperature gradient greenhouse (TGG) located at the same site in Jeju. TGG is a glass house equipped with heaters on the one side of the wall that keeps a small, but constant gradient in the enclosed planting zones. Five planting zones were set up with 2°C to 3°C of temperature difference kept from end to end. ND cultivar was planted on October 7th, 2014 and harvested on May 17th, 2015. Similar measurements to the first dataset were recorded from early vegetative stage until harvest. This dataset was used for evaluating response of the model to modest temperature changes in a similar way that the model would be subject under future climate conditions. Scape was not removed in both datasets. The last third dataset (D3) was obtained from a farm field located at Jeongsil (JS) neighborhood in Jeju, South Korea. ND cultivar was planted on September 9th, 2009 and harvested on June 18th, 2010. Types of recorded variables were

the same as other datasets. Scape was assumed to be removed shortly after its appearance out of the whorl. Only the visible portion of scape was cut and measured for biomass. Hourly time-series of weather data for each dataset were obtained from nearby Jeju station (184) operated by Korean Meteorological Administration.

Table 3: Parameters for leaf development

Symbol	Value	Units	Description
$LTAR_{\max,a}$	0.4421	d^{-1}	Maximal leaf tip appearance rate asymptote
$LER_{\max}$	5.56	$cm\ d^{-1}$	Maximal leaf elongation rate
$LM_{\min}$	100	cm	Minimum length of longest leaf
ILN	6	-	Initial leaf number
SG	1.5	-	Stay green

2.3 Future Climate Projections

For assessing garlic yield changes in future climate conditions, we specifically chose two locations in South Korea (Figure 3). The first site was Gosan, Jeju located in a Southern island with a humid subtropical climate off the Korean peninsula, which is in Zone 9b in terms of USDA Hardiness Zones. Gosan is where Namdo (ND) cultivar is currently grown in a large scale for commercial purpose where stakeholders are interested to see how the cultivar would be performing and if any adjustments in growing practice such as if planting date shifts would be required in the long term. The second site was Chuncheon, Gangwon which is located in the middle of the Korean peninsula with a humid continental climate with cold winter. Chuncheon is in Zone 6b according to USDA Hardiness Zones and the current climate is not favorable for growing a Southern cultivar such as Namdo which has been historically more adapted to warm climate. However, it is not clear whether future climate conditions could allow growing of new crops in a region previously not suited for production. We assumed two RCP (Representative Concentration Pathway) scenarios, RCP4.5 and RCP8.5, for future climate projections (IPCC, 2014). Daily weather dataset for the two locations under RCP4.5 and RCP8.5 scenarios were obtained from AgClimate data portal (RDA, 2019) operated by Rural Development Administration in South Korea. Elevated CO_2 concentrations for RCP scenarios were used accordingly for driving the integrated coupled gas-exchange model. For comparison with current climate conditions of each location, we also obtained 30-years normal weather

274 data from 1980 to 2010 provided by the same data portal. The daily weather
275 dataset was then scaled down to hourly time-series by applying an interpolation
276 method specifically designed for different types of weather variables (Moon et al.,
277 2019). Elevated CO₂ levels for each year under future scenarios were obtained by
278 linear interpolation between CO₂ abundance projection reported in the Climate
279 System Scenario Tables (IPCC, 2013).

Figure 3: Geographic locations of Gosan and Jeju in South Korea. Gosan is located in a Southern island with a humid subtropical climate off the Korean peninsula. Chuncheon is located in the middle of the peninsula with a humid continental climate with cold winter. A color overlay represents average minimum temperature (°C) in January during past 30 years from 1981 to 2010 provided by National Institute of Agricultural Sciences, South Korea. According to USDA Hardiness Zones, Gosan is Zone 9b and Chuncheon is Zone 6b.

280 For each location, multiple runs of simulation were executed with combina-
281 tions of treatments. In the case of future climate projections, 8 time windows
282 from 2020s to 2090s with 10 years interval multiplied by 10 repetitions from differ-
283 ent random seed produced 80 weather datasets for each RCP scenario. Historical
284 weather was organized in a single time window referred to 1980s with 10 random
285 repetitions, leading to 10 weather datasets. Planting date was adjusted from
286 240th to 350th DOY (day of the year) with 10 days interval, requiring 12 runs
287 for each weather condition. In turn, the current normal scenario involved 120
288 runs of simulation while each RCP scenario required 960 runs. Immediate scape
289 removal after appearance was assumed. Harvest date was set fixed to be May

15th considering practical harvest dates of ND cultivar grown in Jeju. Yield was based on fresh biomass of bulb at the harvest date. In model simulation, fresh biomass was calculated with 85 % moisture content in bulb. Optimal plating date was then determined by finding out a planting date which gives maximum yield for a given year. For the sake of yield comparison between planting dates, years were grouped into three time periods, namely 1980–2010s, 2020–2050s, and 2060–2090s. A data point in the 1980–2010 period was composed of 10 samples originated from a set of stochastic weather data pooled for the entire period whereas a data point in the future climate scenario was composed of 50 samples where the result of 10 stochastic weather datasets was combined with 5 periods of 10 years interval.

3 Results

3.1 Model Validation

3.1.1 Dataset 1 (RICCA field)

Our parameter for ND cultivar was first evaluated with measurements from field grown garlic in dataset D1. Green leaf area from the model exhibited consistent overestimation with an average RMSE (root mean square error) of 275 cm² while keeping a similar trajectory of leaf expansion and senescence pattern to the observation (Figure 4). The maximum green leaf area was estimated to be 1193 cm² on April 28th while the maximum value in the dataset was 1150 cm² observed on April 30th. The peak was reached in simulation shortly after scape appeared on April 23rd during early vegetative growth.

Simulated biomass allocation to each part of plant structure, namely bulb, leaf, stalk, and scape, was close to the observation with an average RMSE of 5.0 g, 1.6 g, 2.5 g and 1.7 g, respectively (Figure 5). The maximum biomass of bulb from simulation was 22.6 g reached on June 14th while the observed maximum was 26.0 g measured on June 4th which was the final date recorded in the dataset. The maximum living leaf biomass from simulation was 8.4 g reached on April 27th and the maximum observed value was the same 9.9 g on April 30th. The maximum biomass of stalk, which consists of leaf sheath and scape, from simulation was 15.3 g close to observed maximum of 15.9 g. However, the timing of peak was shifted by almost three weeks later on May 27th compared to the observed May 7th. Since leaf sheath withered away at the end of growth stage,

Figure 4: Comparison of model simulated green leaf area and observation from dataset D1 which was from field measurements and primarily used for evaluating new parameters.

the only remaining part of living stalk would be a scape which was the reason why a convergence between two values occurred in mid-June. The simulated biomass of scape reached its maximum value of 9.5 g on June 20th which was almost identical to the observed maximum which, however, occurred earlier on June 4th. Note that the two slopes of biomass increase for simulation and observation were almost parallel to each other.

Leaf development phenology simulated by the model was in agreement with fresh leaf count recorded in the dataset with an average RMSE of 1.6 leaves (Figure 6). According to the model, leaf initiation was completed on March 22nd with the total number of 16 leaves and the 16th leaf appeared on April 2nd. The final leaf was completely senesced and dropped after about 2.5 months of growth on June 15th.

3.1.2 Dataset 2 (RICCA TGG)

Plants grown in higher temperature zone 1 had larger green leaf area of 1521 cm² and reached its peak earlier on April 20th when compared to plants grown in lower temperature zone 5 whose maximum green leaf area was estimated to be a smaller 1303 cm² reached on May 9th (Figure 7). Simulation result was in

Figure 5: Biomass allocation to leaf, bulb, stalk, and visible portion of scape simulated over time compared with measurements from dataset D1 which was from field measurements and primarily used for evaluating new parameters.

Figure 6: Simulated leaf development phenology compared with fresh leaf count recorded in dataset D1 which was from field measurements and primarily used for evaluating new parameters.

340 agreement with observation recorded in dataset D2 in which zone 1 had larger
 341 leaf area of $1176 \pm 239 \text{ cm}^2$ on April 17th, in contrast to zone 5 having smaller
 342 leaf area of $1124 \pm 90 \text{ cm}^2$ on the same date. After three weeks on May 9th, the
 343 difference was flipped over where plants in zone 1 had senesced faster, resulting
 344 into a smaller green leaf area of $512 \pm 55 \text{ cm}^2$ while zone 5 was still greener with
 345 $701 \pm 70 \text{ cm}^2$. Mean temperature recorded in dataset D2 for zone 1 was 15.5°C
 346 and zone 5 was 12.6°C .

Figure 7: Comparison of model simulated green leaf area and observation for two temperature zones (Zone 1: high, Zone 5: low) recorded in dataset D2 from a temperature-controlled glass house.

347 3.1.3 Dataset 3 (JS field)

348 Simulated biomass allocation compared with another independent dataset D3
 349 came with an average RMSE of 7.4 g, 3.6 g and 4.2 g for bulb, leaf, and stalk,
 350 respectively (Figure 8). The maximum bulb biomass from simulation was 37.4 g
 351 reached on June 22nd while the observed maximum was 28.7 g measured on
 352 May 26th. The maximum living leaf biomass from simulation was 10.0 g reached
 353 on May 1st and the maximum observed value was 9.1 g on April 28th. The
 354 maximum biomass of stalk from simulation was 13.5 g on May 1st when scape
 355 just appeared and subsequently removed. The maximum stalk biomass recorded
 356 in the dataset was 12.3 g observed on April 28th. Once the scape was removed

and no longer a part of stalk composition, stalk biomass gradually decreased as the mature sheath stopped growing and started senescence. According to the dataset, scape was removed during reproductive stage, but an exact date of the removal was not recorded. Thus we assumed scape removal took place as soon as a tip of the scape became visible out of the whorl which occurred on May 1st by model estimation. The scape biomass at the time of removal was 5.7 g.

Figure 8: Biomass allocation to leaf, bulb, stalk, and visible portion of scape simulated over time compared with measurements from dataset D3 which was from field measurements with scape removal.

3.2 Yield Projection

3.2.1 Gosan

In the current climate condition, fresh yield on ND cultivar in Gosan, Jeju was estimated to be maximum at an average of 6.8 kg m^{-2} with standard deviation of 0.8 kg m^{-2} when planted in late August (240 DOY) which was closely followed by early September planting (260 DOY) with an average of $6.7 \pm 0.7 \text{ kg m}^{-2}$ (Figure 9). A similar level of high yield was maintained until late September then yield gradually decreased with later planting dates. The difference between maximum yield from early planting date and minimum yield from later planting date was 3.7 kg m^{-2} . In the near future from 2020s to 2050s, a similar pattern

was observed that high yield was achieved with early planting in September. An overall yield was increased to $7.5 \pm 0.8 \text{ kg m}^{-2}$ under RCP4.5 and $7.6 \pm 0.8 \text{ kg m}^{-2}$ under RCP8.5 scenario when compared to the current climate. There was no clear difference between RCP4.5 and RCP8.5 scenarios in terms of estimated yield for a given planting date. In the distant future from 2060s to 2090s, RCP4.5 scenario still maintained a similar pattern where high yield was achieved in early planting dates, but then the range of potential high yield was expanded to later planting dates. For example, planting dates from late August (240 DOY) to mid-October (290 DOY) all resulted in an average yield closely ranged from 7.6 kg m^{-2} to 7.7 kg m^{-2} . The difference between maximum and minimum yield estimated in the range of planting dates was reduced to 3.0 kg m^{-2} . Under RCP8.5 scenario, the yield curve was more flattened that the difference was only 1.8 kg m^{-2} between all planting dates. The maximum yield was $8.3 \pm 0.6 \text{ kg m}^{-2}$ in early-October (280 DOY).

Figure 9: Yield estimation within a range of planting dates under current and future climate projection in Gosan, Jeju which is a region grows ND cultivar at commercial scale. The current climate indicates simulation results with 30-years normal weather data from 1980–2010. Shades represent a range of ± 1 standard deviation from mean fresh yield.

When optimal plating dates were assessed from current to future climate conditions by 10 year intervals, a clear trend of shifting towards later planting date was found in the future projection although its strength varied depending on scenarios (Figure 11). Both scenarios began with optimal planting date in early to mid-September then showed a strong divergence at the end of the century that the optimal planting date surfaced from late September to early October under RCP4.5 scenario, while mid-November became a possibility under RCP8.5

scenario.

3.2.2 Chuncheon

In the current climate condition, yield of ND cultivar in Chuncheon, Gangwon was estimated close to nil with a maximum yield $0.06 \pm 0.05 \text{ kg m}^{-2}$ due to impeded leaf growth and increased mortality from cold winter temperature during early vegetative growth (Figure 10). However, under future projections, yield became more viable thanks to warmer climate. In the near future from 2020s to 2050s, maximum estimated yield was $1.5 \pm 0.7 \text{ kg m}^{-2}$ under RCP4.5 and $1.3 \pm 0.9 \text{ kg m}^{-2}$ under RCP8.5 scenario when both planted in early September (250 DOY). In the distant future from 2060s to 2090s, more yield was achievable with $2.5 \pm 0.8 \text{ kg m}^{-2}$ under RCP4.5 and $4.3 \pm 1.1 \text{ kg m}^{-2}$ under RCP8.5 scenario with early to mid-September plating dates, 250 DOY and 240 DOY, respectively. The difference between maximum and minimum yield estimated in the range of planting dates was 0.6 kg m^{-2} in early period and 1.0 kg m^{-2} in late period under RCP4.5 scenario. RCP8.5 scenario initially had a similar range of estimated yield over multiple planting dates with 0.5 kg m^{-2} in early period, but then showed much higher variance of 1.7 kg m^{-2} in late period.

Figure 10: Yield estimation within a range of planting dates under current and future climate projection in Chuncheon, Gangwon where current climate is not favorable for growing ND cultivar. The current climate indicates simulation results with 30-years normal weather data from 1980–2010. Shades represent a range of ± 1 standard deviation from mean fresh yield.

In terms of optimal planting dates for each period, planting in around early September consistently turned out to work best regardless of climate scenarios

and estimated yield gradually declined with later planting dates in all scenarios (Figure 11).

Figure 11: Optimal planting dates estimated for each year under current and future climate projections in Gosan and Chuncheon. Year 2010 indicates simulation results with 30-years normal weather data from 1980–2010 representing current climate. For RCP scenarios, each year indicates 10-year average, *e.g.*, 2090 refers to 2090–2100. Error bar and shades represent a range of ± 1 standard deviation from mean optimal planting date.

4 Discussion

4.1 Model Validation

4.1.1 Parameter Calibration

Our parameter set for ND cultivar was mostly derived from existing parameters for Korean Mountain (KM) and Shantung Purple (SP) with small changes (Hsiao et al., 2019). The final parameter set used in our simulation turned out to be similar to one calibrated for SP cultivar. It would not be unexpected given that ND was originated from a Chinese cultivar and hence there is a chance SP might be also related with this original cultivar due to geographical proximity (Kim et al., 2009).

4.1.2 Overestimation

Simulation with the calibrated parameter set often resulted in overestimation in a sense that model output tends to touch upper boundaries of observed data

points (Figure 4,5,7,8). This overestimation is intended and expected because our model estimates for biomass and yield potential for a given environmental condition with assuming no limiting factors such as water and nutrient while the field conditions are likely less than optimal. Some types of overestimation also presumably came from difference between physical measuring methods and modeled algorithms. For example, green leaf area recorded in the dataset was measured by taking only non-withered leaves at the time of observation which is prone to lose leaves with some green parts intact. On the other hand, green leaf area from the model was calculated for individual leaves at a fractional scale by tracking current green portion of the leaf for each time step. These difference could have been one of the reasons leading to overestimation in leaf area simulation (Figure 4,7).

4.1.3 Temperature Regimes

Evaluation of model response under slightly separate temperature regimes as conducted in temperature gradient house provided an insight how elevated air temperature under RCP scenarios would affect plant growth (Figure 7). In short, plant will grow larger and faster and die earlier in warmer conditions. An average of 3 °C difference in temperature led to more than two weeks of shift in growth peak and 15 % change in total green leaf area. Subsequent changes in senescence timing would imply a need for finding out optimal harvest dates which we assumed as a rather constant management decision in our simulation.

4.1.4 Storage Condition

Storage duration (SD) affects maximum leaf tip appearance rate ($LTAR_{max}$) via the dynamic phyllochron model (Equation 1,2,3). Storage temperature (ST) along with storage duration then decides number of leaves initiated inside a seed bulb at the time of planting. Despite their importance on phenology, there often exists no record on when seed bulbs were harvested and in which condition they were stored until the date of planting. With controlled experiments on storage condition, we could have better understanding on how leaves are initiated during storage period and whether more sophisticated approaches like a dynamic plastochron would be necessary.

459 **4.1.5 Other Environmental Cues**

460 While our assessment for yield projection under future condition was primarily
 461 driven by model response to temperature regimes, there is still room for con-
 462 sidering other environmental cues. For instance, we did not have a separate
 463 vernalization process in the model, but assumed winter chilling was always at
 464 an adequate level for triggering continuous development in the following warm
 465 spring. While it worked reasonable well in most cases, some processes like mor-
 466 tality from cold stress could capitalize on this additional cue to make dormant
 467 plants less vulnerable to cold damage, for instance. Soil water availability would
 468 be another important cue in terms of assessing irrigation requirement under
 469 future climate conditions given that garlic plots are often irrigated to prevent
 470 water deficiency during reproductive stage.

471 **4.2 Yield Projection**

472 **4.2.1 Gosan**

473 Jeju, including Gosan, is a region where southern-type garlic cultivars like Namdo
 474 (ND) has been grown at commercial scale thanks to warm climate. Northern
 475 regions in Korean peninsula often have colder winter which prevents growing
 476 southern-type cultivars. Growers instead choose northern-type cultivars for cold
 477 hardiness. As southern cultivars have advantage of higher yield and shorter
 478 growing season, whether they could be adapted to northern regions has been
 479 an important question to many growers and stakeholders. According to our
 480 simulation results using a process-based model, ND cultivar would continue
 481 growing well with a slightly higher yield (Figure 9). The optimal planting date
 482 in early September estimated for the current climate condition was indeed close
 483 to the current practice held in Jeju. This practice is, however, expected to be
 484 delayed in the near future if growers keep trying to maximize yield under both
 485 climate scenarios (Figure 11). The amount of planting date delay depends on
 486 the scenarios and the model estimated that RCP4.5 scenario could see a shift
 487 less than a month while RCP8.5 scenario could potentially result in two months
 488 of late planting. Note that although optimal planting dates were seemingly
 489 changed drastically, yield estimation curve versus planting date was actually
 490 getting more flattened and the difference between maximum and minimum
 491 yield became smaller in warmer climate conditions. Such changes imply that
 492 growers would have more options to choose planting date better suits their

own needs. For example, October planting in the future would still promise a yield level close to potential maximum while taking a relatively shorter growing season to help reducing overall cost of labor and resources. High yield does not necessarily lead to high profitability when longer growing season increases overall water demand entailed by higher irrigation cost for compensation (Lobell, 2014). Looking further in distant future, it was more clear to expect a shift in optimal planting to later dates although yield difference to earlier planting dates was not significant.

4.2.2 Chuncheon

Chuncheon is located in the middle of Korean peninsula and its continental climate frequently experiences below freezing temperature during winter season. Hence only northern-type cultivar adapted to cold climate have been grown in this region. Our simulation results confirmed expected yield for the current climate was almost non-existent as most plants subject to mortality (M) could not survive after episodes of extreme cold. By contrast, future climate conditions were projected to become more favorable in both scenarios so that expecting some tangible yields would be at least feasible (Figure 10). Generally early planting dates were favored in terms of optimal yield as similar in the current growing region like Jeju (Figure 11). Although the level of estimated yield was still lower when compared to the yield currently obtainable in other established regions, growers may be able to take advantage of this new opportunity for expanding crop portfolio during winter season.

4.3 Climate Adaptation

It is important to tease apart an effect of adaptation introduced by planting date shift in the future from an impact of planting date shift in the current condition (Lobell, 2014; Challinor et al., 2018). According to the result from Gosan, planting date shift in established growing region has little effect on adaptation as evidenced by a similar or smaller range between maximum and minimum yield when comparing current and future climate scenarios (Figure 9). Higher estimated yield was a result of increased productivity under more favorable condition primarily due to higher temperature and elevated CO_2 , but not from potential phenological changes occurred by planting date shift. In non-established regions like Chuncheon, however, an effect of adaptation increased virtually from zero to a substantial amount when transitioning to new climate condition as

evidenced by a rising slope of yield estimation curve (Figure 10). In other words, the importance of planting date as a climate adaptation strategy depends on how crops are adapted to the local condition.

Underlying uncertainties of crop models often hinder model driven assessment of climate change adaptation (Rosenzweig et al., 2014; Holzkämper et al., 2015; Corbeels et al., 2018). We adopted a process-based model with coupled gas-exchange to minimize uncertainties in physiological responses to temperature and CO₂, but there still remain many processes that can use improvements. For instance, water stress response was solely dependent on leaf water potential disconnected with soil water status albeit a less practical implication when garlic plants are generally irrigated to meet high water demand during reproductive stage. Nitrogen response is another important factor that the model currently does not take into account and instead assume non-limiting due to extensive use of fertilizers in practice, but will be critical for assessing an exact cost of production in finding optimal yield.

Conflict of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Funding

The information, data, or work presented herein was funded in part by the Cooperative Research Program for Agricultural Science and Technology Development, Rural Development Administration, Republic of Korea under Grant Number PJ015124012021. The views and opinions of authors expressed herein do not necessarily state or reflect those of the funding agency.

References

- Abdalla, M. M. A., Aboul-Nasr, M. H., and Metwaly, A. K. (2011). Growth and yield of fifteen garlic ecotypes. *Assiut Journal of Agricultural Sciences* 36, 52–65

Challinor, A. J., Müller, C., Asseng, S., Deva, C., Nicklin, K. J., Wallach, D.,
et al. (2018). Improving the use of crop models for risk assessment and climate
change adaptation. *Agricultural Systems* 159, 296–306

Choi, S. and Baek, J. (2016). Garlic yields estimation using climate data. *Journal
of the Korean Data and Information Science Society* 27, 969–977

Corbeels, M., Berre, D., Rusinamhodzi, L., and Lopez-Ridaaura, S. (2018). Can
we use crop modelling for identifying climate change adaptation options?
Agricultural and Forest Meteorology 256, 46–52

Cropbox.jl (2021). <https://github.com/cropbox/Cropbox.jl>

[Dataset] FAO (2020). FAOSTAT

Garlic.jl (2021). <https://github.com/cropbox/Garlic.jl>

Giménez, C., Stöckle, C., Suárez-Rey, E., and Gallardo, M. (2016). Crop yields
and n losses tradeoffs in a garlic–wheat rotation in southern spain. *European
Journal of Agronomy* 73, 160–169

Gómez, D., Salvador, P., Sanz, J., and Casanova, J. L. (2021). Regional
estimation of garlic yield using crop, satellite and climate data in mexico.
Computers and Electronics in Agriculture 181, 105943

Heo, I., Kwon, W.-T., Chun, Y., and Lee, S. (2006). The impact of temperature
rising on the distribution of plant-in case of bamboos and garlics. *Journal of
Environmental Impact Assessment* 15, 67–78

Holzkämper, A., Calanca, P., Honti, M., and Fuhrer, J. (2015). Projecting
climate change impacts on grain maize based on three different crop model
approaches. *Agricultural and Forest Meteorology* 214, 219–230

Hsiao, J., Yun, K., Moon, K. H., and Kim, S.-H. (2019). A process-based model
for leaf development and growth in hardneck garlic (*Allium sativum*). *Annals
of Botany* 124, 1143–1160

IPCC (2013). *Climate Change 2013: The Physical Science Basis. Contribution
of Working Group I to the Fifth Assessment Report of the Intergovernmental
Panel on Climate Change*. Tech. rep.

585 IPCC (2014). *Climate Change 2014: Synthesis Report. Contribution of Working*
586 *Groups I, II and III to the Fifth Assessment Report of the Intergovernmental*
587 *Panel on Climate Change*. Tech. rep.

588 Kamenetsky, R. (2007). *Garlic: Botany and Horticulture* (John Wiley & Sons,
589 Ltd), chap. 2. 123–172

590 Kim, C.-H., Seong, K.-C., Lee, J.-S., Kang, K.-H., Um, Y.-C., and Suh, H.-D.
591 (2009). Production of seed garlic by sawing bulbils of southern type garlic in
592 jeju island. *Protected Horticulture and Plant Factory* 18, 74–80

593 Kim, S.-H., Jeong, J. H., and Nackley, L. L. (2013). Photosynthetic and tran-
594 spiration responses to light, CO₂, temperature, and leaf senescence in garlic:
595 Analysis and modeling. *Journal of the American Society for Horticultural*
596 *Science* 138, 149–156

597 Lee, J.-W., Park, G.-A., Joh, H.-K., Lee, K.-H., Na, S.-I., Park, J.-H., et al.
598 (2011). Analysis of relationship between vegetation indices and crop yield
599 using kompsat (koreamulti-purpose satellite)-2 imagery and field investigation
600 data. *Journal of The Korean Society of Agricultural Engineers* 53, 75–82

601 Lobell, D. B. (2014). Climate change adaptation in crop production: Beware of
602 illusions. *Global Food Security* 3, 72–76

603 Marcelis, L., Heuvelink, E., and Goudriaan, J. (1998). Modelling biomass
604 production and yield of horticultural crops: a review. *Scientia Horticulturae*
605 74, 83–111

606 Moon, K., Seo, H., Shin, M., Song, E., and Oh, S. (2019). Generation of hourly
607 weather data using daily data. In *Proceedings of The Korean Society of*
608 *Agricultural and Forest Meteorology Conference*. 195–196

609 Nackley, L. L., Jeong, J. H., Oki, L. R., and Kim, S.-H. (2016). Photosynthetic
610 acclimation, biomass allocation, and water use efficiency of garlic in response to
611 carbon dioxide enrichment and nitrogen fertilization. *Journal of the American*
612 *Society for Horticultural Science* 141, 373–380

613 Põldma, P., Merivee, A., Pae, A., and Justus, K. (2005). Influence of planting
614 time on the development, yield and quality of garlic (*Allium sativum* L.) in
615 estonia. *Acta Horticulturae* , 333–338

616 Portela, J. A., Sidoti, B., Reybet, G., Bellacomo, C., and Astorquiza, R.
617 (2012). Yield stability of ten garlic (*Allium sativum*) clonal cultivars in northern
618 patagonia, argentina. In *VI International Symposium on Edible Alliaceae* 969.
619 107–112

620 [Dataset] RDA (2019). AgClimate data portal

621 Rizzalli, R., Villalobos, F., and Orgaz, F. (2002). Radiation interception,
622 radiation-use efficiency and dry matter partitioning in garlic (*Allium sativum*
623 l.). *European Journal of Agronomy* 18, 33–43

624 Rosenzweig, C., Elliott, J., Deryng, D., Ruane, A. C., Müller, C., Arneth, A.,
625 et al. (2014). Assessing agricultural risks of climate change in the 21st century
626 in a global gridded crop model intercomparison. *Proceedings of the National*
627 *Academy of Sciences* 111, 3268–3273

628 Takagi, H. (1989). *Onions and allied crops* (CRC Press), vol. 3, chap. Garlic
629 *Allium sativum* L. 109–157

630 Villalobos, F., Testi, L., Rizzalli, R., and Orgaz, F. (2004). Evapotranspiration
631 and crop coefficients of irrigated garlic (*Allium sativum* l.) in a semi-arid
632 climate. *Agricultural Water Management* 64, 233–249

633 Yun, K., Timlin, D., and Kim, S.-H. (2020). Coupled gas-exchange model for c_4
634 leaves comparing stomatal conductance models. *Plants* 9, 1358