Original Research Article

Augmenting Recombinant Antibody Production in HEK293E Cells: Optimising Transfection and Culture Parameters

Zealyn Shi-Lin Heng ¹, Joshua Yi Yeo ¹, Darius Wen-Shuo Koh ¹, Samuel Ken-En Gan ^{1,2,3,*} and Wei-Li Ling ^{1,*}

¹Antibody & Product Development Lab, EDDC-BIL, Agency for Science, Technology and Research (A*STAR), Singapore 138672, Singapore, hslzealyn@gmail.com (Z.S.-L.H.); joshua_yeo@eddc.a-star.edu.sg (J.Y.Y.); darius_koh@eddc.a-star.edu.sg (D.W.-S.K.)

²APD SKEG Pte Ltd., Singapore 439444, Singapore.

³James Cook University, Singapore 387380, Singapore.

*Correspondence: samgan@apdskeg.com (S.K.-E.G.); alexling1989@gmail.com (W.-L.L.)

47 **ABSTRACT**

48

49 **Background**

50 Optimising recombinant antibody production is important for cost-effective therapeutics and
 51 diagnostics. With impact on commercialisation, higher productivity beyond laboratory scales
 52 is highly sought, where efficient production can also accelerate antibody characterisations
 53 and investigations.

54 **Methods**

55 Investigating HEK293E cells for mammalian antibody production, various transfection and
 56 culture parameters were systematically analysed for antibody light chain production before
 57 evaluating them for whole antibody production. Transfection parameters investigated include
 58 seeding cell density, the concentration of the transfection reagent and DNA, complexation
 59 time, temperature, and volume, as well as culture parameters such as medium replacement,
 60 serum deprivation, use of cell maintenance antibiotic, incubation temperature, medium
 61 volume, post-transfection harvest day and common nutrient supplements.

62 **Results**

63 Using 2 mL adherent HEK293E cell culture transfections with 25 kDa linear
 64 Polyethylenimine in the most optimised parameters, we demonstrated a ~2-fold production
 65 increase for light chain alone and for whole antibody production reaching 536 and 49 µg
 66 respectively in a cost-effective manner. With the addition of peptone, κ light chain increased
 67 by ~4-fold to 1032 µg while whole antibody increased to a lesser extent by ~2.5-fold to 51 µg,
 68 with benefits potentially for antibodies limited by their light chains in production.

69 **Conclusions**

70 Our optimised findings show promise for a more efficient and convenient antibody
 71 production method through transfection and culture optimisations that can be incorporated to

72 scale up processes and with potential transferability to other mammalian-based recombinant
73 protein production using HEK293E cells.

74

Statement of Significance

Recombinant antibody production is crucial for antibody research and development. Systematically investigating transfection and culture parameters such as PEI/DNA concentrations, complexation time, volume, and temperature, supplements, etc., we demonstrated a ~4-fold light chain alone production increase to 1032 µg and a 2.5-fold whole antibody production increase to 51 µg from 2 mL transfections.

75

76 **Key words:** Recombinant antibody production; Antibody therapeutics manufacturing;

77 Transient transfection; Transfection parameters; HEK293E

78

79 INTRODUCTION

80 The efficient production of recombinant monoclonal antibodies underpins their
81 functional and structural investigations (1) such as toxicology and drug formulation studies (2)
82 and thereby, success of both therapeutic and diagnostic antibodies development. With a direct
83 impact on production costs during development, optimisations of the already expensive
84 mammalian production of recombinant antibodies are highly desired.

85 Currently, whole recombinant antibodies are predominantly produced by mammalian
86 cells, given the need for complex folding, assembly and post-translational modifications to
87 generate functionally active proteins (3, 4). Typically, the protein genes of interest are first
88 transfected into mammalian cells, followed by the selection and generation of stable cell lines
89 expressing the gene to produce the protein of interest (2). As this process is often tedious and
90 laborious, the transient gene expression method on easily transfected cell lines is typically

91 used for research scale investigations. One of the most widely used is the Human Embryonic
 92 Kidney 293 (HEK293) family of cell lines with several genetic variants developed to enhance
 93 protein expressions such as the HEK293E expressing the Epstein-Barr Virus Nuclear
 94 Antigen-1 (5), utilised in this study, and HEK293FT and HEK293T expressing the SV40 T-
 95 antigen (6). Given the scalability of HEK293 (3, 7, 8), they are suitable for producing a wide
 96 range of recombinant proteins including whole viral particles (9).

97 Given that transient recombinant protein production depends on a myriad of
 98 transfection and culture parameters, optimising these parameters directly influences the
 99 scalability and efficiency of recombinant protein production. Transfection parameters include
 100 seeding cell density (10), transfection reagent concentration (11), DNA concentration (12),
 101 complexation time, temperature, and volume (12), together with culture parameters such as
 102 medium replacements (7, 13), serum deprivation (14), use of cell line maintenance antibiotic
 103 (15), incubation temperature (16, 17), culture volume, harvesting time (18–20), and nutrient
 104 supplementation (21–23). Three main strategies are commonly used to optimise these
 105 parameters: the reductionist “one factor at a time” approach (24), stepwise optimisation from
 106 one condition to another (25), and the design of experiment approach (10), of which we took
 107 the single isolated factor approach.

108 In this study, the key drivers were determined by the “one factor at a time” approach
 109 and subsequently combined to maximise antibody production (24). We optimised individual
 110 parameters using Pertuzumab (Perjeta®) antibody light chains (Variable κ -1 or V κ 1), given
 111 that light chains are pre-requisites for whole antibody production (26, 27) and to mitigate the
 112 lopsided effect of dual plasmid transfections. This was followed by validating the combined
 113 parameters using recombinant whole Pertuzumab antibodies produced using two separate
 114 variable kappa (V κ)-1 and variable heavy (VH)-3 plasmids. Based on our previous study
 115 where we generated Pertuzumab variants of VH family 1 to 7 and V κ 1 to 6 frameworks via

complementarity determining regions (CDRs) grafting (28, 29), we analysed light chains known to limit antibody production i.e. the V κ 5 grafted Pertuzumab light chain as a pairing partner.

The final combination of optimised parameters for whole antibody production would complement our previous findings for production that discusses the role of heavy chain constant (30), V κ -VH pairings (29), amino acid usage with leader selection (28) in a holistic manner (1, 31) to significantly boost antibody production for antibody-based therapeutics and diagnostics.

MATERIALS & METHODS

Plasmid DNA

Pertuzumab light chain of V κ 1 region and heavy chain IgG1 of the VH3 region genes, including their CDR-grafted V-region variants, designated V κ 2-6 and VH1, 2, 4-7 were ligated into pTT5 plasmid (YouBio, Cat no. VT2202) and prepared as previously described (28–30, 32, 33). Trastuzumab plasmid variants in Supplementary Figure 1B were prepared similarly. Optimisation of light chain production was performed by single plasmid transfection of Pertuzumab light chain V κ 1 prior to optimising whole V κ 1|VH3 antibody production by dual plasmid transfection with its corresponding heavy chain VH3. To investigate if the optimised parameters can boost production of antibodies limited by light chains, Pertuzumab V κ 5 was paired with VH3 for whole V κ 5|VH3 antibody production. Comparisons of culture incubation temperatures performed using other antibody variants in large flasks are shown in Supplementary Figure 1.

Cell culture conditions

HEK293E adherent cells were routinely cultured in DMEM medium (Gibco, Cat no. 11965-084), supplemented with 10 % FBS (Gibco, Cat no. 10270-106) or 10 % Ultra Low IgG FBS (Gibco, Cat no. 16250078), 1 mM Sodium Pyruvate (Gibco, Cat no. 11360-070), 100 units/mL penicillin, 100 µg/mL Streptomycin (Nacalai tesque, Cat no. 09367-34), G418 (Roche, Cat no. 04727894001) and incubated at 37 °C in 5 % CO₂.

Transfection Parameters

Linear 25 kDa Polyethylenimine (PEI) (Polysciences, Cat no. 23966-1) at pH 7.0 was used in the transfections. 2 mL and 40 mL scale transfections of HEK293E cells were performed in 6-well plates (Costar, Cat no. 3516) and T175 flasks (Greiner Bio-one, Cat no. 660175) respectively, with at least three independent replicates. The conditions tested for the transient gene expression in 6-well plate format included: Cell seeding densities from 1 - 10 × 10⁵ cells/mL; PEI concentrations from 1 - 10 µg/mL; DNA concentrations from 0.25 - 2 µg/mL; PEI-DNA complexation time from 5 - 120 minutes; PEI-DNA complexation temperatures of 18 - 45 °C; PEI-DNA complexation in 1 - 25 % serum free medium (SFM) of total 2 mL volume; PEI-DNA complexation in light or dark environments; culture medium replacements; serum deprivation before transfection; culture with or without G418 antibiotic; culture incubation temperature at 32 or 37 °C; culture volume from 2 - 5 mL; post-transfection harvest of 3 - 21 days. Experimental conditions were compared against control parameters of: Seeding density at 2 × 10⁵ cells/mL; DNA concentration at 1 µg/mL; PEI concentration at 2 µg/mL; PEI-DNA complexation time at 30 minutes in 18 °C; with 10 % SFM of total 2 mL volume, performed in the dark and added into cultures without G418; with the cultures grown at 37 °C and harvesting made on day 14 post-transfection, as were performed routinely in previous work (28–30, 34).

For cell culture conditions, 10 % FBS DMEM was used for light chain production, while 10 % Ultra Low IgG FBS DMEM was used for whole antibody IgG production. The culture conditions used for Supplementary Figure 1 were scaled up proportionately from the 6-well cultures for use at T175 flask scale transfections.

Supplement Preparation and Usage

The nutrient supplements tested were: OmniPur[®] Casamino acids (CA) (EMD Chemicals, Cat no. 65072-00-6), Casein from Bovine Milk (CS) (Sigma-Aldrich, Cat no. C3400), Bacto[™] Peptone (PT) (Becton, Dickinson and Company, Cat no. 21167), Skim Milk (SM) (Millipore, Cat no. 70166), Tryptone (TT) (Bio basic, Cat no. TG217 (G211)), Bacto[™] Yeast Extract (YE) (Becton, Dickinson and Company, Cat no. 212750) and Impact Essential Amino Acids (EA) (MYPROTEIN[™])(28). Supplements were dissolved in water or Phosphate Buffer Saline (PBS), filtered using 0.22 µm filters and stored at 4 °C before use.

The supplements were added post-transfection of Pertuzumab light chain cultures on day 7 at 1 mg/mL final concentration. Further optimisations with peptone were performed at 1, 2, 4 and 6 mg/mL, added on days 1 – 10. Subsequently, the combined optimised parameters of 1) without nutrient supplementation; 2) peptone at 1 mg/mL; 3) essential amino acids at 3.5 mg/mL (28) or 4) both were tested for Pertuzumab whole antibody production (Vκ1|VH3) and light chain limiting variant (Vκ5|VH3) with the supplementations (conditions 2,3, and 4) added on day 7 post-transfection.

Recombinant Antibody Harvest and Titre Quantification

Cell culture supernatants were harvested on various days stipulated by the experimental conditions and at the end of 14 days, and stored at -20 °C before further testing. In the small-scale 2 mL experiment set-ups where evaporation could be a potential

confounding factor, total protein production was calculated from the concentration and remaining volumes measured by a micropipette at the point of harvest. The light chain and whole antibody quantification of the cell culture supernatants were performed using Octet Red 96 instrument as previously described (28–30), with Protein L (PpL) and G (SpG) biosensors (ForteBio, Cat no. 18-5085 and 18-5082) respectively.

Data Analysis

The data were analysed and plotted using Microsoft Excel 365 and GraphPad Prism Version 9.0.0. Protein titre or total protein production data were normalised against the control and expressed as a percentage with the control set at 100 %. One-way ANOVA with Dunnett's comparison test against the control was used when examining three or more different parameters, while a two-tailed unpaired parametric t-test was performed when comparing two parameters.

RESULTS

Optimising Transfection Parameters

Cell seeding densities

Cell seeding densities ranging from 1 - 10×10^5 cells/mL in a 6-well plate were evaluated with respect to their impact on Pertuzumab light chain production and normalised against that of the control condition of 2×10^5 cells/mL set at 100 % in **Figure 1A**. The one-way ANOVA showed no significant differences ($p > 0.05$) in Pertuzumab light chain production across the seeding densities tested despite a decreasing trend above 6×10^5 cells/mL, where production at 10×10^5 cells/mL was only 41 % (70 µg/mL) that of the control (177 µg/mL). The average titre data with accompanying standard error are shown in Supplementary Table 1.

214

215 *PEI and DNA concentrations*

216 PEI concentrations ranging from 1 - 10 µg/mL in 6-well plate transfections were
 217 tested for Pertuzumab light chain production and normalised against that of the control
 218 condition of 2 µg/mL set at 100 % in **Figure 1B**. The results showed a curvilinear
 219 relationship between PEI concentration and light chain production peaking at 145 % (206
 220 µg/mL) for 4 µg/mL of PEI that was significant ($p < 0.05$) with t-test but not the Dunnett's
 221 test against the control (140 µg/mL). PEI concentrations lower or higher than 4 µg/mL
 222 negatively impacted light chain production with 1 µg/mL having the lowest production of 17 %
 223 (22 µg/mL) significant at $p < 0.05$ when analysed with Dunnett's test. Concentrations lower
 224 than 4 µg/mL also had lower production rates despite not being significant by Dunnett's test.

225 DNA concentrations ranging from 0.25 - 2 µg/mL, at 0.25 µg/mL intervals in 6-well
 226 plate transfections, were evaluated for Pertuzumab light chain production and normalised
 227 against that of the control condition of 1 µg/mL set at 100 % in **Figure 1C**. An inverse
 228 relationship between DNA concentration and light chain production was observed with the
 229 lowest and highest DNA concentration (0.25 and 2 µg/mL) corresponding to the best and
 230 poorest light chain production at 190 % (399 µg/mL) and 21 % (64 µg/mL) respectively that
 231 were significant at $p < 0.05$ when analysed with Dunnett's test against the control (208
 232 µg/mL).

233

234 *PEI and DNA complexation*

235 PEI-DNA complexation time from 5 - 120 minutes in 6-well plate transfections was
 236 evaluated with respect to Pertuzumab light chain production and normalised against that of
 237 the control condition of 30 minutes, set at 100 % in **Figure 1D**. An inverse relationship
 238 between complexation time and light chain production showed the shortest and longest

incubation time (5 and 120 minutes) corresponding to the highest and lowest production (166 % and 32 %, equivalent to a titre of 271 and 54 µg/mL respectively), which were both significant at $p < 0.05$ when analysed with Dunnett's test against the control (165 µg/mL) production.

PEI-DNA complexation temperature from 18 to 45 °C in 6-well plate transfections was evaluated with respect to Pertuzumab light chain production and normalised against that of the control condition of 18 °C, set at 100 % in **Figure 1E**. An inverse relationship where the decline in light chain production was found with increasing temperatures. Dunnett's test against the control for 25, 32, and 37 °C showed no significant decrease ($p > 0.05$) despite slight reductions. However, significant decreases ($p < 0.05$) were observed for 40 and 45 °C, at 54 % (105 µg/mL) and 50 % (98 µg/mL) of the control (164 µg/mL) production respectively.

PEI-DNA complexation volume using SFM from 1 to 25 % of the final total culture medium in 6-well plate transfections was evaluated with respect to Pertuzumab light chain production and normalised against that of the control condition of 10 % final cell culture volume (20) set at 100 % in **Figure 1F**. PEI-DNA complexation medium volume was found to increase light chain production with the lowest 1 % SFM yielding the lowest production of 12.7 % (30 µg/mL, $p < 0.05$) increasing to the highest production of 131 % (273 µg/mL) at 25 % SFM ($p > 0.05$). Nonetheless, production was observed to peak at 15 % transfection volume plateauing at higher complexation volumes.

To study photodegradation assumptions of transfection complexes (35), we performed PEI-DNA complexation incubation for light chain using the 6-well plate format under the biosafety cabinet (BSC) light and without (in very dim external room light). There was no significant difference ($p > 0.05$) in antibody light chain production between the two conditions (**Figure 1G**) by t-test.

264

265 **Optimising Culture Conditions**

266 *Culture medium replacements*

267 Fresh medium replacement before or after transfection, as well as both before and
 268 after, were tested for Pertuzumab light chain using the 6-well plates and normalised against
 269 that of the control condition of no medium replacement set at 100 % in **Figure 2A**.
 270 Pertuzumab light chain production was found to increase by 36 % and 9 % (reaching 320 and
 271 250 µg/mL) for before, and both before and after transfection, respectively. A slight
 272 reduction of 7 % to 235 µg/mL was found for a medium replacement after transfection when
 273 compared to the control (226 µg/mL), but the one-way ANOVA showed that these are not
 274 significant ($p > 0.05$). The average titre data with accompanying standard error are shown in
 275 Supplementary Table 1.

276 We next studied serum deprivation by seeding cells in 10% FBS supplemented or
 277 serum free medium a day before transfection in 6-well plate format to test for Pertuzumab
 278 light chain production. An additional condition of seeding cells in serum free medium a day
 279 before transfection followed by replacement with 10% FBS supplemented medium a day
 280 after transfection was included. The serum deprivation condition data was normalised against
 281 that of no serum deprivation set at 100 % in **Figure 2B**. We found the prolonged and short-
 282 term serum deprivation to result in a non-significant ($p > 0.05$) 25 % and 37 % decrease in
 283 light chain production using Dunnett's test against the control.

284

285 *Culture with G418 antibiotics*

286 Studying the effect of cell culture antibiotics on Pertuzumab light chain production in
 287 6-well plates, we did not find G418 to have significant effects ($p > 0.05$) on the light chain
 288 production by t-test (**Figure 2C**).

289

290 *Culture temperature*

291 Culture incubation temperatures at 32 and 37 °C for Pertuzumab light chain 6-well,
292 transfections were shown with t-tests to reveal a significant production decrease at 32 °C ($p <$
293 0.05). 2 mL cultures grown in 32 °C resulted in only 53 % (120 µg) of the light chain
294 produced at 37 °C, set at 100 % (231 µg) (**Figure 2D**). The average total light chain data with
295 standard error are shown in Supplementary Table 2. Further testing with a full panel of whole
296 recombinant Pertuzumab and Trastuzumab variants (VH1-7 and VK1-6 combinations)
297 transfected in T175 flasks did not show a significant difference ($p > 0.05$) between the two
298 temperatures (Supplementary Figure 1).

299

300 *Culture medium volumes*

301 Medium volumes varying from 2 - 5 mL in 6-well plate transfections were evaluated
302 for Pertuzumab light chain production and normalised against that of the control condition of
303 2 mL set at 100 % in **Figure 2E**. The Dunnett's test showed a significantly lower difference
304 ($p < 0.05$) when using a culture volume of 1 mL, producing 148 µg of total light chain
305 compared to the control of 300 µg from a 2 mL culture. While there seems to be a gradual
306 increase in light chain production for higher volumes used, with the highest medium volume
307 of 5 mL showing a rise of 40 % (reaching 413 µg), it was not significantly different ($p > 0.05$)
308 from the control.

309

310 *Culture supernatant harvest day*

311 Post-transfection harvest of culture supernatants across 3 – 21 days from Pertuzumab
312 light chain 6-well plate transfections were normalised against that of the control condition of
313 14 days set at 100 % in **Figure 2F**. We found a direct relationship between the day of

supernatant harvesting and light chain production where total light chain from 2 mL cultures increased from 81 µg on day 3 to plateau at 522 µg from day 14 (Supplementary Table 2). With analysis by Dunnett's test against harvest Day 14, the control, there was an expected significantly lower ($p < 0.05$) light chain production on harvest days 3, 6, and 9. No significant increase ($p > 0.05$) was detected after day 14.

Nutrient Supplements

A variety of culture supplements commonly used to improve recombinant protein production were evaluated for Pertuzumab light chains and normalised with respect to the control condition of no supplement (NS) set at 100 % in **Figure 3A**. Adding the supplements to post-transfected cultures on day 7 at a final concentration of 1 mg/mL, the one-way ANOVA showed no significant difference ($p < 0.05$) across the supplements tested. Given that peptone is the most common supplement of mammalian recombinant protein production (25, 36–38), further investigations were performed using a two-tailed unpaired t-test against the non-supplemented control, revealing that it significantly boosted ($p < 0.05$) light chain production by 25 % (**Figure 3A**) to 351 µg in an initial 2 mL culture (Supplementary Table 2). A similar t-test was performed for yeast extract showing no significant ($p > 0.05$) increases. Further optimisations of concentration and timing of peptone addition showed that 1 mg/mL, added on post-transfection days 5, 6, or 7, gave the highest boost to light chain production (**Figure 3B**). Single supplementation of peptone at 1 mg/mL alone on day 1 or day 7 was found sufficient to enhance light chain production (**data not shown**), showing that repeated addition of low concentrations of peptone was not necessary when a mid-point culture addition on day 7 was sufficient.

Combination of Transfection and Culture Parameters

From 2 mL culture optimisations of the individual transfection and culture parameters in **Table 1**, the combined parameters for Pertuzumab light chain production led to a 217 % significant increase ($p < 0.05$) to 536 μg . This increase was further enhanced ~2-fold to 452 % reaching 1032 μg after peptone supplementation (**Figure 4A** and Supplementary Table 3). Validating this on whole Pertuzumab antibody production in 2 mL cultures, a significant increase ($p < 0.05$) of 236 % to 49 μg was elicited from the control of 22 μg . With peptone supplementation, a subtle increase of 252 % was observed, reaching 51 μg (Supplementary Table 3).

One-way ANOVA showed no significant difference ($p > 0.05$) in total whole antibody produced from initial 2 mL cultures with PEI-DNA complexation incubation times of 5, 15, and 30 minutes in **Figure 4B**. Following on from our previous work where 3.5 mg/ml of essential amino acids (EA) showed improvements in Pertuzumab antibody production (28), EA supplementation was added to our evaluation where we found an unexpected significant decrease ($p < 0.05$) of 13 % by Dunnett's test, from an optimised control of 44 to 38 μg . (**Figure 4C** and Supplementary Table 3). Dual supplementations of both peptone and EA gave a significant increase ($p < 0.05$) of 23 % to 54 μg , with peptone alone giving a significantly higher ($p < 0.05$) increase of 47 % to 65 μg (**Figure 4C** and Supplementary Table 3). Comparing the single and dual addition of peptone and EA to boost light chain limiting antibodies (Pertuzumab V κ 5|VH3), we found peptone alone to significantly improve ($p < 0.05$) production by 28 % but not for other supplements. The average total protein data with accompanying standard error are shown in Supplementary Table 3.

DISCUSSION

The cost-effective and efficient scalable production of recombinant monoclonal antibodies is key to the commercial success of both therapeutic and diagnostic antibodies. For

this purpose, we set out to optimise transfection and culture parameters with relevance beyond antibody production to that of recombinant proteins using mammalian cell culture. Whole recombinant antibody production is affected by many factors such as the constant region (30) and CDRs of the V κ and V H chains (29), their pairings (29), deletions in the V κ (34), the signal peptides and amino acid usage (28). Given the importance of light chains for whole antibody production (26, 27) and to mitigate the possible lopsided effect of dual plasmid transfection, we performed single parameter optimisations on Pertuzumab light chain production. Thereafter, we applied the combined optimised parameters to produce the corresponding Pertuzumab.

Counter-intuitively to the assumption where higher cell seeding density could increase antibody production (39), we did not find significant differences in light chain production levels over the range of tested seeding densities (**Figure 1A**), hinting at the influence of other factors. In fact, high seeding cell densities significantly decreased production, which was consistent with previous literature (25), likely due to the reduction in transfection efficiency. It is important to note that the two cited papers here tested higher cell densities as suspension cultures while we used adherent HEK293E. For this, we found our typical usage of adherent cells in the range of 2×10^5 cells/mL in 2 mL transfection within 6-well plates to be suitable as a baseline for further optimisations. Notably, there would be expected differences between suspension and adherent cell cultures.

With linear PEI previously reported to be a cost-effective (20), scalable (40), and efficient (41, 42) transfection reagent, we managed to obtain a peak light chain titre using up to a PEI concentration of 4 $\mu\text{g/mL}$ with diminishing returns at higher concentrations. This result supports previous findings that increasing PEI to a certain peak improves protein expression but declines thereafter (25, 43). While increasing PEI can improve transfection efficiency, higher levels can be toxic, resulting in cell death, poorer transfection efficiencies

and decreased recombinant protein production (12, 25). Although Dunnett's test did not show a significant increase in light chain production between our control of 2 µg/mL and 4 µg/mL (**Figure 1B**), the t-test showed a significant increase of 45 %, pointing 4 µg/mL as optimal. Interestingly, DNA concentration as low as 0.25 µg/mL augmented light chain production (**Figure 1C**) and was thus incorporated into the optimised parameters. High amounts of DNA were likely to decrease production due to cellular toxicity associated with higher phosphoric acid levels (12).

Next, we optimised PEI-DNA complexation incubation conditions finding good agreement with other studies (12, 24, 44) that longer complexation incubation decreased protein production (**Figure 1D**). This is likely explained by the increase in PEI-DNA complex size with time (45, 46), reducing cell penetration and lowering transfection efficiency. From these findings, we reduced our complexation incubation time to five minutes. With temperature affecting PEI and DNA complexation speed, we also investigated the incubation temperatures, finding 18 °C to be optimal. Higher temperatures reduced light chain production (**Figure 1E**) possibly due to larger complex sizes being formed that decreased entry into cells (12). While these effects may not be the same for co-transfections of more than one type of plasmid, smaller complexes are likely to improve transfection efficiency for single plasmid transfections.

Further examining PEI-DNA complexation volume where complexation typically occurs in a volume of serum free medium (47) at 10 % of the total medium volume (20). Our results (**Figure 1F**) were in agreement with a previous study finding no significant differences between 5 and 10 % (12), and even at higher serum free medium percentages up to 25 %. From this, we retained the recommended 10 %. Since the serum free medium percentage had effects on light chain production, there were hints that reducing complexation

volume could increase the proximity of PEI and DNA by enhancing its complexation to form larger complexes that could lead to poorer transfection efficiency (12).

As PEI-DNA complex is critical in the transfection process, we explored certain lab-culture practices with regards to complexation under light to explore if nucleic acid photodegradation (35) may affect transfection efficiency. Given that our results (**Figure 1 G**) show no significant difference in the light chain production, we ruled out photodegradation as a possible operational concern.

Turning to optimise culture conditions, we studied the effects of medium replacements, serum deprivation, culture in the presence of G418 antibiotic, culture incubation temperature, culture medium volume, post-transfection harvest time and supplements. Interestingly, we did not find medium replacements to cause any significant effects by one-way ANOVA, even though the t-test showed a significant increase in light chain production with fresh medium replacement prior to transfection (**Figure 2A**). This could be due to the transfection efficiency improving with waste product removal (7, 19). While such a practice is reasonable in laboratories, such a need could be costly and laborious (7) for larger-scale conditions. Interestingly, we found that medium replacements after transfection, conventionally performed to reduce transfection reagent toxicity unnecessary, for it reduced light chain production (**Figure 2A**).

With serum deprivation reported to improve transfection efficiency in some cell lines for purposes other than recombinant antibody production (14), we examined if serum deprivation could improve transfection efficiency and enhance light chain production. We found a non-significant decline in light chain production (**Figure 2B**) for the transfected HEK293E cells in agreement with a previous study (37). This could be due to the limited availability of specific essential amino acids from serum deprivation resulting in reduced antibody production (28).

Since G418 antibiotic is a requirement for HEK293E cells maintenance, we investigated if it could potentially increase metabolic stress and affect recombinant protein production after transient transfection (15). Since our results (**Figure 2C**) showed no significant difference in the light chain production with its addition, large-scale protein production can be performed with or without G418.

Based on the assumption that growth rate retardation would retain higher cell viability, lower nutrient consumption rate and reduce waste accumulation, amongst better mRNA stability, improved protein folding, and fidelity of post-translational events (48), cell culture incubation temperature was investigated. Notably, the literature in this area was inconclusive. Some were reportedly better (49), having no effect (50) or decreasing production (51) at lower temperatures such as 32 °C, with only one study on HEK293S suspension cells finding better production of non-antibody recombinant proteins at lower temperatures (13). Controlling for different evaporation rates by performing normalisations using total protein data from 2 mL transfections, we found lower temperatures detrimental to light chain production (**Figure 2D**). An extended experiment using larger 40 mL T175 transfections with a panel of Pertuzumab and Trastuzumab whole antibody variants showed no clear trends for production (Supplementary Figure 1A and B). Thus, we continued using the optimal growth temperature of 37 °C.

On the reasonable assumption that the culture medium would provide the necessary nutrients for protein production, we investigated if increasing the culture medium would overcome nutrition bottlenecks and improve light chain production. Since different culture volumes were used, total protein production data from experimental conditions were normalised to the control. Dunnett's test showed significantly lower light chain production with the usage of 1 mL cultures, but there were no significant increases despite a 40 % increase with the use of 5 mL cultures (**Figure 2F**). Apart from the significantly lower light

chain production, culture volumes below 2 mL are generally not recommended as evaporation becomes an issue. On the contrary, volume increases in 6-well plates with possible spillage and culture medium cost are not practical in larger-scale productions.

Following this, we explored post-transfection harvest, finding a direct relationship between post-transfection harvest time and light chain production which plateaued on day 14. Depending on the downstream application and the quantity of antibody desired, our results suggest that culture supernatant can be reasonably harvested anytime between day 3 to 21 post-transfection (**Figure 2F**), preferably between days 5 to 14 (52, 53). It should also be noted that longer culture times could lead to the accumulation of host-cell proteins, possibly confounding subsequent purification processes. This consideration may also be useful for proteins with poor stability where prolonged culture conditions can cause degradation.

Exploring the addition of supplements, peptone (25, 36–38), tryptone (38), casamino acids (54), yeast extract (55), casein (56), and skim milk (57) were tested. Given that the high amino acid content in the supplements could lead to toxic accumulation of ammonia culminating in cellular toxicity (36), we tested them at 1 mg/mL. Peptone was found to significantly boost light chain production in agreement with previous reports (25, 36–38). However, light chain production was not augmented by other supplements (**Figure 3A**). Determining the optimal time of supplementation, addition of 1 mg/mL peptone on either post-transfection days 5, 6 or 7 were best in boosting production (**Figure 3B**). Repeated additions of supplements did not yield better productions (**data not shown**), possibly due to toxic accumulation of ammonia (36).

Combining the optimised parameters, we improved Pertuzumab light chain production in 2 mL transfections by 217 % to 536 µg, which was further enhanced ~2-fold to 452 % reaching 1032 µg with peptone supplementation (**Figure 4A** and Supplementary Table 3). Applied to whole Pertuzumab antibody in 2 mL transfections, a similar increase of 236 %

from 22 μ g to 49 μ g was observed, with peptone addition further significantly augmenting production to 252 % reaching 51 μ g (**Figure 4A** and Supplementary Table 3). Hypothesising that dual plasmid transfection might require a longer PEI-DNA complexation time, we attempted to optimise it for whole antibody production but did not observe significant differences (**Figure 4B**). From a possible lead in our previous work (28) where 3.5 mg/mL EA supplementation improved whole antibody production, we explored both single and dual supplementations of EA and peptone. We found an unexpected significant whole antibody production decrease upon EA addition which was rescued by the addition of peptone (**Figure 4C**) for the Pertuzumab antibody.

Considering that some whole antibody productions are limited by the light chain (i.e. Pertuzumab V κ 5) from our previous work (29) and that peptone appeared to benefit light chain production in this study, we performed similar experiments using Pertuzumab V κ 5|VH3 whole antibodies for validation. We were able to significantly boost the whole V κ 5|VH3 antibodies production (**Figure 4C**) by 28 % from 2.4 to 3.0 μ g in 2 mL transfections (Supplementary Table 3) supporting the use of peptone supplements for antibodies limited by light chains in transient production. Thereby, peptone could be added for light chain limiting antibody production given its benefits and lack of detrimental effects.

CONCLUSION

Through a detailed systematic selection of antibodies in a holistic (1, 31) manner, the selection of previously published factors of heavy chain constant (30), V κ -VH pairings (29), amino acid usage with signal peptide selection (28), antibody production could be more efficient, mitigating commercial failures due to production costs. Here our optimisations of the parameters led to a ~2-fold increase of light chain only and whole antibody production from HEK293E using 25 kDa linear PEI in a cost-effective and operationally convenient

manner. With peptone supplements, light chain increased by ~4-fold while whole antibody increased to a lesser extent, with most benefits found for antibodies limited by their light chains in production. These findings have potential applications to other mammalian-based recombinant protein production.

DATA AVAILABILITY STATEMENT

The data underlying this article will be shared at reasonable request to the corresponding author.

AUTHOR CONTRIBUTIONS

Conceptualisation, W.-L.L. and S.K.-E.G.; methodology, W.-L.L. and Z.S.-L.H.; formal analysis, W.-L.L., Z.S.-L.H.; S.K.-E.G.; investigation, W.-L.L. and Z.S.-L.H.; validation, D.W.-S.K., J.Y.Y., writing—original draft preparation, Z.S.-L.H.; writing—review and editing, D.W.-S.K., J.Y.Y., S.K.-E.G., W.-L.L. and Z.S.-L.H.; visualisation, S.K.-E.G., W.-L.L. and Z.S.-L.H.; supervision, S.K.-E.G. and W.-L.L.; funding acquisition, S.K.-E.G. All authors have read and agreed to the published version of the manuscript.

FUNDING STATEMENT

This work was initially supported by the Joint Council Office, Agency for Science, Technology, and Research, Singapore, under Grant number JCO1334i00050 and later by the National Research Foundation (NRF) Singapore grant to Experimental Drug Development Centre (EDDC) for the platform AMDED, acquired by S.K.-E.G.

ACKNOWLEDGEMENTS

The authors would also like to thank Jun-Jie Poh for his T175 flasks transfection data contribution.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

ETHICS AND CONSENT STATEMENT

Consent was not required for this study

ANIMAL RESEARCH STATEMENT

This is not applicable to this study.

REFERENCES

1. Ling, W.-L., Lua, W.-H. and Gan, S.K.-E. (2020) Sagacity in antibody humanization for therapeutics, diagnostics and research purposes: considerations of antibody elements and their roles. *Antib. Ther.*, **3**, 71–79.
2. Pham, P.L., Kamen, A. and Durocher, Y. (2006) Large-Scale Transfection of Mammalian Cells for the Fast Production of Recombinant Protein. *Mol. Biotechnol.*, **34**, 225–238.
3. Geisse, S. and Henke, M. (2005) Large-scale Transient Transfection of Mammalian Cells: A Newly Emerging Attractive Option for Recombinant Protein Production. *J. Struct. Funct. Genomics*, **6**, 165–170.
4. Wurm, F.M. (2004) Production of recombinant protein therapeutics in cultivated mammalian cells. *Nat. Biotechnol.*, **22**, 1393–1398.
5. Durocher, Y. (2002) High-level and high-throughput recombinant protein production by transient transfection of suspension-growing human 293-EBNA1 cells. *Nucleic Acids Res.*, **30**, E9.
6. Yuan, J., Xu, W.W., Jiang, S., Yu, H. and Poon, H.F. (2018) The Scattered Twelve Tribes of HEK293. *Biomed. Pharmacol. J.*, **11**, 621–623.
7. Durocher, Y. (2002) High-level and high-throughput recombinant protein production by transient transfection of suspension-growing human 293-EBNA1 cells. *Nucleic Acids Res.*, **30**, 9e–99.
8. Meissner, P., Pick, H., Kulangara, A., Chatellard, P., Friedrich, K. and Wurm, F.M. (2001) Transient gene expression: Recombinant protein production with suspension-adapted HEK293-EBNA cells. *Biotechnol. Bioeng.*, **75**, 197–203.
9. Ho, Y.K. and Too, H.P. (2020) Development of a laboratory scalable process for enhancing lentivirus production by transient transfection of HEK293 adherent cultures. *Gene Ther.*, **27**, 482–494.

- 571 10. Bollin,F., Dechavanne,V. and Chevalet,L. (2011) Design of Experiment in CHO and
572 HEK transient transfection condition optimization. *Protein Expr. Purif.*, **78**, 61–68.
- 573 11. Sou,S.N., Polizzi,K.M. and Kontoravdi,C. (2013) Evaluation of transfection methods for
574 transient gene expression in Chinese hamster ovary cells. *Adv. Biosci. Biotechnol.*, **04**,
575 1013–1019.
- 576 12. Greene,E., Cazacu,D., Tamot,N., Castellano,S., Datar,A., Kronkaitis,A., Gebhard,D.,
577 Reed,J., Mawson,P., Florin,L., *et al.* (2021) Optimization of a transient antibody
578 expression platform towards high titer and efficiency. *Biotechnol. J.*, **16**, e2000251.
- 579 13. Lin,C.-Y., Huang,Z., Wen,W., Wu,A., Wang,C. and Niu,L. (2015) Enhancing Protein
580 Expression in HEK-293 Cells by Lowering Culture Temperature. *PLOS ONE*, **10**,
581 e0123562.
- 582 14. Wallenstein,E.J., Barminko,J., Schloss,R.S. and Yarmush,M.L. (2010) Serum starvation
583 improves transient transfection efficiency in differentiating embryonic stem cells.
584 *Biotechnol. Prog.*, **26**, 1714–1723.
- 585 15. Yallop,C.A. and Svendsen,I. (2001) The effects of G418 on the growth and metabolism
586 of recombinant mammalian cell lines. *Cytotechnology*, **35**, 101–114.
- 587 16. Mason,M., Sweeney,B., Cain,K., Stephens,P. and Sharfstein,S. (2014) Reduced Culture
588 Temperature Differentially Affects Expression and Biophysical Properties of
589 Monoclonal Antibody Variants. *Antibodies*, **3**, 253–271.
- 590 17. Xu,J., Tang,P., Yongky,A., Drew,B., Borys,M.C., Liu,S. and Li,Z.J. (2019) Systematic
591 development of temperature shift strategies for Chinese hamster ovary cells based on
592 short duration cultures and kinetic modeling. *mAbs*, **11**, 191–204.
- 593 18. Baldi,L., Hacker,D.L., Adam,M. and Wurm,F.M. (2007) Recombinant protein production
594 by large-scale transient gene expression in mammalian cells: state of the art and future
595 perspectives. *Biotechnol. Lett.*, **29**, 677–684.
- 596 19. Fang,X.T., Sehlin,D., Lannfelt,L., Syvänen,S. and Hultqvist,G. (2017) Efficient and
597 inexpensive transient expression of multispecific multivalent antibodies in Expi293
598 cells. *Biol. Proced. Online*, **19**, 11.
- 599 20. Schlaeger,E.-J. and Christensen,K. (1999) Transient gene expression in mammalian cells
600 grown in serum-free suspension culture. *Cytotechnology*, **30**, 71–83.
- 601 21. Davami,F., Baldi,L., Rajendra,Y. and Wurm,F.M. (2014) Peptone Supplementation of
602 Culture Medium Has Variable Effects on the Productivity of CHO Cells. *Int. J. Mol.*
603 *Cell Med*, **3**, 146–156.
- 604 22. Pérez-Rodriguez,S., Ramírez-Lira,M. de J., Trujillo-Roldán,M.A. and Valdez-Cruz,N.A.
605 (2020) Nutrient supplementation strategy improves cell concentration and longevity,
606 monoclonal antibody production and lactate metabolism of Chinese hamster ovary
607 cells. *Bioengineered*, **11**, 463–471.
- 608 23. You,M., Liu,Y., Chen,Y., Guo,J., Wu,J., Fu,Y., Shen,R., Qi,R., Luo,W. and Xia,N. (2013)
609 Maximizing Antibody Production in Suspension-Cultured Mammalian Cells by the

610 Customized Transient Gene Expression Method. *Biosci. Biotechnol. Biochem.*, **77**,
611 1207–1213.

612 24. Liu,C., Dalby,B., Chen,W., Kilzer,J.M. and Chiou,H.C. (2008) Transient Transfection
613 Factors for High-Level Recombinant Protein Production in Suspension Cultured
614 Mammalian Cells. *Mol. Biotechnol.*, **39**, 141–153.

615 25. Davami,F., Eghbalpour,F., Barkhordari,F. and Mahboudi,F. (2014) Effect of Peptone
616 Feeding on Transient Gene Expression Process in CHO DG44. *Avicenna J. Med.*
617 *Biotechnol.*, **6**, 147–155.

618 26. Mains,P.E. and Sibley,C.H. (1983) The requirement of light chain for the surface
619 deposition of the heavy chain of immunoglobulin. *J. Biol. Chem.*, **258**, 5027–5033.

620 27. Schlatter,S., Stansfield,S.H., Dinnis,D.M., Racher,A.J., Birch,J.R. and James,D.C. (2005)
621 On the Optimal Ratio of Heavy to Light Chain Genes for Efficient Recombinant
622 Antibody Production by CHO Cells. *Biotechnol. Prog.*, **21**, 122–133.

623 28. Ling,W.-L., Su,C.T.-T., Lua,W.-H., Poh,J.-J., Ng,Y.-L., Wipat,A. and Gan,S.K.-E. (2020)
624 Essentially Leading Antibody Production: An Investigation of Amino Acids,
625 Myeloma, and Natural V-Region Signal Peptides in Producing Pertuzumab and
626 Trastuzumab Variants. *Front. Immunol.*, **11**, 604318.

627 29. Ling,W.-L., Lua,W.-H., Poh,J.-J., Yeo,J.Y., Lane,D.P. and Gan,S.K.-E. (2018) Effect of
628 VH–VL Families in Pertuzumab and Trastuzumab Recombinant Production, Her2 and
629 FcγIIA Binding. *Front. Immunol.*, **9**, 469.

630 30. Lua,W.-H., Ling,W.-L., Yeo,J.Y., Poh,J.-J., Lane,D.P. and Gan,S.K.-E. (2018) The
631 effects of Antibody Engineering CH and CL in Trastuzumab and Pertuzumab
632 recombinant models: Impact on antibody production and antigen-binding. *Sci. Rep.*, **8**,
633 718.

634 31. Phua,S.-X., Chan,K.-F., Su,C.T.-T., Poh,J.-J. and Gan,S.K.-E. (2019) Perspective: The
635 promises of a holistic view of proteins—impact on antibody engineering and drug
636 discovery. *Biosci. Rep.*, **39**, BSR20181958.

637 32. Chan,W.-T., Verma,C.S., Lane,D.P. and Gan,S.K.-E. (2013) A comparison and
638 optimization of methods and factors affecting the transformation of *Escherichia coli*.
639 *Biosci Rep.*, **33**, e00086.

640 33. Poh,J.-J. and Gan,S.K.-E. (2014) The Determination of Factors involved in Column-
641 Based Nucleic Acid Extraction and Purification. *J. Bioprocess. Biotech.*, **4**, 157.

642 34. Su,C.T.-T., Ling,W.-L., Lua,W.-H., Poh,J.-J. and Gan,S.K.-E. (2017) The role of
643 Antibody Vκ Framework 3 region towards Antigen binding: Effects on recombinant
644 production and Protein L binding. *Sci. Rep.*, **7**, 3766.

645 35. Korycka-Dahl,M. and Richardson,T. (1980) Photodegradation of DNA with fluorescent
646 light in the presence of riboflavin, and photoprotection by flavin triplet-state
647 quenchers. *Biochim. Biophys. Acta BBA - Nucleic Acids Protein Synth.*, **610**, 229–234.

- 648 36. Heidemann,R., Zhang,C., Qi,H., Larrick Rule,J., Rozales,C., Park,S., Chuppa,S., Ray,M.,
649 Michaels,J., Konstantinov,K., *et al.* (2000) The use of peptones as medium additives
650 for the production of a recombinant therapeutic protein in high density perfusion
651 cultures of mammalian cells. *Cytotechnology*, **32**, 157–167.
- 652 37. Pham,P.L., Perret,S., Doan,H.C., Cass,B., St-Laurent,G., Kamen,A. and Durocher,Y.
653 (2003) Large-Scale Transient Transfection of Serum-Free Suspension-Growing
654 HEK293 EBNA1 Cells: Peptone Additives Improve Cell Growth and Transfection
655 Efficiency. *Biotechnol. Bioprocess Eng.*, **84**.
- 656 38. Pham,P.L., Perret,S., Cass,B., Carpentier,E., St-Laurent,G., Bisson,L., Kamen,A. and
657 Durocher,Y. (2005) Transient gene expression in HEK293 cells: Peptone addition
658 posttransfection improves recombinant protein synthesis. *Biotechnol. Bioeng.*, **90**,
659 332–344.
- 660 39. Backliwal,G., Hildinger,M., Hasija,V. and Wurm,F.M. (2008) High-density transfection
661 with HEK-293 cells allows doubling of transient titers and removes need for a priori
662 DNA complex formation with PEI. *Biotechnol. Bioeng.*, **99**, 721–727.
- 663 40. Nyamay’Antu,A., Kedinger,V. and Erbacher,P. (2018) Addressing scaling-up limitations:
664 optimized PEI-mediated production of clinical grade viral vectors. *Cell Gene Ther.*
665 *Insights*, **4**, 71–79.
- 666 41. Wiseman,J.W., Goddard,C.A., McLelland,D. and Colledge,W.H. (2003) A comparison of
667 linear and branched polyethylenimine (PEI) with DCChol/DOPE liposomes for gene
668 delivery to epithelial cells in vitro and in vivo. *Gene Ther.*, **10**, 1654–1662.
- 669 42. Hanzlíková,M., Ruponen,M., Galli,E., Raasmaja,A., Aseyev,V., Tenhu,H., Urtti,A. and
670 Yliperttula,M. (2011) Mechanisms of polyethylenimine-mediated DNA delivery: free
671 carrier helps to overcome the barrier of cell-surface glycosaminoglycans: Role of free
672 PEI in transfection. *J. Gene Med.*, **13**, 402–409.
- 673 43. Xie,Q., Xinyong,G., Xianjin,C. and Yayu,W. (2013) PEI/DNA formation affects transient
674 gene expression in suspension Chinese hamster ovary cells via a one-step transfection
675 process. *Cytotechnology*, **65**, 263–271.
- 676 44. Bos,A.B., Duque,J.N., Bhakta,S., Farahi,F., Chirdon,L.A., Junutula,J.R., Harms,P.D. and
677 Wong,A.W. (2014) Development of a semi-automated high throughput transient
678 transfection system. *J. Biotechnol.*, **180**, 10–16.
- 679 45. Han,X., Fang,Q., Yao,F., Wang,X., Wang,J., Yang,S. and Shen,B.Q. (2009) The
680 heterogeneous nature of polyethylenimine-DNA complex formation affects transient
681 gene expression. *Cytotechnology*, **60**, 63–75.
- 682 46. Sang,Y., Xie,K., Mu,Y., Lei,Y., Zhang,B., Xiong,S., Chen,Y. and Qi,N. (2015) Salt ions
683 and related parameters affect PEI–DNA particle size and transfection efficiency in
684 Chinese hamster ovary cells. *Cytotechnology*, **67**, 67–74.
- 685 47. Yang,J.-P. and Huang,L. (1997) Overcoming the inhibitory effect of serum on lipofection
686 by increasing the charge ratio of cationic liposome to DNA. *Gene Ther.*, **4**, 950–960.

687 48. Masterton,R.J., Roobol,A., Al-Fageeh,M.B., Carden,M.J. and Smales,C.M. (2010) Post-
688 translational events of a model reporter protein proceed with higher fidelity and
689 accuracy upon mild hypothermic culturing of Chinese hamster ovary cells. *Biotechnol.*
690 *Bioeng.*, **105**, 215–220.

691 49. Yoon,S.K., Hwang,S.O. and Lee,G.M. (2004) Enhancing Effect of Low Culture
692 Temperature on Specific Antibody Productivity of Recombinant Chinese Hamster
693 Ovary Cells: Clonal Variation. *Biotechnol. Prog.*, **20**, 1683–1688.

694 50. Bloemkolk,J.-W., Gray,M.R., Merchant,F. and Mosmann,T.R. (1992) Effect of
695 temperature on hybridoma cell cycle and MAb production. *Biotechnol. Bioeng.*, **40**,
696 427–431.

697 51. Barnabé,N. and Butler,M. (1994) Effect of temperature on nucleotide pools and
698 monoclonal antibody production in a mouse hybridoma. *Biotechnol. Bioeng.*, **44**,
699 1235–1245.

700 52. Shatz,W., Ng,D., Dutina,G., Wong,A.W., Dunshee,D.R., Sonoda,J., Shen,A. and
701 Scheer,J.M. (2016) An efficient route to bispecific antibody production using single-
702 reactor mammalian co-culture. *mAbs*, **8**, 1487–1497.

703 53. Li,F., Vijayasankaran,N., Shen,A. (Yijuan), Kiss,R. and Amanullah,A. (2010) Cell
704 culture processes for monoclonal antibody production. *mAbs*, **2**, 466–479.

705 54. Kaushik,N., Rohila,D., Arora,U., Raut,R., Lamminmäki,U., Khanna,N. and Batra,G.
706 (2016) Casamino acids facilitate the secretion of recombinant dengue virus serotype-3
707 envelope domain III in *Pichia pastoris*. *BMC Biotechnol.*, **16**, 12.

708 55. Li,M., Liao,X., Zhang,D., Du,G. and Chen,J. (2011) Yeast Extract Promotes Cell Growth
709 and Induces Production of Polyvinyl Alcohol-Degrading Enzymes. *Enzyme Res.*,
710 **2011**, 1–8.

711 56. Russell,J.B., Sniffen,C.J. and Van Soest,P.J. (1983) Effect of Carbohydrate Limitation on
712 Degradation and Utilization of Casein by Mixed Rumen Bacteria. *J. Dairy Sci.*, **66**,
713 763–775.

714 57. Khani,M.-H. and Bagheri,M. (2020) Skimmed milk as an alternative for IPTG in
715 induction of recombinant protein expression. *Protein Expr. Purif.*, **170**, 105593.

716

717 **FIGURES**

Figure 1: Pertuzumab light chain productions in 2 mL transfections from optimising the transfection parameters of (A) Seeding density; (B) PEI concentration; (C) DNA concentration; (D) PEI-DNA complexation time (E); PEI-DNA complexation temperature; (F) PEI-DNA complexation volume; (G) PEI-DNA complexation in light and dark environments. Black coloured bars represent the controls to which the experimental conditions were compared for statistical analysis. *, **, *** denote significance at $p < 0.05$, $p < 0.01$ and $p < 0.001$, respectively from one-way ANOVA Dunnett's post-hoc test, while ⁺ denotes significance at $p < 0.05$ using t-test analysis. (The experiments were carried out in at least three independent replicates with titres and standard errors provided in Supplementary Table 1.)

Figure 2: Pertuzumab light chain production in 2 mL transfections from optimising the culture parameters with respect to A) Medium replacement timing; (B) Serum deprivation; (C) Culturing with G418 antibiotic; (D) Culture temperature; (E) Culture medium volume (F) Post-transfection harvest day. Black coloured bars in A-D and black coloured dots in E and F represent the controls to which the respective experimental conditions were compared against for statistical analysis. *, **, *** denote significance at $p < 0.05$, $p < 0.01$ and $p < 0.001$, respectively from one-way ANOVA Dunnett's post-hoc test. ⁺ denotes significance at $p < 0.05$ using t-test analysis. (The experiments were carried out in at least three independent

replicates with light chain titres for A-D found in Supplementary Table 1 while the total protein for E and F can be found in Supplementary Table 2.)

Figure 3: Antibody light chain production in 2 mL transfections with (A) nutrient supplements and (B) Heat map of one-time additions of different peptone concentrations. Black coloured bars in A represent the control to which the respective supplements were compared for statistical analysis. ⁺, denotes significance at $p < 0.05$ using t-test analysis. (The experiments were carried out in at least three independent replicates where the light chain total protein values are provided in Supplementary Table 2).

Figure 4: Production of recombinant Pertuzumab in 2mL transfections of (A) Light chain (Vκ1) and whole Vκ1|VH3 antibody among standard, optimised and with peptone supplementation conditions; (B) Whole Vκ1|VH3 Pertuzumab production at optimised parameters with PEI-DNA complex time variations; (C) Vκ1|VH3 and Vκ5|VH3 variants of recombinant Pertuzumab among optimised parameters with peptone and/or essential amino acids supplements. Black coloured bars in A and C represent the control to which the respective experiment parameters were compared by one-way ANOVA Dunnett's post-hoc test. *, **, *** denote significance at $p < 0.05$, $p < 0.01$ and $p < 0.001$, respectively. (The experiments were carried out in at least three independent replicates where the total protein values are provided in Supplementary Table 3).

TABLE

Table 1: Control and optimised parameters for transient recombinant production in 2 mL, 6-well plate cultures.

Parameters	Control	Optimal
Cell Density (cells/mL)	2×10^5	2×10^5
PEI Concentration (µg/mL)	2	4
Plasmid DNA Concentration (µg/mL)	1	0.25
Complexation Time (min)	30	5
Complexation Temperature (°C)	18	18
Complexation in SFM (% of transfection volume)	10	10
Culture Medium Volume (mL)	2	2
Post-transfection Harvest (day)	14	14
Culture Incubation Temperature (°C)	37	37
Supplements (Post-transfection)	-	1 mg/mL Peptone on Day 7