1 Identification and molecular characterization of novel viruses in Ugandan cattle.

2 **Authors:** Stephen Balinandi^{1,2}, Juliette Hayer³, Harindranath Cholleti⁴, Michelle Wille^{5,6}, Julius
3 J. Lutwama¹, Maja Malmberg^{3,4}, Lawrence Mugisha^{2,7#}.

4 Affiliations:

5 ¹Uganda Virus Research Institute; Entebbe, Uganda.

6 ²College of Veterinary Medicine, Animal Resources and Biosecurity, Makerere University,
7 Kampala, Uganda.

8 ³Department of Animal Breeding and Genetics, Swedish University of Agricultural Sciences,
9 Uppsala, Sweden.

10 ⁴Department of Biomedical Sciences and Veterinary Public Health, Swedish University of
11 Agricultural Sciences, Uppsala, Sweden.

12 ⁵Sydney Institute for Infectious Diseases, School of Life and Environmental Sciences and School
13 of Medical Sciences, The University of Sydney, Sydney, NSW 2006, Australia.

14 ⁶Department of Microbiology and Immunology, at the Peter Doherty Institute for Infection and
15 Immunity, The University of Melbourne, Melbourne, Victoria, Australia.

16 ⁷Ecohealth Research Group, Conservation & Ecosystem Health Alliance (CEHA), Kampala,
17 Uganda.

18 **#Corresponding Author:** Lawrence Mugisha BVM, MSc, PhD; ORCID: [https://orcid.org/0000-](https://orcid.org/0000-0003-4932-3356)
19 [0003-4932-3356](https://orcid.org/0003-4932-3356). College of Veterinary Medicine, Animal Resources and Biosecurity, Makerere

university, P.O. Box 34153, Kampala, Uganda. Email: mugishalaw@gmail.com;
lawrence.mugisha@mak.ac.ug

Abstract

The risk for the emergence of novel viral zoonotic diseases in animals and humans in Uganda is high given its geographical location with high biodiversity. We aimed to identify and characterize viruses in 175 blood samples from cattle selected in Uganda using molecular approaches. We identified 8 viral species belonging to 4 families (*Flaviviridae*, *Peribunyaviridae*, *Reoviridae* and *Rhabdoviridae*) and 6 genera (*Hepacivirus*, *Pestivirus*, *Orthobunyavirus*, *Coltivirus*, *Dinovernavirus* and *Ephemerovirus*). Four viruses were novel and tentatively named as Zikole virus (Family: *Flaviviridae*), Zeboroti virus (Family: *Reoviridae*), Zebtine virus (Family: *Rhabdoviridae*) and Kokolu virus (Family: *Rhabdoviridae*). In addition, Bovine hepatitis virus, Obodhiang virus, *Aedes pseudoscutellaris* reovirus and Schmallenberg virus were identified for the first time in Ugandan cattle. We report a broad range of viruses including novel ones in the blood of cattle likely as reservoir hosts for emergence of novel viruses with serious public health implications.

Keywords: cattle, zoonoses, pan-PCR, Metagenomics, RNA virus, Uganda

Highlights

- Pan PCR and High Throughput Sequencing Approaches reveal novel viruses in the blood of cattle.
- Identified 8 viral species belonging to 4 families: *Flaviviridae*, *Peribunyaviridae*, *Reoviridae* and *Rhabdoviridae*

- 42 • 60% of the identified viruses were found in the Ankole cattle breed

43

Introduction

Since the mid-1900s, infectious diseases of livestock have been identified, not only as a major threat to animal health, welfare and production, but also to the overall global health security especially in developing countries (Cleaveland et al., 2017; Grace D et al., 2012). Moreover, more than 60% of documented emerging, and re-emerging human diseases are zoonotic in nature. That is, viruses that are endemic within animal populations spill over into humans through vectors and other transmission pathways and cause epidemics (Morse, 2004), or global pandemics as with SARS-CoV-2 (R. Lu et al., 2020)

Uganda is a known hotspot for the emergence of novel pathogens due to its conducive climatic conditions and high biodiversity that include a variety of wild and domestic animals, birds and insects acting as disease vectors, hosts or reservoirs, and the attendant anthropogenic factors such as deforestation (Allen et al., 2017; Jones et al., 2008). Indeed, in just over 10 years, several new high consequence viral pathogens have been identified, including *Bundibugyo ebolavirus* (Family: *Filoviridae*) (Towner et al., 2008), *Sosuga pararubulavirus* (Family: *Paramyxoviridae*) (Albariño et al., 2014), Bukakata orbivirus (*Chobar George virus*; Family: *Reoviridae*) (Fagre et al., 2019) and Ntwetwe virus (Family: *Peribunyaviridae*) (Edridge et al., 2019). Other viruses of public health importance that were previously discovered in Uganda, especially during the middle of the 20th century, include *Zika virus*, *West Nile virus* and *O'nyong-nyong virus* (Dick, Kitchen, & Haddow, 1952; Smithburn, Hughes, Burke, & Paul, 1940; Williams, Woodall, & Gillett, 1965).

Despite enhanced efforts to identify and manage potential disease-causing pathogens in Uganda through improved surveillance, increased public and health sector awareness, and expedited transportation of suspected samples to the testing laboratories (Borchert et al., 2014;

Kemunto et al., 2018; Kiyaga et al., 2013; Shoemaker et al., 2018), the disease burden associated with undifferentiated illnesses of unknown origin remains high (Lamorde et al., 2018). As such, it is critical to implement broad investigations of potential disease-causing agents to include animal hosts and vectors to identify the potential sources of viral pathogens likely to cause diseases in both animals and humans. These investigations would align well with the WHO's Priority 'Disease X', which represents a currently unknown serious global health threat that can emerge in humans (Mehand, Al-Shorbaji, Millett, & Murgue, 2018). In our current study, we aimed at identifying and characterizing viruses from blood samples collected in cattle from selected districts of Uganda. Previous studies have alluded that bovines are a major source of zoonoses (McDaniel, Cardwell, Moeller, & Gray, 2014). More specifically, cattle have been implicated as hosts for 2 human viruses. Firstly, the dated emergence of Human coronavirus OC-43 in 1890 coincides with a pandemic of human respiratory disease in humans between 1889 – 1890, and it is believed that this was after the virus emerged following a cross species transmission event of Bovine coronavirus (Crookshank, 1897; Vijgen et al., 2005). Secondly, measles virus is said to have crossed to humans after it diverged from rinderpest virus, a historical bovine virus, around the 11th to 12th Century (Furuse, Suzuki, & Oshitani, 2010). Therefore, with the high consumption of meat and other dairy products, as the main source of food in Uganda (UBOS, 2017), it is prudent to understand the viral diversity in this animal reservoir to expand on our pandemic preparedness toolkit.

Materials and Methods

Ethics Statement

The authors confirm that the ethical policies of this journal, as noted on the author guidelines page, have been adhered to: sample collection was approved by the Institutional Animal Care and Use Committee (IACUC), School of Veterinary Medicine & Animal Resources (SVAR), Makerere University (Reference Number: SVARREC/03/2017) and the Uganda National Council for Science and Technology (UNCST) (Reference Number: UNCST A580). In addition, written consent was obtained from all animal owners, or their representative(s), following a detailed explanation of the study objectives.

Study Sites

Uganda is an East African country located along the Equator, lying between 4°N and 1°S latitudes, and between 30°E and 35°E longitudes. Thus, many areas within the country have a typical tropical climate; experiencing relatively humid conditions (average of 70%), rainfall (700-1,500 mm) and daily temperatures that are between 20-30°C throughout the year (UBOS, 2017). Its natural vegetation is varied although it is mainly open wooded savannah grassland in most of the country, thus making livestock keeping a major agro-economic activity for the country. Recent data from the Ministry of Agriculture, Animal Industry and Fisheries (MAAIF) and Uganda Bureau of Statistics (UBOS), show that Uganda has over 14 million cattle, 16 million goats, 5 million sheep and 4 million pigs with every household owning at least one type of livestock (UBOS, 2017).

This study was conducted using blood samples from cattle that were sampled from 5 purposively selected districts of Uganda: Kasese (0°N, 30°E), Hoima (1°N, 31°E), Gulu (3°N, 32°E), Soroti (1°N, 33°E) and Moroto (2°N, 34°E) (Figure 1 and Table 2). These districts were

selected as they represent areas with distinct agroecological variations (Drichi & National Biomass, 2003; Wortmann & Eledu, 1999). Briefly, Kasese and Moroto districts lie at the extreme ends of the western and eastern borderlines, respectively, of the Ugandan Cattle Corridor – a large broad zone of semi-arid rangelands, where cattle-rearing is a preoccupation of the local communities. Gulu and Soroti districts lie within the swampy lowlands of the Lake Kyoga basin, while Hoima district is a medium altitude area with mixed agricultural practices.

Figure 1: Map of Uganda showing the sampling sites in this study (Source: This map was created using open-source data in ArcGIS software, v10.2, Environmental Systems Research Institute, Inc., Redlands, CA, USA). Inset is a map of Africa with the enlarged region boxed in red.

Study Samples

In this study, we used samples collected as part of a multidisciplinary baseline investigation on tickborne disease in cattle in Uganda (Balinandi et al., 2021; Malmberg & Hayer, 2019). A subset of 175 cattle samples was randomly selected from a total of 500 sampled animals, comprising 35 samples from each of the 5 study districts (Gulu, Soroti, Moroto, Hoima, and Kasese). In the field, information about each of the sampled animal such as their age, sex, breed, health status, tick burden, grazing system, etc., was collected at the time of sample collection using a standardized biodata form. In addition, about 4ml of whole blood was drawn from the jugular or tail veins of restrained cattle into EDTA tubes (BD Vacutainer®, Franklin Lakes, NJ, USA). Blood was then immediately separated into small aliquots of whole blood and serum, which were immediately frozen in liquid Nitrogen before they were transported to Uganda Virus Research Institute (UVRI), Entebbe, Uganda, and stored at -80°C until RNA extraction and family-wide conventional pan-PCR was undertaken.

pan-PCR

Total nucleic acids were extracted from whole blood samples using the MagMax™ 96 viral Isolation kit (Applied Biosystems, Vilnius, Lithuania), according to the manufacturer's instructions. Extracted RNA was transcribed into cDNA using Superscript III First-strand Synthesis Supermix (Invitrogen, CA, USA). We assayed the samples for 5 viral taxa: *Filoviridae*,

138 *Flaviviridae*, *Alphavirus*, *Rhabdoviridae*, and *Orthobunyavirales*, using protocols previously
139 developed and applied for the USAID/PREDICT II program (Moureau et al., 2007; PREDICT,
140 2014; Sanchez-Seco, Rosario, Quiroz, Guzman, & Tenorio, 2001; Zhai et al., 2007) (Table 1).
141 These viral taxa also represent the main groups of viruses from which most members of
142 arboviruses are frequently identified (Hubálek, Rudolf, & Nowotny, 2014; Woolhouse &
143 Gowtage-Sequeria, 2005).

144 Table 1. Pan-PCR Primer Sequences and Protocols used for Virus Detection in Cattle, Uganda, 2017.

Virus Taxon	Target	1 st Round PCR Primer Sequences ¹	2 nd Round PCR Primer Sequences ¹	Reference
Family: <i>Filoviridae</i>	L gene	F: TITTYTCHVTICAAAAICAYTGGG R: ACCATCATRTRCTIGGRAAKGCTTT	F: TITTYTCHVTICAAAAICAYTGGG R: GCYTCISMIAIIGTTTGIACATT	Modified (Zhai et al., 2007)
Family: <i>Flaviviridae</i>	NS5 gene	F: TGYRBTAYAACATGATGGG R: GTGTCCCAICCNCGNTRTC	- -	(Moureau et al., 2007)
Family: <i>Rhabdoviridae</i>	L gene	F: CCADMCBTTTGYCKYARRCCTTC R: AAGGYAGRTTTTTYKCDYTRATG	F: CCADMCBTTTGYCKYARRCCTTC R: AARTGGAAYAAAYCAYCARMG	Unpublished ²
Genus: <i>Alphavirus</i>	NSP4 gene	F: GAYGCITAYYTIGAYATGGTIGAIGG R: KYTCYTCIGTRTGYYTIGTICIGG	F: GIAAYTGAAAYGTIACICARATG R: GCRAAIARIGCIGCIGCYTYIGGICC	(Sanchez-Seco et al., 2001)
Order: <i>Orthobunyavirales</i>	L gene	F: GARATHAATGCWGAYATGTCHAAAT R: TAIGTYTTYTTTCATRTTDGCTG	F: AAATGGAGTGCHCARGATGT R: TAIGTYTTYTTTCATRTTDGCTG	Unpublished ²

145 ¹Sequences are in 5' to 3' direction

146 ²Assay was developed and validated at the University of California, Davis, USA.

147 *High-throughput sequencing*

148 Ten serum samples (Table 2), corresponding to the whole blood samples that were positive
149 for any viral family, including those that showed non-specific bands on gel electrophoresis, were
150 preserved in TRIzol™ Reagent (Invitrogen, Thermo Scientific) and shipped on dry ice to the
151 Swedish University of Agricultural Sciences (SLU), Uppsala, Sweden, for high-throughput
152 sequencing.

153 RNA was extracted using the GeneJet RNA purification kit (Thermo Scientific™), as per
154 the manufacturer's instructions. The eluted serum RNA was further concentrated by RNeasy
155 MinElute Cleanup Kit (Qiagen) before the RNA was randomly amplified with the Ovation RNA-
156 Seq System v2 (Tecan) following the manufacturer's recommended protocols. The concentration
157 and size distribution of generated double-stranded cDNA products were estimated by the
158 TapeStation 4200 system (Agilent Technologies). Sequencing libraries were prepared from 50ng
159 of DNA according to the manufacturer's preparation guide #112219, using the SMARTer
160 ThruPLEX DNA-Seq kit (R400676, Takara-Clontech). Briefly, DNA was fragmented using a
161 Covairs E220 system, aiming at 350-400bp fragments. The ends of the fragments were end-
162 repaired, and stem-loop adapters were ligated to the 5' ends of the fragments. The 3' end of the
163 stem-loop was subsequently extended to close the nick. Finally, the fragments were amplified, and
164 unique index sequences were introduced using 8 cycles of PCR, followed by purification using
165 AMPure XP beads (Beckman Coulter). The quality of the library was evaluated using the Agilent
166 2200 TapeStation system and a D1000 ScreenTape-kit. The adapter-ligated fragments were
167 quantified by qPCR using the Library quantification kit for Illumina (KAPA Biosystems/Roche)
168 on a CFX384Touch instrument (BioRad), prior to cluster generation and sequencing.

Thereafter, a 300 pM pool of the sequencing libraries in equimolar ratio was subjected to cluster generation and paired-end sequencing, with 150bp read length in an SP flow cell and the NovaSeq6000 system (Illumina Inc.) using the v1 chemistry, according to the manufacturer's protocols. Base-calling was done on the instrument using RTA 3.3.4 and the resulting .bcl files were demultiplexed and converted to FASTQ format with tools provided by Illumina Inc., allowing for one mismatch in the index sequence. Additional statistics on sequence quality were compiled with an in-house script from the Fastq-files, RTA and CASAVA output files. Sequencing was performed by the SNP&SEQ Technology Platform, part of the SciLifeLab, in Uppsala, Sweden. Sequence libraries have been deposited in SRA under the study PRJEB46076 with Accession Numbers: ERR6496566, ERR6496567, ERR6496875, ERR6496876, ERR6496901, ERR6496964, ERR6497048, ERR6497050, ERR6497350 and ERR6497480.

Bioinformatics

Quality control and trimming of the raw sequencing reads was performed using FASTP (Lipman & Pearson, 1985). The good quality reads were then aligned to the cattle genome (Bos_taurus.ARS-UCD1.2.; www.ncbi.nlm.nih.gov/assembly/GCF_002263795.1/) using Bowtie2 (Langmead & Salzberg, 2012) to eliminate reads from the host. The remaining reads were *de novo* assembled using MEGAHIT (Li et al., 2016). The obtained assembled contigs were taxonomically classified using two methods: KRAKEN 2 (Wood, Lu, & Langmead, 2019) and DIAMOND (Buchfink, Xie, & Huson, 2015), both using the NCBI non-redundant protein database *nr* (release of April 2019) (Deng et al., 2006). This analysis was also performed at reads level, on the reads remaining unmapped after the alignment to the cattle genome.

The longest assembled contigs (length>1000 nt) that were classified as viral were retrieved for further analyses. BLASTN and BLASTX (Altschul et al., 1997; Chen, Ye, Zhang, & Xu, 2015) were performed to confirm the closest homologues and to check the alignments with viral sequences from the database. For some identified viruses for which only partial genomes were obtained by de novo assembly, all reads from the dataset of interest were mapped on the reference genome of the closest virus found in the taxonomic classification step. Prokka (Seemann, 2014) was also used to predict and annotate open reading frames from these longest contigs. Viral sequences generated in this study have been deposited in GenBank with Accession numbers as shown in Table 3

Phylogenetic analysis

Reference sequences were downloaded from RefSeq (www.ncbi.nlm.nih.gov/refseq/) and in addition to the top blast hits, amino acid or nucleotide sequences, were aligned using MAFFT (Katoh & Standley, 2013) in Geneious Prime (www.geneious.com/), and gaps trimmed using trimAL (Capella-Gutiérrez, Silla-Martínez, & Gabaldón, 2009). Phylogenetic trees were constructed using IQ-TREE (Nguyen, Schmidt, von Haeseler, & Minh, 2015) with the best model of amino acid or nucleotide substitution.

Based on BLAST results and phylogenetics, viral contigs were not included in the analysis if they were likely contaminants. For example, detecting viruses in clades dominated by arthropod viruses were not included. Furthermore, short contigs that blasted only against highly conserved 3' ends of viruses likely represented misassignment. Finally, we cross-referenced our data with well-known lists of viral contaminants (Asplund et al., 2019). For all virus taxa, trees were

constructed using the ORF containing the RNA-dependant RNA polymerase (RdRp). For the flaviviruses, we also included NS3 non-structural polyprotein, as this previously has also been found suitable (Atif et al., 2016; Rice, 1996).

We considered viruses to be novel if they had a <90% RdRp protein identity, <80% genome identity (Shi et al., 2018; Shi et al., 2016), and/or following cut-off thresholds specified by the International Committee on the Taxonomy of Viruses (ICTV) for each viral family (Lefkowitz et al., 2018).

Results

pan-PCR screening of viruses in blood samples from Ugandan cattle

To understand the degree of viral diversity and presence of specific viruses we screened 175 blood samples obtained from cattle in Uganda using a pan- PCR approach. The cattle ranged in age from <1-12 years (Median, 3.5; Mean, 3.7), and the breeds were Zebu (64.7%), Ankole (27.7%) and others (7.6%). At the time of sample collection, sampled cattle had rectal temperatures ranging from 36.0 – 40.6°C, and an average of 68 ticks per animal (Range: 0 – 450). However, the cattle from which we detected viruses were mainly young ones with an average age of 2.9 years (Range = 1-8), 60% of which were of the Ankole breed. These cattle had a normal rectal temperature (Range: 37.3 – 38.2°C) and a low to moderate tick burden (Range: 2 – 132 ticks).

Using a pan PCR approach, we found that 2.9% (5/175) of the samples were positive for either of 2 viral taxa. These included orthobunyaviruses (n=2) from samples collected in Kasese and Gulu districts, and rhabdoviruses (n=3) from samples collected in Gulu, Soroti and Moroto districts. In addition, we detected non-specific (indeterminate) electrophoretic bands in 5 samples

234 that were obtained from animals in the districts of Hoima (n=2), Kasese (n=2) and Soroti (n=1),
235 particularly for orthobunyaviruses (Figure 2).

236

237 Table 2: Characteristics of Pan-PCR Positive Cattle, Uganda, 2017.

Animal ID#	Sample Collection Area		Animal Characteristics				Results of Pan-PCR
	District	GPS Location	Age ^a	Sex	Breed	Health	
G04B	Gulu	N 03.03243, E 032.30449	2	M	Ankole	Good	Positive (Rhabdovirus)
G11A	Gulu	N 03.01411, E 032.28941	6	F	Ankole	Good	Positive (Orthobunyavirus)
H11A	Hoima	N 01.64420, E 031.29474	5	F	Ankole	Good	Indeterminate (Orthobunyavirus)
H26B	Hoima	N 01.32115, E 031.23425	2	F	Zebu	Good	Indeterminate (Orthobunyavirus)
K09B	Kasese	S 00.10048, E 029.82413	8	F	Ankole	Good	Indeterminate (Orthobunyavirus)
K16B	Kasese	S 00.04747, E 029.82864	2	F	Ankole	Good	Indeterminate (Orthobunyavirus)
K28A	Kasese	N 00.29223, E 030.16044	1	M	Ankole	Good	Positive (Orthobunyavirus)
M15B	Moroto	N 02.53587, E 034.53321	1	F	Zebu	Good	Positive (Rhabdovirus)
S37B	Soroti	N 01.92280, E 033.49659	1	M	Zebu	Good	Indeterminate (Orthobunyavirus)
S41B	Soroti	N 01.93135, E 033.49284	1	M	Zebu	Good	Positive (Rhabdovirus)

239

240 Figure 2. Gel images for some selected positive cattle samples (A) *Rhabdoviridae* and (B)
 241 *Orthobunyavirales*. Animal samples names correspond to G04B, M15B, S41B, G11A and K28A.
 242 LDR, NC, PC and IND comprise of the DNA Ladder, Negative Control, Positive Control, and the
 243 ‘Indeterminate’ pan-PCR samples, respectively.

244

245 *Large diversity of viruses revealed using metagenomics*

To better understand the viral diversity, we performed metagenomic analysis on serum samples corresponding to the 10 positive pan-PCR whole blood samples. Of these 10 serum samples, we found evidence of at least one virus in 9 samples (Table 3). In one sample (G11A in Table 2) that had been collected from Gulu district, and was initially pan-PCR positive for orthobunyaviruses, a 382bp contig had a 98.9% nucleotide similarity to *Crimean-Congo hemorrhagic fever orthonairovirus*. Upon further analysis, these short contigs also had high similarity to reverse genetic vectors (KJ648916, 98.91% identity, p -value $1e-85$), and had low similarity and large gaps when aligned to other *Crimean-Congo hemorrhagic fever orthonairoviruses*, and hence this contig was removed from further data analysis. From the remaining 9 samples, metagenomic sequencing identified a total of 8 viruses, belonging to 4 families and 6 genera (Table 3). Importantly, this metagenomic approach confirmed the presence of virus positive samples generated in the pan-PCR approach.

Despite the pan-PCR approach failing to identify flaviviruses in the samples, metagenomics identified viruses belonging to this family in 3 of the submitted samples. These samples had showed non-specific gel bands on conventional pan-PCR. The identified flavivirus species were Bovine hepatitis virus (Species *Hepacivirus N*, Genus *Hepacivirus*) in samples collected from Gulu and Hoima districts, and a novel flavivirus most closely related to Bat pestivirus (Genus *Pestivirus*) from Kasese district. This new flavivirus, which we have tentatively named Zikole virus, shares 74.5% nucleotide and 81% amino acid similarity to Bat pestivirus (Figure 3). Phylogenetically, Zikole virus is highly divergent to the clades containing pestiviruses from cattle, swine, and other ungulates (Figure 3).

Sequencing revealed three members of *Rhabdoviridae* belonging to the genus *Ephemerovirus*. These 3 viruses were Obodhiang virus (Species *Obodhiang ephemerovirus*)

detected in a sample from Moroto district, and two novel viral species – one related to Puchong virus from a sample collected in Gulu district that we have tentatively named Kokolu virus, and another related to Kotonkan virus (Species *Kotonkan ephemerovirus*) in a sample from Soroti district that we have tentatively named Zebtine virus (Figure 4). The Kokolu virus shares 78.3% identity at the nucleotide level and 90.5% at the amino acid level across the L gene (containing the RdRp) with Puchong virus, while Zebtine virus shares 78.0% identity at the nucleotide level and 86.7% at the amino acid level, across the L gene with Kotonkan ephemerovirus. As per the current ICTV guidelines for species demarcation, rhabdovirus species must have a minimum of 15% amino acid sequence divergence in the L gene (ICTV, 2017). As such, the Zebtine virus identified in this study, falls well outside that threshold, but the status of Kokolu virus is less clear given its high nucleotide divergence, despite its relatively low (9.5%) amino acid divergence. The strain of Obodhiang virus identified in this study represents the first detection in Uganda since its description in mosquitos collected from South Sudan in 1963 (Schmidt, Williams, Lule, Mivule, & Mujomba, 1965), substantially expanding its geographic range.

We also identified Schmallerberg virus (Species *Schmallerberg orthobunyavirus*, Genus *Orthobunyavirus*, Family *Peribunyaviridae*) from a sample collected from Kasese district, with a 91.1% nucleotide and 97.6% amino acid sequence identity in the RdRp gene (Figure 5). Across the L segment, the virus revealed here is sister to the clade containing Shamonda and Schmallerberg viruses, although the M segment falls into the clade comprising Shamonda virus, and more distantly related to Schmallerberg virus (Figure 5).

The final two virus species identified in this study belong to the Family *Reoviridae* (Figure 6); the first, which we have tentatively called Zeboroti virus, is a novel virus closely related to Tarumizu tick virus (Species *Tarumizu Coltivirus*, Genus *Coltivirus*) in a sample from Soroti

district, with 84.5% nucleotide and 94.4% amino acid identity. According to the ICTV species demarcation guidelines for this genus, viruses belong to the same species if they have >89% nucleotide identity in segment 12, or amino acid identities of >55%, >57% and >60% in VP6, VP7 and VP12, respectively (Attoui et al., 2011). Unfortunately, we were only able to find genomic sequences belonging to segment 1 (out of the 12 putative segments), and therefore, ascertaining whether this virus is novel is challenging. Second, we identified *Aedes pseudoscutellaris* reovirus (Species *Aedes pseudoscutellaris reovirus*), the prototype-species for the genus *Dinovernavirus* of the *Reoviridae* family in two samples (one from Kasese district and another from Hoima district) with 98.2% and 100% nucleotide and amino acid identity, respectively. This virus is not well characterized, and it is unclear whether this virus is an arbovirus, an insect-specific virus, or a contaminant.

303 Table 3. Viruses identified by Metagenomics from Pan-PCR Positive Cattle, Uganda, 2017.

Animal ID	Contig Characteristics					Identified Virus		
	No.	Length (bp)	Full Length?	BLASTN Pident (%)	BLASTX Pident (%)	Virus Family (Genus)	Virus Species (Name given from this study)	GenBank Accession Numbers
H11A	3	1,140 - 3,019	-	85.9	-	<i>Flaviviridae, (Hepacivirus)</i>	<i>Bovine hepacivirus</i>	CAJZAA010000001-CAJZAA010000003
G04B	1	8645	-	87.2	-	<i>Flaviviridae, (Hepacivirus)</i>	<i>Bovine hepacivirus</i>	OU592967
K16B	2	5,421 - 6,972	Yes	74.5	-	<i>Flaviviridae, (Pestivirus)</i>	Novel pestivirus (Zikole virus)	OU592965
K28A	4	1,218 - 3,074	No	91.1	97.6	<i>Peribunyaviridae, (Orthobunyavirus)</i>	<i>Schmallenberg virus</i>	CAJZAJ010000001-CAJZAJ010000004
S37B	1	2273	No	84.5	94.4	<i>Reoviridae, (Coltivirus)</i>	Novel Coltivirus (Zeboroti virus)	OU592974
H26B	2	1,031 - 1,410	-	99.2	100	<i>Reoviridae, (Dinovernavirus)</i>	<i>Aedes pseudoscutellaris reovirus</i>	CAJZAK010000001-CAJZAK010000002
K09B	4	1,013 - 1,167	-	98.2	-	<i>Reoviridae, (Dinovernavirus)</i>	<i>Aedes pseudoscutellaris reovirus</i>	CAJZAC010000001-CAJZAC010000004
M15B	1	14618	Yes	96.4	98.3	<i>Rhabdoviridae, (Ephemerovirus)</i>	<i>Obodhiang virus</i>	OU592970
G04B	1	14630	Yes	78.3	90.5	<i>Rhabdoviridae, (Ephemerovirus)</i>	Novel Ephemerovirus (Kokolu virus)	OU592964
S41B	3	1,400 - 12,917	-	78.0	86.7	<i>Rhabdoviridae, (Ephemerovirus)</i>	Novel Ephemerovirus (Zebtine virus)	CAJZAB010000001-CAJZAB010000003

305

306 Figure 3. Phylogeny of select taxa of the *Flaviviridae*. Amino acid phylogeny (A, B) and nucleotide phylogeny (C, D) of *Pestivirus*
 307 (*Zikole virus*) and *Hepacivirus N* (Bovine Hepacivirus). Phylogenies in A, C are of the polypeptide and B, D are of the NS3 Helicase.
 308 For *Pestivirus*, the tree was midpoint rooted for clarity only, and for the Bovine Hepacivirus, the tree was rooted as per Lu *et al* (2019),
 309 including subtype information. Scale bar indicates number of substitutions per site. Sequences in this study are denoted by a filled circle.

310

311 Figure 4. Phylogeny of the RdRp (L) protein of genus *Ephemerovirus* (Family: *Rhabdoviridae*). Ekoma virus 1 and 2 (genus *Tibrovirus*)
 312 are set as the outgroup. Scale bar indicates number of amino acid substitutions per site. Sequences in this study are denoted by a filled
 313 circle.

314

315 Figure 5. Phylogeny of the (A) L segment and (B) M segment of *Schmallenberg orthobunyavirus* including Schmallenberg Virus,
 316 Shamonda Virus, Douglas Virus and Sathuperi virus. Scale bar indicates number of nucleotide substitutions per site. Sequences in this
 317 study are denoted by a filled circle.

318

319 Figure 6. Phylogeny of *Reoviridae*. (A). Segment 1 of Coltivirus. Shelly beach virus was set as the outgroup. Scale bar indicates number
 320 of amino acid substitutions per site. (B) Segment 2 of Dinovernavirus (*Reoviridae*). The tree was mid-point rooted for clarity only. Scale
 321 bar indicates number of nucleotide substitutions per site. Sequences in this study are denoted by a filled circle.

Discussion

This study investigated RNA viruses in blood samples that were collected from cattle in 5 districts in Uganda. Overall, we identified 8 viral species, belonging to 4 distinct viral families 324 (*Flaviviridae*, *Rhabdoviridae*, *Peribunyaviridae* and *Reoviridae*). Four of the identified viral 325 species in this study were novel (Zikole virus, Zeboroti virus, Kokolu virus and Zebtine virus), 326 while the 4 other viruses (Bovine hepatitis virus, Schmallenberg virus, *Aedes pseudoscutellaris* 327 reovirus, Obodhiang virus), are being reported for the first time in Uganda. Thus, our findings, 328 perhaps, are an early indication that cattle in Uganda may be harbouring more viruses than 329 previously known, in line with similar high levels of viral diversity in cattle reported elsewhere 330 (Dacheux et al., 2019; Kato et al., 2016; Kwok, Nieuwenhuijse, Phan, & Koopmans, 2020; 331 McDaniel et al., 2014).

Interestingly, a substantial proportion of the identified and described viruses in this study 333 are transmitted by, or fall into clades, associated with arthropods. Specifically, the coltivirus 334 (Zeboroti virus) described here is most closely related, and fall within a clade dominated by tick- 335 borne viruses, some of which are found in many parts of North America and Europe, where they 336 cause acute febrile human illnesses (Fujita et al., 2017; Geissler et al., 2014; Hubálek & Rudolf, 337 2012). Other viruses detected in this study were mainly those currently associated with mosquito 338 and culicoid flies (Yanase, Murota, & Hayama, 2020). Ticks are a major vector for animal diseases 339 in Uganda. And, we have previously reported a high tick burden, comprising of several tick species 340 on cattle under this study (Balinandi et al., 2020). Taken together, the high tick burden with high 341 diversity of tick species parasitizing cattle provide avenues for likely transmission of several 342 infectious agents, some of which may be novel. 343

Indeed, from our study, we report 4 novel viruses with unknown epidemiology in Uganda. Unfortunately, a major limitation of metagenomics is that, beyond detection, we are unable to determine the pathogenicity or medical-veterinary importance of the identified novel viruses. However, their presence in cattle blood indicates potential for spillover into the environment through vectors, or other biological and mechanical mechanisms. Of particular interest, was the identification of a pestivirus in our study, which we have tentatively named as Zikole virus. Pestiviruses are a group of reemerging viruses that have gained significant interest due to their wide geographical distribution, associated high socioeconomic losses when in livestock, a variety of syndromic manifestations in affected animals, failures in eradication programs, and a growing number of newly identified domestic and wildlife hosts (Blome, Beer, & Wernike, 2017; Postel, Smith, & Becher, 2021). Currently, they are known to infect a variety of artiodactyls, including swine and ruminants, in which they cause mild to severe disease (Schweizer & Peterhans, 2014). The Zikole virus identified in this study, had a close phylogenetic relationship to Bat pestivirus. While this may represent a mere common ancestry between these two viral species, recent studies have implicated bats as a major reservoir from which several known and unknown viral species have spilled over into other animal species (Wang & Anderson, 2019). From our study, the significance and impact of the identified Zikole virus on cattle, as well as human health remains unknown and a subject for further investigation. What is known, however, is that the common species of the genus *Pestivirus*, Bovine viral diarrhea virus type 1, now re-classified as *Pestivirus A* (Smith et al., 2017) causes serious disease in cattle (Liebler-Tenorio, Kenklies, Greiser-Wilke, Makoschey, & Pohlenz, 2006).

As much as some of the viruses identified in this study are being reported for the first time in Uganda, they are known to be widely spread elsewhere in African bovines (Corman et al., 2015;

Sibhat, Ayelet, Gebremedhin, Skjerve, & Asmare, 2018), where their clinical manifestations can range from absent overt disease to abortions, stillbirths and/or congenital malformations (Baechlein et al., 2015; Peperkamp et al., 2015). In this study, all virus-positive animals displayed normal rectal temperatures (Table 2), perhaps, indicating their tolerance to these viral infections (Ayres & Schneider, 2012). Nevertheless, a detailed characterization of these viruses is still crucial in the context of risk analysis, design of public health surveillance systems, and outbreak investigation and response. Of concern is the fact that many, if not all, of the identified viruses in the present study still lack effective vaccines or antiviral treatments. They are of unknown epidemiology, including their circulation in nature, range of potential natural reservoirs and immunological responses in relation to clinical infections, whether in man or in animals. Thus, the main limitation of this study emanated from its cross-sectional nature, which hindered us from obtaining detailed information regarding these epidemiological perspectives.

Funding

This work was supported by funds obtained from the Swedish Research Council to the College of Veterinary Medicine, Animal Resources and Biosecurity, Makerere University, Kampala, Uganda (Grant# 2016-05705). MW is funded by an Australian Research Council Discovery Early Career Award Fellowship (DE200100977).

Ethical Statement

The authors confirm that they read and adhered to journal's ethical policies as stated in the authors guidelines. The required ethical clearances to collect ticks and blood samples from cattle were

obtained from the animal ethical committee and a research permit issued by the Uganda National Council of Science and Technology (UNCST).

Data Availability Statement

Data supporting all components of this study is available on request from the corresponding author. Sequence data was submitted to GeneBank and can be accessed using accession numbers provided in the body of the text.

Declaration of Competing Interest

The authors declare no conflict of interest.

CRedit Authorship Contribution Statement

Stephen Balinandi: investigation, methodology, visualization, writing - original draft **Juliette Hayer:** methodology, data curation, formal analysis, writing - review and editing **Harindranath Cholleti:** methodology, data curation, investigation, writing - review and editing **Michelle Wille:** methodology, data curation, formal analysis, visualization, writing - review and editing **Julius J. Lutwama:** investigation, supervision, writing - review and editing, **Maja Malmberg:** conceptualization, funding acquisition, investigation, methodology, data curation, formal analysis, resources, supervision, project administration, writing - review and editing, **Lawrence Mugisha:** conceptualization, investigation, supervision, project administration, resources, methodology, visualisation, formal analysis, original draft, writing - review and editing, validation.

Acknowledgements

We thank Dr. Bernard Ssebide from the Mountain Gorilla Veterinary and
USAID/PREDICT projects in Uganda, for providing pan-PCR primers that were used in this study.
In addition, sequencing was performed by the SNP&SEQ Technology Platform in Uppsala,
Sweden. The facility is part of the National Genomics Infrastructure and Science for Life
Laboratory. The SNP&SEQ Platform is also supported by the Swedish Research Council and the
Knut and Alice Wallenberg Foundation. We also thank Conservation and Ecosystem Health
Alliance (CEHA) that coordinated the field sample collection activities together with livestock
farmers that availed their animals for the study.

References

- Albariño, C. G., Foltzer, M., Towner, J. S., Rowe, L. A., Campbell, S., Jaramillo, C. M., . . .
Ströher, U. (2014). Novel paramyxovirus associated with severe acute febrile disease,
South Sudan and Uganda, 2012. *Emerging Infectious Diseases*, 20(2), 211-216.
doi:10.3201/eid2002.131620
- Allen, T., Murray, K. A., Zambrana-Torrel, C., Morse, S. S., Rondinini, C., Di Marco, M., . . .
Daszak, P. (2017). Global hotspots and correlates of emerging zoonotic diseases. *Nature
Communications*, 8(1), 1124-1124. doi:10.1038/s41467-017-00923-8
- Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., & Lipman, D. J.
(1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search
programs. *Nucleic Acids Research*, 25(17), 3389-3402. doi:10.1093/nar/25.17.3389

431 Asplund, M., Kjartansdóttir, K. R., Møllerup, S., Vinner, L., Fridholm, H., Herrera, J. A. R., . . .
432 Hansen, A. J. (2019). Contaminating viral sequences in high-throughput sequencing
433 viromics: a linkage study of 700 sequencing libraries. *Clinical Microbiology and Infection*,
434 25(10), 1277-1285. doi:10.1016/j.cmi.2019.04.028

435 Atif, M., Imran, M., Qamar, Z., Javaid, M. U., Irfan, M., Ahmed, A., . . . Waqar, A. (2016).
436 Phylogeny of Dengue Virus 2 based upon the NS3 Gene among USA, Thailand, Singapore,
437 Japan and Philippine. *Journal of Human Virology & Retrovirology*, 3.

438 Attoui, H., Mertens, P., Becnel, Belaganahalli, M., Bergoin, Brussaard, C., . . . Zhou, H. (2011).
439 Family Reoviridae. In A. M. Q. King, M. J. Adams, E. B. Carstens, & E. J. Lefkowitz
440 (Eds.), *Virus Taxonomy: Ninth Report of the International Committee on Taxonomy of*
441 *Viruses*
442 (pp. 541-637).

443 Ayres, J. S., & Schneider, D. S. (2012). Tolerance of infections. *Annual Review of Immunology*,
444 30, 271-294. doi:10.1146/annurev-immunol-020711-075030

445 Baechlein, C., Fischer, N., Grundhoff, A., Alawi, M., Indenbirken, D., Postel, A., . . . Becher, P.
446 (2015). Identification of a Novel Hepacivirus in Domestic Cattle from Germany. *Journal*
447 *of Virology*, 89(14), 7007-7015. doi:10.1128/jvi.00534-15

448 Balinandi, S., Chitimia-Dobler, L., Grandi, G., Nakayiki, T., Kabasa, W., Bbira, J., . . . Mugisha,
449 L. (2020). Morphological and molecular identification of ixodid tick species (Acari:
450 Ixodidae) infesting cattle in Uganda. *Parasitology Research*, 119(8), 2411-2420.
451 doi:10.1007/s00436-020-06742-z

- 452 Balinandi, S., von Brömssen, C., Tumusiime, A., Kyondo, J., Kwon, H., Monteil, V. M., . . .
453 Malmberg, M. (2021). Serological and molecular study of Crimean-Congo hemorrhagic
454 fever virus in cattle from selected districts in Uganda. *Journal of Virological Methods*,
455 114075. doi:10.1016/j.jviromet.2021.114075
- 456 Blome, S., Beer, M., & Wernike, K. (2017). New Leaves in the Growing Tree of Pestiviruses.
457 *Advances in Virus Research*, 99, 139-160. doi:10.1016/bs.aivir.2017.07.003
- 458 Borchert, J. N., Tappero, J. W., Downing, R., Shoemaker, T., Behumbiize, P., Aceng, J., . . .
459 Prevention. (2014). Rapidly building global health security capacity--Uganda
460 demonstration project, 2013. *MMWR. Morbidity and Mortality Weekly Report*, 63(4), 73-
461 76.
- 462 Buchfink, B., Xie, C., & Huson, D. H. (2015). Fast and sensitive protein alignment using
463 DIAMOND. *Nature Methods*, 12(1), 59-60. doi:10.1038/nmeth.3176
- 464 Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. (2009). trimAl: a tool for automated
465 alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, 25(15), 1972-
466 1973. doi:10.1093/bioinformatics/btp348
- 467 Chen, Y., Ye, W., Zhang, Y., & Xu, Y. (2015). High speed BLASTN: an accelerated MegaBLAST
468 search tool. *Nucleic Acids Research*, 43(16), 7762-7768. doi:10.1093/nar/gkv784
- 469 Cleaveland, S., Sharp, J., Abela-Ridder, B., Allan, K. J., Buza, J., Crump, J. A., . . . Halliday, J. E.
470 B. (2017). One Health contributions towards more effective and equitable approaches to
471 health in low- and middle-income countries. *Philosophical Transactions of the Royal*
472 *Society of London. Series B, Biological sciences*, 372(1725), 20160168.
473 doi:10.1098/rstb.2016.0168

- Corman, V. M., Grundhoff, A., Baechlein, C., Fischer, N., Gmyl, A., Wollny, R., . . . Drexler, J. F. (2015). Highly divergent hepaciviruses from African cattle. *Journal of Virology*, 89(11), 5876-5882. doi:10.1128/JVI.00393-15
- Crookshank, E. M. (1897). *Infectious pleuro-pneumonia*. Philadelphia, PA: W. B. Saunders.
- Dacheux, L., Dommergues, L., Chouanibou, Y., Doméon, L., Schuler, C., Bonas, S., . . . Métras, R. (2019). Co-circulation and characterization of novel African arboviruses (genus Ephemerovirus) in cattle, Mayotte island, Indian Ocean, 2017. *Transboundary and Emerging Diseases*, 66(6), 2601-2604. doi:10.1111/tbed.13323
- Deng, Y., Li, J., Wu, S., Zhu, Y., Chen, Y., & He, F. (2006). Integrated nr database in protein annotation system and its localization. *Computer Engineering*, 32(5), 71-74.
- Dick, G. W., Kitchen, S. F., & Haddow, A. J. (1952). Zika virus. I. Isolations and serological specificity. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 46(5), 509-520. doi:10.1016/0035-9203(52)90042-4
- Drichi, P., & National Biomass, S. (2003). *National biomass study : technical report of 1996-2002*. (9970863002 9789970863006). Kampala: National Biomass Study, Forest Dept.
- Edridge, A. W. D., Deijs, M., Namazzi, R., Cristella, C., Jebbink, M. F., Maurer, I., . . . van der Hoek, L. (2019). Novel Orthobunyavirus Identified in the Cerebrospinal Fluid of a Ugandan Child With Severe Encephalopathy. *Clinical Infectious Diseases : an official publication of the Infectious Diseases Society of America*, 68(1), 139-142. doi:10.1093/cid/ciy486

494 Fagre, A. C., Lee, J. S., Kityo, R. M., Bergren, N. A., Mossel, E. C., Nakayiki, T., . . . Kading, R.
 495 C. (2019). Discovery and Characterization of Bukakata orbivirus (Reoviridae:Orbivirus),
 496 a Novel Virus from a Ugandan Bat. *Viruses*, *11*(3), 209. doi:10.3390/v11030209

497 Fujita, R., Ejiri, H., Lim, C. K., Noda, S., Yamauchi, T., Watanabe, M., . . . Sawabe, K. (2017).
 498 Isolation and characterization of Tarumizu tick virus: A new coltivirus from
 499 Haemaphysalis flava ticks in Japan. *Virus Research*, *242*, 131-140.
 500 doi:10.1016/j.virusres.2017.09.017

501 Furuse, Y., Suzuki, A., & Oshitani, H. (2010). Origin of measles virus: divergence from rinderpest
 502 virus between the 11th and 12th centuries. *Virology Journal*, *7*, 52. doi:10.1186/1743-
 503 422x-7-52

504 Geissler, A. L., Thorp, E., Van Houten, C., Lanciotti, R. S., Panella, N., Cadwell, B. L., . . . Staples,
 505 J. E. (2014). Infection with Colorado tick fever virus among humans and ticks in a national
 506 park and forest, Wyoming, 2010. *Vector Borne and Zoonotic Diseases (Larchmont, N.Y.)*,
 507 *14*(9), 675-680. doi:10.1089/vbz.2013.1568

508 Grace D, Mutua F, Ochungo P, Kruska R, Jones K, Brierley L, . . . F., O. (2012). *Mapping of*
 509 *poverty and likely zoonoses hotspots*. Retrieved from <https://hdl.handle.net/10568/21161>

510 Hubálek, Z., & Rudolf, I. (2012). Tick-borne viruses in Europe. *Parasitology Research*, *111*(1), 9-
 511 36. doi:10.1007/s00436-012-2910-1

512 Hubálek, Z., Rudolf, I., & Nowotny, N. (2014). Arboviruses pathogenic for domestic and wild
 513 animals. *Advances in Virus Research*, *89*, 201-275. doi:10.1016/b978-0-12-800172-
 514 1.00005-7

- ICTV. (2017). Genus: Ephemerovirus. Retrieved from https://talk.ictvonline.org/ictv-reports/ictv_online_report/negative-sense-rna-viruses/w/rhabdoviridae/792/genus-ephemerovirus
- Jones, K. E., Patel, N. G., Levy, M. A., Storeygard, A., Balk, D., Gittleman, J. L., & Daszak, P. (2008). Global trends in emerging infectious diseases. *Nature*, 451(7181), 990-993. doi:10.1038/nature06536
- Kato, T., Shirafuji, H., Tanaka, S., Sato, M., Yamakawa, M., Tsuda, T., & Yanase, T. (2016). Bovine Arboviruses in Culicoides Biting Midges and Sentinel Cattle in Southern Japan from 2003 to 2013. *Transboundary and Emerging Diseases*, 63(6), e160-e172. doi:10.1111/tbed.12324
- Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution*, 30(4), 772-780. doi:10.1093/molbev/mst010
- Kemunto, N., Mogoa, E., Osoro, E., Bitek, A., Kariuki Njenga, M., & Thumbi, S. M. (2018). Zoonotic disease research in East Africa. *BMC Infectious Diseases*, 18(1), 545-545. doi:10.1186/s12879-018-3443-8
- Kiyaga, C., Sendagire, H., Joseph, E., McConnell, I., Grosz, J., Narayan, V., . . . Opio, A. (2013). Uganda's new national laboratory sample transport system: a successful model for improving access to diagnostic services for Early Infant HIV Diagnosis and other programs. *PLoS One*, 8(11), e78609. doi:10.1371/journal.pone.0078609

535 Kwok, K. T. T., Nieuwenhuijse, D. F., Phan, M. V. T., & Koopmans, M. P. G. (2020). Virus
536 Metagenomics in Farm Animals: A Systematic Review. *Viruses*, 12(1).
537 doi:10.3390/v12010107

538 Lamorde, M., Mpimbaza, A., Walwema, R., Kamya, M., Kapisi, J., Kajumbula, H., . . . Kugeler,
539 K. J. (2018). A Cross-Cutting Approach to Surveillance and Laboratory Capacity as a
540 Platform to Improve Health Security in Uganda. *Health Security*, 16(S1), S76-s86.
541 doi:10.1089/hs.2018.0051

542 Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature*
543 *Methods*, 9(4), 357-359. doi:10.1038/nmeth.1923

544 Lefkowitz, E. J., Dempsey, D. M., Hendrickson, R. C., Orton, R. J., Siddell, S. G., & Smith, D. B.
545 (2018). Virus taxonomy: the database of the International Committee on Taxonomy of
546 Viruses (ICTV). *Nucleic Acids Research*, 46(D1), D708-D717. doi:10.1093/nar/gkx932

547 Li, D., Luo, R., Liu, C. M., Leung, C. M., Ting, H. F., Sadakane, K., . . . Lam, T. W. (2016).
548 MEGAHIT v1.0: A fast and scalable metagenome assembler driven by advanced
549 methodologies and community practices. *Methods : a Companion to Methods in*
550 *Enzymology*, 102, 3-11. doi:10.1016/j.ymeth.2016.02.020

551 Liebler-Tenorio, E. M., Kenkies, S., Greiser-Wilke, I., Makoschey, B., & Pohlenz, J. F. (2006).
552 Incidence of BVDV1 and BVDV2 infections in cattle submitted for necropsy in Northern
553 Germany. *Journal of Veterinary Medicine. B, Infectious Diseases and Veterinary Public*
554 *Health*, 53(8), 363-369. doi:10.1111/j.1439-0450.2006.00992.x

555 Lipman, D. J., & Pearson, W. R. (1985). Rapid and sensitive protein similarity searches. *Science*,
556 227(4693), 1435-1441. doi:10.1126/science.2983426

557 Lu, G., Ou, J., Zhao, J., & Li, S. (2019). Presence of a Novel Subtype of Bovine Hepacivirus in
558 China and Expanded Classification of Bovine Hepacivirus Strains Worldwide into 7
559 Subtypes. *Viruses*, 11(9), 843.

560 Lu, R., Zhao, X., Li, J., Niu, P., Yang, B., Wu, H., . . . Tan, W. (2020). Genomic characterisation
561 and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor
562 binding. *The Lancet*, 395(10224), 565-574. doi:doi: 10.1016/S0140-6736(20)30251-8

563 Malmberg, M., & Hayer, J. (2019). Ticks and tickborne diseases in Africa. Retrieved from
564 <https://ticksinafrica.org/>.

565 McDaniel, C. J., Cardwell, D. M., Moeller, R. B., Jr., & Gray, G. C. (2014). Humans and cattle: a
566 review of bovine zoonoses. *Vector Borne and Zoonotic Diseases (Larchmont, N.Y.)*, 14(1),
567 1-19. doi:10.1089/vbz.2012.1164

568 Mehnd, M. S., Al-Shorbaji, F., Millett, P., & Murgue, B. (2018). The WHO R&D Blueprint: 2018
569 review of emerging infectious diseases requiring urgent research and development efforts.
570 *Antiviral Research*, 159, 63-67. doi:10.1016/j.antiviral.2018.09.009

571 Morse, S. S. (2004). Factors and determinants of disease emergence. *Revue Scientifique et*
572 *Technique / Office International des épizooties*, 23(2), 443-451.
573 doi:10.20506/rst.23.2.1494

574 Moureau, G., Temmam, S., Gonzalez, J. P., Charrel, R. N., Grard, G., & de Lamballerie, X. (2007).
575 A real-time RT-PCR method for the universal detection and identification of flaviviruses.
576 *Vector Borne and Zoonotic Diseases*, 7(4), 467-477. doi:10.1089/vbz.2007.0206

577 Nguyen, L. T., Schmidt, H. A., von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: a fast and
578 effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular*
579 *Biology and Evolution*, 32(1), 268-274. doi:10.1093/molbev/msu300

580 Peperkamp, N. H., Luttikholt, S. J., Dijkman, R., Vos, J. H., Junker, K., Greijdanus, S., . . . Wouda,
581 W. (2015). Ovine and Bovine Congenital Abnormalities Associated With Intrauterine
582 Infection With Schmallenberg Virus. *Veterinary Pathology*, 52(6), 1057-1066.
583 doi:10.1177/0300985814560231

584 Postel, A., Smith, D. B., & Becher, P. (2021). Proposed Update to the Taxonomy of Pestiviruses:
585 Eight Additional Species within the Genus Pestivirus, Family Flaviviridae. *Viruses*, 13(8).
586 doi:10.3390/v13081542

587 PREDICT. (2014). Reducing pandemic risk, promoting global health. *One Health Institute, Davis,*
588 *CA: University of California, Davis.* Retrieved from
589 <https://www.usaid.gov/sites/default/files/documents/1864/predict-global-flyer-508.pdf>

590 Rice, C. (1996). *Flaviviridae: the viruses and their replication*. Philadelphia, PA: Lippincott-
591 Raven.

592 Sanchez-Seco, M. P., Rosario, D., Quiroz, E., Guzman, G., & Tenorio, A. (2001). A generic
593 nested-RT-PCR followed by sequencing for detection and identification of members of the
594 alphavirus genus. *Journal of Virological Methods*, 95(1-2), 153-161.

595 Schmidt, J., Williams, M., Lule, M., Mivule, A., & Mujomba, E. (1965). Viruses isolated from
596 mosquitoes collected in the southern Sudan and western Ethiopia. *East African Virus*
597 *Research Institute Report*, 15, 24.

598 Schweizer, M., & Peterhans, E. (2014). Pestiviruses. *Annual Review of Animal Biosciences*, 2,
599 141-163. doi:10.1146/annurev-animal-022513-114209

600 Seemann, T. (2014). Prokka: rapid prokaryotic genome annotation. *Bioinformatics*, 30(14), 2068-
601 2069. doi:10.1093/bioinformatics/btu153

602 Shi, M., Lin, X.-D., Chen, X., Tian, J.-H., Chen, L.-J., Li, K., . . . Zhang, Y.-Z. (2018). The
603 evolutionary history of vertebrate RNA viruses. *Nature*, 556(7700), 197-202.
604 doi:10.1038/s41586-018-0012-7

605 Shi, M., Lin, X.-D., Tian, J.-H., Chen, L.-J., Chen, X., Li, C.-X., . . . Zhang, Y.-Z. (2016).
606 Redefining the invertebrate RNA virosphere. *Nature*, 540(7634), 539-543.
607 doi:10.1038/nature20167

608 Shoemaker, T. R., Balinandi, S., Tumusiime, A., Nyakarahuka, L., Lutwama, J., Mbidde, E., . . .
609 Nichol, S. T. (2018). Impact of enhanced viral haemorrhagic fever surveillance on outbreak
610 detection and response in Uganda. *The Lancet. Infectious Diseases*, 18(4), 373-375.
611 doi:10.1016/S1473-3099(18)30164-6

612 Sibhat, B., Ayelet, G., Gebremedhin, E. Z., Skjerve, E., & Asmare, K. (2018). Seroprevalence of
613 Schmallenberg virus in dairy cattle in Ethiopia. *Acta tropica*, 178, 61-67.
614 doi:10.1016/j.actatropica.2017.10.024

615 Smith, D. B., Meyers, G., Bukh, J., Gould, E. A., Monath, T., Scott Muerhoff, A., . . . Becher, P.
616 (2017). Proposed revision to the taxonomy of the genus Pestivirus, family Flaviviridae.
617 *The Journal of General Virology*, 98(8), 2106-2112. doi:10.1099/jgv.0.000873

- 618 Smithburn, K., Hughes, T., Burke, A., & Paul, J. (1940). A neurotropic virus isolated from the
619 blood of a native of Uganda¹. *The American Journal of Tropical Medicine and Hygiene*,
620 *1*(4), 471-492.
- 621 Towner, J. S., Sealy, T. K., Khristova, M. L., Albariño, C. G., Conlan, S., Reeder, S. A., . . . Nichol,
622 S. T. (2008). Newly discovered ebola virus associated with hemorrhagic fever outbreak in
623 Uganda. *PLoS Pathogens*, *4*(11), e1000212-e1000212. doi:10.1371/journal.ppat.1000212
- 624 UBOS. (2017). Uganda Bureau of Statistics. Retrieved October 27, 2019
625 [https://www.ubos.org/wp-](https://www.ubos.org/wp-content/uploads/publications/03_20182017_Statistical_Abstract.pdf)
626 [content/uploads/publications/03_20182017_Statistical_Abstract.pdf](https://www.ubos.org/wp-content/uploads/publications/03_20182017_Statistical_Abstract.pdf)
- 627 Vijgen, L., Keyaerts, E., Moës, E., Thoelen, I., Wollants, E., Lemey, P., . . . Van Ranst, M. (2005).
628 Complete genomic sequence of human coronavirus OC43: molecular clock analysis
629 suggests a relatively recent zoonotic coronavirus transmission event. *Journal of Virology*,
630 *79*(3), 1595-1604. doi:10.1128/jvi.79.3.1595-1604.2005
- 631 Wang, L.-F., & Anderson, D. E. (2019). Viruses in bats and potential spillover to animals and
632 humans. *Current Opinion in Virology*, *34*, 79-89. doi:10.1016/j.coviro.2018.12.007
- 633 Williams, M. C., Woodall, J. P., & Gillett, J. D. (1965). O'nyong-nyong fever: an epidemic virus
634 disease in East Africa. Vii. Virus isolations from man and serological studies up to July
635 1961. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, *59*, 186-197.
636 doi:10.1016/0035-9203(65)90080-5
- 637 Wood, D. E., Lu, J., & Langmead, B. (2019). Improved metagenomic analysis with Kraken 2.
638 *Genome Biology*, *20*(1), 257. doi:10.1186/s13059-019-1891-0

- Woolhouse, M. E. J., & Gowtage-Sequeria, S. (2005). Host range and emerging and reemerging pathogens. *Emerging Infectious Diseases*, 11(12), 1842-1847. doi:10.3201/eid1112.050997
- Wortmann, C. S., & Eledu, C. S. (1999). *Uganda`s agroecological zones: a guide for planners and policy makers*. Kampala: Centro Internacional de Agricultura Tropical (CIAT).
- Yanase, T., Murota, K., & Hayama, Y. (2020). Endemic and Emerging Arboviruses in Domestic Ruminants in East Asia. *Frontiers in Veterinary Science*, 7, 168-168. doi:10.3389/fvets.2020.00168
- Zhai, J., Palacios, G., Towner, J. S., Jabado, O., Kapoor, V., Venter, M., . . . Lipkin, W. I. (2007). Rapid molecular strategy for filovirus detection and characterization. *Journal of Clinical Microbiology*, 45(1), 224-226. doi:10.1128/JCM.01893-06