Intracellular copper storage and delivery in a bacterium

Jaeick Lee and Christopher Dennison*

Biosciences Institute, Newcastle University, Newcastle upon Tyne, NE2 4HH, UK. E-mail: christopher.dennison@ncl.ac.uk

Abstract

A family of cytosolic copper (Cu) storage proteins (the Csp3s) are widespread in bacteria. The Csp3s can bind large quantities of Cu(I) via their Cys-lined four-helix bundles, and the majority are cytosolic (Csp3s). This is inconsistent with the current dogma that bacteria, unlike eukaryotes, have evolved not to maintain intracellular pools of Cu due to its potential toxicity. Sporulation in *Bacillus subtilis* has been used to investigate if a Csp3 can store Cu(I) in the cytosol for a target enzyme. The activity of the Cu-requiring endospore multi-Cu oxidase *BsCotA* (a laccase) increases under Cu-replete conditions in wild type *B. subtilis*, but not in the strain lacking *BsCsp3*. Cuprous ions readily transfer from *BsCsp3*, but not from the cytosolic copper metallochaperone *BsCopZ*, to *BsCotA* *in vitro* producing active enzyme. Both *BsCsp3* and *BsCotA* are upregulated during late sporulation. The hypothesis we propose is that *BsCsp3* acquires and stores Cu(I) in the cytosol for *BsCotA*.

Introduction

Copper (Cu) is essential for most organisms, but use of this metal ion is associated with significant risks due to its potential toxicity. The availability of Cu is restricted by the presence of high-affinity sites in both eukaryotes (1) and prokaryotes (2). Import, cytosolic handling, trafficking to different locations, and storage have all been characterised in eukaryotic cells (3). In bacteria, some of these processes are either not thought to occur, or are not yet fully understood. For example, the plasma membrane protein CcoG, which reduces Cu(II) to the preferred intracellular oxidation state (Cu(I)) has only recently been identified in bacteria as a cytochrome oxidase (COX) assembly factor (4). The reduction of Cu(II) prior to import into eukaryotic cells has been known to occur for many years (3, 5). Excess Cu(I) is removed from the cytosol by probably the best-studied component of bacterial Cu homeostasis; a Cu-transporting P-type ATPase (CopA), which can be assisted by the cytosolic Cu metallochaperone CopZ (Figure 1a) (3, 6-9). Toxicity has been shown to involve Cu(I) binding in place of the native metal in cytosolic iron-sulfur (Fe-S) cluster-containing proteins (10), and Cu catalyses ROS formation (3, 7, 8). The intracellular damage that Cu can cause, and the current dearth of intracellular Cu-requiring enzymes (11), has resulted in a prevailing view that bacteria have evolved not to use Cu in the cytosol (8, 11). However, there is no *a priori* reason why bacteria, like eukaryotes, cannot utilise Cu in this compartment if mechanisms are available to enable its safe handling, i.e. by ensuring tight chelation and specific delivery. The presence of cytosolic Cu storage proteins (Csps) that can bind large quantities of Cu(I) with high affinity (12-15), provides a possible route for intracellular Cu use in bacteria.

The Csps were first identified in Gram-negative bacteria that oxidize methane (12). These methanotrophs can possess different Csp homologues, all having many Cys residues lining the cores of their four-helix bundles, enabling the binding of large quantities of Cu(I) ions (12-14). A Csp exported from the cytosol (Csp1) can store up to 52 Cu(I) ions per tetramer for the particulate (membrane-bound) methane monooxygenase (pMMO) in the model methanotroph *Methylosinus trichosporium* OB3b (*MtCsp1*) (12). *MtCsp1* is upregulated (16) at the Cu concentrations required for methane oxidation by pMMO in switchover methanotrophs, which can use a soluble Fe MMO when Cu is limiting (17). However, a cytosolic Csp homologue (*MtCsp3*) is not upregulated with pMMO in *M. trichosporium* OB3b (16).

Figure 1. Copper handling, a cytosolic Cu(I) storage protein, a Cu-requiring enzyme, and their transcription during sporulation in *B. subtilis*. (a) An overview of Cu homeostasis in *B. subtilis* including Cu (orange circles, oxidation state undefined) export by CopA and CopZ (regulated by CsoR) (19), and import by YcnJ (regulated by YcnK) (20, 22). YcnI is also membrane bound and binds Cu(II) *in vitro*, but its role in Cu homeostasis is unclear (23). The only currently known Cu-requiring enzymes in vegetative *B. subtilis* cells are two cytochrome oxidases (COXs) located on the plasma membrane (11, 24). (b) The crystal structure of Cu(I)-free BsCsp3 (PDB: 5FIG), a tetramer of four-helix bundles each with 19 Cys residues pointing into their cores enabling the binding of up to ~20 Cu(I) ions per monomer (13). (c) The crystal structure of the endospore multi-Cu oxidase (a laccase) BsCotA (PDB: 1W6L, (27)) with domains 1, 2 and 3 coloured green, slate and cyan, respectively (the linking regions are yellow). Substrates are oxidized (Sub_{red} to Sub_{ox}) at the T1 Cu centre with electrons passed to the T2/T3 trinuclear (Cu₃) cluster where oxygen is reduced to water. Detailed views of the T1 Cu site (d) and the Cu₃ cluster (e), with the side chains of coordinating residues represented as sticks, Cu ions as orange spheres and the oxygen atoms of water (bound to the T2 Cu) and hydroxide (bridging the T3 Cu ions) ligands as red spheres in (c-e). (f) Transcription profiles (29) of the *sigK* (σ^K , which facilitates spore coat protein expression, black triangles), *csp3* (red circles), *cotA* (blue squares) and *copZ* (green diamonds) genes during sporulation.

The Gram-positive bacterium *Bacillus subtilis* is an excellent model system for investigating the role of a Csp3, as its Cu homeostasis system is well characterised (Figure 1a) (7, 18-23). The *copZA* operon (Cu efflux machinery, *vide supra*) and its Cu-sensing repressor CsoR (7, 18,

19, 21) are probably the best-studied components. The membrane protein YcnJ is upregulated under Cu-limiting conditions, controlled by the suggested repressor YcnK (20, 22), and has been proposed to play a role in Cu acquisition (Figure 1a). The membrane-anchored YcnI is part of the same (*ycnKJI*) operon and is also regulated by YcnK (22). The soluble domain of YcnI binds Cu(II) *in vitro*, and this protein has been suggested to function as a Cu metallochaperone (23). Cytosolic Cu(I) could be safely stored in *B. subtilis* by the Csp3 homologue (*BsCsp3*) whose core is lined with 19 Cys residues (Figure 1b), which enable the binding of ~80 Cu(I) ions per tetramer *in vitro* (13). Overexpressing *BsCsp3* in the cytosol of a Gram-negative heterologous host (*Escherichia coli*) allows growth at otherwise harmful Cu concentrations, and the protein can acquire Cu(I) in the presence of both CopA and CopZ (15).

Only two families of Cu enzymes are currently known to be present in *B. subtilis*; COXs located on the plasma membrane and the multi-Cu oxidase (MCO; a laccase) *BsCotA* (11, 24-27). The latter is an outer spore-coat (endospore) enzyme (28) that possesses the typical type 1 (T1), 2 (T2) and 3 (T3) Cu sites of an MCO (27), which are involved in the catalytic cycle (see Figure 1c-e). It produces a melanin-like pigment thought to provide spores with protection against hydrogen peroxide and UV light (25, 28). *BsCotA* is upregulated during the latter stages of sporulation, as is *BsCsp3* (Figure 1f) (29).

BsCsp3 does not provide resistance to Cu toxicity. We have therefore tested the hypothesis that *BsCsp3* safely stores Cu(I) ions in the cytosol for a Cu-requiring enzyme by investigating the effect of gene deletion on the activity of *BsCotA* in spores grown under Cu limiting and replete conditions. The data obtained indicate a direct role for *BsCsp3* in ensuring the maximum activity of *BsCotA*. The ability of *BsCsp3* to activate *BsCotA* has been confirmed by the *in vitro* transfer of Cu(I) between these proteins. A model for how *BsCotA* acquires Cu(I) from *BsCsp3* during sporulation is proposed. This is the first example showing that an enzyme can acquire Cu in the cytosol of a bacterium as well as identifying the partner protein.

Results

Is *BsCsp3* involved in combating Cu toxicity in *B. subtilis*?

The presence of a protein with a high capacity for Cu(I) in the cytosol of *B. subtilis* (12-14) would suggest a role in helping to prevent the problems caused by excess Cu (10, 15). The toxicity of Cu to bacteria is highlighted by the influence increasing Cu concentrations in media has on the growth of wild type (WT) *B. subtilis* (Supplementary Figure 1). At higher Cu levels cells grow extremely slowly, with a very small increase in the absorbance/OD observed only after more than 6 h at 2 mM Cu, and this coincides with elevated intracellular Cu concentrations (Supplementary Figure 2). Very similar growth and Cu accumulation results are obtained for the

strain ($\Delta csp3$) lacking the *csp3* gene (Supplementary Figures 1 and 2). These data exhibit an overall likeness to work we reported previously (13), particularly at up to 12 h growth. However, the response to specific Cu concentrations varies, indicating the amounts of Cu added to media must differ. In the present work we are sure of the Cu concentrations in media, having carefully quantified all Cu(II) stocks before their addition, and the reported values in the previous study (13) appear to be too high. The growth studies reported herein demonstrate that *BsCsp3* is not involved in helping prevent the harmful effects on *B. subtilis* caused by elevated Cu levels. Therefore, the protein does not have a function like eukaryotic Cys-rich metallothioneins (3).

Using sporulation to determine the function of *BsCsp3*

The *csp3* and *cotA* genes are both upregulated (29) at similar stages during sporulation (Figure 1f), and *BsCotA* is the only known Cu-requiring enzyme present in spores. We have therefore studied whether *BsCsp3* stores Cu(I) for *BsCotA*. This enzyme binds four Cu ions (Figure 1c-e) and oxidises the laccase substrate 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) *in vitro* and in spores (26). For WT *B. subtilis* spores the ability to oxidise ABTS increases approximately two-fold when 50 μ M Cu is added to media during sporulation (Figure 2a and Supplementary Figure 3a,b). Incubation with Cu(II) only enhances the activity of purified WT spores obtained in the absence of added Cu (Supplementary Figure 4 and Table 1), consistent with previously reported data (26). This indicates that unless supplemented, sporulation media does not contain sufficient Cu ($\sim 0.4 \mu$ M) to fully metallate all of the *BsCotA* produced. However, the addition of 50 μ M Cu, a non-toxic concentration, during growth enables *B. subtilis* to obtain enough of the metal to generate only fully Cu-loaded *BsCotA* in the endospore.

The *BsCotA* activity of $\Delta csp3$ *B. subtilis* spores grown without added Cu is similar to that for WT spores produced under the same conditions (Figure 2b and Supplementary Figure 3c,d). However, unlike for WT *B. subtilis*, supplementing media with Cu during sporulation has no effect on *BsCotA* activity for the $\Delta csp3$ strain. Similar to the WT data, incubation with Cu only significantly increases the activity of purified $\Delta csp3$ spores grown in the absence of added Cu (Supplementary Figure 4 and Table 1). These results demonstrate that *BsCsp3* plays a role in storing Cu for *BsCotA*, particularly under Cu-replete conditions. Some *BsCotA* activity remains for $\Delta csp3$ *B. subtilis* spores, and an alternative mechanism of Cu transfer to *BsCotA* must exist, which could also be responsible for the activity observed in the WT strain under Cu-limiting conditions.

Figure 2. The influence of Cu levels and deleting the *csp3* gene on *BsCotA* activity in *B. subtilis*. Michaelis-Menten plots of *BsCotA* activity for heated purified spores from WT (a), $\Delta csp3$ (b), and the complemented $\Delta csp3$ (c, d) strains. Spores (a, b and c) were produced in Difco sporulation media plus no (open symbols) and 50 μM (black filled symbols) added $\text{Cu}(\text{NO}_3)_2$. (d) For the complemented $\Delta csp3$ strain sporulation was also carried out in the presence of 1 mM IPTG with either no (magenta) or 50 μM (cyan) added $\text{Cu}(\text{NO}_3)_2$. The plots from which the initial rates for WT and $\Delta csp3$ *B. subtilis* spores were obtained are shown in Supplementary Figure 3. Averages and standard deviations from kinetic measurements in 100 mM citrate-phosphate buffer pH 4.0 using three different sets of spores are shown.

To confirm that *BsCsp3* supplies Cu(I) to *BsCotA* *in vivo*, the $\Delta csp3$ strain was complemented by introducing an inducible copy of the *csp3* gene at a different location (the *amyE* locus). The highest *BsCotA* activity is obtained for spores of this strain grown in the presence of isopropyl β -D-thiogalactopyranoside (IPTG, the inducer) and Cu (Figure 2c,d). Activity is almost three-fold greater than without their addition, similar to the increase caused by Cu in WT *B. subtilis* spores (Figure 2a). Elevated activity is observed for spores from the complemented strain grown with Cu, but not IPTG, present in the media (Figure 2c). To ensure this effect is due to leaky expression, a well-established feature of the promoter used [for example, see ref. 30], a control $\Delta csp3$ strain was constructed (see Material and Methods). In this case, the ability to oxidise ABTS is hardly influenced by the presence of Cu and IPTG (Supplementary Figure 5). Therefore, the increased *BsCotA* activity for the complemented $\Delta csp3$ strain is due to the IPTG-inducible copy of the *csp3* gene.

Cu(I) transfer from *BsCsp3* to *BsCotA* *in vitro*

The above data support the hypothesis that *BsCsp3* can store Cu(I) in the cytosol under Cu-replete conditions, which is used to metallate *BsCotA*. To further test this idea, the transfer of Cu(I) from *BsCsp3* to inactive apo(Cu-free)-*BsCotA* has been studied *in vitro*. We have previously found (12, 13) that Cu(I) removal from Csp3s by a large excess of high-affinity Cu(I) ligands is slow (~60% removal in 24 h for *BsCsp3* using bathocuproine disulfonate, see Supplementary Figure 6 and Table 2). The transfer of Cu(I) from *BsCsp3* to *BsCotA* occurs quickly when using a 10-fold excess of Cu(I) in the storage protein over sites in the enzyme, and >50% maximum activity is achieved within 6 h (Figure 3 a,b and Figure 4a). A control experiment was performed studying activation of as-isolated *BsCotA* with Cu(I) in the absence of *BsCsp3*, which is faster, but still takes 4 h to complete.

Figure 3. Kinetic analysis of the activity of *BsCotA* after incubation with Cu from different sources. Plots of absorbance at 420 nm against time for the reaction with 2.4 mM ABTS of mixtures of apo-*BsCotA* incubated with buffer (a), Cu(I)-*BsCsp3* (b), Cu(I) (c), Cu(II) (d and e) and Cu(I)-*BsCspZ* (f) for 0.5, 2, 4, 6 and 24 h. Mixtures were incubated in 20 mM HEPES pH 7.5 plus 200 mM NaCl (the buffer used in a), and *BsCotA* activity was measured 100 mM citrate-phosphate buffer pH 4.0. All data (averages from three independent experiments with error bars showing standard deviations) are from mixtures incubated under anaerobic conditions, apart from in e when *BsCotA* and Cu(II) were incubated in air. The resulting activity data at these and other incubation times are shown in Figure 4.

A further control was carried out using Cu(II) and metalation of the enzyme is slower than for Cu(I) both under anaerobic and aerobic conditions (Figure 3c-e and Figure 4b). Another cytosolic Cu(I)-binding protein with a well-established role (18) in Cu homeostasis (delivering Cu(I) to *BsCspA*) and a similar Cu(I) affinity (31) to *BsCsp3* (13) is *BsCspZ* (see Figure 1a). As this was another potential source of Cu(I) for *BsCotA* we tested to see if *BsCspZ* could transfer Cu(I) to *BsCotA* (Figure 3f and Figure 4a). After the incubation of apo-*BsCotA* with Cu(I)-*BsCspZ* for 48 h very little activity is observed, and Cu(I) transfer does not occur.

Figure 4. Cu(I) transfer from *BsCsp3* to *BsCotA*. (a) Plots of activity (average from three independent experiments with error bars showing standard deviations) against incubation time of as-isolated *BsCotA* incubated with Cu(I)-*BsCsp3* (red circles), Cu(I)-*BsCspZ* (half-orange circles) and buffer alone (half-black circles) for up to 48 h under anaerobic conditions. The activity data for as-isolated *BsCotA* plus buffer alone (black circles) are also shown in (b), as well as the results from control experiments in which the enzyme was incubated with Cu(I) (green circles) and Cu(II) under anaerobic conditions (blue circles), and also with Cu(II) in air (cyan circles). The kinetic data at selected time points are shown in Figure 3.

Discussion

Herein we demonstrate that *BsCsp3* stores Cu(I) in the cytosol during sporulation for the Cu-requiring enzyme *BsCotA*. This is not the only mechanism available to load *BsCotA* with Cu as some activity is still observed in $\Delta csp3$ *B. subtilis*. A possibility we considered was that the cytosolic Cu metallochaperone *BsCspZ*, as well as transferring Cu(I) to *BsCspA*, may also store cuprous ions in the cytosol, which could be transferred to *BsCotA* (the Cu(I) affinity of *BsCspZ* is similar to that of *BsCsp3* (13, 31)). This is supported by the suggestion that *BsCspZ* could play a role in Cu(I) sequestration and recycling in *B. subtilis* (18) as less Cu accumulates in the $\Delta copZ$ strain compared to WT. Furthermore, the activity of *CotA* from a different *Bacillus* strain is enhanced when co-expressed in the cytosol of *E. coli* with its native *CopZ* (32). However, the *in vitro* studies reported here show that *BsCspZ* cannot transfer Cu(I) to *BsCotA* (Figure 3f and Figure 4a). The source(s) of Cu(I) for activating *BsCotA* in the absence of *BsCsp3*, and also at lower intracellular concentrations of the metal ion, remain(s) to be established. Regardless, the lack of Cu(I) transfer from *BsCspZ* highlights the specificity of the *BsCsp3*-*BsCotA* interaction. This is essential in a cell as it ensures protein-mediated Cu(I) transfer to the correct destination, as observed for other Cu-homeostasis proteins (1, 33-39).

Considering the high Cu(I) affinity (13) of *BsCsp3* ($(1.5 \pm 0.4) \times 10^{17} \text{ M}^{-1}$), transfer of cuprous ions to *BsCotA* has to occur via an associative mechanism (unassisted Cu(I) off-rates for *BsCsp3* can be estimated (14) to be $\sim 10^{-9} \text{ s}^{-1}$). For the acquisition of such tightly bound Cu(I) to be possible, metalation must take place once the target enzyme has at least partially folded so the sites where Cu binds have formed. The T1 Cu site is closest to the surface of *BsCotA* (the His497 ligand is solvent exposed), and is $\sim 12.5\text{-}15.5 \text{ \AA}$ from the Cu₃ cluster (Figure 1c). Therefore, *BsCsp3* association at more than one location may be required to metalate all of the sites in folded *BsCotA*. Published Cu(I) affinities of T1 Cu sites (40, 41) are $(2.1\text{-}4.0) \times 10^{17} \text{ M}^{-1}$, similar to the average Cu(I) affinity of *BsCsp3* (13). Therefore, Cu(I) transfer from the storage protein to the enzyme should not be hindered thermodynamically (37, 40). To facilitate access to the more buried Cu₃ cluster the protein may need to be partially unfolded. The MCO CueO from *E. coli* undergoes a transition from an ‘open’ non-metallated folded form with accessible Cu sites, to a more ‘closed’ conformation after Cu has bound (42). A similar change may occur in *BsCotA* to facilitate Cu(I) loading by *BsCsp3*.

Figure 5. The proposed role of *BsCsp3* in Cu(I) acquisition by *BsCotA* during sporulation in *B. subtilis*. The transcription factor Spo0A, along with σ^H , initiates sporulation. A septum asymmetrically divides the cell into the forespore and mother cell, with σ^E and σ^F , respectively, activated within these. The mother cell begins engulfment of the forespore and σ^E directs gene expression and initiation of spore coat (purple) formation. The expression of *BsCsp3* and *BsCotA* now begins, promoted by σ^K (see Figure 1f) and coat

assembly continues. We propose that *BsCsp3* acquires Cu(I) during this stage of sporulation, which is transferred to *BsCotA* prior to insertion of the Cu-enzyme into the spore coat.

The expression of both *BsCsp3* and *BsCotA* are regulated by sigma factor K (SigK or σ^K), which is produced after the forespore has been engulfed by the mother cell (Figure 5). Upregulation of the *csp3* gene occurs prior to *cotA* (Figure 1f), thus allowing the storage protein to acquire Cu(I) before production of the enzyme requiring the metal. As *BsCotA* is one of the last proteins to be added to the spore coat (28), Cu(I) transfer from *BsCsp3* could occur during the very late stages of sporulation. Where *BsCsp3* acquires Cu(I) from is not known, but possible sources are Cu-enzymes that are not required by the spore, with the only currently known possibilities being COXs. The finding that *BsCsp3* provides Cu(I) to an enzyme requiring this metal ion suggests a similar role is performed by Csp3s in the cytosol of other bacteria. The novelty of this finding is further highlighted by there being only one currently known example of Cu acquisition by an enzyme from a partner protein in the cytosol. This is from the eukaryotic Cu metallochaperone CCS to the Cu,Zn-superoxide dismutase, which has been studied in considerable detail (1, 34, 35, 37-39, 43, 44).

Added importance to understanding correct metalation of *BsCotA* is provided by the observation that melanin formation interferes with the phagocytosis of pathogenic yeast, and is required to allow survival in macrophages (45). The related pigment produced by Cu-loaded *BsCotA* is important for spore survival (25, 28), and this may include within a host. *Bacillus* spores, and particularly those from *B. cereus*, cause food poisoning and are a common contaminant in a range of foods (46, 47). The development of more effective inactivation approaches requires a better understanding of enzymes that help protect spores such as CotA. This includes establishing how they acquire essential cofactors including Cu ions.

Materials and Methods

Growth curves for WT and $\Delta csp3$ *B. subtilis* at increasing Cu concentrations

WT and $\Delta csp3$ *B. subtilis* 168 strains were obtained from the Bacillus Genetic Stock Centre library. The disrupted *csp3* gene (*yhjQ*) was amplified by PCR using genomic DNA from the $\Delta csp3$ strain with primers that hybridise ~300 bp upstream and downstream of this region (Supplementary Table 3). The resulting fragment was sequenced with primers designed to hybridise ~20 bp from the ends of the PCR product (Supplementary Table 3) and matches that of the erythromycin resistance gene. To test the influence of Cu on WT and $\Delta csp3$ strains, cultures were grown (agitation at 250 rpm) in LB media at 37 °C overnight, diluted (~100-fold) in LB and LB plus Cu(NO₃)₂ (0.5 to 2.0 mM). The absorbance at 600 nm was measured at regular

intervals for up to 12 h, and also after 24 h growth. The Cu concentration in the stock solution used for these studies was regularly determined by atomic absorption spectrometry (AAS), as described previously (15).

The construction of *B. subtilis* strains

To re-insert the *csp3* gene plus its ribosome binding site (RBS) into the $\Delta csp3$ strain, a region including an additional 28 bp at the 5' end was amplified from *B. subtilis* 168 genomic DNA by PCR using primers; rbs_BsCsp3-F and rbs_BsCsp3-R (Supplementary Table 3). The product was cloned into pGEM-T (Promega) and the resulting *rbs_csp3* fragment sub-cloned into pDR111, which possesses the IPTG-inducible *P_{hyerspank}* promoter (48), using HindIII and NheI to generate pDR111_rbs_csp3. To obtain a strain possessing an IPTG-inducible copy of the *csp3* gene (complemented $\Delta csp3$), $\Delta csp3$ *B. subtilis* was transformed with pDR111_rbs_csp3. Selection was achieved using spectinomycin (50 μ g/mL) and successful integration into the chromosomal *amyE* (α -amylase) gene identified by growing on LB agar containing 1% starch and staining with iodine (49). A strain of $\Delta csp3$ *B. subtilis* in which the region of pDR111 lacking the *csp3* gene was integrated into the genome (control $\Delta csp3$) was also generated. The size of the fragment incorporated was confirmed by PCR using the primers pDR111_int_F and pDR111_int_R (Supplementary Table 3).

The production of *B. subtilis* spores

WT, $\Delta csp3$, complemented $\Delta csp3$ and control $\Delta csp3$ strains were grown overnight (agitation at 250 rpm) in 20 mL Difco sporulation media (DSM). Cultures were diluted 50-fold into 200 mL DSM in a single 1 L Erlenmeyer flask and grown until the absorbance at 580 nm reached ~0.5. This was split into four 50 mL cultures, each in a 250 mL Erlenmeyer flask, and 50 μ M Cu(NO₃)₂ and 1 mM IPTG added when required. The cultures were grown (agitation at 250 rpm) at 37 °C for 48 h and absorbance values at 580 nm measured at regular intervals. To purify spores (50) cultures were centrifuged (4 °C) for 10 min at 5,000 g, pellets re-suspended in 50 mM Tris pH 7.2 plus 50 μ g/mL lysozyme and incubated at 37 °C for 1 h. After incubation and further centrifugation (4 °C) for 10 min at 5,000 g, pellets were washed once in sterile MilliQ water and centrifuged. The pellets were re-suspended in 0.05% SDS by vortexing, centrifuged (4 °C) for 10 min at 5,000 g and subsequently washed three times with sterile MilliQ water. The purified spore stocks were verified by PCR (for example, Supplementary Figure 7) using primers listed in Supplementary Table 3 and stored at 4 °C.

BsCotA activity of purified spores

For kinetic measurements of BsCotA activity, purified spores from the WT, $\Delta csp3$, complemented $\Delta csp3$, and control $\Delta csp3$ strains were diluted with MilliQ water to give an

absorbance at 580 nm of ~1.2 (measured accurately), and heated at 65 °C for 1 h prior to use. To determine the colony forming units per mL (CFUs/mL) for this suspension a 5 x 10⁵-fold dilution in LB was plated (100 µL) onto LB agar. The plates were incubated at 37 °C overnight and colonies counted. An aliquot of the heat-treated spore suspension (100 µL) was added to 900 µL of 100 mM citrate-phosphate buffer pH 4.0 plus 0.1-2.4 mM ABTS, and the absorbance at 420 nm ($\epsilon = 3.5 \times 10^4 \text{ M}^{-1}\text{cm}^{-1}$) measured for 5 min at 37 °C (Supplementary Figure 3). A control using 100 µL of buffer was also measured and showed no change in absorbance at 420 nm. The initial velocity (V_0 ; typically reported in units of µM/min/CFUs/mL) was calculated, and plots of V_0 against ABTS concentration (Figure 2 and Supplementary Figure 5) were fit to the Michaelis-Menten equation to determine V_{max} (the maximum rate) and K_M (the Michaelis constant). Comparing V_{max} values calculated based on the absorbance at 580 nm of the heat-treated spore suspension, rather than using CFUs/mL, has no significant influence on the outcome of the study, but generally produces data with larger errors. The reactivity of heat-treated purified spores from WT and $\Delta csp3$ *B. subtilis* with 2.4 mM ABTS was compared to that of the same spore suspension incubated with 250 µM Cu(NO₃)₂ for 30 min at room temperature.

Cloning and purification of BsCotA

The *cotA* gene was amplified from *B. subtilis* genomic DNA using primers CotA_1F and CotA_1R listed in Supplementary Table 3, and cloned into pGEM-T. After removing the NdeI site in the gene by QuickChange site-directed mutagenesis (with primers CotA_2F and CotA_2R, Supplementary Table 3), the product was excised with NdeI and BamHI and re-cloned into pET11a. BsCotA was overexpressed in BL21 *E. coli* (100 µM IPTG) grown at 20 °C for 24 h. The protein was purified using a modified version of a published procedure (26). Cells from 0.5 to 1.0 L of culture were resuspended in 20 mM Tris pH 8.5, sonicated and centrifuged at 40,000 g for 30 min. The supernatant was diluted five-fold in 20 mM Tris pH 8.5 and loaded onto a HiTrap Q HP column (1 mL) equilibrated in the same buffer. Proteins were eluted with a linear NaCl gradient (0-500 mM, total volume 50 mL) and fractions analysed using 18% SDS-PAGE. BsCotA-containing fractions were diluted with 20 mM Tris pH 7.6 and loaded onto a HiTrap SP HP column (5 mL) and eluted with a linear NaCl gradient (0-500 mM, total volume, 200 mL). The BsCotA-containing fractions were heated at 70 °C for 30 min (BsCotA is a highly thermostable enzyme (26)), centrifuged at 40,000 g for 30 min, and the supernatant exchanged into 20 mM HEPES pH 7.5 plus 200 mM NaCl for further purification on a Superdex 75 10/300 GL gel-filtration column. Purified BsCotA had very little Cu or Zn(II) (the latter, a metal ion that commonly binds to Cu proteins when overexpressed in the cytosol of *E. coli*) associated with it when analysed by AAS (12, 13), and showed almost no ABTS oxidation activity. Samples (~3-12 µM, quantified using an ϵ value of 84,739 M⁻¹cm⁻¹ at 280 nm (51)) were incubated with 250 µM Cu(NO₃)₂, thoroughly exchanged and subsequently washed with a low concentration (~10 µM) of

ethylenediaminetetraacetic acid (EDTA). This gave rise to Cu-loaded *BsCotA* with a k_{cat} value of 20-40 s⁻¹ for the oxidation of ABTS, compared to a literature value of 22 s⁻¹ for Cu-enzyme produced using a similar procedure (51).

Purification of *BsCopZ* and sample preparation

BsCopZ is purified with a small amount of Zn(II) bound, as described previously (13). Samples were therefore incubated with >10 equivalents of EDTA for 1 h and exchanged with 20 mM HEPES pH 7.5 plus 200 mM NaCl. The resulting protein had no Zn(II) associated with it and was reduced with dithiothreitol under anaerobic conditions and desalted as described previously (12, 13).

Analysing Cu(I) transfer from recombinant *BsCsp3* and *BsCopZ* to *BsCotA*

BsCsp3 binding ~18 equivalents of Cu(I) was prepared by adding the appropriate amount of a buffered solution of Cu(I) in an anaerobic chamber (Belle Technology, O₂ << 2 ppm) to apo-protein in 20 mM HEPES pH 7.5 plus 200 mM NaCl, quantified using the 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB) assay (12, 13). Fully-reduced *BsCopZ* was also quantified using the DTNB assay and loaded with ~0.8 equivalents of Cu(I) under anaerobic conditions. Cu(I)-*BsCsp3* (~3 μM binding ~53-54 μM Cu(I)) was mixed with ~1.3 μM of as-isolated *BsCotA*, requiring ~5.6 μM Cu(I) to occupy all Cu sites. Cu(I)-*BsCopZ* (~50-53 μM binding ~40-42 μM Cu(I)) was separately added to ~1 μM of as-isolated *BsCotA*, requiring ~4 μM Cu(I) to fill all Cu sites. Mixtures were incubated at room temperature in the anaerobic chamber for up to 48 h. A similar concentration of *BsCotA* was also incubated anaerobically with Cu(I), prepared as described above, a Cu(NO₃)₂ solution (final concentrations of ~56-58 μM), buffer (20 mM HEPES pH 7.5 plus 200 mM NaCl) alone, and also with Cu(NO₃)₂ in air. To measure activity, 10 μL of each mixture was added to 990 μL of aerated 100 mM citrate-phosphate buffer pH 4.0 plus 2.4 mM ABTS, and the absorbance at 420 nm measured for up to 5 min at 37 °C (Figure 3). The removal of Cu(I) by BCS (~2.5 mM) was analysed for Cu(I)-*BsCsp3* (~0.8-1.2 μM plus ~18 equivalents of Cu(I)) samples used for the transfer experiments, both in the absence (folded *BsCsp3*) and presence (unfolding conditions) of guanidine-HCl (6.64 M) (12, 13). The absorbance increase at 483 nm due to formation of [Cu(BCS)₂]³⁻ ($\epsilon = 12,500 \text{ M}^{-1}\text{cm}^{-1}$) (31) was measured over time at 22 °C in 20 mM HEPES pH 7.5 plus 200 mM NaCl (Supplementary Figure 6).

Acknowledgments

We thank BBSRC for funding (grant BB/K008439/1 to C.D.). We are grateful to Newcastle University for an Overseas Research Scholarship (ORS) award to J.L.. We also thank Prof.

Heath Murray for help with generating *B. subtilis* mutant strains, Dr. Mark Harrison for cloning the *cotA* gene and Dr. Gianpiero Landolfi for purifying BsCopZ.

Author contributions

C.D. and J.L. conceived the project and designed the experiments. J.L. performed the experiments and analysed data with help from C.D. C.D. wrote the manuscript with help from J.L.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper.

References

1. Rae, T. D., Schmidt, P. J., Pufahl, R. A., Culotta, V. C. & O'Halloran, T. V. Undetectable intracellular free copper: The requirement for a copper chaperone for superoxide dismutase. *Science* **284**, 805–808 (1999) doi: 10.1126/science.284.5415.805.
2. Changela, A., Chen, K., Xue, Y., Holschen, J., O'Halloran, T. V. & Mondragón, A. Molecular basis of metal-ion selectivity and zeptomolar sensitivity by CueR. *Science* **301**, 1383–1387 (2003) doi: 10.1126/science.1085950.
3. Festa, R. A. & Thiele, D. J. Copper: an essential metal in biology. *Curr. Biol.* **21**, R877–R883 (2011) doi: 10.1016/j.cub.2011.09.040.
4. Marckmann, D., Trasnea, P. I., Schimpf, J., Winterstein, C., Andrei, A., Schmoller, S., Blaby-Haas, C. E., Friedrich, T., Daldal, F. & Koch, H. G. The *cbb₃*-type cytochrome oxidase assembly factor CcoG is a widely distributed cupric reductase. *Proc. Natl. Acad. Sci. USA* **116**, 21166–21175 (2019) doi: 10.1073/pnas.1913803116.
5. Hassett, R. & Kosman, D. J. Evidence for Cu(II) reduction as a component of copper uptake by *Saccharomyces cerevisiae*. *J. Biol. Chem.* **270**, 128–134 (1995) doi: 10.1074/jbc.270.1.128.
6. Rensing, C., Fan, B., Sharma, R., Mitra, B. & Rosen, B. P. CopA: An *Escherichia coli* Cu(I)-translocating P-type ATPase. *Proc. Natl. Acad. Sci. USA* **97**, 652–656 (2000) doi: 10.1073/pnas.97.2.652.
7. Solioz, M., Abicht, H. K., Mermoud, M. & Mancini, S. Response of Gram-positive bacteria to copper stress. *J. Biol. Inorg. Chem.* **15**, 3–14 (2010) doi: 10.1007/s00775-009-0588-3.
8. Rensing, C. & McDevitt S. F. The copper metallome in prokaryotic cells. *Met. Ions Life Sci.* **12**, 417–450 (2013) doi: 10.1007/978-94-007-5561-1_12.

9. Meydan, S., Klepacki, D., Karthikeyan, S., Margus, T., Thomas, P., Jones, J. E., Khan, Y., Briggs, J., Dinman, J. D., Vázquez-Laslop, N. & Mankin, A. S. Programmed ribosomal frameshifting generates a copper transporter and a copper chaperone from the same gene. *Mol. Cell.* **65**, 207–219 (2017) doi: 10.1016/j.molcel.2016.12.008.
10. Macomber, L. & Imlay, J. A. The iron-sulfur clusters of dehydratases are primary intracellular targets of copper toxicity. *Proc. Natl. Acad. Sci. USA* **106**, 8344–8349 (2009) doi: 10.1073/pnas.0812808106.
11. Ridge, P. G., Zhang, Y. & Gladyshev, V. N. Comparative genomic analyses of copper transporters and cuproproteomes reveal evolutionary dynamics of copper utilization and its link to oxygen. *PLoS One* **3**, e1378 (2008) doi: 10.1371/journal.pone.0001378.
12. Vita, N., Platsaki, S., Baslé, A., Allen, S. J., Paterson, N. G., Crombie, A. T., Murrell, J. C., Waldron, K. J. & Dennison, C. A four-helix bundle stores copper for methane oxidation. *Nature* **525**, 140–143 (2015) doi: 10.1038/nature14854.
13. Vita, N., Landolfi, G., Baslé, A., Platsaki, S., Lee, J., Waldron, K. J. & Dennison, C. Bacterial cytosolic proteins with a high capacity for Cu(I) that protects against copper toxicity. *Sci. Rep.* **6**, 39065 (2016) doi: 10.1038/srep39065.
14. Dennison, C., David, S. & Lee, J. Bacterial copper storage proteins. *J. Biol. Chem.* **293**, 4616–4627 (2018) doi: 10.1074/jbc.TM117.000180.
15. Lee, J. & Dennison, C. Cytosolic copper binding by a bacterial storage protein and interplay with copper efflux. *Int. J. Mol. Sci.* **20**, 4144 (2019) doi: 10.3390/ijms20174144.
16. Gu, W. & Semrau, J. D. Copper and cerium-regulated gene expression in *Methylosinus trichosporium* OB3b. *Appl. Microbiol. Biotechnol.* **101**, 8499–8516 (2017) doi: 10.1007/s00253-017-8572-2.
17. DiSpirito, A. A., Semrau, J. D., Murrell, J. C., Gallagher, W. H., Dennison C. & Vuilleumier, S. Methanobactin and the link between copper and bacterial methane oxidation. *Microbiol. Mol. Biol. Rev.* **80**, 387–409 (2016) doi: 10.1128/MMBR.00058-15.
18. Radford, D. S., Kihlken, M. A., Borrelly, G. P. M., Harwood, C. R., Le Brun, N. E. & Cavet, J. S. CopZ from *Bacillus subtilis* interacts in vivo with a copper exporting CPx-type ATPase CopA. *FEMS Microbiol. Lett.* **220**, 105–12 (2003) doi: 10.1016/S0378-1097(03)00095-8.
19. Smaldone, G. T. & Helmann, J. D. CsoR regulates the copper efflux operon *copZA* in *Bacillus subtilis*. *Microbiology* **153**, 4123–4128 (2007) doi: 10.1099/mic.0.2007/011742-0.
20. Chillappagari, S., Miethke, M., Trip, H., Kuipers, O. P. & Marahiel, M. M. Copper acquisition is mediated by YcnJ and regulated by YcnK and CsoR in *Bacillus subtilis*. *J. Bacteriol.* **191**, 2362–2370 (2009) doi: 10.1128/JB.01616-08.
21. Ma, Z., Cowart, D. M., Scott, R. A. & Giedroc, D. P. Molecular insights into the metal selectivity of the copper(I)-sensing repressor CsoR from *Bacillus subtilis*. *Biochemistry* **48**, 3325–3334 (2009) doi: 10.1021/bi900115w.

22. Hirooka, K., Eda Hiro, T., Kimura, K. & Fujita, Y. Direct and indirect regulation of the *ycnKJl* operon involved in copper uptake through two transcriptional repressors, YcnK and CsoR, in *Bacillus subtilis*. *J. Bacteriol.* **194**, 5675–5687 (2012) doi: 10.1128/JB.00919-12.
23. Damle, M. S., Singh, A. N., Peters, S. C., Szalai, V. A. & Fisher, O. S. The YcnI protein from *Bacillus subtilis* contains a copper-binding domain. *J. Biol. Chem.* **297**, 101078 (2021) doi: 10.1016/j.jbc.2021.101078.
24. Lauraeus, M., Haltia, T., Saraste, M. & Wikström, M. *Bacillus subtilis* expresses two kinds of haem-A-containing terminal oxidases. *Eur. J. Biochem.* **197**, 699–705 (1991) doi: 10.1111/j.1432-1033.1991.tb15961.x.
25. Hullo, M. F., Moszer, I., Danchin, A. & Martin-Verstraete, I. CotA of *Bacillus subtilis* is a copper-dependent laccase. *J. Bacteriol.* **183**, 5426–5430 (2001) doi: 10.1128/JB.183.18.5426-5430.2001.
26. Martins, L. O., Soares, C. M., Pereira, M. M., Teixeira, M., Costa, T., Jones, G. H., Henriques, A. O. Molecular and biochemical characterisation of a highly stable bacterial laccase that occurs as a structural component of the *Bacillus subtilis* endospore coat. *J. Biol. Chem.* **277**, 18849–18859 (2002) doi: 10.1074/jbc.M200827200.
27. Enguita, F. J., Martins, L. O., Henriques, A. O. & Carrondo, M. A. Crystal structure of a bacterial endospore coat component. *J. Biol. Chem.* **278**, 19416–19425 (2003) doi: 10.1074/jbc.M301251200.
28. McKenney, P. T., Driks, A. & Eichenberger, P. The *Bacillus subtilis* endospore: assembly and functions of the multilayered coat. *Nat. Rev. Microbiol.* **11**, 33–44 (2013) doi: 10.1038/nrmicro2921.
29. Nicolas, P. et al. Condition-dependent transcriptome reveals high-level regulatory architecture in *Bacillus subtilis*. *Science* **335**, 1103–1106 (2012) doi: 10.1126/science.1206848.
30. Vavrová, L., Muchová, K. & Barák, I. Comparison of different *Bacillus subtilis* expression systems. *Res. Microbiol.* **161**, 791–797 (2010) doi: 10.1016/j.resmic.2010.09.004.
31. Badarau, A. & Dennison, C. Copper trafficking mechanism of CXXC-containing domains: Insight from the pH dependence of their Cu(I) affinities. *J. Am. Chem. Soc.* **133**, 2983–2988 (2011) doi: 10.1021/ja1091547.
32. Gunne, M., Al-Sultani, D. & Urlacher, V. Enhancement of copper content and specific activity of CotA laccase from *Bacillus licheniformis* by coexpression with CopZ copper chaperone in *E. coli*. *J. Biotechnol.* **168**, 252–255 (2013) doi: 10.1016/j.jbiotec.2013.06.011.
33. Pufahl, R. A., Singer, C. P., Peariso, K. L., Lin, S. J., Schmidt, P. J., Fahrni, C. J., Culotta, V. C., Penner-Hahn, J. E. & O'Halloran, T. V. Metal ion chaperone function of the soluble Cu(I) receptor Atx1. *Science* **278**, 853–856 (1997) doi: 10.1126/science.278.5339.853.
34. Schmidt, P. J., Rae, T. D., Pufahl, R. A., Hamma, T., Strain, J., O'Halloran, T. V. & Culotta, V. C. Multiple protein domains contribute to the action of the copper chaperone for

- 497 superoxide dismutase. *J. Biol. Chem.* **274**, 23719–23725 (1999) doi:
498 10.1074/jbc.274.34.23719.
- 499 35. Lamb, A. L., Torres, A. S., O'Halloran, T. V. & Rosenzweig, A. C. Heterodimeric structure of
500 superoxide dismutase in complex with its metallochaperone. *Nat. Struct. Biol.* **8**, 751–755
501 (2001) doi: 10.1038/nsb0901-751.
- 502 36. Banci, L., Bertini, I., Cantini, F., Felli, I. C., Gonnelli, L., Hadjiladis, N., Pierattelli, R.,
503 Rosato, A. & Voulgaris P. The Atx1-Ccc2 complex is a metal-mediated protein-protein
504 interaction. *Nat. Chem. Biol.* **2**, 367–368 (2006) doi: 10.1038/nchembio797.
- 505 37. Banci, L., Bertini, I., Ciofi-Baffoni, S., Kozyreva, T., Zovo, K. & Palumaa, P. Affinity gradients
506 drive copper to cellular destinations. *Nature* **465**, 645–648 (2010) doi:
507 10.1038/nature09018.
- 508 38. Banci, L., Bertini, I., Cantini, F., Kozyreva, T., Massagni, C., Palumaa, P., Rubino, J. T. &
509 Zovo, K. Human superoxide dismutase 1 (hSOD1) maturation through interaction with
510 human copper chaperone for SOD1 (hCCS). *Proc. Natl. Acad. Sci. USA* **109**, 13555–13560
511 (2011) doi: 10.1073/pnas.1207493109.
- 512 39. Sala, F. A., Wright, G. S. A., Antonyuk, S. V., Garratt, R. C. & Hasnain, S. S. Molecular
513 recognition and maturation of SOD1 by its evolutionary destabilised cognate chaperone
514 hCCS. *PLoS Biol.* **17**, e3000141 (2019) doi: 10.1371/journal.pbio.3000141.
- 515 40. Badarau, A. & Dennison, C. Thermodynamics of copper and zinc distribution in the
516 cyanobacterium *Synechocystis* PCC 6803. *Proc. Natl. Acad. Sci. USA* **108**, 13007–13012
517 (2011) doi: 10.1073/pnas.1101448108.
- 518 41. North, M. L. & Wilcox, D. E. Shift from entropic Cu²⁺ binding to enthalpic Cu¹⁺ binding
519 determines the reduction thermodynamics of blue copper proteins. *J. Am. Chem. Soc.* **141**,
520 14329–14339 (2019) doi: 10.1021/jacs.9b06836.
- 521 42. Strolle, P., Hou, B. & Brüser, T. The Tat substrate CueO is transported in an incomplete
522 folding state. *J. Biol. Chem.* **291**, 13520–13528 (2016) doi: 10.1074/jbc.M116.729103.
- 523 43. Culotta, V. C., Klomp, L. W., Strain, J., Casareno, R. L., Krems, B. & Gitlin, J. D. The copper
524 chaperone for superoxide dismutase. *J. Biol. Chem.* **272**, 23469–23472 (1997) doi:
525 10.1074/jbc.272.38.23469.
- 526 44. Wong, P. C., Waggoner, D., Subramaniam, J. R., Tessarollo, L., Bartnikas, T. B., Culotta, V.
527 C., Price, D. L., Rothstein, J. & Gitlin J. D. Copper chaperone for superoxide dismutase is
528 essential to active mammalian Cu/Zn superoxide dismutase. *Proc. Natl. Acad. Sci. USA* **97**,
529 2886–2891 (2000) doi: 10.1073/pnas.040461197.
- 530 45. Eisenman, H. C. & Casadevall, A. Synthesis and assembly of fungal melanin. *Appl.*
531 *Microbiol. Biotechnol.* **93**, 931–940 (2012) doi: 10.1007/s00253-011-3777-2.
- 532 46. Soni, A., Oey, I., Silcock, P. & Bremer, P. *Bacillus* spores in the food industry: a review on
533 resistance and response to novel inactivation technologies. *Comp. Rev. Food Sci. Food*
534 *Saf.* **15**, 1139–1148 (2016) doi: 10.1111/1541-4337.12231.

47. Jessberger, N., Dietrich, R., Granum, P. E. & Märtlbauer, E. The *Bacillus cereus* food infection as multifactorial process. *Toxins*, **12**, 701 (2020) doi: 10.3390/toxins12110701.
48. Quisel, J. D., Burkholder, W. F. & Grossman, A. D. *In vivo* effects of sporulation kinases on mutant Spo0A proteins in *Bacillus subtilis*. *J. Bacteriol.* **183**, 6573–6578 (2001) doi: 10.1128/JB.183.22.6573-6578.2001.
49. Engman, J., Rogstam, A., Frees, D., Ingmer, H. & von Wachenfeldt, C. The YjbH adaptor protein enhances proteolysis of the transcriptional regulator Spx in *Staphylococcus aureus*. *J. Bacteriol.* **194**, 1186–1194 (2012) doi: 10.1128/JB.06414-11.
50. Tavares, M. B., Souza, R. D., Luiz, W. B., Cavalcante, R. C. M., Casaroli, C., Martins, E. G., Ferreira, R. C. C. & Ferreira, L. C. S. *Bacillus subtilis* endospores at high purity and recovery yields: optimization of growth conditions and purification method. *Curr. Microbiol.* **66**, 279–285 (2013) doi: 10.1007/s00284-012-0269-2.
51. Durão, P., Chen, Z., Fernandes, A. T., Hilderbrandt, P., Murgida, D. H. Todorovic, S., Pereira, M. M., Melo, E. P. & Martins, L. O. Copper incorporation into recombinant CotA laccase from *Bacillus subtilis*: characterisation of fully copper loaded enzymes. *J. Biol. Inorg. Chem.* **13**, 183–193 (2008) doi: 10.1007/s00775-007-0312-0.

SUPPLEMENTARY INFORMATION

CONTENTS:

Supplementary Figure 1. The influence of Cu on the growth of WT and $\Delta csp3$ *B. subtilis*.

Supplementary Figure 2. The influence of Cu levels in media on Cu accumulation by WT and $\Delta csp3$ *B. subtilis*.

Supplementary Figure 3 The influence of deleting the *csp3* gene and Cu levels on *BsCotA* activity in *B. subtilis* strains.

Supplementary Figure 4. Testing spores for enhanced *BsCotA* activity in the presence of Cu.

Supplementary Figure 5. The influence of Cu levels and IPTG on *BsCotA* activity in the control $\Delta csp3$ *B. subtilis* strain.

Supplementary Figure 6. The removal of Cu(I) from *BsCsp3* by the high-affinity ligand BCS.

Supplementary Figure 7. PCR analysis of purified spore stocks from the *B. subtilis* strains used in this study.

Supplementary Table 1. The influence of Cu concentration on the *BsCotA* activity of purified spores from WT and $\Delta csp3$ *B. subtilis*.

Supplementary Table 2. The removal of Cu(I) from *BsCsp3* by BCS over time.

Supplementary Table 3. Primers used in this study.

Supplementary Figure 1. The influence of Cu on the growth of WT and $\Delta csp3$ *B. subtilis*.

Growth (37 °C) of WT (circles) and $\Delta csp3$ (squares) *B. subtilis* in LB media plus 0 (a), 0.5 (b), 1.0 (c), 1.5 (d), and 2.0 (e) mM added $\text{Cu}(\text{NO}_3)_2$. The data obtained at 12 h is compared in (f), and in all cases averages and standard deviations from three independent growth experiments are shown. These results are similar to those we reported previously (13), albeit the influence on growth starts to be observed at a Cu(II) concentration in the media that is ~0.5 mM lower in this work (we have quantified Cu levels in the stock solutions used herein by AAS, but did not previously (13)). The absorbance of these cultures were also measured at 24 h, and at up to 1.0 mM added $\text{Cu}(\text{NO}_3)_2$ a significant decrease was observed compared to the value at 12 h for both strains (consistent with previous data (13)). At higher added $\text{Cu}(\text{NO}_3)_2$ concentrations, the growth data showed no consistent pattern beyond 12 h. This differs to the enhanced cell death that was seen previously for $\Delta csp3$ *B. subtilis* after 12 h growth at reported media Cu concentrations of 1.5-2.0 mM (13).

Supplementary Figure 4. Testing purified spores for enhanced *BsCotA* activity after incubation with Cu(II). The influence of incubating spores in 250 μ M Cu(NO₃)₂ from WT and Δ csp3 *B. subtilis* grown either in the absence or presence (50 μ M) of Cu(NO₃)₂ on their *BsCotA* activity. The activity units are reported based on both CFU values (a) and the absorbance (Abs) at 580 nm (b) to quantify spores. Average values and standard deviations from three independent experiments on the same spores used for the data in Figure 2a,b are shown, and the values are listed in Supplementary Table 1. The reactions with ABTS were measured in 100 mM citrate-phosphate buffer pH 4.0.

Supplementary Figure 5. The influence of Cu levels and IPTG on *BsCotA* activity in the control $\Delta csp3$ *B. subtilis* strain. Michaelis-Menten plots of *BsCotA* activity for heated purified spores from the control $\Delta csp3$ strain. (a) The analysis of spores produced in media plus 0 (open symbols) and 50 (black filled symbols) μM added $\text{Cu}(\text{NO}_3)_2$ with no added IPTG. (b) Data plus 1 mM IPTG and 0 (magenta) and 50 (cyan) μM $\text{Cu}(\text{NO}_3)_2$. Averages and standard deviations from kinetic measurements in 100 mM citrate-phosphate buffer pH 4.0 using three different sets of spores are shown.

Supplementary Figure 6. The removal of Cu(I) from *BsCsp3* by the high-affinity ligand BCS. A plot of $[\text{Cu}(\text{BCS})_2]^{3-}$ concentration against time for *BsCsp3* ($1.08 \mu\text{M}$) plus 18.0 equivalents of Cu(I) mixed with 2.5 mM BCS in 20 mM HEPES pH 7.5 plus 200 mM NaCl (red line). The data at 0.5, 1, 2, 3, 4, 6, 17 and 24 h, which correspond to times at which *BsCotA* activity was measured (Figure 4), are indicated by red circles. Average percentage removal compared to that for the unfolded samples, and the standard deviations, from three independent experiments are shown in Supplementary Table 2. The black line indicates the same experiment but with 6.64 M guanidine-HCl present in the buffer. This helps unfold the protein and removal is much faster giving the end point for the reaction (the value shown was obtained after incubation for 2 h).

Supplementary Figure 7. PCR analyses of purified spore stocks from the *B. subtilis* strains used in this study. (a) The rbs_BsCsp3-F and rbs_BsCsp3-R primers (Supplementary Table 3) give a product of 373 bp for the WT strain, whilst BsCsp3+300-F and BsCsp3+300-R give fragments of 967 and 1663 bp for WT and $\Delta csp3$, respectively. (b) Re-introduction of the *csp3* gene is confirmed in the complemented strain using the rbs_BsCsp3-F and rbs_BsCsp3-F primers (as well as testing with BsCsp3+300-F and BsCsp3+300-R), and the same primers are used to analyse the control $\Delta csp3$ strain (c). Analysis of spores used for the three independent experiments on all four strains shown in Figure 2 and Supplementary Figure 5 gave the same results.

Supplementary Table 1. The influence of Cu concentration on the *BsCotA* activity of purified spores from WT and $\Delta csp3$ *B. subtilis* ^a

<i>B. subtilis</i> strain	Cu(II) in media ^b	Spores incubated with	
		250 μM Cu(II)	No Cu(II)
<i>BsCotA</i> activity nmol/min/CFUs/mL (x 10 ⁻⁸) ^c			
WT	-	11.6 ± 4.29	5.02 ± 1.75
WT	+	13.6 ± 1.37	8.38 ± 3.15
Δ <i>csp3</i>	-	11.0 ± 2.20	5.71 ± 3.32
Δ <i>csp3</i>	+	4.84 ± 1.92	4.71 ± 0.85
<i>BsCotA</i> activity nmol/min/Abs at 580 nm ^d			
WT	-	11.0 ± 1.90	4.33 ± 1.15
WT	+	8.07 ± 1.76	8.00 ± 1.55
Δ <i>csp3</i>	-	6.53 ± 1.40	3.64 ± 2.07
Δ <i>csp3</i>	+	2.87 ± 0.11	3.52 ± 0.57

^a Average values and standard deviations from the three independently grown sets of spores, used for the data in Figure 2a,b are shown (activity measurements measured in 100 mM citrate-phosphate buffer pH 4.0). ^b The minus sign indicates that Cu was not added to media during sporulation, whilst the plus sign indicates that the DSM was supplemented with 50 μ M Cu(NO₃)₂. ^c Values used for Supplementary Figure 4a with the numbers of spores calculated using CFUs. ^d Data shown in Supplementary Figure 4b and calculated by quantifying spores in suspensions using the absorbance (Abs) at 580 nm, to enable direct comparison with a previous study (26). The data for WT *B. subtilis* are very similar to those in the earlier work (26), even though in that case 250 μ M Cu (50 μ M here) was added to the DSM during sporulation. The only discrepancy is that previously (26), WT spores grown in the presence of Cu showed an approximate 2-fold decrease in activity upon incubation with Cu(II). This observation is difficult to explain, but the data reported in ref. 26 appears to be from a single replicate.

Supplementary Table 2. The removal of Cu(I) from *BsCsp3* by BCS over time ^{a,b}

Time (h)	% Cu(I) removal ^c
0.5	12.9 $\pm$ 0.54
1	18.0 $\pm$ 0.71
2	24.7 $\pm$ 1.33
3	28.3 $\pm$ 1.23
4	31.5 $\pm$ 1.39
6	36.6 $\pm$ 1.50
17	52.9 $\pm$ 2.31
24	60.7 $\pm$ 2.35 ^d

^a Shown are the average percentage removal of Cu(I) from *BsCsp3* by BCS over time (and standard deviations) from three independent experiments. ^b The amount of Cu(I) bound to the protein was determined by mixing the Cu(I)-*BsCsp3* sample with 2.5 mM BCS in the presence of 6.64 M guanidine-HCl in 20 mM HEPES pH 7.5 plus 200 mM NaCl, which unfolds the protein giving the maximum possible [Cu(BCS)₂]³⁻ concentration (the value used is after incubation for 2 h). ^c Percentage Cu(I) removal is determined using [Cu(BCS)₂]³⁻ concentration/maximum [Cu(BCS)₂]³⁻ concentration $\times$ 100. ^d On one occasion the experiment was analysed up to 48 h with 70% Cu(I) removal observed.

660 **Supplementary Table 3. Primers used in this study**

Primers	Sequence
BsCsp3+300-F ^a	5'-CATTCATGACAGTGCAGCG-3'
BsCsp3+300-R ^a	5'-CACAAGAGGACTGGACGC-3'
BsCsp3_seq-F ^b	5'-CCGACAGCAGCAAATGCAGAAAACCA-3'
BsCsp3_seq-R ^b	5'-AAACCGTCGACTTTTACTTGCG-3'
rbs_BsCsp3-F ^c	5'-GGAGGACGCA AAGCTT GCCGTGAAACATAAAACC-3'
rbs_BsCsp3-R ^c	5'-GGAGGACGCG GCTAGC <u>TTAC</u> GCTGCCATGCTGCGGC-3'
BsCsp3-F ^d	5'-GCGCATATGGAGCAATATTCTGAGGC-3'
BsCsp3-R ^d	5'-GCGCCATGGTTACGCTGCCATGCTGCGGC-3'
pDR111_int_F ^e	5'-GTGAACGCTCTCCTGAGTAG-3'
pDR111_int_R ^e	5'-GTCGGCTGAAAGATCGTAC-3'
CotA-1F ^f	5'-GGT CATATG ACACTTGAAAAATTTGTGGATGC-3'
CotA-1R ^f	5'-GGT GGATCC <u>TTAT</u> TTTATGGGGATCAGTTATATCC-3'
CotA-2F ^g	5'-CATTGACTTCACAG CGTAT GAAGGAGAATCG-3'
CotA-2R ^g	5'-CGATTCTCCTT CATAC GCTGTGAAGTCAATG-3'

661 ^a Forward and reverse primers designed to hybridise ~300 bp upstream and downstream of the
662 *csp3* gene giving PCR products of 1663 and 967 bp for $\Delta csp3$ and WT *B. subtilis*. ^b Primers used
663 to sequence the PCR product obtained from the $\Delta csp3$ strain using primers BsCsp3+300-F and
664 BsCsp3+300-R. ^c Forward (HindIII site in bold) and reverse (NheI site in bold, stop codon
665 underlined) primers designed to amplify the *csp3* gene plus 28 bp at the 5' end to include its own
666 RBS for cloning into pDR111. ^d Forward and reverse primers that amplify the *csp3* gene giving a
667 PCR fragment of 345 bp for WT *B. subtilis*. ^e Primers that hybridise 294 bp upstream and 118 bp
668 downstream of the region of pDR111 integrated into the *amyE* gene in the *B. subtilis* genome.
669 For the complemented $\Delta csp3$ strain these give a 831 bp fragment, whilst for the control $\Delta csp3$
670 strain the fragment size is 466 bp. ^f Forward (NdeI site in bold) and reverse (BamHI site in bold,
671 stop codon underlined) primers designed to amplify the *cotA* gene. ^g Forward and reverse
672 primers used to mutate the NdeI site (from CATATG to CATACG, highlighted in grey with the
673 site mutation in bold) within the *cotA* gene.