Short title: Development and analysis of tobacco HY5 mutant

Corresponding author:

Prabodh Kumar Trivedi, CSIR-National Botanical Research Institute (CSIR-NBRI), Rana Pratap Marg, Lucknow-226001, INDIA; **Telephone number:** +91 522 2297958; **e-mail address:** prabodht@hotmail.com; prabodht@nbri.res.in

Research area: Biochemistry and metabolism

Tobacco HY5, NtHY5, positively regulates flavonoid biosynthesis and enhances salt stress tolerance

Deeksha Singh^{1,2}, Hiteshwari Sinha^{1,2}, Nivedita Singh^{1,§}, Shambhavi Dwivedi³ & Prabodh Kumar Trivedi^{1,2,3}

¹CSIR-National Botanical Research Institute, Council of Scientific and Industrial Research (CSIR-NBRI), Rana Pratap Marg, Lucknow-226001, INDIA

²Academy of Scientific and Innovative Research (AcSIR), Ghaziabad- 201002, India

³Central Institute of Medicinal and Aromatic Plants (CSIR-CIMAP), Lucknow, India.

&Present address:

Department of Botany, University of Lucknow, Lucknow-226007, India.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (<https://academic.oup.com/plphys/pages/General-Instructions>) is Prabodh K. Trivedi.

One sentence summary: Analysis of *NtHY5* overexpressing and mutant plants suggests role of NtHY5 in flavonoid biosynthesis and salt stress tolerance.

Responsible author for distribution of materials: Prabodh K. Trivedi

Author contributions: P.K.T. designed and supervised this study. D.S. designed and performed most of the experiments; HS, NS and SD provided help in conducting various experiments; P.K.T. analyzed the data; D.S., P.K.T. wrote the article; All authors read, contributed, and approved the article.

Funding information: This research was supported by the Council of Scientific and Industrial Research (CSIR), New Delhi, in the form of NCP project no. MLP026. P.K.T. also acknowledges the Department of Biotechnology, New Delhi, for financial support in the form of projects on pathway engineering, genome editing and a TATA Innovation Fellowship. DS

acknowledge University Grant Commission, New Delhi for Senior and Junior Research Fellowship respectively.

ABSTRACT

Plants have evolved complex signaling networks to regulate their growth and development. Some of these signaling components also play a crucial role in secondary metabolite biosynthesis. Among the signaling components identified to date, ELONGATED HYPOCOTYL 5 (HY5), a bZIP family transcription factor is the most investigated and known as the center of transcriptional network hub. However, HY5 has not been characterized from plants known to synthesize important secondary metabolites. In this study, based on homology search and phylogenetic analysis, HY5 has been identified from *Nicotiana tobaccum*, and characterized for its role in secondary plant product biosynthesis and stress response through developing overexpressing lines and CRISPR/Cas9-based knockout mutant plants. NtHY5 was able to complement the *Arabidopsis thaliana hy5* mutant at molecular, morphological and biochemical levels. Overexpression of *NtHY5* in tobacco led to the up-regulation of the phenylpropanoid pathway genes and enhanced the flavonoid content, whereas mutant plants had the opposite effect. Electrophoretic Mobility Shift Assay (EMSA) suggested that NtHY5 interacts with the promoter of *NtMYB12*, a transcription factor known to regulate flavonoid biosynthesis. In addition, *NtHY5* enhanced the abiotic stress tolerance as evident by the salt tolerance ability of HY5 overexpressing lines by diminishing the ROS accumulation after salt treatment. These data provide credible evidence about the potential role of NtHY5 in light-mediated flavonoid biosynthesis, plant growth and abiotic stress tolerance in tobacco. The photomorphogenic mutant, *Nthy5*, developed in this study, will help in elucidating the role of the HY5 in different biological processes in tobacco.

Keywords: *Arabidopsis thaliana*, flavonoid synthesis, HY5, light-dependent regulation, MYB transcription factor, *Nicotiana tobaccum*, salt stress

INTRODUCTION

Major plant developmental processes such as flowering, seed development, organ development, seed germination and seedling de-etiolation are governed by light (Chory et al., 2010; Vanhaelewyn et al., 2016; Gangappa et al., 2016; Bhatnagar et al., 2020). Many transcription factors of various families, including bHLH, bZIP, MYB, GATA and Zinc-finger get activated after light perception by photoreceptors (Oyama et al., 1997; Chory et al., 2010; Yogev et al., 2020). Constitutive nuclear-localized ELONGATED HYPOCOTYL 5 (HY5), a bZIP family transcription factor, acts downstream to the photoreceptors and plays a central role in seedling development via integrating light and phytohormone signaling pathways (Koornneef et al., 1980; Osterlund et al., 2000; Ulm et al., 2012; Li et al., 2013). As established by ChIP-chip experiments, HY5 regulates the expression of one-third of Arabidopsis genes and binds to promoters of more than 3000 genes directly (Zhang et al., 2011a). Thus, it acts as a master regulator by binding to *cis*-regulatory elements.

HY5 promotes photomorphogenesis in red, far-red, blue and UV-B light and acts downstream of cryptochromes, phytochromes and UVR8 (Jiang et al., 2012; Zheng et al., 2013; Yadav et al., 2020). The *hy5* mutant shows a long-hypocotyl phenotype in light (Koornneef et al., 1980) with an increased number of lateral roots, modified gravitropic and touch responses in roots (Oyama et al., 1997). The content of flavonoids, anthocyanin and chlorophyll was also reduced substantially in *hy5* seedlings compared to wild type in spite of the fact that seedlings of *hy5* mutant had larger cotyledons compared to wild type (Oyama et al., 1997; Sibout et al., 2006; Bhatia et al., 2018; Bhatia et al., 2021). In addition to plant growth and development, there are reports suggesting an involvement of HY5 in the regulation of terpenoid biosynthesis (Michael et al., 2020) and the expression and accumulation of miRNA-encoded peptide, miPEP858a (Sharma et al., 2021).

Flavonoids are low molecular weight polyphenolic secondary metabolites with a C₆-C₃-C₆ backbone. Flavonoids with more than 9000 distinct structures are synthesized via the phenylpropanoid/flavonoid pathway (Falcon Ferreyra et al., 2012). These molecules are divided into several classes, including chalcones, flavanones, dihydroflavonols, flavonols, flavones, isoflavonoids and anthocyanins on the basis of structural diversity (Ververidis et al., 2007; Saito et al., 2013). Sequential action of enzymes like chalcone isomerase (CHI), flavanone 3-

hydroxylase (F3H), flavonoid 3' hydroxylase (F3'H) and flavonol synthase (FLS) leads to the formation of a aglycone flavonoid backbone. Furthermore, various modifications like glycosylation, methylation and acylation lead to huge chemical diversity in flavonoids (Ferrer et al., 2008). Flavonoids have received tremendous attention in the past few years because of their ubiquitous presence, diversity, and diverse roles in plants as well as in the pharmaceutical industry (Cai et al., 2010; Schulz et al., 2015). Flavonoids are known to have diverse roles in plants ranging from tissue and flower pigmentation to provide protection against various kinds of abiotic and biotic stresses (Winkel-Shirley 2002; Misra et al., 2010; Schulz et al., 2016). Overproduction of rutin was found to be responsible for insecticidal activity of AtMYB12 overexpressing transgenic tobacco plants and callus (Misra et al., 2010; Pandey et al., 2012).

Past studies in *Arabidopsis* demonstrated that R2R3 MYBs, bHLHs and WD40 regulatory proteins form the MBW ternary complex that coordinately regulates the flavonoid biosynthesis at the transcriptional level (Hichri et al., 2011). The early biosynthetic genes of the flavonoid biosynthetic pathway are transcriptionally regulated by a set of three closely related R2R3 MYBs; AtMYB12/PFG1, AtMYB11/ PFG2 and AtMYB111/ PFG3 and they belong to “PRODUCTION OF FLAVONOL GLYCOSIDES” family (Stracke et al., 2007; Stracke et al., 2010a). *AtMYB12*, *AtMYB11*, *AtMYB111* and *NtMYB12* are reported to regulate the flavonoid biosynthesis by activating the phenylpropanoid pathway genes (Misra et al., 2010; Pandey et al., 2015; Fu et al., 2016). Studies also suggest that the expression of these MYB transcription factors is regulated by miR858 (Sharma et al., 2016) and miRNA-encoded peptide miPEP858a (Sharma et al., 2020). The light regulatory mechanism of the flavonoid pathway in tobacco is not that much explored, but there are few studies suggesting that light plays an important role in flavonoid biosynthesis in tobacco (Fu et al., 2016; Pandey et al., 2015).

Tobacco is one of the important cash crops worldwide and its leaves contain various secondary plant products, including flavonoids. Therefore, tobacco is a suitable model plant to explore regulatory mechanisms related to various secondary plant products. The extensive characterization of HY5 and flavonoid biosynthesis has been done in *Arabidopsis*, Rice, *Pisum sativum*, *Physcomitrella patens* and *Lotus japonicus* (Burman et al., 2018; Yamawaki et al., 2011, Nishimura et al., 2002; Weller et al., 2009, Li et al., 2015; Zhai et al., 2016; Zhai et al 2019) but, there are no reports of characterization of *Nicotiana tabacum* HY5 and its role in light-regulated photomorphogenesis as well as flavonoid biosynthesis pathway. In this study, we

demonstrated that NtHY5, an important light signaling component, regulates light-mediated flavonoid biosynthesis in tobacco. To decipher the function of NtHY5 in detail, we functionally complemented it in the corresponding Arabidopsis mutants. Additionally, the binding of NtHY5 to the promoter of *NtMYB12* was validated, which suggests that HY5 activates the expression of NtMYB12 in response to light. We also developed the CRISPR/Cas9-based mutant of HY5 in tobacco, which will be a resource to study the involvement of HY5 in various developmental processes, secondary plant product biosynthesis and stress responses. Using NtHY5 overexpressing tobacco lines and developed mutant, we have demonstrated the role of NtHY5 in providing salt stress tolerance in tobacco.

RESULTS

Light modulates the flavonol accumulation in tobacco

In tobacco, the role of light in the regulation of flavonoid biosynthesis has not been studied in detail. To broaden our understanding of the flavonol regulation through light, we analyzed the expression of genes associated with the phenylpropanoid pathway and accumulation of the major flavonol aglycones and glycosides in *Nicotiana* overexpression lines of AtMYB12 (AtMYB12OX), AtMYB11 (AtMYB11OX), and AtMYB111 (AtMYB111OX). An increased expression of pathway genes and flavonoid accumulation was observed in 10-day light-grown transgenic seedlings (AtMYB12OX, AtMYB11OX and AtMYB111OX) compared to WT (**Fig. 1A, B, C, D, E and F**). Additionally, an increase in antioxidant activity in these transgenic lines compared to WT was observed (**Supplemental Fig. S1**). All these data suggested that similar to Arabidopsis, flavonoid biosynthesis in tobacco is also under the transcriptional control of R2R3-MYBs PRODUCTION OF FLAVONOL GLYCOSIDE1 (PFG1 /MYB12, PFG2 /MYB11 and PFG3/MYB111).

Further, 10-day-old light-grown AtMYB12, AtMYB11 and AtMYB111 overexpression transgenic seedlings were incubated in darkness for 24h and 48h. Analysis suggested that dark incubation led to significant down-regulation in the expression of genes (*NtCHS*, *NtCHI*, *NtFLS* and *NtDFR*) involved in the flavonol biosynthesis (Fig. 1A, C and E). This is accompanied by a

decrease in the accumulation of flavonol aglycones like kaempferol, quercetin and glycosides rutin in overexpressing MYB lines as compared to the WT (Fig. 1 B, D and F; Supplemental Fig. S2). These results indicated that light positively regulates flavonoid biosynthesis, and there could be a possible involvement of other light-associated signaling components, in addition to AtMYB12, AtMYB11, and AtMYB111.

Figure 1. Light affect flavonoid biosynthesis in transgenic tobacco. (A),(C),(E) Relative transcript abundance of *AtCHS*, *AtCHI*, *AtFLS* and *AtDFR* that are involved in phenylpropanoid pathway in AtMYB12, AtMYB11 and AtMYB111 overexpressing transgenic lines respectively

by real time PCR using RNA from the 10-d-old light grown seedlings transferred to dark for 24hr and 48hr. Tubulin was used as endogenous control to normalize the relative expression levels. **(B), (D), (F)** Effect of light over flavonol biosynthesis in different transgenic lines expressing AtMYB12, AtMYB11 and AtMYB111 respectively. Compounds (K, Kameferol; Q, Quercetin; R; Rutin) were quantified using methanolic extracts of 10-d-old old seedlings of different transgenic lines through HPLC. Data are derived from the mean of three technical replicates. Error bars represent SE of means (n=3). The statistical analysis was performed using two-tailed Student's t-tests. The asterisks indicate significant difference, *P<0.1; **P<0.01; ***P<0.001.

NtHY5 and light-dependent flavonoid accumulation

The participation of light through associated transcription factors in the regulation of flavonoid biosynthesis has been demonstrated in plants. Studies in *Arabidopsis*, *Oryza sativa*, *Zea mays* and many other species have shown that HY5 is an important light signaling component (Nawkar et al., 2017; Burman et al., 2018); thus, we studied the role of tobacco HY5 in flavonol accumulation. The BLAST search using AtHY5 in the NCBI database identified a homolog in tobacco with more than 90% sequence similarity (XM_009590053). This was named NtHY5 as it showed significant sequence similarity with sequences from other plant species like tomato, capsicum, etc. NtHY5 gene contains 4 exons and 3 introns similar to Arabidopsis **(Supplemental Fig. S3A)**. Phylogenetic and pairwise distance analysis suggests that NtHY5 is closer to CaHY5 and SIHY5 compared to AtHY5 **(Fig. 2A)**. The NtHY5 showed high homology with Arabidopsis and tomato amino acid sequences **(Fig. 2B)** and contains a VPE motif (COP1-interaction motif) and leucine-rich zinc finger motif (as described by Holm et al., 2002). Analysis through the conserved domain database suggested NtHY5 belongs to the bZIP superfamily, like HY5 proteins of other plant species **(Supplemental Fig. S3B)**.

To study the involvement of NtHY5 in flavonol biosynthesis through controlling the expression of genes associated with the phenylpropanoid pathway genes expression, tobacco (WT) seedlings were grown in the light and continuous dark for 10 days. Seedlings grown in dark showed longer hypocotyl length compared to light-grown seedlings **(Fig. 2C)**. The phytochemical analysis and expression of *NtHY5* and phenylpropanoid pathway genes revealed a significantly higher expression and enhanced flavonoid accumulation and expression of genes in 10-day-old light-grown compared to dark-grown seedlings **(Fig. 2D, E and F)**. To study the responsiveness of flavonols towards light, 5-day-old dark-grown seedlings were transferred to light conditions for 3 h. This dark-to-light transition of seedlings showed significant enhancement in the flavonol

content (**Supplemental Fig. S4**), suggesting light-responsiveness of flavonol biosynthesis. To investigate the spatio-temporal expression of endogenous *NtHY5*, a comparative analysis was carried out at different stages of tobacco development and the result showed 4- to 5-fold higher expression in the mature leaf and young leaf, compared with seedling, root, stem, and flower of the young plant (**Fig. 2G**).

Figure 2. NtHY5, the light signaling component, alters flavonoid content in tobacco. (A) Phylogenetic tree of HY5 homologous proteins (sequence obtained from NCBI database) from across species of different families. The numbering at the nodes represents the number of trees that have the same topology as the consensus tree. NtHY5 is denoted through the red branch, and the scale bar indicates the branch length. (B) Amino acids alignment of NtHY5 and its homologs in Tomato and Arabidopsis using multialignment tool. The VPE motif and zinc finger domain is boxed in green and red color respectively. (C) Phenotype of 10-d-old dark and light grown WT seedlings. (D) Expression analysis of phenylpropanoid pathway genes *NtCHS*, *NtCHI*, *NtFLS* and *NtDFR* in WT seedlings grown in light and dark for 10 days. (E) Quantitative estimation of flavonoids (K, Q and R) in 10-d-light and dark grown seedlings through HPLC. (F) Relative expression of NtHY5 in 10 light and dark grown seedlings. (G) Normalized expression of NtHY5 in different tissue [seedling, young (Y) leaf, mature (M) leaf, stem, root and flower]. Data are derived from the mean of three technical replicates. Tubulin was used as endogenous control to normalize the relative expression levels. Error bars represent SE of means (n=3). The statistical analysis was performed using two-tailed Student's t-tests. The asterisks indicate significant difference, **P<0.01; ***P<0.001.

NtHY5 Binds to the promoter of NtMYB12

As the expression of flavonol biosynthetic genes enhanced in response to light, it was of interest to determine whether the Nicotiana transcription factor, NtMYB12, is controlled by the light signaling component, NtHY5. The *in silico* analysis of the promoters of NtMYB12 revealed the presence of putative LREs (GATA, BOX-4, ACE and LTR) upstream of the transcription start

site (**Supplemental Table S1**). Analysis revealed the presence of two G-boxes with core motif (ACGT) within the 1.5-kb region upstream of the transcription start site of NtMYB12 (**Fig. 3A**). Electrophoretic Mobility Shift Assay (EMSA) using NtHY5 suggested that NtHY5 binds to G-box at -220 bp (G-box I) from the transcription start site and to the other G-box at -1249 bp (G-box II) elements (**Fig. 3B**). Results also suggested that NtHY5 binds to the labeled probe containing G-box but not the mutated version (G-mut I and G-mutII) where the G-box had been disrupted (**Fig. 3C and D**). These results indicate that NtHY5 binds directly to the promoter of NtMYB12 and might regulate its expression.

Figure 3. NtHY5 interacts with NtMYB12 promoter. (A) Representation of the probes (G-BOX I, G-mut I, G-BOX II and G-mut II) designed upstream of the transcription start site of NtMYB12 promoter. G-BOX is altered by multiple base substitutions. (B) EMSA (Electrophoretic mobility shift assay) for the binding of 6X-His-HY5 (NtHY5) with core ACE-DIG element (HY5 binding site that is G-BOX I and G-BOX II) present in the LRE motifs of the NtMYB12 promoter. Upper and lower arrows indicate shift and free probe respectively. (C), (D) EMSA with competition between digoxigenin in labelled G-BOX I, unlabeled G-BOX I (cold), G-BOX II and unlabeled G-BOX II (cold) probe to bind to 6X-His-HY5 respectively. Superscript values 10X, 20X, 30X and 50X represent increasing amount of cold probe. Upper and lower arrows indicate shift and free probe respectively.

NtHY5 is a functional ortholog of AtHY5

To investigate the biological function of NtHY5, it was used to complement Arabidopsis *hy5* mutant (**Fig. 4A; Supplemental Fig. 5A**). The homozygous transgenic lines in the *hy5* background, NtHY5OXL-1 and NtHY5OXL-2, were used for detailed analyses. Higher expression of *NtHY5* was observed in complemented lines compared to WT as well as *hy5* mutant (**Fig. 4B**). Since the Arabidopsis, *hy5* mutant is reported to have an elongated hypocotyl length in light (Koornneef et al., 1980; Jing et al., 2013), thus 7-d-old white light-grown homozygous complemented transgenic lines were analyzed for hypocotyl length. The average hypocotyl lengths of 7-d-old wild-type (WT), *hy5* mutant, and NtHY5;*hy5* seedlings were approximately 0.2, 0.7, and 0.4 mm, respectively. The decrease in hypocotyl length in complemented lines indicated that NtHY5 could compensate for the loss of function of HY5 in the *Athy5* mutant (**Fig. 4A and C**).

As it is previously reported that AtHY5 plays a major role in light-regulated root growth in Arabidopsis (Zhang et al., 2017), we analyzed the root length of 7-d-old light-grown WT and *hy5* mutant Arabidopsis seedlings and found root length to be longer in *hy5* mutant compared with WT. The complementation of the *hy5* with NtHY5 regained the root length as of WT (**Fig. 4D**). The average length of *Athy5* mutant rosette diameter was approximately 6 cm whereas, the length of WT and complemented lines were 4 and 3 cm, respectively. Similarly, seeds of *hy5* mutant also regained the size similar to WT after complementation (**Fig. 4E and F; Supplemental Fig. 5B and C**). These results suggested that NtHY5 could complement the phenotypic function of the *Athy5* mutant.

NtHY5 modulates the expression of flavonol biosynthetic genes in Arabidopsis

To understand the function of *NtHY5*, the expression of flavonol biosynthetic pathway genes was studied in WT, *hy5* mutant and NtHY5;*hy5* complemented seedlings. The analysis suggested that overexpression of *NtHY5* in the *hy5* mutant background restored the expression of genes associated with the phenylpropanoid pathway (*AtCHS*, *AtCHI*, *AtFLS* and *AtDFR*) (**Fig. 4G**). Moreover, the levels of flavonols, such as kaempferol and quercetin, were also restored in NtHY5;*hy5* overexpressed lines compared with *hy5* mutant (**Fig. 4H**).

To investigate the importance of NtHY5 in the regulation of genes associated with flavonol biosynthesis, expression of genes was studied in 5-days-old dark-grown seedlings exposed to light for 3 h. The transcript level of structural genes increased on exposure to light for 3h. By contrast, *hy5* mutant showed inhibited light-dependent induction of genes expression (**Fig. 4I**). The expression analysis of MYB family transcription factors revealed a higher expression of these transcription factors in NtHY5 complemented lines after 3h of light induction compared with *hy5* mutant seedlings. These results suggested that *NtHY5*, through regulation of MYB family transcription factors (*AtMYB12*, *AtMYB11* and *AtMYB111*), affects the flavonoid accumulation in Arabidopsis seedlings. These results also suggest that NtHY5 has the potential to complement *Athy5* at the phenotypic as well as the molecular level.

Figure 4. Complementation of Arabidopsis *hy5* mutant by NtHY5. (A) Phenotype of 7-d-old light-grown WT (wild type; Col-0), *hy5* and NtHY5OX; *hy5* seedlings (L1 and L2 are two independent transgenic lines overexpressing NtHY5 in *hy5* mutant background). (B) Relative expression of NtHY5 in 7-d-old light grown WT, *hy5*, L1 and L2. (C) Hypocotyl length of 7-d-old white light grown WT, *hy5*, L1 and L2 seedlings. (D) Root length of 7-d-old white light grown WT, *hy5*, L1 and L2. (E) Rosette diameter of 30-day-old WT, *hy5*, L1 and L2 plants. (F) Representative photographs of the seed size of WT, *hy5* and transgenic lines (L1 and L2). Bars = 1 mm. (G) Relative expression of phenylpropanoid pathway genes (*AtCHS*, *AtCHI*, *AtFLS* and *AtDFR*) in 7-d-light grown WT, *hy5*, L1 and L2 seedlings. (H) Quantitative estimation of flavonol (Kaempferol and Quercetin) content in 7-d-old light grown WT, *hy5*, L1 and L2 seedlings through HPLC. (I) Expression analysis of *AtCHS*, *AtCHI*, *AtFLS* and *AtDFR* in WT, *hy5*, L1 and L2 seedlings grown for 5 days in dark followed by an exposure to light for 3hr. (J) Length of vascular bundle of 30-day-old stem of WT, *hy5*, L1 and L2 plants. Statistical analysis

was performed using two-tailed Student's t-test. Error bars represent SE of means (n=3) For hypocotyl and root length (n=10-12). Tubulin was used as endogenous control to normalize the relative expression levels. Error bars represent standard deviation. Asterisks indicate a significant difference, *P < 0.1, **P < 0.01, ***P < 0.001. ns represent non-significant.

NtHY5 modulates the levels of lignin, anthocyanin and chlorophyll in Arabidopsis

The phenylpropanoid pathway synthesizes various secondary metabolites, including flavonols, flavones, flavonols, isoflavones, flavan-3-ol, proanthocyanidins (PAs), anthocyanin and lignin. Phenylalanine is reported to be a common precursor for all of them (Vogt et al., 2010). Thus, we also studied the effect of NtHY5 on lignin and anthocyanin production in Arabidopsis. The stem cross-sections of WT, *hy5* and NtHY5;*hy5* overexpression lines (NtHY5OXL-1 and NtHY5OXL-2) were used for lignin staining. The results showed a decreased lignification in the interfascicular and vascular tissues. In addition, the length of the vascular bundle decreased in NtHY5;*hy5* compared with WT and *hy5* mutant plants (**Fig. 4J; Supplemental Fig. S5D**). As AtHY5 is known to be involved in anthocyanin and chlorophyll biosynthesis (Oyama et al., 1997; Chattopadhyay et al., 1998), we estimated anthocyanin and chlorophyll contents of WT, *hy5*, and NtHY5;*hy5* seedlings. In 7-d-old light-grown seedlings, the total chlorophyll content of NtHY5;*hy5* was higher compared with WT and *hy5* (**Supplemental Fig. S7A**). The anthocyanin content in NtHY5;*hy5* was higher than *hy5* seedlings but found to be comparatively lesser than WT (**Supplemental Fig. S7B**). Overall, these results suggest that NtHY5 can compensate for the loss in *Athy5* mutant at the transcript level and at the metabolite level.

Overexpression of NtHY5 in tobacco

In addition to the functional complementation of the *hy5* mutant of Arabidopsis by NtHY5, we generated transgenic tobacco lines to study if NtHY5 performs additional or unique functions in tobacco. The expression of *NtHY5* and pathway genes (*NtCHS*, *NtCHI*, *NtFLS* and *NtDFR*), as well as *Nicotiana* MYB family transcription factor (*NtMYB12*), enhanced in 10-d-old light-grown transgenic compared to WT tobacco seedlings (**Fig. 5A, B**). The analysis of metabolites in the NtHY5OXL-1, NtHY5OXL-2 and NtHY5OXL-3 lines suggested a significantly higher accumulation of flavonols (kaempferol, quercetin and rutin) and the CGA levels (**Fig. 5C**). The flower of NtHY5OXL-1, NtHY5OXL-2 and NtHY5OXL-3 lines showed reduced pigmentation compared to WT (**Fig. 5D**). Analysis suggested reduced anthocyanin content in overexpression lines compared to the WT (**Fig. 5F**). As it is known that flavonol content leads to the antioxidant

activity of plants (Pandey et al., 2015; Song et al., 2020), total antioxidant activity was measured in extracts from NtHY5OXL-1, NtHY5OXL-2 and NtHY5OXL-3 tobacco seedlings. Analysis

suggested enhanced antioxidant activity in transgenic lines compared to the WT (**Fig. 5G**). Altogether, these results suggest that overexpression of NtHY5 modulates metabolic content such as flavonol and anthocyanin along with the antioxidant activity of tobacco.

Figure 5. Overexpression of NtHY5 modulates gene expression and metabolite level in tobacco. (A) Relative expression of NtHY5 in 10-d-old light grown WT and NtHY5OX lines of tobacco (L1, L2 and L3). (B) Expression analysis of phenylpropanoid pathway genes (*NtCHS*, *NtCHI*, *NtFLS* and *NtDFR*) in 10-d-old light grown WT and NtHY5OX lines. (C) Quantification of flavonoids (K, Q and R) and polyphenolic compound (CGA) in 10-d-old light grown WT and NtHY5OX lines of tobacco using HPLC. (D) Representative photograph of flower color alterations in petals of NtHY5OX transgenic lines in compared with WT. (E) Total anthocyanin content in petals of WT and NtHY5OX transgenic lines. (F) Estimation of antioxidant activity by

trolox equivalent in WT and NtHY5OX lines. TUBULIN was used as the endogenous control to normalize the relative expression levels. The statistical analysis was performed using two-tailed Student's t-tests. The data are plotted as means $\pm$ SD (n= 3). The error bars represent standard deviations. The asterisks indicate significant difference, *P< 0.1; **P< 0.01; ***P< 0.001.

Development and analysis of NtHY5 mutants

To gain a deeper insight into NtHY5 function, we developed NtHY5 mutant plants using the CRISPR–Cas9 approach. To mutate NtHY5, we used guide RNAs (gRNAs) targeting the coding region (**Supplemental Fig. S8**). The nucleotide sequence analysis of the mutants revealed deletions and insertions in the coding regions of NtHY5 (**Fig. 6A**). The *NtHy5^{CR}* lines showed frame-shift mutations leading to truncated peptides (**Supplemental Fig. S8**). The expression analysis of NtHY5 in mutants revealed a significant reduction in NtHY5 expression (**Fig. 6B**). Expression of *NtMYB12* and other phenylpropanoid pathway genes, like *NtCHS*, *NtCHI*, *NtFLS* and *NtDFR*, was also found to be significantly reduced (**Fig. 6C**). Phytochemical analysis suggested a significantly decreased level of flavonols (kaempferol, quercetin and rutin), CGA and anthocyanin content in *NtHy5^{CR}* lines compared to WT (**Fig. 6D, E, F, G and H**). The decrease in the expression of phenylpropanoid pathway genes and NtMYB12 and reduced

biosynthesis in tobacco.

Figure 6. CRISPR-Cas9-mediated knockout of NtHY5 alters gene expression and accumulation of metabolite content. (A) Representation of mutations in target region in three independent transgenic lines of tobacco. Dots indicate deleted nucleotides. Yellow highlight denotes gRNA region used for targeting the sequence. Red highlights denote indels in nucleotides. (B) Relative expression of NtHY5 in 10-d-old light grown WT and *NtHY5^{CR}* lines of tobacco (L1, L2 and L3). (C) Relative transcript abundance of phenylpropanoid pathway genes (*NtCHS*, *NtCHI*, *NtFLS* and *NtDFR*) in 10-d-old light grown WT and *NtHY5^{CR}* lines. (D) Quantification of flavonoids (K, Q and R) and polyphenolic compound (CGA) in 10-d-old light grown WT and *NtHY5^{CR}* lines of tobacco. (E) Quantification of Anthocyanin content in 10-d-old light grown WT and *NtHY5^{CR}* lines. (F) Quantification of antioxidant activity by trolox equivalent in 10-d-old light grown WT and *NtHY5^{CR}* lines. Tubulin was used as the endogenous control to normalize the relative expression levels. The statistical analysis was performed using two-tailed Student's t-tests. The data are plotted as means $\pm$ SD (n=3). The error bars represent standard deviations. The asterisks indicate significant difference, *P<0.1; **P<0.01; ***P<0.001.

NtHY5 regulates growth and development of tobacco

Several previous studies have demonstrated that AtHY5 regulates the hypocotyl length by repressing the expression of hypocotyl cell elongation-related genes (Liu et al., 2013). To study the difference in hypocotyl length in *NtHY5* overexpression and *NtHy5^{CR}* lines, we measured the hypocotyl length of 10-day-old light- and dark-grown tobacco seedlings. In light-grown seedlings, the hypocotyl length of *NtHy5^{CR}* mutant seedlings was significantly higher compared to overexpression lines (Fig. 7A, B). At the same time, hypocotyl length of the dark-grown overexpressing lines was significantly higher compared to *NtHy5^{CR}* mutant plants (Fig. 7C; Supplemental Fig. S9A).

It is reported that the asymmetric accumulation of flavonoids in root tips in Arabidopsis affects the direction of root growth (Wan et al., 2018). Thus, we determined the effect of NtHY5 on root length and root agravitropic response. For this, the root bending angle of 10-d-old light- and dark-grown, NtHY5 overexpression and *NtHy5^{CR}* seedlings was measured. A significant increase in root length of *NtHy5^{CR}* seedlings compared to overexpression lines was observed (**Fig. 7D**). Furthermore, observations suggested that the root-bending angle of NtHY5OXL-1 and NtHY5OXL-2 seedlings was significantly higher compared to *NtHy5^{CR}*L-1, *NtHy5^{CR}*L-2 and WT plants (**Fig. 7E**). Our analysis suggested that the *NtHy5^{CR}* has a smaller seed size compared to the overexpression plants (**Fig. 7F; Supplemental Fig. S9B**). We also counted the number of seeds per unit (5 mg) weight and *NtHy5^{CR}* had a significantly larger number of seeds compared to overexpression lines (**Fig. 7G**). In addition to these factors, the plant height of 40-days-old *NtHy5^{CR}* plants was significantly higher than NtHY5 overexpression and WT plants (**Fig. 7H and I**). The dark-to-light transition was able to enhance the expression of genes to several folds in NtHY5 overexpressed lines. However, such an increase was not observed in *NtHy5^{CR}* plants (**Fig. 7J and K**). These results suggested that NtHY5 is crucial for plant growth and light-responsive expression of genes associated with the phenylpropanoid pathway.

Figure 7. NtHY5OX and CRISPR–Cas9-derived knockout mutants show altered phenotypes and gene expression. (A) Representative image of 10-d-old light grown WT, NtHY5OX and NtHY5^{CR} seedlings grown on half-strength MS medium. Scale bars, 1 cm. (B), (C) Hypocotyl length of 10-d-old light and dark grown WT, NtHY5OX and NtHY5^{CR} seedlings. (D) Root length of 10-d-old light grown WT, NtHY5OX and NtHY5^{CR} seedlings. (E) Measurement of root bending angle of 10-d-old light grown WT, NtHY5OX and NtHY5^{CR} seedlings. (F) Quantitative analysis of seed size of WT, NtHY5OX and NtHY5^{CR} plants. (G) Quantitative estimation of Number of seeds per 5 mg of WT, NtHY5OX and NtHY5^{CR} plants. (H) Representative image of 40-day-old WT, NtHY5OX and NtHY5^{CR} plants showing plant heights. (I) Data presenting difference in height of 40-day-old WT, NtHY5OX and NtHY5^{CR} plants. (J), (K) Relative expression of NtHY5 and phenylpropanoid pathway genes (*NtCHS*, *NtCHI*, *NtFLS* and *NtDFR*) in 10-d-old light grown seedlings transferred to dark for 3h. Tubulin was used as the endogenous control to normalize the relative expression levels. The statistical analysis was performed using

two tailed Student's t-tests. The data are plotted as means $\pm$ SD (n= 3). For measurement of root bending angle, root length, hypocotyls length and seed size (n=10-12). The error bar represents standard deviations. The asterisks indicate significant difference, *P< 0.1; **P< 0.01; 0.001.

NtHY5 enhanced tobacco tolerance to salt stress

Flavonoids are known to positively regulate the abiotic stress tolerance ability of plants (Wan et al., 2018; Chen et al., 2019; Bhatia et al., 2018). To study the response of NtHY5 under salt stress, leaf disc assay was performed to analyze the comparative effect of NaCl (200 mM) on leaves of NtHY5OX lines and *NtHy5^{CR}* and WT plants. After 3 days of salt stress, higher chlorosis was observed in the leaves of *NtHy5^{CR}* compared with leaves of NtHY5OX and WT plants (**Fig. 8A and B**). Chlorosis affects the survival of leaves by decreasing the chlorophyll content (Pandey et al., 2014). Thus, the survival of leaves was measured by analyzing the total chlorophyll content as well as chlorophyll a and b. Higher chlorophyll content was observed in NtHY5OX leaves compared with *NtHy5^{CR}* and WT leaves after 3 days of salt stress (**Fig. 8C**). It is well known that salt stress leads to excessive accumulation of ROS, leading to cell death (Chen et al., 2019; Ahanger et al., 2017). Higher NBT and DAB staining were observed in *NtHy5^{CR}* compared with NtHY5OX and WT, suggesting that NtHY5 might play a crucial role in salt stress tolerance in tobacco by inhibiting ROS accumulation (**Fig. 8D; Supplemental Fig S10**). A higher ROS accumulation suggests the lower antioxidant activity in *NtHy5^{CR}* lines compared with NtHY5OX and WT.

487 **Discussion**

488 **HY5 participates in light-induced flavonoid biosynthesis in *Nicotiana tobaccum***

489 Studies on flavonol biosynthesis in different plant species suggest a tight regulation by multiple
 490 exogenous and endogenous factors, involving many transcription factors (Stracke et al., 2007;
 491 Stracke et al., 2010a; Pandey et al., 2014; Sharma et al., 2016). In tobacco, regulation of
 492 flavonoids is not studied extensively; however, few reports suggest the importance of
 493 transcription factors in the regulation of flavonoid biosynthesis. According to previous studies,
 494 up-regulation of three functionally redundant transcription factors, namely AtMYB12,
 495 AtMYB11 and AtMYB111, enhance the flavonoid biosynthesis in tobacco by activating a
 496 branch of the phenylpropanoid pathway (Mishra et al., 2010; Pandey et al., 2012; Pandey et al.,
 497 2014; Pandey et al., 2015). Our result suggests that dark exposure inhibits flavonol accumulation
 498 by decreasing the expression of pathway genes in spite of overexpression of AtMYB12,
 499 AtMYB11 and AtMYB111 in tobacco (**Fig. 1**). Similar to our result, a decrease in flavonol
 500 content after dark exposure in AtMYB12 and AtMYB111 overexpressing lines in Arabidopsis
 501 and mature tobacco plants, respectively, was reported (Bhatia et al., 2021; Pandey et al., 2015).
 502 Our data also suggest that the up-regulation of genes involved in the flavonoid biosynthesis
 503 pathway by AtMYB111 is significantly higher compared to AtMYB12 and AtMYB11, although
 504 they are functionally redundant. This might be due to the differential influence of MYB11,
 505 MYB12 and MYB111 on the spatial accumulation of specific flavonol derivatives, as reported
 506 by Stracke et al., (2010a). All these observations gave rise to a fundamental question, i.e.,

whether there is an additional factor other than these transcription factors involved in the regulation of flavonol accumulation in tobacco.

8. salt

transgenic. Fully expanded leaves of 30-d-old plants to floated on 6 ml of 200 mM NaCl solution (for salt stress) for 3 days. **(A), (B)** The phenotypic (chlorosis) changes due to salt treatment on leaf discs were assessed on 0 and 3 days respectively. **(C)** Quantification of chlorophyll content of WT, NtHY5OX and NtHY5^{CR} leaves after 3day of salt stress. **(D)** Detection of Superoxide ($O_2^{\cdot -}$) by NBT staining in 10-d-old light grown WT, NtHY5OX and NtHY5^{CR} tobacco seedlings subjected to 200 mM salt stress for 3 hours. The statistical analysis was performed using two-tailed Student's t-tests. The data are plotted as means $\pm$ SD (n=3). The error bars represent standard deviations. The asterisks indicate significant difference, *P<0.1; **P<0.01; ***P<0.001.

Previous studies in *Arabidopsis* and other plant species suggest HY5 as a regulator of the photomorphogenesis and a key signaling component of light (Ulm et al., 2004; Lee et al., 2007; Nawkar et al., 2017). HY5 is known to interact with LREs present in the minimal promoters of several genes, including *CHS*, *CHI*, *FLS*, and *MYB12* (Chattopadhyay et al., 1998; Shin et al., 2013; Zhou et al., 2017). We hypothesized that HY5 might be one of the potential factors responsible for light-mediated flavonol accumulation in tobacco. Our results suggest that *NtHY5* was light-responsive; therefore, it is possible that *NtHY5* might be a key regulatory factor in flavonoid biosynthesis in tobacco. Expression and metabolite analyses suggest that *NtHY5* together with *NtMYB12* regulate flavonol biosynthesis. The expression profile also suggested the highest expression of *NtHY5* in young and mature leaf, indicating that *NtHY5* is developmentally regulated.

***NtHY5* interacts with *NtMYB12* promoter to promote flavonoid biosynthesis**

HY5 is reported to have a strong binding affinity to G-box (CACGTG), T/G-box (CACGTT), E-box (CAATTG), GATA-box (GATGATA), ACE-box (ACGT), CT-Box (GACGTT), Z-box (ATACGGT), Cbox (GTCANN), A-box (TACGTA), hybrid C/G-(G) and C/A-boxes present in promoters of many genes and transcription factors (Song et al., 2008). Previous studies confirmed that *AtHY5* regulates flavonoid biosynthesis by binding to promoters of specific members of the MYB family transcription factors to promote their expression (Stracke et al., 2010; Shin et al., 2013; Nguyen et al., 2015). Further, *MYB12* and *MYB111* regulate the expression of early flavonoids biosynthetic genes by binding to their promoters, whereas *MYB75* binds to the late flavonoids biosynthetic genes (Stracke et al., 2010a; Shin et al., 2013). We analyzed the promoter sequence of *NtMYB12* and identified several light-responsive elements such as G-box I (at -220 bp) and G-box II (at -1249 bp) (**Supplemental Table S1**), which suggested that *NtHY5* might bind to the promoter of *NtMYB12* to regulate the flavonol content in tobacco. Our EMSA results suggested the binding of *NtHY5* to the G-BOX element of the *NtMYB12* promoter. This result suggests that *NtHY5* binds directly to the conserved light-responsive element that is ACE (ACGT) and promotes *NtMYB12* expression, which in turn up-regulates the expression of genes associated with the phenylpropanoid pathway.

NtHY5 rescue the *Athy5* mutant at phenotypic and metabolite levels

Several HY5 orthologs from different plant species have been identified and characterized (Cheong et al., 1998; Nishimura et al., 2002; Burman et al., 2018). OsZIP48 is reported as a functional ortholog of AtHY5 in rice as it was able to rescue the phenotype of Arabidopsis *hy5* mutant plants after complementing with OsZIP48. Similarly, overexpression of STF1, the HY5 ortholog in soybean, can fully complement *hy5* phenotype for hypocotyl length, chlorophyll, anthocyanin and root gravitropic response (Song et al., 2008). In our study, we demonstrated that NtHY5 could complement Arabidopsis *hy5* mutant for phenotypes such as hypocotyl length, root length, seed size and rosette diameter (**Figure 4 A, C, D, E and F; Supplemental Fig. B, C and D**). Moreover, the RT-qPCR analysis showed that flavonoid biosynthetic structural genes, including *AtCHS*, *AtCHI*, *AtFLS*, *AtDFR* and MYB transcription factors, were significantly up-regulated in *NtHY5;hy5*. The content of metabolites, such as anthocyanin and chlorophyll, was also restored in complemented lines. Cross-sections of stems in Arabidopsis *hy5* mutant showed more lignin-specific staining (**Supplemental Fig. S5D**) compared with *NtHY5;hy5*, indicating the higher accumulation of flavonoids in complement lines due to the diversion of the substrate flux towards flavonoid synthesis at the cost of lignin production. These findings suggest that NtHY5 had a conserved function across different species, and it regulates flavonoid biosynthesis in complemented lines of Arabidopsis.

Modulated expression of genes and metabolite content in *NtHY5OX* and *NtHy5^{CR}* plants

There are plenty of reports, which suggest that the expression of MYBs in various plant species regulates flavonoid content (Misra et al., 2010; Bhatia et al., 2018). In our study, overexpression of NtHY5 in tobacco led to the enhanced expression of genes involved in the phenylpropanoid pathway. Quantitative estimation of flavonol content suggested increased accumulation of kaempferol, quercetin, rutin and CGA in overexpression lines compared with WT. On the contrary, mutant lines showed a decrease in flavonoid content due to the reduced expression of pathway genes as well as *NtMYB12*. Altogether, these results suggest that NtHY5 is functionally similar to HY5 of other plant species in the regulation of flavonoid biosynthesis and NtHY5 up-regulates the flavonol accumulation via increasing the expression of NtMYB12 and other phenylpropanoid pathway genes. Additionally, anthocyanin content reduced in *NtHy5^{CR}*,

indicating the decreased accumulation of precursor substrate common for both flavonoid and anthocyanin biosynthesis.

Role of NtHY5 in plant growth and development

HY5 and specific MYBs are known to be involved in plant growth and development (Zhong et al., 2008; McCarthy et al., 2009; Gangappa et al., 2016; Misra et al., 2010). OsbZIP48 ortholog is reported to cause a reduction in cell size, thus reduces the plant height of OsbZIP48 overexpression transgenic plants (Burman et al., 2018). HY5 negatively regulates hypocotyl elongation by suppressing auxin signaling (Cluis et al., 2004, Sibout et al., 2006). Similar to the above results, we also observed variation in phenotypic characters of the developed NtHY5 overexpression and mutant plants of tobacco. The results suggested that modulation in the NtHY5 expression changes in various morphological characteristics, with contrasting phenotypes between NtHY5OX and *NtHy5^{CR}* plants. Hypocotyls length of mutants was significantly longer than overexpressed and WT seedlings grown in light for 10 days. Interestingly, shorter hypocotyl length was observed in mutants in comparison to overexpressed and WT in dark-grown seedlings (**Fig 7 A, B**). This suggests that mutation in NtHY5 eradicates the light-dependent repression of cell elongation and does not promote cell elongation independent of the light stimulus. Similar to our result, Oyama et al., (1997) also reported the loss of light-dependent cell elongation in *Athy5* mutant plants in the dark.

Previous reports suggest that HY5 negatively regulate the auxin signaling pathway by promoting the expression of auxin signaling inhibitors such as INDOLE ACETIC ACID 14 (SLR/IAA14) and INDOLE ACETIC ACID 7 (AXR2/IAA7), whereas mutation in *hy5* promotes up-regulation of auxin signaling genes (Sibout et al., 2006). Flavonoids negatively regulate plant growth and root length by down-regulating the auxin signaling genes (Brown et al., 2001; Peer et al., 2007; Yin et al., 2014; Sharma et al., 2020). In our study, we found that up-regulated flavonol accumulation in NtHY5OX lines resulted in a decrease in root length, plant height as well as high flavonol content. In addition, increased root bending angle in was observed in NtHY5OX lines compared with tobacco *hy5* mutant and WT (**Fig. 7B**). Additionally, the seed size of the NtHY5 mutant was significantly smaller in comparison to mutant and WT plants (**Fig. 7K**). This significant change in seed size might be due to modulation in hormone signaling mediated by flavonoids. Rapid changes in auxin distribution by ZmPIN1 in the endosperm of maize affect the

seed size (Lur et al., 1993). There are some other reports, which suggest that flavonol accumulation regulates the seed development in Arabidopsis (Doughty et al., 2014; Sharma et al., 2018). Thus, the relation between hormone signaling and NtHY5 could be the subject for further study.

NtHY5 enhances the salt stress tolerance in tobacco

It is well explored that flavonoids have antioxidant properties and have an important role in abiotic stress tolerance. Previous studies showed that AtHY5 plays a role in abiotic stress tolerance by enhancing the flavonol level in plants (Agati et al., 2012; Emiliani et al., 2013 and Muhlemann et al., 2018). In tobacco, several studies reported that high flavonoid content causes high antioxidant properties, which may result in abiotic stress tolerance (Docheva et al., 2014; Pandey et al., 2015). Recent studies indicate that over-expression lines of HDA15 were found to be resistant to salt stress and also displayed increased accumulation of abscisic acid. It was also shown that HY5 directly interacts with HDA15 promoting salt tolerance (Truong et al., 2021). RNAi of NtMYB4 causes up-regulation of NtCHS, leading to enhanced levels of flavonol content and higher antioxidant property and salinity tolerance in tobacco (Chen et al., 2019). Similarly, overexpression of NtMYB12 causes more flavonol accumulation and higher ROS scavenging property thus enhancing tolerance to low Pi stress in tobacco (Song et al., 2020). In our study, the antioxidant activity was higher in overexpression lines in comparison to WT and *NtHY5^{CR}* plants. This leads to lesser chlorosis in salt-treated young leaves of NtHY5OX lines compared with *NtHy5^{CR}*, suggesting that high flavonoid content leads to higher ROS scavenging ability of overexpression lines. As tobacco *NtHy5^{CR}* mutants were salt-sensitive, thus more ROS is accumulated in salt-treated mutant leaves (**Fig. 8D**). These results suggest that NtHY5 promotes salt stress tolerance at least partially through the antioxidant activity of flavonoids.

Conclusion

In conclusion, the present study reveals that NtHY5 is responsible for light-mediated flavonoid biosynthesis as well as for the enhanced antioxidant property of tobacco plants. The CRISPR-edited tobacco *NtHy5^{CR}* plants showed a reduced expression of regulatory and structural genes of the flavonoid biosynthesis pathway, which is accompanied by a lower accumulation of flavonoid and antioxidant properties. The high flavonol content and antioxidant property caused less chlorosis in *NtHY5* overexpressed plants and enhanced ROS scavenging property, which leads to

an increase in salt-tolerance property in NtHY5 overexpressing tobacco leaves. This response likely occurs through NtHY5 binding to the ACE and G-BOXES (cis-elements) present within the promoters of flavonoid biosynthetic genes and NtMYB12, respectively. Thus, NtHY5 and NtMYB12 share common target genes, which are involved in flavonol production and exhibit

combinatorial regulation of the flavonoid biosynthesis genes. Therefore, we propose that these regulatory factors act in concert for light-mediated flavonoid production (Fig. 9). The *NtHy5* mutant developed in this study will be used as an essential tool in elucidating the involvement of NtHY5 in various biological processes in tobacco.

Figure 9. Proposed working model for the role of NtHY5 and NtMYB12 in regulating the expression of flavonoid pathway genes during light exposure. We propose that NtHY5 positively regulates the expression of NtMYB12 by directly binding to promoter of NtMYB12 through G-BOX I and G-BOX II in turn NtMYB12 directly binds to the MRE (MYB recognition element) in the promoters of flavonoid pathway genes and enhances flavonol accumulation whereas NtHY5 also binds independently to ACE element present on the promoter of flavonoid pathway genes and regulate their expression. Thus, together NtHY5 and NtMYB12 lead to flavonol production. Flavonol ultimately leads to salt tolerance by inhibiting ROS production. Arrowheads and tees indicate positive and negative regulation respectively.

MATERIALS AND METHODS

Isolation and sequence analysis of NtHY5. To identify the ortholog of AtHY5 in tobacco, BLASTn searches were performed on National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/BLAST>). The sequence showing 90% similarity with AtHY5 and SlHY5 was retrieved and amino acid sequence was analyzed for the leucine rich bZIP Zinc finger domain and VPE domain using multallignment tool. Conserved domain database (CCD) was also employed to ensure the bZIP family of the identified NtHY5 protein. Promoter sequences (~1.5 kb upstream of translation start site) of NtMYB12 gene was retrieved from the respective databases AGRIS (Palaniswamy et al., 2006) and PlantCare (Lescot et al., 2002), which was utilized for the scanning of light-responsive cis-elements (LREs).

Phylogenetic analysis. Homologs of AtHY5 that belongs to different families were identified by using the National Center for Biotechnology Information BLASTn tool. The genes retrieved were checked for the presence of the requisite domains using InterProScan (<http://www.ebi.ac.uk/Tools/pfa/iprscan/>; Quevillon et al., 2005; Jones et al., 2014; Mitchell et al., 2015). A phylogenetic tree was constructed using the Neighbor-Joining (NJ) algorithm in MEGA6.0 software (Kumar et al., 2008).

Plant material, growth conditions and treatments. Young leaves of *Nicotiana tabacum* cv. *Petit Havana* have been used for raising the transgenic plants. Tobacco plants were grown in glass house at 25°C±2°C and 16 h/8 h light-dark photoperiods for harvesting of tissues belonging to different developmental stage. For dark treatment, seeds were surface sterilized using 70% EtOH for 1 min and then dipped in 50% bleach for 10 min after that washed repeatedly with autoclaved Milli-Q water, and grown for 10 days in culture room with 16 h/8 h light-dark photoperiods, 24°C temperature on one-half-strength Murashige and Skoog (MS) medium (Sigma) plates and then transferred in dark chamber with similar growth conditions as in culture room. AtMYB12, AtMYB11 and AtMYB111-expressing seedlings were harvested after 24 and

48 h of dark treatment for gene expression as well as phytochemical analysis. Samples were frozen in the liquid nitrogen and kept in -80°C deep freezer until further use. The experiment was performed by using three independent replicates. Arabidopsis (ecotype Col-0) was used as the wild-type plant in complementation study. Seeds were surface sterilized and placed on one-half-strength MS medium containing 1% Sucrose. After stratification for 2 to 3 d at 4°C in the dark, seeds were transferred to a growth chamber under controlled conditions of 16-h-light/8-h-dark photoperiod cycle, 22°C temperature, 150 to 180 $\mu\text{mol m}^{-2}\text{s}^{-1}$ light intensity, and 60% relative humidity for 10 d unless mentioned otherwise.

Plasmid construction and generation of Arabidopsis and Tobacco transgenic lines. To overexpress NtHY5, the NtHY5 was amplified from the single-stranded complementary DNA (cDNA) library prepared from 10-d-old tobacco seedlings through PCR using specific primers. The full-length open reading frame of the NtHY5 cDNA, under the control of cauliflower mosaic virus 35S promoter in the binary vector pBI121 (Clontech, USA) was transferred into *Agrobacterium tumefaciens* strain GV3101 and used to transform Arabidopsis (Col0) by floral dip method (Clough and Bent, 1998) and tobacco plants by leaf disk method (Horsch et al., 1985). Several transgenic tobacco lines constitutively expressing NtHY5 cDNA were selected on the basis of RT-PCR. Seeds were harvested, sterilized and plated on solid half strength MS medium supplemented with 100 mg/l kanamycin. Antibiotic resistant plants were shifted to the glass house and grown until maturity.

For generation of mutant plants of NtHY5, CRISPR-Cas9 technology was used. A 20-bp gRNA was selected from the identified coding region. NtHY5 gRNAs was cloned into the binary vector pHSE401 using the BsaI restriction site (Xing et al., 2014). This vector contains Cas9 endonuclease-encoding gene under dual CaMV35S promoter as well as genes encoding neomycin phosphotransferase and hygromycin phosphotransferase as selection markers. All the constructs were sequenced from both the orientations using plasmid-specific forward and vector reverse primers after that transferred to GV3101 and then transformed to tobacco.

Gene expression analysis. Total RNA was isolated using a Spectrum Plant Total RNA kit (Sigma Aldrich) following the manufacturer's instruction. For qRT-PCR analysis, 2 μg of DNase-free RNA was reverse transcribed using a RevertAid H minus first-strand cDNA synthesis Kit (Fermentas) according to the manufacturer's instructions. qRT-PCR was performed using

Fast Syber Green mix (Applied Biosystems) in a Fast 7500 Thermal Cycler instrument (Applied Biosystems). Expression was normalized using Tubulin and analyzed through the comparative CT method (Livak and Schmittgen 2001). The oligonucleotide primers used to study expression of different genes were designed using the Primer Express 3.0.1 tool (Applied Biosystems) and information is provided in Supplementary Table S1.

Extraction and quantitative estimation of flavonols. Flavonols were determined either as aglycones or as flavonol glycosides by preparing acid-hydrolyzed or non-hydrolyzed extracts respectively. Extraction of flavonols was carried out by grinding the plant material (300mg) into the fine powder in liquid N₂ and then extracted with 80% methanol overnight at room temperature with brief agitation. The extract was hydrolyzed with an equal amount of 6 N HCl at 70°C for 40 min followed by addition of an equal amount of methanol to prevent the precipitation of the aglycones (Jiang et al. 2015). Extracts were filtered through 0.2 mm filter (Millipore, USA) before HPLC. For non-hydrolyzed extracts, samples were extracted as described previously (Pandey et al., 2014). Breeze 2 software (Waters) was used for the quantification of various metabolites through HPLC. The mobile phase was a gradient prepared from 0.05% (v/v) ortho-phosphoric acid in HPLC-grade water (component A) and methanol (component B). Before use, the components were filtered through 0.45-mm nylon filters and de-aerated in an ultrasonic bath. The gradient from 25 to 50% B in 0–3 min, 50 to 80% B in 3–18 min, 80 to 25% B in 25 min and 25% B in 30 min was used for conditioning of the column with a flow rate of 1 ml/min. All the samples were analysed by HPLC–PDA with a Waters 1525 Binary HPLC Pump system comprising PDA detector following the method developed by (Niranjan et al., 2011).

Extraction of total anthocyanin, chlorophyll and lignin staining. Anthocyanin was measured according to the method described by Li et al., (2016). Briefly, 300mg tissue were boiled for 3 min in extraction solution (propanol:HCl:H₂O, 18:1:81), and incubated at room temperature in the dark for at least 2 h. Samples were centrifuged, and the absorbance of the supernatants was measured at 535 and 650 nm. Anthocyanin content was calculated as

$$\text{Anthocyanin content} = (A_{535} - 2.2 \cdot A_{650}) / \text{g FW}.$$

Chlorophyll content was measured by using the 300mg tissue of 10-d-old WT and transgenic seedlings. Sample was crushed in 2 ml (80 %, v/v) chilled acetone (Arnon 1949). Homogenized tissues were centrifuged at 7,800g for 10 min. Supernatant was collected and absorbance was recorded at 663, 645, 510 and 480 nm. Total chlorophyll content was calculated in mg g⁻¹ fresh wt using formula of Machlachalan and Zalik (1963).

$$\text{Total chlorophyll} = [34.5(A_{652} * V) / d * 1000 * W]$$

Where, V=volume of the final extract (ml), W= weight of the plant sample (g) and d=width of the cuvette(1cm).

To visualize lignified cells in stems, hand-cut sections were stained using phloroglucinol (Sigma-Aldrich) for 1min and visualized on a Leica DM2500 microscope (Sharma et al., 2020).

Determination of total antioxidant activity. A method by Wong et al., (2006) with slight modification was used to determine the antioxidant activity. DPPH (1, 1-diphenyl-2-picrylhydrazyl free radical) assay was used in determining the free radical scavenging activity. In brief, 0.1 mM solution of DPPH was prepared in methanol. The initial absorbance of DPPH in methanol was measured at 515 nm and did not change throughout the period of assay. An aliquot of the samples (50–100 µl) was used to make up volume to 1.5 ml with DPPH solution. The sample was incubated at room temperature for 30 min. The absorbance of the samples was recorded at 515 nm using Ultraspec 3000 UV/Vis spectrophotometer (Pharmacia Biotech Ltd., Cambridge, CB4, 4FJ, UK). Trolox was used to make the calibration curve using different dilutions. The antioxidant capacity based on the DPPH free radical scavenging ability of the extract was expressed as mM Trolox equivalents per gm of plant material on fresh weight basis.

Expression and purification of NtHY5. The complete open reading frame of NtHY5 was amplified and cloned into pET-28b(+) vector (Novagen, Germany). The construct was transformed into *E.coli* BL21 (DE3) pLysS (Invitrogen, USA) for prokaryotic expression, induced by IPTG. Recombinant 6X-Histagged HY5 was purified using a Ni-NTA column (Nucleopore, India) and the eluted and was quantified through Bradford assay.

Electrophoretic mobility shift assay. The 2nd generation DIG Gel Shift EMSA kit (Roche, USA) was used to label the probes with digoxigenin as per manufacturer's instructions. Labelled

probes were incubated at 21 °C for 30 min in binding buffer [100 mM HEPES (pH 7.6), 5 mM EDTA, 50 mM (NH₄)₂SO₄, 5 mM DTT, Tween 20, 1% (w/v), 150 mM KCl] with or without recombinant protein. Unlabeled probes were added to the reaction solution in increasing concentrations to test specific binding. The binding reaction was resolved on a 6% polyacrylamide gel in 0.5X TBE (pH 8.0) buffer and was semi-dry blotted (Transblot, BIO-RAD, USA) onto a positively charged nylon membrane (BrightStar, Invitrogen, USA) followed by UV cross-linking. The membrane was finally incubated with CSPD chemiluminescent solution and exposed to X-ray blue film (Retina, India).

Salt stress treatment. Salt stress analysis was performed by using the young leaves from 2nd and 3rd nodes from 1-month-old *NtHY5OX*, *NtHy5CR* and WT plants to determine the comparative response during the leaf discs' salt assay. These leaves were transferred into salt solution of 200 mM. The treatment was carried out at 25 °C for 3 days. In order to determine the effect of salinity stress, survival of leaves and chlorophyll content of NaCl treated leaves were analysed. All the data were recorded from 0 to 3 day of NaCl treatment. The experiment was repeated three times.

Nitro blue tetrazolium (NBT) and 3,3'-diaminobenzidine (DAB) staining. Superoxide (O₂⁻) staining was performed by infiltration by the nitro blue tetrazolium (NBT) method (Liu et al., 2018; Song et al., 2019) with some modification. WT as well as transgenic seedlings were grown for 10 days on half strength MS media then transferred to 200mM salt solution for 3hours. After 3 hours of salt stress seedlings were dipped in 0.5mg/ml NBT solution for 2h followed by removal of chlorophyll by heating at 70°C temperature until chlorophyll is removed. Leaves were directly visualized for NBT staining.

Hydrogen peroxide (H₂O₂) staining of 10-day light grown tobacco seedlings was performed by staining with 3,3'-diaminobenzidine (DAB) using an adaptation of previous methods (Zhao et al 2016). DAB is oxidized by hydrogen peroxide in the presence of some haem-containing proteins, such as peroxidases, to generate a dark brown precipitate. WT as well as transgenic seedlings were grown for 10 days on half strength MS media then transferred to 200mM salt solution for 3hours. After 3 hours of salt stress seedlings were dipped in 1mg/ml DAB solution for 5 hours. Following the incubation, DAB staining solution was replaced with bleaching solution (ethanol:

acetic acid: glycerol = 3:1:1). After that chlorophyll was removed by heating at 90-95°C temperature. Leaves were directly visualized for DAB staining.

Statistical analysis. Data are plotted as means $\pm$ SE (n=3) in the figures. Data of flavonol accumulation were evaluated for statistical significance using the Two-tailed Student's t-test using Graphpad prism 5.01 software. Asterisks indicate significance levels with (* P < 0.1; ** P < 0.01; *** P < 0.001) as confidence intervals.

Supplemental Data

Figure S1. Antioxidant activity by trolox equivalent. Antioxidant activity was analysed in 10-day-old AtMYB12, AtMYB11 and AtMYB111 overexpressing tobacco transgenic lines grown in light. Statistical analysis was performed using two-tailed Student's t-test. Data are plotted as means $\pm$ SD (n=3). Error bars represent standard deviation. Asterisks indicate a significant difference***P < 0.001.

Figure S2. Metabolite analysis of MYB12OX lines of tobacco through HPLC. (A) Chromatograms showing peak of standard (Kaempferol, Quercetin and Rutin). (B) Chromatograms showing peak of Kaempferol and Quercetin at 0 and 48 h of dark treatment different in AtMYB12OX line of tobacco. (C) Chromatograms showing peak of Rutin at 0 and 48 h of dark treatment different in AtMYB12OX line of tobacco.

Figure S3. Estimation of flavonol content in WT tobacco. Flavonol aglycones (kaempferol and quercetin) were measured in 10-d-old dark-grown WT tobacco seedlings and WT seedlings transferred to light for 3 h after growing 10 days in darkness. Statistical analysis was performed using two-tailed Student's t-test. Data are plotted as means $\pm$ SD (n=3). Error bars represent standard deviation. Asterisks indicate a significant difference, **P < 0.01, ***P < 0.001.

Figure S4. Sequence analysis of NtHY5 gene. (A) Gene structure of NtHY5 in comparison to AtHY5. The black boxes represent exons, lines represent introns and numbers represent the length of the exons and introns (bp). (B) Representation of NtHY5 conserved domain that is basic leucine zipper (bZIP) domain through conserved domain database.

Figure S5. Phenotypic analysis of different transgenic lines. (A) A schematic representation of NtHY5 construct under the control of CaMV35S promoter. (B) Representative photographs of rosette diameter of transgenic WT, *hy5* and *NtHY5;hy5* plants. (C) Measurement of seed size of *NtHY5;hy5* lines of Arabidopsis in comparison to WT and *hy5* plants. (D) Transverse section of stem to show lignin content in *NtHY5; hy5* in comparison to *hy5* and WT. Statistical analysis was performed using two-tailed Student's t-test. Data are plotted as means $\pm$ SD (n=10-12) for rosette diameter. Seed size, and lignin staining experiments. Error bars represent standard deviation. *hy5* was taken as control for significance. Asterisks indicate a significant difference, *P < 0.1, **P < 0.01.

Figure S6. Relative expression of MYB transcription factors associated with phenylpropanoid pathway. Expression analysis of *AtMYB12*, *AtMYB11* and *AtMYB111* through real time PCR in *NtHY5;hy5*, *hy5* and WT after growing for 7-d-dark condition and then transferred to light for 3 hours. Statistical analysis was performed using two-tailed Student's t-test. Tubulin was used as endogenous control to normalize the relative expression levels. Data are plotted as means $\pm$ SD (n=3). Dark grown seedlings were used as control for significance. Error bars represent standard deviation. Asterisks indicate a significant difference, **P < 0.01, ***P < 0.001.

Figure S7. Estimation of total chlorophyll and anthocyanin content. (A) Chlorophyll content in 30-d-old rosette leaves of WT, *hy5* and *NtHY5; hy5* plants. (B) Total Anthocyanin content in WT, *hy5* and *NtHY5OX;hy5* in 7-d-old light-grown seedlings. Statistical analysis was performed using two-tailed Student's t-test. Data are plotted as means $\pm$ SD (n=3). Error bars represent standard deviation. Asterisks indicate a significant difference, **P < 0.01, ***P < 0.001.

Figure S8. Position of gRNA in NtHY5 sequences. (A) Coding sequence of NtHY5 showing location of gRNA in red color, PAM sequences (NGG) are green in color. (B) Amino acid sequence of WT and *NtHY5^{CR}* edited plants.

Figure S9. Comparative analysis of HY5OX vs. *NtHy5^{CR}*. (A) Phenotype of WT, *NtHYR^{CR}* and NtHY5OX 10-d-dark grown seedling. (B) Measurement of seed size of WT, NtHY5OX and *NtHY5^{CR}* plants of tobacco. (C) Quantitative analysis of chlorophyll content of 1-month-old WT, NtHY5OX and *NtHY5^{CR}* plants of tobacco. Statistical analysis was performed using two-tailed

Student's t-test. Data are plotted as means $\pm$ SD (n=3). Error bars represent standard deviation. Asterisks indicate a significant difference, ***P < 0.001.

Figure S10. Detection of hydrogen peroxide by DAB staining in 10-d-old light grown WT, NtHY5OX and *NtHY5^{CR}* tobacco seedlings subjected to 200 mM salt stress for 3 hours. The statistical analysis was performed using two-tailed Student's t-tests. The data are plotted as means $\pm$ SD (n=3). The error bars represent standard deviations. The asterisks indicate significant difference, *P< 0.1; **P< 0.01; ***P< 0.001.

Table S1. Putative *cis*-acting elements in NtMYB12 promoter

Table S2. List of Primers used in the study

Figure legends

Figure 1. Light affect flavonoid biosynthesis in transgenic tobacco. (A),(C),(E) Relative transcript abundance of *AtCHS*, *AtCHI*, *AtFLS* and *AtDFR* that are involved in phenylpropanoid pathway in AtMYB12, AtMYB11 and AtMYB111 overexpressing transgenic lines respectively by real time PCR using RNA from the 10-d-old light grown seedlings transferred to dark for 24hr and 48hr. Tubulin was used as endogenous control to normalize the relative expression levels. **(B), (D), (F)** Effect of light over flavonol biosynthesis in different transgenic lines expressing AtMYB12, AtMYB11 and AtMYB111 respectively. Compounds (K, Kamepferol; Q, Quercitin; R; Rutin) were quantified using methanolic extracts of 10-d-old old seedlings of different transgenic lines through HPLC. Data are derived from the mean of three technical replicates. Error bars represent SE of means (n=3). The statistical analysis was performed using two-tailed Student's t-tests. The asterisks indicate significant difference, *P< 0.1; **P< 0.01; ***P< 0.001.

Figure 2. NtHY5, the light signaling component, alters flavonoid content in tobacco. (A) Phylogenetic tree of HY5 homologous proteins (sequence obtained from NCBI database) from across species of different families. The numbering at the nodes represents the number of trees that have the same topology as the consensus tree. NtHY5 is denoted through the red branch, and the scale bar indicates the branch length. **(B)** Amino acids alignment of NtHY5 and its homologs in Tomato and Arabidopsis using multallignment tool. The VPE motif and zinc finger domain is

boxed in green and red color respectively. **(C)** Phenotype of 10-d-old dark and light grown WT seedlings. **(D)** Expression analysis of phenylpropanoid pathway genes *NtCHS*, *NtCHI*, *NtFLS* and *NtDFR* in WT seedlings grown in light and dark for 10 days. **(E)** Quantitative estimation of flavonoids (K, Q and R) in 10-d-light and dark grown seedlings through HPLC. **(F)** Relative expression of NtHY5 in 10 light and dark grown seedlings. **(G)** Normalized expression of NtHY5 in different tissue [seedling, young (Y) leaf, mature (M) leaf, stem, root and flower]. Data are derived from the mean of three technical replicates. Tubulin was used as endogenous control to normalize the relative expression levels. Error bars represent SE of means (n=3). The statistical analysis was performed using two-tailed Student's t-tests. The asterisks indicate significant difference, **P<0.01; ***P<0.001.

Figure 3. NtHY5 interacts with NtMYB12 promoter. **(A)** Representation of the probes (G-BOXI, G-mut I, G-BOXII and G-mut II) designed upstream of the transcription start site of NtMYB12 promoter. G-BOX is altered by multiple base substitutions. **(B)** EMSA (Electrophoretic mobility shift assay) for the binding of 6X-His-HY5 (NtHY5) with core ACE-DIG element (HY5 binding site that is G-BOXI and G-BOXII) present in the LRE motifs of the NtMYB12 promoter. Upper and lower arrows indicate shift and free probe respectively. **(C), (D)** EMSA with competition between digoxigenin in labelled G-BOXI, unlabeled G-BOX I (cold), G-BOXII and unlabeled G-BOX II (cold) probe to bind to 6X-His-HY5 respectively. Superscript values 10X, 20X, 30X and 50X represent increasing amount of cold probe. Upper and lower arrows indicate shift and free probe respectively.

Figure 4. Complementation of Arabidopsis *hy5* mutant by NtHY5. **(A)** Phenotype of 7-d-old light-grown WT (wild type; Col-0), *hy5* and NtHY5OX; *hy5* seedlings (L1 and L2 are two independent transgenic lines overexpressing NtHY5 in *hy5* mutant background). **(B)** Relative expression of NtHY5 in 7-d-old light grown WT, *hy5*, L1 and L2. **(C)** Hypocotyl length of 7-d-old white light grown WT, *hy5*, L1 and L2 seedlings. **(D)** Root length of 7-d-old white light grown WT, *hy5*, L1 and L2. **(E)** Rosette diameter of 30-day-old WT, *hy5*, L1 and L2 plants. **(F)** Representative photographs of the seed size of WT, *hy5* and transgenic lines (L1 and L2). Bars = 1 mm. **(G)** Relative expression of phenylpropanoid pathway genes (*AtCHS*, *AtCHI*, *AtFLS* and *AtDFR*) in 7-d-light grown WT, *hy5*, L1 and L2 seedlings. **(H)** Quantitative estimation of

flavonol (Kaempferol and Quercetin) content in 7-d-old light grown WT, *hy5*, L1 and L2 seedlings through HPLC. **(I)** Expression analysis of *AtCHS*, *AtCHI*, *AtFLS* and *AtDFR* in WT, *hy5*, L1 and L2 seedlings grown for 5 days in dark followed by an exposure to light for 3hr. **(J)** Length of vascular bundle of 30-day-old stem of WT, *hy5*, L1 and L2 plants. Statistical analysis was performed using two-tailed Student's t-test. Error bars represent SE of means (n=3) For hypocotyl and root length (n=10-12). Tubulin was used as endogenous control to normalize the relative expression levels. Error bars represent standard deviation. Asterisks indicate a significant difference, *P < 0.1, **P < 0.01, ***P < 0.001. ns represent non-significant.

Figure 5. Overexpression of NtHY5 modulates gene expression and metabolite level in tobacco. **(A)** Relative expression of NtHY5 in 10-d-old light grown WT and NtHY5OX lines of tobacco (L1, L2 and L3). **(B)** Expression analysis of phenylpropanoid pathway genes (*NtCHS*, *NtCHI*, *NtFLS* and *NtDFR*) in 10-d-old light grown WT and NtHY5OX lines. **(C)** Quantification of flavonoids (K, Q and R) and polyphenolic compound (CGA) in 10-d-old light grown WT and NtHY5OX lines of tobacco using HPLC. **(D)** Representative photograph of flower color alterations in petals of NtHY5OX transgenic lines in compared with WT. **(E)** Total anthocyanin content in petals of WT and NtHY5OX transgenic lines. **(F)** Estimation of antioxidant activity by trolox equivalent in WT and NtHY5OX lines. TUBULIN was used as the endogenous control to normalize the relative expression levels. The statistical analysis was performed using two-tailed Student's t-tests. The data are plotted as means $\pm$ SD (n= 3). The error bars represent standard deviations. The asterisks indicate significant difference, *P< 0.1; **P< 0.01; ***P< 0.001.

Figure 6. CRISPR-Cas9-mediated knockout of NtHY5 alters gene expression and accumulation of metabolite content. **(A)** Representation of mutations in target region in three independent transgenic lines of tobacco. Dots indicate deleted nucleotides. Yellow highlight denotes gRNA region used for targeting the sequence. Red highlights denote indels in nucleotides. **(B)** Relative expression of NtHY5 in 10-d-old light grown WT and *NtHY5^{CR}* lines of tobacco (L1, L2 and L3). **(C)** Relative transcript abundance of phenylpropanoid pathway genes (*NtCHS*, *NtCHI*, *NtFLS* and *NtDFR*) in 10-d-old light grown WT and *NtHY5^{CR}* lines. **(D)** Quantification of flavonoids (K, Q and R) and polyphenolic compound (CGA) in 10-d-old light grown WT and *NtHY5^{CR}* lines of tobacco. **(E)** Quantification of Anthocyanin content in 10-d-old light grown WT and *NtHY5^{CR}* lines. **(F)** Quantification of antioxidant activity by trolox equivalent in 10-d-old light grown WT and *NtHY5^{CR}* lines. Tubulin was used as the endogenous

control to normalize the relative expression levels. The statistical analysis was performed using two-tailed Student's t-tests. The data are plotted as means $\pm$ SD (n=3). The error bars represent standard deviations. The asterisks indicate significant difference, *P<0.1; **P<0.01; ***P<0.001.

Figure 7. NtHY5OX and CRISPR–Cas9-derived knockout mutants show altered phenotypes and gene expression. **(A)** Representative image of 10-d-old light grown WT, NtHY5OX and *NtHY5^{CR}* seedlings grown on half-strength MS medium. Scale bars, 1 cm. **(B)** Measurement of root bending angle of 10-d-old light grown WT, NtHY5OX and *NtHY5^{CR}* seedlings. **(C), (D)** Hypocotyl length of 10-d-old light and dark grown WT, NtHY5OX and *NtHY5^{CR}* seedlings. **(E)** Root length of 10-d-old light grown WT, NtHY5OX and *NtHY5^{CR}* seedlings. **(F)** Quantitative analysis of seed size of WT, NtHY5OX and *NtHY5^{CR}* plants. **(G)** Quantitative estimation of Number of seeds per 5 mg of WT, NtHY5OX and *NtHY5^{CR}* plants. **(H), (I)** Relative expression of NtHY5 and phenylpropanoid pathway genes (*NtCHS*, *NtCHI*, *NtFLS* and *NtDFR*) in 10-d-old light grown seedlings transferred to dark for 3h. **(J)** Representative image of 40-day-old WT, NtHY5OX and *NtHY5^{CR}* plants showing plant heights. **(K)** Data presenting difference in height of 40-day-old WT, NtHY5OX and *NtHY5^{CR}* plants. Tubulin was used as the endogenous control to normalize the relative expression levels. The statistical analysis was performed using two tailed Student's t-tests. The data are plotted as means $\pm$ SD (n= 3). For measurement of root bending angle, root length, hypocotyls length and seed size (n=10-12). The error bar represents standard deviations. The asterisks indicate significant difference, *P<0.1; **P<0.01; 0.001.

Figure 8. Effect of salt stress on different transgenic. Fully expanded leaves of 30-d-old plants to floated on 6 ml of 200 mM NaCl solution (for salt stress) for 3 days. **(A), (B)** The phenotypic (chlorosis) changes due to salt treatment on leaf discs were assessed on 0 and 3 days respectively. **(C)** Quantification of chlorophyll content of WT, NtHY5OX and *NtHY5^{CR}* leaves after 3day of salt stress. **(D)** Detection of Superoxide ($O_2^{\cdot-}$) by NBT staining in 10-d-old light grown WT, NtHY5OX and *NtHY5^{CR}* tobacco seedlings subjected to 200 mM salt stress for 3 hours. The statistical analysis was performed using two-tailed Student's t-tests. The data are plotted as means $\pm$ SD (n=3). The error bars represent standard deviations. The asterisks indicate significant difference, *P<0.1; **P<0.01; ***P<0.001.

Figure 9. Proposed working model for the role of NtHY5 and NtMYB12 in regulating the expression of flavonoid pathway genes during light exposure. We propose that NtHY5 positively regulates the expression of NtMYB12 by directly binding to promoter of NtMYB12 through G-BOX I and G-BOX II in turn NtMYB12 directly binds to the MRE (MYB recognition element) in the promoters of flavonoid pathway genes and enhances flavonol accumulation whereas NtHY5 also binds independently to ACE element present on the promoter of flavonoid pathway genes and regulate their expression. Thus, together NtHY5 and NtMYB12 lead to flavonol production. Flavonol ultimately leads to salt tolerance by inhibiting ROS production. Arrowheads and tees indicate positive and negative regulation respectively.

Parsed Citations

Agati G, Azzarello E, Pollastri S, Tattini M (2012) Flavonoids as antioxidants in plants: location and functional significance. Plant Sci 196: 67–76.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ahanger MA, Tomar NS, Tittal M, Argal S, Agarwal RM (2017) Plant growth under water/salt stress: ROS production; antioxidants and significance of added potassium under such conditions. Physiol Mol Biol Plants 23:731-744.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ang LH, Chattopadhyay S, Wei N, Oyama T, Okada K, Batschauer A, Deng XW (1998) Molecular interaction between COP1 and HY5 defines a regulatory switch for light control of Arabidopsis development. Molecular Cell 1: 213–222.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bhatia C, Pandey A, Gaddam SR, Hoecker U, Trivedi PK (2018) Low temperature-enhanced flavonol synthesis requires light-associated regulatory components in Arabidopsis thaliana. Plant and Cell Physiol. 59:2099–2112.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bhatia C, Gaddam SR, Pandey A, Trivedi PK (2021) COP1 mediates light-dependent regulation of flavonol biosynthesis through HY5 in Arabidopsis. Plant Science, 303: 110760.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bhatnagar A, Singh S, Khurana JP (2020) HY5-COP1: the central module of light signaling pathway. J. Plant Biochem. Biotechnol. 29, 590–610.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Brown BA, Jenkins GI (2008) UV-B signaling pathways with different fluence-rate response profiles are distinguished in mature Arabidopsis leaf tissue by requirement for UVR8, HY5, and HYH. Plant Physiol. 146:576–588.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Brown DE, Rashotte AM, Murphy AS, Normanly J, Tague BW, Peer WA, Taiz L, Muday GK (2001) Flavonoids act as negative regulators of auxin transport in vivo in Arabidopsis. Plant Physiol 126: 524–535.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Burbulis IE, Winkel-Shirley B (1999) Interactions among enzymes of the Arabidopsis flavonoid biosynthetic pathway. Proc Natl Acad Sci USA 96: 12929-12934.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Burman N, Bhatnagar A, Khurana JP (2018) OsbZIP48, a HY5 transcription factor ortholog, exerts pleiotropic effects in light-regulated development Plant Physiol. 176: 1262-1285.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cai W, Gu X, Tang J (2010) Extraction, purification, and characterisation of the flavonoids from Opuntia milpa alta skin. Czech J. Food Sci. 28: 108–116.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chattopadhyay S, Ang LH, Puente P, Deng XW, Wei N (1998) Arabidopsis bZIP protein HY5 directly interacts with light-responsive promoters in mediating light control of gene expression. Plant Cell 10: 673–683.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chen S, Wu F, Li Y, Qian Y, Pan X, Li F (2019) NtMYB4 and NtCHS1 are critical factors in the regulation of flavonoid biosynthesis and are involved in salinity responsiveness. Front Plant Sci 10: 178.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chen S, Wu F, Li Y, Qian Y, Pan X, Li F, Wang Y, Wu Z, Fu C, Lin H, Yang A (2019) NtMYB4 and NtCHS1 are critical factors in the regulation of flavonoid biosynthesis and are involved in salinity responsiveness. Front Plant Sci 10: 178.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chory J (2010) Light signal transduction: an infinite spectrum of possibilities. Plant J 61: 982-991.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cluis CP, Mouchel CF, Hardtke CS (2004) The Arabidopsis transcription factor HY5 integrates light and hormone signaling pathways. 38: 332-347.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Czemmel S, Heppel SC, Bogs J (2012) R2R3 MYB transcription factors: key regulators of the flavonoid biosynthetic pathway in grapevine. Protoplasma 249 Suppl 2:S109-18.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Docheva M, Dagnon S, Statkova-Abeghe S (2014) Flavonoid content and radical scavenging potential of extracts prepared from tobacco cultivars and waste. Nat Prod Res. 28: 1328-34.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

- Doughty J, Aljabri M, Scott RJ (2014) Flavonoids and the regulation of seed size in Arabidopsis. Biochem Soc Trans 42: 364–369.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Emiliani J, Grotewold E, Falcone Ferreyra ML, Casati P (2013) Flavonols protect Arabidopsis plants against UV-B deleterious effects. Mol Plant 6:1376-1379.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Falcone Ferreyra ML, Rius SP, Casati P (2012) Flavonoids: biosynthesis, biological functions, and biotechnological applications. Front in plant sci 3: 222.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Funk M, Weisshaar B, Ulm R (2010) The Arabidopsis bZIP transcription factor HY5 regulates expression of the PFGs1/MYB12 gene in response to light and ultraviolet-B radiation. Plant Cell Environ 33: 88–103.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Fu B, Ji X, Zhao M, He F, Wang X, Wang Y, Liu P, Niu L (2016) The influence of light quality on the accumulation of flavonoids in tobacco (Nicotiana tabacum L.) leaves. J Photochem Photobiol B 162:544-549.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Gangappa SN, Botto JF (2016) The multifaceted roles of HY5 in plant growth and development. Molecular Plant 9: 1353–1365.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Gou JY, Felippes FF, Liu CJ, Weigel D, Wang JW (2011) Negative regulation of anthocyanin biosynthesis in Arabidopsis by a miR156-targeted SPL transcription factor. The Plant Cell 23: 1512–1522.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Hartmann U, Sagasser M, Mehrtens F, Stracke R, Weisshaar B (2005) Differential combinatorial interactions of cis-acting elements recognized by R2R3-MYB, BZIP, and BHLH factors control light-responsive and tissue-specific activation of phenylpropanoid biosynthesis genes. Plant Molecular Biology 57: 155-171.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Heijde M, Ulm R (2012) UV-B photoreceptor-mediated signalling in plants. Trends Plant Sci. 17:230–237.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Hichri I, Barrieu F, Bogs J, Kappel C, Delrot S, Lauvergeat V (2011) Recent advances in the transcriptional regulation of the flavonoid biosynthetic pathway. J Exp Bot 62: 2465-2483.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Holton TA, Edwina CC (1995) Cornish ECGenetics and biochemistry of anthocyanin biosynthesis. Plant Cell 7: 1071–1083.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Horsch RB (1985) A simple method for transferring genes into plants. Science 227: 1229–123.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Huang X, Ouyang X, Yang P, Lau OS, Li G, Li J, Chen H, Deng XW (2012) Arabidopsis FHY3 and HY5 positively mediate induction of COP1 transcription in response to photomorphogenic UV-B light. Plant Cell 24: 4590–4606.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Jakoby M, Weisshaar B, Dröge-Laser W, Vicente-Carbajosa J, Tiedemann J, Kroj T, Parcy F (2002) bZIP transcription factors in Arabidopsis. Trends Plant Sci 7:106-111.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Jiang L, Wang Y, Li QF, Björn LO, He JX, Li SS (2012) Arabidopsis STO/ BBX24 negatively regulates UV-B signaling by interacting with COP1 and repressing HY5 transcriptional activity. Cell Res 22: 1046–1057.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Jiang, W, Yin Q, Wu R, Zheng G, Liu J, Dixon RA (2015) Role of a chalcone isomerase-like protein in flavonoid biosynthesis in Arabidopsis thaliana. J Exp Bot 66: 7165–7179.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Jing Y, Zhang D, Wang X, Tang W, Wang W, Huai J, Xu G, Chen D, Li Y, Lin R (2013) Arabidopsis chromatin remodeling factor PICKLE interacts with transcription factor HY5 to regulate hypocotyl cell elongation. Plant Cell 25: 242–256.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Kim JI, Dolan W, Anderson N, Chapple C (2015) Indole glucosinolate biosynthesis limits phenylpropanoid accumulation in Arabidopsis thaliana. Plant Cell 27: 1529–1546.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Koornneef M, Rolff E, Spruit CJP (1980) Genetic control of light inhibited hypocotyl elongation in Arabidopsis thaliana (L.) Heynh. Z Pflanzenphysiol 100: 147–160.**
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Kumar S, Nei M, Dudley J, Tamura K (2008) MEGA: a biologist-centric software for evolutionary analysis of DNA and protein sequences. Brief. Bioinform. 9: 299–306.**

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lau OS, Deng XW (2012) The photomorphogenic repressors COP1 and DET1: 20 years later. Trends Plant Sci. 17: 584–593.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lee J, He K, Stolc V, Lee H, Figueroa P, Gao Y, Zhao H, Lee I, Deng XW (2007) Analysis of transcription factor HY5 genomic binding sites revealed its hierarchical role in light regulation of development. Plant Cell 19: 731–749.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lescot M, Dehais G, Thisj K, Marchal Y, Peer Y, Rouze S, Rombauts S (2020) RombautsPlantCARE, a database of plant cis-acting regulatory elements and a portal to tools for in silico analysis of promoter sequences. Nucleic Acids Res 30: 325-327.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li H, Yang Z, Zeng Q, Wang S, Huang Y, Xin Y, He N (2020) Abnormal expression of bHLH3 disrupts a flavonoid homeostasis network, causing differences in pigment composition among mulberry fruits. Hortic Res. 7: 83.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li J, Yang L, Jin D, Nezames, CD, Terzaghi W, Deng XW (2013) UV-B-induced photomorphogenesis in Arabidopsis. Protein Cell 4:485–492.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li S, Wang W, Gao J, Yin K, Wang R, Wang C, Petersen M, Mundy J, Qui J (2016) MYB75 phosphorylation by MPK4 is required for light-Induced anthocyanin accumulation in Arabidopsis. Plant Cell 28: 2866–2883.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li Y, Chen M, Wang S L, Ning J, Ding XH, Chu ZH (2015) AtMYB11 regulates caffeoylquinic acid and flavonol synthesis in tomato and tobacco. Plant Cell 122: 309–319.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu X, Qin T, Ma Q (2013) Light-regulated hypocotyl elongation involves proteasome-dependent degradation of the microtubule regulatory protein WDL3 in Arabidopsis. Plant Cell 25:1740-1755.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu ZY, Wang PL, Zhang TQ, Li YB, Wang YY, Gao CQ (2018) Comprehensive analysis of BpHSP genes and their expression under heat stresses in Betula platyphylla. Environ Exp Bot 152: 167–176.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liva KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. Methods 25, 402–408

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lur HS, Setter TL (1993) Role of auxin in maize endosperm development: timing of nuclear-DNA endoreduplication, zein expression, and cytokinin. Plant Physiol 103: 273–280.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Martinez V, Mestre TC, Rubio F, Moreno AD, Mittler R (2016) Accumulation of flavonols over hydroxycinnamic acids favors oxidative damage protection under abiotic stress. Front Plant Sci 7: 838.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

McCarthy RL, Zhong R, Ye ZH (2009) MYB83 is a direct target of SND1 and acts redundantly with MYB46 in the regulation of secondary cell wall biosynthesis in Arabidopsis. Plant Cell Physiol 50: 1950–1964.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mehrtens F, Kranz H, Bednarek P, Weisshaar B (2005) The Arabidopsis transcription factor MYB12 is a flavonol-specific regulator of phenylpropanoid biosynthesis. Plant Physiology 138: 1083-1096.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Michael R, Ranjan A, Kumar RS, Parthak PK, Trivedi PK (2020) Light-regulated expression of terpene synthase gene, AtTPS03, is controlled by the bZIP transcription factor, HY5, in Arabidopsis thaliana. Biochem Biophys Res Commun 529: 437-443.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Misra P, Pandey A, Tiwari M, Chandrashekar K, Sidhu OP, Asif MH, Chakrabarty D, Singh PK, Trivedi PK (2010) Modulation of transcriptome and metabolome of tobacco by Arabidopsis transcription factor, AtMYB12, leads to insect resistance. Plant Physiol. 152: 2258–2268.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Muhlemann JK, Younts TLB, Muday GK (2018) Flavonols control pollen tube growth and integrity by regulating ROS homeostasis during high temperature stress. Proc. Natl. Acad. Sci. USA 115: E11188-E11197.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nawkar GM, Kang CH, Maibam P, Park JH, Jung, YJ, Chae HB, Chi, YH, Jung IJ, Kim, WY, Yun DJ (2017) HY5, a positive regulator of light signaling, negatively controls the unfolded protein response in Arabidopsis. Proc. Natl. Acad. Sci. USA 114:2084–2089.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nguyen NH, Jeong CY, Kang GH, Yoo SD, Hong SW, Lee H (2015) MYBD employed by HY5 increases anthocyanin accumulation via repression of MYBL2 in Arabidopsis. Plant J 84: 1192-1205.

Google Scholar: [Author Only Title Only Author and Title](#)

Niggeweg R, Michael AJ, Martin C (2004) Engineering plants with increased levels of the antioxidant chlorogenic acid. Nat. Biotechnol. 22: 746–754.

Google Scholar: [Author Only Title Only Author and Title](#)

Niranjana A (2011) Development and optimization of HPLC-PDA-MS-MS method for simultaneous quantification of three classes of flavonoids in legume seeds, vegetables, fruits and medicinal plants. J. Liq. Chromatogr. Relat. Technol 34, 1729–1742

Google Scholar: [Author Only Title Only Author and Title](#)

Nishimura R, Ohmori M, Fujita H, Kawaguchi M (2002) A Lotus basic leucine zipper protein with a RING-finger motif negatively regulates the developmental program of nodulation. Proc Natl Acad Sci USA 99: 15206–15210.

Google Scholar: [Author Only Title Only Author and Title](#)

Nishimura R, Ohmori M, Fujita H, Kawaguchi M (2002) A Lotus basic leucine zipper protein with a RING finger motif negatively regulates the developmental program of nodulation. Proc. Natl Acad Sci 99: 15206-15210.

Google Scholar: [Author Only Title Only Author and Title](#)

Osterlund MT, Hardtke CS, Wei N, Deng XW (2000) Targeted destabilization of HY5 during light-regulated development of Arabidopsis. Nature 405: 462–466.

Google Scholar: [Author Only Title Only Author and Title](#)

Oyama T, Shimura Y, Okada K (1997) The Arabidopsis HY5 gene encodes a bZIP protein that regulates stimulus-induced development of root and hypocotyl. Genes Dev 11: 2983–2995.

Google Scholar: [Author Only Title Only Author and Title](#)

Palaniswamy SK, James S, Sun H, Lamb RS, Davuluri RV, Grotewold E (2006). AGRIS and AtRegNet: a platform to link cis-regulatory elements and transcription factors into regulatory networks. Plant Physiol 140: 818-829.

Google Scholar: [Author Only Title Only Author and Title](#)

Pandey A, Misra P, Bhambhani S, Bhatia C, Trivedi PK (2014) Expression of Arabidopsis MYB transcription factor, AtMYB111, in tobacco requires light to modulate flavonol content. Sci Rep 4: 5018.

Google Scholar: [Author Only Title Only Author and Title](#)

Pandey A, Misra P, Chandrashekhar K, Trivedi PK (2012) Development of AtMYB12-expressing transgenic tobacco callus culture for production of rutin with biopesticidal potential. Plant Cell Rep 31: 1867–1876.

Google Scholar: [Author Only Title Only Author and Title](#)

Pandey A, Misra P, Trivedi PK (2015) Constitutive expression of Arabidopsis MYB transcription factor, AtMYB11, in tobacco modulates flavonoid biosynthesis in favor of flavonol accumulation. Plant Cell Rep 34: 1515–1528.

Google Scholar: [Author Only Title Only Author and Title](#)

Peer WA, Murphy AS (2007) Flavonoids and auxin transport: modulators or regulators? Trends Plant Sci 12: 556–563.

Google Scholar: [Author Only Title Only Author and Title](#)

Quail PH (2002) Phytochrome photosensory signalling networks. Nat. Rev. Mol. Cell Biol. 3:85–93.

Google Scholar: [Author Only Title Only Author and Title](#)

Ram H, Chattopadhyay S (2013) Molecular interaction of bZIP domains of GBF1, HY5 and HYH in Arabidopsis seedling development. Plant Signal Behav 8: e22703.

Google Scholar: [Author Only Title Only Author and Title](#)

Saito K, Yonekura-Sakakibara K, Nakabayashi R, Higashi Y, Yamazaki M, Tohge T, Fernie AR (2013) The flavonoid biosynthetic pathway in Arabidopsis: structural and genetic diversity. Plant Physiol Biochem 72: 21-34.

Google Scholar: [Author Only Title Only Author and Title](#)

Scalbert A, Johnson I, Saltmarsh M (2005) Polyphenols: antioxidants and beyond. Am J Clin Nutr 81: 215–217.

Google Scholar: [Author Only Title Only Author and Title](#)

Schulz E, Tohge T, Zuther E, Fernie AR, Hinch DK (2015). Natural variation in flavonol and anthocyanin metabolism during cold acclimation in Arabidopsis thaliana accessions. Plant, Cell & Environment 38: 1658- 1672.

Google Scholar: [Author Only Title Only Author and Title](#)

Schulz E, Tohge T, Zuther E, Fernie AR, Hinch DK (2016) Flavonoids are determinants of freezing tolerance and cold acclimation in Arabidopsis thaliana. Scientific Reports 6: 34027.

Google Scholar: [Author Only Title Only Author and Title](#)

Sharma, D, Tiwari M, Pandey A, Bhatia C, Sharma A, Trivedi, PK 2016 MicroRNA858 is a potential regulator of phenylpropanoid pathway and plant development. Plant Physiol 171, 944-959.

Sharma A, Badola PK, Bhatia C, Sharma D, Trivedi PK (2020) Primary transcript of miR858 encodes regulatory peptide and controls flavonoid biosynthesis and development in Arabidopsis. Nat. Plants 6: 1262–1274.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sharma A, Badola PK, Gautam H, Gaddam SR, Trivedi PK (2021) HY5 regulates light-dependent expression and accumulation of miR858a encoded peptide, miPEP858a. Biochem Biophys Res Commun <https://doi.org/10.1016/j.bbrc.2021.12.020>

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sharma Y, Nagar A, Srivastava NS, Bala K (2017) Antioxidant activity of polyphenolic flavonoid of stem of Nicotiana tabacum. Am J Drug Disc Devel 25: 32.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Shin DH, Choi M, Kim K, Bang G, Cho M, Choi SB, Choi G, Park YI (2013). HY5 regulates anthocyanin biosynthesis by inducing the transcriptional activation of the MYB75/PAP transcription factor in Arabidopsis. FEBS Lett 587,1543-1547.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sibout R, Su P, Hettiarachchi C, Holm M, Muday GK, Hardtke CS (2006) Opposite root growth phenotypes of hy5 versus hy5 hyh mutants correlate with increased constitutive auxin signaling. PLoS Genet 2: e202.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Song YH, Yoo CM, Hong AP, Kim SH, Jeong HJ, Shin SY, Kim HJ, Yun DJ, Lim CO, Bahk JD (2008) DNA-binding study identifies C-box and hybrid C/G-box or C/A-box motifs as high-affinity binding sites for STF1 and LONG HYPOCOTYL5 proteins. Plant Physiol 146: 1862-1877.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Song ZP, Fan NB, Jiao G Z, Liu MH, Wang XY, Jia HF (2019) Overexpression of OsPT8 increases auxin content and enhances tolerance to high-temperature stress in Nicotiana tabacum. Genes 10: 809.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Song Z, Luo Y, Wang W, Fan N, Wang D, Yang C and Jia H (2020) NtMYB12 positively regulates flavonol biosynthesis and enhances tolerance to low Pi stress in Nicotiana tabacum. Front Plant Sci 10: 1683.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Stracke R, Favory JJ, Gruber H, Bartelniewoehner L, Bartels S, Binkert M, Funk M, Weissshaar B, Ulm R (2010a) The Arabidopsis bZIP transcription factor HY5 regulates expression of the PFG1/MYB12 gene in response to light and ultraviolet-B radiation. Plant Cell Environ 33: 88–103.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Stracke R, Ishihara H, Huep G, Barsch A, Mehrtens F, Niehaus K, Weissshaar (2007) Differential regulation of closely related R2R3-MYB transcription factors controls flavonol accumulation in different parts of the Arabidopsis thaliana seedling. Plant J 50: 660–677.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Truong HA, Lee S, Trinh CS, Lee WJ, Chung EH, Hong SW, Lee H (2021) Overexpression of the HDA15 Gene Confers Resistance to Salt Stress by the Induction of NCED3, an ABA Biosynthesis Enzyme. Front Plant Sci 12:640443

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ulm R, Baumann A, Oravecz A, Máté Z, Adám E, Oakeley EJ, Schäfer E, Nagy F (2004) Genome-wide analysis of gene expression reveals function of the bZIP transcription factor HY5 in the UV-B response of Arabidopsis. Proc Natl Acad Sci USA 101: 1397–1402.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vanhaelewyn L, Schumacher P, Poelman D, Fankhauser C, Van Der Straeten D, Vandenbussche F. (2016) REPRESSOR OF ULTRAVIOLET-B PHOTOMORPHOGENESIS function allows efficient phototropin mediated ultraviolet-B phototropism in etiolated seedlings. Plant Science 252: 215-221.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ververidis F, Trantas E, Douglas C, Vollmer G, Kretschmar G, Panopoulos N (2007) Biotechnology of flavonoids and other phenylpropanoid-derived natural products. Part I: Chemical diversity, impacts on plant biology and human health. Biotechnol J 2:1214-1234.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vogt T (2010) Phenylpropanoid biosynthesis. Mol. Plant 3: 2–20.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wan J, Zhang P, Wang R, Sun L, Wang W, Zhou H and Xu J (2018) UV-B Radiation Induces Root Bending Through the Flavonoid-Mediated Auxin Pathway in Arabidopsis. Front. Plant Sci 9:618

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wei YZ, Hu FC, Hu GB, Li XJ, Huang XM, Wang HC (2011) Differential expression of anthocyanin biosynthetic genes in relation to anthocyanin accumulation in the pericarp of Litchi chinensis Sonn. PLoS ONE 6: e19455.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Weller JL, Hecht V, Vander Schoor JK, Davidson SE, Ross JJ (2009) Light regulation of gibberellin biosynthesis in pea is mediated through the COP1/HY5 pathway. Plant Cell 21: 800–813.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Winkel-shirley B (2002) Biosynthesis of flavonoids and effects of stress. Curr Opin Plant Biol 5:218-223.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wong SP, Leong LP, Koh JHW (2006). Antioxidant activities of aqueous extracts of selected plants. Food Chem. 99: 775–783.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Xing HL, Dong L, Wang ZP, Zhang HY, Han CY, Liu B, Wang XC, Chen QJ (2014). CRISPR/Cas9 toolkit for multiplex genome editing in plants. BMC Plant Biol 14: 327.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yadav A, Singh D, Lingwan M, Yadukrishnan P, Masakapalli, SK, Datta S (2020). Light signaling and UV-B-mediated plant growth regulation. J. Integr. Plant Biol 62: 1270-1292.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yamawaki S, Yamashino T, Nakanishi H, Mizuno T (2011) Functional characterization of HY5 homolog genes involved in early light-signaling in *Physcomitrella patens*. Biosci Biotechnol Biochem 75: 1533–15.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yin, R, Han K, Heller W, Albert A, Dobrev PI, Schaffner AR (2014) kaempferol 3-O-rhamnoside-7-O-rhamnoside is an endogenous flavonol inhibitor of polar auxin transport in *Arabidopsis* shoots. New Phytol. 201: 466–475.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yoo CM, Park JM, Ryu GR, Goekjian VH, Nagao RT, Hong JC (1998) STF1 is a novel TGACG-binding factor with a zinc-finger motif and a bZIP domain which heterodimerizes with GBF proteins. Plant J 15:199-209.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhai R, Wang Z, Zhang S, Meng G, Song, Wang Z (2016) Two MYB transcription factors regulate flavonoid biosynthesis in pear fruit (*Pyrus bretschneideri* Rehd.). J Exp Bot 67: 1275–1284.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhai R, Zhao Y X, Wu M, Yang J, Li XY, Liu HT (2019) The MYB transcription factor PbMYB12b positively regulates flavonol biosynthesis in pear fruit. BMC Plant Biol 19 (1).

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang H, He H, Wang X, Wang X, Yang X, Li L, Deng XW (2011a) Genome-wide mapping of the HY5-mediated gene networks in *Arabidopsis* that involve both transcriptional and post-transcriptional regulation. Plant J 65: 346–358.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang Y, Li C, Zhang J, Wang J, Yang J, Lv Y (2017) Dissection of HY5/HYH expression in *Arabidopsis* reveals a root autonomous HY5-mediated photomorphogenic pathway. PLoS ONE 12(7): e0180449.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhao Q, Ren YR, Wang QJ, Wang XF, You CX, Hao YJ (2016) Ubiquitination-related MdbT scaffold proteins target a bHLH transcription factor for iron homeostasis. Plant Physiol 172, 1973–1988.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zheng X, Wu S, Zhai H, Zhou P, Song M, Su L, Xi Y, Li Z, Cai Y, Meng F (2013) *Arabidopsis* phytochrome B promotes SPA1 nuclear accumulation to repress photomorphogenesis under far-red light. Plant Cell 25: 115–133.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhong R, Lee C, Zhou J, McCarthy RL, Ye ZH (2008) A battery of transcription factors involved in the regulation of secondary cell wall biosynthesis in *Arabidopsis*. Plant Cell 20: 2763–2782.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhou, Z., Schenke, D., Miao, Y. and Cai, D (2017) Investigation of the crosstalk between the flg22 and the UV-B-induced flavonol pathway in *Arabidopsis thaliana* seedlings. Plant Cell Environ 40: 453–458.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)