A Large-Scale Investigation of White Matter Microstructural Associations with Reading Ability

Steven L. Meisler *

Program in Speech and Hearing Bioscience and Technology

Harvard University

43 Vassar Street, Bldg. 46, Room 4033

Cambridge, MA 02139, USA

`smeisler@g.harvard.edu`

John D.E. Gabrieli

Department of Brain and Cognitive Sciences and McGovern Institute for Brain Research

Massachusetts Institute of Technology

43 Vassar Street, Bldg. 46, Room 4033

Cambridge, MA 02139, USA

`gabrieli@mit.edu`

*: Corresponding Author

January 6, 2022

Abstract

Reading involves the functioning of a widely distributed brain network, and white matter tracts are responsible for transmitting information between constituent network nodes. Several studies have analyzed fiber bundle microstructural properties to shed insights into the neural basis of reading abilities and disabilities. Findings have been inconsistent, potentially due to small sample sizes and varying methodology. To address this, we analyzed a large data set of 686 children ages 5-18 using state-of-the-art neuroimaging acquisitions and processing techniques. We searched for associations between fractional anisotropy (FA) and single-word and single-nonword reading skills in children with diverse reading abilities across multiple tracts previously thought to contribute to reading. We also looked for group differences in tract FA between typically reading children and children with reading disabilities. FA of the white matter increased with age across all participants. There were no significant correlations between overall reading abilities and tract FAs across all children, and no significant group differences in tract FA between children with and without reading disabilities. There were associations between FA and nonword reading ability in older children (ages 9 and above). Higher FA in the right superior longitudinal fasciculus (SLF) and left inferior cerebellar peduncle (ICP) correlated with better nonword reading skills. These results suggest that letter-sound correspondence skills, as measured by nonword reading, are associated with greater white matter coherence among older children in these two tracts, as indexed by higher FA.

Keywords— Reading, DTI, Fractional anisotropy, White matter

1 Introduction

The development of reading skills is vital for progress in education and communication. Reading disability is the most prevalent learning disability and is characterized by difficulty with word reading accuracy and/or fluency (Roongpraiwan et al., 2002; Lyon et al., 2003). Multiple studies have reported structural and functional brain differences in children and adults with reading disability (Pugh et al., 2000; Eckert, 2004; Maisog et al., 2008; Linkersdörfer et al., 2012; Cattinelli et al., 2013; Richlan et al., 2013; Jednorog et al., 2015), including differences in white matter pathways as measured by diffusion tensor imaging (DTI) (Vandermosten et al., 2012b). Studies of this nature often report the relationship between reading scores and tract fractional anisotropy (FA), which quantifies how direction-dependent water movement is in a given area (Basser and Pierpaoli, 1996; Hagmann et al., 2006) and has been interpreted as an index of white matter coherence (Beaulieu, 2002). Such associations may be important to identify because these fiber bundles of myelinated axons connect the gray matter regions that support reading, and especially the left-hemisphere reading network comprised of the angular gyrus, precuneus, middle temporal gyrus, superior temporal gyrus (including Wernicke’s area), fusiform gyrus (including the visual word form area), and inferior frontal gyrus (including Broca’s area) (Cattinelli et al., 2013; Wandell and Yeatman, 2013; Murphy et al., 2019).

Studies of white matter microstructural properties as they relate to reading abilities in children and adults have yielded interesting, albeit sometimes inconsistent, results. The most commonly implicated tract in DTI studies of reading skills is the superior longitudinal fasciculus (SLF), particularly in the left hemisphere, which connects frontal and temporoparietal brain regions (Wang et al., 2016). Multiple studies have found that higher FA in left and/or right SLF was associated with better reading outcomes, whether that manifested from

group comparisons between dyslexic and typically reading individuals (Richards et al., 2008; Steinbrink et al., 2008; Carter et al., 2009; Marino et al., 2014) or correlations with reading test scores on a continuous scale (Steinbrink et al., 2008; Carter et al., 2009; Hoeft et al., 2011; Feldman et al., 2012; Lebel et al., 2013; Zhang et al., 2014; Horowitz-Kraus et al., 2015; Borchers et al., 2019). However, other studies that have investigated the SLF, whether from a whole-brain or targeted approach, have failed to replicate this (Odegard et al., 2009; Welcome and Joanisse, 2014; Arrington et al., 2017), and a few studies have even reported negative FA-reading associations in the SLF (Carter et al., 2009; Frye et al., 2011).

The temporal sub-component of the SLF, the arcuate fasciculus (AF), has also been linked to reading performance (Rauschecker et al., 2009), and has often been analyzed separately from other SLF components due to its unique contributions to the language network (Catani et al., 2005). Similarly, FA reductions in dyslexia have been reported in the left AF (Klingberg et al., 2000; Deutsch et al., 2005; Vandermosten et al., 2012a; Marino et al., 2014; Christodoulou et al., 2017; Su et al., 2018), and its FA has been positively associated with reading skills (Klingberg et al., 2000; Deutsch et al., 2005; Yeatman et al., 2012a; Horowitz-Kraus et al., 2014; Christodoulou et al., 2017; Borchers et al., 2019). However, bilateral AF regions of higher FA in dyslexia have been identified (Žarić et al., 2018), and separate studies have reported negative FA-reading associations in the left AF (Yeatman et al., 2012a; Christodoulou et al., 2017; Huber et al., 2018).

Two fiber bundles that run under the SLF, the inferior longitudinal fasciculus (ILF) and inferior fronto-occipital fasciculus (IFO), serve to connect occipital and temporal-occipital areas to anterior temporal and frontal regions, respectively (Martino et al., 2010; Herbet et al., 2018), and have been identified as candidate reading tracts (Vandermosten et al., 2012a; Yeatman et al., 2013). The left ILF has exhibited increased FA in typically developing readers

compared to dyslexic readers (Steinbrink et al., 2008; Marino et al., 2014; Su et al., 2018), and bilateral ILF FA has been positively related to reading performance (Steinbrink et al., 2008; Odegard et al., 2009; Feldman et al., 2012; Yeatman et al., 2012a; Lebel et al., 2013; Horowitz-Kraus et al., 2014; Zhang et al., 2014; Horowitz-Kraus et al., 2015). However, a few studies have found negative associations between FA and reading scores in the left ILF (Yeatman et al., 2012a; Huber et al., 2018), and one study has found increased left ILF FA in dyslexic individuals compared to their typically reading counterparts (Banfi et al., 2019). While studies have only identified positive correlations between FA and reading in bilateral IFO (Steinbrink et al., 2008; Odegard et al., 2009; Feldman et al., 2012; Lebel et al., 2013; Welcome and Joanisse, 2014; Zhang et al., 2014; Arrington et al., 2017), only a single study has found a reduction in left IFO FA in dyslexia (Steinbrink et al., 2008). It is worth noting that other investigations have also yielded null results in these tracts (Klingberg et al., 2000; Frye et al., 2011; Borchers et al., 2019).

The uncinate fasciculus (UF) is thought to contribute to the ventral orthographic pathways of reading, connecting temporal and orbitofrontal regions (Catani et al., 2002; Schlaggar and McCandliss, 2007). Despite reports of positive associations between FA and reading skills in bilateral UF (Odegard et al., 2009; Feldman et al., 2012; Welcome and Joanisse, 2014; Zhang et al., 2014; Arrington et al., 2017), the only significant group difference in FA that has been reported favored higher FA in dyslexia (Arrington et al., 2017). The same study also reported a negative correlation between FA and one of their reading measures in the right UF (Arrington et al., 2017). The splenium of the corpus callosum is also thought to contribute to reading, as it subserves interhemispheric communication between visual cortices (Putnam et al., 2010). Opposing results for both group (Frye et al., 2008; Marino et al., 2014) and

continuous analyses (Frye et al., 2008; Odegard et al., 2009; Feldman et al., 2012; Lebel et al., 2013; Zhang et al., 2014; Huber et al., 2018) have been reported.

One theory of the etiology of dyslexia is the cerebellar hypothesis, which implicates cerebellar dysfunction in deficits of procedural learning and reading fluency (Nicolson et al., 2001; Nicolson and Fawcett, 2007; Stoodley and Stein, 2011). To this end, the contributions of the superior (SCP), inferior (ICP), and middle (MCP) cerebellar peduncles to reading have been investigated. The SCP contains efferent fibers that connect the deep cerebellum to inferior prefrontal regions involved in reading (Bruckert et al., 2020). The ICP and MCP contain primarily afferent fibers which connect the brainstem with the cerebellum. Such connections may facilitate automation of articulatory and oculomotor control (Bruckert et al., 2020). Bilateral SCP FA has exhibited negative relations to reading skill (Travis et al., 2015; Bruckert et al., 2020), while the left ICP has shown both positive (Borchers et al., 2019) and negative (Travis et al., 2015) associations. In the MCP, separate studies have paradoxically reported a higher FA in dyslexic readers compared to typically developing readers (Richards et al., 2008) as well as a positive association between FA and reading skills (Lebel et al., 2013; Travis et al., 2015).

Despite the general trend of higher FA relating to better reading and modest agreement in reading tract outcomes, a meta-analysis showed no evidence for systemic FA disruptions in dyslexia (Moreau et al., 2018). Small cohort sizes, inhomogeneous acquisition parameters, employment of different reading measures, variety of age groups, and diversity in processing and analytical methods may underlie the inconsistencies in past results (Moreau et al., 2018; Ramus et al., 2018; Schilling et al., 2021a). To address this, we leveraged a large database, the Healthy Brain Network (Alexander et al., 2017), to investigate white matter microstructural correlates of individual differences in single-word and single-nonword aptitude in children

with diverse reading abilities. We additionally looked for tract-specific differences in FA between groups of children with and without reading disabilities. Considering the limited sensitivity of group difference analyses and the meta-analysis by (Moreau et al., 2018), we did not expect to find significant FA group differences. However, with the added specificity of using tract-based ROIs and sensitivity from an individual differences approach, we hypothesized that several tracts, particularly in the left hemisphere, would exhibit positive associations between FA and reading scores. Further, we performed exploratory analyses considering two possible factors that may have contributed to variable prior findings. First, we divided children into younger (ages 8 and below) and older (ages 9 and above) groups who are, respectively, learning to read versus reading to learn. Second, we examined whether findings differed between scores based on reading words, which includes memorized knowledge of specific words, versus reading pronounceable nonwords (pseudowords), which is a pure measure of knowledge of letter-sound correspondence.

2 Materials and Methods

2.1 Participants

We downloaded data from 1221 participants across the first 8 data releases of the Healthy Brain Network project (Alexander et al., 2017). Participants were all scanned at Rutgers University. All data were accessed in accordance with a data use agreement provided by the Child Mind Institute. The Healthy Brain Network project was approved by the Chesapeake Institutional Review Board (now called Advarra, Inc.; <https://www.advarra.com/>). The research team obtained written informed consent from participants ages 18 or older. For younger participants, written informed consent was collected from their legal guardians,

while written assent was obtained from the participant. Full inclusion and exclusion criteria are described in the project's publication (Alexander et al., 2017). Of note, each participant was fluent in English, had an IQ over 66, and did not have any physical or mental disorder precluding them from completing the full battery of scanning and behavioral examinations. Several behavioral and cognitive evaluations were collected as part of the project. Relevant to this study, participants completed the Test of Word Reading Efficiency 2nd edition (TOWRE) (Torgesen et al., 1999) and the Edinburgh Handedness Inventory (EHI) (Oldfield, 1971).

The TOWRE consists of two subtests, Sight Word Efficiency (SWE) and Phonemic Decoding Efficiency (PDE). For Sight Word Efficiency, each participant is shown a list of words and asked to read the words aloud as quickly as possible. Raw scores are based on the number of words read correctly within the 45-second time limit and then converted to a standard score (population mean = 100, SD = 15). For Phonemic Decoding Efficiency, each participant is shown a list of pseudowords (pronounceable nonwords) and asked to read the pseudowords aloud as quickly as possible. Raw scores are based on the number of pseudowords read correctly within the 45-second time limit and then converted to a standard score (population mean = 100, SD = 15). The composite TOWRE score is the mean of the two standardized scores.

After quality control (QC), there were 686 participants ages 5-18 years old. We split these participants into two groups based on clinical diagnoses following the 5th edition of the Diagnostic and Statistical Manual for Mental Disorders (Edition et al., 2013). There were 104 participants who were diagnosed with a "specific learning disability with impairment in reading" and assigned to the reading disability RD group, while the remaining 582 participants were assigned to the typically-reading (TR) group. It is worth noting that the specific criteria for diagnosing reading disabilities were not provided, and several participants

with clinically low TOWRE scores - lower than 85 (Heath et al., 2006; Pugh et al., 2014; Johnston et al., 2016) - were not diagnosed with a reading disability (Figure 1).

We compared the ages, sex distribution, handedness, intracranial volume (ICV), and global FA (gFA) between the groups (Table 1). The TR group was slightly but significantly older than the RD group ($p < 0.05$; two-tailed Welch's t -test). There were overall more males than females in the cohort, but the proportions of sexes did not significantly differ between groups ($p > 0.05$; χ^2 test). Handedness did not differ between groups ($p > 0.3$; two-tailed Welch's t -test). ICV in TR participants was greater than in RD participants ($p < 0.05$; two-tailed Welch's t -test). The RD group had a greater global FA than the TR group ($p < 0.05$; two-tailed Welch's t -test).

Metric	All	TR	RD
<i>n</i>	686	582	104
Sex [M / F]	447 / 239	387 / 195	60 / 44
Age [years]	10.69 (3.09)	10.79 (3.18)*	10.16 (2.52)*
Handedness [EHI]	61.78 (50.69)	60.92 (50.17)	66.61 (53.74)
ICV [mm ³]	1500184 (160011)	1505734 (160747)*	1469124 (153688)*
Global FA	0.38 (0.05)	0.375 (0.052)*	0.385 (0.043)*

Table 1: Phenotypic information in the different reading proficiency groups. Values are listed as mean (standard deviation). For group comparisons, * denotes $p < 0.05$ (two-tailed Welch's t -tests). EHI: Edinburgh Handedness Inventory, FA: fractional anisotropy, ICV: intracranial volume.

Figure 1: Distribution of standardized TOWRE composite scores across all participants. Bar colors denote whether the participant was formally diagnosed with a specific learning disability with impairment in reading (RD; teal) or not (TR; red). The black dotted line marks a TOWRE score of 85 which is conventionally used for diagnosing reading disabilities. TOWRE: Tests of Word Reading Efficiency.

2.2 Neuroimaging Acquisition

Participants were scanned using a Siemens 3T Tim Trio scanner while wearing a standard Siemens 32-channel head coil. A high resolution T1-weighted (T1w) sequence was collected with the following parameters: TR = 2500 ms, TE = 3.15 ms, Flip Angle = 8°, 0.8 mm isotropic voxel resolution. A diffusion kurtosis imaging scan was administered with the following parameters: TR = 3320 ms, TE = 100.2 ms, Flip Angle = 90°, 1.8 mm isotropic voxel resolution, 1 b = 0 image, 64 noncollinear directions collected at b = 1000 s/mm² and b = 2000 s/mm². A pair of PEPolar fieldmaps were collected before the diffusion scan to quantify magnetic field inhomogeneity. Detailed scanner protocols are published on the Healthy Brain Network project website (http://fcon_1000.projects.nitrc.org/indi/chni/healthy_brain_network/File/mri/HBN_RU_Protocol.pdf).

2.3 Neuroimaging Preprocessing

Results included in this manuscript come from preprocessing performed using *QSIprep* 0.13.0RC1 (Cieslak et al., 2021, RRID:SCR_016216) (<https://qsiprep.readthedocs.io/en/latest/>) which is based on *Nipype* 1.6.0 (Gorgolewski et al., 2011, 2018, RRID:SCR_002502). Many internal operations of *QSIprep* use *Nilearn* 0.7.0 (Abraham et al., 2014, RRID:SCR_001362) and *Dipy* 1.3.0 (Garyfallidis et al., 2014, RRID:SCR_000029). Much of the text in the following two sections was provided by *QSIprep* under a CC0 license so it may be included in a manuscript for the sake of transparency and reproducibility. We made minor changes for succinctness.

2.3.1 Anatomical Preprocessing

The T1w image was corrected for intensity non-uniformity (INU) with `N4BiasFieldCorrection` (Tustison et al., 2010), distributed with *ANTs* 2.3.3 (Avants et al., 2008, RRID:SCR_004757), and used as the T1w-reference throughout the workflows. The T1w-reference was then skull-stripped with a *Nipype* implementation of the `antsBrainExtraction.sh` workflow (from *ANTs*), using OASIS30ANTs as the target template. Brain tissue segmentation of cerebrospinal fluid (CSF), white matter (WM) and gray-matter (GM) was performed on the brain-extracted T1w using `fast` (Zhang et al., 2001, *FSL* 5.0.9, RRID:SCR_002823). Additionally, brain surfaces were reconstructed using `recon-all` (*FreeSurfer* 6.0.1) (Dale et al., 1999, RRID:SCR_001847).

2.3.2 Diffusion MRI Preprocessing

MP-PCA denoising as implemented in *MRtrix3*'s `dwidenoise` (Veraart et al., 2016) was applied with a 5-voxel window. After MP-PCA, Gibbs unringing was performed using *MRtrix3*'s `mrdegibbs` (Kellner et al., 2016). Following unringing, B1 field inhomogeneity was corrected using `dwibiascorrect` from *MRtrix3* with the N4 algorithm (Tustison et al., 2010). After B1 bias correction, the mean intensity of the diffusion-weighted imaging (DWI) series was adjusted so all the mean intensity of the $b = 0$ images matched across each separate DWI scanning sequence.

FSL's (version 6.0.3:b862cdd5) `eddy` function was used for head motion correction and Eddy current correction (Andersson and Sotiropoulos, 2016). The function was configured with a q -space smoothing factor of 10, a total of 5 iterations, and 1000 voxels used to estimate hyperparameters. A linear first level model and a linear second level model were used to characterize Eddy current-related spatial distortion. q -space coordinates were forcefully

assigned to shells. Field offset was attempted to be separated from participant movement. Shells were aligned post-eddy. eddy's outlier replacement was run (Andersson et al., 2016). Data were grouped by slice, only including values from slices determined to contain at least 250 intracerebral voxels. Groups deviating by more than 4 standard deviations from the prediction had their data replaced with imputed values. Data were collected with reversed phase-encoded blips, resulting in pairs of images with distortions going in opposite directions. Here, $b = 0$ reference images with reversed phase encoding directions were used along with an equal number of $b = 0$ images extracted from the DWI scans. From these pairs the susceptibility-induced off-resonance field was estimated using a method similar to that described in (Andersson et al., 2003). These susceptibility maps were ultimately incorporated into the Eddy current and head motion correction interpolation. Final interpolation was performed using the jac method. Slicewise cross correlation was calculated. The DWI time-series were resampled to ACPC with 1.2mm isotropic voxels.

2.4 Tract Segmentation

We used *TractSeg* version 2.3 (Wasserthal et al., 2018, 2019) (<https://github.com/MIC-DKFZ/TractSeg>), a deep-learning based white matter segmentation method, to reconstruct fiber bundles. We chose this method due to its favorable balance between the accuracy of manual fiber tracking and objectivity of atlas-based methods (Genc et al., 2020). This involved the following steps: First, we reoriented images (preprocessed T1w, DWI, and brain mask) to the *FSL* standard space with *fslreorient2std* and accordingly corrected the diffusion gradient table with *MRtrix3*'s *dwigradcheck*. We used the *MRtrix3* commands *dwi2tensor* and *tensor2metric* to fit the diffusion tensor with an iterative weighted least-squares algorithm (Basser et al., 1994; Veraart et al., 2013) and produce the

FA map. Multi-tissue fiber response functions were estimated using *MRtrix3*'s *dhollander* algorithm (Dhollander et al., 2016, 2019). Fiber orientation densities (FODs) were estimated via multi-shell multi-tissue constrained spherical deconvolution (CSD) (Tournier et al., 2004, 2008; Jeurissen et al., 2014). FODs were intensity-normalized using *mtnormalize* (Raffelt et al., 2017). The first three principal FOD peaks were extracted and used as inputs into *TractSeg*'s convolutional neural network. We produced tract segmentations for the following 9 bilateral tracts: AF, SLF (I, II, and III), ILF, IFO, UF, SCP, and ICP. We also segmented the MCP and splenium of the corpus callosum (CC 7 by *TractSeg* naming convention), leading to a total of 20 fiber bundles to analyze. We extracted the average FA in each tract by calculating the average intensity of the intersection between the tract's segmentation mask with the participant's FA map. As a *post-hoc* analysis, we analyzed FA at several points along the fiber bundle (Yeatman et al., 2012b) instead of using a tract average. For this procedure, streamlines were generated for each tract. This method and its results are described in the Supplementary Materials.

2.5 Statistical Analysis

Several covariates were included in the public data set as phenotypic information, including sex (binary 0 or 1), age (in years), and handedness (EHI score; from -100 to 100). We additionally extracted estimated total intracranial volume (ICV) from *FreeSurfer* reconstructions, as well as the global white matter FA (gFA). We calculated gFA by binarizing the white matter probabilistic segmentation from *QSIprep*'s anatomical workflow at a threshold of 0.5, and averaging the FA intensity within the resulting white matter mask. We did not define gFA on a whole-brain scale as that would have introduced variance from potential differences in white matter volumetric proportions. Given the novelty of our analytical tools

and recency of the data set, we first ran Spearman correlations between these phenotypic and neuroimaging-derived metrics, probing well-established relationships in our data as a way of validating our approach and informing our choice of model parameters (Figure 2).

For each tract, we computed both a correlation between mean FA and composite TOWRE scores (skipped correlation (Wilcox, 2004; Rousselet and Pernet, 2012) with Spearman's ρ) and group difference (Welch's t -test) between mean FA values in the RD and TR groups. Before running statistical tests, we removed any participant whose tract FA was less than 0.2, as this indicated either a tract segmentation containing significant amount of non-white matter or a tract being out of the field-of-view (this happened most often for the UF in orbitofrontal cortex and the ICP in the cerebellum). This accounted for no more than 4 participants in any given tract. For group difference analyses, we linearly regressed out the effects of sex, handedness, age, global FA, and the interaction between group and sex on mean tract FA. For correlation analyses, we regressed out sex, handedness, and ICV. Standardization of TOWRE scores accounted for differences in age. Covariates were demeaned and rescaled to unit variance before regression. 40 statistical inferences were made (20 tracts $\times$ 2 measures per tract). We used Benjamini-Hochberg FDR correction to account for multiple comparisons (Benjamini and Hochberg, 1995). Statistical tests were executed by the Python package Pingouin (0.5.0) (Vallat, 2018) and visualized with the Seaborn package (0.11.2) (Waskom, 2021).

We conducted several *post-hoc* exploratory analyses to probe additional factors that may contribute to variation in prior DTI studies of reading. Recognizing that reading processes may differ between early and late-stage readers, we divide the cohort into two age brackets based on a threshold of 9 years old. This resulted in 455 older participants and 231 younger participants. Additionally, to account for differences in neurocognitive mechanisms between

Figure 2: Correlations between phenotypic and neuroimaging-derived metrics. The color of each cell corresponds to the Spearman's ρ of the correlation between the metrics in the respective row and column. * denotes $p < 0.01$, ** denotes $p < 0.001$, adjusting for multiple comparisons (Benjamini-Hochberg FDR). gFA: global fractional anisotropy, HAND: Edinburgh Handedness Inventory score, ICV: intracranial volume, TOWRE: Tests of Word Reading Efficiency (age-standardized composite score).

single-word and single-nonword reading, we also ran correlations against the SWE and PDE sub-scores. We used Benjamini-Hochberg FDR correction within each of these secondary analyses to account for multiple testing. Finally, we ran analyses of tract profiles (see Supplementary Materials), in which inferences are calculated on small segments along the length of a bundle.

2.6 Data Inclusion and Quality Control

Of the original 1221 participants, 862 participants had all the necessary neuroimaging data (T1w, diffusion, and fieldmap) and were able to be run through *QSIprep* and *TractSeg* without errors. Three of those participants were excluded for having ubiquitously sparse fiber bundle reconstructions. Two additional participants had misaligned anatomical outputs from *QSIprep*, and were excluded because gFA could not be calculated for them. Additionally, 152 of the remaining participants were missing either *FreeSurfer* reconstructions or necessary phenotypic data. An additional 16 participants were excluded for being over 18 years old. For the remaining participants, due to the volume of images included in the cohort, visual inspection of each DWI image was not practical. We instead adopted an automated QC procedure described in (Yeh et al., 2019), which has been integrated into the outputs of *QSIprep*. This involves rejecting a scan if it had different or incomplete scanning acquisitions (no participants excluded) or if over 0.1% of slices (9 slices at 72 slices/volume $\times$ 128 diffusion volumes) had significant signal dropout based on changes in slice-wise correlation (3 participants excluded). Therefore, a total of $n = 686$ participants were analyzed.

2.7 Data and Code Availability

Neuroimaging and phenotypic data can be collected following directions on the Healthy Brain Network data portal (http://fcon_1000.projects.nitrc.org/indi/cmi_healthy_brain_network/index.html) after signing a data use agreement. We cannot distribute this data publicly. All code and instructions for preprocessing neuroimaging data and running the statistical models can be found at https://github.com/smeisler/Meisler_ReadingFA_Associations. With minimal modification, the preprocessing code should be able to run on most BIDS-compliant data sets using the SLURM job scheduler (Yoo et al., 2003). Some softwares we used were distributed as Docker (Merkel, 2014) containers, then compiled and run with Singularity (3.6.3) (Kurtzer et al., 2017):

- *QSIprep* 0.13.0RC1 (singularity build qsiprep.simg
docker://pennbbl/qsiprep:0.13.0RC1)
- *TractSeg* 2.3 (singularity build tractseg.simg
docker://wasserth/tractseg:master)
- *MRtrix* 3.0.3 (singularity build mrtrix.simg
docker://mrtrix3/mrtrix3:3.0.3)
- *FSL* 6.0.4 (singularity build fsl.simg
docker://brainlife/fsl:6.0.4-patched)

We encourage anyone to use the latest stable releases of these softwares.

3 Results

3.1 Correlations Between Phenotypic and Neuroimaging Measures

We examined relationships between phenotypic and neuroimaging metrics (Figure 2). There was a significant and positive association between ICV and composite TOWRE scores (Spearman's $\rho = 0.21$, $p < 0.001$). There was also a significant and positive association between ICV and gFA (Spearman's $\rho = 0.51$, $p < 0.001$). In relation to age (development), there were strong positive correlations between age and gFA (Spearman's $\rho = 0.42$, $p < 10^{-30}$) and between age and ICV (Spearman's $\rho = 0.31$, $p < 0.001$). A negative correlation between sex (M = 0, F = 1) and ICV suggests that females had, on average, smaller brain volume than males in this cohort (Spearman's $\rho = -0.45$, $p < 10^{-30}$). Interestingly, we found a modest negative correlation between handedness quotients and TOWRE scores (Spearman's $\rho = -0.11$, $p < 0.01$), suggesting that right-handedness was related to worse reading ability. However, this comparison was unbalanced due to the high preponderance of right-handed participants (Table 1). SWE and PDE scores were highly correlated (Spearman's $\rho = 0.79$, $p < 10^{-30}$).

3.2 Mean FA-TOWRE Associations and Group Differences

Among all participants, no tract exhibited either significant associations between FA and TOWRE composite scores or significant FA differences between RD and TR groups. Correlation coefficients approximately ranged from -0.036 to 0.036, and t-statistics ranged from -0.92 to 0.24 (with a positive statistic indicating TR > RD). No significant correlations or group differences were observed when restricting analyses to specific age brackets. However,

associations tended to be stronger and more positive, albeit still modest, among the older readers, with the right SLF I exhibiting a correlation coefficient of 0.09, corresponding to a p-value of 0.07.

3.3 TOWRE Sub-Score FA Correlations

No correlations between SWE scores and mean tract FA were significant within either age bracket. Significant associations between PDE scores and FA were significant only among children ages 9 years or older (Figure 3). In this age bracket, FA in the right SLF I (Spearman's $\rho = 0.10$, $p = 0.038$) and left ICP (Spearman's $\rho = 0.09$, $p = 0.048$) were positively associated with PDE scores. Neither of the results remained significant after multiple comparison correction.

4 Discussion

This study is the first large-scale investigation of tract segmentation-derived white matter microstructural associations with single-word and single-nonword reading abilities across a diverse pediatric data set. The size of our participant cohort was large relative to other DWI studies, particularly those relating to reading aptitude. We used high-quality publicly available data and state-of-the-art analytical methods, showcasing the rapid advances in diffusion imaging acquisition and processing techniques. No tracts exhibited group differences in average FA between children with and without reading disabilities, even when considering more specific age brackets. Positive correlations between FA and reading abilities were only significant in the right SLF and left ICP when considering older participants and PDE scores, and even then, effect sizes were modest and did not survive multiple comparison correction.

Figure 3: Significant correlations between mean tract FA and TOWRE Phonemic Decoding Efficiency (PDE) scores among older participants ($n = 455$). The dots map participants' age-standardized PDE scores against average tract FA after regressing out sex, handedness, and intracranial volume. Teal denotes a RD participant, and red denotes TR participants. The black line represents the best fit across all participants, and the shaded region surrounding it depicts the 95% confidence interval of the fit. Summaries of the statistical tests are written out below each plot, including test coefficients, 95% confidence intervals surrounding the test coefficient, and uncorrected p -values. Tract shapes and anatomical image come from a single subject. Tract color relates to streamline direction. SLF: superior longitudinal fasciculus, ICP: inferior cerebellar peduncle, TR: Typically reading group, RD: reading disability group.

Our hypotheses were supported in that, overall, there were no significant group differences in tract-specific FA values between children with versus without reading disabilities or and no significant correlations between reading scores and tract-specific FA values.

The present findings are surprising in several ways, especially given the exceptional statistical power of the present sample. Firstly, we found that children with reading disability had higher global FA values than their typically reading counterparts. This differs from the frequent focus in dyslexia research on reduced FA in white matter tracts connecting the major occipito-temporal, temporal-parietal, and frontal nodes of the reading network. Secondly, while the use of age-standardized reading scores controlled for simple age-related gains in reading, but there was little relation between composite TOWRE scores and white matter pathway FA. Using G*Power 3.1 (Faul et al., 2009), we conducted a power analysis to see how many participants would be needed for the strongest correlation achieved in the primary analysis to be significant. At that effect size (0.036), with a desired $\alpha = 0.05$ and power of 0.8, 4766 participants would be required. Even if one could recruit that many participants, such a low effect size presents challenges in interpretation. Significant effect sizes were only found in our *post-hoc* exploratory analyses, and even these were modest. These secondary analyses revealed significant correlations only when using PDE scores, despite PDE and SWE scores reading being highly related, and only when examining older children (ages 9-18). At the very least, the significant correlations between FA and PDE scores were positive, which is consistent with most prior neuroanatomical studies of reading.

Given the left-lateralized dominance of reading and language brain circuitry, we did not expect FA associations to be present in the right hemisphere, at least when considering a cohort that is predominantly composed of typical readers. There has been some evidence that poor readers utilize right-hemisphere pathways to compensate for limitations of the left-

hemisphere pathways that are typically most associated with single-word reading. Several studies have reported enhanced right-hemisphere brain activation in individuals with dyslexia while reading, which has been interpreted as a compensatory mechanism (Milne et al., 2002; Temple et al., 2003; Hoeft et al., 2011; Waldie et al., 2013). Greater FA in right-hemisphere white matter pathways, in particular the right SLF, has been associated with better reading outcomes in children with dyslexia (Hoeft et al., 2011) and in young pre-reading children with low scores on assessments that predict future reading difficulty (Zuk et al., 2021). Positive correlations between right SLF FA and reading abilities have also been observed among typical readers (Horowitz-Kraus et al., 2015). However, opposing results have been reported for correlations between right SLF FA and pseudoword reading ability in particular (Steinbrink et al., 2008; Frye et al., 2011).

The TOWRE composite score was the average between real-word (SWE) and nonword (PDE) reading ability. These two measures are typically correlated, and both are impaired in dyslexia, but there are distinctions between the two measures. Single-word reading partially reflects reading experience with real words, including memorization of words. Single nonword reading, although less directly related to reading experience, provides a more pure measure of letter-sound decoding of print. We found significant, albeit small, positive correlations between FA and nonword reading scores only. This finding suggests that white matter microstructure may be especially salient for letter-sound decoding of print.

We found that age was related to finding associations between FA and nonword reading skill such that only in older children (ages 9 and older) did some tracts show a positive correlation between PDE scores and FA. Such brain-behavior relations might differ in younger children who are learning to read single words and older children who have mastered single-word reading skills and are more focused on reading comprehension of advanced texts. Indeed,

there is evidence that learning a skill leads to an initial increase in FA in putative white matter regions followed by a decline when reaching proficiency (Scholz et al., 2009), and similar early changes in white matter have been reported in mathematical abilities in dyslexia (Koerte et al., 2016), musicianship (Schmithorst and Wilke, 2002), and dance (Hänggi et al., 2010).

We, however, observed the opposite relation such that higher white matter FA was associated with nonword reading skill only in older children with more advanced reading skills. It is possible that we did not observe any significant relationships in younger readers due to reduced statistical power. In addition, our results may suggest that variation of microstructural properties of white matter does not relate to the etiology of reading ability, but rather to the long-term consequence of one's reading ability. That is, in older participants, lower FA among poor readers could be the result of having lived for years with reading difficulties (Protopapas and Parrila, 2018, 2019). A similar association of gray matter volume and reading skills was observed only in older participants in a different large-scale study (Torre and Eden, 2019). In that study, reading ability was primarily correlated with right superior temporal gyrus volume among males. Alternatively, differential reading instruction in early grades may confound reading experience and reading skill in the younger children with reading disability.

There is debate surrounding whether dyslexia can properly be described as a neurodevelopmental disorder. Population reading skills tend to follow a normal distribution, even when including those with developmental dyslexia (Figure 1). One may argue that dyslexia is not a neurodevelopmental disorder, but rather a way to group those on the lower end of the reading skills distribution (or bell curve). From this perspective, both reading differences and associated brain differences should lie along a continuum (Protopapas and Parrila, 2018, 2019).

At the simplest interpretation of this argument, being that the behavioral and neuroimaging indices lie on the same continuum, findings that would most strongly align with this view would be if reading-FA correlations were positively correlated across the entire participant cohort. We found no such association, either globally or within tracts. This suggests that the continua relating reading aptitude and FA are more complex than a shared linear relationship. Indeed, FA-behavior relationships may not be generalizable across domains and may index various neural and behavioral measures (Lazari et al., 2021).

This is the first study of reading skills that has used *TractSeg*, which has been shown to outperform several white matter segmentation methods (Wasserthal et al., 2018; Schilling et al., 2021b). Newer tract-based approaches, as employed here, represent a paradigm shift from several earlier DTI studies. Many previous papers of FA-reading relationships, particularly before the publication of Vandermosten and colleagues' review of DTI applications to reading (Vandermosten et al., 2012b), performed whole-brain voxel-based analyses (VBA). VBA sensitivity suffers from stricter multiple-comparison correction across the entire brain. In addition, VBA methods tend to be less precise due to their being performed on a group-averaged image because the shape of long-range fiber bundles varies among people (Yeatman et al., 2011; Wassermann et al., 2011). Spatial smoothing and affine transformation to MNI before group analysis in VBA may obfuscate unique properties of a participant's anatomy (Christensen et al., 1997). Significant findings from VBA are not always assigned to tracts. In fact, a single voxel may contain multiple fiber bundles, so it is possible early studies may have incorrectly ascribed significant voxels to fiber bundles due to not also considering the primary diffusion directionality in these areas. Tract-Based Spatial Statistics (TBSS) (Smith et al., 2006), a method that improves upon traditional VBA by restraining analyses to a skeletonized white matter voxel map, also suffers in tract localization (Tsang et al.,

2010). Methodological examinations of TBSS have revealed bias in the FA map used for spatial normalization, sensitivity to pathologies and noise that affect brain anatomy, incorrect voxel-to-tract assignments (Bach et al., 2014), and instability dependent on tensor-fitting methods (Maximov et al., 2015). VBA can still be valuable as a data exploratory tool, but tract-based methods likely provide for superior sensitivity and reduced ambiguity of fiber bundle localization (Ramus et al., 2018). These benefits have already been observed in multiple clinical populations (Kamagata et al., 2013; Kuchling et al., 2018; Wasserthal et al., 2020; Forkel et al., 2021).

With these limitations in mind, it may not be surprising that tract-based methods have yielded some opposing findings to voxel-based methods. For example, in the left AF, tract-based studies have yielded observations of lower FA being associated with better reading scores (Yeatman et al., 2012a; Christodoulou et al., 2017; Huber et al., 2018), as well as findings of higher FA in dyslexic readers compared to typical readers (Žarić et al., 2018). However, tractography applications to reading are still in their infancy, and different tractography measures may provide different insights. For example, CSD methods are better at resolving fiber bundles that pure DTI methods have difficulty with, such as the right AF (Catani et al., 2007; Yeatman et al., 2011; Zhao et al., 2016). The field may best be served by harmonizing diffusion acquisition and analysis protocols (Ramus et al., 2018; Cieslak et al., 2021; Schilling et al., 2021a,b), particularly scanning with multiple shells including high b-values (at least $b = 1300$) to optimally use CSD.

While our largely null results stand in opposition to previous studies reported significant FA-reading associations or group differences, a recent meta-analysis of VBA studies has suggested that these findings are not robust (Moreau et al., 2018). The present study extends this in a tract-based paradigm, using a single data set to mitigate concerns of variability from

scanning protocol confounding results (Schilling et al., 2021b). Our findings also underscore the importance including global confounds to isolate local from global effects (Ramus et al., 2018). Intracranial volume and TOWRE scores were significantly related (Figure 2), and groups differed in global fractional anisotropy (Table 1). If we had not included these factors in our models, we would have likely reported significant effects in several bilateral tracts, despite the effects being driven by global differences.

Unrelated to reading skill, we found strong correlations between age and global FA. Indeed, the present study is the largest investigation of linear white matter development patterns to date (for a review, see (Lebel et al., 2019); for a similarly-sized group-based analysis, see (Chiang et al., 2011)). Our findings are consistent with prior reports of linear FA-age relationships during child development (Mukherjee et al., 2002; Schmithorst et al., 2002; Barnea-Goraly et al., 2005; Bonekamp et al., 2007; Muetzel et al., 2008), although more recent studies suggest an exponential trajectory (Lebel et al., 2008; Tamnes et al., 2010; Taki et al., 2013; Simmonds et al., 2014) which we did not evaluate in this study. Several of our other findings unrelated to reading skill are also consistent with well-established trends, including worse readers having smaller brain volumes (Ramus et al., 2018), males having larger brain volumes than females (Courchesne et al., 2000), and brain volumes correlating positively with global FA (Takao et al., 2014). These agreements with prior studies support the quality of the DWI acquisition and the validity of our analytic method.

The results of this study should be interpreted in the context of some limitations. Firstly, the composition of the participant cohort is unique in that most participants in the Healthy Brain Network have at least one psychological, neurodevelopmental, or learning disorder (Alexander et al., 2017). Incentives for participating in the study included a cash reward as well as a free behavioral examination, and advertisements were targeted towards parents who

may have been concerned about their child's psychological state. RD participants comprised approximately 15% of the cohort. This figure exceeds estimates of dyslexia prevalence which tend to range from 5-10% (Roongpraiwan et al., 2002). Secondly, we used FA as our tract metric of interest since it has largely been interpreted as a measure of structural integrity. There is often an implicit assumption that a higher FA may relate to more myelination, which in turn may increase synaptic efficiency (Lebel and Deoni, 2018). While myelination certainly modulates FA, other factors such as axonal diameter, density, and coherence also influence it (Beaulieu, 2009; Shemesh, 2018; Friedrich et al., 2020). The dueling mechanisms underlying FA may confound results, as both axonal pruning and increased myelination may relate to typical brain development but have opposing effects on FA (Yeatman et al., 2012a). FA is also prone to underestimation and noise in voxels that contain both gray and white matter or crossing fibers (Oouchi et al., 2007), which would especially confound measurements of the corpus callosum. Analysis of T1 relaxation time in tracts using quantitative MRI may better quantify the degree of myelination per se (Lutti et al., 2014; Schurr et al., 2018). Thirdly, we could not account for FA alterations related to nonverbal IQ, since this information, in the form of the Kaufman Brief Intelligence Test (Kaufman, 1990), Second Edition, was not available for most participants. Next, our significant results should be interpreted with some caution, as they originated from exploratory analyses for which we did not develop *a priori* hypotheses. Finally, while our results lend themselves to theories of the etiology of dyslexia, we cannot interpret our findings to make causal statements of white matter microstructural properties' connection to reading disabilities. Longitudinal interventional studies should continue to be performed to better understand this relationship (Huber et al., 2018). Multivariate pattern analysis could shed insight into the relative contributions of

different tract microstructural properties to classifying reading aptitude, similar to what was done in the study by (Cui et al., 2016).

The tract-profile approach employed in the Supplementary Materials, in which FA is sampled at several points across the length of the tract, allows researchers to detect group differences and correlations that may be localized to certain portions of a tract, even if such tests using the mean FA may be null (Yeatman et al., 2012a). Using this method, we found segments of greater FA in the TR group within the left SLF and UF (Figure S1). However, it is unclear if these localized phenomena are a result of true structural differences in a given tract, or more global structural differences such as inhomogeneities in the location of a crossing fiber (Ramus et al., 2018). It also is not well-established what effect, if any, a local disruption in FA might have on synaptic efficiency between disparate areas on a fiber bundle. This presents a trade-off between anatomical specificity and interpretability. For these reasons, we keep the tract profile analyses in the Supplementary Materials.

5 Conclusion

In several ways, this study of the relation between white matter microstructure and single-word reading ability had features that ought to have yielded conclusive findings, but the study raises some open questions. The number of participants was exceptionally large, the TOWRE is a widely used and well-established measure of single-word reading ability, and the analyses took advantage of progress in DWI analytic methods. Our largely null findings, while not consistent with several individual DTI studies of reading, are consistent with recent larger-scale meta-analyses of neuroanatomical studies of dyslexia (Moreau et al., 2018; Ramus et al., 2018). Without replicating individual processing pipelines, it is difficult

to identify precise reasons previous results were or were not consistent with this study. In this respect, we believe that large-scale population studies and longitudinal studies with reproducible methods should be emphasized to account for inter-participant variability not reliably represented in small cross-sectional samples. Only future research can resolve the apparent contradictions among findings, but one possibility is that there is greater diversity among readers than has been heretofore imagined, perhaps combined with developmental variation from infancy to beginning readers to more advanced readers.

CRedit Authorship Contributions

Steven L. Meisler: Conceptualization, Formal analysis, Investigation, Methodology, Software, Visualization, Data Curation, Writing - original draft, Writing - review & editing. **John D.E. Gabrieli:** Conceptualization, Supervision, Writing - original draft, Writing - review & editing.

Acknowledgements

We thank the Healthy Brain Network team for their diligence in collecting the data and generosity in sharing it. We thank all of the participants and their families for volunteering their time to participate in the study. We thank Evelina Fedorenko, Nadine Gaab, and Tyler Perrachione for their constructive criticism of the manuscript. Finally, we thank Amanda O'Brien for proofreading the manuscript.

Funding

This work was funded by the National Institutes of Health (grant numbers 5T32DC000038-29 and 5T32DC000038-30), the Halis Family Foundation, and Reach Every Reader, a grant supported by the Chan Zuckerberg Foundation.

Conflicts of Interest

Declarations of interest: none

References

- Alexandre Abraham, Fabian Pedregosa, Michael Eickenberg, Philippe Gervais, Andreas Mueller, Jean Kossaifi, Alexandre Gramfort, Bertrand Thirion, and Gaël Varoquaux. Machine learning for neuroimaging with scikit-learn. *Frontiers in neuroinformatics*, 8:14, 2014.
- Lindsay M Alexander, Jasmine Escalera, Lei Ai, Charissa Andreotti, Karina Febre, Alexander Mangone, Natan Vega-Potler, Nicolas Langer, Alexis Alexander, Meagan Kovacs, et al. An open resource for transdiagnostic research in pediatric mental health and learning disorders. *Scientific data*, 4(1):1–26, 2017.
- Jesper LR Andersson and Stamatios N Sotiropoulos. An integrated approach to correction for off-resonance effects and subject movement in diffusion mr imaging. *Neuroimage*, 125:1063–1078, 2016.

Jesper LR Andersson, Stefan Skare, and John Ashburner. How to correct susceptibility distortions in spin-echo echo-planar images: application to diffusion tensor imaging. *Neuroimage*, 20(2):870–888, 2003.

Jesper LR Andersson, Mark S Graham, Enikő Zsoldos, and Stamatios N Sotiropoulos. Incorporating outlier detection and replacement into a non-parametric framework for movement and distortion correction of diffusion mr images. *Neuroimage*, 141:556–572, 2016.

C Nikki Arrington, Paulina A Kulesz, Jenifer Juraneck, Paul T Cirino, and Jack M Fletcher. White matter microstructure integrity in relation to reading proficiency. *Brain and language*, 174:103–111, 2017.

Brian B. Avants, Charles L. Epstein, Murray Grossman, and James C. Gee. Symmetric diffeomorphic image registration with cross-correlation: Evaluating automated labeling of elderly and neurodegenerative brain. *Medical Image Analysis*, 12(1):26–41, 2008. ISSN 1361-8415. doi: 10.1016/j.media.2007.06.004. URL <http://www.sciencedirect.com/science/article/pii/S1361841507000606>.

Michael Bach, Frederik B Laun, Alexander Leemans, Chantal MW Tax, Geert J Biessels, Bram Stieltjes, and Klaus H Maier-Hein. Methodological considerations on tract-based spatial statistics (tbss). *Neuroimage*, 100:358–369, 2014.

Chiara Banfi, Karl Koschutnig, Kristina Moll, Gerd Schulte-Körne, Andreas Fink, and Karin Landerl. White matter alterations and tract lateralization in children with dyslexia and isolated spelling deficits. *Human brain mapping*, 40(3):765–776, 2019.

Naama Barnea-Goraly, Vinod Menon, Mark Eckert, Leanne Tamm, Roland Bammer, Asya Karchemskiy, Christopher C Dant, and Allan L Reiss. White matter development during

- childhood and adolescence: a cross-sectional diffusion tensor imaging study. *Cerebral cortex*, 15(12):1848–1854, 2005.
- Peter J Basser and Carlo Pierpaoli. Microstructural and physiological features of tissues elucidated by quantitative-diffusion-tensor mri. *Journal of magnetic resonance*, 213(2): 560–570, 1996.
- Peter J Basser, James Mattiello, and Denis LeBihan. Estimation of the effective self-diffusion tensor from the nmr spin echo. *Journal of Magnetic Resonance, Series B*, 103(3):247–254, 1994.
- Christian Beaulieu. The basis of anisotropic water diffusion in the nervous system—a technical review. *NMR in Biomedicine: An International Journal Devoted to the Development and Application of Magnetic Resonance In Vivo*, 15(7-8):435–455, 2002.
- Christian Beaulieu. *Diffusion MRI: from quantitative measurement to in vivo neuroanatomy*. Academic Press, 2009.
- Yoav Benjamini and Yosef Hochberg. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)*, 57(1):289–300, 1995.
- David Bonekamp, Lidia M Nagae, Mahaveer Degaonkar, Melissa Matson, Wael MA Abdalla, Peter B Barker, Susumu Mori, and Alena Horská. Diffusion tensor imaging in children and adolescents: reproducibility, hemispheric, and age-related differences. *Neuroimage*, 34(2):733–742, 2007.
- Lauren R Borchers, Lisa Bruckert, Cory K Dodson, Katherine E Travis, Virginia A Marchman, Michal Ben-Shachar, and Heidi M Feldman. Microstructural properties of white

matter pathways in relation to subsequent reading abilities in children: a longitudinal analysis. *Brain Structure and Function*, 224(2):891–905, 2019.

Lisa Bruckert, Katherine E Travis, Aviv A Mezer, Michal Ben-Shachar, and Heidi M Feldman. Associations of reading efficiency with white matter properties of the cerebellar peduncles in children. *The Cerebellum*, 19(6):771–777, 2020.

John C Carter, Diane C Lanham, Laurie E Cutting, Amy M Clements-Stephens, Xuejing Chen, Muhamed Hadzipasic, Joon Kim, Martha B Denckla, and Walter E Kaufmann. A dual dti approach to analyzing white matter in children with dyslexia. *Psychiatry Research: Neuroimaging*, 172(3):215–219, 2009.

Marco Catani, Robert J Howard, Sinisa Pajevic, and Derek K Jones. Virtual in vivo interactive dissection of white matter fasciculi in the human brain. *Neuroimage*, 17(1):77–94, 2002.

Marco Catani, Derek K Jones, and Dominic H Ffytche. Perisylvian language networks of the human brain. *Annals of Neurology: Official Journal of the American Neurological Association and the Child Neurology Society*, 57(1):8–16, 2005.

Marco Catani, Matthew PG Allin, Masud Husain, Luca Pugliese, Marsel M Mesulam, Robin M Murray, and Derek K Jones. Symmetries in human brain language pathways correlate with verbal recall. *Proceedings of the National Academy of Sciences*, 104(43):17163–17168, 2007.

Isabella Cattinelli, N Alberto Borghese, Marcello Gallucci, and Eraldo Paulesu. Reading the reading brain: a new meta-analysis of functional imaging data on reading. *Journal of neurolinguistics*, 26(1):214–238, 2013.

Ming-Chang Chiang, Katie L McMahon, Greig I de Zubicaray, Nicholas G Martin, Ian Hickie, Arthur W Toga, Margaret J Wright, and Paul M Thompson. Genetics of white matter development: a dti study of 705 twins and their siblings aged 12 to 29. *Neuroimage*, 54(3):2308–2317, 2011.

Gary E Christensen, Sarang C Joshi, and Michael I Miller. Volumetric transformation of brain anatomy. *IEEE transactions on medical imaging*, 16(6):864–877, 1997.

Joanna A Christodoulou, Jack Murtagh, Abigail Cyr, Tyler K Perrachione, Patricia Chang, Kelly Halverson, Pamela Hook, Anastasia Yendiki, Satrajit Ghosh, and John DE Gabrieli. Relation of white-matter microstructure to reading ability and disability in beginning readers. *Neuropsychology*, 31(5):508, 2017.

Matthew Cieslak, Philip A Cook, Xiaosong He, Fang-Cheng Yeh, Thijs Dhollander, Azeez Adebimpe, Geoffrey K Aguirre, Danielle S Bassett, Richard F Betzel, Josiane Bourque, et al. Qsirep: an integrative platform for preprocessing and reconstructing diffusion mri data. *Nature Methods*, pages 1–4, 2021.

Eric Courchesne, Heather J Chisum, Jeanne Townsend, Angilene Cowles, James Covington, Brian Egaas, Mark Harwood, Stuart Hinds, and Gary A Press. Normal brain development and aging: quantitative analysis at in vivo mr imaging in healthy volunteers. *Radiology*, 216(3):672–682, 2000.

Zaixu Cui, Zhichao Xia, Mengmeng Su, Hua Shu, and Gaolang Gong. Disrupted white matter connectivity underlying developmental dyslexia: a machine learning approach. *Human brain mapping*, 37(4):1443–1458, 2016.

Anders M Dale, Bruce Fischl, and Martin I Sereno. Cortical surface-based analysis: I. segmentation and surface reconstruction. *Neuroimage*, 9(2):179–194, 1999.

Gayle K Deutsch, Robert F Dougherty, Roland Bammer, Wai Ting Siok, John DE Gabrieli, and Brian Wandell. Children's reading performance is correlated with white matter structure measured by diffusion tensor imaging. *Cortex*, 41(3):354–363, 2005.

Thijs Dhollander, David Raffelt, and Alan Connelly. Unsupervised 3-tissue response function estimation from single-shell or multi-shell diffusion mr data without a co-registered t1 image. In *ISMRM Workshop on Breaking the Barriers of Diffusion MRI*, volume 5, page 5, 2016.

Thijs Dhollander, Remika Mito, David Raffelt, and Alan Connelly. Improved white matter response function estimation for 3-tissue constrained spherical deconvolution. In *Proc. Intl. Soc. Mag. Reson. Med*, page 555, 2019.

Mark Eckert. Neuroanatomical markers for dyslexia: a review of dyslexia structural imaging studies. *The neuroscientist*, 10(4):362–371, 2004.

Fifth Edition et al. Diagnostic and statistical manual of mental disorders. *Am Psychiatric Assoc*, 21, 2013.

Franz Faul, Edgar Erdfelder, Axel Buchner, and Albert-Georg Lang. Statistical power analyses using g* power 3.1: Tests for correlation and regression analyses. *Behavior research methods*, 41(4):1149–1160, 2009.

Heidi M Feldman, Eliana S Lee, Jason D Yeatman, and Kristen W Yeom. Language and reading skills in school-aged children and adolescents born preterm are associated with white matter properties on diffusion tensor imaging. *Neuropsychologia*, 50(14):3348–3362, 2012.

Stephanie J Forkel, Patrick Friedrich, Michel Thiebaut de Schotten, and Henrietta Howells.

White matter variability, cognition, and disorders: a systematic review. *Brain Structure and Function*, pages 1–16, 2021.

Patrick Friedrich, Christoph Fraenz, Caroline Schlüter, Sebastian Ocklenburg, Burkhard

Mädler, Onur Güntürkün, and Erhan Genç. The relationship between axon density, myelination, and fractional anisotropy in the human corpus callosum. *Cerebral Cortex*, 30(4):2042–2056, 2020.

Richard E Frye, Khader Hasan, Lian Xue, David Strickland, Benjamin Malmberg, Jacque-

line Liederman, and Andrew Papanicolaou. Splenium microstructure is related to two dimensions of reading skill. *Neuroreport*, 19(16):1627, 2008.

Richard E Frye, Jacqueline Liederman, Khader M Hasan, Alexis Lincoln, Benjamin Malm-

berg, John McLean III, and Andrew Papanicolaou. Diffusion tensor quantification of the relations between microstructural and macrostructural indices of white matter and reading. *Human brain mapping*, 32(8):1220–1235, 2011.

Eleftherios Garyfallidis, Matthew Brett, Bagrat Amirbekian, Ariel Rokem, Stefan Van

Der Walt, Maxime Descoteaux, and Ian Nimmo-Smith. Dipy, a library for the analysis of diffusion mri data. *Frontiers in neuroinformatics*, 8:8, 2014.

Sila Genc, Chantal MW Tax, Erika P Raven, Maxime Chamberland, Greg D Parker, and

Derek K Jones. Impact of b-value on estimates of apparent fibre density. *Human brain mapping*, 41(10):2583–2595, 2020.

Krzysztof Gorgolewski, Christopher D. Burns, Cindee Madison, Dav Clark, Yaroslav O.

Halchenko, Michael L. Waskom, and Satrajit Ghosh. Nipype: a flexible, lightweight and

extensible neuroimaging data processing framework in python. *Frontiers in Neuroinformatics*, 5:13, 2011. doi: 10.3389/fninf.2011.00013.

Krzysztof J. Gorgolewski, Oscar Esteban, Christopher J. Markiewicz, Erik Ziegler, David Gage Ellis, Michael Philipp Notter, Dorota Jarecka, Hans Johnson, Christopher Burns, Alexandre Manhães-Savio, Carlo Hamalainen, Benjamin Yvernault, Taylor Salo, Kesshi Jordan, Mathias Goncalves, Michael Waskom, Daniel Clark, Jason Wong, Fred Loney, Marc Modat, Blake E Dewey, Cindee Madison, Matteo Visconti di Oleggio Castello, Michael G. Clark, Michael Dayan, Dav Clark, Anisha Keshavan, Basile Pinsard, Alexandre Gramfort, Shoshana Berleant, Dylan M. Nielson, Salma Bougacha, Gael Varoquaux, Ben Cipollini, Ross Markello, Ariel Rokem, Brendan Moloney, Yaroslav O. Halchenko, Demian Wassermann, Michael Hanke, Christian Horea, Jakub Kaczmarzyk, Gilles de Hollander, Elizabeth DuPre, Ashley Gillman, David Mordom, Colin Buchanan, Rosalia Tungaraza, Wolfgang M. Pauli, Shariq Iqbal, Sharad Sikka, Matteo Mancini, Yannick Schwartz, Ian B. Malone, Mathieu Dubois, Caroline Frohlich, David Welch, Jessica Forbes, James Kent, Aimi Watanabe, Chad Cumba, Julia M. Huntenburg, Erik Kastman, B. Nolan Nichols, Arman Eshaghi, Daniel Ginsburg, Alexander Schaefer, Benjamin Acland, Steven Giavasis, Jens Kleesiek, Drew Erickson, René Küttner, Christian Haselgrove, Carlos Correa, Ali Ghayoor, Franz Liem, Jarrod Millman, Daniel Haehn, Jeff Lai, Dale Zhou, Ross Blair, Tristan Glatard, Mandy Renfro, Siqi Liu, Ari E. Kahn, Fernando Pérez-García, William Triplett, Leonie Lampe, Jörg Stadler, Xiang-Zhen Kong, Michael Hallquist, Andrey Chetverikov, John Salvatore, Anne Park, Russell Poldrack, R. Cameron Craddock, Souheil Inati, Oliver Hinds, Gavin Cooper, L. Nathan Perkins, Ana Marina, Aaron Mattfeld, Maxime Noel, Lukas Snoek, K Matsubara, Brian Cheung, Simon Rothmei, Sebastian Urchs, Joke Durnez, Fred Mertz, Daniel Geisler, Andrew Floren, Stephan

- Gerhard, Paul Sharp, Miguel Molina-Romero, Alejandro Weinstein, William Broderick, Victor Saase, Sami Kristian Andberg, Robbert Harms, Kai Schlamp, Jaime Arias, Dimitri Papadopoulos Orfanos, Claire Tarbert, Arielle Tambini, Alejandro De La Vega, Thomas Nickson, Matthew Brett, Marcel Falkiewicz, Kornelius Podranski, Janosch Linkersdörfer, Guillaume Flandin, Eduard Ort, Dmitry Shachnev, Daniel McNamee, Andrew Davison, Jan Varada, Isaac Schwabacher, John Pellman, Martin Perez-Guevara, Ranjeet Khanuja, Nicolas Pannetier, Conor McDermottroe, and Satrajit Ghosh. Nipype. *Software*, 2018. doi: 10.5281/zenodo.596855.
- Patric Hagmann, Lisa Jonasson, Philippe Maeder, Jean-Philippe Thiran, Van J Wedeen, and Reto Meuli. Understanding diffusion mr imaging techniques: from scalar diffusion-weighted imaging to diffusion tensor imaging and beyond. *Radiographics*, 26(suppl_1): S205–S223, 2006.
- Jürgen Hänggi, Susan Koenke, Ladina Bezzola, and Lutz Jäncke. Structural neuroplasticity in the sensorimotor network of professional female ballet dancers. *Human brain mapping*, 31(8):1196–1206, 2010.
- Steve M Heath, Dorothy VM Bishop, John H Hogben, and Neil W Roach. Psychophysical indices of perceptual functioning in dyslexia: A psychometric analysis. *Cognitive neuropsychology*, 23(6):905–929, 2006.
- Guillaume Herbet, Ilyess Zemmoura, and Hugues Duffau. Functional anatomy of the inferior longitudinal fasciculus: from historical reports to current hypotheses. *Frontiers in neuroanatomy*, 12:77, 2018.
- Fumiko Hoeft, Bruce D McCandliss, Jessica M Black, Alexander Gantman, Nahal Zakerani, Charles Hulme, Heikki Lyytinen, Susan Whitfield-Gabrieli, Gary H Glover, Allan L Reiss,

- et al. Neural systems predicting long-term outcome in dyslexia. *Proceedings of the National Academy of Sciences*, 108(1):361–366, 2011.
- Tzipi Horowitz-Kraus, Jennifer J Vannest, Darren Kadis, Nicole Cicchino, Yingying Y Wang, and Scott K Holland. Reading acceleration training changes brain circuitry in children with reading difficulties. *Brain and Behavior*, 4(6):886–902, 2014.
- Tzipi Horowitz-Kraus, Molly Grainger, Mark DiFrancesco, Jennifer Vannest, and Scott K Holland. Right is not always wrong: Dti and fmri evidence for the reliance of reading comprehension on language-comprehension networks in the right hemisphere. *Brain imaging and behavior*, 9(1):19–31, 2015.
- Elizabeth Huber, Patrick M Donnelly, Ariel Rokem, and Jason D Yeatman. Rapid and widespread white matter plasticity during an intensive reading intervention. *Nature communications*, 9(1):1–13, 2018.
- Katarzyna Jednorog, Artur Marchewka, Irene Altarelli, Ana Karla Monzalvo Lopez, Muna van Ermingen-Marbach, Marion Grande, Anna Grabowska, Stefan Heim, and Franck Ramus. How reliable are gray matter disruptions in specific reading disability across multiple countries and languages? insights from a large-scale voxel-based morphometry study. *Human brain mapping*, 36(5):1741–1754, 2015.
- Ben Jeurissen, Jacques-Donald Tournier, Thijs Dhollander, Alan Connelly, and Jan Sijbers. Multi-tissue constrained spherical deconvolution for improved analysis of multi-shell diffusion mri data. *NeuroImage*, 103:411–426, 2014.
- Richard Johnston, Nicola J Pitchford, Neil W Roach, and Timothy Ledgeway. Why is the processing of global motion impaired in adults with developmental dyslexia? *Brain and Cognition*, 108:20–31, 2016.

Koji Kamagata, Yumiko Motoi, Hiroyuki Tomiyama, Osamu Abe, Kenji Ito, Keigo Shimoji, Michimasa Suzuki, Masaaki Hori, Atsushi Nakanishi, Tamotsu Sano, et al. Relationship between cognitive impairment and white-matter alteration in parkinson's disease with dementia: tract-based spatial statistics and tract-specific analysis. *European radiology*, 23(7):1946–1955, 2013.

Alan S Kaufman. *Kaufman brief intelligence test: KBIT*. AGS, American Guidance Service Circle Pines, MN, 1990.

Elias Kellner, Bibek Dhital, Valerij G Kiselev, and Marco Reisert. Gibbs-ringing artifact removal based on local subvoxel-shifts. *Magnetic resonance in medicine*, 76(5):1574–1581, 2016.

Torkel Klingberg, Maj Hedehus, Elise Temple, Talya Salz, John DE Gabrieli, Michael E Moseley, and Russell A Poldrack. Microstructure of temporo-parietal white matter as a basis for reading ability: evidence from diffusion tensor magnetic resonance imaging. *Neuron*, 25(2):493–500, 2000.

Inga K Koerte, Anna Willems, Marc Muehlmann, Kristina Moll, Sonia Cornell, Silvia Pixner, Denise Steffinger, Daniel Keeser, Florian Heinen, Marek Kubicki, et al. Mathematical abilities in dyslexic children: a diffusion tensor imaging study. *Brain imaging and behavior*, 10(3):781–791, 2016.

Joseph Kuchling, Yael Backner, Frederike C Oertel, Noa Raz, Judith Bellmann-Strobl, Klemens Ruprecht, Friedemann Paul, Netta Levin, Alexander U Brandt, and Michael Scheel. Comparison of probabilistic tractography and tract-based spatial statistics for assessing optic radiation damage in patients with autoimmune inflammatory disorders of the central nervous system. *NeuroImage: Clinical*, 19:538–550, 2018.

Gregory M Kurtzer, Vanessa Sochat, and Michael W Bauer. Singularity: Scientific containers for mobility of compute. *PloS one*, 12(5):e0177459, 2017.

Alberto Lazari, Piergiorgio Salvan, Michiel Cottaar, Daniel Papp, Olof Jens van der Werf, Ainslie Johnstone, Zeena-Britt Sanders, Cassandra Sampaio-Baptista, Nicole Eichert, Kentaro Miyamoto, et al. Reassessing associations between white matter and behaviour with multimodal microstructural imaging. *Cortex*, 2021.

Catherine Lebel and Sean Deoni. The development of brain white matter microstructure. *Neuroimage*, 182:207–218, 2018.

Catherine Lebel, Lindsay Walker, Alexander Leemans, Linda Phillips, and Christian Beaulieu. Microstructural maturation of the human brain from childhood to adulthood. *Neuroimage*, 40(3):1044–1055, 2008.

Catherine Lebel, Bennett Shaywitz, John Holahan, Sally Shaywitz, Karen Marchione, and Christian Beaulieu. Diffusion tensor imaging correlates of reading ability in dysfluent and non-impaired readers. *Brain and language*, 125(2):215–222, 2013.

Catherine Lebel, Sarah Treit, and Christian Beaulieu. A review of diffusion mri of typical white matter development from early childhood to young adulthood. *NMR in Biomedicine*, 32(4):e3778, 2019.

Janosch Linkersdörfer, Jan Lonnemann, Sven Lindberg, Marcus Hasselhorn, and Christian J. Fiebach. Grey matter alterations co-localize with functional abnormalities in developmental dyslexia: An ale meta-analysis. *PLOS ONE*, 7(8):1–10, 08 2012. doi: 10.1371/journal.pone.0043122. URL <https://doi.org/10.1371/journal.pone.0043122>.

- Antoine Lutti, Frederic Dick, Martin I Sereno, and Nikolaus Weiskopf. Using high-resolution quantitative mapping of r1 as an index of cortical myelination. *Neuroimage*, 93:176–188, 2014.
- G Reid Lyon, Sally E Shaywitz, and Bennett A Shaywitz. A definition of dyslexia. *Annals of dyslexia*, 53(1):1–14, 2003.
- José M Maisog, Erin R Einbinder, D Lynn Flowers, Peter E Turkeltaub, and Guinevere F Eden. A meta-analysis of functional neuroimaging studies of dyslexia. *Annals of the New York Academy of Sciences*, 1145(1):237–259, 2008.
- Cecilia Marino, Paola Scifo, Pasquale A Della Rosa, Sara Mascheretti, Andrea Facoetti, Maria L Lorusso, Roberto Giorda, Monica Consonni, Andrea Falini, Massimo Molteni, et al. The dcdc2/intron 2 deletion and white matter disorganization: focus on developmental dyslexia. *Cortex*, 57:227–243, 2014.
- Juan Martino, Christian Brogna, Santiago G Robles, Francesco Vergani, and Hugues Duffau. Anatomic dissection of the inferior fronto-occipital fasciculus revisited in the lights of brain stimulation data. *cortex*, 46(5):691–699, 2010.
- Ivan I Maximov, Heike Thönneßen, Kerstin Konrad, Laura Amort, Irene Neuner, and N Jon Shah. Statistical instability of tbss analysis based on dti fitting algorithm. *Journal of Neuroimaging*, 25(6):883–891, 2015.
- Dirk Merkel. Docker: lightweight linux containers for consistent development and deployment. *Linux journal*, 2014(239):2, 2014.

- R Duncan Milne, Ari Syngieniotis, Graeme Jackson, and Michael C Corballis. Mixed lateralization of phonological assembly in developmental dyslexia. *Neurocase*, 8(3): 209–209, 2002.
- David Moreau, Josephine E Stonyer, Nicole S McKay, and Karen E Waldie. No evidence for systematic white matter correlates of dyslexia: an activation likelihood estimation meta-analysis. *Brain research*, 1683:36–47, 2018.
- Ryan L Muetzel, Paul F Collins, Bryon A Mueller, Ann M Schissel, Kelvin O Lim, and Monica Luciana. The development of corpus callosum microstructure and associations with bimanual task performance in healthy adolescents. *Neuroimage*, 39(4):1918–1925, 2008.
- Pratik Mukherjee, Jeffrey H Miller, Joshua S Shimony, Joseph V Philip, Deepika Nehra, Abraham Z Snyder, Thomas E Conturo, Jeffrey J Neil, and Robert C McKinstry. Diffusion-tensor mr imaging of gray and white matter development during normal human brain maturation. *American Journal of Neuroradiology*, 23(9):1445–1456, 2002.
- Kimberly A Murphy, Jigar R Jogia, and Joel B Talcott. On the neural basis of word reading: A meta-analysis of fmri evidence using activation likelihood estimation. *Journal of Neurolinguistics*, 49:71–83, 2019.
- Roderick I Nicolson and Angela J Fawcett. Procedural learning difficulties: reuniting the developmental disorders? *TRENDS in Neurosciences*, 30(4):135–141, 2007.
- Roderick I Nicolson, Angela J Fawcett, and Paul Dean. Developmental dyslexia: the cerebellar deficit hypothesis. *Trends in neurosciences*, 24(9):508–511, 2001.

Timothy N Odegard, Emily A Farris, Jeremiah Ring, Roderick McColl, and Jeffrey Black.

Brain connectivity in non-reading impaired children and children diagnosed with developmental dyslexia. *Neuropsychologia*, 47(8-9):1972–1977, 2009.

Richard C Oldfield. The assessment and analysis of handedness: the edinburgh inventory.

Neuropsychologia, 9(1):97–113, 1971.

Hiroyuki Oouchi, Kei Yamada, Koji Sakai, Osamu Kizu, Takao Kubota, Hiroto Ito, and

Tsunehiko Nishimura. Diffusion anisotropy measurement of brain white matter is affected by voxel size: underestimation occurs in areas with crossing fibers. *American Journal of Neuroradiology*, 28(6):1102–1106, 2007.

Athanassios Protopapas and Rauno Parrila. Is dyslexia a brain disorder? *Brain sciences*, 8

(4):61, 2018.

Athanassios Protopapas and Rauno Parrila. Dyslexia: Still not a neurodevelopmental disorder.

Brain sciences, 9(1):9, 2019.

Kenneth R Pugh, W Einar Mencl, Annette R Jenner, Leonard Katz, Stephen J Frost, Jun Ren

Lee, Sally E Shaywitz, and Bennett A Shaywitz. Functional neuroimaging studies of reading and reading disability (developmental dyslexia). *Mental retardation and developmental disabilities research reviews*, 6(3):207–213, 2000.

Kenneth R Pugh, Stephen J Frost, Douglas L Rothman, Fumiko Hoeft, Stephanie N Del Tufo,

Graeme F Mason, Peter J Molfese, W Einar Mencl, Elena L Grigorenko, Nicole Landi, et al. Glutamate and choline levels predict individual differences in reading ability in emergent readers. *Journal of Neuroscience*, 34(11):4082–4089, 2014.

Mary Colvin Putnam, Megan S Steven, Karl W Doron, Adam C Riggall, and Michael S Gazzaniga. Cortical projection topography of the human splenium: hemispheric asymmetry and individual differences. *Journal of cognitive neuroscience*, 22(8):1662–1669, 2010.

David Raffelt, Thijs Dhollander, J-Donald Tournier, Rami Tabbara, Robert E Smith, Eric Pierre, and Alan Connelly. Bias field correction and intensity normalisation for quantitative analysis of apparent fibre density. In *Proc. Intl. Soc. Mag. Reson. Med*, volume 25, page 3541, 2017.

Franck Ramus, Irene Altarelli, Katarzyna Jednoróg, Jingjing Zhao, and Lou Scotto Di Covella. Neuroanatomy of developmental dyslexia: Pitfalls and promise. *Neuroscience & Biobehavioral Reviews*, 84:434–452, 2018.

Andreas M Rauschecker, Gayle K Deutsch, Michal Ben-Shachar, Armin Schwartzman, Lee M Perry, and Robert F Dougherty. Reading impairment in a patient with missing arcuate fasciculus. *Neuropsychologia*, 47(1):180–194, 2009.

Todd L Richards, Jim Stevenson, J Crouch, L Clark Johnson, Kenneth R Maravilla, Pat Stock, Robert D Abbott, and Virginia W Berninger. Tract-based spatial statistics of diffusion tensor imaging in adults with dyslexia. *American Journal of Neuroradiology*, 29(6):1134–1139, 2008.

Fabio Richlan, Martin Kronbichler, and Heinz Wimmer. Structural abnormalities in the dyslexic brain: A meta-analysis of voxel-based morphometry studies. *Human brain mapping*, 34(11):3055–3065, 2013.

Rawiwan Roongpraiwan, Nichara Ruangdaraganon, Pongsak Visudhiphan, and Kanitta Santikul. Prevalence and clinical characteristics of dyslexia in primary school students.

- Journal of the Medical Association of Thailand= Chotmai het thangphaet*, 85:S1097–103, 2002.
- Guillaume A Rousselet and Cyril R Pernet. Improving standards in brain-behavior correlation analyses. *Frontiers in human neuroscience*, 6:119, 2012.
- Kurt G Schilling, François Rheault, Laurent Petit, Colin B Hansen, Vishwesh Nath, Fang-Cheng Yeh, Gabriel Girard, Muhamed Barakovic, Jonathan Rafael-Patino, Thomas Yu, et al. Tractography dissection variability: what happens when 42 groups dissect 14 white matter bundles on the same dataset? *NeuroImage*, page 118502, 2021a.
- Kurt G Schilling, Chantal MW Tax, Francois Rheault, Colin Hansen, Qi Yang, Fang-Cheng Yeh, Leon Cai, Adam W Anderson, and Bennett A Landman. Fiber tractography bundle segmentation depends on scanner effects, vendor effects, acquisition resolution, diffusion sampling scheme, diffusion sensitization, and bundle segmentation workflow. *NeuroImage*, 242:118451, 2021b.
- Bradley L Schlaggar and Bruce D McCandliss. Development of neural systems for reading. *Annu. Rev. Neurosci.*, 30:475–503, 2007.
- Vincent J Schmithorst and Marko Wilke. Differences in white matter architecture between musicians and non-musicians: a diffusion tensor imaging study. *Neuroscience letters*, 321(1-2):57–60, 2002.
- Vincent J Schmithorst, Marko Wilke, Bernard J Dardzinski, and Scott K Holland. Correlation of white matter diffusivity and anisotropy with age during childhood and adolescence: a cross-sectional diffusion-tensor mr imaging study. *Radiology*, 222(1):212–218, 2002.

Jan Scholz, Miriam C Klein, Timothy EJ Behrens, and Heidi Johansen-Berg. Training induces changes in white-matter architecture. *Nature neuroscience*, 12(11):1370–1371, 2009.

Roey Schurr, Yiran Duan, Anthony M Norcia, Shumpei Ogawa, Jason D Yeatman, and Aviv A Mezer. Tractography optimization using quantitative t1 mapping in the human optic radiation. *Neuroimage*, 181:645–658, 2018.

Noam Shemesh. Axon diameters and myelin content modulate microscopic fractional anisotropy at short diffusion times in fixed rat spinal cord. *Frontiers in Physics*, 6:49, 2018.

Daniel J Simmonds, Michael N Hallquist, Miya Asato, and Beatriz Luna. Developmental stages and sex differences of white matter and behavioral development through adolescence: a longitudinal diffusion tensor imaging (dti) study. *Neuroimage*, 92:356–368, 2014.

Stephen M Smith, Mark Jenkinson, Heidi Johansen-Berg, Daniel Rueckert, Thomas E Nichols, Clare E Mackay, Kate E Watkins, Olga Ciccarelli, M Zaheer Cader, Paul M Matthews, et al. Tract-based spatial statistics: voxelwise analysis of multi-subject diffusion data. *Neuroimage*, 31(4):1487–1505, 2006.

Claudia Steinbrink, Katrin Vogt, A Kastrup, H-P Müller, FD Juengling, J Kassubek, and A Riecker. The contribution of white and gray matter differences to developmental dyslexia: insights from dti and vbm at 3.0 t. *Neuropsychologia*, 46(13):3170–3178, 2008.

Catherine J Stoodley and John F Stein. The cerebellum and dyslexia. *Cortex: A Journal Devoted to the Study of the Nervous System and Behavior*, 2011.

- Mengmeng Su, Jingjing Zhao, Michel Thiebaut de Schotten, Wei Zhou, Gaolang Gong, Franck Ramus, and Hua Shu. Alterations in white matter pathways underlying phonological and morphological processing in chinese developmental dyslexia. *Developmental cognitive neuroscience*, 31:11–19, 2018.
- Hidemasa Takao, Naoto Hayashi, and Kuni Ohtomo. Sex dimorphism in the white matter: fractional anisotropy and brain size. *Journal of Magnetic Resonance Imaging*, 39(4): 917–923, 2014.
- Yasuyuki Taki, Benjamin Thyreau, Hiroshi Hashizume, Yuko Sassa, Hikaru Takeuchi, Kai Wu, Yuka Kotozaki, Rui Nouchi, Michiko Asano, Kohei Asano, et al. Linear and curvilinear correlations of brain white matter volume, fractional anisotropy, and mean diffusivity with age using voxel-based and region-of-interest analyses in 246 healthy children. *Human brain mapping*, 34(8):1842–1856, 2013.
- Christian K Tamnes, Ylva Østby, Anders M Fjell, Lars T Westlye, Paulina Due-Tønnessen, and Kristine B Walhovd. Brain maturation in adolescence and young adulthood: regional age-related changes in cortical thickness and white matter volume and microstructure. *Cerebral cortex*, 20(3):534–548, 2010.
- Elise Temple, Gayle K Deutsch, Russell A Poldrack, Steven L Miller, Paula Tallal, Michael M Merzenich, and John DE Gabrieli. Neural deficits in children with dyslexia ameliorated by behavioral remediation: evidence from functional mri. *Proceedings of the National Academy of Sciences*, 100(5):2860–2865, 2003.
- Joseph K Torgesen, Carol Alexander Rashotte, and Richard K Wagner. *TOWRE: Test of word reading efficiency*. Psychological Corporation Toronto, Ontario, 1999.

Gabrielle-Ann A Torre and Guinevere F Eden. Relationships between gray matter volume and reading ability in typically developing children, adolescents, and young adults. *Developmental cognitive neuroscience*, 36:100636, 2019.

J-Donald Tournier, Fernando Calamante, David G Gadian, and Alan Connelly. Direct estimation of the fiber orientation density function from diffusion-weighted mri data using spherical deconvolution. *Neuroimage*, 23(3):1176–1185, 2004.

J-Donald Tournier, Chun-Hung Yeh, Fernando Calamante, Kuan-Hung Cho, Alan Connelly, and Ching-Po Lin. Resolving crossing fibres using constrained spherical deconvolution: validation using diffusion-weighted imaging phantom data. *Neuroimage*, 42(2):617–625, 2008.

Katherine E Travis, Yael Leitner, Heidi M Feldman, and Michal Ben-Shachar. Cerebellar white matter pathways are associated with reading skills in children and adolescents. *Human brain mapping*, 36(4):1536–1553, 2015.

JM Tsang, RF Dougherty, and BA Wandell. Tract alignment errors decrease detection power in group analyses of diffusion data with tbss. *Society for Neuroscience. San Diego, CA*, 2010.

Nicholas J. Tustison, Brian B. Avants, Philip A. Cook, Yuanjie Zheng, Alexander Egan, Paul A. Yushkevich, and James C. Gee. N4itk: Improved n3 bias correction. *IEEE Transactions on Medical Imaging*, 29(6):1310–1320, 2010. ISSN 0278-0062. doi: 10.1109/TMI.2010.2046908.

Raphael Vallat. Pingouin: statistics in python. *The Journal of Open Source Software*, 3(31): 1026, November 2018.

Maaïke Vandermosten, Bart Boets, Hanne Poelmans, Stefan Sunaert, Jan Wouters, and Pol Ghesquiere. A tractography study in dyslexia: neuroanatomic correlates of orthographic, phonological and speech processing. *Brain*, 135(3):935–948, 2012a.

Maaïke Vandermosten, Bart Boets, Jan Wouters, and Pol Ghesquière. A qualitative and quantitative review of diffusion tensor imaging studies in reading and dyslexia. *Neuroscience & Biobehavioral Reviews*, 36(6):1532–1552, 2012b.

Jelle Veraart, Jan Sijbers, Stefan Sunaert, Alexander Leemans, and Ben Jeurissen. Weighted linear least squares estimation of diffusion mri parameters: strengths, limitations, and pitfalls. *Neuroimage*, 81:335–346, 2013.

Jelle Veraart, Dmitry S Novikov, Daan Christiaens, Benjamin Ades-Aron, Jan Sijbers, and Els Fieremans. Denoising of diffusion mri using random matrix theory. *NeuroImage*, 142:394–406, 2016.

Karen E Waldie, Charlotte E Haigh, Gjurgjica Badzakova-Trajkov, Jude Buckley, and Ian J Kirk. Reading the wrong way with the right hemisphere. *Brain Sciences*, 3(3):1060–1075, 2013.

Brian A Wandell and Jason D Yeatman. Biological development of reading circuits. *Current Opinion in Neurobiology*, 23(2):261–268, 2013.

Xuhui Wang, Sudhir Pathak, Lucia Stefanescu, Fang-Cheng Yeh, Shiting Li, and Juan C Fernandez-Miranda. Subcomponents and connectivity of the superior longitudinal fasciculus in the human brain. *Brain Structure and Function*, 221(4):2075–2092, 2016.

Michael L Waskom. Seaborn: statistical data visualization. *Journal of Open Source Software*, 6(60):3021, 2021.

- Demian Wassermann, Yogesh Rathi, Sylvain Bouix, Marek Kubicki, Ron Kikinis, Martha Shenton, and Carl-Fredrik Westin. White matter bundle registration and population analysis based on gaussian processes. In *Biennial International Conference on Information Processing in Medical Imaging*, pages 320–332. Springer, 2011.
- Jakob Wasserthal, Peter Neher, and Klaus H Maier-Hein. Tractseg-fast and accurate white matter tract segmentation. *NeuroImage*, 183:239–253, 2018.
- Jakob Wasserthal, Peter F Neher, Dusan Hirjak, and Klaus H Maier-Hein. Combined tract segmentation and orientation mapping for bundle-specific tractography. *Medical image analysis*, 58:101559, 2019.
- Jakob Wasserthal, Klaus H Maier-Hein, Peter F Neher, Georg Northoff, Katharina M Kubera, Stefan Fritze, Anais Harneit, Lena S Geiger, Heike Tost, Robert C Wolf, et al. Multiparametric mapping of white matter microstructure in catatonia. *Neuropsychopharmacology*, 45(10):1750–1757, 2020.
- Suzanne E Welcome and Marc F Joanisse. Individual differences in white matter anatomy predict dissociable components of reading skill in adults. *Neuroimage*, 96:261–275, 2014.
- Rand Wilcox. Inferences based on a skipped correlation coefficient. *Journal of Applied Statistics*, 31(2):131–143, 2004.
- Jason D Yeatman, Robert F Dougherty, Elena Rykhlevskaia, Anthony J Sherbondy, Gayle K Deutsch, Brian A Wandell, and Michal Ben-Shachar. Anatomical properties of the arcuate fasciculus predict phonological and reading skills in children. *Journal of cognitive neuroscience*, 23(11):3304–3317, 2011.

Jason D Yeatman, Robert F Dougherty, Michal Ben-Shachar, and Brian A Wandell. Development of white matter and reading skills. *Proceedings of the National Academy of Sciences*, 109(44):E3045–E3053, 2012a.

Jason D Yeatman, Robert F Dougherty, Nathaniel J Myall, Brian A Wandell, and Heidi M Feldman. Tract profiles of white matter properties: automating fiber-tract quantification. *PloS one*, 7(11):e49790, 2012b.

Jason D Yeatman, Andreas M Rauschecker, and Brian A Wandell. Anatomy of the visual word form area: adjacent cortical circuits and long-range white matter connections. *Brain and language*, 125(2):146–155, 2013.

Fang-Cheng Yeh, Islam M Zaydan, Valerie R Suski, David Lacomis, R Mark Richardson, Joseph C Maroon, and Jessica Barrios-Martinez. Differential tractography as a track-based biomarker for neuronal injury. *Neuroimage*, 202:116131, 2019.

Andy B Yoo, Morris A Jette, and Mark Grondona. Slurm: Simple linux utility for resource management. In *Workshop on Job Scheduling Strategies for Parallel Processing*, pages 44–60. Springer, 2003.

Gojko Žarić, Inge Timmers, Patty Gerretsen, Gorka Fraga González, Jurgen Tijms, Maurits W van der Molen, Leo Blomert, and Milene Bonte. Atypical white matter connectivity in dyslexic readers of a fairly transparent orthography. *Frontiers in psychology*, 9:1147, 2018.

Mingxia Zhang, Chuansheng Chen, Gui Xue, Zhong-lin Lu, Leilei Mei, Hongli Xue, Miao Wei, Qinghua He, Jin Li, and Qi Dong. Language-general and-specific white matter microstructural bases for reading. *NeuroImage*, 98:435–441, 2014.

- Yongyue. Zhang, Michael. Brady, and Stephen Smith. Segmentation of brain MR images through a hidden markov random field model and the expectation-maximization algorithm. *IEEE Transactions on Medical Imaging*, 20(1):45–57, 2001. ISSN 0278-0062. doi: 10.1109/42.906424.
- Jingjing Zhao, Michel Thiebaut de Schotten, Irene Altarelli, Jessica Dubois, and Franck Ramus. Altered hemispheric lateralization of white matter pathways in developmental dyslexia: Evidence from spherical deconvolution tractography. *Cortex*, 76:51–62, 2016.
- Jennifer Zuk, Jade Dunstan, Elizabeth Norton, Xi Yu, Ola Ozernov-Palchik, Yingying Wang, Tiffany P Hogan, John DE Gabrieli, and Nadine Gaab. Multifactorial pathways facilitate resilience among kindergarteners at risk for dyslexia: A longitudinal behavioral and neuroimaging study. *Developmental science*, 24(1):e12983, 2021.