The diversity and functional capacity of microbes associated with coastal phototrophs

Khashiff Miranda^{1,2*}, Brooke L. Weigel^{3,4}, Emily C. Fogarty^{5,6}, Iva A. Veseli⁷, Anne E. Giblin⁸,
A. Murat Eren^{5,9}, Catherine A. Pfister^{3,10}

* Corresponding author email: khashiff.miranda.1@ulaval.ca

¹ The College, The University of Chicago, Chicago, IL, 60637, USA

² Département de biologie, Université Laval, Québec, QC, G1V 0A6, Canada

³ Committee on Evolutionary Biology, The University of Chicago, Chicago, IL, 60637, USA

⁴ Friday Harbor Laboratories, University of Washington, Friday Harbor, WA 98250, USA

⁵ Department of Medicine, The University of Chicago, Chicago, IL 60637, USA

⁶ Committee on Microbiology, The University of Chicago, Chicago, IL 60637, USA

⁷ Biophysical Sciences Program, The University of Chicago, Chicago, IL, 60637, USA

⁸ The Ecosystems Center, The Marine Biological Laboratory, Woods Hole, MA, 02543, USA

⁹ Josephine Bay Paul Center, Marine Biological Laboratory, Woods Hole, MA 02543, USA

¹⁰ Department of Ecology & Evolution, The University of Chicago, Chicago, IL, 60637 USA

Abstract

Coastal marine phototrophs exhibit some of the highest rates of primary productivity in the world. They have been found to host a diverse set of microbes, many of which may impact the biology of their phototroph hosts through metabolisms that are unique to microbial taxa. Here we characterized the metabolic functions of phototroph-associated microbial communities using metagenomes collected from 2 species of kelp (*Laminaria setchellii* and *Nereocystis luetkeana*)

and 3 marine angiosperms (*Phyllospadix scouleri*, *P. serrulatus* and *Zostera marina*), including the rhizomes of two surfgrass species (*Phyllospadix* spp.) and the seagrass *Zostera marina*, and the sediments surrounding *P. scouleri* and *Z. marina*. Using metagenomic sequencing, we describe 72 metagenome assembled genomes (MAGs) that potentially benefit from being associated with macrophytes and may contribute to macrophyte fitness through their metabolic gene content. All host-associated metagenomes contained genes for the use of dissolved organic matter from hosts and vitamin (B₁, B₂, B₇, B₁₂) biosynthesis. Additionally, we found a range of nitrogen metabolism genes that transform dissolved inorganic nitrogen into forms that may be more available to the host. The rhizosphere of surfgrass and seagrass contained genes for anaerobic microbial metabolisms, including *nifH* genes associated with nitrogen fixation, despite residing in a well-mixed and oxygenated environment. The range of oxygen environments engineered by macrophytes likely explains the diversity of both oxidizing and reducing microbial metabolisms, and contributes to the functional capabilities of microbes and their influence on carbon and nitrogen cycling in nearshore ecosystems.

Importance

Kelps, seagrasses and surfgrasses are ecosystem engineers on rocky shorelines where they show remarkably high levels of primary production. Through analysis of their associated microbial communities, we found a variety of microbial metabolisms that may benefit the host, including nitrogen metabolisms and the production of B vitamins. In turn, these microbes have the genetic capability to assimilate the dissolved organic compounds released by their phototroph hosts. We describe a range of oxygen environments associated with surfgrass, including low-oxygen microhabitats in their rhizomes that host genes for nitrogen fixation. The tremendous

productivity of coastal phototrophs is likely due in part to the activities of associated microbes and an increased understanding of these associations is needed.

Introduction

We are experiencing a paradigm shift in biology with the recognition that many species exist as a consortium with microbes (1). These microbial associations are nearly ubiquitous, spanning a diversity of hosts across ecosystems. In coastal marine environments, phototrophic microbial hosts are diverse and range from marine angiosperms to large eukaryotic protists (macroalgae). Different macroalgal host species (2, 3) and different phototroph tissues (4, 5) host distinct microbial communities numbering in the millions per cm² of host tissue (6), yet we still know little about the functional role the microbiome plays in host fitness or how the host influences the microbiome. The microbiome of phototroph species has been shown to have metabolisms that provide nitrogen to the host (7, 8). Bacteria also supply B vitamins (9) and affect development of their host (10). Further, the contributions that marine phototrophs make to host carbon and nitrogen cycling have largely ignored the role that microbes play. Even as we begin to describe their microbiome, we are discovering that environmental change affects these communities (11). For many of the foundational phototrophic species in the coastal ocean, our understanding of the diversity and role of their microbiome is nascent.

A unique aspect of host-associated microbes are the strong gradients in oxygen that they experience due to the biological activities of the host. The photosynthetic and respiratory activities of the host can generate a ‘phycosphere’ (12) where the host influences the physical environment experienced by microbes, sometimes over micron or mm scales. For example, the

basal leaf meristem of the seagrass *Zostera* ranges from oxic to anoxic conditions over a scale of 300 microns when measured with oxygen microsensors (13). This range of oxygen concentrations likely selects for a diversity of microbial metabolisms in association with macrophytes.

Another factor important to the microbial metabolisms associated with coastal macrophytes is nutrient availability. In coastal systems, nitrogen can limit primary production and microbial associates that aid in accessing nitrogen might be selected. Microbial metabolisms that can increase the available dissolved inorganic nitrogen (DIN) for the host (14) include pathways that cleave carbon-nitrogen bonds to generate ammonium. This ammonification in biological systems can result from a diversity of hydrolases, including ureases and other enzymes that cleave C-N bonds (15). Further, microbes that fix atmospheric nitrogen have been discovered in an increasing number of taxa (8, 16), now recognized to include heterotrophic as well as phototrophic taxa (17–19). Nitrogen fixation was previously assumed to be restricted to nitrogen-poor environments, but has been quantified recently in systems thought to be nitrogen-rich (8, 20), an enigmatic finding given that nitrogen fixation is a costly metabolic process that consumes 16 ATPs per N₂ fixed (21). Sediments where oxygen is low and nutrients can be depleted by macrophytes, such as the rhizosphere of seagrasses have provided evidence of nitrogen fixation (22–26). The recent discovery that nitrogen fixation takes place on particles in the coastal ocean where nitrate is relatively abundant (8, 20) suggests that *nifH* genes could be abundant in other nearshore systems.

Microbial metabolisms that synthesize compounds and vitamins needed by seaweeds and seagrasses may also underlie host-microbe exchanges. The active form of Vitamin B1 (thiamin) is essential for all organisms and is involved in carbohydrate and amino acid metabolisms. Vitamin B2 (riboflavin)-binding proteins are co-enzymes in various oxidases and are involved in photosynthesis and phototropism (27). Vitamin B7 (biotin) is a cofactor for acetyl coenzyme A (coA) which is essential for fatty acid synthesis. Vitamin B12 (cobalamin) is required as a coenzyme in the mitochondria for many algae, yet they depend upon prokaryotes to produce it (9, 28). Thus, marine macrophytes may be auxotrophic for key vitamins, and their production by host-associated bacteria may be another basis for phototroph-microbiome interactions in nature.

Hosts might reciprocally benefit microbes, especially if heterotrophic microbes benefit from the dissolved organic carbon that is released by their hosts. Of the carbon that is fixed, kelp have been demonstrated to release 15-16% of it as dissolved organic matter (29, 30), and seagrasses too provide a constant source of dissolved organic carbon (31, 32), likely stimulating heterotrophic bacterial processes (33). These rates of organic carbon release, often involving highly labile organic carbon compounds (34), could provide the basis for reciprocal benefits between microbes and their associated hosts.

Here, we analyzed microbial metagenomes collected from 5 different coastal phototrophs to determine if there is functional genomic evidence of microbial metabolisms that could reciprocally benefit host and microbes. We analyzed the surface microbiome on the blade of two kelp species (*Laminaria setchellii* and *Nereocystis luetkeana*) and the surfgrass *Phyllospadix scouleri*, the rhizomes of *P. scouleri*, *P. serrulatus*, and the seagrass species, *Zostera marina*, and

the sediment surrounding the rhizomes of *P. scouleri* and *Z. marina*. We quantified the variable oxygen environment in the rhizomes of *Phyllospadix spp.* to determine if they allow for aerobic as well as anaerobic metabolisms. We analyzed the microbial taxa present and examined their gene content to estimate their functional and metabolic capacities. We hypothesized that microbial partners: 1) enhance host access to dissolved inorganic nitrogen through nitrogen recycling, ammonification and nitrogen fixation, 2) provision vitamins B₁, B₂, B₇, B₁₂, and 3) use a diversity of abundant dissolved organic carbon exudates from the host. We tested whether microbial taxonomy and function differed across hosts and host tissue types, and whether anaerobic metabolisms were present in low-O₂ environments (e.g., rhizomes and sediment). Through this study, we find that the range of oxygen environments engineered by host phototrophs likely explains the diversity of both oxidizing and reducing microbial metabolisms, and contributes to the functional capabilities of microbes and their influence on carbon and nitrogen cycling in nearshore ecosystems.

Methods

Sampling and DNA Extraction

We collected metagenome samples from the surfaces of 5 different phototroph species (Table S1). The surface of *Phyllospadix scouleri* blades, *Laminaria setchellii* fronds and the inner bulbs of *Nereocystis luetkeana* were swabbed with a sterile swab and brushed with an interdental brush (GUM Proxabrush Go-Betweens). We preserved sections of the rhizomes of *Phyllospadix scouleri*, *P. serrulatus* and *Zostera marina*. Sediment surrounding *P. scouleri* and *Z. marina* was also collected. All samples were collected from Tatoosh Island, WA, USA (48.393679, -124.734617) on 16-17 Jul 2019, except for *Z. marina* samples which were sampled from West

Falmouth Bay, MA, USA (41.60708333, -70.64527778) on 19 Sept 2019. We included samples from the rhizosphere of *Z. marina* from the Atlantic Ocean as a known positive control for nitrogen fixation (22, 23). Swabs, tissue and sediment were immediately frozen at 20° C and shipped to storage at -80° C. DNA from these collections was extracted with a Qiagen PowerSoil Kit and multiple samples were pooled for each metagenome sample to increase DNA quantity and possible discovery: *P. scouleri* blade, rhizome and sediment (3 pooled individuals each), *P. serrulatus* rhizome (3 individuals), *L. setchellii* blade (3 individuals), *N. luetkeana* interior bulb (4 individuals), *Z. Marina* rhizomes and sediment (2 individuals).

Shotgun metagenomic sequencing, assembly, and read recruitment

The above 8 samples were run over 2 lanes on a HiSeq 2500 (2x150) with TruSeq DNA library preps at Argonne National Laboratory. For each sample, resulting DNA sequences were first quality filtered (35)(Minoche et al. 2011), then assembled with IDBA-UD v1.1.3 (36) (Peng et al. 2012) with a minimum scaffold length of 1 kbp. Metagenomic short reads from each sample were then recruited back to their corresponding assembled contigs using Bowtie2 (37). Samtools (38) was used to generate sorted and indexed BAM files. Anvi'o v7.0 (39) was used as the command line environment for all downstream analyses. 'anvi-gen-contigs-database' was used to generate anvi'o contigs databases, during which Prodigal v2.6.3 (40) identified open reading frames, and 'anvi-run-hmms' was used to identify genes matching to archaeal and bacterial single-copy core gene collections using HMMER (41).

Reconstructing metagenome-assembled genomes (MAGs)

To reconstruct genomes from the assembled metagenomes, we used a combination of automatic binning via CONCOCT v1.1.0 (42), followed by a manual curation of each MAG as outlined by Shaiber et al. 2020 (43). Genome taxonomy was determined using GTDB v.1.3.0 (44), and 'anvi-run-scg-taxonomy'. We also inferred gene-level taxonomy using Centrifuge v1.0.4 (45) to aid manual curation.

Phylogenomic analysis of MAGs

To perform a phylogenomic analysis of our MAGs, we recovered amino acid sequences for bacterial single-copy core genes (SCGs) from each genome (except the only archaeal genome in our collection) using the program 'anvi-get-sequences-for-hmm-hits' with the parameter '--hmm-source 'Bacteria_71' on the ribosomal gene set 'Ribosomal L1-L6' and the flag '--concatenate', which independently aligned each SCG independently using Muscle v3.8.1 (46) before concatenating them into a final superalignment. We then refined the alignment using trimAl v1.4.rev15 (47) to remove any position in the alignment if more than 50% of the residues were gap characters. A maximum-likelihood phylogeny was inferred using IQTree (48) with 1,000 bootstrap replicates, and a LG+R6 model best fit our data using ModelFinder (49).

Functional analysis of microbial communities

To address the metabolic capabilities of host-associated microbes, we annotated genes in each anvi'o contigs database with 3 different databases using 'anvi-run-kegg-kofams', 'anvi-run-ncbi-cogs', and 'anvi-run-pfams', which used the databases of Kyoto Encyclopedia of Genes and Genomes (KEGG) (50), NCBI's Clusters of Orthologous Genes (COGs) (51) and EBI's Pfam database (52) respectively. We used these annotated genes to test for 1) nitrogen cycling

metabolisms, especially those within the nitrogen-fixation pathway, 2) hydrolases, including ureases, as well as ammonia-lyases, to cleave the C-N bonds in amino acids and make ammonium available to the host, 3) vitamin production, namely vitamins B₁, B₂, B₇ and B₁₂ and 4) a set of dissolved organic matter (DOM) transporter genes identified by Poretsky et al. (34) that indicate the ability of the microbial community to assimilate DOM exudates from kelps and surfgrasses. The list of genes used is indicated in Table S4. We additionally developed and used a graph-based algorithm on KEGG definitions for vitamins B₁, B₂, B₇ and B₁₂ to detect the presence of these biosynthetic pathways (Supplementary Code 1). To expand our functional analysis of kelp blade genes, we included 32 MAGs from the surface of *N. luetkeana* blades that were collected from the same location at the same time using similar methods as those described above (53).

Phylogenetic analysis of nifH genes

To search for *nifH* amino acid sequences in our environmental samples, we identified 9 MAGs which contained *nifH* genes using the KEGG identifier K02588 with e-value < 1e-20. We aligned the AA sequences for these genes against 89 well-characterized reference *nifH* AA sequences (Table S6) using Muscle v3.8.1 (46) and refined the alignment using trimAl (gap-threshold: 0.5) and ‘anvi-script-reformat-fasta’ (max-percentage-gap: 50%). A maximum-likelihood phylogeny was inferred using IQTree (48) with 1,000 bootstrap replicates, and a LG+R5 model best fit our data using ModelFinder (49). *nifH* genes from the *Zostera* samples served as positive controls to detect nitrogen fixation genes in other samples. Figures 2, 3, 4 were generated using iTol v5 (54), R v4.0.3 and FigTree respectively. We additionally took tissue samples from *P. scouleri* rhizome (n = 16), basal meristematic region just distal to the sheath

(n = 12) and blade 35 cm above the rhizome (n = 12) to quantify stable isotopes of $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ to look for signatures of nitrogen fixation (methods described in Appendix 1).

Quantifying the Oxygen Environment

We quantified the oxygen concentrations in proximity to *Phyllospadix* spp. rhizomes by comparing dissolved oxygen (DO) concentrations in the surrounding seawater and in the sediment around the rhizome. We used a Pyro Science Robust Oxygen Probe (OXROB10, Firesting™, Pyroscience), and repeated measurements around 0900h across 4 days (7-9 June 2019, 13 June 2021) within *P. scouleri* (n = 18) and *P. serrulatus* (n = 11) rhizomes. Each reading first measured the surrounding seawater after which we gently pushed the tip of the oxygen probe into the sediment and rhizome mass to a depth of 1-3 mm, the typical thickness (*pers. observation*). We let the probe equilibrate and took a reading at 150 sec. This allowed the rhizome oxygen environment to equilibrate after we disturbed the intact rhizome. We compared surrounding water and within-rhizome oxygen using paired t-tests in R.

Data Availability

In addition to the code available on GitHub (____), the final MAG database files generated in anvi'o are available on the FigShare repository: (____). Metagenomic sequence data are available at the NCBI's Sequence Read Archive under accession no. (submission in progress).

Results

Surfgrass rhizomes have lower oxygen concentrations than surrounding seawater

The oxygen environment in the rhizomes differed significantly from that of the surrounding seawater (Fig. 1). Rhizomes maintained a lower dissolved oxygen (DO) concentration than the surrounding seawater for both *P. scouleri* (n=18, pairwise t-test: $p < 0.001$) and *P. serrulatus* (n=11, pairwise t-test: $p < 0.001$). *P. serrulatus* maintained a slightly lower DO concentration in the rhizome at 2.11 mg l^{-1} , compared to 5.61 mg l^{-1} for *P. scouleri*. However, the nature of sampling likely introduced more oxygenated water from the surrounding water column to the rhizome-sediment microenvironment, suggesting that the actual DO concentration within the sediment is lower than the value reported.

Diversity of MAGs assembled across hosts

Following filtering, we obtained an average of 41 million reads per sample (range 6.48 to 67.73 million), with 79.8% of raw reads retained on average. When these reads were assembled into contigs of at least 1000 nucleotides, a mean of 42,026 contigs and a mean of 110,054 genes were present across samples (Table S1).

Across 8 metagenomes we manually binned 33 high quality MAGs, defined as having a completion score $> 90\%$ and contamination (or redundancy) $< 10\%$ (Table 2). We also identified 39 lower quality MAGs that had completion scores between 38 and 93% and redundancy scores between 0 and 21% (Table S3). All MAGs were bacterial except for a single archaeal MAG on the rhizome of *P. scouleri*. The bacterial MAGs spanned 7 phyla, including *Proteobacteria* (n=34), *Bacteroidota* (n=19), *Verrumicrobia* (n=2), *Campylobacterota* (n=3), *Desulfobacterota* (n=5), and a single MAG in each of *Desulfomonadota*, *Acidobacteriota*, and *Spirochaetota*. The Archaea belonged to the phylum *Chrenarchaeota*. There were 46 MAGs resolved to the species

level, with 8 to the genus level, 9 to family, 2 to order, and 2 to class level. Five MAGs were resolved only as Bacteria (Table S2).

The 72 MAGs belong to diverse microbial phyla, which were distributed across the 5 host species and tissue types (Fig. 2). In some cases, bacterial taxa from kelp blade tissues were most closely related to bacteria collected from the rhizome or sediment of a seagrass, suggesting that closely related bacterial taxa can associate with diverse hosts. Known anaerobic sulfur cyclers like *Desulfobulbia*, *Desulfobacteria*, *Desulfuromonadia* and *Campylobacteria* (*Sulfurovum* sp000296775 and *Sulfurimonas autotrophica*) were exclusively found in the low oxygen rhizome and sediment samples of *Zostera marina* and *Phyllospadix* spp. Conversely, *Alphaproteobacteria*, were exclusively found on surfaces exposed to the water column. *Gammaproteobacteria* was the only class found across the range of tissue types (6 out of 8 host environments). We did not include the only well-resolved archaeal taxon found in our samples, *Crenarchaea* (*P. scouleri* rhizome), as our analysis compared single-copy core genes specific to bacterial phyla.

Host-associated microbial genomes contain pathways to synthesize vitamins, recycle nitrogen, and use host-generated carbon

We found evidence for a number of metabolic pathways that are likely important for exchanges between host phototrophs and their microbial partners (Fig. 3). Microbes on hosts had genes for diverse carbohydrate and carboxylic acid assimilation via cell membrane transport proteins. Host-associated microbes also had genes for a diversity of nitrogen metabolisms, including ureases and hydrolases that could regenerate ammonium. Nitrogen metabolisms were most

diverse in rhizome and sediment samples where we identified both oxidizing (nitrification) and reducing (nitrate reduction, nitrogen fixation, denitrification) metabolisms, as well as metabolisms that both oxidize and reduce (annamox).

Every sample had at least one gene from B-vitamins biosynthesis pathways. Using a simple-path based algorithm on KEGG definitions (Supplementary Code 1), we determined that all microbial communities had the metabolic pathways to synthesize vitamins B₁ (with the exception of the *P. scouleri* rhizome), B₂ and B₇ (except inside the bulb of *N. luetkeana*). The Vitamin B₁₂ anaerobic biosynthesis pathway, however, was only present in MAGs found on the blades of *L. setchellii* (2) and *P. scouleri* (3) and the rhizomes of *P. serrulatus* (2) and *Z. marina* (1). Additionally, all three MAGs on the blade of *P. scouleri* that had this anaerobic pathway had the genes necessary to synthesize Vitamin B₁₂ aerobically as well.

Novel detection of nifH genes in surfgrass

We identified the nitrogenase gene (*nifH*) in 9 MAGs with e-value support < 1.3e-120 (KEGG) and < 1.1e-135 (COG). These 9 MAGs were assembled from *P. serrulatus* rhizomes (n = 2) and *Z. marina* rhizomes (n = 3) and the surrounding sediment (n = 4). Of these 9 MAGs, 5 were resolved to the genus level, while others were resolved to the order and family level, including *Campylobacteriales*, *Desulfobacteriales* and 2 *Flavobacteriaceae* (Fig. 4, Table S5). *nifH* genes identified in the rhizomes of *P. serrulatus* and *Z. marina* belonging to the class *Desulfobacteria* and family *Flavobacteriaceae*, clustered within Cluster III: anaerobic nitrogen-fixers that are often coupled with sulfate-reduction metabolisms. Samples from *Z. marina* sediment and rhizome also contained 3 *nifH* genes in *Campylobacterial* MAGs that clustered together in a

sister clade to the aerobic nitrogen-fixers of Cluster I. The COG gene identified as *nifH* (COG1348) also includes the homologous protochlorophyllides, which are involved in photosynthetic pigment synthesis but have high sequence similarity to the *nifH* gene (21, 55). Instead, we used the KEGG gene (K02588) that does not detect these homologs. When we inspected genes on the same contig with *nifH*, we found a number of genes related to nitrogen fixation (Table S5), including *nifD* (COG 2710) in 7 of the 9 contigs, nitrogen regulatory protein PII (COG 347), *nifB* (COG 535), and multiple iron containing proteins including ferredoxin and Fe-Mo cluster-binding proteins (Table S5).

Discussion

Phototroph tissues and sediment host distinct microbial taxa and functions

The phototroph species we sampled in this study are foundational in coastal ecosystems (56–59), yet a description of the diversity and function of their microbiomes have been lacking. All MAGs were bacterial, except for a single archaeal MAG (*Crenarchaeota*) in the rhizome of *Phyllospadix scouleri*, which was identified as *Nitrosopumilis*, a genus associated with nitrification (Table S3). Together, these 5 phototrophs hosted bacteria from 9 phyla. The only low diversity sample was the interior of the bulb of *Nereocystis*, where we assembled only a single MAG (*UBA7415 sp002470515*) suggesting that this environment of high carbon monoxide and nitrogen gas (60) may inhibit microbial activity or pose a highly selective environment. Blades of kelp and surfgrass, in contrast, were a locus of microbial diversity and function, a finding that is similar to many recent studies of macroalgal and seagrass microbiomes reporting high microbial diversity (2, 4, 5, 61–63). The functional attributes of microbial taxa associated with marine macrophytes include pathogen resistance (64), the ability to provision the

host with B vitamins (9), and enhanced host algal fitness (65), perhaps through some of the nitrogen metabolisms we documented here (14, 66).

Host-microbe interactions in a dynamic oxygen microenvironment

Grouping MAGs by microbial metabolisms (Fig. 3) showed key functional differences among phototroph hosts. Blade tissues that interacted directly with the water column were associated with microbial nitrogen metabolisms that were mostly oxidizing. The abundance of dissolved organic carbon from phototroph hosts (29–31, 59) might select for heterotrophic metabolisms. Indeed, we found an abundance of genes for dissolved organic matter assimilation and transport in all metagenomes, suggesting that hosts may stimulate heterotrophy in their associated microbial community similar to findings by Poretsky et al. (34). Improved characterization of the components of dissolved organic matter and the genomes of hosts will allow us to better assess complementarity in resource supply by hosts and resource use by microbes.

The host tissue types in this study differed in surface oxygen concentrations. Blade tissue interacts with the water column and is likely more oxygenated than rhizome tissue or sediments, though a previous study suggests there can also be a 60% reduction in oxygen along the immediate surface of kelp blades (67), and along the mucus layer where some kelp-associated bacteria reside (6). Over two-thirds of the bacterial taxa on blades of *N. luetkeana* belonged to families associated with obligately aerobic metabolisms, demonstrating the role of oxygen in shaping phototroph-associated microbial communities (68). The sediment surrounding the rhizomes of *Phyllospadix spp.* contained low oxygen microenvironments (Fig. 1) likely maintained by macroinvertebrate respiration (69)(Moulton and Hacker 2011), similar to the

biological processes in the anaerobic sediment surrounding *Zostera* (13). Low rhizosphere oxygen concentrations likely structured the taxonomic composition of *Z. marina* to include anaerobic taxa such as *Campylobacteria*, *Desulfatitaley* and *Desulfobulbus*. The presence of anaerobes like *Desulfuromonadia*, *Desulfobacteria*, *Spirochaeta* and *Aminicenantia* in *P. serrulatus* rhizomes suggests sulfate reduction also occurs, possibly coupled to dissolved organic carbon use as an energy source (e.g. (70) Howarth & Hobbie 1982). Additionally, *Campylobacteria* and the genus *Thiodiazotropha* were associated with *Z. marina* and may remove detrimental sulfide accumulation through sulfur oxidation (71, 72).

Nitrogen metabolisms that were both oxidizing and reducing were found in MAGs associated with rhizomes of both *Z. marina* and *Phyllospadix* (Fig. 3), suggesting the potential for temporal niches when, for example, ammonium oxidation to nitrate occurs during high-O₂ daylight periods, followed by nitrate reduction or nitrogen fixation during O₂-depleted nighttime hours. Additionally, all MAGs in this study contained hydrolases that cleave carbon-nitrogen bonds to produce ammonium (14), recycling nitrogen compounds for host uptake. Oxidizing and reducing metabolisms are likely separated only by microns in the hosts studied here.

We detected biosynthetic pathways for vitamins B₁, B₂, B₇ and B₁₂ that are required by the auxotrophic phototroph hosts in this study (9, 10, 73). We found that only the blades of *P. scouleri* harbored MAGs with both anaerobic and aerobic biosynthetic pathways for Vitamin B₁₂, suggesting that the variable oxygen environment driven by host-metabolism creates diverse metabolic niches for associated microbes. Strong gradients in oxygen and metabolically diverse microbial metabolisms are present in a diversity of animal hosts such as corals and sponges as a

result of host metabolism (74–76). Fluctuating oxygen microenvironments might also promote cross-feeding, where microbial taxa produce a metabolite that can be consumed by other taxa. Cross-feeding is potentially important for nitrogen (77) and carbon metabolisms (78, 79) in microbial communities.

Characteristics of previously undescribed nitrogen fixation in surfgrass

Building on recent studies that illustrate the association of nitrogen fixing microbes with a diversity of macroalgae (80) and seagrasses (22, 23, 81, 82), we found a previously undescribed diversity of nitrogenase genes associated with the surfgrass *Phyllospadix*. We detected *nifH* genes in *P. serrulatus* rhizomes that resolved into the Cluster I group of *nifH* genes, which are characterized by aerobic nitrogen fixers. *P. serrulatus*, in comparison to *P. scouleri*, is found higher up in the intertidal zone and often in sheltered tidepools that tend to undergo dramatic daily fluctuations in oxygen, possibly allowing for a temporal low-O₂ niche during the night (83). Conversely, we did not detect nitrogenase genes in the microbiome of *P. scouleri*, which inhabits more wave-exposed and thus better oxygenated environments (Fig. 1). However, stable isotope analyses across *P. scouleri* samples show a lower nitrogen isotopic signature in the rhizome compared to the rest of the plant, a possible indication of nitrogen from an atmospheric source (Fig. S1), though *in situ* experiments with stable isotope tracers are needed to confirm the presence of nitrogen fixation.

Nitrogen fixation by microbial associates provides a key means of increasing the availability of ammonium, possibly supporting primary productivity. *P. scouleri* biomass reaches 12.7 kg of wet mass per square meter of shore and exudes 0.93 mg C per hour per gram dry mass as

dissolved organic carbon that may fuel microbial activity (59). There is evidence that nitrogen fixation can contribute to seagrass productivity (66, 84), a possible adaptation to low nitrogen environments. Our finding that nitrogen fixing microbes are associated with a rocky intertidal surfgrass is especially surprising given that Tatoosh Island is in an area of upwelling and high DIN (86) at the more northerly end of the California Current Large Marine Ecosystem. Whether nitrogen fixation forms the basis for reciprocal host-microbe exchange is still unknown.

The metagenomic analyses we present here suggest that phototroph-associated microbiomes may be involved in carbon, nitrogen and vitamin metabolisms important to their hosts, likely generating commensal or mutualistic interactions. Future experiments should test these hypothesized interactions between host and microbiome. The importance of seaweeds and seagrasses to coastal productivity, and the demonstrated sensitivity of both host and microbes to increasing temperatures and pH (11, 62, 85), pathogens (61), and other anthropogenic stressors, underline the importance of further studying phototroph-microbiome interactions.

Acknowledgements

Our gratitude to the Makah Tribal Nation for access to Tatoosh Island. We thank The University of Chicago's Microbiome Center for pilot award funding, and Washington Department of Natural Resources grants 93099282, 93100399 (CAP) and NSF-DEB grant (#1556874) awarded to JT Wootton. We appreciate the work of C Saucedo in the isotope analysis, and A Wootton, A Wood and K Foreman in the field sampling. KM was supported by an EE Fellowship from The University of Chicago. S Owens and S Greenwald at Argonne National Lab provided expertise in sequencing.

References

1. McFall-Ngai M, Hadfield MG, Bosch TCG, Carey HV, Domazet-Lošo T, Douglas AE, Dubilier N, Eberl G, Fukami T, Gilbert SF, Hentschel U, King N, Kjelleberg S, Knoll AH, Kremer N, Mazmanian SK, Metcalf JL, Nealson K, Pierce NE, Rawls JF, Reid A, Ruby EG, Rumpho M, Sanders JG, Tautz D, Wernegreen JJ. 2013. Animals in a bacterial world, a new imperative for the life sciences. *Proc Natl Acad Sci USA* 110:3229–3236.
2. Weigel BL, Pfister CA. 2019. Successional Dynamics and Seascape-Level Patterns of Microbial Communities on the Canopy-Forming Kelps *Nereocystis luetkeana* and *Macrocystis pyrifera*. *Frontiers in Microbiology* 10:346.
3. Lemay MA, Martone PT, Keeling PJ, Burt JM, Krumhansl KA, Sanders RD, Wegener Parfrey L. 2018. Sympatric kelp species share a large portion of their surface bacterial communities: Kelp-associated bacterial diversity. *Environmental Microbiology* 20:658–670.
4. Quigley CTC, Capistrant-Fossa KA, Morrison HG, Johnson LE, Morozov A, Hertzberg VS, Brawley SH. 2020. Bacterial Communities Show Algal Host (*Fucus* spp.)/Zone Differentiation Across the Stress Gradient of the Intertidal Zone. *Front Microbiol* 11:563118.
5. Lemay MA, Davis KM, Martone PT, Parfrey LW. 2021. Kelp-associated Microbiota are Structured by Host Anatomy¹. *J Phycol* 57:1119–1130.
6. Ramirez-Puebla ST, Weigel BL, Jack L, Schlundt C, Pfister CA, Mark Welch JL. 2020. Spatial organization of the kelp microbiome at micron scales. preprint, *Microbiology*.
7. Jacoby R, Peukert M, Succurro A, Koprivova A, Kopriva S. 2017. The Role of Soil Microorganisms in Plant Mineral Nutrition—Current Knowledge and Future Directions. *Front Plant Sci* 8:1617.
8. Mills MM, Turk-Kubo KA, van Dijken GL, Henke BA, Harding K, Wilson ST, Arrigo

435 KR, Zehr JP. 2020. Unusual marine cyanobacteria/haptophyte symbiosis relies on N₂ fixation
436 even in N-rich environments. *The ISME Journal* <https://doi.org/10.1038/s41396-020-0691-6>.

437 9. Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, Smith AG. 2005. Algae acquire
438 vitamin B12 through a symbiotic relationship with bacteria. *Nature* 438:90–93.

439 10. Wichard T, Charrier B, Mineur F, Bothwell JH, Clerck OD, Coates JC. 2015. The green
440 seaweed *Ulva*: a model system to study morphogenesis. *Front Plant Sci* 6.

441 11. Qiu Z, Coleman MA, Provost E, Campbell AH, Kelaher BP, Dalton SJ, Thomas T,
442 Steinberg PD, Marzinelli EM. 2019. Future climate change is predicted to affect the microbiome
443 and condition of habitat-forming kelp. *Proceedings of the Royal Society B: Biological Sciences*
444 286:20181887.

445 12. Bell W, Mitchell R. 1972. Chemotactic and Growth Responses of Marine Bacteria to
446 Algal Extracellular Products. *Biological Bulletin* 143:265–277.

447 13. Brodersen KE, Siboni N, Nielsen DA, Pernice M, Ralph PJ, Seymour J, Kühl M. 2018.
448 Seagrass rhizosphere microenvironment alters plant-associated microbial community
449 composition. *Environ Microbiol* 20:2854–2864.

450 14. Tarquinio F, Bourgouire J, Koenders A, Laverock B, Sävström C, Hyndes GA. 2018.
451 Microorganisms facilitate uptake of dissolved organic nitrogen by seagrass leaves. *The ISME*
452 *Journal* <https://doi.org/10.1038/s41396-018-0218-6>.

453 15. Ladd JN, Jackson RB. 1982. Biochemistry of Ammonification, p. 173–228. *In* Stevenson,
454 FJ (ed.), *Agronomy Monographs*. American Society of Agronomy, Crop Science Society of
455 America, Soil Science Society of America, Madison, WI, USA.

456 16. Delmont TO, Quince C, Shaiber A, Esen ÖC, Lee ST, Rappé MS, McLellan SL, Lückers
457 S, Eren AM. 2018. Nitrogen-fixing populations of Planctomycetes and Proteobacteria are

abundant in surface ocean metagenomes. *Nature Microbiology* 3:804–813.

17. Sohm JA, Webb EA, Capone DG. 2011. Emerging patterns of marine nitrogen fixation. *Nat Rev Microbiol* 9:499–508.

18. Bombar D, Paerl RW, Riemann L. 2016. Marine Non-Cyanobacterial Diazotrophs: Moving beyond Molecular Detection. *Trends in Microbiology* 24:916–927.

19. Harding K, Turk-Kubo KA, Sipler RE, Mills MM, Bronk DA, Zehr JP. 2018. Symbiotic unicellular cyanobacteria fix nitrogen in the Arctic Ocean. *Proc Natl Acad Sci USA* 115:13371–13375.

20. Cabello AM, Turk-Kubo KA, Hayashi K, Jacobs L, Kudela RM, Zehr JP. 2020. Unexpected presence of the nitrogen-fixing symbiotic cyanobacterium UCYN-A in Monterey Bay, California. *J Phycol* 56:1521–1533.

21. Raymond J, Siefert JL, Staples CR, Blankenship RE. 2004. The Natural History of Nitrogen Fixation. *Molecular Biology and Evolution* 21:541–554.

22. Patriquin D, Knowles R. 1972. Nitrogen fixation in the rhizosphere of marine angiosperms. *Marine Biology* 16:49–58.

23. Capone DG. 1982. Nitrogen Fixation (Acetylene Reduction) by Rhizosphere Sediments of the Eelgrass *Zostera marina*. *Marine Ecology Progress Series* 10:67–75.

24. Cole L, McGlathery K. 2012. Nitrogen fixation in restored eelgrass meadows. *Marine Ecology Progress Series* 448:235–246.

25. Agawin N, Ferriol P, Sintes E. 2019. Simultaneous measurements of nitrogen fixation in different plant tissues of the seagrass *Posidonia oceanica*. *Mar Ecol Prog Ser* 611:111–127.

26. Aoki L, McGlathery K. 2019. High rates of N fixation in seagrass sediments measured via a direct $^{30}\text{N}_2$ push-pull method. *Mar Ecol Prog Ser* 616:1–11.

- 481 27. Massey V. 2000. The chemical and biological versatility of riboflavin. 28:283–296.
- 482 28. Grossman A. 2016. Nutrient Acquisition: The Generation of Bioactive Vitamin B 12 by
483 Microalgae. Current Biology 26:R319–R321.
- 484 29. Reed DC, Carlson CA, Halewood ER, Nelson JC, Harrer SL, Rassweiler A, Miller RJ.
485 2015. Patterns and controls of reef-scale production of dissolved organic carbon by giant kelp *M*
486 *acrocystis pyrifera*: DOC production by giant kelp. Limnology and Oceanography 60:1996–
487 2008.
- 488 30. Weigel BL, Pfister CA. 2021. The dynamics and stoichiometry of dissolved organic
489 carbon release by kelp. Ecology 102.
- 490 31. Wetzel RG, Penhale PA. 1979. Transport of carbon and excretion of dissolved organic
491 carbon by leaves and roots/rhizomes in seagrasses and their epiphytes. Aquatic Botany 6:149–
492 158.
- 493 32. Barrón C, Apostolaki ET, Duarte CM. 2014. Dissolved organic carbon fluxes by seagrass
494 meadows and macroalgal beds. Frontiers in Marine Science 1.
- 495 33. Pfister CA, Altabet MA. 2019. Enhanced microbial nitrogen transformations in
496 association with macrobiota from the rocky intertidal. Biogeosciences 16:193–206.
- 497 34. Poretsky RS, Sun S, Mou X, Moran MA. 2010. Transporter genes expressed by coastal
498 bacterioplankton in response to dissolved organic carbon. Environmental Microbiology 12:616–
499 627.
- 500 35. Minoche AE, Dohm JC, Himmelbauer H. 2011. Evaluation of genomic high-throughput
501 sequencing data generated on Illumina HiSeq and Genome Analyzer systems. Genome Biol
502 12:R112.
- 503 36. Peng Y, Leung HCM, Yiu SM, Chin FYL. 2012. IDBA-UD: a de novo assembler for

single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics* 28:1420–1428.

37. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359.

38. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The sequence alignment/map format and SAMtools. *Bioinformatics* 25:2078–2079.

39. Eren AM, Kiefl E, Shaiber A, Veseli I, Miller SE, Schechter MS, Fink I, Pan JN, Yousef M, Fogarty EC, Trigodet F, Watson AR, Esen ÖC, Moore RM, Clayssen Q, Lee MD, Kivenson V, Graham ED, Merrill BD, Karkman A, Blankenberg D, Eppley JM, Sjödin A, Scott JJ, Vázquez-Campos X, McKay LJ, McDaniel EA, Stevens SLR, Anderson RE, Fuessel J, Fernandez-Guerra A, Maignien L, Delmont TO, Willis AD. 2021. Community-led, integrated, reproducible multi-omics with anvi'o. *Nat Microbiol* 6:3–6.

40. Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119.

41. Eddy SR. 2011. Accelerated Profile HMM Searches. *PLoS Comput Biol* 7:e1002195.

42. Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, Lahti L, Loman NJ, Andersson AF, Quince C. 2014. Binning metagenomic contigs by coverage and composition. *Nat Methods* 11:1144–1146.

43. Shaiber A, Willis AD, Delmont TO, Roux S, Chen L-X, Schmid AC, Yousef M, Watson AR, Lolans K, Esen ÖC, Lee STM, Downey N, Morrison HG, Dewhirst FE, Mark Welch JL, Eren AM. 2020. Functional and genetic markers of niche partitioning among enigmatic members

527 of the human oral microbiome. *Genome Biol* 21:292.

528 44. Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil P-A, Hugenholtz P. 2021.

529 GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically

530 consistent, rank normalized and complete genome-based taxonomy. *Nucleic Acids Research*

531 gkab776.

532 45. Kim D, Song L, Breitwieser FP, Salzberg SL. 2016. Centrifuge: rapid and sensitive

533 classification of metagenomic sequences. *Genome Res* 26:1721–1729.

534 46. Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high

535 throughput. *Nucleic Acids Research* 32:1792–1797.

536 47. Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. 2009. trimAl: a tool for automated

537 alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25:1972–1973.

538 48. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A Fast and

539 Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular*

540 *Biology and Evolution* 32:268–274.

541 49. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017.

542 ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods* 14:587–

543 589.

544 50. Kanehisa M. 2000. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids*

545 *Research* 28:27–30.

546 51. Galperin MY, Makarova KS, Wolf YI, Koonin EV. 2015. Expanded microbial genome

547 coverage and improved protein family annotation in the COG database. *Nucleic Acids Research*

548 43:D261–D269.

549 52. Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL,

550 Tosatto SCE, Paladin L, Raj S, Richardson LJ, Finn RD, Bateman A. 2021. Pfam: The protein
551 families database in 2021. *Nucleic Acids Research* 49:D412–D419.

552 53. Weigel BL, Miranda KK, Fogarty EC, Watson AR, Pfister CA. in review. Functional
553 insights into the kelp microbiome from metagenome assembled genomes.

554 54. Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for
555 phylogenetic tree display and annotation. *Nucleic Acids Research* 49:W293–W296.

556 55. Kapili BJ, Dekas AE. 2021. PPIT: an R package for inferring microbial taxonomy from
557 *nifH* sequences. *Bioinformatics* 37:2289–2298.

558 56. Shelton AO. 2010. Temperature and community consequences of the loss of foundation
559 species: Surfgrass (*Phyllospadix* spp., Hooker) in tidepools. *Journal of Experimental Marine*
560 *Biology and Ecology* 391:35–42.

561 57. Lefcheck JS, Wilcox DJ, Murphy RR, Marion SR, Orth RJ. 2017. Multiple stressors
562 threaten the imperiled coastal foundation species eelgrass (*Zostera marina*) in Chesapeake Bay,
563 USA. *Glob Change Biol* 23:3474–3483.

564 58. Pfister CA, Altabet MA, Weigel BL. 2019. Kelp beds and their local effects on seawater
565 chemistry, productivity, and microbial communities. *Ecology* <https://doi.org/10.1002/ecy.2798>.

566 59. Miranda KK, Weigel BL, McCoy SJ, Pfister CA. 2021. Differential impacts of alternate
567 primary producers on carbon cycling. *Ecology* <https://doi.org/10.1002/ecy.3455>.

568 60. Liggan LM, Martone PT. 2018. Under pressure: biomechanical limitations of developing
569 pneumatocysts in the bull kelp (*Nereocystis luetkeana* , Phaeophyceae). *Journal of Phycology*
570 54:608–615.

571 61. Egan S, Harder T, Burke C, Steinberg P, Kjelleberg S, Thomas T. 2013. The seaweed
572 holobiont: understanding seaweed–bacteria interactions. *FEMS Microbiology Reviews* 37:462–

573 476.

574 62. Minich JJ, Morris MM, Brown M, Doane M, Edwards MS, Michael TP, Dinsdale EA.

575 2018. Elevated temperature drives kelp microbiome dysbiosis, while elevated carbon dioxide

576 induces water microbiome disruption. PLOS ONE 13:e0192772.

577 63. Capistrant-Fossa KA, Morrison HG, Engelen AH, Quigley CTC, Morozov A, Serrão EA,

578 Brodie J, Gachon CMM, Badis Y, Johnson LE, Hoarau G, Abreu MH, Tester PA, Stearns LA,

579 Brawley SH. 2021. The microbiome of the habitat-forming brown alga *Fucus vesiculosus*

580 (Phaeophyceae) has similar cross-Atlantic structure that reflects past and present drivers¹. J

581 Phycol jpy.13194.

582 64. Li J, Weinberger F, Saha M, Majzoub ME, Egan S. 2021. Cross-Host Protection of

583 Marine Bacteria Against Macroalgal Disease. Microb Ecol [https://doi.org/10.1007/s00248-021-](https://doi.org/10.1007/s00248-021-01909-2)

584 01909-2.

585 65. Burgunter-Delamare B, KleinJan H, Frioux C, Fremy E, Wagner M, Corre E, Le Salver

586 A, Leroux C, Leblanc C, Boyen C, Siegel A, Dittami SM. 2020. Metabolic Complementarity

587 Between a Brown Alga and Associated Cultivable Bacteria Provide Indications of Beneficial

588 Interactions. Front Mar Sci 7:85.

589 66. Mohr W, Lehnen N, Ahmerkamp S, Marchant HK, Graf JS, Tschitschko B, Yilmaz P,

590 Littmann S, Gruber-Vodicka H, Leisch N, Weber M, Lott C, Schubert CJ, Milucka J, Kuypers

591 MMM. 2021. Terrestrial-type nitrogen-fixing symbiosis between seagrass and a marine

592 bacterium. Nature <https://doi.org/10.1038/s41586-021-04063-4>.

593 67. Noisette F, Hurd C. 2018. Abiotic and biotic interactions in the diffusive boundary layer

594 of kelp blades create a potential refuge from ocean acidification. Functional Ecology 32:1329–

595 1342.

68. Weigel BL, Pfister CA. 2020. Oxygen metabolism shapes microbial settlement on photosynthetic kelp blades compared to artificial kelp substrates. *Environmental Microbiology Reports* 1758-2229.12923.
69. Moulton O, Hacker S. 2011. Congeneric variation in surfgrasses and ocean conditions influence macroinvertebrate community structure. *Mar Ecol Prog Ser* 433:53–63.
70. Howarth RW, Hobbie JE. 1982. THE REGULATION OF DECOMPOSITION AND HETEROTROPHIC MICROBIAL ACTIVITY IN SALT MARSH SOILS: A REVIEW, p. 183–207. *In* *Estuarine Comparisons*. Elsevier.
71. Martin BC, Middleton JA, Fraser MW, Marshall IPG, Scholz VV, Hausl B, Schmidt H. 2020. Cutting out the middle clam: lucinid endosymbiotic bacteria are also associated with seagrass roots worldwide. *ISME J* 14:2901–2905.
72. Keller AH, Schleinitz KM, Starke R, Bertilsson S, Vogt C, Kleinstaub S. 2015. Metagenome-Based Metabolic Reconstruction Reveals the Ecophysiological Function of Epsilonproteobacteria in a Hydrocarbon-Contaminated Sulfidic Aquifer. *Front Microbiol* 6.
73. Helliwell KE. 2017. The roles of B vitamins in phytoplankton nutrition: new perspectives and prospects. *New Phytol* 216:62–68.
74. Babbin AR, Tamasi T, Dumit D, Weber L, Rodríguez MVI, Schwartz SL, Armenteros M, Wankel SD, Apprill A. 2021. Discovery and quantification of anaerobic nitrogen metabolisms among oxygenated tropical Cuban stony corals. *ISME J* 15:1222–1235.
75. Fiore CL, Jarett JK, Olson ND, Lesser MP. 2010. Nitrogen fixation and nitrogen transformations in marine symbioses. *Trends in Microbiology* 18:455–463.
76. Morris RL, Schmidt TM. 2013. Shallow breathing: bacterial life at low O₂. *Nature Reviews Microbiology* 11:205–212.

619 77. Lilja EE, Johnson DR. 2019. Substrate cross-feeding affects the speed and trajectory of
620 molecular evolution within a synthetic microbial assemblage. BMC Evol Biol 19:129.

621 78. de Jesús Astacio LM, Prabhakara KH, Li Z, Mickalide H, Kuehn S. 2020. Closed
622 microbial communities self-organize to persistently cycle carbon. preprint, Ecology.

623 79. Goldford JE, Lu N, Bajić D, Estrela S, Tikhonov M, Sanchez-Gorostiaga A, Segrè D,
624 Mehta P, Sanchez A. 2018. Emergent simplicity in microbial community assembly. Science
625 361:469–474.

626 80. Hamersley MR, Sohm JA, Burns JA, Capone DG. 2015. Nitrogen fixation associated
627 with the decomposition of the giant kelp *Macrocystis pyrifera*. Aquatic Botany 125:57–63.

628 81. McGlathery K, Risgaard-Petersen N, Christensen P. 1998. Temporal and spatial variation
629 in nitrogen fixation activity in the eelgrass *Zostera marina* rhizosphere. Mar Ecol Prog Ser
630 168:245–258.

631 82. Garcias-Bonet N, Arrieta JM, Duarte CM, Marbà N. 2016. Nitrogen-fixing bacteria in
632 Mediterranean seagrass (*Posidonia oceanica*) roots. Aquatic Botany 131:57–60.

633 83. Steunou A-S, Jensen SI, Brecht E, Becraft ED, Bateson MM, Kilian O, Bhaya D, Ward
634 DM, Peters JW, Grossman AR, Kühl M. 2008. Regulation of nif gene expression and the
635 energetics of N₂ fixation over the diel cycle in a hot spring microbial mat. ISME J 2:364–378.

636 84. Cardini U, van Hoytema N, Bednarz VN, Al-Rshaidat MMD, Wild C. 2018. N₂ fixation
637 and primary productivity in a red sea *Halophila stipulacea* meadow exposed to seasonality: N₂
638 fixation in *Halophila stipulacea*. Limnol Oceanogr 63:786–798.

639 85. Agawin NSR, Gil Atorrasagasti MG, Frank Comas A, Fernández-Juárez V, López-
640 Alforja X, Hendriks IE. 2021. Response of the seagrass *POSIDONIA OCEANICA* and its associated
641 N₂ fixers to high business-as-usual climate change scenario in winter. Limnol Oceanogr

642 lno.11758.

643

644 Table 1. Summary of the features of 8 metagenomes. More information is in Table S1 and the
645 taxonomy based on single copy genes is in Table S2.

<i>Phyllospadix</i> <i>scouleri</i>			<i>Phyllospadix</i> <i>serrulatus</i>	<i>Laminaria</i> <i>setchellii</i>	<i>Nereocystis</i> <i>luetkeana</i>	<i>Zostera</i> <i>marina</i>	
Sediment	Rhizome	Blade	Rhizome	Blade	Inner bulb	Sediment	Rhizome
# quality reads (in millions)							
43.68	67.73	38.41	37.99	48.58	6.48	19.37	65.76
Bacteria							
63.7%	58.3%	63.6%	63.0%	63.9%	62.2%	33.6%	60.6%
Archaea							
34.2%	38.3%	33.1%	33.3%	32.7%	35.9%	62.1%	34.2%

646

647

Table 2. Metagenome assembled genomes across all samples and their representation across phyla. More detailed information on the MAGs can be found in Table S3.

<i>Phyllospadix scouleri</i>			<i>Phyllospadix serrulatus</i>	<i>Laminaria setchellii</i>	<i>Nereocystis luetkeana</i>	<i>Zostera marina</i>	
Sediment	Rhizome	Blade	Rhizome	Blade	Inner bulb	Sediment	Rhizome
High Quality MAGs							
3	1	7	6	9	1	2	5
Other MAGs							
2	2	8	7	7	0	5	7
Proteobacteria							
2	-	9	2	10	1	5	5
Bacteroidota							
3	2	5	4	4	-	-	1
Verrucomicrobia							
-	-	-	-	2	-	-	-
Campylobacterota							
-	-	-	-	-	-	2	1
Desulfobacterota							
-	-	-	2	-	-	-	3
Desulfuromonadota							
-	-	-	1	-	-	-	-
Acidobacteriota							
-	-	-	1	-	-	-	-
Spirochaetota							
-	-	-	1	-	-	-	-
No ID							
-	-	1	2	-	-	-	2
Crenarchaeota							
-	1	-	-	-	-	-	-

Figure Captions

Figure 1. Boxplot comparing the dissolved oxygen concentrations of water column (blue) and the sediment-rhizome environment (red) of *P. scouleri* (pairwise t-test: $p < 0.001$) and *P. serrulatus* (pairwise t-test: $p < 0.001$). Sampling dates are represented by different colors.

Figure 2. A phylogenomic tree of 6 concatenated bacterial single-copy core ribosomal genes from 71 bacterial MAGS across 8 samples, showing the results from 33 high quality MAGs and 38 lower quality ones. One MAG, PSC_RHZ_Bin_00003, from the rhizome of *P. scouleri*, was identified as an archaeal genome and was thus omitted from this tree. Gaps in class, family and genus indicate the level to which taxonomic classification was resolved in each MAG. All blade tissues have 'water column exposure', while rhizome and sediment samples do not.

Figure 3. Microbial Metabolisms in the MAGs reported in Fig. 2 and Table S3 across all hosts and grouped as those that might benefit the host ("hosts benefit") and microbial metabolisms that might utilize host provisioned metabolites ("microbes benefit"). Each tick along the x-axis corresponds to a MAG. *N. luetkeana* blade MAGs are from Weigel et al. (in review). The metabolisms for **DOC Uptake** that benefit microbes are shown as a heatmap of the count of the number of genes that can metabolize *Compatible Solutes*, *Carboxylic Acids*, *Carbohydrate Pentoses* and *General Carbohydrates*. Microbial metabolisms that benefit the host are **Ammonification Hydrolases**, where the heatmap provides a count of the hydrolases acting on C-N bonds other than peptide bonds, **Nitrogen Metabolisms** and **Vitamin Synthesis**, both shown as the presence or absence of a gene in a pathway. The genes used in this are in Table S4.

675 **Figure 4.** A phylogenomic tree of *nifH* genes found on the rhizomes of *P. serrulatus* (PSE,
676 n = 3) and the rhizomes and surrounding sediment of *Z. marina* (ZMA, n = 2 and 5,
677 respectively). Some *nifH* genes group into Cluster I, including a sulfur oxidizing taxon on the
678 rhizome of *Z. marina*, and other taxa in *Campylobacterota*, including *Sulfurovum*. Cluster III
679 contains taxa associated with rhizomes including rhizomes including *Desulfobulbus*
680 *mediterraneus* on *P. serrulatus* and a Desulfobacterales associated with *Z. marina* rhizomes.
681

Figure 1

Figure 3

Figure 4

Supplementary Files

Table S1. A summary of eight metagenomes from five macrophyte taxa.

Table S2. A summary of taxonomy of MAGs

Table S3. The features of 72 metagenome assembled genomes (MAGs).

Table S4. Genes used to generate Fig. 3

Table S5. The features of *nifH* genes found in MAGs.

Table S6. *nifH* reference amino acid sequences

Appendix 1. Additional methods to quantify carbon and nitrogen stable isotopes in *P. scouleri*

Figure S1. Stable isotope analysis of $\delta^{13}C$ and $\delta^{15}N$ at blade tip, meristem, rhizome of *P.*

scouleri. From blade tip to rhizome, water flow and thus elemental mixing reduces due to

attenuation and boundary layer effects of surfgrass canopy. Assuming elemental uptake occurs

from the same pools of C and N, the lower the extent of mixing, the heavier the isotopic

signature should be at that point of the plant. This is observed with $\delta^{13}C$ which gets heavier

from the tip to the blade. This is observed with $\delta^{15}N$ till the meristem after which it lightens.

This is probably occurring as nitrogen is taken up from a different pool of nitrogen from that

around the blade/meristem. This different pool is probably made available through n-fixation