Critical-like bistable dynamics in the resting-state human brain

Sheng H. Wang^{1,2,3,4,†}, Gabriele Arnulfo^{1,5}, Vladislav Myrov^{1,4}, Felix Siebenhühner¹, Lino Nobili^{6,7,8}, Michael Breakspear⁹, Satu Palva^{1,10}, J. Matias Palva^{1,4,10,†}

1. Neuroscience Center, Helsinki Institute of Life Science (HiLIFE), University of Helsinki, Finland.
2. Doctoral Programme Brain & Mind, University of Helsinki, Finland.
3. BioMag laboratory, HUS Medical Imaging Center, Helsinki, Finland.
4. Department of Neuroscience and Biomedical Engineering, Aalto University, Finland.
5. Department of Informatics, Bioengineering, Robotics and System engineering, University of Genoa, Genoa, Italy.
6. Department of Neurosciences, Rehabilitation, Ophthalmology, Genetics and Maternal and Children's Sciences, University of Genoa, Genoa, Italy.
7. Child Neuropsychiatry Unit, IRCCS Istituto Giannina Gaslini, Genoa, Italy.
8. Centre of Epilepsy Surgery "C. Munari", Department of Neuroscience, Niguarda Hospital, Milan, Italy.
9. College of Engineering, Science and Environment; College of Health and Medicine, University of Newcastle, Callaghan, Australia.
10. Centre for Cognitive Neuroimaging, Institute of Neuroscience & Psychology, University of Glasgow, United Kingdom.

† Correspondence should be addressed to: sheng.wang@helsinki.fi and matias.palva@helsinki.fi

Abstract

Brain activity exhibits scale-free avalanche dynamics and power-law long-range temporal correlations (LRTCs) across the nervous system. This has been thought to reflect “brain criticality”, *i.e.*, brains operating near a critical phase transition between disorder and excessive order. Neuronal activity is, however, metabolically costly and may be constrained by activity-limiting mechanisms and resource depletion, which could make the phase transition discontinuous and bistable. Observations of bistability in awake human brain activity have nonetheless remained scarce and its functional significance unclear. First, using computational modelling where bistable synchronization dynamics emerged through local positive feedback, we found bistability to occur exclusively in a regime of critical-like dynamics. We then assessed bistability *in vivo* with resting-state magnetoencephalography and stereo-encephalography. Bistability was a robust characteristic of cortical oscillations throughout frequency bands from δ (3–7 Hz) to high- γ (100–225 Hz). As predicted by modelling, bistability and LRTCs were positively correlated. Importantly, while moderate levels of bistability were positively correlated with executive functioning, excessive bistability was associated with epileptic pathophysiology and predictive of local epileptogenicity. Critical bistability is thus a salient feature of spontaneous human brain dynamics in awake resting-state and is both functionally and clinically significant. These findings expand the framework of brain criticality and show that critical-like neuronal dynamics *in vivo* involves both continuous and discontinuous phase transitions in a frequency-, neuroanatomy-, and state-dependent manner.

Abbreviations

BiS: bistability index

DFA: the scaling exponent obtained with detrended fluctuation analysis is an estimate of LRTCs

EZ: epileptogenic zone

nEZ: non-EZ, areas outside of the epileptogenic zone

κ : coupling strength between oscillators in the Kuramoto model

LRTCs: long-range temporal correlations

MEG: magnetoencephalography

ρ : the strength of the state-dependent noise in the Kuramoto model

R: the order parameter of the Kuramoto model

SEEG: stereo-EEG

1 Introduction

2 Since Newton and Leibniz, differential equations have been used to describe natural
3 phenomena that manifest *continuous* and smooth temporal evolution. Nonetheless, this classic
4 approach fails in modelling many dynamics, particularly in biology and neuroscience, that show
5 *discontinuity* and abrupt divergence into discrete states over time^{1,2}. Catastrophic events
6 emerging in complex systems, such as disasters in ecosystems or epileptic seizures in the brain,
7 comprise an important subcategory of discontinuous phenomena and attract inter-disciplinary
8 research to mitigate their detrimental consequences and to identify the underlying
9 mechanisms¹⁻³.

10 Neuronal population oscillations and their synchronization reflect rhythmical fluctuations in
11 cortical excitability and regulate neuronal communication^{4,5}. The “brain criticality hypothesis”
12 posits that the brain, like many complex systems, operate near a “critical” point of a *continuous*
13 transition⁶ between asynchronous and fully synchronous activity⁷⁻¹⁰. Operation near such a
14 critical point endows the system with moderate mean synchronization, emergent power-law
15 spatio-temporal correlations, and many functional benefits such as maximal dynamic range¹¹,
16 communication¹², processing¹³, and representational capacity^{14,15}. Conversely, inadequate or
17 excessive synchrony are incompatible with healthy brain functions^{6,16} and represent coma⁻¹⁷ and
18 seizure-like states¹⁸, respectively.

19 However, the classic brain criticality hypothesis does not offer an explanation to neuronal
20 bistability, *i.e.*, discontinuous transitions between asynchronous and fully synchronous activity.
21 Bistability *per se* is a well known phenomenon in neurophysiological dynamics and is salient, for
22 example, in slow oscillations with up- and down-states observable across scales from intra-
23 cellular^{19,20} to local-field potentials (LFP)^{21,22} in animal brains. In the human brain, while there
24 are several lines of *in vivo* evidence for “critical-like” brain dynamics near a continuous phase
25 transition^{7,8,10,23}, evidence for discontinuous transitions, *i.e.*, bistable criticality, in awake resting-
26 state brain dynamics has remained scarce.

27 Neuronal bistability in awake humans has only been reported by in a single series of
28 electroencephalography (EEG) studies that reported bistable switching of alpha oscillations
29 between putatively quiescent and a hyper-synchronized states²⁴⁻²⁶. Studies of whole-brain
30 cortical activity in resting-state functional magnetic resonance imaging data (fMRI) also suggest
31 spontaneous bistable switching between synchronous and asynchronous, or between integrated
32 and segregated dynamics, respectively^{27,28}. The underlying neuronal activity substrates at these
33 multi-second time scales have, however, remained unclear.

34 Theoretical studies posit that a high degree of bistability is universally indicative of catastrophic
35 shifts^{1,2,29,30}. Hence, even if moderate bistability could characterize healthy brain dynamics, we
36 hypothesize that high bistability in neuronal synchrony would be indicative of a shift from
37 healthy to a pathological regime where neuronal populations abruptly jump between
38 asynchronous and hyper-synchronized, seizure-prone states.

39 In this study, we asked whether awake resting-state human brain exhibits critical-like bistable
40 dynamics. We first used generative modelling to establish how varying degree of bistable

dynamics emerges as a consequence of introducing a slow positive local feedback²⁵ that is conceptually equivalent to increasing demands for limited resources³¹. We then analyzed a large body of magnetoencephalography (MEG) and intracerebral stereo-EEG (SEEG) recordings of resting-state human brain activity. In both MEG and SEEG, we found that anatomically and spectrally widespread bistability characterized neuronal oscillations from δ (3–7 Hz) to high- γ (100–225 Hz) frequencies. In MEG, moderate resting-state bistability was correlated positively with executive functions. In SEEG, conversely, excessive resting-state bistability was co-localized with the epileptogenic zone and thereby associated with the pathophysiology underlying epilepsy. Bistable criticality thus constitutes a pervasive and functionally significant feature of awake resting-state brain dynamics.

Results

State-dependent noise induces bistable criticality *in silico*

To assess the emergence of bistability and its relationship with critical-like dynamics, we simulated a variant of the classic Kuramoto model, a simple generative model of synchronization dynamics³² (Supplementary Methods). Briefly, the conventional Kuramoto model has a single control parameter, κ , that defines the coupling strength between neuronal oscillators. Higher κ leads to stronger synchrony among the oscillators that is typically quantified with “order”, R . Here, we introduced a second parameter ρ that scales state-dependent noise via local positive feedback in a manner that is conceptually equivalent to the state-dependency in the stochastic Hopf bifurcation^{25,33,34}, or the slow resource-loading mechanisms leading to self-organized bistability³⁵.

At small values of ρ , the model behaved similarly to a conventional critical-like system with a continuous second-order phase transition where a gradual increase of κ results in a monotonic increase of order (Fig 1A). At moderate order, *i.e.*, at the phase transition between low and high order, power-law long-range temporal correlations (LRTCs)³⁶ emerged in model order fluctuations and delineated a critical regime (Fig 1B). Here, LRTCs were quantified using the detrended fluctuation analysis (DFA) of the order time series (Supplementary Methods).

With increasing ρ values, the model synchronization dynamics became progressively bistable (Fig 1C) as evidenced by increasing values of the bistability index (BiS), an index of the relative fit of a bimodal versus a unimodal probability distribution (*pdf*) to the time series of squared order (R^2 , comparable to oscillation powers, see Supplementary). We found bistable dynamics exclusively within the critical regime (Fig. 1D). The presence of a bistable/discontinuous transition was also evident in the sudden increase in the order parameter at the critical value and the sharp peak in the DFA in contrast to the continuous transition (Fig 1E–F) and the representative time series (Fig 1H). Bistable dynamics at high ρ values thus likely reflect a first- rather than second-order phase transition. These *in silico* findings show that even in a minimal model, synchronization of oscillators may exhibit a continuum between classic and bistable critical dynamics under the influence of state-dependent noise via local feedback.

Fig 1. Bistability is caused by elevated state-dependent noise. (A) Kuramoto model order parameter (R), (B) Detrended fluctuation analysis exponent (DFA) – an estimate of LRTC, and (C) Bistability index (BiS) as functions of noise state-dependency (ρ) and the intrinsic control parameter (κ). Each pixel is the mean of 50 independent model realizations. (D) Summary of overlapping regimes based on observation from (B–D), i.e., the classic criticality is associated with small ρ (black dashed line) whereas bistable criticality is caused by mid-to-high degree of ρ (red dashed line); (E) Probability density of R in both normal and bistable criticality is in line with the Hopf bifurcation (see Supplementary). DFA peaks (black line) coincide with the phase transition. (F) Probability density (pdf) and (G) power-law scaling of the DFA fluctuation functions in classic and bistable critical regime marked in (E), colour coded. (H) Exemplary order parameter time series; insets in are the moments of Kuramoto oscillators (black dots) in low-, mid- and high-synch state (red vectors).

79 **Bistable criticality characterizes brain dynamics *in vivo***

80 We next assessed the presence of bistability and critical dynamics in meso- and macroscopic
81 human brain activity in 10-minute resting-state recordings intracranially via stereo-
82 electroencephalography (SEEG, $N = 64$) and source-reconstructed magnetoencephalography
83 (MEG, $N = 18$), respectively. We first restricted analysis of the SEEG to neocortical grey matter
84 contacts outside of the epileptogenic zone (EZ) (Fig 2–4). Although the anatomical sampling
85 with SEEG is heterogenous across patients, the present cohort size yielded essentially a full
86 coverage of the cerebral cortex (Supplementary Fig 2). We estimated LRTCs using DFA and
87 bistability with BiS for narrow-band SEEG and MEG source amplitude time series that
88 predominantly reflect local cortical synchronization dynamics.

89 **Bistability is anatomically widespread and spectrally prevalent**

90 Visual inspection of narrow-band MEG and SEEG amplitude time series revealed salient
91 examples of bistability as intermittent switching between low- and high-amplitude oscillations
92 (Fig 2 A–B, for examples of model fitting for DFA and BiS estimates see Supplementary Fig 3).
93 Statistical testing showed that both MEG-source signals and SEEG-electrode-contact LFP signals
94 exhibited significant ($p < 0.05$, see Supplementary Fig 4) bistability and LRTCs across broad

95 frequencies (Fig 2C–E). MEG showed a peak DFA and BiS estimates in the alpha (~11 Hz)
 96 frequency band whereas in SEEG, the BiS peak extended over δ (2–4 Hz), θ (4–7.6 Hz), and α
 97 (10–13 Hz) bands (Fig 2F–G). In SEEG, DFA and BiS estimates were overall stronger and occurred
 98 across more frequencies than in MEG (Fig 2E, F–H).

Fig 2. Bistability and LRTCs are robust, large-scale phenomena in MEG and SEEG. (A) Five minutes of broad band (top) and narrow-band filtered (11 Hz bottom) power (R^2) time series from a MEG parcel located in visual area (Vis) in one subject; (B) Comparable output from five minutes of SEEG contact recorded from middle frontal gyrus (mFG) in one patient; insets: evidence of bistability as narrow-band traces switching between “up” and “down” states. (C) Group-level probability (z-axis) distribution of narrow-band (y-axis) mean amplitude (R), (D) DFA exponents, and (E) BiS estimates; data were pooled over all non-EZ SEEG contacts and MEG parcels; subject and contact/parcel number indicated in (C); black lines indicate mean of real data and red dashed lines are 99%-tile of surrogate observation. (F–H) Examples of narrow-band DFA and BiS probability distribution as indicated by colored arrows in (D–E).

Neuroanatomical structure of bistability and LRTCs

We next characterized the neuroanatomical structure of bistability and inspected its anatomical relationship with LRTCs across frequencies. We first collapsed narrow-band BiS and DFA estimates of MEG parcels (400) and SEEG contacts into a standard atlas of 100 cortical parcels (see Supplementary Fig 4). Next, the neuroanatomical similarity within and between bistability and LRTCs were assessed by computing all-to-all Spearman's correlations between narrow-band parcel BiS and DFA estimates.

Both MEG and SEEG showed high anatomical similarity between neighbouring frequencies. Correlations between slow and fast rhythms were negative in MEG (Fig 3 A) and weak in SEEG (Fig 3B). This indicates that regions tended to show bistability and criticality in a cluster of high or low frequencies, but not both. Based on these neuroanatomical similarities (red boxes, Fig 3 A–B, see also Supplementary Fig 5–6), we collapsed narrow-band BiS and DFA estimates into θ – α (5.4–11Hz) and γ -band (45–225 Hz) for further analyses (Fig 3C). The partitioning of β (15–30 Hz) band was not consistent and thus was not included (Supplementary Fig 5C).

MEG and SEEG cortical maps of θ – α band bistability revealed distinct neuroanatomical features. In MEG, visual (VIS), somatomotor (SM) and dorsal attention network (DAN) (Fig 3 C–D) exhibited greater BiS than expected by chance ($p < 0.05$, two-tailed permutation test, 10^5 permutations, not corrected for multiple comparisons, Supplementary Fig 9A–B). SEEG show high BiS in fronto-parietal (FP), ventral attention (VAN), default network (DEF), and limbic (LIM) systems (Fig 3C, E). Although comparable to the values in found in MEG, VIS showed the lowest BiS in SEEG ($p < 0.05$, two-tailed permutation test, 10^5 permutations, not corrected for multiple comparisons, Supplementary Fig 9C–D).

A Kruskal-Wallis test for variance among subjects' BiS and DFA estimates within each Yeo system revealed that in SEEG, individuals showed different levels of BiS and DFA estimates between systems (Fig 3F) with bistability greater in DEF, FP, and LIM than in VIS and SM (unpaired t-test, $p < 0.05$, FDR corrected, Supplementary Fig9 C–D). There was no statistically significant regional variation in MEG data.

In both MEG and SEEG, group-average parcel bistability was correlated with LRTCs (Fig 3 G–H, see also Supplementary Fig 7). We validated this analysis in narrow-band frequencies and found the results to converge well (Supplementary Fig 6). To further validate this relationship, we averaged parcel BiS and DFA within subjects for each Yeo system and found that the subject BiS were indeed correlated with their DFA estimates on systems-level (Fig 3I, Supplementary Fig 8).

Fig 3. Bistability and LRTC were coexisting, correlated phenomena in MEG and SEEG. Neuroanatomical similarity (Spearman's correlation) between group-average narrow-band BiS and DFA estimates of (A) MEG and (B) SEEG in Schaefer 100-parcel atlas; red boxes indicate frequency clusters showing high similarity. (C) Narrow-band group-averaged estimates were collapsed into θ - α (5.4–11Hz) and γ (40–225Hz) band based on similarity shown in (A); white-out columns in SEEG data indicate excluded parcels due to insufficient sampling (Supplementary). (D) Parcel-wise MEG group-average θ - α band BiS estimates presented in the cortex. (E) The same for SEEG group-average θ - α band BiS estimates. (F) Kruskal-Wallis one-way analysis of variance for group-level differences in DFA and BiS estimates between Yeo systems. Dashed line indicates $-\log_{10}(p \text{ value}) > 1.3$, i.e., $p < 0.05$. Correlations between group-average parcel BiS and DFA estimates in θ - α (cross) and γ band (circles) in (G) MEG and (H) SEEG, $-\log_{10}(p) > 5$, FDR corrected (Supplementary Fig 7). (I) Spearman's correlations between within-subject-average BiS and DFA estimates in Yeo systems (subject N_{MEG} per system = 18; N_{SEEG} per system = 50 ± 9.4 , range: 36–60, variable SEEG subject N per system due to heterogeneous spatial sampling).

Bistability is functionally significant in healthy subjects

We next asked whether bistability and LRTCs would predict individual differences in cognition. We assessed working memory, attention, and executive functions with neuropsychological tests (Methods). We averaged the BiS and DFA estimates across the cortical parcels to obtain four subject-specific neurophysiological estimates: $DFA_{\theta-\alpha}$, DFA_{γ} , $BiS_{\theta-\alpha}$, and BiS_{γ} (Supplementary Fig 10B), and correlated these against neuropsychological test scores. We found that $\theta-\alpha$ band BiS and DFA estimates were negatively correlated ($p < 0.05$, FDR corrected) with the Zoo Map Test Execution Time (Fig 4A–D, see also Supplementary Fig 10C–D). Thus the subjects with greater $\theta-\alpha$ band bistability and stronger LRTCs executed faster in this flexible planning task, which is well in line with prior observations linking LRTCs with cognitive flexibility³⁷.

To inspect the neuro-behavioural correlations in greater anatomical detail, we computed Spearman's correlations between neuropsychological scores and individual parcel BiS and DFA estimates. A large fraction of the cortex showed significant neuro-behavioural correlations of $\theta-\alpha$ band BiS and DFA estimates with the Zoo Map Time test but not with other neuropsychological scores (Fig 4E and G, see also Supplementary Fig 10E–F). The correlations of Zoo Map Time test with DFA estimates were most pronounced in fronto-parietal, limbic, somatosensory and, and visual areas (Fig 4F), whereas the correlations with BiS estimates were widespread across the cortex (Fig 4H).

Fig 4. Executive functions were correlated with $\theta-\alpha$ band DFA and BiS estimates in MEG subjects. (A) Spearman correlation between subject neuropsychological test scores and within subject mean parcel $\theta-\alpha$ band DFA and (B) BiS estimates collapsed over parcels; dashed lines indicate 5% and 95%-tile of correlations for surrogate data ($N_{\text{surrogate}} = 10^5$, FDR corrected, p -values see Supplementary Fig 10). (C) Scatter plots showing subject Zoo map time test and corresponding $\theta-\alpha$ band parcel-collapsed DFA and (D) BiS estimates. Each marker in (C–D) stands for one subject. (E) Fraction of significant parcels that showed significant correlation between neuropsychological test scores and individual parcel $\theta-\alpha$ band DFA and (G) BiS Estimates ($p < 0.05$, FDR corrected, details see S.Fig 10). (F) Parcels showing significant correlations between Zoom map time scores and $\theta-\alpha$ band DFA and (H) BiS estimates.

Excessive bistability characterizes the epileptogenic zone

Excessive bistability may predispose complex systems to catastrophic events^{29,38}. Under the influence of strong state-dependent noise, our model demonstrated increased sensitivity to coupling strength (Fig 1E), which suggests that strong bistability could be an early sign of shift towards supercritical hypersynchronization events, *i.e.*, epileptic seizures. We thus asked whether bistability estimated from seizure-free, inter-ictal-activity-free resting-state SEEG recording could be informative about epileptic pathophysiology. In particular, we addressed whether bistability could delineate the epileptogenic zone (EZ) and dissociate EZ signals from signals in nEZ contacts that reflect more healthy forms of brain activity.

Representative time series (Fig 5A–B) showed that the EZ contacts did not show conspicuous epileptic inter-ictal events (IIE), and the sparse IIEs were removed from analysis where found (Supplementary Methods). Interestingly, elevated > 80 Hz bistability of the EZ contact was already a visually salient characteristic and stronger in EZ than in a nearby nEZ contact from the same region (DFA, bistability fitting see Supplementary Fig 3 F–J). We assessed bistability and LRTCs in narrow-bands frequencies at the group level for nEZ- versus EZ-electrode contacts (Fig 5C–D). Collapsing narrow-band DFA and BiS estimates into broader frequency bands revealed significant differences between nEZ- and EZ-electrode contacts in β - and γ -band BiS estimates with effect sizes of 0.5 and 0.65 (Cohen's *d*), respectively (Fig 5E). There was also a difference between nEZ- and EZ-electrode contacts in the δ -band DFA exponent with a Cohen's *d* of 0.2.

These group-level findings suggest that both bistability and LRTCs could constitute informative features for classifying nEZ- and EZ-electrode contacts. We thus conducted an EZ-vs-nEZ classification analysis using random forest algorithm³⁹ and with frequency-collapsed BiS and DFA estimates as neuronal features, with the electrode contact location in Yeo systems as an additional feature. The cross-validation for the classification was a 80:20-partition (training:test) with 500 iterations. This revealed a reliable outcome with the area under curve (AUC) for the receiver operating characteristic reaching $AUC = 0.8 \pm 0.002$ (mean $\pm$ std) (Supplementary Fig 11C). To identify the most informative components to the classifier, we assessed global and within-subject feature importance with the Shapley Additive exPlanations (SHAP) values⁴⁰. The SHAP values corroborated that γ - and β -band BiS estimates were indeed the most important features, followed by contact location (Yeo system), and δ -band DFA (Fig 5F).

Given these encouraging group-level and contact-classification results, we quantified the within-subject accuracy of neuronal bistability in localizing the epileptogenic area. We used leave-one-out validation so that the EZ-vs-nEZ contact classification was performed for each patient with the rest of the patient serving as training data. Additionally, to independently evaluate the contributions of BiS and DFA estimates to classification accuracy, we implemented the classification with four feature sets: DFA alone, BiS alone, combining DFA and BiS (D&B), and combining D&B and SEEG contact location in Yeo systems (D&B(Y)).

Fig 5. Bistability showed strong predictive power for pathophysiology. (A–B) Five minutes of broad-band traces and narrow-band power (R^2) time series of an EZ (A) and a nEZ (B) cortical location recorded with two distinct electrode shafts in one subject. Both contacts were 19.7 mm apart within supervisor frontal gyrus (sFG) and were referenced with the same nearest white matter contact (Arnulfo et al., 2015). (C) Average normalized narrow-band BiS and (D) DFA estimates for all EZ (pink) and nEZ contacts (green). Shaded areas indicate 25% and 75%-tile. (E) The effect size of differences between EZ and nEZ contacts in frequency-collapsed BiS (red) and DFA (black). Dashed line indicates 99%-tile observation from surrogate data ($N_{\text{surrogate}}=1000$). (F) Feature importance estimated using SHAP values. (G) The area under curve (AUC) of receiver operating characteristics (ROC) averaged across subjects (black) and the AUC of pooled within-subject classification results (blue) when using (i) DFA alone, (ii) BiS alone, (iii) combining DFA and BiS (D&B), and (iv) D&B plus contact loci in Yeo systems (D&B(Y)). Dashed lines indicate 99%-tile of AUC observed from 1,000 surrogates created independently for each of the four feature sets. (H–J) Post-hoc inspection of results derived using D&B(Y) feature set (the black marker in G). (H) Spearman's correlation ($p < 10^{-6}$, $n=55$) between individual AUC and within-subject mean Cohen's d between EZ and nEZ in band-collapsed DFA and BiS. (I) Receiver operating characteristics of classification within subjects (thin lines) and mean ROC (thick). (J) Within-patient prediction precision as a function of TPR indicated by the magenta box from (I); the red marker indicates the population mean. $\text{Precision} = \text{true positive} \div \text{reported positive}$.

Overall, the within-subject classification accuracy for EZ contacts was higher than chance level across all feature combinations (Fig 5G). Classification with all features yielded the best performance at an average AUC of 0.7 ± 0.14 (Black marker, Fig 5G). BiS alone yielded a greater AUC than DFA alone. Including the contact-brain system as an additional feature to D&B

increased the AUC by 0.06. The subject AUC values were correlated with the subject-specific differences in DFA and BiS estimates between EZ and nEZ ($r = -0.53$, $p < 10^{-6}$) (Fig 5H) and were not affected by either the total numbers of contacts, EZ contacts, nor the ratio of EZ and nEZ contacts (Pearson's correlation coefficient, $r = -0.06$, $p = 0.66$; $r = -0.07$, $p = 0.61$; $r = -0.09$, $p = 0.50$; respectively). Finally, the classifier yielded an average precision of 0.74 ± 0.30 (mean $\pm$ SD). While the true positive rate was 0.24 ± 0.17 , the false-positive rate was only 0.03 ± 0.03 (Fig 5G and H), which shows that most EZ contacts identified with the bistability-based classification were correct even though the classifier did not identify all true EZ contacts.

Discussion

We found here that bistable criticality is a pervasive characteristic of human brain activity and is both functionally significant in healthy brain dynamics and clinically informative as a putative pathophysiological mechanism in epilepsy. In a generative model of synchronization dynamics with positive feedback, bistability occurred exclusively within a critical-like regime. In both MEG and SEEG, we found significant bistability and LRTCs in spontaneous amplitude fluctuations of cortical oscillations widely across the neocortex and from δ - (2–4 Hz) to γ - (40–225 Hz) frequency bands. Moreover, as predicted by the model, bistability was positively correlated with LRTCs. As key evidence for functional significance, resting-state bistability was a trait-like predictor⁴¹ of healthy cognitive performance in MEG and clearly associated with epileptogenic pathophysiology in SEEG. These findings indicate that bistable criticality is an important novel facet of large-scale brain dynamics in both healthy and diseased human brain. Moreover, these observations strongly suggest that the brain criticality framework—currently focused on continuous phase transitions—should be expanded to include both continuous and discontinuous phase transitions (see Supplementary Theory).

In the model, we found bistability to occur exclusively within the critical regime so that the level of bistability increased monotonically with increasing positive feedback ρ . This state-dependent feedback thus acted as a continuous control parameter for shifting the model between a continuous and a discontinuous phase transition, at low or high levels of feedback, respectively. In both MEG and SEEG, band-collapsed θ – α (5.4–11 Hz) and γ (40–225 Hz) frequency cortical BiS and DFA estimates were correlated on group average level (see Fig 3 G–H) and among individuals within functional systems (see Fig 3I). These widespread correlations collectively suggest that a gradient from “low LRTCs and weakly bistable” to “high LRTCs and strongly bistable” is a systematic characteristic of the human brain dynamics.

A positive feedback loop is thought to be a generic mechanism^{31,35,42–44} leading to bistability in a wide range of modeled and real-world complex systems including the canonical sand-pile model³⁵ and its variations⁴², ecosystems^{29,45}, gene regulatory networks^{25,46,47}, intra-cellular signaling^{48,49}, and network models of spiking neurons^{31,50}. In our model, the positive local feedback was implemented as state-dependent phase noise^{25,34}. Three mechanisms have been proposed to account for feedback and state-dependency in microscopic neuronal dynamics⁵¹, whereas the exact neuronal mechanism for meso- and macroscopic state-dependency remains unclear. We postulate that the state-dependency be a slowly fluctuating physiological parameter conceivably reflecting the cortical excitability and corresponding resource demand^{31,50,52}.

Bistable dynamics, in general, could be associated a dichotomy of both beneficial and detrimental outcomes. Organisms can operate in bistable mode that is thought to reflect a dynamic motif favourable to adaptation and survival^{125,46,49,53}. However, a high degree of bistability characterizes many complex systems prone to catastrophic shifts^{43,44} such as sudden violent vibrations in aerodynamic systems⁵⁴, irrevocable environmental changes^{29,55,56}, wars and conflicts⁵⁷. In healthy MEG subjects, BiS and DFA were correlated, and higher θ – α band BiS and DFA estimates predicted better cognitive performance. In the SEEG from epileptic patients, excessive β - and γ -band BiS, but not DFA, characterized EZ (see Fig 5E–F), which suggests a functional gradient wherein moderate bistability reflect functional advantages but high degree of bistability be a sign of pathological hyper-excitability; this pathological bistability could be associated with hyper-excitability, excessive synchrony, high resource demands, and likely subsequent oxidative stress and tissue damage⁵⁸. This speculation is in accordance with biophysical models of seizures that suggest a crucial role of a discontinuous transition (a sub-critical Hopf bifurcation)^{25,33} in generalized seizures^{52,53}.

With invasive SEEG, we found consistent and accurate performance of the BiS estimates in EZ localization, which suggests a great potential for broader clinical utility, *e.g.*, using non-invasive MEG or EEG. Future work could exploit the presence of widespread bistability to large-scale biophysical models of neural dynamics, building on the analytic link between the simplified model employed here and physiologically derived neural mass and mean field models. Whereas the simple model yields dynamical insights, the large-scale biophysical models are crucial for understanding biological mechanisms³³, including those that describe seizure propagation in individual patient brain networks⁵⁹.

Materials and Methods

The canonical Hopf bifurcation

The canonical model of sub- or supercritical bifurcation is ²⁵:

$$\dot{r} = -r^5 + \lambda r^3 + \beta r + \eta[(1 - \rho)\zeta_a(t) + \rho r \zeta_m(t)], \quad (1)$$

where $\dot{r}$ is the time derivative of a local neuronal activity r (a real number); λ is the shape parameter and β the bifurcation parameter; η scales the overall influence of noise; where $\zeta_a(t)$ and $\zeta_m(t)$ are additive and state-dependent noise respectively, and they are two uncorrelated Wiener processes; the parameter ρ weights the influence of state-dependent noise. Different combinations of λ and β result in either super-critical or sub-critical bifurcation (details in ²⁵), which are associated with continuous or discontinuous (or second- or first-order) phase transition, respectively ^{23,35,60}. When r described the amplitude of a two-dimensional system with phase θ , then (eq. 1) describes a Normal form stochastic Hopf bifurcation.

The Kuramoto model

We studied first- and second-order phase transitions in a Kuramoto model with a modified noise term. The Kuramoto model is a generative model that can be used for studying the collective behaviours of a number of interconnected phase oscillators due to weak interactions ^{32,61}. In a

272 Kuramoto model, the dynamics of each oscillator i is a scalar phase time series θ_i ($\theta \in 0:2\pi$),
 273 coupled into a population ensemble θ as,

$$274 \quad \dot{\theta}_i = \omega_i + K_i + Z_i, \quad (2)$$

275 where, for any oscillator i , $\dot{\theta}_i$ is the rotation of its phase θ_i ; ω_i is the natural (uncoupled)
 276 frequency of i ; $K_i = K_i(\theta)$ the coupling between i and the rest oscillators of the ensemble, and
 277 Z_i is a stochastic term. The degree of synchrony of the ensemble (*i.e.*, order parameter or mean
 278 field) is the outcome of tripartite competition for controlling the collective behaviours of all
 279 oscillators: ω_i and Z_i are desynchronizing factors whereas K_i is the synchronizing factor. Here,
 280 ω_i follows a normal distribution with mean of zero (Hz), meaning without loss of generality, the
 281 system can be observed on a rotating phase plan with arbitrary angular velocity. In the classic
 282 model, the coupling term K_i is defined as the i -th oscillator adjusts its phase according to
 283 interactions with all other oscillators in the system through a pair-wise phase interaction
 284 function:

$$285 \quad K_i = \frac{\kappa}{N} \sum_{j=1}^N \sin(\theta_i - \theta_j), \quad (3)$$

286 here, κ is a scalar number representing coupling strength, $N=200$ is the number of oscillators in
 287 the ensemble. For simplicity, here we used a fully coupled network to avoid other families of
 288 emerging dynamics due to nodal or network structural disorders, *e.g.*, Griffiths phase^{62,63}. In
 289 addition, we found that with a Gaussian nodal-weight distribution, the model behaved
 290 identically to the fully coupled networks. We modified the noise term in line with the Hopf
 291 bifurcation (*eq. 1*) as:

$$292 \quad Z_i = \eta[(1 - \rho)\zeta_a(t) + \rho(R_{MAX} - R)\zeta_m(t)], \quad (4)$$

293 here, $\zeta_a(t)$ and $\zeta_m(t)$ are additive and multiplicative noise, respectively – as described in (*eq. 1*),
 294 however, in (*eq. 4*) $\zeta_a(t)$ and $\zeta_m(t)$ are uncorrelated and independent Gaussian phase noise
 295 with zero mean and unit variance; ρ scales the influence of $\zeta_m(t)$; note that the 2 bracketed
 296 terms are offset (*i.e.*, $(1 - \rho)$ and ρ) such that their combined effect stays approximately
 297 constant in magnitude; R_{MAX} is the maximal order the population can reach (*e.g.*, slightly below
 298 1 due to the presence of noise) and R is the current mean field (a scalar) that quantifies the
 299 degree of synchrony of the population at time t ,

$$300 \quad R(t) = \left| \frac{1}{N} \sum_{n=1}^N e^{i\theta_n(t)} \right| \quad (5)$$

301 when viewing R from the complex phase plan, it essentially is the centroid vector of the
 302 population phase distribution: if the whole population is in full synchrony, $R = R_{MAX} \rightarrow 1$; when
 303 there is no synchrony, $R \rightarrow 0$ (see insets, Fig 1H).

304 **MEG recording and subjects**

305 We recorded 10 minutes resting-state magnetoencephalographic (MEG) data from 18 subjects
 306 (11 males, 31.7 ± 10.5 , mean $\pm$ std, years of age) at the BioMag Laboratory, HUS Medical Imaging
 307 Center, Helsinki Finland. Subjects were seated in a dimly lit room and instructed to focus on a
 308 cross on the screen in front of them. Recordings were carried out at Meilahti hospital in Helsinki

(detailed in Supp. Material). All subjects were screened for neurological conditions. The study protocol for MEG and MRI data obtained at the University of Helsinki was approved by the Coordinating Ethical Committee of Helsinki University Central Hospital (HUCH) (ID 290/13/03/2013) and was performed according to the Declaration of Helsinki.

We also assessed working memory, attention, and executive functions in these subjects with a battery of neuropsychological tests. These included (see Fig 4 x-axis): Zoo Map Time, Toulouse-Pieron test (TP), Digit Symbol Coding test, Zoo Map Plan, Digit Span forward and backward (BackDigits and ForwDigits, respectively), Letter-Number Sequencing (LNS), Trail Making Test parts A and B (TMT-A, TMT-B). Some subjects had missing/invalid behavioural scores, and we studied the neuro-behavioural correlations with dataset that at least had 16 valid subjects' scores.

SEEG recording and subjects

We acquired 10 minutes of uninterrupted, seizure-free resting-state brain activity with eyes closed from 64 drug resistant focal epilepsy patients (28 females, 30.1 ± 9.1 , mean $\pm$ SD, years of age, see Supplementary Table 1) undergoing pre-surgical assessments (detailed in Supp. Material). The subjects gave written informed consent for participation in research studies. The study protocol for SEEG, computerized tomography (CT), and MRI data obtained in the La Niguarda Hospital were approved by the ethical committee of the Niguarda "Ca Granda" Hospital, Milan (ID 939), Italy, and was performed according to the Declaration of Helsinki.

Prior to surgery, medical doctors identified epileptogenic and seizure propagation zone by visual analysis of the SEEG traces^{64,65}. Epileptogenic areas (generators) are the region of interest that are necessary and sufficient for the origin and early organization of the epileptic activities⁶⁶. SEEG contacts recorded from such generators often show low voltage fast discharge or spike and wave events at seizure onset. Seizure propagation areas (receivers) are recruited during seizure propagation, but they do not initialize seizures^{59,67}. Contacts recorded from receivers show delayed, rhythmic modifications after seizure initiation in the generators. It is common to see regions demonstrating both generator and receiver dynamics, thus they were identified as generator-receiver. In this study, we refer to generator, receiver, and generator-receiver collectively as epileptogenic zone (EZ) to distinguish them from those that were tentatively identified as healthy non-EZ regions (nEZ).

339 References

- 340 1. Thom, R. Structural Stability and Morphogenesis. (CRC Press, 1972).
- 341 2. Zeeman, E. C. Catastrophe Theory. Sci. Am. (1976).
- 342 3. Sornette, D. Dragon-Kings, Black Swans and the Prediction of Crises. Int. J. Terrasp. Sci. Eng. **2**, 1–18
- 343 (2009).
- 344 4. Uhlhaas, P. J. et al. Neural synchrony in cortical networks: History, concept and current status. Front.
- 345 Integr. Neurosci. **3**, (2009).
- 346 5. Fries, P. A mechanism for cognitive dynamics: Neuronal communication through neuronal coherence.
- 347 Trends Cogn. Sci. **9**, 474–480 (2005).
- 348 6. Chialvo, D. R. Emergent complex neural dynamics. Nat. Phys. **6**, 744–750 (2010).
- 349 7. Palva, S. & Palva, J. M. Roles of Brain Criticality and Multiscale Oscillations in Temporal Predictions
- 350 for Sensorimotor Processing. Trends Neurosci. **41**, 729–743 (2018).
- 351 8. Beggs, J. M. & Timme, N. Being critical of criticality in the brain. Front. Physiol. (2012).
- 352 9. Muñoz, M. A. Colloquium: Criticality and dynamical scaling in living systems. Rev. Mod. Phys. **90**,
- 353 (2018).
- 354 10. Wiltling, J. & Priesemann, V. 25 years of criticality in neuroscience — established results, open
- 355 controversies, novel concepts. Curr. Opin. Neurobiol. **58**, 105–111 (2019).
- 356 11. Kinouchi, O. & Copelli, M. Optimal dynamical range of excitable networks at criticality. Nat. Phys. **2**,
- 357 348–352 (2006).
- 358 12. Shew, W. L., Yang, H., Yu, S., Roy, R. & Plenz, D. Information capacity and transmission are
- 359 maximized in balanced cortical networks with neuronal avalanches. J. Neurosci. **31**, 55–63 (2011).
- 360 13. Boedecker, J., Obst, O., Lizier, J. T., Mayer, N. M. & Asada, M. Information processing in echo state
- 361 networks at the edge of chaos. Theory Biosci. **131**, 205–213 (2012).
- 362 14. Bertschinger, N. & Natschläger, T. Real-Time Computation at the Edge of Chaos in Recurrent Neural
- 363 Networks. Neural Comput. **16**, 1413–1436 (2004).
- 364 15. Shriki, O. & Yellin, D. Optimal Information Representation and Criticality in an Adaptive Sensory
- 365 Recurrent Neuronal Network. PLoS Comput. Biol. **12**, (2016).
- 366 16. Zimmern, V. Why Brain Criticality Is Clinically Relevant: A Scoping Review. Front. Neural Circuits
- 367 **14**, (2020).
- 368 17. Liu, X., Ward, B. D., Binder, J. R., Li, S. J. & Hudetz, A. G. Scale-free functional connectivity of the
- 369 brain is maintained in anesthetized healthy participants but not in patients with unresponsive
- 370 wakefulness syndrome. PLoS One **9**, (2014).
- 371 18. Cerf, R., El Ouasdad, E. H. & Kahane, P. Criticality and synchrony of fluctuations in rhythmical brain
- 372 activity: Pretransitional effects in epileptic patients. Biol. Cybern. **90**, 239–255 (2004).
- 373 19. Wilson, H. R. & Cowan, J. D. Excitatory and Inhibitory Interactions in Localized Populations of Model
- 374 Neurons. Biophys. J. **12**, 1–24 (1972).
- 375 20. Wilson, C. Up and down states. Scholarpedia **3**, 1410 (2008).
- 376 21. Steriade, M., Nunez, A. & Amzica, F. A novel slow (< 1 Hz) oscillation of neocortical neurons in
- 377 vivo: Depolarizing and hyperpolarizing components. Journal of Neuroscience vol. 13 (1993).
- 378 22. Holcman, D. & Tsodyks, M. The emergence of Up and Down states in cortical networks. PLoS
- 379 Comput. Biol. **preprint**, e23 (2005).
- 380 23. Cocchi, L., Gollo, L. L., Zalesky, A. & Breakspear, M. Criticality in the brain: A synthesis of
- 381 neurobiology, models and cognition. Progress in Neurobiology (2017).
- 382 24. Roberts, J. A., Boonstra, T. W. & Breakspear, M. The heavy tail of the human brain. Current Opinion in
- 383 Neurobiology (2015).
- 384 25. Freyer, F., Roberts, J. A., Ritter, P. & Breakspear, M. A Canonical Model of Multistability and Scale-
- 385 Invariance in Biological Systems. PLoS Comput. Biol. (2012).
- 386 26. Freyer, F., Aquino, K., Robinson, P. A., Ritter, P. & Breakspear, M. Bistability and Non-Gaussian
- 387 Fluctuations in Spontaneous Cortical Activity. J. Neurosci. (2009).
- 388 27. Hansen, E. C. A., Battaglia, D., Spiegler, A., Deco, G. & Jirsa, V. K. Functional connectivity dynamics:
- 389 Modeling the switching behavior of the resting state. Neuroimage **105**, 525–535 (2015).
- 390 28. Zalesky, A., Fornito, A., Cocchi, L., Gollo, L. L. & Breakspear, M. Time-resolved resting-state brain

networks. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 10341–10346 (2014).

29. Villa Martín, P., Bonachela, J. A., Levin, S. A. & Muñoz, M. A. Eluding catastrophic shifts. *Proc. Natl. Acad. Sci.* **112**, E1828–E1836 (2015).

30. Agu, M. & Teramachi, Y. Prediction of catastrophes in bistable systems using externally applied random force. *J. Appl. Phys.* **49**, 3645–3648 (1978).

31. di Santo, S., Villegas, P., Burioni, R. & Muñoz, M. A. Landau–Ginzburg theory of cortex dynamics: Scale-free avalanches emerge at the edge of synchronization. *Proc. Natl. Acad. Sci.* **115**, E1356–E1365 (2018).

32. Breakspear, M., Heitmann, S. & Daffertshofer, A. Generative Models of Cortical Oscillations: Neurobiological Implications of the Kuramoto Model. *Front. Hum. Neurosci.* (2010).

33. Breakspear, M. Dynamic models of large-scale brain activity. *Nature Neuroscience* (2017).

34. Izhikevich, E. M. *Dynamical systems in neuroscience: the geometry of excitability and bursting*. (MIT Press, 2007).

35. Di Santo, S., Burioni, R., Vezzani, A. & Muñoz, M. A. Self-Organized Bistability Associated with First-Order Phase Transitions. *Phys. Rev. Lett.* (2016).

36. Linkenkaer-Hansen, K., Nikouline, V. V., Matias Palva, J. & Ilmoniemi, R. J. Long-Range Temporal Correlations and Scaling Behavior in Human Brain Oscillations. *J. Neurosci.* (2001).

37. Simola, J., Zhigalov, A., Morales-Muñoz, I., Palva, J. M. & Palva, S. Critical dynamics of endogenous fluctuations predict cognitive flexibility in the Go/NoGo task. *Sci. Rep.* **7**, (2017).

38. Pal, M., Pal, A. K., Ghosh, S. & Bose, I. Early signatures of regime shifts in gene expression dynamics. *Phys. Biol.* **10**, 036010 (2013).

39. Breiman, L. Random Forests. in *Machine Learning* vol. 45 5–32 (Kluwer Academic Publishers, 2001).

40. Lundberg, S. M., Allen, P. G. & Lee, S.-I. A Unified Approach to Interpreting Model Predictions. in *Conference on Neural Information Processing Systems (NIPS) 31st* (2017).

41. Palva, J. M. et al. Neuronal long-range temporal correlations and avalanche dynamics are correlated with behavioral scaling laws. *Proc. Natl. Acad. Sci.* (2013).

42. Buendía, V., di Santo, S., Bonachela, J. A. & Muñoz, M. A. Feedback Mechanisms for Self-Organization to the Edge of a Phase Transition. *Frontiers in Physics* vol. 8 (2020).

43. Hugo, H. L. D., Oriá, M., Sornette, D., Ott, E. & Gauthier, D. J. Predictability and suppression of extreme events in a chaotic system. *Phys. Rev. Lett.* **111**, 1–5 (2013).

44. Sornette, D. & Ouillon, G. Dragon-kings Mechanisms, statistical methods and empirical evidence. *Eur. Phys. J. Spec. Top.* **205**, 1–26 (2012).

45. Kéfi, S., Rietkerk, M., van Baalen, M. & Loreau, M. Local facilitation, bistability and transitions in arid ecosystems. *Theor. Popul. Biol.* **71**, 367–379 (2007).

46. Dubnau, D. & Losick, R. Bistability in bacteria. *Mol. Microbiol.* **61**, 564–572 (2006).

47. Kuwahara, H. & Soyer, O. S. Bistability in feedback circuits as a byproduct of evolution of evolvability. *Mol. Syst. Biol.* **8**, (2012).

48. Mitrophanov, A. Y. & Groisman, E. A. Positive feedback in cellular control systems. (2008).

49. Bednarz, M., Halliday, J. A., Herman, C. & Golding, I. Revisiting bistability in the lysis/lysogeny circuit of bacteriophage lambda. *PLoS One* **9**, (2014).

50. Cowan, J. D., Neuman, J. & van Drongelen, W. Wilson–Cowan Equations for Neocortical Dynamics. *J. Math. Neurosci.* **6**, 1–24 (2016).

51. Doiron, B., Litwin-Kumar, A., Rosenbaum, R., Ocker, G. K. & JosiC, K. The mechanics of state-dependent neural correlations. *Nature Neuroscience* vol. 19 383–393 (2016).

52. Jirsa, V. K., Stacey, W. C., Quilichini, P. P., Ivanov, A. I. & Bernard, C. On the nature of seizure dynamics. *Brain* (2014).

53. Breakspear, M. et al. A unifying explanation of primary generalized seizures through nonlinear brain modeling and bifurcation analysis. *Cereb. Cortex* (2006).

54. Qi, X. & Zhongke, S. Bifurcation analysis and stability design for aircraft longitudinal motion with high angle of attack. *Chinese J. Aeronaut.* **28**, 250–259 (2015).

55. Boerlijst, M. C., Oudman, T. & de Roos, A. M. Catastrophic Collapse Can Occur without Early Warning: Examples of Silent Catastrophes in Structured Ecological Models. *PLoS One* **8**, (2013).

56. Barnosky, A. D. et al. Approaching a state shift in Earth’s biosphere. *Nature* **486**, 52–58 (2012).

57. Díaz, F. A. Inequality, Social Protests and Civil War. *OASIS* 25–39 (2017).

- 445 58. Salim, S. Oxidative stress and the central nervous system. *Journal of Pharmacology and Experimental*
446 *Therapeutics* vol. 360 201–205 (2017).
- 447 59. Proix, T., Bartolomei, F., Guye, M. & Jirsa, V. K. Individual brain structure and modelling predict
448 seizure propagation. *Brain* (2017).
- 449 60. Kim, S., Park, S. H. & Ryu, C. S. Noise-enhanced multistability in coupled oscillator systems. *Phys.*
450 *Rev. Lett.* **78**, 1616–1619 (1997).
- 451 61. Rodrigues, F. A., Peron, T. K. D. M., Ji, P. & Kurths, J. The Kuramoto model in complex networks.
452 *Physics Reports* (2016).
- 453 62. Moretti, P. & Muñoz, M. A. Griffiths phases and the stretching of criticality in brain networks. *Nat.*
454 *Commun.* (2013).
- 455 63. Muñoz, M. A., Juhász, R., Castellano, C. & Ódor, G. Griffiths phases on complex networks. *Phys. Rev.*
456 *Lett.* (2010).
- 457 64. Cardinale, F. et al. Stereoelectroencephalography: Surgical methodology, safety, and stereotactic
458 application accuracy in 500 procedures. *Neurosurgery* (2013).
- 459 65. Cossu, M. et al. Stereoelectroencephalography in the presurgical evaluation of focal epilepsy: a
460 retrospective analysis of 215 procedures. *Neurosurgery* **706–18**, 706–18 (2005).
- 461 66. Luders, H., Engel, J. J. & Munari, C. General principles. in *Surgical Treatment of the Epilepsies*. 2nd
462 ed. 137–153 (Raven Press, 1993).
- 463 67. Bartolomei, F., Guye, M. & Wendling, F. Abnormal binding and disruption in large scale networks
464 involved in human partial seizures. *EPJ Nonlinear Biomed. Phys.* (2013).
- 465 68. Taulu, S. & Simola, J. Spatiotemporal signal space separation method for rejecting nearby interference
466 in MEG measurements. *Phys. Med. Biol.* **51**, 1759–1768 (2006).
- 467 69. Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J. M. FieldTrip: Open source software for advanced
468 analysis of MEG, EEG, and invasive electrophysiological data. *Comput. Intell. Neurosci.* **2011**, (2011).
- 469 70. Schaefer, A. et al. Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional
470 Connectivity MRI. *Cereb. Cortex* **28**, 3095–3114 (2017).
- 471 71. Yeo, B. T. et al. The organization of the human cerebral cortex estimated by intrinsic functional
472 connectivity. *J. Neurophysiol.* **106**, 1125–1165 (2011).
- 473 72. Hämmäläinen, M. S. R. & Ilmoniemi, R. J. Interpreting magnetic fields of the brain: minimum norm
474 estimates. *Med. Biol. Eng. Comput.* **32**, 35–42 (1994).
- 475 73. Hamalainen, M. S. & Sarvas, J. Realistic Conductivity Geometry Model of the Human Head for
476 Interpretation of Neuromagnetic Data. *IEEE TRANSACTIONS ON BIOMEDICAL ENGINEERING*
477 vol. 36 (1989).
- 478 74. Korhonen, O., Palva, S. & Palva, J. M. Sparse weightings for collapsing inverse solutions to cortical
479 parcellations optimize M/EEG source reconstruction accuracy. *J. Neurosci. Methods* **226**, 147–160
480 (2014).
- 481 75. Fischl, B. et al. Neurotechnique Whole Brain Segmentation: Automated Labeling of Neuroanatomical
482 Structures in the Human Brain. *Neuron* **33**, 341–355 (2002).
- 483 76. Arnulfo, G., Hirvonen, J., Nobili, L., Palva, S. & Palva, J. M. Phase and amplitude correlations in
484 resting-state activity in human stereotactical EEG recordings. *Neuroimage* 114–127 (2015).
- 485 77. Arnulfo, G., Narizzano, M., Cardinale, F., Fato, M. M. & Palva, J. M. Automatic segmentation of deep
486 intracerebral electrodes in computed tomography scans. *BMC Bioinformatics* 16: 99 (2015).
- 487 78. Monto, S., Vanhatalo, S., Holmes, M. D. & Palva, J. M. Epileptogenic neocortical networks are
488 revealed by abnormal temporal dynamics in seizure-free subdural EEG. *Cereb. Cortex* **17**, 1386–1393
489 (2007).
- 490 79. Eke, A. et al. Physiological time series: Distinguishing fractal noises from motions. *Pflugers Arch. Eur.*
491 *J. Physiol.* (2000).
- 492 80. Poil, S.-S., Hardstone, R., Mansvelder, H. D. & Linkenkaer-Hansen, K. Critical-State Dynamics of
493 Avalanches and Oscillations Jointly Emerge from Balanced Excitation/Inhibition in Neuronal
494 Networks. *J. Neurosci.* (2012).
- 495 81. Ihlen, E. A. F. Introduction to multifractal detrended fluctuation analysis in Matlab. *Front. Physiol.*
496 (2012).
- 497 82. Hardstone, R. et al. Detrended fluctuation analysis: A scale-free view on neuronal oscillations. *Front.*
498 *Physiol.* (2012).

83. Wit, E., van den Heuvel, E. & Romeijn, J. W. 'All models are wrong.': An introduction to model uncertainty. *Stat. Neerl.* (2012).

84. Lancaster, G., Iatsenko, D., Pidde, A., Ticcinelli, V. & Stefanovska, A. Surrogate data for hypothesis testing of physical systems. *Physics Reports* vol. 748 1–60 (2018).

85. Sokal, R. & Michener, C. A statistical method for evaluating systematic relationships. *Univ. Kansas Sci. Bull.* **38**, 1409–1438 (1958).

86. Hastie, T., Tibshirani, R. & Friedman, J. *The Elements of Statistical Learning Data Mining, Inference, and Prediction.* (Springer, 2009).

87. Wilting, J. & Priesemann, V. Between Perfectly Critical and Fully Irregular: A Reverberating Model Captures and Predicts Cortical Spike Propagation. *Cereb. Cortex* **29**, 2759–2770 (2019).

88. Bonachela, J. A. & Muñoz, M. A. Self-organization without conservation: True or just apparent scale-invariance? *J. Stat. Mech. Theory Exp.* **2009**, (2009).

89. Bonachela, J. A., De Franciscis, S., Torres, J. J. & Muñoz, M. A. Self-organization without conservation: Are neuronal avalanches generically critical? *J. Stat. Mech. Theory Exp.* **2010**, (2010).

90. Cowan, J. D., Neuman, J. & van Drongelen, W. Self-Organized Criticality and Near-Criticality in Neural Networks. in *Criticality in Neural Systems* 465 (Wiley-VCH, 2014).

91. Bak, P., Tang, C. & Wiesenfeld, K. Self-Organized Criticality: An Explanation of 1/f Noise. *Phys. Rev. Lett.* **59**, 381 (1987).

92. Mandelbrot, B. Self-affinity and fractal dimension. in *Physica Scripta* 257–260 (1985).

93. Mandelbrot, B. B. & Van Ness, J. W. Fractional Brownian Motions, Fractional Noises and Applications. *SIAM Rev.* (1968).

94. Kinouchi, O., Brochini, L., Costa, A. A., Campos, J. G. F. & Copelli, M. Stochastic oscillations and dragon king avalanches in self-organized quasi-critical systems. *Sci. Rep.* **9**, (2019).

95. Grinstein, G. Generic scale invariance and self-organized criticality. in *Scale Invariance, Interfaces, and Non-Equilibrium Dynamics* 261–293 (1995).

96. Drossel, B., Clar, S. & Schwabl, P. Exact Results for the One-Dimensional Self-Organized Critical Forest-Fire Model. vol. 71.

97. Levina, A., Herrmann, J. M. & Geisel, T. Dynamical synapses causing self-organized criticality in neural networks. *Nat. Phys.* **3**, 857–860 (2007).

98. Herculano-Houzel, S. Scaling of brain metabolism with a fixed energy budget per neuron: Implications for neuronal activity, plasticity and evolution. *PLoS One* **6**, (2011).

99. Kuzawa, C. W. et al. Metabolic costs and evolutionary implications of human brain development. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 13010–13015 (2014).

100. Roh, E. & Kim, M. S. Brain regulation of energy metabolism. *Endocrinology and Metabolism* vol. 31 519–524 (2016).

101. Ikeda, K. & Bekkers, J. M. Counting the number of releasable synaptic vesicles in a presynaptic terminal. (2009).

102. Markram, H. & Tsodyks, M. Redistribution of synaptic efficacy between pyramidal neurons. *Nature* 807–10 (1996).

103. Tsodyks, M. V. & Markram, H. The neural code between neocortical pyramidal neurons depends on neurotransmitter release probability. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 719–723 (1997).

104. Robinson, P. A., Rennie, C. J. & Rowe, D. L. Dynamics of large-scale brain activity in normal arousal states and epileptic seizures. *Phys. Rev. E - Stat. Physics, Plasmas, Fluids, Relat. Interdiscip. Top.* **65**, 9 (2002).

105. Protachevicz, P. R. et al. Bistable firing pattern in a neural network model. *Front. Comput. Neurosci.* **13**, (2019).

106. Lundstrom, B. N., Boly, M., Duckrow, R., Zaveri, H. P. & Blumenfeld, H. Slowing less than 1 Hz is decreased near the seizure onset zone. *Sci. Rep.* **9**, (2019).

107. Englot, D. J. et al. Impaired consciousness in temporal lobe seizures: Role of cortical slow activity. *Brain* **133**, 3764–3777 (2010).

108. Arnulfo, G. et al. Long-range phase synchronization of high-frequency oscillations in human cortex. *Nat. Commun.* **11**, (2020).

109. Freyer, F. et al. Biophysical Mechanisms of Multistability in Resting-State Cortical Rhythms. *J. Neurosci.* (2011).

Acknowledgements

We would like to thank our Italian collaborators from the Centre of Epilepsy Surgery “C. Munari” at Niguarda Hospital, in particular Annalisa Rubino and Dr. Francesco Cardinale, for their support. We thank Drs. James Roberts for sharing code, and Serena di Santo, Stewart Heitmann, and Alexander Zhigalov for insightful discussions.

Funding:

The Academy of Finland SA 266402, 303933, and SA 325404 (SP)

The Academy of Finland SA 253130 and 296304 (JMP)

Author contributions:

Conceptualization: SHW, JMP

Funding acquisition: JMP, SP

Supervision: JMP

Methodology: SHW, MB, JMP

Software: SHW, MB, VM

Formal analysis: SHW

Resources: LN, GA

Data Curation: GA, FS

Visualization: SHW

Writing - Original Draft: SHW, JMP, MB, and SP

Competing Interest Statement:

All authors declare no competing interests. The funders had no role in study design, data acquisition, analyses, decision to publish, and preparation of the manuscript.

Data and materials availability:

Raw data and patient details cannot be shared due to Italian governing laws and Ethical Committee restrictions. Intermediate data, final processed data, and all code that support the findings of this study are available from the corresponding authors upon reasonable request.

Supplementary Materials

Bistable critical dynamics in the resting-state human brain

Sheng H. Wang, Gabriele Arnulfo, Vladislav Myrov, Felix Siebenhühner, Lino Nobili, Michael Breakspear, Satu Palva, J. Matias Palva

sheng.wang@helsinki.fi

This PDF file includes:

Supplementary Methods
Supplementary materials
Figs. S1 to S11
Tables S1

Supplementary methods

MEG data acquisition

A 306-channel MEG system (204 planar gradiometers and 102 magnetometers) with a Vectorview-Triux (Elekta-Neuromag) was used to record 10 minutes eyes-open resting-state brain activity from 18 healthy adult subjects at the BioMag Laboratory (HUS Medical Imaging Center, Helsinki Finland). For cortical surface reconstruction, T1-weighted anatomical MRI scans were obtained at a resolution of 1x1x1 mm with a 1.5 T MRI scanner (Siemens, Germany). This study was approved by the ethical committee of Helsinki University Central hospital and was performed according to the Declaration of Helsinki. Written informed consent was obtained from each subject prior to the experiment.

MEG data preprocessing and filtering

The Maxfilter software with temporal signal space separation (tSSS) (Elekta Neuromag Ltd., Finland) was used to suppress extra-cranial noise in sensors and to interpolate bad channels⁶⁸. Independent component analysis (Matlab Fieldtrip toolbox, <http://fieldtrip.fcdonders.nl>) was next used to identify and remove components that were correlated with ocular (identified using the EOG signal), heart-beat (identified using the magnetometer signal as a reference) or muscle artefacts⁶⁹. The FreeSurfer software (<http://surfer.nmr.mgh.harvard.edu/>) was used for subject MEG sources reconstruction, volumetric segmentation of MRI data, surface reconstruction, flattening, cortical parcellation, and neuroanatomical labeling with the Schaefer-2017 atlas⁷⁰. Each of the Schaefer-parcel belongs to a functional system⁷¹ which informed later systems-level analysis.

The MNE software package was used to create head conductivity models and cortically constrained source models with 5000-7500 sources per subject and for the MEG-MRI co-registration and for the preparation of the forward and inverse operators^{72,73}. For each MEG

subject, a cortical parcellation of 400 Schaefer-parcels was obtained using reconstruction-accuracy optimized source-vertex-to-parcel collapsing method⁷⁴. The broadband time series of these parcels were then filtered into narrow-band time series using a bank of 20 Morlet filters with $m = 5$ and log-linearly spaced center frequencies ranging from 2 to 225 Hz. For group-level analyses, subject DFA and BiS estimates were morphed from 400 Schaefer-parcels into 100 Schaefer-parcels.

SEEG data acquisition

Resting-state brain activity from 64 drug resistant focal epilepsy patients (28 females, 30.1 ± 9.1 , mean $\pm$ SD, years of age, see S.Table 1) was acquired as monopolar local field potentials (LFPs) from brain tissue with platinum-iridium, multi-lead electrodes using a 192-channel SEEG amplifier system (NIHON-KOHDEN NEUROFAX-110) at 1 kHz sampling rate. Each penetrating electrode shaft has 8 to 15 contacts, and the contacts were 2 mm long, 0.8 mm thick and had an inter-contact border-to-border distance of 1.5 mm (DIXI medical, Besancon, France). The anatomical positions and amounts of electrodes varied according to surgical requirements⁶⁴. On average, each subject had 17 ± 3 (mean $\pm$ standard deviation) shafts (range 9-23) with a total of 153 ± 20 electrode contacts (range 122-184, left hemisphere: 66 ± 54 , right hemisphere: 47 ± 55 contacts, grey-matter contacts: 110 ± 25). The subjects gave written informed consent for participation in research studies and for publication of their data. This study was approved by the ethical committee (ID 939) of the Niguarda “Ca’ Granda” Hospital, Milan, and was performed according to the Declaration of Helsinki.

SEEG preprocessing and filtering

Cortical parcels were extracted from pre-surgically scanned T1 MRI 3D-FFE (used for surgical planning) using the Freesurfer package⁷⁵. A novel nearest-white-matter referencing scheme (its merits discussed in⁷⁶ was employed for referencing the monopolar SEEG LFP signals. An automated SEEG-electrode localization method was next used to assign each SEEG contact to a cortical parcel of Schaefer 100-parcel atlas with sub-millimeter accuracy⁷⁷. The SEEG electrodes were implanted to probe the suspected epileptogenic zones (EZ) while inevitably passing through healthy cortical structures. Contacts located at EZ are known to pick up frequent inter-ictal spikes and generate abnormally large DFA⁷⁸. Therefore, EZ contacts and contacts recorded from subcortical regions such as thalamus, hippocampus and basal ganglia were excluded from analysis.

Nevertheless, inter-ictal events (IIE) such as spikes can be occasionally observed at non-EZ locations in some subjects during rests. These IIE are characterized by high-amplitude fast temporal dynamics as well as by widespread spatial diffusion, which need to be excluded to avoid bias to DFA and BiS estimates. We followed approach used in to identify such IIEs. Briefly, each SEEG contact broad-band signal was partitioned into non-overlapping windows of 500 ms in length; a window was tagged as “spiky” and discarded from LRTCs and bistability analyses when at least 3 consecutive samples exceeding 7 times the standard deviation above the channel mean amplitude. Last, narrow-band frequency amplitude time series was obtained by convoluting the broad-band SEEG contact time series with Morlet wavelets that were identical to that of MEG data.

Estimating LRTCs using detrend fluctuation analysis

LRTCs in 1D time series can be assessed with several metrics⁷⁹, and in this study, linear detrend fluctuation analysis (DFA) was used to assess specifically how fast the overall root mean square of local fluctuations grows with increasing sampling period^{36,80}. An estimated DFA exponent reflects the finite-size power-law scaling in narrow-band amplitude fluctuations based on the assumption that the gradual evolution of a mono-fractal process time series would result in a normal distribution where the fluctuation can be captured by the second order statistical moments such as variance⁸¹. The computation of DFA briefly as follows (for rationales and technical details of the algorithm see⁸²):

- 1) The signal profile ($\mathbf{X}$) of a signal was computed by computing the cumulative sum of a demeaned narrow-band amplitude of a MEG parcel or SEEG contact time series.
- 2) A vector of window widths ($\mathbf{T}$) was defined in which the widths were linearly spaced on $\log_{10}$ scale between 10 and 90 seconds. The same scaling range was used across frequencies and for both MEG and SEEG, *i.e.*, identical vector of $\mathbf{T}$. The lower boundary of 10 s was set to safely avoid high non-stationarity and the filter artifacts, *i.e.*, 20 cycles of the slowest rhythm of 2 Hz; the upper bound of 90 s was 15% of total sample of the resting-state recording.
- 3) For each window width $t \in \mathbf{T}$, $\mathbf{X}$ was partitioned into an array of temporal windows in which each window was of length t and with 25% overlap between windows $W(t)$.
 - a. For each window $w \in \mathbf{W}(t)$, detrended signal $\mathbf{w}_{detrend}$ was obtained by removing the linear trend, *i.e.*, subtracting the least-squares fit of samples of w from the samples of w , and then obtained the root mean square of $\mathbf{w}_{detrend}$ ($\text{RMS}(\mathbf{w}_{detrend})$).
 - b. Finally, $F(t)$, the detrended fluctuation of window size t , was obtained by computing the mean of $\text{RMS}(\mathbf{w}_{detrend})$.
- 4) By repeating step (3) for all window lengths of $\mathbf{T}$ defined in step (2), $\mathbf{F}$, a vector of $F(t)$, $t \in \mathbf{T}$, was obtained. The DFA exponent is the slope of the trend line of $\mathbf{F}$ as a function of $\mathbf{T}$ on log-log scale (Supplementary Fig 3 G, J).

Estimating bistability index (BiS)

The BiS index of a power time series (R^2) derives from model comparison between a bimodal or unimodal fit of its probability distribution function (pdf); a large BiS means that the observed pdf is better described as bimodal, and when $\text{BiS} \rightarrow 0$ the pdf is better described as unimodal. We followed the approach used in^{24,25} to compute BiS. First, to find the pdf of power time series R^2 , the empirical R^2 was partitioned into 200 equal-distance bins and the number of observations in each bin was tallied. Next, maximum likelihood estimate (MLE) was used to fit a single-exponential function (*i.e.*, the square of a Gaussian process follows an exponential pdf):

$$P_x(x) = \gamma e^{-\gamma x} \quad (1)$$

and a bi-exponential function:

$$P_x(x) = \delta \gamma_1 e^{-\gamma_1 x} - (1 - \delta) \gamma_2 e^{-\gamma_2 x} \quad (2)$$

where γ_1 and γ_2 are two exponents and δ is the weighting factor.

Next, the Bayesian information criterion (BIC) was computed for the single- and bi-exponential fitting:

$$BIC = \ln(n)k - 2\ln(\hat{L}) \quad (3)$$

where n is the number of samples; $\hat{L}$ is the likelihood function; k is the number of parameters: $k = 1$ for single-exponential BIC_{Exp} and $k = 3$ for bi-exponential model BIC_{biE} . Thus BIC imposes a penalty to model complexity of the bi-exponential model⁸³ because it has two more degrees of freedom (second exponents and the weight δ) than the single exponential model.

Last, the BiS estimate is computed as the $\log_{10}$ transform of difference between the two BIC estimates as

$$dBIC = BIC_{Exp} - BIC_{biE} \quad (4)$$

$$BiS = \log_{10}(dBIC), \text{ if } dBIC > 0,$$

$$BiS = 0, \text{ if } dBIC \leq 0$$

Thus, a better model yields a small BIC value, and BiS will be large if the bi-exponential model is a more likely model for the observed power time series (Supplementary Fig 3 F, I).

Constructing surrogate data

To determine the chance level observations of DFA and BiS in null hypothesis data, *i.e.*, without embedded non-linear critical-like structures but with the same power spectrum as the real physiological signals⁸⁴, phase randomized Fourier transform surrogates of the broad-band time series was constructed for each MEG parcel and SEEG contact, $N_{MEG} = 6,800$ and $N_{SEEG} = 4,142$ (Supplementary Fig. 4). The surrogate broad-band data were filtered into narrow-band data and their DFA and BiS estimates were subsequently computed. Thus, the real observation can be compared against the significance thresholds that were derived from the probability distribution of the surrogate data across frequencies (Fig. 2).

Morphing MEG and SEEG data into a standard atlas

The MEG and SEEG group-level analyses were conducted in a 100-parcel standardized Schaefer atlas⁷⁰. The group-level MEG data were obtained in two steps (top, Supplementary Fig 4). First, narrow-band DFA and BiS estimates were computed within subjects using a finer parcellation of 400-parcel, and the resulting estimates were morphed into 100-parcel within subjects by averaging children parcels; group level cortical maps were next obtained by collapsing subjects' 100-parcel metrics.

Due to the variability in electrode location and other constraints in SEEG subjects, the morphing was done and verified differently. First, narrow-band DFA and BiS estimates of individual SEEG contacts were morphed directly into the Schaefer 100-parcel atlas (bottom, Supplementary Fig 4). The resulting group-level parcel estimates were thus heterogeneous in terms of sampling. For example, one parcel may contain observations from a varying number of electrodes and/or subjects. Hence, the group-level estimate of each parcel was the median of all observations, and the estimate for each of the seven Yeo sub-systems was the median of its constituent parcels. Furthermore, only the parcels ($n=90$) sampled by at least 3 subjects and 10 SEEG contacts were kept for group-level analysis. The group mean parcel metrics (Supplementary Fig. 5B) were identical to that of raw data (overlay curves, Fig. 2D–E) thus confirming that heterogeneous

SEEG sampling (Supplementary Fig. 2 and Supplementary Fig.4) did not bias the parcel-level metrics.

Clustering narrow-band frequency data

Frequency clustering analyses were conducted to reduce redundancy and for better interpretability of the narrow-band data (20 frequencies $\times$ DFA/BiS $\times$ 100 parcels). Spearman's rank correlation coefficients were computed between the group-level DFA/BiS estimates of 100-parcel for all pairs of frequencies. This resulted in a set of 20×20 adjacency matrices ($A_{f1,f2}$) representing the cortical topological similarity between frequencies. Next, the similarity matrix $A_{f1,f2}$ was partitioned using the unweighted pair group method with arithmetic mean algorithm⁸⁵, an agglomerative hierarchical clustering method, to obtain frequency clusters. The algorithm first builds a hierarchical tree through an iterative procedure to represent the distance between pairs of objects in $A_{f1,f2}$ (Supplementary Fig. 5C). In each iteration, two objects p and q with nearest distance $d(p, q)$ were merged into a cluster, and p and q can be either an element from $A_{f1,f2}$ or a cluster of elements from $A_{f1,f2}$; The distance function is defined as:

$$d(p, q) = \frac{1}{n_p n_q} \sum_{x \in p} \sum_{y \in q} d(x, y), \quad (5)$$

where n_p and n_q are the number of elements in p and q respectively, $d(x, y)$ is the distance between x , y , and x , y are matrix elements from $A_{f1,f2}$. The hierarchical tree was then used to partition the elements from $A_{f1,f2}$ into separated clusters, *e.g.*, if the height of p is close to the height of q , then their constituent elements are similar and therefore could be considered as a cluster (dashed boxes in Supplementary Fig. 5C, and solid boxes in Supplementary Fig. 5D).

Classifying pathophysiological SEEG contacts

The group-level frequency clustering analysis revealed that much of the narrow-band data were topologically correlated (Fig. 3). Hence, for the classification task, twenty narrow-band metrics were also collapsed into four frequency clusters as δ , θ - α , β , and γ band (Fig. 5A). As subjects varied greatly in their DFA and BiS estimates, band-collapsed data was normalized within subjects as $[X - \text{median}(X)] / \max(X - \text{median}(X))$, and thereby the differences between EZ and nEZ within subjects remained. The effect size of differences between band-collapsed and normalized DFA and BiS estimates were assessed with Cohen's d and compared with the 99%-tile of Cohen's d observed from 1,000 EZ-nEZ label-shuffled surrogate data (Fig. 5C).

The feature importance of these neuronal estimates were assessed with the SHapley Additive exPlanations (SHAP) values⁴⁰. In addition to the neuronal scores, the contact location in Yeo systems was also included as an additional feature (Fig. 4D). The SHAP values is a generic metric to explain any tree-based model by explicating the local and global interpretability of features, which advances the transparency that conventional classifications approaches lack of. For solving the EZ-classification problem, the non-parametric random-forest method⁸⁶ was employed. The random-forest algorithm is a machine learning method uses bootstrapped training dataset and combines the simplicity of decision trees with extended flexibility to handle new data⁸⁶. The random-forest method allows multiple target class-labels (*e.g.*, nEZ plus three distinct EZ subtypes) over binary classifiers, and here the primary interest was to separate EZ and nEZ contacts.

Supplementary materials

Theory: continuous and discontinuous phase transition in the brain

The classic Brain Criticality framework hypothesizes that, across the brain, neurons operate in a regime of continuous transitions between asynchronous and hyper-synchronous activity, which resembles the phase transition seen in numerous complex systems and is commonly known as the “critical point” or a “critical region”^{7–10,23,87}. Recent theoretical research suggests that the brains could rather be a “quasi-critical” than a “true critical” system^{50,88–90}. True criticality arises in idealized systems where energy is conserved and, with a small and constant drive, the system self-organizes into dynamics with one critical point⁹¹. True criticality is characterized by a continuous (second-order) phase transition between disorder and order³⁵. At the phenomenological level, the ensuing critical dynamics are characterized by stationary fractional-Gaussian statistics⁷⁹ and the emergence of spatial⁹², temporal⁹³, and spatio-temporal power-law behaviour^{88,91}.

On the other hand, quasi-critical systems^{42,87,94} differ categorically from the true critical systems by energy dissipation, as exemplified by forest fires and earthquakes^{95,96}. The loss of energy in quasi-critical systems requires a “loading” mechanism to keep them from becoming quiescent. Simulations suggest that quasi-critical systems require external fine tuning to operate near the critical point^{89,97}. The brain is energetically expensive, accounting for ~20% of the human energy consumption in adults⁹⁸ and of up to ~66% in children⁹⁹ with many physiological mechanisms serving energy replenishment and metabolic regulation¹⁰⁰. In addition to glucose metabolism *per se*, the notion of “loading” mechanisms also encompasses resources and mechanisms that limit neuronal activity levels, for example synaptic vesicle depletion¹⁰¹ and post-synaptic depression^{102,103}, respectively.

Theoretical studies show that neuronal systems with resource-consuming activity and slow resource loading^{19,31,50,90} may indeed exhibit dynamics with a discontinuous (first-order) phase transition when resource demands exceed the loading capacity. This gives rise to spontaneous neuronal *bistability*^{50,90}. For example, in computational models of local populations operating near a marginally stable critical point³⁴, excitatory neurons exhibit bistable firing rates when resource demands are high^{50,90}. Likewise, in networks of such populations, the balance of resource depletion and recovery^{102,103} determines the switching between continuous and bistable transitions³¹.

Slow state-dependent noise ρ controls fast mean field

When investigating the behaviours in Kuramoto model (Fig 1), ρ and κ were held as constant values (eq. 4), which reflect that both variables fluctuate at a much slower rate than the population order R . The variable ρ is a key parameter that controls the degree of bistability in R time series. Under the influence of a weak ρ (eq. 4), the Kuramoto model could demonstrate dynamics resembling that of a *supercritical* stochastic Hopf bifurcation^{25,34} which is also controlled by a weak ρ (eq. 1). Specifically, by gradually increasing κ , a subcritical Kuramoto ensemble would reach a critical point, where the subcritical fixed point (quiescent state) loses stability and a smooth transition to a critical phase takes place. The time course of R in this classic critical scenario follows a Gaussian distribution^{24–26}.

On the other hand, when the Kuramoto model is controlled by a high ρ , the ensemble would express bistable criticality in R that resembles the dynamics of a *subcritical* stochastic Hopf bifurcation. In this scenario, a seemingly quiescent ensemble suddenly shows supercritical hypersynchrony – before the quiescent fixed-point loses stability following further increases in κ . Thereby, the time course of R is characterized as bimodal, supporting erratic switching between low and high activity modes. Our modeling results confirmed this prediction. When ρ is held at high value, the pdf of R as a function of κ (the colour plot in top panel, Fig. 1E) coincides with the prediction of the first order phase transition with a moderate width of bistable regime (top, Supplementary Fig. 1A) – as comparing to theoretical possibility of a much wider bistable regime^{25,35}. When ρ is held at low value (bottom, Fig 1E), the model dynamics accords with the prediction of supercritical Hopf bifurcation as the classic criticality^{7,23}. When the coupling is too strong, the ensemble only dwells on the supercritical state of hypersynchrony. To better demonstrate the effect of ρ on R , we also simulated slowly fluctuating ρ with κ held constant and found that various waveforms of ρ can result in rich bistable dynamics in R , even when the temporal average of ρ was approximately the same (Supplementary Fig. 1B).

The model shows high degree of bistability under the influence of strong state-dependent noise, which results in a tendency to stay in either an up-state or a down-state thus avoiding moderate level of synchrony as expected in the classic criticality models. This up state corresponds to resource-demanding large amplitude oscillations (limit cycle), which eventually leads to depletion and ensuing returning of a low amplitude, subcritical down-state fixed point attractor for recovery. On the extreme spectrum of such bifurcation and the underlying slow variable is epileptic seizure^{52,53,104}.

SEEG cortical sampling statistics

Preprocessing of the SEEG data yielded 7,019 SEEG contacts in various cortical and subcortical gray matter locations (Supplementary Fig. 2A). For investigating the LRTCs and bistability of cortical dynamics that were tentatively considered as normal, contacts recorded from subcortical structures and epileptogenic zones were excluded. Contacts with more than 2.5% samples identified as “spiky” were also excluded (Supplementary Fig. 2A–B, see suppl. Methods). Thereby, 3/7 of available contacts were excluded, and in the resulted 4,142 contacts, a small fraction cannot be reliably assign a parcel by the segmentation software⁷⁷ and therefore were also excluded. This resulted in 4,122 contacts (66.8 ± 24.5 per patient, range: 4 to 123) for analyses. Although the cortical sampling was heterogeneous across patients, with 4,124 cortical nEZ contacts, we were able to cover 90 out of the 100 Schaefer parcels with each parcel sampled by at least 3 subjects and 10 contacts (see also Supplementary Fig 2C–D).

Narrow-band DFA and BiS estimates in MEG and SEEG

SEEG and MEG demonstrated differentiated spectral peaks and magnitude of DFA and BiS estimates (Fig 2). We speculated that these discrepancies between SEEG and MEG might be attributable to two factors (or the combination of both):

First, it was due to different brain states in healthy vs epileptic brains. High degree of bistability has been suggested as an early sign of shift to catastrophic events in ecosystems²⁹, considering the likely universal nature of bistability^{31,35}, high bistability outside of the visual system (e.g., default model and limbic systems) in SEEG data could be a sign of transition to

catastrophic events (*i.e.*, seizures) in the epileptic brain. It could be a great clinical interest to further this line of work to find solutions to reverse high bistability to mildly bistability or smooth phase transitions (*e.g.*, dampening) for eluding the catastrophic events in the brain¹⁰⁵. Moreover, epilepsy is known to affect brain rhythms especially the δ and θ band synchrony^{106,107}, which might be an explanation to the presence of high bistability in slow rhythms below α band in SEEG.

Second, mesoscopic SEEG recording is able to pick up highly local signals and thus more sensitive to specificity such as difference between functional systems, which MEG is unable to. SEEG signals is highly localized to the sampled tissue surrounding the electrode contacts (2mm in length) and minimally affected by signal mixing¹⁰⁸, whereas MEG sensors are at least several centimeters away from the cortex, and its signals are therefore the linear summation of unknown number of cortical sources. According to the central limit theorem, even though individual processes are non-Gaussian, the linear combination of multiple such processes will appear Gaussian.

Topological similarity between LRTCs and bistability

As a preliminary inspection of topological similarity, all-to-all correlations were computed indiscriminately between group-level parcel DFA and BiS estimates, between MEG and SEEG, and across all frequencies. Within metrics, well delineated clusters of a slow and a fast frequency band were observed (along diagonal, Supplementary Fig. 6). There were positive correlations between DFA and BiS in both MEG and SEEG data, and some negative correlations between MEG and SEEG. The frequency-collapsed θ - α and γ -band DFA and BiS estimates were next inspected (Fig. 3C). The similarity between band-clustered cortical maps of 100 Schaefer parcels (Supplementary Fig. 7) converged with narrow-band observation (Supplementary Fig. 6). On systems-level (Supplementary Fig. 8), subjects' band-clustered DFA and BiS were correlated in almost all functional systems except for SEEG θ - α band limbic system (Spearman's rank correlation, $p < 0.01$).

Anatomical specificity of bistability and LRTCs

While there were no systems-wise differences in MEG, SEEG θ - α band BiS and DFA estimates, γ band DFA estimates between Yeo functional systems were different (Kruskal-Wallis test, $p < 0.05$, Fig. 3F).

In MEG, the mean θ - α and γ -band DFA and BiS estimates of Yeo systems appeared similar (thick black lines, Supplementary Fig. 9A–B), confirming high correlations between DFA and BiS estimates observed in both group and systems-level in individuals (Supplementary Fig. 7–8, respectively). With these source-modeled MEG data, we replicated θ - α band resting-state bistability in visual areas that was reported previously in EEG sensors^{26,109}. Furthermore, the visual (VIS) and somatosensory (SM) systems showed higher BiS and DFA, whereas fronto-parietal (FP) and default-model (DEF) systems showed lower BiS and DFA estimates than the null-hypothesis observations (line plots, Supplementary Fig. 9A). The surrogate data of no systems-wise differences were constructed by shuffling Yeo system labels of the parcels. In MEG γ -band, DFA estimates were similar in magnitude to that of θ - α band, but BiS estimates were about half of the magnitude of θ - α band. In particular, the dorsal-attention (DAN) and the limbic (LIM) systems showed the highest γ -band DFA, whereas VIS had the lowest DFA. There were no differences in DFA and BiS estimates between Yeo systems (Wilcoxon's signed-rank

test, $\alpha=0.05$, FDR corrected) confirming the negative findings of the Kruskal-Wallis tests (Fig. 3F).

In SEEG, the θ - α and γ band mean DFA and BiS estimates of Yeo systems appeared different (Supplementary Fig. 9 C–D), which confirmed low correlations in group and systems-level in individuals (Supplementary Fig. 7 & 8). Although visual systems in SEEG and MEG data were comparable in θ - α band BiS and DFA estimates, in SEEG, the visual system showed lowest θ - α band BiS and DFA estimates.

The post-hoc tests (unpaired t-test, $p < 0.05$, FDR corrected) for between-system differences in DFA and BiS estimates (see also Fig. 3F) revealed that DAN showed higher θ - α band DFA, FP and DEF showed higher θ - α band BiS, ventral-attention (VAN) showed lower γ band BiS than most of other systems (interaction matrices, Supplementary Fig. 9 C–D). On the other hand, the VIS and VAN show low DFA estimates. System-wise difference in DFA was only observed between LIM and VAN, which validates the negative finding of systems' effect on mean DFA (γ band pink dot in Fig 3F). On the other hand, the VIS, DAN, and LIM system showed high γ band BiS estimates, and VAN and FP systems have low BiS estimates. The VIS system had highest γ band BiS estimates.

Classifying epileptogenic zones (EZ)

Across subjects, there was large variability in the number of EZ contacts, *i.e.*, the target variables of the classifier (Supplementary Fig. 11A). To ensure there were enough data for the classifier within subjects, a selection criterion was imposed such that each patient should have at least five EZ and five nEZ contacts (red dashed lines, Supplementary Fig. 11A) and at least a total of 30 contacts. Thus, 55 subjects met these criteria and were selected for the classification analysis. On average, each subject had 28.5 ± 17.0 (mean, std) EZ and 66.4 ± 21.2 nEZ, and on population the ratio of EZ:nEZ = 1/2.3 with some variability across Yeo systems, among which subcritical contacts were ten times more likely to be EZ than nEZ (Supplementary Fig. 11B).

On population level, between EZ and nEZ contacts, several bands showed differences in normalized DFA and BiS estimates (Fig. 5A–C). However, we were more interested in classifying EZ contacts in individuals. As a liability check, we classified pooled individual contacts using all nine features, *i.e.*, δ , θ - α , β , and γ band DFA and BiS estimates plus SEEG contact loci in Yeo systems. The classification was performed using the random forest algorithm³⁹ with a total of 5,217 contacts from 55 patients (Supplementary Fig. 11B) with randomly split 20% and 80% as testing and training set, respectively. The analysis of the receiver operating characteristics (ROC) of classification outcome revealed an area under the curve (AUC) of 0.8, and thus confirming useful information among these features for within individual classification.

Global and local feature importance to the random forest classifier were next assessed with SHAP values. The results supported the hypotheses that γ - and β -band BiS, contact-locus and δ -band DFA were indeed the most important features to tell EZ apart from nEZ contacts (Fig. 5D). To better understand the impact of the features on the classification outcome, the within-subject EZ classifications were carried out with four incremental feature sets as using *i*) DFA alone, *ii*) BiS alone, *iii*) combining DFA and BiS, *iv*) combining DFA, BiS, and contact-locus in Yeo systems.

940 The overall classification outcome was variable across Yeo systems and classifying EZ contacts in
 941 limbic systems yielded best outcome (large AUC, Supplementary Fig. 11 D). Using DFA alone, the
 942 algorithm performed poorly (gray line, Supplementary Fig. 11E). The individual subject ROC
 943 curves showed large variability (Supplementary Fig. 11 F–I), and overall combining DFA, BiS, and
 944 contact-locus yielded best outcome (black curve, Supplementary Fig. 11J). Last, the mean AUC of
 945 ROC in Yeo systems (Supplementary Fig. 11E) and in individual patients (Supplementary Fig. 11J)
 946 were compared against 1,000 label-shuffled surrogate data (Fig. 5E).

947 **Supplementary (S.) Figs**

948 **Supplementary Fig. 1**

949 **Fluctuations at two different time-scale in Kuramoto Model.** (A) With coupling strength (κ) held
950 constant and just below the critical point (see Fig 1), slow fluctuations in ρ (the thick gray band) result in
951 (B) diverse patterns of bistability in the fast fluctuating mean field of the Kuramoto model (red).

Supplementary Fig. 2

SEEG electrode selection criteria and population sampling statistics. (A) SEEG contact selection criteria, in brackets are the number of contacts after a specific criterion was applied. (B) The cumulative distribution function (cdf) of contact number per subject after specific criteria were applied; colour code is the same as in (A); inset is the cdf of spike samples out of 10 min resting from 4,693 SEEG contacts, red dashed line indicates the threshold in box 5 in (A). (C) Contact number as a function of number of distinct subjects per parcel (after morphing SEEG contacts into Schaefer 100-parcels); one marker represents one Schaefer 100-parcel, red dashed lines are exclusion criteria, *i.e.*, at least 3 subjects and 10 contacts per parcel. (D) Visualization of distinct subject number and SEEG contact number per Schaefer parcel as shown in (C). cWM-ref: closest white-matter contact reference scheme (Arnulfo et al., 2015).

962 **Supplementary Fig. 3**

963 **SEEG time series, bistability and DFA fitting.** (A–D) Examples of bistability in nEZ SEEG contact signals
964 from four distinct subjects. mFG: middle frontal gyrus; sOG: superior occipital gyrus; mOG: middle occipital
965 gyrus; sTS: superior temporal sulcus. (E–J) Examples of bistable time series and model fitting of an EZ (E) and
966 a nEZ (H) contact. These two contact locations were 19.7 mm apart and recorded with two distinct electrodes
967 from the superior frontal gyrus (sFG) and were referenced with the same nearest white matter contact (Arnulfo et
968 al., 2015). (F) Examples of bi-exponential model fitting for BiS estimates and (G) DFA power-law fitting of 4
969 Hz and 40 Hz narrow-band real and surrogate time series of the EZ contact from (E). (I) Bi-exponential fitting
970 and (J) DFA power-law fitting of 4 Hz and 40 Hz narrow-band real and surrogate time series of the nEZ contact
971 from (H). The DFA fitting plot reads as, when the observation window size doubles (by narrow-band cycle
972 length – irrespective of frequency), the detrended fluctuation increase by a constant rate of log(DF).

973 **Supplementary Fig. 4**

974 **Analysis pipeline for MEG and SEEG data, respectively.**

975 **Supplementary Fig. 5**

976 **The cortical origins of DFA and BiS are similar between neighbouring frequencies but different**
 977 **between slow and fast rhythms demarcated at 40 Hz.** (A) BiS and DFA of each Schaefer parcel (y axis)
 978 observed at narrow-band frequency (x axis) ranging from 2 to 225 Hz for SEEG and MEG data. The 10
 979 gray-out rows are the excluded SEEG parcels due to under sampling. (B) Parcel mean across frequencies
 980 of real data (solid lines) and surrogate (dashed lines). (C) The distance between each frequency's spatial
 981 similarity. (A–C) share the same x-axis (Frequency); (D) Cross-frequency adjacency matrices of
 982 topological similarity, the same as Fig 3A; correlation is the Spearman's r between Schaefer Atlas parcels'
 983 DFA or BiS of two different frequencies.

984 **Supplementary Fig. 6**

985 **All-to-all cross correlations between frequencies, metrics and datasets.** (A) Normalized group-level
986 Schaefer 100-parcel metrics afo frequencies. Normalization $y = (x(i) - \min(x)) / (\max(x) - \min(x))$. (B) All-to-
987 all topological correlations between R, DFA, and BiS and between MEG and SEEG data (Spearman's rank
988 order r , $p > 0.01$, not controlled for FDR).

989 **Supplementary Fig. 7**

990 **DFA and BiS estimates correlated in MEG and SEEG cortical parcels.** (A) The adjacency matrix of
 991 significant Spearman's correlation between band-collapsed DFA and BiS ($p < 10^{-6}$, FDR corrected). (B)
 992 The effect size of the differences between DFA and BiS of MEG and SEEG data in Schaefer parcels
 993 ($n=90$, due to exclusion of 10 SEEG parcels). (C–J) Scatter plots showing correlation between DFA
 994 and BiS estimates in band-clustered all-to-all correlation matrix (top); each data point corresponds to
 995 the group average metrics in one Schaefer 100-parcel ($N_{MEG}=100$; $N_{SEEG} = 90$).

996 **Supplementary Fig. 8**

997 **DFA and BiS are correlated in MEG and SEEG subsystems.** Top left panel is from Fig 3H; top right is
998 the number of subjects observed in each Yeo system: MEG N=18, and SEEG N=50±9, range: 36–60,
999 variable subject N per system in SEEG due to heterogeneous spatial sampling. In scatter plots, each dot
1000 corresponds to the observation from one subject.

1001 **Supplementary Fig. 9**

1002 **Systems-level differences and anatomical localization of DFA and BiS.** These are post hoc tests for
1003 differences in DFA and BiS between Yeo systems (Fig 3I). (A–D) Neuro-anatomical localization of
1004 frequency-collapsed BiS and DFA; marker (*) in colour-bar label indicates significant differences between
1005 systems (Fig 3I). The line plots below the brains are median DFA/BiS of Yeo systems, dashed lines are
1006 5%- and 95%-tile of permutation median ($N_{\text{permutation}} = 10^5$). The adjacency matrices are the significant –
1007 $\log_{10}(p)$ values of pairwise tests for between-system differences (SEEG: unpaired t-test; MEG: Wilcoxon's
1008 signed-rank test, $p < 0.05$, FDR corrected).

1009 **Supplementary Fig. 10**

1010 **Behaviourally relevant DFA and BiS in MEG.** (A) Left: percentage of parcels showed significant
1011 correlation between executive function score (normalized zoo map time) and parcel DFA and BiS; right:
1012 θ - α band BiS as a function of DFA. (B) The behavioural correlations (primary y-axis) and probability of
1013 parcels showing significant correlations (secondary y-axis) on functional systems-level (corrected for FDR).
1014 (C) Top row: cortical maps of behavioural correlations; bottom row: subject neuronal estimates as a
1015 function of executive score from exemplary regions of interest where each marker is the observation from
1016 one subject (C-D) The corresponding p values of correlation tests in Fig 4A and C, respectively; dashed
1017 lines indicate FDR (Benjamini-Hochberg procedure with FDR at 0.05 for 8 neuropsychological scores)
1018 adjusted p values for eight neuropsychological tests. (E) Sorted p -values (thick line) of individual parcel
1019 behavioural correlation for θ - α band DFA and (F) BiS estimates. Thin lines indicate Benjamini-Hochberg
1020 procedure with FDR at 0.05 adjusted p values. (G) Examples of parcel-level behavioural correlations to
1021 Zoom map time rank for a-band DFAs and (H) θ - α band BiS estimates, where each marker indicates one
1022 subject.

1023 **Supplementary Fig. 11**

1024 **Classification of EZ and nEZ SEEG contacts (both cortical and subcortical) using band clustered**
1025 **DFA and BiS estimates.** (A) Number of EZ and nEZ contacts in each subject; solid dots indicate subjects
1026 who met selection criteria for classification analysis; red dashed lines indicate selection criterion: each
1027 subject must have great than five EZ and nEZ contacts. (B) EZ and nEZ contacts located in Yeo's systems
1028 and subcortical (subC) regions, data pooled over 55 subjects from (A). (C) An example of the receiver
1029 operator curve (ROC) of preliminary population-level classification using all contacts and all features with
1030 randomly split test and train set (20% and 80%, respectively); TPR: true positive rate; FPR: false positive
1031 rate; AUC: area under the ROC curve. (D–J) Leave-one(subject)-out classification results using random
1032 forest algorithm. (D) The ROC of each subsystem, classification with the full feature set, D&B(Y). (E) The
1033 AUC yielded from classification using DFA only, BiS estimates only, combining DFA and BiS (D&B), and
1034 full feature set D&B(Y), *i.e.*, DFA, BiS and SEEG contact loci. (F–I): individual (thin lines) and group
1035 average ROC (thick) yielded from classification using varying feature sets. (J) The AUC of ROCs shown
1036 in (F–J), subjects were sorted by the area under the ROC curve of D&B (Y) feature set.