A pleiotropic chemoreceptor facilitates the functional coupling of pheromone production and perception

Cassandra Vernier^{1,2†}, Kathleen M. Zelle^{1,†}, Nicole Leitner¹, Xitong Liang³, Sean Halloran⁴, Jocelyn G. Millar⁴, Yehuda Ben-Shahar^{1,5,*}

¹Department of Biology, Washington University in Saint Louis, 1 Brookings Drive, Saint Louis, MO 63130, USA.

²Carl R. Woese Institute for Genomic Biology, University of Illinois, 1206 W. Gregory Dr., Urbana, IL 61801, USA

³Department of Neuroscience, Washington University School of Medicine, 660 South Euclid Avenue, St. Louis, MO 63110, USA

⁴Department of Entomology, University of California, Riverside, 900 University Avenue, Riverside, CA 92521, USA

⁵Lead Contact

[†]Denotes equal contribution

*Correspondence: benshahary@wustl.edu (Y.B.)

SUMMARY

Optimal mating decisions depend on the robust coupling of signal production and perception because independent changes in either could carry a fitness cost. However, since the perception and production of mating signals are often mediated by different tissues and cell types, the mechanisms that drive and maintain their coupling remain unknown for most animal species. Here, we show that in *Drosophila*, sensory perception and production of an inhibitory mating pheromone are co-regulated by *Gr8a*, a member of the *Gustatory receptor* gene family. Specifically, we found that the pleiotropic action of *Gr8a* independently regulates the perception of pheromones by the chemosensory systems of males and females, as well as their production in the fat body and oenocytes of males. These findings provide a relatively simple molecular explanation for how pleiotropic receptors maintain robust mating signaling systems at the population and species levels.

Keywords: *Drosophila melanogaster*; Vinegar fly; Fruit fly; Cuticular hydrocarbons; *Gr8a*; Oenocyte.

INTRODUCTION

The majority of sexually-reproducing animals use intricate mating signaling systems, which rely on a robust physiological coupling between the production and perception of species-specific signals since any independent changes in either the signal or the capacity to sense it would carry a fitness cost (Boake, 1991; Brooks et al., 2005; Hoy et al., 1977; Shaw et al., 2011; Shaw and Lesnick, 2009; Steiger et al., 2011; Sweigart, 2010; Symonds and Elgar, 2008; Wyatt, 2014). Previously published theoretical models have postulated that the maintenance of robust coupling between the production and perception of mating signals is driven by strong genetic linkage between the cellular and physiological processes that regulate mating-signal production and its perception, or alternatively, via the action of pleiotropic genes that control both processes (Boake, 1991; Butlin and Ritchie, 1989; Butlin and Trickett, 1997; Shaw et al., 2011; Shaw and Lesnick, 2009). Consequently, both mechanisms provide plausible explanations for how mating-signaling systems could remain stable and reliable at the population level while still retaining their capacity for future diversification, as necessitated for speciation (Chebib and Guillaume, 2021; Hoy et al., 1977; Kirkpatrick and Hall, 2004; Lande, 1980; Shaw et al., 2011; Shaw and Lesnick, 2009; Wiley et al., 2012).

Empirical data in support of the contribution of gene-linkage or pleiotropy to the maintenance of coupling between mating signal production and perception at the population level are rare (Chebib and Guillaume, 2021; Hoy et al., 1977; Shaw et al., 2011; Shaw and Lesnick, 2009; Wiley et al., 2012). Additionally, the complex characteristics of mating behaviors, and the species-specific signals that drive them, present a major barrier for identifying the actual molecular mechanisms and candidate pleiotropic genes that support the coupling between the production and perception of specific mating signals (Chenoweth and Blows, 2006; Singh and Shaw, 2012). How the functional coupling of the physiological processes responsible for the production and perception of mating signals remains robust is particularly puzzling since their perception is mediated by the peripheral sensory nervous system, while their production is restricted to specialized, non-neuronal pheromone producing cells (Chung and Carroll, 2015; Chung et al., 2014; McKinney et al., 2015). Notwithstanding, a previous *Drosophila* study has implied that the gene *desat1*, which encodes a fatty acid desaturase, directly contributes to both the perception and production of pheromones (Bousquet et al., 2012). However, subsequent studies have shown that *desat1* expression is enriched in central neurons, and that the effect of *desat1* mutations on the behavioral response to pheromones is not likely to be directly mediated via the modulation of pheromone perception by sensory neurons (Billeter et al., 2009). Furthermore, the effects of *desat1* mutations on the overall CHC profiles of both males and females are broad and lack specificity (Labeur et al., 2002). Together, these data suggest that *desat1* is not likely to act as a pleiotropic factor that directly couples the production and perception of mating pheromones in *Drosophila*. Consequently, the molecular identities of genes that may mediate the genetic and functional linkage between the production of insect mating pheromones by the coenocytes, and their perception by the chemosensory system, remained unknown.

Here we show that some pheromone-driven mating behaviors in *Drosophila* depend on the pleiotropic action of *Gr8a*, a member of the *Gustatory receptor* gene family (Lee et

al., 2012; Shim et al., 2015), which contributes to both the perception of inhibitory mating signals in pheromone-sensing neurons, and independently, to the production of inhibitory mating pheromones in non-neuronal abdominal pheromone-producing oenocytes. Together, these data provide a relatively simple molecular explanation for how genetic linkage could maintain functional coupling between the independent cellular and physiological processes that drive pheromone perception and production.

RESULTS

Some gustatory-like receptors exhibit enriched expression in abdominal tissues

Similar to other insect species, *Drosophila* cuticular hydrocarbons (CHCs), or long-chain fatty acids synthesized by the fat body and oenocytes (Billeter et al., 2009; Gutierrez et al., 2007), provide a hydrophobic desiccation barrier, as well as play an important role as pheromones in regulating diverse behaviors, including mating (Blomquist and Bagnères, 2010; Chung and Carroll, 2015; Ferveur, 2005; McKinney et al., 2015). Specifically, complex blends of CHCs are often utilized by insects to communicate sex identity and female mating status, as well as to define the behavioral reproductive boundaries between closely related species (Ben-Shahar, 2015; Billeter et al., 2009; Chung and Carroll, 2015; Chung et al., 2014; Dweck et al., 2015; Lu et al., 2012; Lu et al., 2014; Yew and Chung, 2015).

While some of the genes and pathways that contribute to CHC synthesis in *Drosophila* are known (Blomquist and Bagnères, 2010; Chung et al., 2014; Ferveur, 2005; Howard and Blomquist, 2005; McKinney et al., 2015), the molecular identities of most CHC receptors remain unknown. Current models stipulate that the perception of volatile CHCs is mediated by olfactory sensory neurons (ORNs) located in the antennae and maxillary palps, while less volatile CHCs are sensed by specialized gustatory-like receptor neurons (GRNs) in the appendages (legs and wings), female genitalia, and the proboscis (Benton et al., 2007; Clowney et al., 2015; Datta et al., 2008; Koh et al., 2014; Kurtovic et al., 2007; Lebreton et al., 2014; Lu et al., 2012; Lu et al., 2014; Pikielny, 2012; Thistle et al., 2012; Toda et al., 2012; van der Goes van Naters and Carlson, 2007; Vijayan et al., 2014).

Consequently, we chose to examine members of the *Gustatory receptor* (*Gr*) gene family as candidate pleiotropic genes that might contribute to both the perception and production of pheromonal mating signals in *Drosophila*. Because several family members have already been implicated in the detection of specific excitatory and inhibitory pheromones (Bray and Amrein, 2003; Miyamoto and Amrein, 2008; Moon et al., 2009; Watanabe et al., 2011), and the majority of genes that encode family members are already known to be enriched in GRNs (Clyne et al., 2000; Dunipace et al., 2001; Scott et al., 2001; Wang et al., 2004), we reasoned that any pleiotropic *Gr* genes should be also expressed in the abdominal oenocytes (Billeter et al., 2009). We tested this by using an RT-PCR screen, which revealed that 24 out of the 60 members of the *Gr* family are expressed in abdominal tissues of adult *Drosophila* (Table 1). This suggests that at least some *Gr* genes may contribute to both the perception and production of mating signals in *Drosophila*.

Gr8a is a chemosensory receptor with sexually dimorphic expression in abdominal cells

Although several members of the *Gr* gene family, including *Gr68a*, *Gr32a*, *Gr66a*, *Gr39a*, and *Gr33a*, were previously linked to the sensory perception of mating pheromones (Bray and Amrein, 2003; Lacaille et al., 2007; Miyamoto and Amrein, 2008; Moon et al., 2009; Watanabe et al., 2011), none of these candidate genes were identified in our initial RT-PCR screen for *Gr* genes expressed in abdominal tissues of either males or females (Table 1). However, *Gr8a*, which was indicated by our screen as being a male-specific abdomen-enriched receptor (Table 1) (Park and Kwon, 2011), was previously shown to play a role in the chemosensation of the non-proteinogenic amino acid L-Canavanine (Lee et al., 2012; Shim et al., 2015). Because our initial expression screen was based on whole-abdomen RNAs, we next used a GAL4 transgenic driver to determine which abdominal cells express *Gr8a*. We found that, as was previously reported (Lee et al., 2012), *Gr8a* is expressed in 14-16 GRNs in the proboscis (Figure 1A-B), as well as in two paired GRNs in the pretarsus of the prothoracic legs in males (Figure 1C) and females (Figure 1D). We also observed *Gr8a* expression in abdominal oenocyte-like cells in males (Figure 1E) but not females (Figure 1F). The male-biased expression in the abdomen was further supported by qRT-PCR analysis (Figure 1G). These data further indicate that in addition to its chemosensory functions, *Gr8a* may also contribute to oenocyte physiology.

To further examine the spatial expression of *Gr8a* in males, we used a membrane bound GFP reporter to trace the axonal projection patterns of *Gr8a*-expressing GRNs in the prothoracic legs. We found that in contrast to the primary, sexually dimorphic *ppk23*-expressing pheromone-sensing GRNs (Lu et al., 2012; Lu et al., 2014; Thistle et al., 2012; Toda et al., 2012), the axons of tarsal *Gr8a*-expressing neurons ascend to the brain and do not cross the midline of the ventral nerve cord (VNC) in males (Figure 1H). Likewise, we found that *Gr8a*-expressing GRNs do not co-express the sex determination factor *fru* (Figure 1I) or the ion channel *ppk23* (Figure 1J), which are were previously assumed to be expressed in all pheromone-sensing GRNs in the fly appendages. These data indicate that *Gr8a*-expressing GRNs in the prothoracic tarsal segments possibly represent a distinct subclass of pheromone-sensing GRNs.

In the male abdomen, we found that *Gr8a* is co-expressed with the oenocyte specific *desat1* driver (Billeter et al., 2009), as well as possibly in *desat1*-negative fat-body-like cells (Figure 1K-M). To better understand how *Gr8a* might function in non-neuronal oenocytes, we next characterized the subcellular localization of the native *Gr8a* protein in abdominal tissues, by using CRISPR/Cas9 genome editing to generate an endogenous GFP-tagged allele of *Gr8a*. Subsequently, immunohistochemical staining of abdominal tissues from *Gr8a*-GFP males with an anti-GFP antibody revealed that the receptor protein is enriched in vacuolar membranes in some oenocyte clusters (Figure 1N). Together, these data indicate that in addition to its possible role in the perception of L-Canavanine, *Gr8a* also contributes to the perception, and possibly production, of mating pheromones in the male.

***Gr8a* activity contributes to mating decisions in females**

We next hypothesized that if *Gr8a* is a pleiotropic gene that independently contributes to the production of a mating pheromone in males, and its chemosensory perception in females, then the knockdown of *Gr8a* in either males or females should have similar effects on female mating behavior. Therefore, we first investigated whether *Gr8a*, and the GRNs that express it, are required for sensory functions associated with female mate choice by using single-pair courtship assays (Lu *et al.*, 2012; Lu *et al.*, 2014). We found that blocking neuronal transmission in female *Gr8a*-expressing GRNs by the transgenic expression of tetanus toxin (TNT) shortens copulation latency relative to wild-type females, when courted by wild-type males (Figure 2A). Similarly, homozygous (Figure 2B) and hemizygous (Figure 2C) *Gr8a*-null females exhibited shorter copulation latencies when courted by wild type males, which can be rescued by driving the expression of the *Gr8a* cDNA by *Gr8a*-GAL4 (Figure 2D). In contrast, genetic manipulations of *Gr8a* did not affect male courtship behavior as measured by courtship latency and index towards wild-type females (Supplemental Figure 1). These data suggest that *Gr8a* is required for regulating female mating receptivity via the chemosensory detection of male-borne inhibitory mating pheromones.

Because *Gr8a* expression is specifically enriched in male oenocytes (Figure 1K-M), we next tested the hypothesis that *Gr8a* also plays a role in the production and/or release of inhibitory mating signals by males. We found that wild-type virgin females exhibited shorter copulation latencies towards *Gr8a* mutant males relative to wild-type controls (Figure 2E). These data indicate that the *Gr8a* mutant males produce and/or release lower levels of inhibitory mating pheromones relative to wild type controls. Together, these behavioral studies suggest that *Gr8a* is a pleiotropic gene that regulates both the production of an inhibitory mating signal in the male oenocytes, and its perception by the chemosensory system in females.

***Gr8a* regulates the copulatory transfer, and the post-mating perception, of inhibitory pheromones by males**

Mating decisions in *D. melanogaster* rely on a balance between excitatory and inhibitory drives (Billeter *et al.*, 2009; Clowney *et al.*, 2015; Kallman *et al.*, 2015; Krupp *et al.*, 2008; Laturney and Billeter, 2016). Therefore, male-borne inhibitory signals may help females optimize mate choices by delaying their decision to copulate with specific males. Additionally, previous studies showed that, in order to increase their fitness, *Drosophila* males transfer inhibitory mating pheromones to females during copulation, which subsequently lowers the overall attractiveness of mated females to other males (Averhoff and Richardson, 1974; Datta *et al.*, 2008; Jin *et al.*, 2008; Kurtovic *et al.*, 2007; Miyamoto and Amrein, 2008; Yang *et al.*, 2009). We found that *Gr8a* mutant males were more likely to court mated females than wild-type controls (Figure 2F), suggesting that *Gr8a* is also required in males for the sensory recognition of the inhibitory signals that label the post-mating status of females. We also found that wild-type males failed to recognize the mating status of wild-type females that were previously mated with *Gr8a* mutant males (Figure 2F). These data indicate that *Gr8a* is also important for the production of inhibitory pheromones that are transferred from

males to females during copulation. Together, these findings suggest that *Gr8a* is responsible for the production and perception of transferrable inhibitory mating signals that advertise post-mating status in females. The simplest overall interpretation of these data is that *Gr8a* is a pleiotropic factor, which independently contributes to the production/ transfer of male inhibitory mating pheromones, as well as their sensory perception in both males and females.

***Gr8a* contributes to quantitative and qualitative attributes of the pheromone profiles of males and mated females**

Because our data indicate that the *Gr8a* mutation has a dramatic effect on the copulation latency of mated females and the ability of males to detect the mating status of females, we hypothesized that *Gr8a* is contributing to the production and/or transfer of an inhibitory pheromone in males. Therefore, we next examined whether the *Gr8a* mutation has a direct effect on qualitative and quantitative aspects of male and mated-female CHC profiles. We found that the overall CHC profile of *Gr8a* mutant males is both qualitatively (Figure 3A) and quantitatively different from that of wild-type males (Figure 3B-C and Table 2). In particular, the *Gr8a* mutation affects the levels of several alkenes and methyl-branched alkanes, which have been implicated in mate choice behaviors in diverse *Drosophila* species (Billeter *et al.*, 2009; Billeter and Levine, 2013; Chung and Carroll, 2015; Chung *et al.*, 2014; Dyer *et al.*, 2014; Shirangi *et al.*, 2009).

Although the exact mechanism by which *Gr8a* might be regulating the levels of specific CHCs remains unknown, we found that the expression levels of the desaturases *desat1* and *CG8630*, which play a role in the biosynthesis of alkenes (Chung and Carroll, 2015), are affected by the *Gr8a* mutation in the male abdomen (Figure 3D). We also found that the overall qualitative aspects of the CHC profiles of wild-type females were not affected by mating with either *Gr8a* mutant or wild-type males (Figure 3E). However, quantitative analyses of individual CHCs revealed that nonacosane (C₂₉) is higher in females that mated with *Gr8a* mutant males relative to those that mated with wild-type males (Figure 3F). Together, these data suggest that *Gr8a* action in oenocytes contributes to the production of some cuticular alkenes and methyl-branched alkanes in males, which possibly function as inhibitory mating pheromones.

Since the *Gr8a* mutation is not spatially restricted in *Gr8a* mutant males, it is possible that at least some of the effects of the *Gr8a* mutation on the pheromone profiles of males are indirectly mediated via its action in pheromone-sensing GRNs, instead of directly mediated via its action in oenocytes. Therefore, we next examined the effect of oenocyte-specific *Gr8a* knockdown on the production of male CHCs. We found that oenocyte-specific *Gr8a* RNAi knockdown in males leads to significant changes in their overall CHC profile relative to control males (Figure 3G). In contrast, fat-body-specific knockdown of *Gr8a* has no effect on the CHC profiles of males (Figure 3H). These data suggest that *Gr8a* is likely to play an oenocyte-specific role in the production of male CHCs. Together, our behavioral and pheromonal data indicate that *Gr8a* action contributes to mating decisions in females by co-regulating the perception of an inhibitory mating pheromone by females and males, as well as its production by males. This is consistent with a pleiotropic function for *Gr8a*.

Gr8a-associated CHCs inhibit normal courtship behaviors

To further characterize whether any of the individual CHCs regulated by *Gr8a* actually function as inhibitory mating pheromones, we tested the effect of perfuming naïve males with individual candidate CHCs on the copulation latency of wild-type females (Ben-Shahar et al., 2010; Ben-Shahar et al., 2007; Leitner and Ben-Shahar, 2020; Lu et al., 2012; Lu et al., 2014). We found that wild-type females did not copulate with *Gr8a* mutant males that were perfumed with the alkenes 9-C₂₅, 7-C₂₅, and 7-C₂₇ (Figure 4A). Similarly, we found that wild-type males exhibited a longer courtship latency and lower courtship index towards wild-type females perfumed with 9-C₂₅ (Figure 4B-D), and exhibited longer copulation latency towards wild-type females perfumed with 7-C₂₅ (Figure 4E-G). In contrast, perfuming wild-type females with 7-C₂₇ had no effect on male courtship or female mating latency (Figure 4H-J). These data suggest that at least some of the CHCs regulated by *Gr8a* activity in the male oenocytes are inhibitory mating pheromones.

Variations in Gr8a contribute to species-specific male pheromonal profiles across the Drosophila genus

As populations diversify, pheromonal signals and their receptors often have to co-evolve to maintain behavioral species boundaries (Boake, 1991; Khallaf et al., 2021; Symonds and Elgar, 2008; Symonds and Wertheim, 2005). One possible mechanism for maintaining the functional coupling of coevolving signal-receptor pairs during speciation is pleiotropy (Boake, 1991; Shaw et al., 2011; Singh and Shaw, 2012). Because our data suggest that *Gr8a* is a pleiotropic pheromone receptor, we tested the hypothesis that cross-species variations in the *Gr8a* coding sequence may have contributed to the rapid evolution of mating pheromones in the *Drosophila* species group (Khallaf et al., 2021; Shahandeh et al., 2018; Shirangi et al., 2009). To test this hypothesis, we first performed a phylogenetic analysis of *Gr8a* orthologs across *Drosophila* species, which indicated that *Gr8a* is a conserved, sexually dimorphic receptor across the *Drosophila* genus (Figure 5A-B). Furthermore, alignment of *Gr8a* proteins across all the major *Drosophila* clades revealed that, in spite of its high overall sequence conservation, the *Gr8a* receptor has at least one phylogenetically variable domain (magenta frame, Figure 5C), which includes the second intracellular and extracellular domains (Figure 5D).

Although the ligand-binding domains of the insect *Gr* gene family have not been identified yet, these data suggest that this phylogenetically variable protein domain in *Gr8a* may contribute to species-specific shifts in ligand-binding specificity and/or sensitivity across the *Drosophila* genus. Therefore, we next tested whether the transgenic rescue of the *Gr8a* null allele via ectopic expression of *Gr8a* cDNAs from different *Drosophila* species is sufficient to drive changes in the CHC profile of *D. melanogaster* males. By using a cross-species male mate-choice assay, we found that while *D. melanogaster* males are generally promiscuous, they do court *D. mojavensis* females at a significantly lower proportion than conspecific females. Because these assays are performed under red light, which eliminates visual mating cues, these data suggested that the lower sex drive towards *D. mojavensis* females is pheromone-

dependent (Figure 5E). Subsequently, we generated transgenic *D. melanogaster* lines which express either the *D. mojavensis* or the *D. melanogaster Gr8a* cDNAs driven by an oenocyte-specific GAL4 in the background of the *Gr8a* null allele. Comparison of male CHC profiles across the two genotypes revealed that rescuing the *Gr8a* mutation by *Gr8a* cDNAs from these two distantly related species resulted in significantly different male CHC profiles (Figure 5F). These data indicate that species-specific *Gr8a* coding variations are sufficient to drive differential CHC production by the male oenocytes, and suggest that pleiotropic pheromone receptors may have played a role in driving the rapidly evolving behavioral mating boundaries in *Drosophila*.

DISCUSSION

The data presented here demonstrate that *Gr8a* is a pleiotropic chemoreceptor that co-regulates the perception and production of an inhibitory pheromonal signal that plays an important role in mating behaviors of both *D. melanogaster* sexes. How *Gr8a*, a member of a canonical chemoreceptor family, might also contribute to the production of pheromonal signals is not obvious. In some better understood secretory cell types, autoreceptors are essential for the regulation of synthesis and secretion rates. For example, dopaminergic and serotonergic cells regulate rates of synthesis and release of their respective neuromodulators by the action of autoreceptors, which act via signaling feedback in response to changes in the extracellular concentrations of the secreted molecule (Ford, 2014; Stagkourakis et al., 2016). Therefore, one possible explanation for how *Gr8a* might regulate the synthesis and/or secretion of specific CHCs is by acting as an oenocyte-intrinsic autoreceptor, which regulates the synthesis of specific CHCs by providing feedback information about their levels in internal stores and/or extracellularly (Figure 6).

Recent studies have indicated that *Drosophila* bitter receptor neurons typically express multiple *Gr* genes, and that bitter receptor ligand specificity is determined via combinatorial heteromeric receptor complexes (Dweck and Carlson, 2020; Shim et al., 2015; Sung et al., 2017). *Gr8a* is specifically required for the sensory perception of the feeding deterrent L-canavanine (Lee et al., 2012; Shim et al., 2015), but not for the detection of other bitter feeding deterrents such as caffeine, strychnine, and umbelliferone (Lee et al., 2009; Poudel et al., 2015). Our data indicate that similar to other *Drosophila* “bitter” taste receptors (Lacaille et al., 2007; Moon et al., 2009), *Gr8a* contributes to inhibitory sensory inputs in the contexts of both feeding and mating decisions. In the context of feeding, *Gr8a*-dependent perception of L-canavanine is mediated via its heterotrimeric interaction with *Gr66a* and *Gr98b* in bitter sensing neurons in the proboscis (Shim et al., 2015). However, although both *Gr66a* and *Gr98b* were also identified in our initial screen for receptors enriched in the adult abdomen, we found that *Gr66a* is expressed in both sexes and *Gr98b* is specifically enriched in females (Table 1). Therefore, we conclude that *Gr8a*-dependent contributions to sensory functions associated with mating decisions are independently driven via its heteromerization with different *Gr* genes than those that drive feeding-specific decisions.

Although we do not yet know the specific chemical identity of the ligand of *Gr8a*, previous studies indicated that at least two inhibitory mating pheromones, 11-cis-vaccenyl acetate (cVA) and CH503, are transferred from males to females during copulation. While our data suggest that the *Gr8a* mutation affects the level of cVA expressed by males, it is unlikely that either cVA or CH503 are the putative *Gr8a* ligands because the volatile cVA acts primarily via the olfactory receptor *Or67d* (Benton *et al.*, 2007; Datta *et al.*, 2008; Kurtovic *et al.*, 2007), and CH503 has been reported to signal via *Gr68a*-expressing neurons, which are anatomically distinct from the *Gr8a* GRNs we describe here (Figure 1A-B) (Shankar *et al.*, 2015; Yew *et al.*, 2009). Instead, our analyses of the effect of the *Gr8a* mutation on the CHC profile (Figure 3), and our results of the perfuming behavioral studies (Figure 4), suggest that the alkenes 5-C₂₅, 7-C₂₅, and 7-C₂₇, which seem to act as inhibitory mating signals as well, are potentially the ligands of *Gr8a*.

Overall, our studies indicate that pleiotropic receptors, such as *Gr8a*, contribute to the physiological coupling between the production and perception of some mating pheromones by acting as both a sensory receptor in pheromone-sensing neurons, and possibly as an autoreceptor for the same chemical in the pheromone-producing oenocytes. Our finding that *Gr8a* is also a sexually dimorphic receptor that is conserved across the *Drosophila* genus, with at least one phylogenetically variable domain (Figure 5A-C), suggests that it might also drive the divergence of mating signaling systems in association with rapid speciation. This is supported by our finding that rescuing the *Gr8a* mutation in *D. melanogaster* with a *Gr8a* cDNA from a distant species, *D. mojavensis*, leads to the development of a male CHC profile that is different from the profile of mutant males rescued with the *D. melanogaster Gr8a* cDNA (Figure 5F).

Studies in other animal species suggest that receptor pleiotropy likely plays a role in mating signaling via other sensory modalities including auditory communication in crickets (Heinen-Kay *et al.*, 2020; Hoy *et al.*, 1977; Wiley *et al.*, 2012) and visual communication in fish (Fukamachi *et al.*, 2009). While the specific genes and signaling pathways that mediate the coupling of the mating signals and their receptors in these mating systems remain mostly unknown, these data suggest that genetic linkage in signal-receptor pairs important for mating communication is likely to be more common than previously thought. Therefore, the genetic tractability of *D. melanogaster*, in combination with the diversity of mating communication systems in this species-rich phylogenetic group, provide a unique opportunity for understanding the evolution and mechanisms that drive and maintain the robustness of mating systems at the genetic, molecular, and cellular levels.

ACKNOWLEDGMENTS

We thank members of the Ben-Shahar lab for comments on earlier versions of the manuscript. We thank Joshua Krupp (University of Toronto) for assistance with perfuming studies, Nabeel Chowdhury and Deanna Simon for assistance with qRT-PCR analysis, and Paula Kiefel for technical help with generating transgenic flies. This work was supported by NSF grants 1322783, 1754264, and 1707221, and NIH grant NS089834 awarded to Y. B-S. Stocks obtained from the Bloomington *Drosophila* Stock

Center (NIH P40OD018537) were used in this study. Wild-type *Drosophila* species were obtained from the National *Drosophila* Species Stock Center at Cornell University.

AUTHOR CONTRIBUTIONS

K.M.Z., C.V., J.G.M. and Y.B-S designed experiments. K.M.Z., C.V., N.L., X.L., S.H., J.G.M. and Y.B-S collected and analyzed data. K.M.Z., C.V., N.L. and Y.B-S wrote the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

398

399

400

401

402

Figure 1. *Gr8a* is a sexually dimorphic chemosensory receptor. (A-F) *Gr8a* is expressed in the proboscis (A-B) and prothoracic legs (C-D) of both males (top) and females (bottom), but is only expressed in the abdomen of males (E-F). Cells labeled by *Gr8a*-GAL4. (G) *Gr8a* has sexually dimorphic mRNA expression in the bodies of flies.

403 Relative mRNA levels were measured by real-time quantitative RT-PCR. **, $p < 0.01$
404 Mann Whitney Rank Sum Test, $n = 3/\text{group}$. **(H-J)** *Gr8a*-expressing GRNs represent a
405 distinct subclass of pheromone sensing neurons. (H) Axonal projection patterns in the
406 T1 VNC neuromere in a *Gr8a*-GAL4>UAS-CD8::GFP male (green). Arrowheads,
407 individual axons; dashed line, midline of the VNC. Magenta, neuropil marker (nc82). (I)
408 Confocal z-stack of a male *fruP1*-LexA>LexAop-myrGFP (green); *Gr8a*-GAL4>UAS-
409 Red-Stinger (magenta) prothoracic leg. (J) Confocal z-stack of a male *ppk23*-
410 LexA>LexAop-CD8::GFP (green); *Gr8a*-GAL4>UAS-Red-Stinger (magenta) prothoracic
411 leg. **(K-M)** *Gr8a* is expressed in oenocytes and other abdominal cells. Confocal z-stack
412 images of oenocytes in a *Gr8a*-GAL4>UAS-CD8::GFP; *desat1*>*luciferase* male: (K)
413 *desat1* (green); (L) *Gr8a* (magenta); (M) Merge. Blue, DAPI. White arrow, expression of
414 *Gr8a* in oenocytes; yellow arrows, expression of *Gr8a* in other abdominal cells. **(N)**
415 GR8A protein is enriched in abdominal cells. Confocal z-stack of a GFP-tagged *Gr8a*
416 allele in male abdominal cells; green, anti-GFP; blue, DAPI. Scale bars = 50 μm .

Figure 2. *Gr8a* activity contributes to the perception and production of an inhibitory signal associated with mating decisions in males and females. (A) Blocking neural activity in female *Gr8a*-expressing sensory neurons (*Gr8a>TNT*) shortens copulation latency relative to wild-type controls (*Gr8a>TNT^{inactive}*). **(B-C)** Homozygous (B) or hemizygous (C) *Gr8a* null females show shortened copulation latency relative to wild-type controls. Df(1)BSC663 is a deficiency that covers the *Gr8a* locus. Df(1)BSC754 was used as a control. **(D)** Expression of *Gr8a* cDNA with the *Gr8a* promoter (*Gr8a⁺;Gr8a^{tg}*) rescues the copulation latency phenotype in *Gr8a* mutant females. **(E)** Wild-type females exhibit shortened copulation latency when courted by *Gr8a* mutant males relative to wild-type males. **(F)** *Gr8a* mutant males do not recognize the mating status of females, and have a reduced transfer of inhibitory mating pheromones during copulations. Female, female genotype; Sperm donor, genotype of males mated first with focal females; Focal male, genotypes of experimental males presented with mated females. Different letters above bars indicate statistically significant Tukey's HSD *post hoc* contrasts between groups. Panels C, D, and F: p<0.05 ANOVA, n>15/group. Panels A, B, E: *, p<0.05, Mann Whitney Rank Sum Test, n>15/group. All assays performed under red light conditions.

Figure 3. The *Gr8a* mutation affects the pheromone profiles of males and mated females. (A) Wild-type (wt) and *Gr8a* mutant (*Gr8a*⁻) males differ in CHC profile. p<0.001, Permutation MANOVA. **(B-C)** The *Gr8a* mutation affects the levels of individual CHCs in males. (B) CHCs found at high proportions in males. (C) CHCs found at low proportions in males. Only affected CHCs are shown. See Table 2 for the complete list. *, p<0.05, **, p<0.001, Student's t-test or Mann Whitney Rank Sum Test, n=6 (*Gr8a*⁻) or 7 (wt). **(D)** The *Gr8a* mutation affects the expression level of several desaturase genes. Only affected genes are shown. See Table 3 for the complete list. *, p<0.05, Student's t-test, n=4/group. **(E)** Females mated with wild-type or *Gr8a* mutant males do not differ in CHC profile. p=0.570, Permutation MANOVA. **(F)** Nonacosane (C₂₉) differs between females mated with wild-type and *Gr8a* mutant males. See Table 4 for complete list of mated-female CHCs. *, p<0.05, Student's t-test, n=6/group. **(G)** Control (*desat1* > *GFP*-RNAi) and oenocyte-specific *Gr8a* knockdown (*desat1* > *Gr8a*-RNAi) males differ in CHC profile. p<0.001, Permutation MANOVA. **(H)** Control (*r4* > *GFP*-RNAi) and fat body-specific *Gr8a* knockdown (*r4* > *Gr8a*-RNAi) males do not differ in CHC profile. p = 0.298, Permutation MANOVA. Panels A, E, G, and H depicted as Nonmetric Multidimensional Scaling (NMDS) plots with Bray-Curtis dissimilarity.

Figure 4. *Gr8a*-associated alkenes inhibit normal courtship behaviors. (A) Perfuming males with exaggerated amounts of several alkenes increases copulation latency compared to control males. **(B-D)** Perfuming females with 9-C₂₅ increases courtship latency (B), decreases courtship index (C), but does not affect copulation latency (D) compared to control females. **(E-G)** Perfuming females with 7-C₂₅ does not affect courtship latency (E) or index (F), but increases copulation latency (G) compared to control females. **(H-J)** Perfuming females with 7-C₂₇ does not affect courtship latency (H), courtship index (I), or copulation latency (J) compared to control females. Asterisks above bars indicate statistically significant contrasts compared to control flies, $p < 0.05$, Kruskal-Wallis Test followed by Dunn's Test (A) or Mann Whitney Rank Sum Test (B-J), $n = 15/\text{group}$.

Figure 5. Sexually dimorphic *Gr8a* expression across the *Drosophila* genus may contribute to species-specific differences in male CHC profiles. (A) Phylogenetic tree of *Drosophila* *Gr8a* proteins. Substitution rate = 0.2. (B) *Gr8a* mRNA expression is enriched in males relative to females across *Drosophila*. Black, males; white, females. *, $p < 0.05$; **, $p < 0.01$; Mann Whitney Rank Sum Test, $n = 4/\text{group}$. Live *D. grimshawi* was not analyzed because live specimens were not available at the *Drosophila* Species Stock Center (DSSC). (C) Multiple aligned amino acid sequences of *Gr8a* protein sequences from 12 species across *Drosophila*. The magenta dashed box highlights a putative hypervariable protein domain. Numbers on top of alignment indicate amino acid number. Black, 100% identical; Dark Gray, 80-100% similar; Light Gray, 60-80% similar; White, less than 60% similar (Blosum62 score matrix, threshold=1). Bars below consensus represent overall level of amino acid conservation. (D) *Gr8a* protein topology. Boxes, transmembrane domains; Red lines, intracellular domain; Blue lines, extracellular domains. (E) In female choice assays, *D. melanogaster* males court females from most other *Drosophila* species first at an equal proportion as *D. melanogaster* females, but court *D. mojavensis* females first at a lower proportion than *D. melanogaster* females. Assays performed under red light. *, $p < 0.05$, Pearson's Chi-squared test. (F) *Gr8a* mutant *D. melanogaster* males with oenocyte-specific *D. melanogaster* *Gr8a* rescue differ in CHC profile from *Gr8a* mutant *D. melanogaster* males with oenocyte-specific *D. mojavensis* *Gr8a* rescue. Depicted as NMDS plot with Bray-Curtis dissimilarity; *Gr8a*⁻; *desat1* > *Gr8a*^{*melanogaster*}, *D. melanogaster* *Gr8a* oenocyte rescue; *Gr8a*⁻; *desat1* > *Gr8a*^{*mojavensis*}, *D. mojavensis* *Gr8a* oenocyte rescue. Bold letters in legend denote statistical significance, $p < 0.05$, permutation MANOVA.

Figure 6. Model for the pleiotropic action of *Gr8a* in the perception and production of pheromones. (A) *Drosophila* male. The location of CHC-producing oenocytes is shown in magenta. (B) *Gr8a*-expressing GRNs are located at the last tarsal segment of the prothoracic legs. (C) *Gr8a* functions as an inhibitory pheromone receptor in a specific subset of leg GRNs. (D) Oenocytes are the primary CHC-producing cells in the male abdomen. (E) *Gr8a* functions as an autoreceptor in oenocytes, which regulates CHC synthesis [I-II] and/or CHC secretion [III] via signaling feedback loops.

501 **TABLES**

Gene	Male	Female
<i>Gr2a</i>	-	+
<i>Gr8a</i>	+	-
<i>Gr10a</i>	+	+
<i>Gr21a</i>	-	+
<i>Gr22a</i>	+	-
<i>Gr22e</i>	+	+
<i>Gr36c</i>	+	-
<i>Gr58c</i>	+	+
<i>Gr59a</i>	+	+
<i>Gr59b</i>	+	+
<i>Gr63a</i>	+	-
<i>Gr64a</i>	+	-
<i>Gr64b</i>	+	+
<i>Gr64c</i>	+	+
<i>Gr64d</i>	+	-
<i>Gr66a</i>	+	+
<i>Gr89a</i>	+	+
<i>Gr93a</i>	-	+
<i>Gr93d</i>	+	+
<i>Gr97a</i>	+	+
<i>Gr98a</i>	+	+
<i>Gr98b</i>	-	+
<i>Gr98c</i>	+	+
<i>Gr98d</i>	+	+

Table 1. Candidate *Gr* genes expressed in male and/or female abdomens. Plus and minus signs indicate whether RT-PCR products were detected. Only genes with positive PCR products in at least one sex are shown.

R.T.	Compound	wt % total	<i>Gr8a</i> ⁻ % total	p value
12.31	C ₂₁	0.589	1.102	<0.001
13.24	Unknown	0.071	0.191	<0.001
14.2	C ₂₂	0.893	0.949	0.339
14.34	7-C ₂₂	0.362	0.490	<0.001
15.25	Unknown	0.128	0.226	<0.001
16.22	C ₂₃ & 9-C ₂₃	13.682	12.671	<0.001
16.4	7-C ₂₃	30.201	35.478	0.002
16.53	5-C ₂₃	2.772	2.389	<0.001
16.71	CvA	10.240	15.700	<0.001
18.03	C ₂₄ & 9-C ₂₄	0.578	0.367	<0.001
18.19	8-C ₂₄	0.742	0.493	<0.001
18.27	7-C ₂₄	0.579	0.402	<0.001
18.37	6-C ₂₄	0.395	0.233	<0.001
18.46	5-C ₂₄	0.040	0.047	0.943
19.09	2Me-C ₂₄	2.426	3.240	<0.001
19.95	C ₂₅	0.000	2.038	0.001
20.02	C ₂₅ & 9-C ₂₅	6.195	1.793	<0.001
20.18	7-C ₂₅	14.781	4.344	<0.001
20.42	5-C ₂₅	0.296	0.000	<0.001
22.89	2Me-C ₂₆	6.007	6.933	0.002
23.7	C ₂₇	1.127	0.719	0.052
23.94	7-C ₂₇	0.308	0.083	<0.001
26.46	2Me-C ₂₈	4.992	5.775	0.078
27.25	C ₂₉	0.351	0.317	0.574
29.89	2Me-C ₃₀	1.245	2.353	<0.001

Table 2. Male CHCs. Retention time (R.T.), compound, percent total (% total), and p-value (Student's t-test or Mann Whitney Rank Sum Test) of each compound as part of the total pheromonal bouquet for wild-type (wt) and *Gr8a* mutant (*Gr8a*⁻) males.

Gene	wt mRNA fold difference	<i>Gr8a</i> ⁻ mRNA fold difference	p value
<i>desat1</i>	1.282	0.931	0.037
<i>desat2</i>	1.413	1.270	0.506
<i>CG8630</i>	0.951	1.429	0.012
<i>CG9747</i>	0.838	0.525	0.343
<i>CG9743</i>	1.060	0.959	0.373
<i>CG15331</i>	0.774	1.000	0.21

Table 3. Desaturase gene expression. Relative mRNA expression of each desaturase gene for wild-type (wt) and *Gr8a* mutant (*Gr8a*⁻) males. Statistics via Student's t-test.

R.T.	Compound	WT % total	<i>Gr8a</i> % total	p value
12.31	C ₂₁	0.234	0.242	0.712
14.2	C ₂₂	0.330	0.352	0.512
14.34	7-C ₂₂	0.029	0.029	0.958
15.25	Unknown	0.238	0.200	0.128
16.09	C ₂₃	6.481	6.580	0.748
16.4	7-C ₂₃	2.454	2.729	0.149
16.39	7,11-C ₂₃	0.473	0.426	0.208
16.53	5-C ₂₃	0.254	0.235	0.416
16.71	CvA	1.349	1.135	0.233
17.99	C ₂₄	0.660	0.732	0.209
18.19	8-C ₂₄	0.156	0.174	0.334
19.09	2Me-C ₂₄	1.214	1.104	0.155
19.95	C ₂₅	5.404	5.683	0.336
20.02	C ₂₅ & 9-C ₂₅	3.141	3.851	0.699
20.18	7-C ₂₅	3.144	2.999	0.422
20.25	7,11-C ₂₅	1.822	1.684	0.315
20.42	5-C ₂₅	0.526	0.555	0.375
20.47	5,9-C ₂₅	0.686	0.704	0.589
22.89	2Me-C ₂₆	10.010	9.247	0.1554
23.7	C ₂₇	3.616	3.393	0.24
23.8	9-C ₂₇	2.498	2.739	0.937
23.94	7-C ₂₇	3.819	4.885	0.24
24.1	7,11-C ₂₇	21.464	19.770	0.18
24.28	5,9-C ₂₇	2.557	2.438	0.12
25.85	7,11-C ₂₈	0.658	0.645	0.573
26.46	2Me-C ₂₈	4.779	5.022	0.306
27.25	C ₂₉	0.577	0.693	0.01
27.7	7,11-C ₂₉	19.411	19.589	0.824
29.89	2Me-C ₃₀	1.359	1.465	0.17
31.03	7,11-C ₃₁	0.562	0.606	1

510 **Table 4.** Mated-female CHCs. Retention time (R.T.), compound, percent total (% total),
511 and p-value (Student's t-test or Mann Whitney Rank Sum Test) of each compound as
512 part of the total pheromonal bouquet for females mated with wild-type (wt) or *Gr8a*
513 mutant (*Gr8a⁻*) males.

METHODS

Animals. Flies were maintained on a standard cornmeal medium under a 12:12 light-dark cycle at 25 Celsius. Unless specifically stated, the *D. melanogaster* Canton-S (CS) strain served as wild-type control animals. UAS-TNT-E, UAS-TNT-IMP-V1-A, UAS-mCD8::GFP, UAS-myr::GFP, UAS-Red Stinger, Df(1)BSC663, Df(1)BSC754, *Gr8a-GAL4*, *Gr8a*¹, *desat1-Gal4*, *r4-Gal4* and *fruP1-LexA* fly lines were from the Bloomington Stock center. Originally in the *w*¹¹¹⁸ background, the *Gr8a*¹ null allele was outcrossed for six generations into the CS wild-type background, which was used as a control.

Likewise, the *desat1-Gal4* allele was outcrossed for six generations into this *Gr8a* null background. *PromE(800)-GAL4* and *PromE(800)>Luciferase* were from Joel Levine (The University of Toronto, Canada). The following *Drosophila* species were obtained from the San Diego Stock Center: *D. simulans* 14011-0251.192, *D. sechellia* 14021-0248.03, *D. yakuba* 14021-0261.01, *D. erecta* 14021-0224.00, *D. ananassae* 14024-0371.16, *D. pseudoobscura* 14011-0121.104, *D. persimilis* 14011-0111.50, *D. willistoni* 14030-0811.35, *D. mojavensis* 15081-1352.23, and *D. virilis* 15010-1051.118. The UAS-*Gr8a* transgenic lines were generated by cloning the *D. melanogaster* and *D. mojavensis* *Gr8a* cDNAs into pUAST-attB vector by using 5' EcoRI and 3' NotI restriction sites, followed by Φ C31 integrase-dependent transgenesis at a Chromosome 2 attP landing site (2L:1476459), as previously described (Zheng et al., 2014). Subsequently, both UAS-*Gr8a*^{CDNA} lines were transgressed into the *Gr8a*¹ background, resulting in complete substitution of the endogenous *Gr8a* with expression of a *Gr8a* ortholog. The *ppk23-LexA* line was generated by integrating our previously described *ppk23* promoter DNA fragment (Lu et al., 2012) into the pBPnlsLexA::p65Uw plasmid (Pfeiffer et al., 2010), followed by Φ C31 integrase-dependent transgenesis as above.

The GFP-tagged allele of *Gr8a* was generated via CRISPR/Cas9-dependent editing using a modified “scarless” strategy by using the sgRNA CGAGCAAGGCGGGAACGATT and a 3XP3>dsRed in the donor plasmid as a reporter for edited animals as previously described (Hill et al., 2017; Hill et al., 2019). Control lines with matching genetic backgrounds were established by selecting DsRed-negative injected animals. The final tagged *Gr8a* allele was generated by removing the DsRed cassette via the introduction of the *piggyBac* transposase (Hill et al., 2019).

Immunohistochemistry. To visualize the expression pattern of *Gr8a* in males and females, *Gr8a-GAL4* flies (Lee et al., 2012) were crossed to UAS-CD8::EGFP and live-imaged at 5 days old using a Nikon-A1 confocal microscope. To demonstrate *Gr8a* expression in oenocytes, abdomens from *Gr8a-GAL4/UAS-myr::GFP*; *PromE(800)>Luciferase* flies were dissected and immunostained as previously described (Lu et al., 2012; Zheng et al., 2014) by using a Rabbit anti-GFP (1:1000; A-11122, Thermo Fisher Scientific) and a mouse anti-luciferase (1:100; 35-6700, Thermo Fisher Scientific) antibodies followed by AlexaFluor 488 anti-rabbit and AlexaFluor 568 anti-mouse secondary antibodies (Both at 1:1000; Thermo Fisher Scientific). To visualize the GR8A protein, abdomens of control flies and flies with CRISPR/Cas9 generated GFP-tagged GR8A were dissected and immunostained as previously described (Lu et al., 2012; Zheng et al., 2014) using a Rabbit anti-GFP antibody (1:1000; A-11122, Thermo Fisher Scientific) followed by AlexaFluor 488 anti-rabbit secondary antibody (1:1000; Thermo Fisher Scientific).

mRNA expression. Newly eclosed flies were separated by sex under CO₂ and aged for 5 days on standard cornmeal medium. On day 6, flies were placed in a -80°C freezer until RNA extraction. To separate body parts, frozen flies were placed in 1.5ml microcentrifuge tubes, dipped in liquid nitrogen, and then vortexed repeatedly until heads, appendages, and bodies were clearly separated. Total RNA was extracted using the Trizol Reagent (Thermo Fisher Scientific) separately from heads, bodies, and appendages for *Gr8a* expression and from bodies for desaturase enzyme genes. cDNAs were synthesized using SuperScript II reverse transcriptase (Thermo Fisher Scientific) with 500 ng total RNA in a 20 uL reaction. Real-time quantitative RT-PCR was carried out as previously described with *Rp49* as the loading control gene (Hill *et al.*, 2017; Hill *et al.*, 2019; Lu *et al.*, 2012; Lu *et al.*, 2014; Zheng *et al.*, 2014). Primer sequences are described in Supplemental Tables 1-3.

Courtship Behavior Assays. Single-pair assays were performed as we have previously published (Lu *et al.*, 2012, 2014). In short, newly eclosed males were kept individually on standard fly food in plastic vials (12 x 75mm). Newly eclosed virgin females were kept in groups of 10 flies. All behaviors were done with 4-7 day-old animals, which were housed under constant conditions of 25° C and a 12h:12h light-dark cycle. Courtship was video recorded for 10 min for male courtship and 15 min for female mating receptivity. Male courtship latency and index were measured as previously described (Lu *et al.*, 2012; Lu *et al.*, 2014). Female receptivity index was defined as the time from the initiation of male courtship until copulation was observed. Unless otherwise indicated, assays were performed under normal light conditions.

Male mate-choice assays were performed in round courtship arenas. Briefly, one *D. melanogaster* virgin female and one interspecific virgin female was decapitated under CO₂ and placed in the arena. One virgin male *D. melanogaster* was then aspirated into the arena and behavior was video recorded for 10 minutes. The first female courted (by male wing extension) was noted. Male mate-choice assays were performed under red light conditions.

Perfuming studies. Synthetic compounds were synthesized by J.G.M. Perfuming studies were performed using a modified protocol from (Billeter *et al.*, 2009). In short, 3 mg of each compound was dissolved in 6 mL hexane (Sigma-Aldrich #139386-500ML) and 0.5 mL was pipetted into individual 2 mL glass vials fitted with 9mm PTFE lined caps (Agilent Crosslab, Santa Clara, CA, USA). The hexane was evaporated under a nitrogen gas flow, such that a residue of the compound was left around the bottom one-third of the vial. Control vials were prepared using hexane without a spiked compound. Vials were kept at -20°C until use. Flies used in these trials were collected as described above, kept in single sex groups and aged for 4 days on standard cornmeal medium at 25°C. 24 hours before perfuming, 20 flies of one or the other sex were placed in glass vials containing standard cornmeal medium (12 x 75mm). To perfume the flies, these groups of 20 flies were dumped without anesthesia into each 2 mL vial containing the compound of interest, and were vortexed at medium-low speed for 3 pulses of 20 seconds punctuated by 20 second rest periods. Flies were transferred to new food vials and were allowed to recover for one hour. Perfumed flies were then used in courtship behavior assays as described above and the remaining flies were used in pheromone analyses to verify compound transfer. The genotype of flies that were perfumed differed

based upon the genotype with the lower amount of each compound as determined in Figure 3 (B, C, F). In all cases, compound transfer was verified by CHC extraction and GC/MS (Supplemental Table 4).

Phylogenetic analysis. Protein sequences of GR8A orthologs from the 12 sequenced *Drosophila* reference genomes were aligned by using the ClustalW algorithm in the Omega package (Sievers et al., 2011), followed by ProtTest (v2.4) to determine the best model of protein evolution (Abascal et al., 2005). Subsequently, Akaike and Bayesian information criterion scores were used to select the appropriate substitution matrix. We then used a maximum likelihood approach and rapid bootstrapping within RAXML v 7.2.8 Black Box on the Cipres web portal to make a phylogenetic tree (Miller et al., 2010). Visualizations of the bipartition files were made using FigTree v1.3.1 (<http://tree.bio.ed.ac.uk/software/figtree/>).

Pheromone Analysis. Virgin flies were collected upon eclosion under a light CO² anesthesia and kept in single-sex vials in groups of 10 with 6 biological replications for each genotype and sex. Virgin flies were aged for 5 days on standard cornmeal medium at 25°C. To collect mated flies, both females and males were aged for 3 days before single mating pairs were placed in a standard fly vial with standard cornmeal food for 24 hours. The pair was then separated for 24 hours before collection. Copulation was confirmed by the presence of larvae in the vials of mated females several days later. On the morning of day 5, flies were anesthetized under light CO² and groups of five flies were placed in individual scintillation vials (VWR 74504-20). To extract CHCs, each group of flies was covered by 100 μ L hexane (Sigma-Aldrich #139386-500ML) containing 50 μ g/mL hexacosane (Sigma-Aldrich #241687-5G) and was washed for ten minutes. Subsequently, hexane washes were transferred into a new 2 ml glass vial containing a 350 μ L insert (Thermo Scientific C4000-LV-1W) and were stored at -20°C until shipment to the Millar laboratory.

Analyses of CHC profiles were done by gas chromatography and mass spectroscopy (GC-MS) in the Millar laboratory at UC Riverside as previously described (Chung et al., 2014). Peak areas were measured, and data was normalized to known quantity of internal standard hexacosane (Sigma-Aldrich #241687-5G). The relative proportion of each compound in each sample was calculated and used in further statistical analysis.

Statistical Analysis. All statistical analyses were performed in R (v 3.6.2). The following functions were used in the base statistics package: `t.test()` (t-test), `wilcox.test()` (Mann-Whitney Rank Sum Test), `aov()` (ANOVA), `TukeyHSD()` (Tukey's HSD post hoc test), `Kruskal.test()` (Kruskal-Wallis test), `chisq.test()` (Pearson's Chi-squared test). Kruskal-Wallis post hoc was performed using the `dunn.test.control` function in the PMCMR package (Pohlert, 2014). Qualitative CHC data were analyzed through a permutation MANOVA using the `adonis` function in the `vegan` package of R with Bray-Curtis dissimilarity measures (Oksanen, 2011). CHC profile data were visualized using non-metric multidimensional scaling (`metaMDS`) function in the `vegan` package of R (Oksanen, 2015) using Bray-Curtis dissimilarity, and either 2 or 3 dimensions in order to minimize stress to <0.1.

649

650 **Supplemental Figure 1.** *Gr8a* has no effect on male courtship latency or index toward
651 wild-type females. (A) Courtship latency (s) and index of control *Gr8a-gal4/UAS-IMP-*
652 *TNT-V1A* (*Gr8a>TNT^{inactive}*) and *Gr8a-gal4/UAS-TNT-E* (*Gr8a>TNT*) mutant males
653 towards wild-type females. (B) Courtship latency (s) and index of wild-type (CS) and
654 *Gr8a null* (*Gr8a⁻*) males toward wild-type decapitated females. Mann Whitney Rank
655 Sum Test, not significant ($p>0.05$), $n=15$ /group.

Gene	Forward Primer	Reverse Primer
<i>Gr10a</i>	AGCGCGAGAGCTTTCGGATGC	AAAGTCCACACAGCATGGGCGG
<i>Gr10b</i>	GCGACTGTCCAGAGGATGTGGC	CAGCTGATCAAAGCGAATGCGC
<i>Gr21a</i>	AACTACCGGCTCGCCGATCG	AGGTTGCCCCGTGGTCGATGA
<i>Gr22a</i>	TGGGACCTTTGGCAGGGAATCG	TGCGATGACTGCAGAAGAGGGT
<i>Gr22b</i>	GGTTGTGCATTGCCGCATGCG	CTGCAAAGCCAAGCGAACTCGT
<i>Gr22c</i>	GCCGCCAGCGATCTAGCTGAG	CAATGACCGCACAGCAATGCGA
<i>Gr22e</i>	TGGTCTTCGTTCAAGCGCTGGT	GCCGGCGATGGCTGCAGAATAAT
<i>Gr22f</i>	GATCGCCTGCTGAAGCTGAGTGA	GTGGGTGCACAGCCATGCGAAT
<i>Gr23a</i>	AGCGATTTGTGGTGACCGCCA	CATCCGCTGCCTCGCGTTGA
<i>Gr28a</i>	TCGCCCTGCCGCACATCAATA	GGGTGAGAGTTCCAAGGCGCT
<i>Gr28b</i>	ACTGGAGGCATAGTGACTCCCT	AGTGGTCAAGGCCCCGCTGAT
<i>Gr2a</i>	TTTGCGACCGAACGGCCCAG	GTTGAAGAAGCCAGCGGCGC
<i>Gr32a</i>	ACACGGTCACCACCAACTGCT	ACCAAGTATGTTGTGACGGCCGAA
<i>Gr33a</i>	ATCGCCACGCACCCCTTACC	GCTCACCACAAAGCAGGCCG
<i>Gr36a</i>	GAGGTCGTGTCGTTGCAGCCC	GTGCTCGCTGACGCCATCTGT
<i>Gr36b</i>	TGGCATGCAAGGGCTTATGGCT	CCGCAGTTGAGCCACGCGTG
<i>Gr36c</i>	CGGCAGAGCAAACATGTTGCACG	ACTGCAAGCCCAGTCCCAGGT
<i>Gr39a</i>	TACCCCGAACC GG GACTGGG	CCAGAGCGAAAAATCCATAAGCGGT
<i>Gr39b</i>	GGGCGCCAAC TG CACATTGG	TCCAATCGAGGGATGGCAGGACA
<i>Gr43a</i>	GCTGGCCCTGGCTCCATATGC	AACCCCTTGTTCTTGGCTGCGGC
<i>Gr47a</i>	GGTCGGGCCAGAGAAATCGGC	CCCGCTTGGGATGGAGGGCC
<i>Gr47b</i>	GCCCTGGGATCCCGAGTACG	TGCCCCGATGTCAACTGGTGATCT
<i>Gr57a</i>	GGCAAGGAGATGCCTTCGCCT	GCCCATCTTCTGGGTCATGGCG
<i>Gr58a</i>	TGTGGATGACGTGACGCGGA	ACGGATGACAACGAAGGTGATCCGA
<i>Gr58b</i>	ACCTCCTGCAACAACACTGGCC	CCGATTCTGCCTCAGCTGCATGG
<i>Gr58c</i>	ACCGCTGTAAGTGGTTGGGACT	TCCAACGATGTGCAGACGTCTCA
<i>Gr59a</i>	GCCAAAGTTCTTCGCCCCCGA	TCCAATCGCGTGCTGCTCTCG
<i>Gr59b</i>	CAGCCAAGCACTCGTGAGTGA	TGGCAAACCACATGCTGCGATT
<i>Gr59c</i>	CCGAGTGCTGGCGGATCGAAC	CTGGCCAGTTGCAGTTGGAAGT
<i>Gr59d</i>	CGTCGATTGCTGGATCAACGCG	TCGACCAAAC TC GAAAAGGCCG
<i>Gr59e</i>	CCTCTCTGGGAGGAGTCCGTGC	AGGGCAACGCCCATTAGCGACT

<i>Gr59f</i>	GCCATGCACGTGGGAAAGGTGT	ACATTGCCCTGAACCGACTTGGA
<i>Gr5a</i>	GCCTGCAACGTGCTCGTCCT	TTGAGCGGGACAAGGCGCAG
<i>Gr61a</i>	AAAGTGCGGCGCCAGAAGCA	GTGCGACCAGGAGCGAGCAA
<i>Gr63a</i>	CGCAAGAAGGGTGACGCGGTG	AAGGGTCCACTCAGGGAGCGC
<i>Gr64a</i>	CGCGTGCAATTGGCTGCCTTG	GTCCGCCGCCGTGAGGAATAC
<i>Gr64b</i>	GCCCGCGGTCTTTTGGACAGA	CATGGCCAGGACGAGTGAGCG
<i>Gr64c</i>	ATGCACTGGCGCATCAGGCA	CCGCCCTCGCAGAAGCTCTT
<i>Gr64d</i>	ACCGCGCCTGGATGGTGTTC	TGCAACCACGGCACCTTGCAT
<i>Gr64e</i>	TGGCCACCACTTGCTCGCTG	ACGCCGCGTGTTAGCCGAAA
<i>Gr64f</i>	TCTGGCTAGGCAGTGGCCCC	GCATGGGCCACCGATGGCAT
<i>Gr66a</i>	GAGGACCGCAGCCTAAAGGCC	CGCGGTCAACTGGTCGGAGAC
<i>Gr77a</i>	CGCAGTTCAGGCCGTCTGCTT	CCACAGGCGCTGACGGCAAATA
<i>Gr85a</i>	CTCTCGCCTACATGGCACGC	ATAGTGACGTGCGTGTGCTGC
<i>Gr89a</i>	GCTGCATGGAGTTCCCCTGACC	ACATTGCCAGTGGGCGTGAGC
<i>Gr8a</i>	ACATACGCATCGCCGTGGACTG	ATGTAGGTGCCCACGGAACAGG
<i>Gr92a</i>	TGTCTCCGTTGAGGGAGCGGT	TGGGTGACATCGAATAGGCCCA
<i>Gr93a</i>	GGTGAGGCAGTCAGAAGTGCC	CCTCGGACGCCTCAAAGCCAC
<i>Gr93b</i>	ACCACGCCATTCTGCGAAGAC	GCCCAAAGGAAACACTCGCAGC
<i>Gr93c</i>	TGCGACGCCTGAGTCTGGAGA	CCCAAGGGTCTAACGCGAAACTCG
<i>Gr93d</i>	ACTCGGGAACGTGCTCTGGAT	AGAGGCCCAACAAGTTAACCTGA
<i>Gr97a</i>	TGCGTCGCTTCTGCGAGTTAGC	GCGTGCTGCAGATCGAATCGC
<i>Gr98a</i>	GTTGAGCACACAAGGGTGGCTG	GCTGTTGCTTGCTGCTGCTCTGA
<i>Gr98b</i>	TTGGAGGCTGGAGGGCGATGT	ATGGTGACAAGCAACCCGCCA
<i>Gr98c</i>	GCCAGGATTCAGGAGCTGTGCG	CGCTGGTATTTTGTCTGTAGCCGA
<i>Gr98d</i>	CTGGCTGGACGCATTTGGGGC	ATGTCAAAGAGGCCACCGCATGT
<i>Gr9a</i>	TGTCCTCTGTCTGGGCGACTGG	ACAAGGGACACGAACTGCAGGA

Supplemental Table 1. Nucleotide sequences for qRT-PCR primers for *D. melanogaster Gr* genes.

Species	Gr8a Forward Primer	Gr8a Reverse Primer
<i>D. melanogaster</i>	TGACCATCAACATACGCATCG	CGTATATGAAGGCGGGAATCTC
<i>D. simulans</i>	GAACCTTTTCGCTGCAACTCC	ACTTCGGTATAAACTGGATGGTG
<i>D. sechellia</i>	GAGATTCCCGCCTTCATATACG	GAGTTGCAGCGAAAAGTTCTG
<i>D. erecta</i>	CAGATTCAGAACTTTTCGCTGC	GGTGTAGATCATGTAGGTGCC
<i>D. yakuba</i>	TGCCTCGGACTAACAATTCTG	GTGTAGATCATGTAGGTGCCC
<i>D. ananassae</i>	AATGTACCGAAGTTTCCAGGG	GCGGGTATGATCAGGAAATAGTC
<i>D. pseudoobscura</i>	CCCGTTTCCGTGACAATATTG	ACCATCTACATATCCGTTGCC
<i>D. persimilis</i>	TTTCGCTTCTCCACACTGAC	AGGCGGGCAATATCAAAGAG
<i>D. willistoni</i>	GAAATGTTGCCCAGAATAGCC	CCCAAAGCATGTATAACCACTG
<i>D. virilis</i>	TCTTCAGATCCAAAACCTTTTCGC	TTGGGCATCAGTTGTACGG
<i>D. mojavensis</i>	CATATACCCGCCTTTCTCTACAC	GTTCGTGCAGAATTTGTAGCG

Species	Rp49 Forward Primer	Rp49 Reverse Primer
<i>D. melanogaster</i>	ATCTTGGGCCTGTATGCTG	TGTGATGGGAATTCGTGGG
<i>D. simulans</i>	GTCGGATCGATATGCTAAGCTG	CAGATACTGTCCCTTGAAGCG
<i>D. sechellia</i>	CATACAGGCCCAAGATCGTG	CAGATACTGTCCCTTGAAGCG
<i>D. erecta</i>	GTCGGATCGATATGCTAAGCTG	CAGATACTGTCCCTTGAAGCG
<i>D. yakuba</i>	CATACAGGCCCAAGATCGTG	GGCATCAGATACTGTCCCTTG
<i>D. ananassae</i>	TACAGGCCCAAGATCGTTAAG	GTACTGACCCTTGAAGCGAC
<i>D. pseudoobscura</i>	CCAGCTCCAAAATGACGATTC	TCAATACCCTTAGGCTTGCG
<i>D. persimilis</i>	AAGCACTTCATCCGTCACC	TCAATACCCTTAGGCTTGCG
<i>D. willistoni</i>	AAGCACTTCATCCGTCACC	GTTGGGCATCAGATATTGGC
<i>D. virilis</i>	AGTCGGATCGTTATGCTAAGTTG	TGGAGGGTACGCTTGTTTG
<i>D. mojavensis</i>	ACCATTTCGTCCAGCATACAG	TTGGCCCTTGAAGCGAC

Supplemental Table 2. Nucleotide sequences for qRT-PCR primers for *D. melanogaster Gr8a*, *Rp49* and orthologs.

Gene	Forward Primer	Reverse Primer
<i>desat1</i>	TTCTGAACGTAACCTGGCTG	GGGAAGACGTGATGGTAGTTATG
<i>desat2</i>	TTTACTGGAATGAGACGCTGG	CTGGGTAGGATTCATGGTCTTG
<i>Fad2</i>	GTGCCTCAGTCTCAATCTCATC	GTGGTAGTTGTGGTATCCCTC
<i>CG8630</i>	ATGTGATGGCTAAGATCGGAC	TGGTCGGAATGGCACTATTG
<i>CG9747</i>	TCAAGTGGGACAAAGTCATCC	GCCGTAAC TCCGAATCCG
<i>CG9743</i>	GGTACTTCTGGAACGAGGATC	TCAGTGGACATGAGGTTCTTG
<i>CG15531</i>	CTCAAGGATGTGGATATGTCGG	ATGGAGGTGGCTAGGGAG
<i>rp49</i>	CACCAAGCACTTCATCCG	TCGATCCGTAACCGATGT

Supplemental Table 3. Nucleotide sequences for qRT-PCR primers for *D. melanogaster* desaturase enzyme genes.

Compound	Genotype	Sex	ng/sample	ng/fly
C ₂₅	wt	male	1350	270
C ₂₅	wt	male	1170	234
C ₂₅	wt	male	27730	5546
9-C ₂₅	Gr8a-	male	67390	13478
9-C ₂₅	Gr8a-	male	59380	11876
9-C ₂₅	Gr8a-	male	1000	200
7-C ₂₅	Gr8a-	male	43790	8758
7-C ₂₅	Gr8a-	male	31330	6266
7-C ₂₅	Gr8a-	male	24130	4826
5-C ₂₅	Gr8a-	male	12050	2410
5-C ₂₅	Gr8a-	male	14160	2832
5-C ₂₅	Gr8a-	male	1870	374
C ₂₇	Gr8a-	male	890	178
C ₂₇	Gr8a-	male	720	144
C ₂₇	Gr8a-	male	263	52.6
7-C ₂₇	Gr8a-	male	36710	7342
7-C ₂₇	Gr8a-	male	21250	4250
7-C ₂₇	Gr8a-	male	15910	3182
C ₂₉	wt	male	260	52
C ₂₉	wt	male	340	68
C ₂₉	wt	male	890	178
9-C ₂₅	wt	female	14060	2812
9-C ₂₅	wt	female	13830	2766
7-C ₂₅	wt	female	23200	4640
7-C ₂₅	wt	female	12780	2556
7-C ₂₇	wt	female	1010	202
7-C ₂₇	wt	female	1080	216

Supplemental Table 4. Amount (ng) of each perfumed compound measured in each sample of perfumed flies (5 flies per sample).

Supplemental Data Legends:

Figure 1 Data: Average qRT-PCR C_t scores across 3 technical replicates for each fly sample for *Gr8a* and *rp49* (control).

Figure 2 Data: Copulation latency (s) or courtship index of single-pair courtship trials corresponding to Figure 2.

Figure 3 Data: Amount (ng) of each compound extracted from each sample (5 flies/sample) in Figure 3. Average qRT-PCR C_t scores across 3 technical replicates for each fly sample for every desaturase gene measured and *rp49* (control). Genes were run on separate qPCR plates, indicated by plate number in parentheses.

Figure 4 Data: Copulation latency (s) of single-pair courtship trials with perfumed males. Courtship latency (s), courtship index, and copulation latency (s) of single-pair courtship trials with perfumed females.

Figure 5 Data: Average qRT-PCR C_t scores across 3 technical replicates for each fly sample for *Gr8a* and *rp49* (control) across *Drosophila* species. Number of flies courted first by *D. melanogaster* males in choice assays. Amount (ng) of each compound extracted from each sample (5 flies/sample) in Figure 5.

Figure S1 Data: Courtship latency (s) and index of single-pair courtship trials corresponding to Supplemental Figure 1.

REFERENCES

- Abascal, F., Zardoya, R., and Posada, D. (2005). ProtTest: selection of best-fit models of protein evolution. *Bioinformatics* 21, 2104-2105. bti263 [pii] 10.1093/bioinformatics/bti263.
- Averhoff, W.W., and Richardson, R.H. (1974). Pheromonal control of mating patterns in *Drosophila melanogaster*. *Behav Genet* 4, 207-225.
- Ben-Shahar, Y. (2015). Editorial overview: Neuroscience: How nervous systems generate behavior: lessons from insects. *Current Opinion in Insect Science* 12, v-vii. 10.1016/j.cois.2015.10.005.
- Ben-Shahar, Y., Lu, B., Collier, D.M., Snyder, P.M., Schnizler, M., and Welsh, M.J. (2010). The *Drosophila* gene *CheB42a* is a novel modifier of Deg/ENaC channel function. *PLoS One* 5, e9395. 10.1371/journal.pone.0009395.
- Ben-Shahar, Y., Nannapaneni, K., Casavant, T.L., Scheetz, T.E., and Welsh, M.J. (2007). Eukaryotic operon-like transcription of functionally related genes in *Drosophila*. *Proc Natl Acad Sci U S A* 104, 222-227. 10.1073/pnas.0609683104.
- Benton, R., Vannice, K.S., and Vosshall, L.B. (2007). An essential role for a CD36-related receptor in pheromone detection in *Drosophila*. *Nature* 450, 289-293. 10.1038/nature06328.
- Billeter, J.C., Atallah, J., Krupp, J.J., Millar, J.G., and Levine, J.D. (2009). Specialized cells tag sexual and species identity in *Drosophila melanogaster*. *Nature* 461, 987-991. 10.1038/nature08495.
- Billeter, J.C., and Levine, J.D. (2013). Who is he and what is he to you? Recognition in *Drosophila melanogaster*. *Curr Opin Neurobiol* 23, 17-23. 10.1016/j.conb.2012.08.009.
- Blomquist, G.J., and Bagnères, A.-G. (2010). *Insect hydrocarbons : biology, biochemistry, and chemical ecology* (Cambridge University Press).

718 Boake, C.R. (1991). Coevolution of senders and receivers of sexual signals: Genetic
 719 coupling and genetic correlations. *Trends Ecol Evol* 6, 225-227. 10.1016/0169-
 720 5347(91)90027-U.

721 Bousquet, F., Nojima, T., Houot, B., Chauvel, I., Chaudy, S., Dupas, S., Yamamoto, D.,
 722 and Ferveur, J.F. (2012). Expression of a desaturase gene, *desat1*, in neural and
 723 nonneural tissues separately affects perception and emission of sex pheromones in
 724 *Drosophila*. *Proc Natl Acad Sci U S A* 109, 249-254. 10.1073/pnas.1109166108.

725 Bray, S., and Amrein, H. (2003). A putative *Drosophila* pheromone receptor expressed
 726 in male-specific taste neurons is required for efficient courtship. *Neuron* 39, 1019-1029.

727 Brooks, R., Hunt, J., Blows, M.W., Smith, M.J., Bussiere, L.F., and Jennions, M.D.
 728 (2005). Experimental evidence for multivariate stabilizing sexual selection. *Evolution* 59,
 729 871-880. Doi 10.1111/J.0014-3820.2005.Tb01760.X.

730 Butlin, R.K., and Ritchie, M.G. (1989). Genetic coupling in mate recognition systems:
 731 what is the evidence? *Biological Journal of the Linnean Society* 37, 237-246.
 732 10.1111/j.1095-8312.1989.tb01902.x.

733 Butlin, R.K., and Trickett, A.J. (1997). Can Population Genetic Simulations Help to
 734 Interpret Pheromone Evolution? In *Insect Pheromone Research*, R.T. Cardé, and A.K.
 735 Minks, eds. (Springer US), pp. 548-562. 10.1007/978-1-4615-6371-6_47.

736 Chebib, J., and Guillaume, F. (2021). Pleiotropy or linkage? Their relative contributions
 737 to the genetic correlation of quantitative traits and detection by multitrait GWA studies.
 738 *Genetics* 219. 10.1093/genetics/iyab159.

739 Chenoweth, S.F., and Blows, M.W. (2006). Dissecting the complex genetic basis of
 740 mate choice. *Nat Rev Genet* 7, 681-692. 10.1038/nrg1924.

741 Chung, H., and Carroll, S.B. (2015). Wax, sex and the origin of species: Dual roles of
 742 insect cuticular hydrocarbons in adaptation and mating. *Bioessays* 37, 822-830.
 743 10.1002/bies.201500014.

744 Chung, H., Loehlin, D.W., Dufour, H.D., Vaccarro, K., Millar, J.G., and Carroll, S.B.
 745 (2014). A single gene affects both ecological divergence and mate choice in *Drosophila*.
 746 *Science* 343, 1148-1151. 10.1126/science.1249998.

747 Clowney, E.J., Iguchi, S., Bussell, J.J., Scheer, E., and Ruta, V. (2015). Multimodal
 748 Chemosensory Circuits Controlling Male Courtship in *Drosophila*. *Neuron* 87, 1036-
 749 1049. 10.1016/j.neuron.2015.07.025.

750 Clyne, P.J., Warr, C.G., and Carlson, J.R. (2000). Candidate Taste Receptors in
 751 *Drosophila*. *Science* 287, 1830-1834. 10.1126/science.287.5459.1830.

752 Datta, S.R., Vasconcelos, M.L., Ruta, V., Luo, S., Wong, A., Demir, E., Flores, J.,
 753 Balonze, K., Dickson, B.J., and Axel, R. (2008). The *Drosophila* pheromone cVA
 754 activates a sexually dimorphic neural circuit. *Nature* 452, 473-477. nature06808 [pii]
 755 10.1038/nature06808.

756 Dunipace, L., Meister, S., McNealy, C., and Amrein, H. (2001). Spatially restricted
 757 expression of candidate taste receptors in the *Drosophila* gustatory system. *Curr Biol*
 758 11, 822-835. S0960-9822(01)00258-5 [pii].

759 Dweck, H.K., Ebrahim, S.A., Thoma, M., Mohamed, A.A., Keeseey, I.W., Trona, F.,
 760 Lavista-Llanos, S., Svatos, A., Sachse, S., Knaden, M., and Hansson, B.S. (2015).
 761 Pheromones mediating copulation and attraction in *Drosophila*. *Proc Natl Acad Sci U S*
 762 *A* 112, E2829-2835. 10.1073/pnas.1504527112.

763 Dweck, H.K.M., and Carlson, J.R. (2020). Molecular Logic and Evolution of Bitter Taste
 764 in *Drosophila*. *Curr Biol* 30, 17-30 e13. 10.1016/j.cub.2019.11.005.

765 Dyer, K.A., White, B.E., Sztepanacz, J.L., Bewick, E.R., and Rundle, H.D. (2014).
 766 Reproductive character displacement of epicuticular compounds and their contribution
 767 to mate choice in *Drosophila subquinaria* and *Drosophila recens*. *Evolution* 68, 1163-
 768 1175. 10.1111/evo.12335.

769 Ferveur, J.F. (2005). Cuticular hydrocarbons: their evolution and roles in *Drosophila*
 770 pheromonal communication. *Behav Genet* 35, 279-295. 10.1007/s10519-005-3220-5.

771 Ford, C.P. (2014). The role of D2-autoreceptors in regulating dopamine neuron activity
 772 and transmission. *Neuroscience* 282, 13-22. 10.1016/j.neuroscience.2014.01.025.

773 Fukamachi, S., Kinoshita, M., Aizawa, K., Oda, S., Meyer, A., and Mitani, H. (2009).
 774 Dual control by a single gene of secondary sexual characters and mating preferences in
 775 medaka. *BMC Biol* 7, 64. 10.1186/1741-7007-7-64.

776 Gutierrez, E., Wiggins, D., Fielding, B., and Gould, A.P. (2007). Specialized hepatocyte-
 777 like cells regulate *Drosophila* lipid metabolism. *Nature* 445, 275-280.
 778 10.1038/nature05382.

779 Heinen-Kay, J.L., Nichols, R.E., and Zuk, M. (2020). Sexual signal loss, pleiotropy, and
 780 maintenance of a male reproductive polymorphism in crickets. *Evolution* 74, 1002-1009.
 781 10.1111/evo.13952.

782 Hill, A., Zheng, X., Li, X., McKinney, R., Dickman, D., and Ben-Shahar, Y. (2017). The
 783 *Drosophila* Postsynaptic DEG/ENaC Channel ppk29 Contributes to Excitatory
 784 Neurotransmission. *J Neurosci* 37, 3171-3180. 10.1523/JNEUROSCI.3850-16.2017.

785 Hill, A.S., Jain, P., Folan, N.E., and Ben-Shahar, Y. (2019). The *Drosophila* ERG
 786 channel seizure plays a role in the neuronal homeostatic stress response. *PLoS Genet*
 787 15, e1008288. 10.1371/journal.pgen.1008288.

788 Howard, R.W., and Blomquist, G.J. (2005). ECOLOGICAL, BEHAVIORAL, AND
 789 BIOCHEMICAL ASPECTS OF INSECT HYDROCARBONS. *Annual Review of*
 790 *Entomology* 50, 371-393. 10.1146/annurev.ento.50.071803.130359.

791 Hoy, R.R., Hahn, J., and Paul, R.C. (1977). Hybrid cricket auditory behavior: evidence
 792 for genetic coupling in animal communication. *Science* 195, 82-84.
 793 10.1126/science.831260.

794 Jin, X., Ha, T.S., and Smith, D.P. (2008). SNMP is a signaling component required for
 795 pheromone sensitivity in *Drosophila*. *Proc Natl Acad Sci U S A* 105, 10996-11001.
 796 0803309105 [pii]
 797 10.1073/pnas.0803309105.

798 Kallman, B.R., Kim, H., and Scott, K. (2015). Excitation and inhibition onto central
799 courtship neurons biases *Drosophila* mate choice. *eLife* 4. 10.7554/elife.11188.

800 Khallaf, M.A., Cui, R., Weissflog, J., Erdogmus, M., Svatos, A., Dweck, H.K.M.,
801 Valenzano, D.R., Hansson, B.S., and Knaden, M. (2021). Large-scale characterization
802 of sex pheromone communication systems in *Drosophila*. *Nat Commun* 12, 4165.
803 10.1038/s41467-021-24395-z.

804 Kirkpatrick, M., and Hall, D.W. (2004). Sexual selection and sex linkage. *Evolution* 58,
805 683-691. 10.1111/j.0014-3820.2004.tb00401.x.

806 Koh, T.W., He, Z., Gorur-Shandilya, S., Menuz, K., Larter, N.K., Stewart, S., and
807 Carlson, J.R. (2014). The *Drosophila* IR20a clade of ionotropic receptors are candidate
808 taste and pheromone receptors. *Neuron* 83, 850-865. 10.1016/j.neuron.2014.07.012.

809 Krupp, J.J., Kent, C., Billeter, J.C., Azanchi, R., So, A.K., Schonfeld, J.A., Smith, B.P.,
810 Lucas, C., and Levine, J.D. (2008). Social experience modifies pheromone expression
811 and mating behavior in male *Drosophila melanogaster*. *Curr Biol* 18, 1373-1383. S0960-
812 9822(08)01034-8 [pii]
813 10.1016/j.cub.2008.07.089.

814 Kurtovic, A., Widmer, A., and Dickson, B.J. (2007). A single class of olfactory neurons
815 mediates behavioural responses to a *Drosophila* sex pheromone. *Nature* 446, 542-546.
816 nature05672 [pii]
817 10.1038/nature05672.

818 Labeur, C., Dallerac, R., and Wicker-Thomas, C. (2002). Involvement of *desat1* gene in
819 the control of *Drosophila melanogaster* pheromone biosynthesis. *Genetica* 114, 269-
820 274. 10.1023/a:1016223000650.

821 Lacaille, F., Hiroi, M., Twele, R., Inoshita, T., Umemoto, D., Manière, G., Marion-Poll, F.,
822 Ozaki, M., Francke, W., Cobb, M., et al. (2007). An Inhibitory Sex Pheromone Tastes
823 Bitter for *Drosophila* Males. *PLoS ONE* 2, e661. 10.1371/journal.pone.0000661.

824 Lande, R. (1980). The Genetic Covariance between Characters Maintained by
825 Pleiotropic Mutations. *Genetics* 94, 203-215. 10.1093/genetics/94.1.203.

826 Laturney, M., and Billeter, J.C. (2016). *Drosophila melanogaster* females restore their
827 attractiveness after mating by removing male anti-aphrodisiac pheromones. *Nat*
828 *Commun* 7, 12322. 10.1038/ncomms12322.

829 Lebreton, S., Grabe, V., Omondi, A.B., Ignell, R., Becher, P.G., Hansson, B.S., Sachse,
830 S., and Witzgall, P. (2014). Love makes smell blind: mating suppresses pheromone
831 attraction in *Drosophila* females via Or65a olfactory neurons. *Scientific reports* 4, 7119.
832 10.1038/srep07119.

833 Lee, Y., Kang, M.J., Shim, J., Cheong, C.U., Moon, S.J., and Montell, C. (2012).
834 Gustatory receptors required for avoiding the insecticide L-canavanine. *J Neurosci* 32,
835 1429-1435. 10.1523/JNEUROSCI.4630-11.2012.

836 Lee, Y., Moon, S.J., and Montell, C. (2009). Multiple gustatory receptors required for the
837 caffeine response in *Drosophila*. *Proc Natl Acad Sci U S A* 106, 4495-4500.
838 0811744106 [pii]
839 10.1073/pnas.0811744106.

840 Leitner, N., and Ben-Shahar, Y. (2020). The neurogenetics of sexually dimorphic
841 behaviors from a postdevelopmental perspective. *Genes Brain Behav* 19, e12623.
842 10.1111/gbb.12623.

843 Lu, B., LaMora, A., Sun, Y., Welsh, M.J., and Ben-Shahar, Y. (2012). ppk23-Dependent
844 chemosensory functions contribute to courtship behavior in *Drosophila melanogaster*.
845 *PLoS Genet* 8, e1002587. 10.1371/journal.pgen.1002587.

846 Lu, B., Zelle, K.M., Seltzer, R., Hefetz, A., and Ben-Shahar, Y. (2014). Feminization of
847 pheromone-sensing neurons affects mating decisions in *Drosophila* males. *Biology*
848 *open* 3, 152-160. 10.1242/bio.20147369.

849 McKinney, R.M., Vernier, C., and Ben-Shahar, Y. (2015). The neural basis for insect
850 pheromonal communication. *Curr Opin Insect Sci* 12, 86-92.
851 10.1016/j.cois.2015.09.010.

852 Miller, M.A., Pfeiffer, W., and Schwartz, T. (2010). Creating the CIPRES Science
853 Gateway for inference of large phylogenetic trees. 14-14 Nov. 2010. pp. 1-8.

854 Miyamoto, T., and Amrein, H. (2008). Suppression of male courtship by a *Drosophila*
855 pheromone receptor. *Nat Neurosci* 11, 874-876. nn.2161 [pii]
856 10.1038/nn.2161.

857 Moon, S.J., Lee, Y., Jiao, Y., and Montell, C. (2009). A *Drosophila* gustatory receptor
858 essential for aversive taste and inhibiting male-to-male courtship. *Curr Biol* 19, 1623-
859 1627. S0960-9822(09)01487-0 [pii]
860 10.1016/j.cub.2009.07.061.

861 Oksanen, J. (2011). Multivariate analysis of ecological communities in R: vegan tutorial.
862 R package version 1, 1-43.

863 Oksanen, J. (2015). Vegan: an introduction to ordination. URL <http://cran.r-project.org/web/packages/vegan/vignettes/introvegan.pdf> 8, 19.

865 Pfeiffer, B.D., Ngo, T.T., Hibbard, K.L., Murphy, C., Jenett, A., Truman, J.W., and Rubin,
866 G.M. (2010). Refinement of tools for targeted gene expression in *Drosophila*. *Genetics*
867 186, 735-755. 10.1534/genetics.110.119917.

868 Pikielny, C.W. (2012). Sexy DEG/ENaC channels involved in gustatory detection of fruit
869 fly pheromones. *Sci Signal* 5, pe48. 10.1126/scisignal.2003555.

870 Pohlert, T. (2014). The pairwise multiple comparison of mean ranks package (PMCMR).

871 Poudel, S., Kim, Y., Kim, Y.T., and Lee, Y. (2015). Gustatory receptors required for
872 sensing umbelliferone in *Drosophila melanogaster*. *Insect Biochem Mol Biol* 66, 110-
873 118. 10.1016/j.ibmb.2015.10.010.

874 Scott, K., Brady, R., Jr., Cravchik, A., Morozov, P., Rzhetsky, A., Zuker, C., and Axel, R.
875 (2001). A chemosensory gene family encoding candidate gustatory and olfactory
876 receptors in *Drosophila*. *Cell* 104, 661-673. S0092-8674(01)00263-X [pii].

877 Shahandeh, M.P., Pischedda, A., and Turner, T.L. (2018). Male mate choice via
878 cuticular hydrocarbon pheromones drives reproductive isolation between *Drosophila*
879 species. *Evolution* 72, 123-135. 10.1111/evo.13389.

880 Shankar, S., Chua, J.Y., Tan, K.J., Calvert, M.E.K., Weng, R., Ng, W.C., Mori, K., and
881 Yew, J.Y. (2015). The neuropeptide tachykinin is essential for pheromone detection in a
882 gustatory neural circuit. *eLife* 4. 10.7554/elife.06914.

883 Shaw, K.L., Ellison, C.K., Oh, K.P., and Wiley, C. (2011). Pleiotropy, “sexy” traits, and
884 speciation. *Behavioral Ecology* 22, 1154-1155.

885 Shaw, K.L., and Lesnick, S.C. (2009). Genomic linkage of male song and female
886 acoustic preference QTL underlying a rapid species radiation. *Proc Natl Acad Sci U S A*
887 106, 9737-9742. 10.1073/pnas.0900229106.

888 Shim, J., Lee, Y., Jeong, Y.T., Kim, Y., Lee, M.G., Montell, C., and Moon, S.J. (2015).
889 The full repertoire of *Drosophila* gustatory receptors for detecting an aversive
890 compound. *Nat Commun* 6, 8867. 10.1038/ncomms9867.

891 Shirangi, T.R., Dufour, H.D., Williams, T.M., and Carroll, S.B. (2009). Rapid evolution of
892 sex pheromone-producing enzyme expression in *Drosophila*. *PLoS Biol* 7, e1000168.
893 10.1371/journal.pbio.1000168.

894 Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R.,
895 McWilliam, H., Remmert, M., Söding, J., et al. (2011). Fast, scalable generation of high-
896 quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems*
897 *Biology* 7.

898 Singh, N.D., and Shaw, K.L. (2012). On the scent of pleiotropy. *Proc Natl Acad Sci U S*
899 *A* 109, 5-6. 10.1073/pnas.1118531109.

900 Stagkourakis, S., Kim, H., Lyons, D.J., and Broberger, C. (2016). Dopamine
 901 Autoreceptor Regulation of a Hypothalamic Dopaminergic Network. *Cell reports*.
 902 10.1016/j.celrep.2016.03.062.

903 Steiger, S., Schmitt, T., and Schaefer, H.M. (2011). The origin and dynamic evolution of
 904 chemical information transfer. *Proc Biol Sci* 278, 970-979. 10.1098/rspb.2010.2285.

905 Sung, H.Y., Jeong, Y.T., Lim, J.Y., Kim, H., Oh, S.M., Hwang, S.W., Kwon, J.Y., and
 906 Moon, S.J. (2017). Heterogeneity in the *Drosophila* gustatory receptor complexes that
 907 detect aversive compounds. *Nat Commun* 8, 1484. 10.1038/s41467-017-01639-5.

908 Sweigart, A.L. (2010). The genetics of postmating, prezygotic reproductive isolation
 909 between *Drosophila virilis* and *D. americana*. *Genetics* 184, 401-410.
 910 10.1534/genetics.109.111245.

911 Symonds, M.R., and Elgar, M.A. (2008). The evolution of pheromone diversity. *Trends*
 912 *Ecol Evol* 23, 220-228. 10.1016/j.tree.2007.11.009.

913 Symonds, M.R., and Wertheim, B. (2005). The mode of evolution of aggregation
 914 pheromones in *Drosophila* species. *J Evol Biol* 18, 1253-1263. 10.1111/j.1420-
 915 9101.2005.00971.x.

916 Thistle, R., Cameron, P., Ghorayshi, A., Dennison, L., and Scott, K. (2012). Contact
 917 Chemoreceptors Mediate Male-Male Repulsion and Male-Female Attraction during
 918 *Drosophila* Courtship. *Cell* 149, 1140-1151. S0092-8674(12)00523-5 [pii]
 919 10.1016/j.cell.2012.03.045.

920 Toda, H., Zhao, X., and Dickson, B.J. (2012). The *Drosophila* Female Aphrodisiac
 921 Pheromone Activates ppk23+ Sensory Neurons to Elicit Male Courtship Behavior. *Cell*
 922 *reports doi:10.1016/j.celrep.2012.05.007*, 599-607. 10.1016/j.celrep.2012.05.007.

923 van der Goes van Naters, W., and Carlson, J.R. (2007). Receptors and neurons for fly
 924 odors in *Drosophila*. *Curr Biol* 17, 606-612. S0960-9822(07)01020-2 [pii]
 925 10.1016/j.cub.2007.02.043.

926 Vijayan, V., Thistle, R., Liu, T., Starostina, E., and Pikielny, C.W. (2014). *Drosophila*
 927 pheromone-sensing neurons expressing the ppk25 ion channel subunit stimulate male
 928 courtship and female receptivity. *PLoS Genet* 10, e1004238.
 929 10.1371/journal.pgen.1004238.

930 Wang, Z., Singhvi, A., Kong, P., and Scott, K. (2004). Taste representations in the
 931 *Drosophila* brain. *Cell* 117, 981-991. 10.1016/j.cell.2004.06.011
 932 S009286740400577X [pii].

933 Watanabe, K., Toba, G., Koganezawa, M., and Yamamoto, D. (2011). *Gr39a*, a highly
 934 diversified gustatory receptor in *Drosophila*, has a role in sexual behavior. *Behav Genet*
 935 41, 746-753. 10.1007/s10519-011-9461-6.

936 Wiley, C., Ellison, C.K., and Shaw, K.L. (2012). Widespread genetic linkage of mating
 937 signals and preferences in the Hawaiian cricket *Laupala*. *Proc Biol Sci* 279, 1203-1209.
 938 10.1098/rspb.2011.1740.

939 Wyatt, T.D. (2014). *Pheromones and animal behavior : chemical signals and signatures*,
 940 Second edition. Edition (Cambridge University Press).

941 Yang, C.-h., Rumpf, S., Xiang, Y., Gordon, M.D., Song, W., Jan, L.Y., and Jan, Y.-N.
 942 (2009). Control of the Postmating Behavioral Switch in *Drosophila* Females by Internal
 943 Sensory Neurons. *Neuron* 61, 519-526.

944 Yew, J.Y., and Chung, H. (2015). Insect pheromones: An overview of function, form,
 945 and discovery. *Prog Lipid Res* 59, 88-105. 10.1016/j.plipres.2015.06.001.

946 Yew, J.Y., Dreisewerd, K., Luftmann, H., Muthing, J., Pohlentz, G., and Kravitz, E.A.
 947 (2009). A new male sex pheromone and novel cuticular cues for chemical
 948 communication in *Drosophila*. *Curr Biol* 19, 1245-1254. S0960-9822(09)01313-X [pii]
 949 10.1016/j.cub.2009.06.037.

950 Zheng, X., Valakh, V., DiAntonio, A., and Ben-Shahar, Y. (2014). Natural antisense
 951 transcripts regulate the neuronal stress response and excitability. *eLife* 3, e01849.
 952 10.7554/eLife.01849.