Short title: Photosystem II-PsbS/Psb27 Activated by Bicarbonate

Bicarbonate Activation of Monomeric Photosystem II-PsbS/Psb27 Complex

**Andrea Fantuzzi^{a1}, Patrycja Haniewicz^{bc1}, Domenica Farci^d, M. Cecilia Loi^e, Keunha Park^a,
Claudia Büchel^f, Matthias Bochtler^{cg}, A. William Rutherford^{a*†}, Dario Piano^{ce†}**

^a*Department of Life Sciences, Imperial College London, London SW7 2AZ, United Kingdom;*

^b*Department of Plant Physiology, Warsaw University of Life Sciences - SGGW, Nowoursynowska Str. 159, 02-776, Warsaw, Poland;*

^c*International Institute of Molecular and Cell Biology, Trojdena 4, 02-109 Warsaw, Poland;*

^d*Department of Chemistry, Umeå University, Linnaeus väg 6, Umeå, Sweden;*

^e*Department of Life and Environmental Sciences, University of Cagliari, V.le S. Ignazio da Laconi 13, 09123 Cagliari, Italy;*

^f*Institute of Molecular Biosciences, University of Frankfurt, Max von Laue Straße 9, 60438 Frankfurt am Main, Germany;*

^g*Polish Academy of Science, Institute of Biochemistry and Biophysics, Pawinskiego 5a, 02-106 Warsaw, Poland.*

¹ these authors contributed equally to this work.

* corresponding author: a.rutherford@imperial.ac.uk

†Senior author

Keywords: photosynthesis, photoprotection, photoassembly, photoactivation, thylakoid membranes, lamellae region, *Nicotiana tabacum*

One sentence summary

A photosystem II monomer with PsbS and Psb27 as additional subunits, is inactive as isolated but activated by bicarbonate, and is attributed to be a late-stage intermediate in photoassembly.

34

35

36

37 **ORCID**

38 AF: 0000-0002-7026-2175

39 AWR: 0000-0002-3124-154X

40 DP: 0000-0003-2980-7572

41 PH: 0000-0002-6298-7184

42 DF: 0000-0002-3691-2699

43 MCL: 0000-0002-2708-4019

44 CL: 0000-0001-6640-7610

45 MB: 0000-0001-7884-4463

46

47

48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68

Abstract

In thylakoid membranes, Photosystem II monomers from the stromal lamellae contain the subunits PsbS and Psb27 (PSII_m-S/27), while Photosystem II monomers from granal regions (PSII_m) lack these subunits. Here, we have isolated and characterised these two types of Photosystem II complexes. The PSII_m-S/27 showed enhanced fluorescence, the near-absence of oxygen evolution, as well as limited and slow electron transfer from Q_A to Q_B compared to the near-normal activities in the granal PSII_m. However, when bicarbonate was added to the PSII_m-S/27, water splitting and Q_A to Q_B electron transfer rates were comparable to those in granal PSII_m. The findings suggest that the binding of PsbS and/or Psb27 inhibits forward electron transfer and lowers the binding affinity for the bicarbonate. This can be rationalized in terms of the recently discovered photoprotection role played by bicarbonate binding via the redox tuning of the Q_A/Q_A^{•-} couple, which controls the charge recombination route, and this limits chlorophyll triplet mediated ¹O₂ formation (Brinkert K et al. (2016) Proc Natl Acad Sci U S A. 113(43):12144-12149). These findings suggest that PSII_m-S/27 is an intermediate in the assembly of PSII in which PsbS and/or Psb27 restrict PSII activity while in transit, by using a bicarbonate-mediated switch and protective mechanism.

69

70

71 **Introduction**

72 Oxygenic photosynthesis is a light-driven biochemical process providing the biosphere with organic
73 carbon, energy, and molecular oxygen (Johnson, 2016). In higher plants, this process takes place in
74 the chloroplast, a specialized organelle consisting of outer and inner membranes forming a network
75 of photosynthetic membranes named thylakoids (Albertsson, 2001; Vothknecht and Westhoff,
76 2002). The protein composition of the different portions of these membranes are distinct, showing
77 segregation of the photosystems. Photosystem II (PSII) is mainly present in the appressed granal
78 regions, while Photosystem I (PSI) is found in the non-appressed regions of the granal margins and
79 in the stromal lamellae (Andersson and Anderson, 1980). Dynamic responses to various
80 environmental factors have been shown to change the ultrastructure and composition of the
81 membranes and photosystems (Ruban and Johnson, 2015; Kirchoff, 2019).

82 In higher plants, the organization of thylakoid membranes also reflects the partition of
83 different kinds of PSII, which are assembled and repaired in the stromal lamellae, while the fully
84 functional PSII complexes are located in the grana (Andersson and Anderson, 1980; Danielsson et
85 al., 2006) (Fig. 1). The small fraction of PSII complexes that are found in stromal lamellae are
86 mainly PSII monomers (PSII_m) and a series of smaller assembly intermediates. In contrast, the
87 grana are dominated by PSII dimers that can form a range of complexes with chlorophyll antenna
88 proteins, including Light Harvesting Complex II (LHCII), forming PSII-LHCII (Danielsson et al.,
89 2006; Watanabe et al., 2009; Haniewicz et al., 2013).

90 PSII, the water/plastoquinone photo-oxidoreductase, uses the energy of light to drive charge
91 separation, oxidise water and reduce plastoquinone. The photochemistry occurs as a one-
92 photon/one-electron reaction four times sequentially to accumulate the 4 oxidising equivalents
93 necessary for water oxidation and oxygen release at the Mn₄CaO₅ active site on the luminal side of
94 PSII (Dau and Zaharieva, 2009). An exchangeable quinone, Q_B, accepts 2 electrons and 2 protons
95 sequentially, before it is released as Q_BH₂ from the stromal side of PSII into the PQ/PQH₂ pool in
96 the membrane (De Causmaecker et al., 2019). The sequential electron transfer steps involve the
97 formation of a stable intermediate, Q_B^{•-}, that can back-react via Q_A^{•-} with the semi-stable, charge
98 accumulation intermediates of the Mn₄CaO₅ cluster (Rutherford et al., 1982). This back reaction
99 occurs via the thermal repopulation of the P^{•+}Pheo^{•-} state, which recombines mainly by a route
100 forming the chlorophyll triplet state ³P₆₈₀ (Rutherford et al., 1981). This triplet state reacts with
101 oxygen to form singlet oxygen ¹O₂ (Krieger-Liszkay, 2005; Rutherford et al., 2012). The ¹O₂
102 generated causes damage to PSII (Krieger-Liszkay, 2005). Other reactive oxygen species, generated

by reductive and oxidative processes in PSII, might also contribute to damage (Pospíšil, 2016).

Repairing the damage is an energetically costly process since proteins and cofactors must be synthesized and replaced to maintain efficient photosynthetic activity (Komenda et al., 2012; Tikkanen and Aro, 2012). This takes place in the stromal lamellae via a stepwise assembly of subcomplexes (Komenda et al., 2012; Nickelsen and Rengstl, 2013; Tomizioli et al., 2014; Puthiyaveetil et al., 2014). A large variety of PSII protein complexes are present in the stroma lamellae, with partially assembled systems co-existing with the PSII in different oligomeric states and different levels of activity (Danielsson et al., 2006; Haniewicz et al., 2013; Tomizioli et al., 2014; Puthiyaveetil et al., 2014). Inactive PSII from the stromal lamellae have been studied for decades (Melis, 1985; Lavergne and Leci 1993), and it was reported that 10-20% of PSII in the chloroplast were inactive and this was due to blocked forward electron transfer and not due to a lack of oxidised plastoquinone (Lavergne and Leci, 1993).

The complexity of the assembly/repair cycle, together with the low abundance of most of the intermediate complexes, means that our understanding of it is still evolving (Komenda et al., 2012; Nickelsen and Rengstl, 2013). Due to the low concentration, instability and intrinsically transient nature of these assembly intermediates, their isolation has required specific strategies: 1) the generation of mutants that lack either specific assembly factors or small PSII subunits, resulting in the accumulation of assembly intermediates (Komenda et al., 2002; Roose and Pakrasi, 2008; Zabert et al., 2021; Huang et al., 2021); and/or 2) tagging one of the PSII subunits to allow isolation of low concentration intermediates by affinity chromatography (Nowaczyk et al., 2006; Liu et al., 2011). Differential fractionation (Danielsson et al., 2006) and more recently differential solubilisation (Fey et al., 2008; Haniewicz et al., 2013) allowed the isolation of some of the subpopulations of PSII. Fractions originating from the lamellae and the granal margins (Haniewicz et al., 2013; Haniewicz et al., 2015) yielded a monomeric PSII containing two additional subunits, PsbS and Psb27, which are absent in functional granal PSII (Haniewicz et al., 2015). While Psb27 has been shown to bind to PSII sub-complexes and to play a role in PSII assembly (Nowaczky et al., 2006; Roose et al., 2008), PsbS has been associated, directly or indirectly, with photoprotection mechanisms via non-photochemical fluorescence quenching (Niyogi and Truong, 2013; Ruban et al., 2016; Bassi and Dall'Osto, 2021).

Since its discovery in 1984 (Ljungberg et al., 1984), the role of PsbS has been controversial (Niyogi and Truong, 2013; Fan et al., 2015; Ware et al., 2015; Ruban et al., 2016; Dall'Osto et al., 2017; Bassi and Dall'Osto, 2021). Despite the availability of a PsbS crystallographic structure (Fan et al., 2015), there is still a debate about its basic components. The presence of chlorophyll, xanthophyll (Correa-Galvis et al., 2016; Gachek et al., 2019) and its role as a luminal pH sensor (Bergantino et al., 2003; Li et al., 2004; Roach and Krieger-Liszkay, 2012; Liguori et al., 2019), are

still debated. The primary role of PsbS is thought to be a protective one, as a key player in some aspects of non-photochemical quenching (NPQ) (Niyogi and Truong, 2013; Ruban et al., 2016; Bassi and Dall'Osto, 2021). Several reports showed the involvement, either direct or indirect, of PsbS, in quenching the excess of energy in free-LHCII complexes and/or in LHCII associated with PSII (Sacharz et al., 2017). A photo-protective role has also been suggested to act via CP47 and the minor external antennas of PSII (Correa-Galvis et al., 2016). However, to date, there is no consensus on a mechanism linking PsbS with NPQ and the xanthophyll cycle that would explain PsbS-mediated photoprotection and its role as a pH sensor.

PsbS has been found to be bound stoichiometrically to purified PSII cores only in samples originating from the lamellae and granal margins of *Tobacco* thylakoids (Haniewicz et al., 2013). Indirect evidence of its presence in PSII dimers and monomers, and its association to the PSII-LHCII complexes in grana have also been reported (Bergantino et al., 2003; Caffari et al., 2009; Correa-Galvis et al., 2016).

Psb27 is present in eukaryotes and prokaryotes, though most of the information relates to the cyanobacterial form. Important differences, such as the eukaryotic Psb27 lacking the covalently bound lipid moiety that is present in the cyanobacteria, raise doubts on whether they have the same location and function. In cyanobacteria, Psb27 is involved in the assembly of the Mn_4CaO_5 cluster and was found to be associated with inactive PSII lacking the three extrinsic proteins PsbO, PsbU and PsbV (Roose et al., 2004; Nowaczyk et al., 2006). It was found to play a significant role during PSII D1 repair, where it was suggested to bind to CP43 and facilitate the assembly of the Mn-cluster by providing greater accessibility and preventing premature association of the other extrinsic proteins (Nowaczyk et al., 2006; Roose and Pakrasi, 2008). Its location close to the CP43 loop E and its allosteric role in weakening the binding of the extrinsic proteins (Liu et al., 2011) was confirmed and clarified in two recent cryo-EM structures (Zabert et al., 2021; Huang et al., 2021). Its role in facilitating the photoactivation of the Mn-cluster was found to be more complex than simply displacing the extrinsic proteins from the apo-PSII (Avramov et al., 2020). In higher plants, this subunit is found to exist in two isoforms, Psb27-1 and Psb27-2 (Chen et al., 2006; Wei et al., 2010). Their specific function is still under investigation, with Psb27-1 found to be required for the efficient repair of photo-damaged PSII (Chen et al., 2006) but also to play a role in the state transition mechanism (Dietzel et al., 2011), while Psb27-2 is suggested to play an important role in the processing of the precursor form of D1 (Wei et al., 2010).

In the present study we have characterized a PSII monomer containing PsbS and Psb27, which was isolated from the stromal lamellae and the grana margins (PSII_m-S/27). We compared this complex with the PSII monomers isolated from the grana stacks (PSII_m). The data indicate an unexpected role for PsbS and/or the Psb27 protecting newly assembled and photoactivated PSII by

inhibiting electron transfer, an inhibition that is reversed by bicarbonate binding.

Results

Association of PsbS and Psb27 to PSII monomers

Two types of purified Photosystem II core complexes, PSII_m and the PSII_m-S/27, were isolated according to the procedures previously described in Haniewicz *et al.* (2013) and in Haniewicz *et al.* (2015), where it was demonstrated that the PSII_m and the PSII_m-S/27 originate from the grana stacks and margins/stromal lamellae, respectively (Fig 1). The two types of monomeric PSII were compared by SDS-PAGE and we confirmed the previous observation that they had similar composition in terms of protein subunits, except for the two additional bands in the PSII_m-S/27 at 20 and 13 kDa (Fig. 2A), attributed to the PsbS and Psb27 subunits, respectively (Haniewicz *et al.*, 2013). These subunits were found to be stoichiometric with the other core subunits of PSII. Small differences in the sub-stoichiometric content of external antennas, which are mainly ascribed to CP26 and CP29, were also observed (see SI Appendix).

Spectroscopic characterisation of PSII_m and the PSII_m-S/27

UV-Vis absorption spectra of the PSII_m and the PSII_m-S/27 were recorded (Fig. 2B). When normalising the spectra at 675 nm, the comparison showed only minimal differences localised between 350 and 550 nm with the PSII_m-S/27 showing slightly higher absorbance. As neither the Psb27 and PsbS have been shown to contain chromophores, these differences are more likely to be due to the small differences in CP26 and CP29 content of the samples.

Room-temperature fluorescence emission spectra were recorded for both PSII samples and showed a single peak at 681 and 682.5 nm for PSII_m-S/27 and PSII_m, respectively (Fig. 2C). In equally concentrated samples, the intensity of the emission peak was much higher (nearly double) for the PSII_m-S/27 sample when compared with the PSII_m (Fig. 2C).

The comparison of the circular dichroism spectra (Fig. 2D) for the PSII_m and the PSII_m-S/27 samples showed differences that can be related to the small changes in the absorption spectra shown in the Fig. 2B. Overall, both spectra resemble a typical PSII spectrum (Alfonso *et al.*, 1994; Kraus *et al.*, 2005), suggesting that no major changes in the position or number of the cofactors is induced by the presence of PsbS and Psb27. However, in the PSII_m-S/27 sample the spectral region between 350 and 550 nm showed changes in peak intensity and position, with 3 minor bands at 427, 475, and 484 nm, respectively. The Q_y band showed the typical PSII double peak in both samples (Alfonso *et al.*, 1994; Kraus *et al.*, 2005). In the PSII_m-S/27 sample, these features were red-shifted by about 2 nm with respect to PSII_m (Fig. 2D).

208

209 **Water oxidation catalytic activity**

210 The enzymatic activities of both PSII_m and PSII_m-S/27 were compared by measuring their oxygen
211 evolution rates. PSII_m showed good activity with rates of $1030 \pm 13 \mu\text{mol O}_2 \text{ mgChl}^{-1} \text{ h}^{-1}$. In
212 contrast, PSII_m-S/27 showed a drastically reduced activity of $52 \pm 5 \mu\text{mol O}_2 \text{ mgChl}^{-1} \text{ h}^{-1}$ (Table 1).

213

214 **Electron transfer from $\text{Q}_\text{A}^{\bullet-}$ to Q_B or $\text{Q}_\text{B}^{\bullet-}$**

215 PSII photochemistry was tested by measuring flash-induced chlorophyll fluorescence and $\text{Q}_\text{A}^{\bullet-}$
216 oxidation kinetics. The illumination with a single-saturating flash of a dark-adapted sample induces
217 the reduction of Q_A to $\text{Q}_\text{A}^{\bullet-}$ in most of the centres, with a resulting increase in the prompt
218 fluorescence yield. Subsequent re-oxidation of $\text{Q}_\text{A}^{\bullet-}$, either by electron transfer to Q_B or $\text{Q}_\text{B}^{\bullet-}$ or by
219 recombination with S_2 , results in the decay of the fluorescence yield (Croft and Wraight, 1983).

220 Figure 3A shows that both minimal fluorescence (F_0) and maximal fluorescence (F_m) yields were
221 higher in the PSII_m-S/27 sample (see SI Appendix). PSII_m was found to be unstable during the
222 measurements at room temperature, therefore experiments were performed at 15°C (Fig. 3B and D).
223 This is likely to slow down some rates when compared to other measurements done at room
224 temperature and when comparing them to the literature. The decay rates of the fluorescence yield
225 are shown in Fig. 3B. PSII_m kinetics are like those expected for functional PSII, while in PSII_m-
226 S/27 the fluorescence decayed ~10 times more slowly, indicating a marked inhibition of forward
227 electron transfer. The kinetics were fitted with three decay phases (Fig. 3B and Table 2) and the
228 origins of the decay phases were assigned to forward and backward electron transfer reactions
229 according to the literature (Vass et al., 1999).

230 The initial fast phase arises from $\text{Q}_\text{A}^{\bullet-}$ forward electron transfer to either Q_B or $\text{Q}_\text{B}^{\bullet-}$. It has a
231 rate of 1.9 ms and an amplitude of 31% for the PSII_m, but a drastically reduced rate and amplitude
232 of 17 ms and 7% amplitude for PSII_m-S/27.

233 The middle phase is often assigned to be electron transfer from $\text{Q}_\text{A}^{\bullet-}$ when the Q_B site is
234 either empty or occupied by $\text{Q}_\text{B}\text{H}_2$ at the time of the flash, and therefore the electron transfer rate is
235 determined by the arrival of PQ into the Q_B site. For PSII_m this phase shows an amplitude of 18%
236 and a half-time of 85 ms, kinetics compatible with the usual assignment of this phase although on
237 the slow side of the range and could indicate a contribution to this phase of charge recombination
238 with TyrZ• in damaged centres. However, for PSII_m-S/27, while the amplitude is like that in PSII_m,
239 the decay is 9-fold slower, with a $t_{1/2} = 770$ ms. This is a very slow value for quinone exchange
240 although slow quinone exchange is a feature of bicarbonate loss from the non-heme iron or
241 bicarbonate replacement by other carboxylic acids (Shevela et al., 2012). This range of fluorescence
242 decay is within those seen for S_2 recombination with $\text{Q}_\text{A}^{\bullet-}$ in PSII monomers (Zimmermann et al.,

2006) but it seems that in this material and at this temperature, this $S_2Q_A^{\bullet-}$ recombination takes place more slowly (see below).

The third and slowest phase is usually attributed to the back reaction of $Q_A^{\bullet-}$ with the S_2 state of the Mn-cluster and thus this phase is seen when forward electron transfer is blocked due to reduction of the pool, modification of the Q_B site or binding of an herbicide in the Q_B site. This phase with a $t_{1/2} = 23$ s seems to be present in about half of the centres in PSII_m, this is in common with other reports from PSII_m in the literature (Table 2). The slower rates compared with the literature values is due to the lower temperature (15°C) used here. The PSII_m-S/27 sample showed even more of this phase, 68%, and a marked slow-down of the half-time to 180s. Thermoluminescence measurements (TL) of PSII_m showed that upon a single saturating flash the PSII_m presents a single peak at 22°C, while PSII_m-S/27 presents a peak at 10°C with a shoulder at 25°C (Fig. 3C and S1). These peaks may be attributed to $S_2Q_A^{\bullet-}$ and $S_2Q_B^{\bullet-}$ recombination respectively based on the typical TL peak temperatures (Rutherford et al 1982). The TL data suggest that the presence of PsbS and Psb27 block the electron transfer from $Q_A^{\bullet-}$ to Q_B . This appears to agree with fluorescence kinetics presented above on the $Q_A^{\bullet-}$ re-oxidation kinetic considering the temperature of the measurement.

Effect of the inhibitor DCMU on the $Q_A^{\bullet-}$ oxidation kinetics

To investigate the possible interference of PsbS and/or Psb27 with Q_B binding, we measured the fluorescence yield relaxation kinetics in the presence of the PSII inhibitor 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU) (Fig. 3D). This inhibitor binds to the Q_B binding site and blocks the electron transfer from $Q_A^{\bullet-}$, leaving the recombination to S_2 as the only possible route for the electrons. The kinetics of $Q_A^{\bullet-}$ oxidation will, therefore, be dominated by the slow phase associated with the recombination with S_2 . Addition of DCMU to both the PSII_m and the PSII_m-S/27 resulted in kinetics with very similar half-times of approximately 30-40 s in 50-60% of the centres, while the remaining 40-50% appear to show longer decaying times. This observation suggests that the presence of PsbS and/or Psb27 does not interfere with the DCMU binding nor the resulting inhibition. These rates are longer than those typically measured in fully functional plant PSII, where it is ~1 second, but this is at least partially explained by the experiments being done at 15°C to preserve the intactness of PSII_m. The slow phases of $S_2Q_A^{\bullet-}$ decay measured with DCMU are similar in PSII_m and PSII_m-S/27, while in the absence of DCMU the slow phase of fluorescence decay was significantly slower in PSII_m-S/27 (Fig. 3B and D).

A notable difference in the kinetics is shown for the PSII_m in the presence of DCMU, where an additional faster phase is present (Fig. 3D) with ~20% amplitude and $t_{1/2} = 50$ ms. This phase could correspond to $Tyr_Z \bullet Q_A^{\bullet-}$ recombination, a reaction reflecting PSII centres that lack the Mn-

cluster. This is consistent with the observed instability of the PSII_m.

Effect of bicarbonate on PSII_m-S/27

Given the recently discovered protective role of bicarbonate and the demonstration that it can be lost under physiological conditions (Brinkert et al., 2016), we investigated the effect of bicarbonate on PSII_m-S/27. These experiments were done at 20°C, a temperature at which the effect of bicarbonate has been characterised and the PSII_m-S/27 was stable. The addition of 5 mM bicarbonate to the PSII_m-S/27 sample resulted in it becoming activated to a level comparable to the functional PSII_m (Fig. 4A). The kinetics of $Q_A^{\bullet-}$ oxidation after addition of bicarbonate showed an acceleration of all three fluorescence decay phases to rates like those measured in the PSII_m and typical of a functional PSII monomer (Zimmermann et al., 2006). In the presence of bicarbonate, the fast phase $t_{1/2}$ decreased from 67 ms to 3 ms, the middle phase decreased from 982 ms to 115 ms, and the slow phase from 43 to 13 s (Table 2). The amplitude of the fast phase also appeared to increase when bicarbonate was present, but this is less certain because of the influence of the poorer fitting of the equivalent phase but with slower kinetic, in the PSII_m-S/27 lacking bicarbonate.

The addition of DCMU to the bicarbonate containing PSII_m-S/27 complex yielded almost the same kinetic profile as seen in the absence of the added 5 mM bicarbonate (Fig. 4B). The fitting of the data showed two main phases, the first with a half-time of approximately 1 s and an amplitude of 70%, consistent with $S_2Q_A^{\bullet-}$ recombination, the second with a half-time of approximately 60 s and an amplitude of 30%. The longer half-life of this second phase is consistent with the oxidation of a relatively stable donor (e.g. from Mn^{2+} , TyrD, the side path donors) giving rise to a long-lived $Q_A^{\bullet-}$ (Nixon et al., 1992; Debus et al., 2000).

Oxygen evolution assays performed in presence of 5 mM bicarbonate showed a ~15-fold increase in the activity of PSII_m-S/27, reaching levels ($795 \pm 8 \mu\text{mol O}_2 \text{ mg Chl}^{-1} \text{ h}^{-1}$) that are ~70% of the values recorded for the PSII_m (Table 1). These results show that the PSII_m-S/27 samples contain a fully functional Mn-cluster in at least 70% of the centers. Both kinetics and oxygen evolution measurements show that approximately 30% of the centers lack catalytic activity and these show a kinetic profile that is consistent with either partial Mn occupancy or Mn-free PSII.

The kinetics of oxidation was studied as a function of bicarbonate concentration. The kinetics accelerated as the bicarbonate concentration was increased. When the fluorescence value at 0.2 s were plotted, and the value obtained prior to bicarbonate addition was subtracted (Fig. 4C), the curve showed hyperbolic behavior saturating at 5 mM. Data fitting with a hyperbolic model for ligand binding yielded an apparent dissociation constant for the bicarbonate of ~ 600 μM .

TL (Fig. 4D) of PSII_m-S/27 with no addition showed a peak centered at 10°C attributed to $S_2Q_A^{\bullet-}$ recombination, while in the presence of bicarbonate the TL intensity increased with a

dominant peak at 27°C, typical of $S_2Q_B^{\bullet-}$ recombination, and an increase in $S_2Q_A^{\bullet-}$ TL at 10°C, which is seen as shoulder. The TL results show that PSII_m-S/27 has inhibited forward electron transfer from $Q_A^{\bullet-}$ and only a low level of luminescence arising from $S_2Q_A^{\bullet-}$. Addition of bicarbonate resulted in the recovery of near-normal behavior with the formation of $S_2Q_B^{\bullet-}$ recombination in most of the centres. In a fraction of centres the bicarbonate did not reconstitute electron transfer to $Q_B^{\bullet-}$ but did result in more $S_2Q_A^{\bullet-}$ recombination.

Finally, it was also observed that the peak intensity and position of the room temperature fluorescence was essentially unaffected by the addition of 5 mM bicarbonate (Fig. S2).

Discussion

Here, we compared two types of PSII monomers, i) those isolated from grana, PSII_m, and ii) those isolated from the stromal lamellae and granal margins, which contain stoichiometric amounts of PsbS and Psb27, PSII_m-S/27 (Haniewicz *et al.*, 2013; Haniewicz *et al.*, 2015). Comparison of the UV-Vis absorption and CD spectra (Figs. 2B and 2D) in the two types of PSII monomers, showed only minimal differences (see SI Appendix). The most significant difference between the two complexes is the near absence of activity in PSII_m-S/27 compared to the high activity in the PSII_m. Both the oxygen evolution rates (Table 1) and the kinetics of $Q_A^{\bullet-}$ oxidation (Fig. 3 and Table 2) were strongly inhibited in PSII_m-S/27. The addition of DCMU to PSII_m-S/27 blocked oxygen evolution and shut down the residual, sub-second fluorescence decay, due the near-complete block of forward electron transfer from $Q_A^{\bullet-}$. The DCMU-treated PSII_m-S/27 showed 70% of the centres with the typical seconds-timescale kinetics of $S_2Q_A^{\bullet-}$ recombination (Fig. 3D and 4B), while the rest of the centres showed much slower rates of $Q_A^{\bullet-}$ decay, presumably due to the electron donation from a more stable electron donor in a fraction of centres. These observations are comparable to those made by Lavergne and Leci (1993) when investigating the fraction of inactive PSII that is normally present in green algal cells. The inactive PSII in algal cells reported earlier (Lavergne and Leci, 1993) could be the algal equivalent of the PSII_m-S/27 described here.

The kinetic characteristics of the impaired $Q_A^{\bullet-}$ oxidation in PSII_m-S/27 indicate heterogeneity and suggest that forward electron transfer, $Q_A^{\bullet-}$ to Q_B and to $Q_B^{\bullet-}$, and the exchange of Q_BH_2 , are all inhibited. These events all involve protonation. The thermoluminescence of PSII_m-S/27 had a low intensity but the peak positions of the residual TL were consistent with inhibition of electron transfer from $Q_A^{\bullet-}$ to Q_B or $Q_B^{\bullet-}$, with the main peak at 10°C, typical of $S_2Q_A^{\bullet-}$ recombination in inhibited centres, and only a very small shoulder at 25°C corresponding to the $S_2Q_B^{\bullet-}$ recombination in a small number of functional centres (Fig. 3C).

The inhibition of $Q_A^{\bullet-}$ oxidation in PSII_m-S/27 cannot be explained by the loss of Q_B (except in a small fraction of the centres), as the addition of bicarbonate activated forward electron

transfer in most of the centres. Similarly, the weak TL intensity cannot be explained by the absence of the Mn cluster, as most of the centres were capable of water splitting when bicarbonate was added. The low luminescence of PSII_m-S/27 prior to the addition of bicarbonate, and despite the presence of the Mn cluster and both quinones, could be due to an increase in the redox potential $Q_A^{\bullet-}$, as occurs upon loss of the bicarbonate from granal PSII dimers (Brinkert et al., 2016). A sufficiently high redox potential is expected to result in the loss of radiative recombination (Rutherford et al., 2012).

The remarkable activation of the seemingly inactive PSII_m-S/27 by millimolar concentrations of bicarbonate was manifest as the appearance of normal rates of forward electron transfer and water oxidation activity in at least 70% of the centres, as monitored by fluorescence kinetics, TL and O₂ evolution. The remaining 30% inactive centres did show a slow-down of the rate of $Q_A^{\bullet-}$ reoxidation in the ~10 s range (Fig. 4A, Table 2). As these kinetics are much slower than typically found for electron transfer to Q_B and $Q_B^{\bullet-}$, this observation could indicate the absence of Q_B in the site in this fraction of the centers. However, as the 10 s phase is eliminated in the presence of DCMU and replaced by a much slower rate, ~100 s (Fig. 4B), this behaviour could indicate an unusually slow rate of electron transfer from $Q_A^{\bullet-}$ to Q_B . This could originate from a situation where the reduction potentials of $Q_A/Q_A^{\bullet-}$ and $Q_B/Q_B^{\bullet-}$ are similar. The reduction potential of $Q_A/Q_A^{\bullet-}$ is reported to shift toward that of $Q_B/Q_B^{\bullet-}$ when the Mn cluster is absent (Johnson et al., 1995), and it could be shifted even further when modified by the binding of the PsbS and Psb27 subunits.

The major difference between the two types of PSII monomers studied here is that the PSII_m-S/27 almost completely lacks activity until activated by the addition of bicarbonate. This difference is presumably due to the binding of PsbS and/or Psb27. It is not clear that this difference is due to one or the other of these polypeptides or to a combination of both. This uncertainty is shared with the recent structural work on other assembly intermediates in cyanobacterial systems (Huang et al., 2021; Zebert et al., 2021). Below we discuss the potential roles of the Psb27 and PsbS.

Psb27 in higher plants is relatively poorly studied, and is known to exist in two isoforms, Psb27-1 and Psb27-2 (Chen et al., 2006; Wei et al., 2010). In the present study, it was not possible to determine which of the two isoforms is bound to PSII_m-S/27. Its suggested functions are associated with responses to photodamage and maturation of D1 in newly synthesised PSII (Chen et al., 2006; Wei et al., 2010). In cyanobacteria, Psb27 is involved in the assembly of the Mn₄CaO₅ cluster, where it is suggested to facilitate photoassembly by allosterically regulating the binding of the extrinsic proteins, PsbO, PsbU, and PsbV (Nowaczyk et al., 2006; Roose and Pakrasi, 2008). In both *Synechocystis* sp PCC 6803 and *Thermosynechococcus elongatus*, Psb27 was found to be

associated with inactive PSII monomers and dimers in which the three extrinsic proteins were absent and either no Mn or sub-stoichiometric amounts of Mn were reported (Roose and Pakrasi, 2004; Nowaczyk et al., 2006; Mamedov et al., 2007; Roose and Pakrasi, 2008). Nevertheless, PSII complexes with Psb27 bound in the presence of either PsbO alone, or the full complement of extrinsic proteins, were found in a range of conditions: 1) as PSII dimers in cold-stressed *T. elongatus* (Grasse et al., 2011), 2) in affinity purified His-tagged Psb27 in *Synechocystis* sp PCC 6803 (Liu et al., 2011), and 3) as PSII monomers in a *psbJ* deletion mutant of *T. elongatus* (Zabert et al., 2021).

Two recent structures of PSII complexes with bound Psb27 (Zabert et al., 2021; Huang et al., 2021), confirmed the previously suggested (Liu et al., 2011; Komenda et al 2012) binding site for Psb27 close to the loop E in CP43. The structures also indicate that binding of Psb27 does not directly interfere with the binding of the extrinsic proteins, in agreement with the range of different Psb27-bound forms of PSII reported in the literature, and therefore reflecting the dynamic process of assembly and repair (Komenda et al., 2002; Roose and Pakrasi, 2004; Nowaczyk et al., 2006; Mamedov et al., 2007; Roose and Pakrasi, 2008; Grasse et al., 2011; Liu et al., 2011; Komenda et al., 2012; Avramov et al., 2020; Huang et al., 2021; Zabert et al., 2021). This agrees with the observation in the present work that in the PSII_m-S/27, the Psb27 is bound, the extrinsic polypeptides are also bound, and the Mn₄CaO₅ cluster is fully assembled in the majority (70%) of the centres. It is not clear if the PSII_m-S/27 is an early-stage intermediate in the repair/assembly process, as suggested in Grasse *et al.* (2011) for a Mn-containing but inactive Psb27-bound PSII dimer in *T. elongatus* (Grasse et al., 2011), or a late stage intermediate, following photoassembly of Mn cluster, prior to joining the fully functional PSII population in the grana. However, the high activity seen in PSII_m-S/27 when the centres were activated by bicarbonate, points to a lack of photodamage and favours its assignment as a late stage post-photoactivation intermediate.

The recent cryo-EM structure of the PSII monomer from *T. elongatus*, showing Psb27 bound to the CP43 and no Mn cluster, also showed significant modifications to the structure around the non-heme iron and the Q_B site. The acceptor side modifications appear to be related to the binding of two other polypeptides, the Psb28 and the Psb34, that cause a conformational change of the D-E loop of the D1 protein that forms a stabilising interaction with Psb28. Part of the C-terminus of CP47 is also displaced by Psb34 forming a stabilising interaction with Psb28. Perhaps the most remarkable result of this conformational change was that the bicarbonate ligand to the non-heme iron was displaced by the carboxylic group of Glu241-D2 (Xiao et al., 2021; Zabert et al., 2021).

Kinetics of Q_A^{•-} oxidation measured for this complex (Zabert et al., 2021), and in other related complexes (Liu et al., 2011; Mamedov et al., 2011), all show large fractions of centres with slow Q_A^{•-} decay (greater than 10 s). This slow decay of Q_A^{•-} has been attributed to a situation in

which forward electron transfer is blocked and $Q_A^{\bullet-}$ is trapped due to a stable electron donor to either $Tyr_Z^{\bullet}$ or P_{D1}^+ , such as a Mn^{2+} or a side-path donor (Nixon et al., 1992; Debus et al., 2000). This situation resembles that observed here in the fraction (~30%) of centres of PSII_m-S/27 lacking the intact Mn cluster. This fraction could represent either centres that have yet to undergo photoactivation, similarly to the intermediates presented in Zabert *et al.* (2021), or centres that have lost the Mn-cluster during isolation from the thylakoid membrane.

Given the clear association of the Psb27 with the electron donor side in the cyanobacterial system, it is tempting to suggest that the plant Psb27 binds in a similar location and plays a similar role/s. It has been known for decades that changes on the electron donor side can have major effects on the electron acceptor side (Johnson et al., 1995). It is thus possible that the binding of the Psb27 has a long-range effect on the electron acceptor side. Indeed, in the crystal structure of the PSII dimer with Psb27, the Psb27 binding site overlaps with the binding site of PsbQ in plants, and PsbQ' in red algae and diatoms (Huang et al., 2021). It has been reported that the PsbQ' binding to PSII shifts the reduction potential of the $Q_A/Q_A^{\bullet-}$ couple to more positive values (Yamada et al., 2018). This suggests the possibility that the binding of Psb27 might result in a shift in the reduction potential of $Q_A/Q_A^{\bullet-}$.

There is much less relevant information for the PsbS, as there are no examples of isolated PSII cores with bound PsbS other than the PSII monomer studied here (Haniewicz *et al.*, 2013; Haniewicz *et al.*, 2015). Previous work on PsbS has been aimed at understanding its role, directly or indirectly associated with non-photochemical quenching (Niyogi and Truong, 2013; Ruban, 2016; Bassi and Dall'Osto, 2021). However, the PSII_m-S/27 was isolated from plants that had not been subjected to light stress leading to the induction of non-photochemical quenching. Furthermore, the spectroscopic characterisation of PSII_m-S/27 (Fig. 2 and 3) shows no quenching of the fluorescence due to PsbS binding. Therefore, a different functional role for PsbS in the PSII_m-S/27 complex should be considered. Cross-linking experiments in thylakoid membranes upon induction of quenching, indicated that monomeric PsbS was associated with CP47 and D2, in addition to its expected association with LHCII (Correa-Galvis et al., 2016). The N-terminal and C-terminal loops, which are both on the stromal side of PsbS, could interact with the electron acceptor side of PSII and affect its function. In the absence of further structural information, it remains possible that the PsbS, in PSII_m-S/27 is located as suggested in the cross-linking experiments (Correa-Galvis et al., 2016).

Now we turn to the effect of bicarbonate. Here, we found that the addition of bicarbonate activated PSII_m-S/27 giving normal rates of $Q_A^{\bullet-}$ oxidation (Fig. 4A and Table 2) and oxygen evolution (Table 1) and near normal TL (4D, Fig. S2). This unexpected bicarbonate-dependent activation showed that the PSII_m-S/27 complex as isolated was essentially intact and capable of

normal function but was “switched off”. The study of the kinetics of $Q_A^{\bullet-}$ oxidation as a function of the bicarbonate concentration showed a hyperbolic dependence, typical of the binding of a ligand to a discrete binding site and an apparent dissociation constant of $\sim 600 \mu\text{M}$. Brinkert *et al.* (2016) showed that the bicarbonate binding site on the non-heme Fe in functional granal PSII dimers has two binding affinities: a high affinity when Q_A is present and a lower affinity when long-lived $Q_A^{\bullet-}$ is present. Based on the thermodynamic relationship between the effect of bicarbonate binding and the reduction potential (E_m) of $Q_A/Q_A^{\bullet-}$, and the literature K_d of $80 \mu\text{M}$ (Stemler and Murphy, 1983) taken as the high affinity value, Brinkert *et al.* (2016) calculated the K_d for the low affinity state to be 1.4 mM . However, they pointed out that based on their observations, the literature K_d value, $80 \mu\text{M}$, appeared to be overestimated and suggested that the actual value was in the low μM range, i.e., that it had a significantly higher affinity. This would mean that the K_d for the low affinity conformation would be smaller than the 1.4 mM , and therefore close to, or smaller than the $600 \mu\text{M}$ K_d measured here for bicarbonate activation of PSII_m-S/27.

Brinkert *et al.* (2016) argued that the increase in the E_m of $Q_A/Q_A^{\bullet-}$ that occurred upon the loss of the bicarbonate, would increase the energy gap between Pheo and Q_A , and this would disfavour the back-reaction route for charge recombination and favour direct charge recombination (Johnson et al, 1995). As described in the introduction, the back-reaction route via the pheophytin leads to chlorophyll triplet formation and thence to $^1\text{O}_2$ -mediated photodamage, while the direct recombination of the slower $P^+Q_A^{\bullet-}$ radical pair is considered safe (Rutherford et al., 1982; Johnson et al., 1995). The bicarbonate-mediated redox tuning of Q_A was thus considered to be a regulatory and protective mechanism (Brinkert et al., 2016). It seems quite likely that a similar protective mechanism exists in the PSII_m-S/27, as protection is needed during and after synthesis or repair.

The increased stability of the PSII_m-S/27 conferred by the presence of PsbS and/or Psb27 was manifest by its resilience to long incubations and photochemical measurements at room temperature. This resilience contrasted with the fragility of the PSII_m, which could not be studied even for short periods at room temperature without loss of activity.

A bicarbonate-controlled, redox-tuning-based, protective mechanism in PSII_m-S/27 would appear to be beneficial for this complex. A non-functional PSII, like PSII_m-S/27 with an intact electron donor-side but with inhibited forward electron transfer and a low-potential Q_A (Johnson et al., 1995), would be hypersensitive to backreaction-associated photodamage, just as occurs in herbicide-treated PSII (Rutherford and Krieger-Liszky, 2001). It is known that before photoactivation of water oxidation in PSII, the E_m of $Q_A/Q_A^{\bullet-}$ is high and thus PSII is photo-protected, and at some point during the photoassembly of the Mn complex, the E_m is shifted to a functional, low potential value (Johnson et al., 1995). If the PSII_m-S/27 is a late-stage intermediate of photoactivation, then the present work would indicate that the donor-side-induced switching of

the E_m of $Q_A/Q_A^{\bullet-}$ is overridden by modifications to the acceptor side that maintain Q_A in a safe, high-potential form, until the fully assembled PSII is delivered into the granal stack and is dimerised. The binding of the PsbS and Psb27, can be considered as exerting conformational restrictions to the assembled but non-functional PSII, protecting it until it is in the right place and dimerised. At that point, presumably the PsbS and Psb27 dissociate, allowing the PSII to adopt its functional conformation, allowing the high affinity bicarbonate site to form. The bicarbonate duly binds, shifting the E_m of $Q_A/Q_A^{\bullet-}$ to low potential and allowing optimal function. When considering the measured dissociation constant for bicarbonate in PSII_m-S/27 (600 μ M), it seems clear that the physiological concentration of CO_2 could control, to some extent, the activity of this complex. At pH 8.0 the equilibrium concentration of bicarbonate will be sufficiently high to allow more than 50% of this complex to show normal forward electron transfer kinetics.

Another less likely explanation for the lower activity of PSII_m-S/27 is that it represents an early-stage intermediate in the repair cycle. In this model, photodamage would be manifest as an electron acceptor side restriction, and the binding of Psb27 and PsbS and the higher potential form of Q_A , due to the loss of bicarbonate, would protect the system from photodamage during its transit to the repair site in the stromal lamellae.

In the literature there are several examples of other PSII subunits which seem to exhibit similar or related effects to those described here for PsbS and Psb27, though none of them are as marked as observed in this work. These are listed in the supplementary information (See SI Appendix). This evidence in the literature and the present work point to a broader picture in which there is an interplay between the binding of small subunits to PSII with the resulting conformational effects, and the binding of bicarbonate with the resulting redox tuning effects. This interplay appears to control electron transfer rates and thermodynamic equilibria between the different quinones in all their forms, thereby regulating and safeguarding PSII during the diverse steps of its life cycle.

Conclusions

We show that the PSII monomer from the stromal lamellae/stromal margins, which has PsbS and Psb27 bound to it, has very low activity but is activated upon binding bicarbonate. These findings indicate that PSII_m-S/27 is a switched-off state that is protected from photodamage presumably due to the changes induced by the binding of the two extra polypeptides. The nature of the protection mechanism appears to be complex, not least because of sample heterogeneity, but the dominant one in the PSII_m-S/27 sample appears to involve a modification of the Q_B site, affecting its proton-coupled electron transfer properties and its exchange with the PQ pool. Another important feature of this complex is the diminished affinity for bicarbonate and the significant positive redox shift of the

$Q_A/Q_A^{\bullet-}$ reduction potential that appears to be present when the bicarbonate is not bound. Just such a shift occurs in standard granal PSII dimers when bicarbonate is released upon $Q_A^{\bullet-}$ accumulation (Brinkert et al., 2016). The redox shift protects against the well-characterised photodamage arising from chlorophyll triplet-mediated, singlet-oxygen generation (Johnson et al., 1995). This kind of protection is expected to be important in a near-intact PSII that is switched-off in transit, either after photoactivation or prior to repair.

Material and Methods

Growth and cultivation of tobacco plants

Transplastomic plants of *Nicotiana tabacum*, which have a hexa-histidine tag sequence at the 5' end of the gene coding for the PsbE subunit were used for this work (Fey et al., 2008). Plants were kept at a constant temperature of 25°C, at 50% relative humidity, and grown for 10–12 weeks under a light regime of 12 h/day, with a light intensity of 150–200 $\mu\text{mol photons s}^{-1} \text{ m}^{-2}$.

Thylakoids preparation and PSII core solubilizations

Thylakoid membranes and PSII cores were prepared as previously reported (Haniewicz et al., 2013; Haniewicz et al., 2015), with only minimal modifications in the solubilization step. Briefly, PSII core complexes retaining the subunits PsbS and Psb27 were obtained from thylakoid membranes solubilized for 5 min at 4°C at a final chlorophyll concentration of 3 mg/mL. After solubilization, the unsolubilized fraction was separated by centrifugation at 35000 g for 10 min at 4°C. The unsolubilized fraction underwent a second solubilization step to isolate the PSII core complexes lacking PsbS and Psb27. This second solubilization took place for 15 min at 4°C at a final chlorophyll concentration of 1 mg/mL. Also, after this second solubilization, the unsolubilized fraction was separated by centrifugation at 35000 g for 10 min at 4°C. In both cases, the solubilization was carried out in the dark, adjusting the chlorophyll concentration with Grinding buffer (20 mM MES–NaOH, pH 6.5, 5 mM MgCl_2) and using 20 mM (1.02 %) n-Dodecyl- β -D-maltoside (β -DDM).

PSII core complexes isolation

Photosystem II samples were prepared using Ni-affinity chromatography and a subsequent step of size exclusion chromatography as reported in Haniewicz et al. (2013) for PSII complexes retaining the subunit PsbS, and according to Haniewicz et al. (2015) for PSII complex lacking the subunit PsbS. For the size exclusion chromatography step, buffer containing 20 mM MES–NaOH, pH 6.5, 5

mM MgCl₂ and with 0.02% (~0.39 mM) β -DDM (buffer A). Previously a slightly higher detergent content was used of 0.03% (~ 0.59 mM) (Haniewicz *et al.*, 2013; Haniewicz *et al.*, 2015). In these studies, all chromatography columns were subjected to the ReGenFix procedure (<https://www.regenfix.eu/>) for regeneration and calibration prior use.

Polyacrylamide gel electrophoresis

Denaturing Sodium Dodecyl Sulphate-Polyacrylamide Gel Electrophoresis (SDS-PAGE) consisted in 10% (w/v) separating polyacrylamide/urea gels with 4% (w/v) stacking gels (Piano *et al.*, 2010; Collu *et al.*, 2017). Samples were denatured with Rotiload (Roth) at room temperature before loading, and, after electrophoresis gels were stained with Coomassie brilliant blue G250.

Absorption, CD spectroscopy and chlorophyll determination

The protein content of thylakoids was assessed through three independent measurements based on the concentrations of Chl *a* and Chl *b*. The absorption of chlorophylls extracted in 80% (v/v) acetone, in a dilution factor of 200 or 500, was measured with a Pharmacia Biotech Ultrospec 4000 spectrophotometer, and their relative concentrations were calculated according to Porra *et al.* (1989). CD spectra were the average of three accumulations recorded at a sensitivity of 100 mdeg and a scan speed of 100 nm/min using a CD spectrometer JASCO J-810. Absorption and CD spectra were recorded at room temperature in the range of 370-750 nm, with an optical path length of 1 cm and a band-pass of 2 nm. Spectra were recorded on an absorption Ultra Micro quartz cell with 10 mm light path (Hellma Analytics). In all cases, measurements were performed in a range between 0.01 and 0.2 mg/mL Chls, and samples were diluted in buffer A.

Fluorescence spectroscopy and kinetics

Emission and excitation spectra were recorded on a Jasco FP-8200 spectrofluorometer at 4°C in 0.1 nm steps and 3 nm band-pass. Spectra were corrected for the photomultiplier sensitivity using a calibrated lamp spectrum. Emission spectra in the range of 600-750 nm were recorded using the main absorption bands as excitation wavelength (437 nm in Fig 2). Fluorescence spectra were recorded on a fluorescence Ultra Micro quartz cell with 3 mm light path (Hellma Analytics). The flash-induced increase and the subsequent decay of chlorophyll fluorescence yield and the values of F₀, F_m and F_v were measured with a fast double modulation fluorimeter (FL 3000, PSI, Czech Republic). The sample concentration was 5 µg Chl/ml in buffer A. Samples were subjected to a pre-illumination in room light for 10 seconds followed by a period of 5 to 10 minutes of dark adaptation.

Multicomponent deconvolution of the measured curves was done by using a fitting function with three components based on the widely used model of the two-electron gate (Croft and Wraight, 1983; Vass et al., 1999). The fast and middle phases were simulated with exponential components. However, slow recombination of $Q_A^{\bullet-}$ via charge recombination has been shown to obey hyperbolic kinetics corresponding to an apparently second order process (Bennoun, 1994), most probably the result of stretched exponentials indicative of inhomogeneity in this time-range. Therefore, the data were fitted with a linear combination of two exponentials and a hyperbolic component, where $F(t)$ is the variable fluorescence yield, F_0 is the basic fluorescence level before the flash, A_1 – A_3 are the amplitudes, T_1 – T_3 are the time-constants, from which the half-lives were calculated via $t_{(1/2)} = \ln 2 \cdot T$ for the exponential components, and $t_{(1/2)}$ is the T for the hyperbolic component.

$$F(t) - F_0 = A_1 \exp(-t/T_1) + A_2 \exp(-t/T_2) + A_3 / (1 + t/T_3) \quad \text{Eq.1}$$

When the herbicide 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU) was added, 10 μ M of an ethanolic solution were added in the dark to 1 mL of protein solution in buffer A, prior to the 5 minutes dark adaptation step.

The kinetics measured as a function of increasing concentrations of bicarbonate (0.5, 1, 5, 10 mM) were fitted with a hyperbolic curve (Eq. 2) from which the apparent dissociation constant ($K_{d \text{ app}}$) was calculated.

$$\Delta \text{Fluorescence at } 0.2s = f_{\max} / (K_{d \text{ app}} + [\text{bicarbonate}]) \quad \text{Eq2}$$

where $\Delta \text{Fluorescence at } 0.2 \text{ s}$ is the difference in fluorescence value at 0.2 s subtracted of the value before any addition of bicarbonate, and $f_{\max}$ is the fluorescence difference value when the binding site is fully occupied. The pH was monitored upon addition of bicarbonate to make sure that no shifts in pH occurred.

Oxygen evolution

The oxygen evolution was measured with a Clark-type electrode (Hansatech, England) at 20°C, with 1 mM 2,6-dichloro-p-benzoquinone and 1 mM ferricyanide as electron acceptors in the reaction mixture. Measurements were carried on samples with a chlorophyll concentration of 1 mg/mL diluted 20 times in buffer A to a final concentration of 50 μ g Chl/mL. Three independent measurements were done on the same preparation to test the activity. Reactions were started with illumination from a white light source (400-700 nm) with a Photosynthetically Active Photon Flux Density (PPFD) of 700-800 μ moles of photons $\text{m}^{-2} \text{ s}^{-1}$. For the effect of bicarbonate on PSII_m-S/27

625 samples, 5 mM NaHCO₃ was added to the reaction mixture.

626

627 **Thermoluminescence**

628 Thermoluminescence (TL) was measured with a lab-built apparatus, essentially as described in
 629 (Ducruet and Vass, 2009) but using a GaAsP photomultiplier H10769A-50 (Hamamatsu). Samples
 630 were pre-illuminated with room light ($\sim 20 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 10 s, dark-adapted for 5 to 10 min and
 631 then cooled to 5°C. After 2 min, samples were excited with a single turnover saturating flash.
 632 Finally, samples were rapidly cooled to -15°C and luminescence was recorded with a $20^\circ\text{C min}^{-1}$
 633 heating rate. The sample concentration was 5 $\mu\text{g Chl/ml}$ in buffer A.

634

635 **Acknowledgments**

636 This work was carried out with the support of the program Homing Plus (Foundation for Polish
 637 Science) grant No: 2012-6/10 co-financed by the European Union under the European Regional
 638 Development Funds (to D.P.), the PRELUDIUM programme (National Science Centre) grant
 639 number DEC-2012/05/05/N/NZ1/01922 (to P.H.), Biotechnology and Biological Sciences Research
 640 Council (BBSRC) Grants BB/K002627/1 and BB/R00921X (to A.W.R.). We thank J-M Ducruet
 641 for help with setting up the thermoluminescence.

642 **Author contributions:** A.F., A.W.R., D.P., and C.B. designed research; A.F., K.P., D.P., P.H., and
 643 D.F. performed research; D.P. and D.F. contributed new reagents/analytic tools; A.F., K.P.,
 644 A.W.R., C.B., M.B., and D.P. analyzed data; A.F. A.W.R., D.P., P.H., D.F., C.B., M.B., and M.C.L.
 645 wrote the paper.

646

Tables

Table 1: Rates of oxygen evolution for PSII_m and PSII_m-S/27 with or without added bicarbonate. Data represent mean $\pm$ SD, n = 3.

PSII type	Added [NaHCO ₃]	Oxygen evolution rates (μ mol O ₂ / mg Chls·h)
PSII _m	0	1030 $\pm$ 13
PSII _m -S/27	0	52 $\pm$ 5
	5 mM	795 $\pm$ 8

Table 2: Kinetic parameters of flash-induced chlorophyll fluorescence decay in PSII_m, PSII_m-S/27 samples in absence or presence of additional bicarbonate in solution. The values of the kinetic half-lives (t) and the amplitudes of each phase are compared to the literature values for PSII monomer from *T. elongatus* (Zimmermann et al., 2006)(*). Data represent mean $\pm$ SD, n = 3.

PSII type	Added NaHCO ₃ [mM]	T °C	Fast phase		Middle phase		Slow phase	
			t (ms)	Amp (%)	t (ms)	Amp (%)	t (s)	Amp (%)
PSII _m	0	15	1.9 $\pm$ 0.3	31 $\pm$ 2	85 $\pm$ 10	18 $\pm$ 3	23 $\pm$ 4	51 $\pm$ 1
PSII _m -S/27	0	15	17 $\pm$ 3	7 $\pm$ 3	770 $\pm$ 40	25 $\pm$ 3	180 $\pm$ 35	68 $\pm$ 2
	0	20	67 $\pm$ 10	21 $\pm$ 4	982 $\pm$ 130	34 $\pm$ 2	41 $\pm$ 5	45 $\pm$ 3
	5	20	3 $\pm$ 0.5	27 $\pm$ 2	115 $\pm$ 21	25 $\pm$ 2	13 $\pm$ 2	48 $\pm$ 2
PSII monomer ^(*)	0	20	3.7 $\pm$ 0.5	30 $\pm$ 7	37 $\pm$ 15	11 $\pm$ 2	9.9 $\pm$ 0.7	59 $\pm$ 5

Figure Legends

Figure 1: The location within the thylakoid membrane of the protein complexes investigated in this work. PSII_m-S/27 is in the lamella membranes (red inset) and suggested to be an intermediate in the assembly and repair pathway, probably after photoactivation. This complex migrates (as indicated by the black arrows) to the grana membranes (blue inset) where upon dissociation of PsbS and Psb27, it forms the active monomer, PSII_m. The PSII_m will then form active dimers and bind antenna proteins to form fully functional complexes.

671 **Figure 2:** Comparisons of the PSII_m and PSII_m-S/27. A) Coomassie Blue Stained SDS-PAGE of
 672 the PSII_m (A) and PSII_m-S/27 (B). The additional presence of a ~ 22 kDa (PsbS) and a ~ 14 kDa
 673 (Psb27) bands in the PSII_m-S/27 sample are indicated together with the main PSII subunits. The
 674 lanes labelled as M indicate the molecular weight markers. B) UV-Vis absorption spectra of PSII_m
 675 (black line) and PSII_m-S/27 (red line). Spectra were taken at 20°C in buffer A and were normalized
 676 at 675 nm. C) Fluorescence emission spectra of PSII_m (black line) and PSII_m-S/27 (red line).
 677 Spectra, recorded at 4°C in buffer A with excitation at 437 nm. D) Circular dichroism spectra at
 678 20°C in buffer A of PSII_m-S/27 (red line) and PSII_m (black line).

680 **Figure 3:** A) Fluorescence relaxation kinetics data presented without normalization to show the
 681 values of F_0 and F_m for PSII_m (squares symbols, black line) and PSII_m-S/27 (circles symbols, red
 682 line). Both F_0 and F_m were found to be higher in the sample with bound PsbS and Psb27. B)
 683 Fluorescence relaxation kinetics of PSII_m (squares, black line) and PSII_m-S/27 (circles, red line),
 684 measured at 15°C in buffer A. Data were normalized using the initial amplitudes. Fittings were
 685 carried out with equation 1 (see methods). C) Thermoluminescence measurements for PSII_m (black
 686 squares and line) and PSII_m-S/27 (red circles and line), in buffer A. Single saturating flash was
 687 given at 5°C followed by rapid cooling to -5°C. Scan rate was 0.5°C/s. D) Fluorescence relaxation
 688 kinetics upon a single saturating flash for the PSII_m (squares, black line) and PSII_m-S/27 (circles,
 689 red line), measured at 15°C in buffer A in the presence of 10 µM DCMU. Data were normalized
 690 using the initial amplitudes. Fittings were carried out with equation 1 (see methods).

692 **Figure 4:** A) Fluorescence relaxation kinetics for PSII_m-S/27, measured at 20 °C in buffer A,
 693 without (circles symbols, red line) and with (triangles symbols, blue line) added 5 mM bicarbonate.
 694 Data were normalized using the initial amplitudes. Fittings were carried out with equation 1 (see
 695 methods). Error bars represent the standard errors calculated from 4 independent measurements. B)
 696 Fluorescence relaxation kinetics for PSII_m-S/27, measured at 20 °C in buffer A and 10 µM DCMU,
 697 without (circles symbols, red line) and with (triangles symbols, blue line) added 5 mM bicarbonate.
 698 Data were normalized using the initial amplitudes. Fittings were carried out with equation 1 (see
 699 methods). Error bars represent the standard errors calculated from 4 independent measurements. C)
 700 Plot of the fluorescence intensity at 0.2 s (triangles symbols) from the fluorescence relaxation
 701 kinetics, at 20 °C in buffer A, from different samples to which increasing concentrations of
 702 bicarbonate were added. The data were fitted with the hyperbole in equation 2 (see methods) (blue
 703 line). Error bars represent the standard errors calculated from 4 independent measurements; D)
 704 Thermoluminescence measurements of PSII_m-S/27 in buffer A, in the absence (red circles) and

presence (blue triangles) of added 5 mM bicarbonate. Single saturating flash was given at 5 °C followed by rapid cooling to -5 °C. Scan rate was 0.5 °C/s.

References

- Albertsson P-A** (2001) A quantitative model of the domain structure of the photosynthetic membrane. *Trends in Plant Sci.* **6**: 349-354
- Alfonso M, Montoya G, Cases R, Rodríguez R, Picorel R** (1994) Core antenna complexes, CP43 and CP47, of higher plant photosystem II. Spectral properties, pigment stoichiometry, and amino acid composition. *Biochem.* **33**: 10494-10500
- Andersson B, Anderson J-M** (1980) Lateral heterogeneity in the distribution of chlorophyll-protein complexes of the thylakoid membranes of spinach chloroplasts. *Biochim. Biophys. Acta* **593**: 427-440
- Avramov AP, Hwang HJ, Burnap RL** (2020) The role of Ca²⁺ and protein scaffolding in the formation of nature's water oxidizing complex. *Proc. Natl. Ac. Sci. USA* **117**: 28036-28045
- Bassi R, Dall'Osto L** (2021) Dissipation of light energy absorbed in excess: the molecular mechanisms. *Annu. Rev. Plant Biol.* **72**: 47-76
- Bennoun P** (1994) Chlororespiration revisited: Mitochondrial-plastid interactions in *Chlamydomonas*. *Biochim. Biophys. Acta* **1186**: 59-66
- Bergantino E, Segalla A, Brunetta A, Teardo E, Rigoni F, Giacometti GM, Szabò I** (2003) Light- and pH-dependent structural changes in the PsbS subunit of photosystem II. *Proc. Natl. Acad. Sci. USA* **100**: 15265–15270
- Brinkert K, De Causmaecker S, Krieger-Liszkay A, Fantuzzi A, Rutherford AW** (2016) Bicarbonate-induced redox tuning in Photosystem II for regulation and protection. *Proc. Natl. Acad. Sci. U S A.* **113**: 12144-12149
- Caffarri S, Kouril R, Kereiche S, Boekema EJ, Croce R** (2009) Functional architecture of higher plant photosystem II supercomplexes. *EMBO J.* **28**: 3052-3063
- Chen H, Zhang D, Guo J, Wu H, Jin M, Lu Q, Lu C, Zhang L** (2006) A Psb27 homologue in *Arabidopsis thaliana* is required for efficient repair of photodamaged photosystem II. *Plant Mol. Biol.* **61**: 567-575
- Collu G, Farci D, Esposito F, Pintus F, Kirkpatrick J, Piano D** (2017) New insights into the operative network of FaEO, an enone oxidoreductase from *Fragaria x ananassa* Duch. *Plant Mol. Biol.* **94**:125-136
- Correa-Galvis V, Poschmann G, Melzer M, Stühler K, Jahns P** (2016) PsbS interactions involved in the activation of energy dissipation in *Arabidopsis*. *Nat. Plants.* **2**: 15225

740 **Crofts AR, Wraight CA** (1983) The electrochemical domain of photosynthesis. *Biochim. Biophys.*
741 *Acta* **726**: 149–185

742 **Dall'Osto L, Cazzaniga S, Bressan M, Paleček D, Židek K, Niyogi KK, Fleming GR,**
743 **Zigmantas D, Bassi R** (2017) Two mechanisms for dissipation of excess light in monomeric and
744 trimeric light-harvesting complexes. *Nat. Plants.* **3**:17033

745 **Danielsson R, Suorsa M, Paakkarinen V, Albertsson P-A, Styring S, Aro E-M, Mamedov F**
746 (2006) Dimeric and monomeric organization of photosystem II. *J. Biol. Chem.* **281**: 14241–14249

747 **Dau H, Zaharieva I** (2009) Principles, efficiency, and blueprint character of solar-energy
748 conversion in photosynthetic water oxidation. *Acc Chem Res.* **42**: 1861-1870

749 **De Causmaecker S, Douglass JS, Fantuzzi A, Nitschke N, Rutherford AW** (2019) Energetics of
750 the exchangeable quinone, Q_B, in Photosystem II. *Proc. Natl. Acad. Sci.* **116**: 9458-9463

751 **Debus RJ, Campbell KA, Pham DP, Hays A-MA, Britt RD** (2000). Glutamine 189 of the D1
752 polypeptide modulates the magnetic and redox properties of the manganese cluster and tyrosine Yz
753 in Photosystem II. *Biochem.* **39**: 6275-6287

754 **Dietzel L, Bräutigam K, Steiner S, Schöffler K, Lepetit B, Grimm B, Schöttler MA,**
755 **Pfannschmidt T** (2011) Photosystem II supercomplex remodelling serves as an entry mechanism
756 for state transitions in *Arabidopsis*. *Plant Cell.* **23**: 2964-2977

757 **Ducruet J, Vass I** (2009) Thermoluminescence: experimental. *Photosynth. Res.* **101**: 195–204

758 **Fan M, Li M, Liu Z, Cao P, Pan X, Zhang H, Zhao X, Zhang J, Chang W** (2015) Crystal
759 structures of the PsbS protein essential for photoprotection in plants. *Nat. Struct. Mol. Biol.* **22**:
760 729-735

761 **Fey H, Piano D, Horn R, Fischer D, Schmidt M, Ruf S, Schröder WP, Bock R, Büchel C**
762 (2015) Isolation of highly active photosystem II core complexes with a His-tagged Cyt *b*₅₅₉ subunit
763 from transplastomic tobacco plants. *Biochim. Biophys. Acta* **1777**: 1501–1509

764 **Gachek DA, Holleboom CP, Tietz S, Kirchhoff H, Walla PJ** (2019) PsbS-dependent and -
765 independent mechanisms regulate carotenoid-chlorophyll energy coupling in grana thylakoids.
766 *FEBS Lett.* **593**: 3190-3197

767 **Grasse N, Mamedov F, Becker K, Styring S, Rögner M, Nowaczyk MM** (2011) Role of novel
768 dimeric Photosystem II (PSII)-Psb27 protein complex in PSII repair. *J. of Biol. Chem.* **286**: 29548-
769 29555

770 **Haniewicz P, De Sanctis D, Büchel C, Schröder W P, Loi M C, Kieselbach T, Bochtler M,**
771 **Piano D** (2013) Isolation of monomeric photosystem II that retains the subunit PsbS. *Photosynth*
772 *Res.* **118**: 199-207

773 **Haniewicz P, Floris D, Farci D, Kirkpatrick J, Loi MC, Büchel C, Bochtler M, Piano D** (2008)
774 Isolation of Plant Photosystem II Complexes by Fractional Solubilization. *Front. Plant Sci.* **6**:1100

775 **Huang G, Xiao Y, Pi X, Zhao L, Zhu Q, Wang W, Kuang T, Han G, Sui SF, Shen JR** (2021)
776 Structural Insights into a dimeric Psb27-photosystem II complex from a cyanobacterium
777 *Thermosynechococcus vulcanus*. Proc. Natl. Acc. Sci. USA **118**: e2018053118

778 **Johnson GN, Rutherford AW, Krieger A** (1995) A change in the midpoint potential of the quinone
779 Q(A) in Photosystem II associated with photoactivation of oxygen evolution. Biochim. Biophys.
780 Acta. **1229**: 202-207

781 **Johnson M P** (2016) Photosynthesis. Essays Biochem. **60**: 255–273

782 **Kirchhoff H** (2019) Chloroplast ultrastructure in plants. New Phytologist **223**: 565-574

783 **Komenda J, Lupinkova L, Kopecky J** (2002) Absence of the psbH gene product destabilizes
784 Photosystem II complex and bicarbonate binding on its acceptor side in *Synechocystis* PCC 6803.
785 Eur. J. Biochem. **269**: 610-619

786 **Komenda J, Sobotka R, Nixon PJ** (2012) Assembling and maintaining the Photosystem II
787 complex in chloroplasts and cyanobacteria. Curr. Op. in Plant Biol. **15**: 245-251

788 **Krausz E, Hughes JL, Smith P, Pace R, Årsköld SP** (2005) Oxygen-evolving Photosystem II
789 core complexes: a new paradigm based on the spectral identification of the charge-separating state,
790 the primary acceptor and assignment of low-temperature fluorescence. Photochem. Photobiol. Sci.
791 **4**: 744-753

792 **Krieger-Liszkay A** (2005) Singlet oxygen production in photosynthesis. J. of Exp. Bot. **56**: 337–
793 346

794 **Lavergne J, Leci E** (1993) Properties of inactive Photosystem II centres. Photosynth. Res. **35**: 323-
795 343

796 **Li XP, Gilmore AM, Caffarri S, Bassi R, Golan T, Kramer D, Niyogi KK** (2004) Regulation of
797 photosynthetic light harvesting involves intrathylakoid lumen pH sensing by the PsbS protein. J
798 Biol Chem. **279**: 22866-22874

799 **Liguori N, Campos SRR, Baptista AM, Croce R** (2019) Molecular Anatomy of Plant
800 Photoprotective Switches: The Sensitivity of PsbS to the Environment, Residue by Residue. J Phys.
801 Chem. Lett. **10**:1737-1742

802 **Liu H, Roose JL, Cameron JC, Pakrasi HB** (2011) A genetically tagged Psb27 protein allows
803 purification of two consecutive Photosystem II (PSII) assembly intermediates in *Synechocystis*
804 6803, a cyanobacterium. J. of Biol. Chem. **286**: 24865-24871

805 **Ljungberg U, Åkerlund HE, Larsson C, Andersson B** (1984) Identification of polypeptides
806 associated with the 23 and 33 kDa proteins of photosynthetic oxygen evolution. Biochim. Biophys.
807 Acta. **767**: 145-152

808 **Mamedov F, Nowaczyk MM, Thapper A, Rogner M, Styring S** (2007) Functional
809 characterization of monomeric Photosystem II core preparations from *Thermosynechococcus*

810 *elongatus* with or without the Psb27 protein. Biochem. **46**: 5542-5551

811 **Melis A** (1985) Functional properties of Photosystem II_β in spinach chloroplasts. Biochim Biophys
812 Acta. **808**: 334-342

813 **Nickelsen J, Rengstl B** (2013) Photosystem II assembly: from cyanobacteria to plants. Annu. Rev.
814 Plant Biol. **64**: 609-635

815 **Nixon PN, Trost JT, Diner BA** (1992) Role of carboxyl terminus of polypeptide D1 in the
816 assembly of a functional water-oxidizing manganese cluster in Photosystem II of the
817 cyanobacterium *Synechocystis* sp. PCC 6803: assembly requires a free carboxyl group at C-terminal
818 position 344. Biochem. **31**:10859-10871

819 **Niyogi KK, Truong TB** (2013) Evolution of flexible non-photochemical quenching mechanisms
820 that regulate light harvesting in oxygenic photosynthesis. Curr. Op. Plant Biol. **16**: 307-314

821 **Nowaczyk MM, Hebel R, Schlodder E, Meyer HE, Warscheid B, Rögner M** (2006) Psb27, a
822 cyanobacterial lipoprotein, is involved in the repair cycle of photosystem II. Plant Cell. **18**: 3121-
823 3131

824 **Piano D, El Alaoui S, Korza HJ, Filipek R, Sabala I, Haniewicz P, Buechel C, De Sanctis D,**
825 **Bochtler M** (2010) Crystallization of the photosystem II core complex and its chlorophyll binding
826 subunit CP43 from transplastomic plants of *Nicotiana tabacum*. Photosynth. Res. **106**: 221–226

827 **Porra RJ, Thompson WA, Kriedmann PE** (1989) Determination of accurate extinction
828 coefficients and simultaneous equations for assaying chlorophylls *a* and *b* with four different
829 solvents: verifications of the concentration of chlorophyll standards by atomic absorption
830 spectroscopy. Biochim. Biophys. Acta **975**: 384–394

831 **Pospíšil P** (2016) Production of Reactive Oxygen Species by Photosystem II as a Response to Light
832 and Temperature Stress. Front Plant Sci. **7**: 1950

833 **Puthiyaveetil S, Tsabari O, Lowry T, Lenhart S, Lewis RR, Reich Z, Kirchhoff H** (2014)
834 Compartmentalization of the protein repair machinery in photosynthetic membrane. Proc. Natl.
835 Acad. Sci. USA **111**: 15839-15844

836 **Roach T, Krieger-Liszkay A** (2012) The role of the PsbS protein in the protection of photosystems
837 I and II against high light in *Arabidopsis thaliana*. Biochim. Biophys. Acta. **1817**: 2158-2165

838 **Roose JL, Pakrasi HB** (2004) Evidence that D1 processing is required for manganese binding and
839 extrinsic protein assembly into Photosystem II. J. Biol. Chem. **279**: 45417-45422

840 **Roose JL, Pakrasi HB** (2008) The Psb27 protein facilitates manganese cluster assembly in
841 photosystem II. J. Biol. Chem. **283**: 4044-4050

842 **Ruban A V, Johnson M P** (2015) Visualizing the dynamic structure of the plant photosynthetic
843 membrane. Nat. Plants **1**: 15161

844 **Ruban AV** (2016) Nonphotochemical chlorophyll fluorescence quenching: mechanism and

effectiveness in protecting plants from photodamage. *Plant Physiol.* **170**: 1903-1916

Rutherford AW, Crofts AR, Inoue Y (1982) Thermoluminescence as a probe of Photosystem II photochemistry: the origin of the flash induced glow peaks. *Biochim. Biophys. Acta* **682**: 457-465.

Rutherford AW, Krieger-Liszkay A (2001) Herbicide-induced oxidative stress in Photosystem II. *Trends In Biochem. Sc.* **26**: 648-653

Rutherford AW, Osyczka A, Rappaport F (2012) Back-reactions, short-circuits, leaks and other energy wasteful reactions in biological electron transfer: Redox tuning to survive life in O₂. *FEBS Lett.* **586**: 603-616

Rutherford AW, Paterson DR, Mullet JE (1981) A light-induced spin-polarized triplet detected by EPR in Photosystem II reaction centers. *Biochim. Biophys. Acta* **635**: 205-214

Sacharz J, Giovagnetti V, Ungerer P, Mastroianni G, Ruban AV (2017) The xanthophyll cycle affect reversible interactions between PsbS and light-harvesting complex II to control non-photochemical quenching. *Nat. Plant.* **3**: 16225

Shevela D, Eaton-Rye JJ, Shen JR, Govindjee (2012) Photosystem II and the unique role of bicarbonate: a historical perspective. *Biochim. Biophys. Acta.* **1817**: 1134-1151

Stemler A, Murphy J (1983) Determination of the binding constant of H¹⁴CO₃⁻ to the photosystem II complex in maize chloroplasts - effects of inhibitors and light. *Photochem. Photobiol.* **38**: 701-707

Tikkanen M, Aro EM (2012) Thylakoid protein phosphorylation in dynamic regulation of photosystem II in higher plants. *Biochim. Biophys. Acta* **1817**: 232-238

Tomizioli M, Lazar C, Brugière S, Burger T, Salvi D, Gatto L, Moyet L, Breckels LM, Hesse A-M, Lilley K-S, Seigneurin-Berny D, Finazzi G, Rolland N, Ferro N (2014) Deciphering thylakoid sub-compartments using a mass spectrometry-based approach. *Mol Cell Proteomics.* **13**: 2147-2167

Vass I, Kirilovsky D, Etienne A-L (1999) UV-B radiation-induced donor- and acceptor-side modifications of Photosystem II in the cyanobacterium *Synechocystis* sp. PCC 6803 *Biochem.* **38**: 12786-12794

Vothknecht U C, Westhoff P (2002) Biogenesis and origin of thylakoid membranes. *Biochim. Biophys. Acta* **1541**: 91-101

Ware MA, Giovagnetti V, Belgio E, Ruban AV (2015) PsbS protein modulates non-photochemical chlorophyll fluorescence quenching in membranes depleted of photosystems. *J. Photochem. Photobiol. B.* **152**: 301-307

Watanabe M, Iwai M, Narikawa R, Ikeuchi M (2009) Is the photosystem II complex a monomer or a dimer? *Plant Cell Physiol.* **50**: 1674-1680

Wei L, Guo J, Ouyang M, Sun X, Ma J, Chi W, Lu C, Zhang L (2010) LPA19, a Psb27 homolog

880 in *Arabidopsis thaliana*, facilitates D1 protein precursor processing during PSII biogenesis. *J. Biol.*
881 *Chem.* **285**: 21391-21398

882 **Xiao Y, Huang G, You X, Zhu Q, Wang W, Kuang T, Han G, Sui SF, Shen JR** (2021) Structural
883 insights into cyanobacterial photosystem II intermediates associated with Psb28 and Tsl0063. *Nat.*
884 *Plants* **7**: 1132–1142

885 **Yamada M, Nagao R, Iwai M, Arai Y, Makita A, Ohta H, Tomo T** (2018) The PsbQ'protein
886 affects the redox potential of the Q_A in photosystem II. *Photosynthetica* **56**: 185–191

887 **Zabert J, Bohn S, Schuller SK, Arnolds O, Möller M, Meier-Credo J, Liauw P, Chan A,**
888 **Tajkhorshid E, Langer JD, Stoll R, Krieger-Liszkay A, Engel BD, Rudack T, Schuller JM,**
889 **Nowaczyk MM** (2021) Structural insights into a photosystem II assembly. *Nat. Plants* **7**: 524–538

890 **Zimmermann K, Heck M, Frank J, Kern J, Vass I, Zouni A** (2006) Herbicide binding and
891 thermal stability of photosystem II isolated from *Thermosynechococcus elongatus*. *Biochim.*
892 *Biophys. Acta* **1757**: 106-114

893
894

LAMELLAL MEMBRANE

GRANAL MEMBRANE

From assembly and repair pathway

To dimerization and antenna binding

Figure 1

Figure 2

Figure 3

Figure 4

Parsed Citations

- Albertsson P-A (2001)** A quantitative model of the domain structure of the photosynthetic membrane. *Trends in Plant Sci.* 6: 349-354
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Alfonso M, Montoya G, Cases R, Rodríguez R, Picorel R (1994)** Core antenna complexes, CP43 and CP47, of higher plant photosystem II. Spectral properties, pigment stoichiometry, and amino acid composition. *Biochem.* 33: 10494-10500
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Andersson B, Anderson J-M (1980)** Lateral heterogeneity in the distribution of chlorophyll-protein complexes of the thylakoid membranes of spinach chloroplasts. *Biochim. Biophys. Acta* 593: 427-440
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Avramov AP, Hwang HJ, Burnap RL (2020)** The role of Ca²⁺ and protein scaffolding in the formation of nature's water oxidizing complex. *Proc. Natl. Acad. Sci. USA* 117: 28036-28045
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bassi R, Dall'Osto L (2021)** Dissipation of light energy absorbed in excess: the molecular mechanisms. *Annu. Rev. Plant Biol.* 72: 47-76
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bennoun P (1994)** Chlororespiration revisited: Mitochondrial-plastid interactions in *Chlamydomonas*. *Biochim. Biophys. Acta* 1186: 59-66
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bergantino E, Segalla A, Brunetta A, Teardo E, Rigoni F, Giacometti GM, Szabò I (2003)** Light- and pH-dependent structural changes in the PsbS subunit of photosystem II. *Proc. Natl. Acad. Sci. USA* 100: 15265-15270
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Brinkert K, De Causmaecker S, Krieger-Liszkay A, Fantuzzi A, Rutherford AW (2016)** Bicarbonate-induced redox tuning in Photosystem II for regulation and protection. *Proc. Natl. Acad. Sci. U S A* 113: 12144-12149
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Caffarri S, Kouril R, Kerešič S, Boekema EJ, Croce R (2009)** Functional architecture of higher plant photosystem II supercomplexes. *EMBO J.* 28: 3052-3063
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Chen H, Zhang D, Guo J, Wu H, Jin M, Lu Q, Lu C, Zhang L (2006)** A Psb27 homologue in *Arabidopsis thaliana* is required for efficient repair of photodamaged photosystem II. *Plant Mol. Biol.* 61: 567-575
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Collu G, Farci D, Esposito F, Pintus F, Kirkpatrick J, Piano D (2017)** New insights into the operative network of FaEO, an enone oxidoreductase from *Fragaria x ananassa* Duch. *Plant Mol. Biol.* 94:125-136
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Correa-Galvis V, Poschmann G, Melzer M, Stühler K, Jahns P (2016)** PsbS interactions involved in the activation of energy dissipation in *Arabidopsis*. *Nat. Plants.* 2: 15225
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Crofts AR, Wraight CA (1983)** The electrochemical domain of photosynthesis. *Biochim. Biophys. Acta* 726: 149-185
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Dall'Osto L, Cazzaniga S, Bressan M, Paleček D, Židek K, Niyogi KK, Fleming GR, Zigmantas D, Bassi R (2017)** Two mechanisms for dissipation of excess light in monomeric and trimeric light-harvesting complexes. *Nat. Plants.* 3:17033
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Danielsson R, Suorsa M, Paakkarinen V, Albertsson P-A, Styring S, Aro E-M, Mamedov F (2006)** Dimeric and monomeric organization of photosystem II. *J. Biol. Chem.* 281: 14241-14249
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Dau H, Zaharieva I (2009)** Principles, efficiency, and blueprint character of solar-energy conversion in photosynthetic water oxidation. *Acc Chem Res.* 42: 1861-1870
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- De Causmaecker S, Douglass JS, Fantuzzi A, Nitschke N, Rutherford AW (2019)** Energetics of the exchangeable quinone, QB, in Photosystem II. *Proc. Natl. Acad. Sci.* 116: 9458-9463
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Debus RJ, Campbell KA, Pham DP, Hays A-MA, Britt RD (2000)** Glutamine 189 of the D1 polypeptide modulates the magnetic and

redox properties of the manganese cluster and tyrosine Yz in Photosystem II. *Biochem.* 39: 6275-6287

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Dietzel L, Bräutigam K, Steiner S, Schöffler K, Lepetit B, Grimm B, Schöttler MA, Pfannschmidt T (2011) Photosystem II supercomplex remodelling serves as an entry mechanism for state transitions in *Arabidopsis*. *Plant Cell.* 23: 2964-2977

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ducruet J, Vass I (2009) Thermoluminescence: experimental. *Photosynth. Res.* 101: 195–204

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fan M, Li M, Liu Z, Cao P, Pan X, Zhang H, Zhao X, Zhang J, Chang W (2015) Crystal structures of the PsbS protein essential for photoprotection in plants. *Nat. Struct. Mol. Biol.* 22: 729-735

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fey H, Piano D, Horn R, Fischer D, Schmidt M, Ruf S, Schröder WP, Bock R, Büchel C (2015) Isolation of highly active photosystem II core complexes with a His-tagged Cyt b559 subunit from transplastomic tobacco plants. *Biochim. Biophys. Acta* 1777: 1501–1509

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gachek DA, Holleboom CP, Tietz S, Kirchhoff H, Walla PJ (2019) PsbS-dependent and -independent mechanisms regulate carotenoid-chlorophyll energy coupling in grana thylakoids. *FEBS Lett.* 593: 3190-3197

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Grasse N, Mamedov F, Becker K, Styring S, Rögner M, Nowaczyk MM (2011) Role of novel dimeric Photosystem II (PSII)-Psb27 protein complex in PSII repair. *J. of Biol. Chem.* 286: 29548-29555

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Haniewicz P, De Sanctis D, Büchel C, Schröder WP, Loi M C, Kieselbach T, Bochtler M, Piano D (2013) Isolation of monomeric photosystem II that retains the subunit PsbS. *Photosynth Res.* 118: 199-207

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Haniewicz P, Floris D, Farci D, Kirkpatrick J, Loi MC, Büchel C, Bochtler M, Piano D (2008) Isolation of Plant Photosystem II Complexes by Fractional Solubilization. *Front. Plant Sci.* 6:1100

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Huang G, Xiao Y, Pi X, Zhao L, Zhu Q, Wang W, Kuang T, Han G, Sui SF, Shen JR (2021) Structural Insights into a dimeric Psb27-photosystem II complex from a cyanobacterium *Thermosynechococcus vulcanus*. *Proc. Natl. Acc. Sci. USA* 118: e2018053118

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Johnson GN, Rutherford AW, Krieger A (1995) A change in the midpoint potential of the quinone Q(A) in Photosystem II associated with photoactivation of oxygen evolution. *Biochim. Biophys. Acta.* 1229: 202-207

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Johnson M P (2016) Photosynthesis. *Essays Biochem.* 60: 255–273

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kirchhoff H (2019) Chloroplast ultrastructure in plants. *New Phytologist* 223: 565-574

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Komenda J, Lupinkova L, Kopecky J (2002) Absence of the psbH gene product destabilizes Photosystem II complex and bicarbonate binding on its acceptor side in *Synechocystis* PCC 6803. *Eur. J. Biochem.* 269: 610-619

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Komenda J, Sobotka R, Nixon PJ (2012) Assembling and maintaining the Photosystem II complex in chloroplasts and cyanobacteria. *Curr. Op. in Plant Biol.* 15: 245-251

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Krausz E, Hughes JL, Smith P, Pace R, Årsköld SP (2005) Oxygen-evolving Photosystem II core complexes: a new paradigm based on the spectral identification of the charge-separating state, the primary acceptor and assignment of low-temperature fluorescence. *Photochem. Photobiol. Sci.* 4: 744-753

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Krieger-Liszkay A (2005) Singlet oxygen production in photosynthesis. *J. of Exp. Bot.* 56: 337–346

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lavergne J, Leci E (1993) Properties of inactive Photosystem II centres. *Photosynth. Res.* 35: 323-343

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li XP, Gilmore AM, Caffarri S, Bassi R, Golan T, Kramer D, Niyogi KK (2004) Regulation of photosynthetic light harvesting involves intrathylakoid lumen pH sensing by the PsbS protein. *J Biol Chem.* 279: 22866-22874

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liguori N, Campos SRR, Baptista AM, Croce R (2019) Molecular Anatomy of Plant Photoprotective Switches: The Sensitivity of PsbS to the Environment, Residue by Residue. J Phys. Chem. Lett. 10:1737-1742

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu H, Roose JL, Cameron JC, Pakrasi HB (2011) A genetically tagged Psb27 protein allows purification of two consecutive Photosystem II (PSII) assembly intermediates in Synechocystis 6803, a cyanobacterium. J. of Biol. Chem. 286: 24865-24871

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ljungberg U, Åkerlund HE, Larsson C, Andersson B (1984) Identification of polypeptides associated with the 23 and 33 kDa proteins of photosynthetic oxygen evolution. Biochim. Biophys. Acta. 767: 145-152

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mamedov F, Nowaczyk MM, Thapper A, Rogner M, Styring S (2007) Functional characterization of monomeric Photosystem II core preparations from Thermosynechococcus elongatus with or without the Psb27 protein. Biochem. 46: 5542-5551

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Melis A (1985) Functional properties of Photosystem II β in spinach chloroplasts. Biochim Biophys Acta. 808: 334-342

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nickelsen J, Rengstl B (2013) Photosystem II assembly: from cyanobacteria to plants. Annu. Rev. Plant Biol. 64: 609-635

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nixon PN, Trost JT, Diner BA (1992) Role of carboxyl terminus of polypeptide D1 in the assembly of a functional water-oxidizing manganese cluster in Photosystem II of the cyanobacterium Synechocystis sp. PCC 6803: assembly requires a free carboxyl group at C-terminal position 344. Biochem. 31:10859-10871

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Niyogi KK, Truong TB (2013) Evolution of flexible non-photochemical quenching mechanisms that regulate light harvesting in oxygenic photosynthesis. Curr. Op. Plant Biol. 16: 307-314

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nowaczyk MM, Hebel R, Schlodder E, Meyer HE, Warscheid B, Rögner M (2006) Psb27, a cyanobacterial lipoprotein, is involved in the repair cycle of photosystem II. Plant Cell. 18: 3121-3131

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Piano D, El Alaoui S, Korza HJ, Filipek R, Sabala I, Haniewicz P, Buechel C, De Sanctis D, Bochtler M (2010) Crystallization of the photosystem II core complex and its chlorophyll binding subunit CP43 from transplastomic plants of Nicotiana tabacum. Photosynth. Res. 106: 221-226

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Porra RJ, Thompson WA, Kriedmann PE (1989) Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophylls a and b with four different solvents: verifications of the concentration of chlorophyll standards by atomic absorption spectroscopy. Biochim. Biophys. Acta 975: 384-394

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pospíšil P (2016) Production of Reactive Oxygen Species by Photosystem II as a Response to Light and Temperature Stress. Front Plant Sci.7: 1950

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Puthiyaveetil S, Tsabari O, Lowry T, Lenhert S, Lewis RR, Reich Z, Kirchhoff H (2014) Compartmentalization of the protein repair machinery in photosynthetic membrane. Proc. Natl. Acad. Sci. USA 111: 15839-15844

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Roach T, Krieger-Liszkay A (2012) The role of the PsbS protein in the protection of photosystems I and II against high light in Arabidopsis thaliana. Biochim. Biophys. Acta. 1817: 2158-2165

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Roose JL, Pakrasi HB (2004) Evidence that D1 processing is required for manganese binding and extrinsic protein assembly into Photosystem II. J. Biol. Chem. 279: 45417-45422

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Roose JL, Pakrasi HB (2008) The Psb27 protein facilitates manganese cluster assembly in photosystem II. J. Biol. Chem. 283: 4044-4050

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ruban AV, Johnson M P (2015) Visualizing the dynamic structure of the plant photosynthetic membrane. Nat. Plants 1: 15161

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ruban AV (2016) Nonphotochemical chlorophyll fluorescence quenching: mechanism and effectiveness in protecting plants from photodamage. Plant Physiol. 170: 1903-1916

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rutherford AW, Crofts AR, Inoue Y (1982) Thermoluminescence as a probe of Photosystem II photochemistry: the origin of the flash induced glow peaks. Biochim. Biophys. Acta 682: 457-465.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rutherford AW, Krieger-Liszkay A (2001) Herbicide-induced oxidative stress in Photosystem II. Trends In Biochem. Sc. 26: 648–653

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rutherford AW, Osyczka A, Rappaport F (2012) Back-reactions, short-circuits, leaks and other energy wasteful reactions in biological electron transfer: Redox tuning to survive life in O₂. FEBS Lett. 586: 603–616

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rutherford AW, Paterson DR, Mullet JE (1981) A light-induced spin-polarized triplet detected by EPR in Photosystem II reaction centers. Biochim. Biophys. Acta 635: 205–214

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sacharz J, Giovagnetti V, Ungerer P, Mastroianni G, Ruban AV (2017) The xanthophyll cycle affect reversible interactions between PsbS and light-harvesting complex II to control non-photochemical quenching. Nat. Plant. 3: 16225

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Shevela D, Eaton-Rye JJ, Shen JR, Govindjee (2012) Photosystem II and the unique role of bicarbonate: a historical perspective. Biochim. Biophys. Acta. 1817: 1134-1151

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Stemler A, Murphy J (1983) Determination of the binding constant of H¹⁴CO₃⁻ to the photosystem II complex in maize chloroplasts - effects of inhibitors and light. Photochem. Photobiol. 38: 701–707

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tikkanen M, Aro EM (2012) Thylakoid protein phosphorylation in dynamic regulation of photosystem II in higher plants. Biochim. Biophys. Acta 1817: 232-238

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tomizioli M, Lazar C, Brugière S, Burger T, Salvi D, Gatto L, Moyet L, Breckels LM, Hesse A-M, Lilley K-S, Seigneurin-Berny D, Finazzi G, Rolland N, Ferro N (2014) Deciphering thylakoid sub-compartments using a mass spectrometry-based approach. Mol Cell Proteomics. 13: 2147-2167

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vass I, Kirilovsky D, Etienne A-L (1999) UV-B radiation-induced donor- and acceptor-side modifications of Photosystem II in the cyanobacterium Synechocystis sp. PCC 6803 Biochem. 38: 12786-12794

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vothknecht U C, Westhoff P (2002) Biogenesis and origin of thylakoid membranes. Biochim. Biophys. Acta 1541: 91-101

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ware MA, Giovagnetti V, Belgio E, Ruban AV (2015) PsbS protein modulates non-photochemical chlorophyll fluorescence quenching in membranes depleted of photosystems. J. Photochem. Photobiol. B. 152: 301-307

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Watanabe M, Iwai M, Narikawa R, Ikeuchi M (2009) Is the photosystem II complex a monomer or a dimer? Plant Cell Physiol. 50: 1674–1680

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wei L, Guo J, Ouyang M, Sun X, Ma J, Chi W, Lu C, Zhang L (2010) LPA19, a Psb27 homolog in Arabidopsis thaliana, facilitates D1 protein precursor processing during PSII biogenesis. J. Biol. Chem. 285: 21391-21398

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Xiao Y, Huang G, You X, Zhu Q, Wang W, Kuang T, Han G, Sui SF, Shen JR (2021) Structural insights into cyanobacterial photosystem II intermediates associated with Psb28 and Tsl0063. Nat. Plants 7: 1132–1142

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yamada M, Nagao R, Iwai M, Arai Y, Makita A, Ohta H, Tomo T (2018) The PsbQ' protein affects the redox potential of the QA in photosystem II. Photosynthetica 56: 185–191

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zabert J, Bohn S, Schuller SK, Arnolds O, Möller M, Meier-Credo J, Liauw P, Chan A, Tajkhorshid E, Langer JD, Stoll R, Krieger-Liszkay A, Engel BD, Rudack T, Schuller JM, Nowaczyk MM (2021) Structural insights into a photosystem II assembly. Nat. Plants 7: 524–538

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zimmermann K, Heck M, Frank J, Kern J, Vass I, Zouni A (2006) Herbicide binding and thermal stability of photosystem II isolated from *Thermosynechococcus elongatus*. *Biochim. Biophys. Acta* 1757: 106-114

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)