1 **Rapid transgenerational adaptation in response to intercropping increases**
2 **facilitation and reduces competition**

3 **Laura Stefan**¹, Nadine Engbersen¹, Christian Schöb¹

4 ¹Institute of Agricultural Sciences, ETH, Zürich, Switzerland

5

6

Summary:

By capitalising on positive biodiversity-productivity relationships, intercropping provides opportunities to improve agricultural sustainability¹. However, intercropping is generally implemented using commercial seeds that were bred for maximal productivity in monocultures, which might limit the benefits of crop diversity on yield^{2,3}. Plants can adapt over generations to the level of surrounding plant diversity, notably through increases in niche differentiation⁴. However, this adaptation potential and the corresponding yield benefit potential have not been explored in annual crop systems. Here we show that plant–plant interactions among annual crops evolved towards increased facilitation and reduced competition when the plants’ coexistence history matched their current diversity setting, which led to an increase in overyielding of up to 58%. These higher yield benefits were linked to character convergence between species sharing the same coexistence history for two generations. Notably, the six crop species tested converged towards taller phenotypes with lower leaf dry matter content when grown in mixtures. This study provides the first empirical evidence for the importance of parental diversity affecting plant–plant interactions and ecosystem functioning of the following generations in annual cropping systems. These results have important implications for diversified agriculture as they demonstrate the yield potential of targeted cultivars for intercropping, which can be achieved through specific breeding for mixtures.

Following decades of studies demonstrating the positive relationship between species diversity and plant primary productivity in natural systems^{5,6}, intercropping, i.e. growing more than two species in the same field during the same period, has been increasingly considered as a promising option to increase agricultural sustainability^{1,7}. The productivity benefits of increasing species diversity rely on two main mechanisms, namely selection effects and complementarity effects, the latter encompassing both facilitation and niche differentiation^{8,9}. In perennial natural grasslands, complementarity effects have been shown to increase over time due to evolutionary processes^{4,10,11}. Notably, greater species complementarity can result from evolutionary changes¹² – i.e. changes in gene frequency – or from heritable epigenetic changes¹³ affecting species traits in response to surrounding plant diversity, which either increases niche differentiation (i.e. reduces competition) or facilitation¹⁴. The evolutionary potential of plant–plant interactions in diverse communities has tremendous implications for the diversification of agricultural systems¹⁵. This is of particular relevance for mixed cropping systems, where the use of commercial seeds domesticated and bred for maximum yield in monoculture is the norm, which may compromise the diversity benefits^{2,3,16–18}. Despite the paramount importance of this question, the yield potential of mixture-adapted varieties is, to our knowledge, unknown, as are the character differences of monoculture- compared to mixture-adapted crops. Therefore, in this project, we determined whether and how crop species adapt over three generations to the level of plant

diversity that they are surrounded by. We investigated how plant–plant interactions, i.e. competition and facilitation, and plant traits changed and evolved within different coexistence histories over time, and whether these changes translated into yield benefits. To that end, we conducted an intercropping experiment in Switzerland with six different crop species commonly cultivated in Europe and belonging to four functionally different phylogenetic groups. The mesocosms included monocultures, 13 different 2-species mixtures, four different 4-species mixtures, and isolated single plants, and was replicated in two different fertilizing conditions. We selected open-pollinated varieties as seed source to provide the genetic variability needed for evolutionary processes to occur. To assess potential transgenerational changes, we repeated the experiment over the course of three years with seeds from plants grown from either monocultures, mixtures, or single individual plants of the previous year (Fig. 1, Fig. 5).

Figure 1. Experimental design. Six crop species were used to sow single plant individuals (6), monocultures (6), 2-species mixtures (13) and 4-species mixtures (4) in 2018 (Year 1); seeds were collected at the end of the growing season and resown in 2019 (Year 2) in the same diversity setting as their previous generation. Seeds were collected again and resown in 2020 (Year 3), this time either in the same community their seeds were collected from [same coexistence history], or in a community different to the one of their parents [different coexistence history] ($n = 468$ plots). This process was replicated in two different fertilizing conditions. We expected that crops growing in the same community as their parents would have adapted over the two generations, and therefore would exhibit less competition and have higher productivity than crops growing in a community different to the one of their parents.

Results from the third year showed that plant–plant interactions shifted towards stronger facilitation and weaker competition when the plants were growing in the same community conditions than their two previous generations (Fig. 2, Extended Data Fig. 1, Extended Data Table 1). More precisely, net

interaction index, as well as competition and facilitation indexes, were significantly higher when the crops were grown in the community their seeds were collected from than when they were growing in a community different to the one of their parents (Fig. 2, Extended Data Fig. 1; +54% for the net index, +9% for the competition index, +93% for the facilitation index). Pairwise comparisons further showed that this effect of coexistence history was particularly true in mixtures and only a trend in monocultures, for both fertilizing conditions (Extended Data Table 2). This notably demonstrates that in mixtures, mixture-adapted communities (i.e. with the same coexistence history) exhibited more facilitation and less competition than monoculture-adapted communities or single plant-adapted

communities (i.e. with a different coexistence history).

Figure 2: Plant interaction index in response to coexistence history

Net interaction index of monocultures, 2- and 4-species mixtures in response to coexistence history, for fertilized and unfertilized conditions. $n = 276$. This index compares the performance of plants growing in communities to the performance of single plants (see Methods). Negative interaction index indicates

competition within a community, positive interaction index indicates facilitation. The closer this index gets to 1, respectively -1, the stronger the facilitation, respectively competition. “Same coexistence history” indicates that crops were grown in the community their seeds were collected from. “Different coexistence history” refers to crops grown in a community different to the one of their parents. The effect of fertilization and coexistence history were highly significant. See Extended Data Table 1 for the complete statistical analysis, and Extended Data Fig. 1 for competition and facilitation indexes.

Horizontal lines represent the median of the data, boxes represent the lower and upper quartiles (25% and 75%), with vertical lines extending from the hinge of the box to the smallest and largest values, no further than $1.5 \times$ the interquartile range. Data beyond the end of the whiskers are outlying and plotted individually.

This shift in plant–plant interactions was accompanied by a similar shift in net biodiversity effect (NE) in fertilized plots (Fig. 3a). Net biodiversity effect was calculated following the method of Loreau & Hector (2001) and represents the deviation from the expected yield in the mixture, based on the yield of the corresponding monocultures⁸. We observed that under fertilized conditions, NE was on average 58% higher with the same coexistence history than with a different coexistence history (Fig. 3a, Extended Data Table 3), which corresponded to an increase in total yield ranging from 8 to 22% in mixtures (Fig. 3b). This indicates that in fertilized plots, the yield benefits of crop mixtures were higher with mixture-adapted individuals compared to monoculture-adapted and single-adapted individuals. Interestingly, in unfertilized plots we did not observe the same trend. When looking at the

partitioning of net effects into complementarity and selection effects⁸, we only observed a significant effect of coexistence history on selection effects under fertilized conditions for 4-species mixtures (Extended Data Fig. 2b).

Figure 3: Effects of coexistence history on net biodiversity effects (a) and total yield per plot (b)

Effects of coexistence history and crop species number on (a) net biodiversity effect – reflecting the yield advantage of mixtures compared to monocultures – and (b) total yield per plot (square-root transformed) in fertilized and unfertilized plots. (a) $n=276$; (b) $n=204$. “Same coexistence history” indicates that crops were grown in the community their seeds were collected from. “Different coexistence history” refers to crops grown in a community different to the one of their parents. See Extended Data Table 3 & 4 for the complete statistical analysis, and Extended Data Fig. 2 for complementarity and selection effects.

Horizontal lines represent the median of the data, boxes represent the lower and upper quartiles (25% and 75%), with vertical lines extending from the hinge of the box to the smallest and largest values, no further than 1.5 * the interquartile range. Data beyond the end of the whiskers are outlying and plotted individually.

To investigate the ecological mechanisms behind the shift in plant–plant interactions and biodiversity effects with coexistence history, we measured standard above-ground plant traits and compared the average values as well as coefficients of variation at the species and community levels of single-, monoculture- and mixture-adapted varieties. Following traditional niche theory, we expected that the observed reduction in competition would be linked to an increase in functional trait variation, thereby reflecting an increase in niche differentiation. Surprisingly, we did not observe character displacement – i.e. increased trait variation⁴ – in our intercrop systems, but rather character convergence – i.e. reduced trait variation (Fig. 4). More specifically, we found a reduction in trait variation at the community level, notably of height and leaf dry matter content: the coefficient of variation of height was lower in the same coexistence history treatment compared to a different coexistence history (-9%) (Fig. 4d), and for leaf dry matter content it was 15% lower with the same history compared to a different history (Fig. 4e). Furthermore, the coefficient of variation of mass per seed was also lower

under the same history compared to a different history, but this effect was only significant in monocultures (-33%) (Fig. 4f). The community-weighted means of plant traits (CWM, calculated at the community level) further suggest that when growing in the same coexistence history, plants seem to converge towards taller individuals with lower leaf dry matter content. Indeed, the community-weighted mean of leaf dry matter content was significantly lower with the same coexistence history compared to a different history (-3%, Fig 4); height community-weighted mean was – although non-significantly – higher in the case of the same history compared to different coexistence history (Fig 4a). We observed similar responses of height and leaf dry matter content at the species level (Extended Data Fig. 5 & 6, Extended Data Tables 5-9).

Figure 4: Plot-level traits response to coexistence history

Effects of coexistence history and crop species number on community-weighted mean (CWM) of height (in cm) (a), Leaf Dry Matter Content (LDMC) (b), and mass per seed (in g) (c), and on coefficient of variation at the community level of height (d), Leaf Dry Matter Content LDMC (e), and mass per seed (f). $n = 271$. "Same coexistence history" indicates that crops were grown in the community their seeds were collected from. "Different coexistence history" refers to crops grown in a community different to the one of their parents. Dots represent the mean values across plots; lines

represent the standard error. See Extended Data Tables 10-14 for the complete statistical analysis, and Extended Data Fig. 5-6 as well as Extended Data Tables 5-9 for responses at the species level.

Our research demonstrates that, after only two generations, annual crop plants growing in the same diversity setting as their preceding generations showed reduced competition and increased facilitation compared to plants growing in a different diversity setting as their parents, which led to increased overyielding. We further investigated whether character displacement was responsible for this evolution of plant–plant interactions; contrary to our hypotheses, results did not show evidence for character displacement, but rather for character convergence in plant aboveground traits.

The observed shift in plant–plant interactions are consistent with several grassland studies investigating the effects of community evolution on community productivity and niche differentiation, where it was found that common rapid evolution in plant communities can lead to increases in ecosystem functioning^{4,10,11,19}. We indeed observed a positive effect of common community history on the net biodiversity effect (i.e. overyielding), which means that the yield benefit of mixtures compared to monocultures was higher when the plants had been adapted to growing in mixtures (Fig. 3). This can explain why diversity effects generally increase over time^{20,21}. Here we did not observe a significant increase in complementarity effect in response to common community history (Extended Data Fig. 2a). However, we observed a similar trend – although nonsignificant – for CE as for NE (Extended Data Fig. 2a): in fertilized plots, CE tended to be higher in the case of a mixture coexistence history, notably in 2-species mixtures. We suggest that the limited timeframe of this study – two generations – might be the reason for the lack of more significant changes in CE and emphasizes the need for longer-term research to confirm or infirm this trend. Surprisingly, selection effects also increased in 4-species mixtures in response to coexistence history (Extended Data Fig. 2b). This is unexpected, as selection effects have not, to our knowledge, been shown to increase over time²². However, it might be that this short common community history has favoured a specific species or a specific trait that was particularly plastic or beneficial for fitness^{23,24}.

The above-mentioned increases in biodiversity, complementarity and selection effects were only present in fertilized conditions, which could indicate that the benefits of common community history might be dependent on the abiotic conditions. This is nonetheless consistent with several recent studies demonstrating that biodiversity effects are higher in high-inputs systems^{2,25,26}, and emphasizes the role of fertilization in driving these effects. Indeed, by promoting crop growth and, consequently, higher competition between plants, fertilization may foster higher benefits of niche differentiation^{27–29}.

Overall, increases in biodiversity effects are associated with changes in species traits in response to surrounding plant diversity^{4,14,30}. Traditional hypotheses of trait and niche theory indeed predict that when several species co-occur closely together, selection over generations would favour character displacement that would reduce resource overlap and consequently increase niche differentiation^{31,32}.

Surprisingly, here we found the reverse and observed that a reduction in trait variation favoured increased yield benefits in mixtures. Furthermore, functional diversity – calculated as the volume occupied in the space of the considered traits in this study³³ – did not respond to common coexistence history (Extended Data Fig. 7 & Table 15). While surprising, this result is not unheard of²³, and suggests that our plants might have adapted to express the phenotype that would maximise their fitness^{34–36}. This ideal phenotype is, in our mixture communities, characterized by taller plants with lower leaf dry matter content, the latter indicating soft leaves associated with rapid biomass production³⁷, and consequently less resource-conservative strategies³⁸. Lower leaf dry matter content has recently been associated with lower parental or ambient competition³⁹, which is consistent with our results of plant–plant interaction intensities. The traits examined here did not allow to understand the mechanisms behind the observed reduction in competition; we suggest that other traits or processes not measured in this experiment might have responded to the coexistence history treatment. Notably, there could be a shift in below-ground traits, such as root-associated traits³⁹, or temporal differentiation of resource capture⁴⁰, such as light. We indeed observed a significant increase in light capture ability in plants coming from the same diversity setting compared to the same communities but a different coexistence history (Extended Data Fig. 8 & Table 16), which indicates that plants used to growing in the same diversity setting during several generations might capture the resources more fully than plants coming from a different diversity setting. However, here we rely on our light interception measurements and suggest more longer-term studies to understand changes in the use of other resources, such as nutrients or water, and how this is associated to plant traits. Furthermore, the scope of this study did not allow us to investigate the mechanisms behind these changes in plant–plant interactions and traits in response to coexistence history. The adaptation response might be genetically-based and due to natural selection¹¹, as we specifically selected open-pollinated varieties in order to ensure a minimum amount of genetic variability. Furthermore, outcrossing could have occurred in the first year of this experiment, as we had a similar experiment running in the same experimental garden with Spanish varieties from the same species^{27,41}. However, considering the short timeframe of this study and the low rate of outcrossing in most of our species, epigenetic changes – i.e, stable heritable changes in cytosine methylation – might also have played an important role as potential evolutionary mechanisms^{13,42–46}.

For the first time, our study provides empirical evidence for rapid transgenerational adaptation in response to diversity history in annual crop communities. Notably, we demonstrated that when plants were coming from the same diversity setting as their parents, plant–plant interactions shifted towards reduced competition and increased facilitation. This effect was particularly true for mixtures and translated into enhanced overyielding under fertilized conditions. This reduction in competition was surprisingly not linked to character displacement, but we instead observed character convergence towards taller plants with lower leaf dry matter content. This research emphasizes the importance of considering transgenerational effects of diversity for crop mixtures. This is particularly relevant for

breeding programs and highlights the need of including diversity when breeding for crop mixtures, in order to design varieties specifically adapted for intercropping¹⁷.

References

1. Brooker, R. W. *et al.* Improving intercropping: a synthesis of research in agronomy, plant physiology and ecology. *New Phytol.* **206**, 107–117 (2015).
2. Chen, J. *et al.* Diversity increases yield but reduces reproductive effort in crop mixtures. *bioRxiv* 2020.06.12.149187 (2020) doi:10.1101/2020.06.12.149187.
3. Thrall, P. H. Darwinian Agriculture: How understanding evolution can improve agriculture R. Ford Denison 2012. Princeton University Press. *Evolutionary Applications* vol. 6 408–410 (2013).
4. Zuppinger-Dingley, D. *et al.* Selection for niche differentiation in plant communities increases biodiversity effects. *Nature* **515**, 108–111 (2014).
5. Tilman, D. *et al.* Diversity and productivity in a long-term grassland experiment. *Science* (80-.). **294**, 843–845 (2001).
6. Spehn, E. M. *et al.* Ecosystem effects of biodiversity manipulations in european grasslands. *Ecol. Monogr.* **75**, 37–63 (2005).
7. Gurr, G. M. *et al.* Multi-country evidence that crop diversification promotes ecological intensification of agriculture. *Nat. plants* **2**, 16014 (2016).
8. Loreau, M. & Hector, A. Partitioning selection and complementarity in biodiversity experiments. *Nature* **412**, 72–76 (2001).
9. Hooper, D. U. *et al.* Effects of biodiversity on ecosystem functioning: a consensus of current knowledge. *Ecol. Monogr.* **75**, 3–35 (2005).
10. van Moorsel, S. J. *et al.* Community evolution increases plant productivity at low diversity. *Ecol. Lett.* **21**, 128–137 (2018).
11. van Moorsel, S. J. *et al.* Evidence for rapid evolution in a grassland biodiversity experiment.

- 279 *Mol. Ecol.* **28**, 4097–4117 (2019).
- 280 12. Anderson, J. T., Willis, J. H. & Mitchell-Olds, T. Evolutionary genetics of plant adaptation.
281 *Trends Genet.* **27**, 258–266 (2011).
- 282 13. Verhoeven, K. J. F., VonHoldt, B. M. & Sork, V. L. Epigenetics in ecology and evolution:
283 what we know and what we need to know. *Mol. Ecol.* **25**, 1631–1638 (2016).
- 284 14. Schöb, C., Brooker, R. W. & Zuppinger-Dingley, D. Evolution of facilitation requires diverse
285 communities. *Nat. Ecol. Evol.* **2**, 1381–1385 (2018).
- 286 15. Isbell, F. *et al.* Benefits of increasing plant diversity in sustainable agroecosystems. *Journal of*
287 *Ecology* vol. 105 871–879 (2017).
- 288 16. Chacón-Labella, J., García Palacios, P., Matesanz, S., Schöb, C. & Milla, R. Plant
289 domestication disrupts biodiversity effects across major crop types. *Ecology Letters* vol. 22
290 1472–1482 (2019).
- 291 17. Wuest, S. E., Peter, R. & Niklaus, P. A. Ecological and evolutionary approaches to improving
292 crop variety mixtures. *Nat. Ecol. Evol.* 1–10 (2021) doi:10.1038/s41559-021-01497-x.
- 293 18. Annicchiarico, P. *et al.* Do we need specific breeding for legume-based mixtures? *Adv. Agron.*
294 **157**, 141–215 (2019).
- 295 19. van Moorsel, S. J. *et al.* Co-occurrence history increases ecosystem stability and resilience in
296 experimental plant communities. *Ecology* **102**, (2021).
- 297 20. Meyer, S. T. *et al.* Effects of biodiversity strengthen over time as ecosystem functioning
298 declines at low and increases at high biodiversity. *Ecosphere* **7**, e01619 (2016).
- 299 21. Allan, E. *et al.* More diverse plant communities have higher functioning over time due to
300 turnover in complementary dominant species. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 17034–17039
301 (2011).
- 302 22. Cardinale, B. J. *et al.* Impacts of plant diversity on biomass production increase through time
303 because of species complementarity. *Proc. Natl. Acad. Sci.* **104**, 18123–18128 (2007).
- 304 23. Colom, S. M. & Baucom, R. S. Belowground competition favors character convergence but not
305 character displacement in root traits. *New Phytol.* (2020) doi:10.1111/nph.17100.
- 306 24. Turcotte, M. M. & Levine, J. M. Phenotypic Plasticity and Species Coexistence. *Trends in*
307 *Ecology and Evolution* vol. 31 803–813 (2016).
- 308 25. Li, C. *et al.* Syndromes of production in intercropping impact yield gains. *Nat. Plants* **6**, 653–
309 660 (2020).

- 310 26. Stefan, L., Hartmann, M., Engbersen, N., Six, J. & Schöb, C. Positive Effects of Crop Diversity
311 on Productivity Driven by Changes in Soil Microbial Composition. *Front. Microbiol.* **12**, 808
312 (2021).
- 313 27. Stefan, L., Engbersen, N. & Schöb, C. Crop–weed relationships are context□dependent and
314 cannot fully explain the positive effects of intercropping on yield. *Ecol. Appl.* (2021)
315 doi:10.1002/eap.2311.
- 316 28. Bertness, M. D. & Callaway, R. Positive interactions in communities. *Trends Ecol. Evol.* **9**,
317 191–193 (1994).
- 318 29. Goldberg, D. & Novoplansky, A. *On the Relative Importance of Competition in Unproductive*
319 *Environments*. Source: *Journal of Ecology* vol. 85 <https://www.jstor.org/stable/2960565>
320 (1997).
- 321 30. Abakumova, M., Zobel, K., Lepik, A. & Semchenko, M. Plasticity in plant functional traits is
322 shaped by variability in neighbourhood species composition. *New Phytol.* **211**, 455–463
323 (2016).
- 324 31. Pfennig, K. S. & Pfennig, D. W. Character displacement: ecological and reproductive
325 responses to a common evolutionary problem. *Quarterly Review of Biology* vol. 84 253–276
326 (2009).
- 327 32. Losos, J. B. Ecological character displacement and the study of adaptation. *Proceedings of the*
328 *National Academy of Sciences of the United States of America* vol. 97 5693–5695 (2000).
- 329 33. Petchey, O. L. & Gaston, K. J. Functional diversity (FD), species richness and community
330 composition. *Ecol. Lett.* **5**, 402–411 (2002).
- 331 34. Fox, J. W. & Vasseur, D. A. Character convergence under competition for nutritionally
332 essential resources. *Am. Nat.* **172**, 667–680 (2008).
- 333 35. Grant, P. R. Convergent and divergent character displacement. *Biol. J. Linn. Soc.* **4**, 39–68
334 (1972).
- 335 36. Weedon, D. O. & Finckh, R. M. Response of wheat composite cross populations to disease and
336 climate variation over 13 generations. *Front. Agr. Sci. Eng* **2021**, 400–415.
- 337 37. Díaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529**, 167–171 (2016).
- 338 38. Reich, P. B. The world-wide ‘fast-slow’ plant economics spectrum: A traits manifesto. *J. Ecol.*
339 **102**, 275–301 (2014).
- 340 39. Puy, J. *et al.* Competition-induced transgenerational plasticity influences competitive
341 interactions and leaf decomposition of offspring. *New Phytol.* **7**, 3497–3507 (2020).

40. Engbersen, N., Brooker, R. W., Stefan, L., Studer, B. & Schöb, C. Temporal differentiation of resource capture and biomass accumulation as a driver of yield increase in intercropping. *Front. Plant Sci.* **12**, 926 (2021).
41. Chen, J. *et al.* Diversity increases yield but reduces harvest index in crop mixtures. *Nat. Plants* **2021** 1–6 (2021) doi:10.1038/s41477-021-00948-4.
42. Cortijo, S. *et al.* Mapping the epigenetic basis of complex traits. *Science (80-.).* **343**, 1145–1148 (2014).
43. Van Der Graaf, A. *et al.* Rate, spectrum, and evolutionary dynamics of spontaneous epimutations. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 6676–6681 (2015).
44. Saze, H., Scheid, O. M. & Paszkowski, J. Maintenance of CpG methylation is essential for epigenetic inheritance during plant gametogenesis. *Nat. Genet.* **34**, 65–69 (2003).
45. Springer, N. M. Epigenetics and crop improvement. *Trends Genet.* **29**, 241–247 (2013).
46. Puy, J., Carmona, C. P., Dvořáková, H., Latzel, V. & De Bello, F. Diversity of parental environments increases phenotypic variation in Arabidopsis populations more than genetic diversity but similarly affects productivity. *Ann. Bot.* **127**, 425–436 (2021).

Methods

Study sites

The Crop Diversity Experiment took place in 2018, 2019, and 2020 in an outdoor experimental garden located at the Irchel campus of the University of Zurich, Switzerland (47.3961 N, 8.5510 E, 508 m a.s.l). Zurich is characterized by a temperate climate²⁷. The experimental garden was irrigated during the growing season with the aim of maintaining a sufficient amount of water for optimal plant growth. The dry threshold of soil moisture was set at 50% of field capacity, with a target soil moisture of 90% of field capacity. Whenever dry thresholds were reached (measured through PlantCare soil moisture sensors (PlantCare Ltd., Switzerland), irrigation was initiated, and water added until reaching the target value.

Each experimental garden consisted of square plots of 0.25 m². The uppermost 30 cm were filled with standard, not enriched, agricultural soil coming from the local region. This soil consisted of 45 % sand, 45 % silt, and 10 % clay, and initially contained 0.19 % nitrogen (N), 3.39 % carbon (C), and 332 mg total phosphorous (P)/kg, with a mean pH of 7.25. Beneath that, there was local soil of uncharacterized properties that allowed unlimited root growth. The plots were embedded into larger beds of 7 x 1 m, each bed containing 28 plots. Inside a bed, plots were separated from each other by metal frames. While the relatively small plot sizes allowed us to undertake a large experiment under environmentally highly controlled but realistic outdoor conditions, some variables can suffer edge effects and interferences with neighbouring plots. However, such effects would probably increase residual variation more than between-treatment variation, because randomization was used to prevent confounding of between-plot interactions with treatments. In the only relevant study of which we are aware, the biodiversity–productivity relationship in herbaceous communities was not affected by plot size⁴⁷ while a recent theoretical study showed that, if anything, biodiversity effects should increase with plot size⁴⁸.

We therefore assume that effect size in our experiment, if anything, is probably rather conservatively estimated compared with that in studies using larger plot sizes.

Every year, we fertilized half of the beds with N, P and potassium (K) at the concentration of 120 kg/ha N, 205 kg/ha P, and 120 kg/ha K. Fertilizers were applied three times per year, namely once just before sowing (50 kg/ha N, 85 kg/ha P, 50 kg/ha K), once when wheat was at the tillering stage (50 kg/ha N, 85 kg/ha P, 50 kg/ha K), and once when wheat was flowering (20 kg/ha N, 34 kg/ha P, 20 kg/ha K). The other half of the beds served as unfertilized controls. In 2018, we randomly allocated individual beds to a fertilized or non-fertilized control treatment. In the following years, we kept the initial fertilization treatment allocation.

Crop species

Experimental communities were constructed with six annual crop species of agricultural interest. We selected only seed crops with similar growth requirements in terms of climate and length of growing season, and with similar plant sizes to fit at least 40 individuals in the rather small plots. The six species belong to four different phylogenetic groups with varying functional characteristics: we first separated monocots [*Triticum aestivum* (wheat, C3 grass, Poaceae) and *Avena sativa* (oat, C3 grass, Poaceae)] and dicots. Among the dicots, we differentiated between suparasterids [*Coriandrum sativum* (coriander, herb, Apiaceae)] and superrosids. Among the superrosids, we separated legumes [*Lens culinaris* (lentil, legume, Fabaceae)] from non-legumes [*Linum usitatissimum* (flax, herb, Lineaceae) and *Camelina sativa* (false flax, herb, Brassicaceae)]. Furthermore, we chose crop varieties that were locally adapted and commercially available in Switzerland (Table 1).

Species	Switzerland		413
	Ecotype	Supplier	
<i>Avena sativa</i>	Canyon	Sativa Rheinau	414
<i>Triticum aestivum</i>	Fiorina	DSP, Delley	
<i>Coriandrum sativum</i>	Indian	Zollinger Samen, Les Evouettes	415
<i>Lens culinaris</i>	Anicia	Agroscope, Reckenholz	416
<i>Camelina sativa</i>	n.a.	Zollinger Samen, Les Evouettes	
<i>Linum usitatissimum</i>	Lirina	Sativa Rheinau	417

418

419 **Table 1.** List of crop species ecotypes and their suppliers.

420 *Avena Sativa* (oat) is mainly self-pollinating, with outcrossing rates of around 1%⁵². The
421 variety Canyon was acquired in 2014 through conventional selection processes.

422 *Triticum aestivum* (wheat) is principally self-pollinating, with outcrossing rates generally
423 between 1 and 4%^{49,50}, although some cultivars have been shown to have outcrossing rates up
424 to 8%⁵¹. Fiorina is an accession originating from Switzerland, acquired in 2015, specifically
425 for organic agriculture.

426 *Coriandrum sativum* (coriander) has a generally high genetic variability, with studies showing
427 up to 70.46% polymorphism, indicating the presence of high degree of molecular
428 variation in the studied coriander varieties^{58,59}. The variety that we used originally came
429 from an Indian market and was not a fixed variety, which ensured a minimum of genetic
430 variability. The flowers of coriander are self-incompatible but plants are self-compatible.

431 Geitonogamy is therefore common. Cross-pollination is facultative but can reach up to 20%⁶⁰.

432 *Lens culinaris* (lentil) is mainly self-pollinating; depending on the cultivar, outcrossing rates
433 reach between 1 and 5%⁵³.

434 *Camelina sativa* (camelina) is mainly self-pollinating, with outcrossing rates of less than
435 1%^{61,62}. In the study, we used a local landrace that was not a fixed variety.

Linum usitatissimum (flax) is mainly self-pollinating but outcrossing does occur, at a rate of 1-5%⁵⁴. Lirina, the variety of *Linum* that we used has been defined by ProSpecieRara as a rare or ancient variety. ProSpecieRara ensures the preservation of rare traditional varieties⁵⁵. Furthermore, studies have shown that linseed varieties have higher genetic variability than fiber flax and should therefore be considered as valuable genetic resources^{56,57}.

Experimental crop communities

Experimental communities consisted of single plots with one individual, monocultures, 2- and 4-species mixtures (Fig. 5). We planted every possible combination of 2-species mixtures with two species from different phylogenetic groups and every possible 4-species mixture with a species from each of the four different phylogenetic groups present (Table 2). We replicated the experiment two times with the exact same species composition, except for single individuals which were replicated 4 times. Monoculture and mixture plots were randomized among plots and beds within each fertilizer treatment, while single plant plots were randomly allocated to plots in separate beds in order to minimize interference among neighbouring plots. Each monoculture and mixture community consisted of one, two or four species planted in four rows. Two species mixtures were organized following a speciesA|speciesB|speciesA|speciesB pattern. The order of the species was chosen randomly. For 4-species mixtures, the order of the species was also randomized. Density of sowing differed among species groups and was based on current cultivation practices: 160 seeds/m² for legumes, 240 seeds/m² for superasterids, 400 seeds/m² for cereals, and 592 seeds/m² for superrosids. Each year, seeds were sown by hand in early April.

460

461

462 **Table 2.** List of species mixture combinations.
463

<i>Monoculture</i>	<i>2-species mixtures</i>	<i>2-species mixtures</i>	<i>4-species mixtures</i>
Avena	Avena-Lens	Lens- <i>Linum</i>	Avena-Lens- <i>Linum</i> -Coriandrum
Triticum	Avena- <i>Linum</i>	Lens-Camelina	Avena-Lens-Camelina-Coriandrum
Lens	Avena-Camelina	Lens-Coriandrum	Triticum-Lens- <i>Linum</i> -Coriandrum
<i>Linum</i>	Avena-Coriandrum	<i>Linum</i> -Coriandrum	Triticum-Lens-Camelina-Coriandrum
Camelina	Triticum-Lens	Camelina-Coriandrum	
Coriandrum	Triticum- <i>Linum</i>	Triticum-Coriandrum	
	Triticum-Camelina		

464

Figure 5: Pictures of the experimental plots. Top-left: part of the experimental garden, showing the plots within beds, and planted with single individuals. Top-right: a plot is outlined in red, showing a 2-species mixtures, with a cereal (wheat or oat) alternated with camelina. Bottom-left: a plot is outlined in red, showing a 2-species mixtures, with flax alternated with coriander.

Adaptation treatment

In 2019, we used the seeds collected in 2018 to add a coexistence history treatment: we repeated the experiment with seeds coming from single individuals, monocultures, and mixtures, respectively. This means that each plot described above was repeated three times: once with seeds coming from single plants, once with seeds coming from monoculture plants, and once with seeds coming from mixture plants. We respected the fertilizing treatment, i.e. there was a history treatment for each fertilizing condition. When planting the mixtures with a mixture history, we specifically used seeds coming from the same species combination. When planting the monocultures and singles with a mixture history, we used seeds coming from a common pool combining all 4-species mixtures.

In 2020, we repeated this process and selected seeds from 2019 to sow the single and community plots. We only selected seeds that had a “pure” history, i.e. that were always grown in the same coexistence history (for instance, for single history seeds in 2020 we selected only seeds that were grown as singles also in 2018 and 2019).

Data collection

Photosynthetically Active Radiation (PAR): Interception of PAR by the plant canopy was measured weekly with a LI-1500 (LI-COR Biosciences GmbH, Germany). In each plot, three PAR measurements were taken around noon by placing the sensor on the soil surface in the center of each of the three in-between rows. Light measurements beneath the canopy were compared to ambient radiation through simultaneous PAR measurements of a calibration sensor, which was mounted on a vertical post at 2 m above ground in the middle of the experimental garden. FPAR (%) indicates the percentage of PAR that was intercepted by the crop canopy.

Traits measurements: At the time of flowering, three individuals per crop species per plot were randomly marked. We measured the height of each individual with a ruler from the soil surface to the highest photosynthetically active tissue. We then measured plant width with a ruler by taking the largest horizontal distance between two photosynthetically active tissues. We sampled one healthy leaf from each marked individual and immediately wrapped this leaf in moist cotton; this was stored overnight at room temperature in open plastic bags. The following day, we removed any excess surface water on the leaf and weighed it to obtain its water saturated weight ⁶³. Then this leaf was scanned with a flatbed scanner (CanoScan LiDE 120, Canon), oven-dried in a paper envelope at 80°C for 72 hours, and subsequently reweighed to obtain its dry weight. We calculated Leaf Dry Matter Content (LDMC) as the ratio of leaf dry mass (g) to water saturated leaf mass (g). Using the leaf scans, we measured leaf area with the image processing software ImageJ ⁶⁴. Specific Leaf Area (SLA) was then calculated as the ratio of leaf area (cm²) to dry mass (g).

Plot grain yield and biomass: Grain yield and aboveground biomass of each crop species was determined per plot at maturity. This corresponded to July/August. As time of maturity slightly varied among the different crop species, we conducted harvest species by species. We clipped plants right above the soil surface and separated seeds from the vegetative parts. Seeds were sun-dried for five days and weighed. Biomass was oven-dried at 80 °C until constant weight and weighed.

Individual yield and biomass: We harvested the three marked individuals for the trait measurements separately; we separated seeds from aboveground biomass and they were both dried and weighed as previously mentioned. Furthermore, for each marked individual we weighed ten randomly selected seeds to obtain the mass per seed.

Data analyses

All analyses were performed using R version 4.1.0⁶⁵. **Plant Interaction Index:** Plant interaction intensity in the plots was calculated for each marked individual by means of the neighbor-effect intensity index with commutative symmetry NIntC⁶⁶:

$$NIntC = 2 \times \frac{yield_{comm} - yield_{single}}{yield_{comm} + yield_{single} + |yield_{comm} - yield_{single}|} \quad (1)$$

, where $yield_{single}$ is the yield of a single plant grown in isolation, and $yield_{comm}$ is the yield of an individual of the same species when grown in a community. NIntC values of all species (a, b, c, d) composing the community (i.e. species a in case of a monoculture and species a to d in case of a mixture of four species) were averaged and subsequently weighted by their proportional abundance $r_i = \frac{1}{\text{number of species}}$ to calculate the mean net interaction in the community (NIntCnet):

$$NIntCnet = \sum_{i=a}^d (NIntC_i r_i) \quad (2)$$

We then partitioned this net interaction index into its facilitation and competition components: NIntC facilitation was obtained subsetting those individuals with a positive NIntC value (i.e. with increased performance in communities compared to single plant individuals without neighbour interactions) and calculating the mean of all the species per plot weighed by their relative abundance; NIntC competition was obtained by subsetting those individuals with a negative NIntC value (i.e. with reduced performance in communities compared to single plant individuals without neighbour interactions) and calculating the mean of all the species per plot weighed by their relative abundance.

$$NIntC_{facil} = \sum_{i=a}^d (NIntC_i > 0 \times r_i) \quad (3)$$

$$NIntC_{comp} = \sum_{i=a}^d (NIntC_i < 0 \times r_i) \quad (4)$$

Net biodiversity effect: For all mixture communities we quantified the net biodiversity effect (NE) and its two components, the complementarity and selection effects according to Loreau and Hector⁸:

$$NE = N \cdot \overline{\Delta RY} \cdot \bar{M} + N \cdot cov(\Delta RY, M) \quad (5)$$

where N is the number of species in the plot, ΔRY is the deviation from expected relative yield of the species in mixture in the respective plot, which is calculated as the ratio of observed relative yield of the species in mixture to the yield of the species in monoculture, and M is the yield of the species in monoculture. The first component of the biodiversity effect equation ($N \cdot \overline{\Delta RY} \cdot \bar{M}$) is the complementarity effect (CE), while the second component ($N \cdot cov(\Delta RY, M)$) is the selection effect (SE).

Total crop yield: To assess crop performance, we calculated total crop yield per plot as the sum of total seed mass per species.

Trait analyses: Traits were analysed both at the species-level and at the plot-level. At the species level, we calculated the mean and coefficient of variation (CV) per species for each trait per plot. At the plot-level, we calculated Community-Weighted-Means (CMW) based on biomass per species, and coefficient of variation per plot for each trait.

Functional richness (FRic) was calculated in each plot using the function *dbFD* from the package *FD*⁶⁷, by measuring the convex hull volume occupied by the individuals of a plot in the space of the considered traits.

To analyze the effects of the experimental treatments on NIntCnet, NIntCfacil, NIntCcomp, NE, CE, LER, total crop yield, FRic, and CWM and CV per plot, we used generalized linear mixed models using the function *lmer*. Fixed factors included fertilizing condition (yes or no), coexistence history (considered as “same” or “different”), crop species number (2 vs 4) nested

in monoculture vs mixture, as well as the interactions between them. Species composition and bed were set as random factors. Effect sizes were calculated from marginal means obtained using the function *emmeans*, and pairwise comparisons were calculated using Tukey tests from the *emmeans* function⁶⁸. To analyze the effects of the experimental treatments on the mean and coefficient of variation of the different traits per species (height, width, SLA, LDMC, mass per seed, respectively), we used generalized linear mixed models using *lmer* with the same fixed factors as previously described. Species, species composition and bed were set as random factors. The response variables were log-transformed or square-root-transformed where needed. To analyse the response of FPAR, we used similar linear mixed models as described above, but added day of year as a random factor. For all models, we tested for normality of the residuals using a Shapiro-Wilk test and homogeneity of the variance using a Levene test.

Reference Methods

47. Roscher, C. *et al.* Overyielding in experimental grassland communities – irrespective of species pool or spatial scale. *Ecol. Lett.* **8**, 419–429 (2005).
48. Isbell, F. *et al.* Quantifying effects of biodiversity on ecosystem functioning across times and places. *Ecol. Lett.* **21**, 763–778 (2018).
49. Hai, L., Wagner, C. & Friedt, W. Quantitative structure analysis of genetic diversity among spring bread wheats (*Triticum aestivum* L.) from different geographical regions. *Genetica* **130**, 213–225 (2007).
50. Loureiro, I., Escorial, M. C., González, Á. & Chueca, M. C. Pollen-mediated gene flow in wheat (*Triticum aestivum* L.) in a semiarid field environment in Spain. *Transgenic Res.* **21**, 1329–1339 (2012).
51. Lawrie, R. G., Matus-Cádiz, M. A. & Hucl, P. Estimating Out-Crossing Rates in Spring Wheat Cultivars Using the Contact Method. *Crop Sci.* **46**, 247–249 (2006).
52. Shorter, R., Gibson, P. & Frey, K. J. Outscoring Rates in Oat Species Crosses (*Avena sativa* L. $\times$ *A. sterilis* L.)¹. *Crop Sci.* **18**, 877–878 (1978).
53. Horneburg, B. & Weber, W. E. Outcrossing in lentil (*Lens culinaris*) depends on cultivar,

596 location and year, and varies within cultivars. *Plant Breed.* **125**, 638–640 (2006).

597 54. Jhala, A. J., Bhatt, H., Topinka, K. & Hall, L. M. Pollen-mediated gene flow in flax (*Linum*
598 *usitatissimum* L.): can genetically engineered and organic flax coexist? *Heredity (Edinb)*. **106**,
599 557 (2011).

600 55. BLAUES WUNDER, LIN » Acheter des semences bio en ligne / Engrais verts - SATIVA
601 Online Shop. https://www.sativa.bio/fr_ch/engrais-verts/blaues-wunder-lin-prospecierara-gr31.

602 56. Vromans, J., van de Bilt, E., Pieper, B., Stam, P. & van Eck, H. J. Molecular genetic studies in
603 flax. (2006).

604 57. Hoque, A., Fiedler, J. D. & Rahman, M. Genetic diversity analysis of a flax (*Linum*
605 *usitatissimum* L.) global collection. *BMC Genomics* **21**, 1–13 (2020).

606 58. Choudhary, S., Sharma, R., Nehru, J., Vishwavidyalaya, K. & Vishal, M. K. Molecular
607 diversity in coriander (*Coriandrum sativum*) using RAPD and ISSR markers. *Indian J. Agric.*
608 *Sci.* **89**, 193–198 (2019).

609 59. Singh, S. K. *et al.* Studies on genetic divergence among Indian varieties of a spice herb,
610 *Coriandrum sativum*. *J. Environ. Biol.* **33**, 781–789 (2012).

611 60. Diederichsen, A. *Coriander (Coriandrum sativum L.). Promoting the conservation and use of*
612 *underutilized and neglected crops.* (1996).

613 61. Walsh, K. D. *et al.* Short communication: First report of outcrossing rates in camelina
614 [*camelina sativa* (L.) crantz], a potential platform for bioindustrial oils. *Can. J. Plant Sci.* **92**,
615 681–685 (2012).

616 62. Walsh, K. D., Hills, M. J., Martin, S. L. & Hall, L. M. Pollen-mediated Gene Flow in Camelina
617 *sativa* (L.) Crantz. *Crop Sci.* **55**, 196–202 (2015).

618 63. Cornelissen, J. H. C. *et al.* A handbook of protocols for standardised and easy measurement of
619 plant functional traits worldwide. *Aust. J. Bot.* **51**, 335–380 (2003).

620 64. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image
621 analysis. *Nature Methods* vol. 9 671–675 (2012).

622 65. R Core Team. A language and environment for statistical computing. Vienna, Austria: R
623 Foundation for Statistical Computing. (2019).

624 66. Díaz-Sierra, R., Verwijmeren, M., Rietkerk, M., de Dios, V. R. & Baudena, M. A new family
625 of standardized and symmetric indices for measuring the intensity and importance of plant
626 neighbour effects. *Methods in Ecology and Evolution* vol. 8 580–591 (2017).

627 67. Laliberte, E. & Legendre, P. A distance-based framework for measuring functional diversity

from multiple traits. *Ecology* **91**, 299–305 (2010).

68. Lenth, R. V. emmeans: Estimated Marginal Means, aka Least-Squares Means. (2021)
doi:10.1080/00031305.1980.10483031.

Acknowledgements

We thank Elisa Pizarro Carbonell, Carlos Barriga Cabanillas, Anja Schmutz, Sandra Gonzalez Sanchez, Lukas Meile, Carlos Federico Ingala, Roman Hüppi, Simon Baumgartner, Benjamin Wilde, Manon Longepierre, Marijn Van de Broek, Leonhard Späth, Inea Lehner, Anna Bugmann, Jianguo Chen, Nicola Haggenmacher and Zita Sartori for their help in the field, and Johan Six for comments on the experimental design. We also thank the Aprisco de Las Corchuelas Field Station and the University of Zurich for the use of their facilities. The study was funded by the Swiss National Science Foundation (PP00P3_170645).

Author contributions

LS, NE, and CS conceptualised the study; LS and CS designed the experiment; LS, NE, and CS carried out the experiment, LS and CS analysed the data; LS and CS wrote the paper with input from NE.

Competing interests

The authors declare no competing financial interests.

Materials & Correspondence

Correspondence and requests for materials should be addressed to Laura Stefan.

Data availability statement

The data that support the findings of this study are available on Zenodo:
<https://doi.org/10.5281/zenodo.5223410>

651 **Code availability statement**

652 The R code is available on Zenodo: <https://doi.org/10.5281/zenodo.5223410>

653

654 Extended Data

Extended Data Figure 1: Effects of coexistence history and crop species number on competition (a), and facilitation (b) indexes, for fertilized and unfertilized conditions. “Same coexistence history” indicates that crops were grown in the community their seeds were collected from. “Different coexistence history” refers to crops grown in a community different to the one of their parents. See methods for the index calculations.

Horizontal lines represent the median of the data, boxes represent the lower and upper quartiles (25% and 75%), with vertical lines extending from the hinge of the box to the smallest and largest values, no further than 1.5 * the interquartile range. Data beyond the end of the whiskers are outlying and plotted individually. n =276

655

Extended Data Table 1. Type-I Analysis of Variance table of the experimental treatment effects on net, competition and facilitation indexes, in 2020

DenDF, degrees of freedom of error term; NumDF, degrees of freedom of term; F-value, variance ratio; Pr(>F), error probability. P-values in bold are significant at $\alpha = 0.05$; * (P < 0.05), ** (P < 0.01), *** (P < 0.001). n =276

	NumDF	Net			Competition			Facilitation		
		DenDF	F value	Pr(>F)	DenDF	F value	Pr(>F)	DenDF	F value	Pr(>F)
Fertilizer	1	7.02	36.269	0.0005 ***	6.82	18.658	0.0037 **	6.74	39.646	0.0005 ***
History	1	238.10	31.901	4.61E-08 ***	232.57	9.071	0.0029 **	236.53	38.318	2.63E-09 ***
Monocultures vs. mixtures	1	19.99	0.571	0.4587	20.00	0.343	0.5647	19.97	0.945	0.3425
Diversity	1	19.97	0.110	0.7438	20.00	0.084	0.7756	19.90	0.159	0.6942
Fertilizer x history	1	237.87	0.205	0.6509	232.63	0.279	0.5982	236.49	0.921	0.3383
Fertilizer x mono vs. mix	1	242.37	0.695	0.4054	243.37	0.036	0.8493	242.65	1.269	0.2611
Fertilizer x diversity	1	240.31	0.305	0.5816	241.70	0.075	0.7852	240.70	0.374	0.5415
History x mono vs. mix	1	240.38	1.319	0.2518	241.43	0.253	0.6152	240.79	2.149	0.1440
History x diversity	1	240.50	0.912	0.3405	241.93	0.279	0.5977	240.88	1.111	0.2928
Fertilizer x history x mono vs. mix	1	240.34	0.022	0.8836	241.39	0.000	0.9864	240.75	0.003	0.9567
Fertilizer x history x diversity	1	240.42	0.100	0.7523	241.89	0.296	0.5868	240.81	0.003	0.9542

656

Extended Data Table 2. Pairwise comparisons of the effect of net interaction index between fertilizer (yes, no), coexistence history (diff [different], same), and monoculture vs mixture (mix [mixture], mono [monoculture]).

Net interaction index	estimate	SE	df	t.ratio	p.value
no diff mix - yes diff mix	2.54E-01	0.0604	33.1	4.206	0.0041
no diff mix - no same mix	-2.73E-01	0.0705	237.2	-3.877	0.0034
no diff mix - yes same mix	2.33E-02	0.0725	60.5	0.321	1
no diff mix - no diff mono	1.83E-01	0.2298	22.8	0.796	0.9917
no diff mix - yes diff mono	3.58E-01	0.2303	23	1.553	0.7718
no diff mix - no same mono	2.34E-02	0.2365	25.5	0.099	1
no diff mix - yes same mono	2.48E-01	0.2371	25.9	1.047	0.9621
yes diff mix - no same mix	-5.28E-01	0.0729	57.2	-7.233	<.0001
yes diff mix - yes same mix	-2.31E-01	0.0694	242	-3.328	0.0222
yes diff mix - no diff mono	-7.13E-02	0.2308	23.1	-0.309	1
yes diff mix - yes diff mono	1.04E-01	0.2293	22.7	0.452	0.9998
yes diff mix - no same mono	-2.31E-01	0.2372	25.9	-0.973	0.9744
yes diff mix - yes same mono	-5.87E-03	0.2362	25.5	-0.025	1
no same mix - yes same mix	2.97E-01	0.0832	84.3	3.565	0.0134
no same mix - no diff mono	4.56E-01	0.2336	24.3	1.954	0.5303
no same mix - yes diff mono	6.31E-01	0.2339	24.4	2.698	0.1706
no same mix - no same mono	2.97E-01	0.2395	26.9	1.24	0.9124
no same mix - yes same mono	5.22E-01	0.2406	27.3	2.168	0.4002
yes same mix - no diff mono	1.60E-01	0.2342	24.5	0.681	0.9968
yes same mix - yes diff mono	3.34E-01	0.2329	24.1	1.436	0.832
yes same mix - no same mono	9.22E-05	0.2406	27.3	0	1
yes same mix - yes same mono	2.25E-01	0.2395	26.9	0.939	0.979
no diff mono - yes diff mono	1.75E-01	0.0848	90.2	2.062	0.4474
no diff mono - no same mono	-1.59E-01	0.1	242.7	-1.595	0.7527
no diff mono - yes same mono	6.54E-02	0.1018	144.2	0.642	0.9982
yes diff mono - no same mono	-3.34E-01	0.101	144.6	-3.31	0.0253
yes diff mono - yes same mono	-1.09E-01	0.0984	238.1	-1.111	0.9539
no same mono - yes same mono	2.25E-01	0.1157	182.9	1.944	0.5228

Extended Data Figure 2: Effects of coexistence history and crop species number on complementarity effect (a) and selection effect (b) in fertilized and unfertilized plots.

Horizontal lines represent the median of the data, boxes represent the lower and upper quartiles (25% and 75%), with vertical lines extending from the hinge of the box to the smallest and largest values, no further than 1.5 * the interquartile range. Data beyond the end of the whiskers are outlying and plotted individually. n=204

659

Extended Data Table 3. Type-I Analysis of Variance table of the experimental treatment effects on net, complementarity, and selection effects in 2020

DenDF, degrees of freedom of error term; *NumDF*, degrees of freedom of term; *F-value*, variance ratio; *Pr(>F)*, error probability. P-values in bold are significant at $\alpha = 0.1$; . ($P < 0.1$); * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$). n=204

	<i>NumDF</i>	<i>Net effect</i>			<i>Complementarity effect</i>			<i>Selection effect</i>		
		<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>
<i>Fertilizer</i>	1	7.64	1.005	0.3468	7.38	2.684	0.14312	7.69	1.295	0.2894
<i>History</i>	1	177.34	0.162	0.6876	179.89	1.567	0.2123	178.62	1.512	0.2204
<i>Diversity</i>	1	14.90	2.583	0.1290	14.87	5.887	0.0285 *	15.06	4.009	0.0636 .
<i>Fertilizer x history</i>	1	177.08	9.595	0.0023 **	179.80	2.719	0.1009	178.66	2.498	0.1158
<i>Fertilizer x diversity</i>	1	173.48	0.197	0.6581	174.92	5.579	0.0193 *	178.22	6.092	0.0145 *
<i>History x diversity</i>	1	173.66	0.052	0.8191	175.06	1.399	0.2385	178.42	3.165	0.0769 .
<i>Fertilizer x history x diversity</i>	1	173.36	0.046	0.8297	174.85	2.026	0.1564	178.35	4.093	0.0446 *

660

661

662

663

664

665

666

Extended Data Table 4. Type-I Analysis of Variance table of the experimental treatment effects on total crop yield per plot (square-root transformed)

DenDF, degrees of freedom of error term; *NumDF*, degrees of freedom of term; *F-value*, variance ratio; *Pr(>F)*, error probability. P-values in bold are significant at $\alpha = 0.1$; . ($P < 0.1$); * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$). n=276

	<i>NumDF</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>	
<i>Fertilizer</i>	1	7.682	18.5184	0.002862 **	
<i>History</i>	1	241.392	0.1812	0.670703	
<i>Mono vs. mixtures</i>	1	19.973	3.5836	0.072934 .	
<i>Diversity</i>	1	19.956	0.5134	0.481957	
<i>Fertilizer x history</i>	1	241.338	0.0003	0.986084	
<i>Fertilizer x mono vs. mix</i>	1	238.145	0.2223	0.637735	
<i>Fertilizer x diversity</i>	1	237.548	0.0887	0.766152	
<i>History x mono vs. mix</i>	1	237.451	0.0013	0.97136	
<i>History x diversity</i>	1	237.596	0.0991	0.753181	
<i>Fertilizer x history x mono vs. mix</i>	1	237.347	0.0326	0.856797	
<i>Fertilizer x history x diversity</i>	1	237.487	0.0385	0.844517

Extended Data Figure 3: Effects of coexistence history of total yield per plot, per species combination. “Same coexistence history” indicates that crops were grown in the community their seeds were collected from. “Different coexistence history” refers to crops grown in a community different to the one of their parents.

Extended Data Figure 5: Effects of coexistence history and crop species number on mean height (in cm) (a) and LDMC (b). Dots represent the averaged values across species and plots; lines represent the standard error. $n = 1726$

Extended Data Figure 6: Mean height (cm) (a) and LDMC (b) according to their coexistence history, for the six species considered in our study. Dots represent the averaged values across species and plots; lines represent the standard error. $n = 1726$

Extended Data Table 5. Type-I Analysis of Variance table of the experimental treatment effects on mean and coefficient of variation of height, per species per plot (species level)

DenDF, degrees of freedom of error term; *NumDF*, degrees of freedom of term; *F-value*, variance ratio; *Pr(>F)*, error probability. P-values in bold are significant at $\alpha = 0.05$; * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$), n=1726

	Mean				Coefficient of variation		
	<i>NumDF</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>
<i>Fertilizer</i>	1	7.9	0.2292	0.645078	7.04	16.8065	4.51E-03 **
<i>History</i>	1	169.16	4.2929	0.039789 *	197.26	0.077	0.781684
<i>Mono vs. mixtures</i>	1	23.57	0.1617	0.691188	45.01	0.325	0.571472
<i>Diversity</i>	1	10.68	0.121	0.73467	45.48	0.2586	0.61356
<i>Fertilizer x history</i>	1	168.93	0.1986	0.656442	197.18	5.8068	0.016883 *
<i>Fertilizer x mono vs. mix</i>	1	416.7	9.2129	0.002554 **	487.47	5.0102	0.025648 *
<i>Fertilizer x diversity</i>	1	118.78	0.0013	0.971368	127.23	0.0009	0.976698
<i>History x mono vs. mix</i>	1	418.29	1.435	0.231632	483.31	0.2095	0.647359
<i>History x diversity</i>	1	119.33	0.6601	0.418149	128.24	0.1385	0.710429
<i>Fertilizer x history x mono vs. mix</i>	1	417.97	0.0046	0.945955	483.3	0.5825	0.445694
<i>Fertilizer x history x diversity</i>	1	119.15	0.065	0.79923	128.25	0.0665	0.796946

699

700

Extended Data Table 6. Type-I Analysis of Variance table of the experimental treatment effects on mean and coefficient of variation of width, per species per plot (species level)

DenDF, degrees of freedom of error term; *NumDF*, degrees of freedom of term; *F-value*, variance ratio; *Pr(>F)*, error probability. P-values in bold are significant at $\alpha = 0.1$; . ($P < 0.1$); * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$), n=1726

	Mean				Coefficient of variation		
	<i>NumDF</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>
<i>Fertilizer</i>	1	7.59	8.5571	0.02027 *	7.34	4.8793	6.12E-02
<i>History</i>	1	531.5	2.0724	0.15057	233.05	2.5024	0.11503
<i>Mono vs. mixtures</i>	1	19.68	0.9369	0.34482	534.07	0.0141	0.90552
<i>Diversity</i>	1	10.47	0.0639	0.80529	144.7	0.8817	0.3493
<i>Fertilizer x history</i>	1	530.95	1.6511	0.19937	232.84	0.0033	0.9541
<i>Fertilizer x mono vs. mix</i>	1	523.5	2.5201	0.11301	533.17	0.7796	0.37767
<i>Fertilizer x diversity</i>	1	526.73	3.905	0.04866 *	142.23	0.2794	0.59791
<i>History x mono vs. mix</i>	1	522.6	0.1858	0.66657	528.21	1.4807	0.2242
<i>History x diversity</i>	1	526.98	1.1295	0.28837	144.44	1.1028	0.2954
<i>Fertilizer x history x mono vs. mix</i>	1	522.34	0.054	0.81631	528.23	1.7197	0.1903
<i>Fertilizer x history x diversity</i>	1	526.26	0.9072	0.3413	144.54	0.8736	0.35153

701

702

703

704

Extended Data Table 7. Type-I Analysis of Variance table of the experimental treatment effects on mean and coefficient of variation of SLA, per species per plot (species level)

DenDF, degrees of freedom of error term; *NumDF*, degrees of freedom of term; *F-value*, variance ratio; *Pr(>F)*, error probability. P-values in bold are significant at $\alpha = 0.1$; . ($P < 0.1$); * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$), n=1726

	Mean				Coefficient of variation		
	<i>NumDF</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>
<i>Fertilizer</i>	1	7.8	9.3229	0.01623 *	7.7	7.4375	0.026891 *
<i>History</i>	1	203.35	0.0678	0.79477	224.44	0.141	0.707662
<i>Mono vs. mixtures</i>	1	27.81	2.1781	0.15122	39.2	1.8674	0.179558
<i>Diversity</i>	1	10.76	0.0715	0.79425	12.52	3.0841	0.103467
<i>Fertilizer x history</i>	1	203.8	0.2212	0.6386	224.52	0.0117	0.914055
<i>Fertilizer x mono vs. mix</i>	1	428.56	0.0664	0.79679	415.79	7.1378	0.007844 **
<i>Fertilizer x diversity</i>	1	144.51	0.2563	0.61342	168.99	3.165	0.077029 .
<i>History x mono vs. mix</i>	1	427.99	2.4804	0.11601	412.72	0.0641	0.80022
<i>History x diversity</i>	1	146.35	1.0093	0.31674	169.58	0.0099	0.920833
<i>Fertilizer x history x mono vs. mix</i>	1	427.9	0.0372	0.84708	412.49	0.2757	0.599836
<i>Fertilizer x history x diversity</i>	1	146.35	0.0089	0.92507	169.62	0.1557	0.693661

Extended Data Table 8. Type-I Analysis of Variance table of the experimental treatment effects on mean and coefficient of variation of LDMC, per species per plot (species level)

DenDF, degrees of freedom of error term; *NumDF*, degrees of freedom of term; *F-value*, variance ratio; *Pr(>F)*, error probability. P-values in bold are significant at $\alpha = 0.1$; . ($P < 0.1$); * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$), n=1726

	Mean				Coefficient of variation		
	<i>NumDF</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>
<i>Fertilizer</i>	1	7.86	8.0352	0.02239 *	186.4	12.783	4.46E-04 ***
<i>History</i>	1	183.84	3.5956	0.05950 .	190.46	0.0447	0.832786
<i>Mono vs. mixtures</i>	1	19.67	0.1589	0.69442	516.86	0.0353	0.851015
<i>Diversity</i>	1	5.24	0.1083	0.75481	116.92	0.6369	0.426449
<i>Fertilizer x history</i>	1	181.59	0.0432	0.83566	190.48	3.1314	0.078397 .
<i>Fertilizer x mono vs. mix</i>	1	467.36	0.0294	0.86399	515.71	0.6107	0.434889
<i>Fertilizer x diversity</i>	1	115.57	0.3376	0.56232	114.6	3.8577	0.051939 .
<i>History x mono vs. mix</i>	1	468.83	0.0043	0.94775	515.4	0.2779	0.598296
<i>History x diversity</i>	1	116.59	1.0418	0.30953	114.82	0.9337	0.335923
<i>Fertilizer x history x mono vs. mix</i>	1	468.38	0.0166	0.89752	515.36	1.6235	0.203176
<i>Fertilizer x history x diversity</i>	1	116.48	0.0016	0.96772	114.88	1.3649	0.24511

705

706

707

708

709

Extended Data Table 9. Type-I Analysis of Variance table of the experimental treatment effects on mean and coefficient of variation of mass per seed, per species per plot (species level)

DenDF, degrees of freedom of error term; *NumDF*, degrees of freedom of term; *F-value*, variance ratio; *Pr(>F)*, error probability. P-values in bold are significant at $\alpha = 0.1$; . ($P < 0.1$); * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$), n=1726

	<i>Mean</i>				<i>Coefficient of variation</i>		
	<i>NumDF</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>
<i>Fertilizer</i>	1	210.65	4.3651	0.03788 *	180.24	0.1253	7.24E-01
<i>History</i>	1	211.89	0.0491	0.82487	182.21	0.0853	0.770551
<i>Mono vs. mixtures</i>	1	20.08	10.0297	0.00483 **	33.75	17.0854	0.000223 ***
<i>Diversity</i>	1	10.77	1.3367	0.27261	8.88	0.1203	0.736758
<i>Fertilizer x history</i>	1	211.88	0.6633	0.4163	182.12	1.6601	0.199224
<i>Fertilizer x mono vs. mix</i>	1	493.44	0.602	0.43818	474.87	1.16	0.282008
<i>Fertilizer x diversity</i>	1	137.39	0.3993	0.52848	118.13	1.4557	0.230024
<i>History x mono vs. mix</i>	1	493.59	1.4337	0.23174	470.19	4.9519	0.026536 *
<i>History x diversity</i>	1	138.08	0.2159	0.64289	119.33	0.2756	0.600592
<i>Fertilizer x history x mono vs. mix</i>	1	493.59	2.7072	0.10053	470.29	0.0587	0.808609
<i>Fertilizer x history x diversity</i>	1	138.1	0.6308	0.42842	119.29	0.0329	0.856454

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

Extended Data Table 10. Type-I Analysis of Variance table of the experimental treatment effects on community-weighted mean and coefficient of variation of height, per plot (community level)

DenDF, degrees of freedom of error term; *NumDF*, degrees of freedom of term; *F-value*, variance ratio; *Pr(>F)*, error probability. P-values in bold are significant at $\alpha = 0.1$; . ($P < 0.1$); * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$). n=271

	<i>CWM</i>				<i>Coefficient of variation</i>		
	<i>NumDF</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>
<i>Fertilizer</i>	1	7.804	2.687	1.41E-01	6.983	7.6601	2.79E-02 *
<i>History</i>	1	231.321	0.5067	0.47728	234.457	3.9279	0.04866 *
<i>Mono vs. mixtures</i>	1	19.976	0.0555	0.81612	20.095	16.9763	0.000527 ***
<i>Diversity</i>	1	19.985	0.1513	0.70145	19.944	3.0898	0.094124 .
<i>Fertilizer x history</i>	1	228.963	0.0912	0.76295	234.705	0.6664	0.415143
<i>Fertilizer x mono vs. mix</i>	1	227.04	5.6668	0.01812 *	238.629	5.7944	0.016838 *
<i>Fertilizer x diversity</i>	1	226.105	0.7338	0.39255	235.462	0.0078	0.929608
<i>History x mono vs. mix</i>	1	226.029	0.4327	0.51136	235.769	2.0403	0.154507
<i>History x diversity</i>	1	226.301	0.1589	0.69052	235.616	0.1045	0.746762
<i>Fertilizer x history x mono vs. mix</i>	1	226.108	0.0055	0.9408	236.013	0.0007	0.97901
<i>Fertilizer x history x diversity</i>	1	226.151	0.0112	0.91576	235.639	0.3324	0.564822

Extended Data Table 11. Type-I Analysis of Variance table of the experimental treatment effects on community-weighted mean and coefficient of variation of width, per plot (community level)

DenDF, degrees of freedom of error term; *NumDF*, degrees of freedom of term; *F-value*, variance ratio; *Pr(>F)*, error probability. P-values in bold are significant at $\alpha = 0.1$; . ($P < 0.1$); * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$), n =271

	<i>CWM</i>				<i>Coefficient of variation</i>		
	<i>NumDF</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>
<i>Fertilizer</i>	1	7.484	6.0869	4.09E-02 *	7.352	10.7862	1.25E-02 *
<i>History</i>	1	233.262	0.001	0.974539	239.502	0.2976	0.585869
<i>Mono vs. mixtures</i>	1	19.917	1.7012	0.207003	20.153	15.6397	0.000773 ***
<i>Diversity</i>	1	19.935	0.383	0.543019	19.819	2.4097	0.13641
<i>Fertilizer x history</i>	1	231.77	1.6122	0.205462	239.1	0.0345	0.852701
<i>Fertilizer x mono vs. mix</i>	1	228.97	11.0067	0.001056 **	237.11	8.7687	0.003376 **
<i>Fertilizer x diversity</i>	1	227.323	3.9302	0.048631 *	233.942	1.1993	0.274588
<i>History x mono vs. mix</i>	1	227.213	0.1337	0.714952	234.231	4.7012	0.031149 *
<i>History x diversity</i>	1	227.671	0.1338	0.714862	234.017	0.029	0.864848
<i>Fertilizer x history x mono vs. mix</i>	1	227.362	0.391	0.532421	234.403	2.0867	0.149924
<i>Fertilizer x history x diversity</i>	1	227.418	0.3123	0.576817	233.993	0.6311	0.427744

727

728

729

730

731

Extended Data Table 12. Type-I Analysis of Variance table of the experimental treatment effects on community-weighted mean and coefficient of variation of SLA, per plot (community level)

DenDF, degrees of freedom of error term; *NumDF*, degrees of freedom of term; *F-value*, variance ratio; *Pr(>F)*, error probability. P-values in bold are significant at $\alpha = 0.1$; . ($P < 0.1$); * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$), n=271

	<i>CWM</i>				<i>Coefficient of variation</i>		
	<i>NumDF</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>
<i>Fertilizer</i>	1	7.612	0.8181	3.93E-01	7.426	3.6336	9.59E-02 .
<i>History</i>	1	227.776	2.0837	0.15025	225.431	2.4299	0.120442
<i>Mono vs. mixtures</i>	1	19.754	0.0616	0.80648	19.682	15.332	0.000879 ***
<i>Diversity</i>	1	19.858	0.0067	0.93546	20.339	8.4912	0.008482 **
<i>Fertilizer x history</i>	1	227.607	0.0158	0.89994	224.689	0.8382	0.360883
<i>Fertilizer x mono vs. mix</i>	1	225.127	0.4507	0.50268	222.442	7.7108	0.005957 **
<i>Fertilizer x diversity</i>	1	223.255	0.3865	0.53476	222.197	3.5889	0.059465 .
<i>History x mono vs. mix</i>	1	222.96	3.4724	0.06371 .	220.028	0.1602	0.689401
<i>History x diversity</i>	1	223.809	0.2463	0.62019	222.407	0.1505	0.698459
<i>Fertilizer x history x mono vs. mix</i>	1	223.168	0.2017	0.65379	220.233	0.5343	0.465566
<i>Fertilizer x history x diversity</i>	1	223.828	0.0913	0.76284	221.495	0.0051	0.943381

732

733

Extended Data Table 13. Type-I Analysis of Variance table of the experimental treatment effects on community-weighted mean and coefficient of variation of LDMC, per plot (community level)

DenDF, degrees of freedom of error term; *NumDF*, degrees of freedom of term; *F-value*, variance ratio; *Pr(>F)*, error probability. P-values in bold are significant at $\alpha = 0.1$; . ($P < 0.1$); * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$), n=271

	<i>CWM</i>				<i>Coefficient of variation</i>		
	<i>NumDF</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>
<i>Fertilizer</i>	1	7.809	10.5893	1.20E-02 *	7.368	2.1001	1.88E-01
<i>History</i>	1	225.249	4.3323	0.03853 *	233.338	4.1789	0.042053 *
<i>Mono vs. mixtures</i>	1	19.998	0.0614	0.80688	20.123	10.747	0.003737 **
<i>Diversity</i>	1	20.012	0.0293	0.86579	19.46	0.7439	0.39891
<i>Fertilizer x history</i>	1	223.935	0.598	0.44016	233.38	0.9977	0.318908
<i>Fertilizer x mono vs. mix</i>	1	222.756	0.3654	0.54615	236.325	4.6325	0.032385 *
<i>Fertilizer x diversity</i>	1	221.841	0.0235	0.87843	232.776	0.0154	0.901256
<i>History x mono vs. mix</i>	1	221.728	0.1477	0.70108	233.192	0.4468	0.504525
<i>History x diversity</i>	1	222.051	0.0037	0.95178	233	0.3198	0.572283
<i>Fertilizer x history x mono vs. mix</i>	1	221.903	0.1505	0.69841	233.579	2.708	0.101189
<i>Fertilizer x history x diversity</i>	1	221.981	0.3306	0.56587	233.206	0.8702	0.351866

734

735

736

737

Extended Data Table 14. Type-I Analysis of Variance table of the experimental treatment effects on community-weighted mean and coefficient of variation of mass per seed, per plot (community level)

DenDF, degrees of freedom of error term; *NumDF*, degrees of freedom of term; *F-value*, variance ratio; *Pr(>F)*, error probability. P-values in bold are significant at $\alpha = 0.1$; . ($P < 0.1$); * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$), n=271

	<i>CWM</i>				<i>Coefficient of variation</i>		
	<i>NumDF</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>
<i>Fertilizer</i>	1	7.319	2.4734	0.15792	6.612	1.0304	3.46E-01
<i>History</i>	1	240.762	0.3199	0.57217	225.443	1.4213	0.234444
<i>Mono vs. mixtures</i>	1	20.012	0.1269	0.72535	20.021	14.3694	0.001145 **
<i>Diversity</i>	1	19.996	0.004	0.9499	19.95	0.5302	0.474995
<i>Fertilizer x history</i>	1	235.451	0.0844	0.77174	226.003	0.0292	0.864583
<i>Fertilizer x mono vs. mix</i>	1	238.14	0.9245	0.33727	239.091	0.1291	0.719721
<i>Fertilizer x diversity</i>	1	235.395	0.3647	0.54651	236.191	0.2304	0.631662
<i>History x mono vs. mix</i>	1	235.632	1.5019	0.22161	236.507	5.4785	0.020084 *
<i>History x diversity</i>	1	235.537	0.0009	0.9767	236.403	0.1301	0.718673
<i>Fertilizer x history x mono vs. mix</i>	1	235.858	3.5455	0.06094 .	236.836	0.3515	0.553841
<i>Fertilizer x history x diversity</i>	1	235.548	0.3137	0.57596	236.483	0.1133	0.736717

738

739

740

741

742

743

744

745

746

747

748

749

750

751

Extended Data Figure 7: Functional richness in response to crop species diversity, in fertilized and unfertilized plots. N=271

Extended Data Table 15. Type-I Analysis of Variance table of the experimental treatment effects on functional richness

DenDF, degrees of freedom of error term; *NumDF*, degrees of freedom of term; *F-value*, variance ratio; *Pr(>F)*, error probability. P-values in bold are significant at $\alpha = 0.1$; . ($P < 0.1$); * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$). n=271

	<i>NumDF</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>
<i>Fertilizer</i>	1	240.322	12.1182	0.000593 ***
<i>History</i>	1	239.331	0.4763	0.490791
<i>Diversity</i>	2	20.095	224.893	1.76E-14 ***
<i>Fertilizer x history</i>	1	239.229	0.3573	0.550593
<i>Fertilizer x diversity</i>	2	240.165	1.6086	0.202324
<i>History x diversity</i>	2	239.277	0.1769	0.837967
<i>Fertilizer x history x diversity</i>	2	239.303	0.5828	0.559107

Extended Data Figure 8: Fraction of PAR absorbed (in %) according to the day of year, for plants with the same or different coexistence history. The lines represent local polynomial regression fittings, with the grey area representing the 0.95 confidence interval. n=2484.

Extended Data Table 16. Type-I Analysis of Variance table of the experimental treatment effects on FPAR

DenDF, degrees of freedom of error term; *NumDF*, degrees of freedom of term; *F-value*, variance ratio; *Pr(>F)*, error probability. P-values in bold are significant at $\alpha = 0.05$; * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$). n=2484

	<i>NumDF</i>	<i>DenDF</i>	<i>F value</i>	<i>Pr(>F)</i>
<i>Fertilizer</i>	1	7.76	18.986 15.396	0.002604 **
<i>History</i>	1	2420.53	2	8.96E-05 ***
<i>Mono vs. mixtures</i>	1	20.03	1.3841	0.253196
<i>Diversity</i>	1	19.98	0.4482	0.510868
<i>Fertilizer x history</i>	1	2421.07	1.0275	0.310837
<i>Fertilizer x mono vs. mix</i>	1	2445.01	2.2105	0.137204
<i>Fertilizer x diversity</i>	1	2443.87	0.0309	0.86054
<i>History x mono vs. mix</i>	1	2443.67	0.4901	0.483969
<i>History x diversity</i>	1	2444.07	0.0881	0.7666
<i>Fertilizer x history x mono vs. mix</i>	1	2442.98	0.0041	0.948804
<i>Fertilizer x history x diversity</i>	1	2443.39	0.1576	0.691409