Structural insights into the galanin receptors signaling

Authors and Affiliations:

Wentong Jiang^{1,2}, Sanduo Zheng^{1,3*}

¹National Institute of Biological Sciences, Beijing, China

²Graduate School of Peking Union Medical College, Beijing, China

³Tsinghua Institute of Multidisciplinary Biomedical Research, Tsinghua University, Beijing, China

Abstract

Galanin is a biologically active neuropeptide, and functions through three distinct G protein-coupled receptors (GPCRs), namely GALR1, GALR2 and GLAR3. GALR signaling plays important roles in regulating various physiological processes such as energy metabolism, neuropathic pain, epileptic activity, and sleep homeostasis. GALR1 and GALR3 signal through the $G_{i/o}$ pathway, whereas GALR2 signals mainly through the $G_{q/11}$ pathway. However, the molecular basis for galanin recognition and G protein selectivity of GALRs remains poorly understood. Here, we report the cryoelectron microscopy structures of the GALR1- G_o and the GALR2- G_q complexes bound to the endogenous ligand galanin or spexin. The galanin peptide mainly adopts an alpha helical structure, which binds at the extracellular vestibule of the receptors, nearly parallel to the membrane plane without penetrating deeply into the receptor core. Structural analysis combined with functional studies reveals important structural determinants for the G protein selectivity of GALRs as well as other class A GPCRs. In addition, we show that the zinc ion is a negative allosteric regulator of GALR1 but not GALR2. Our studies provide insight into the mechanisms of G protein selectivity of GPCRs and highlight potential novel function of the neuromodulator zinc ion as a modulator of GPCR signaling in the central nervous system.

Significance Statement

Galanin exerts various physiological functions through galanin receptors, including antinociceptive activity, depression and sleep. Here, we reveal a distinct binding site and binding pose of galanin peptide in galanin receptors from that of the published structures of peptide-bound GPCRs. Moreover, our work show that the neuromodulator zinc ion negatively modulates galanin signaling in the central nervous system, and further advances our understanding of mechanisms of G protein selectivity of GPCRs. These unique features of galanin receptors can be exploited for rational design of subtype selective ligands for treatments of neurological disorders.

Introduction

Galanin is a 29 or 30-amino-acid peptide that was isolated from pig intestine in 1983 (1). Through its wide distribution in the nervous system and the endocrine system, galanin is involved in a variety of physiological functions, including regulation of hormones and neurotransmitters release, antinociceptive activity, depression and sleep/wake homeostasis (2, 3). The endogenous action of galanin is mediated through activation of three distinct receptor subtypes (GALR1-3), which belong to the class A of G protein-coupled receptors (GPCRs) family (4, 5).

GALR subtypes vary in their downstream signaling pathways and the tissue distribution. GALR1 and GLAR3 mainly couple to the inhibitory $G_{\alpha_{i/o}}$ pathway, leading to the inhibition of the adenylyl cyclase activity and the decrease of the intracellular adenosine 3',5'-cyclic monophosphate (cAMP) level. By contrast, GALR2 mainly couples to the stimulatory pathway of $G_{q/11}$, inducing the formation of inositol triphosphate (IP3), which in turn increases the cytosolic Ca^{2+} level (3) (**Fig. 1A**). While GALR1 is particularly enriched in the nervous system, GALR2 and GALR3 are broadly distributed in brain as well as peripheral tissues. GALRs activation via overexpression or administration of galanin in the nervous system of animals suppresses seizure development and neuropathic pain behavior, and show anxiolytic and antidepressant effect (6-10). A missense mutation in galanin peptide was identified as a cause of temporal lobe epilepsy (TLE) (11). Moreover, galanin expression is upregulated in the injured neurons, and galanin has been shown to play a role in neuroprotection and neuronal regeneration (12, 13). Accumulating evidence indicate that GALRs signaling is a key regulator of both sleep time and sleep/awake homeostasis in model organisms such as zebrafish and mouse (14, 15). Therefore, GALRs are potential therapeutic targets for the treatment of pain, epilepsy, depression, neuron injury and sleep disorders.

In addition to galanin, the endogenous galanin-like peptide (GALP) and spexin have been

shown to activate GALRs (16, 17). Although these peptides share high sequence similarity, they show distinct receptor binding preference. In contrast to galanin that interacts with all three receptor subtypes, spexin specifically activate GALR2 and GALR3. Homology modeling and site directed mutagenesis studies revealed the essential residues of galanin involved in the receptor binding and activation, and the potential galanin binding site of GALRs (18, 19). However, the molecular details of galanin binding and the peptide selectivity of GALRs remain poorly defined at the molecular level. Moreover, little is known about the molecular basis of G protein coupling specificity of GALRs. To gain insight into the molecular basis of ligand recognition and ligand selectivity of GALRs and extend our understanding of G protein selectivity, we sought to determine the cryoelectron microscopy (cryo-EM) structures of GALR1 and GALR2 in complex with G_o and G_q heterotrimer respectively.

79

80

Results

Structure determination

To obtain stable GPCR-G protein complexes, we used engineered thermostable mini-G proteins, which only contain the GTPase domain of $G\alpha$ but still bind to $G\beta\gamma$ heterodimer and recapitulate the pharmacological and structural changes in GPCRs induced by the full-length $G\alpha$ proteins (20). Moreover, the N-terminal residues of αN in mini-Gao and mini-Gaq were replaced by the equivalent residues of $G\alpha i$ to acquire the ability to bind the antibody fragment scFv16 that stabilizes the nucleotide-free GPCR-G protein complex (21). Furthermore, we introduced a linker that contains a 3C protease cleavage site, between the C-terminus of the receptor and the N-terminus of the mini- $G\alpha$ to create a GPCR-G fusion protein. The GALR1-mini-Gao or GALR2-mini-Gaq fusion protein was transiently expressed in Expi293 cell, and was assembled with purified $G\beta 1\gamma 2$ and scFv16 in the presence of galanin (**SI Appendix, Fig. S1**). The resulting GALR1-mini-Go complexes were co-eluted and mono-disperse with or without 3C protease treatment from the size exclusion chromatography, indicating that GALR1 forms a stable complex with mini-Go (**SI Appendix, Figs. S1A-1C**). The peak fraction corresponding to the complexes were concentrated and subjected to cryo-EM single particle analysis. 2D class average analysis showed that the GALR1-mini-G fusion protein complex gives more orientations than the GALR1-mini-G complex without a linker between the receptor and mini- $G\alpha$ (**SI Appendix, Figs. S1D and S1E**). Combination of the two datasets enables us to obtain a final cryo-EM map of the GALR1-mini-Go complex at a global nominal resolution of 3.3 Å (**SI Appendix, Fig. S2 and Table S1**). The structure of the galanin- and spexin-bound GALR2-minGq fusion complex was determined to a nominal resolution of 3.3 Å and 3.5 Å, respectively (**SI Appendix, Fig. S3 and Table S1**). The high quality EM map allowed us to unambiguously assign side chains of the galanin peptide 1-17 and the most amino acids of the receptors except the extreme terminal residues and some intracellular loops because of their high flexibility (**Fig. 1 B and C**). The overall structure of the GALR1-Go complex resembles that of the GALR2-Gq complex, with root-mean-square deviation values of 0.886 Å for the C α atoms of

the receptors and 0.604 Å for the Ca atoms of the G proteins.

Comparison of galanin binding pockets of GALR1 and GALR2

The existence of bulky aromatic amino acids and the high quality EM density map allowed us to unambiguously assign side chains of galanin (**Fig. 2A**). The N-terminal portion of galanin (residue 1-15) was well resolved due to its direct contact with the receptors, which is consistent with previous studies showing that the binding affinity of N-terminal region of galanin (1-16) for the receptors is comparable to the full-length galanin (22, 23). Moreover, the N-terminal region (1-16) but not the remaining part is highly conserved in GALP and spexin peptides (**SI Appendix, Fig. S4A**), both of which are able to activate the receptors, further supporting our structural observation. Galanin mainly forms an alpha helical structure when bound to the receptor as well as in solution itself (24, 25). It occupies at the extracellular vestibule of GALRs that is equivalent to the binding site of a positive allosteric agonist LY2119620 in M2R (26) (**SI Appendix, Fig. S4B**). It lays on top of the receptor, nearly parallel to the membrane plane and distant from the toggle switch W^{6.48}, the conformational change of which is essential for the receptor activation. By contrast, most neuropeptide agonists of class A GPCRs such as endothelin, orexin and opioid peptides binds nearly perpendicular to the membrane plane with one end buried in the helical cavity and the other end interacting with the extracellular loops, and these peptides penetrate in proximity to the toggle switch (27-29) (**SI Appendix, Fig. S4B**). Galanin contacts all seven TM helices as well as extracellular loops ECL2 and ECL3, burying a surface area of 866 Å², which accounts for the high affinity binding of galanin for GALRs in the sub-nanomolar range (30). GALR1 and GALR2 use overlapping but distinct set of residues to contact galanin, mostly via hydrophobic and hydrogen bond interactions (**Fig. 2**). The first N-terminal residue of G1 lies between TM1 and TM7, and is closer to TM1 of GALR1 than that of GALR2 (**Fig. 2F**), which may explain that removal of G1 in galanin reduced its binding affinity for GALR1 but not GALR2 (24). W2 is sandwiched between L277^{ECL3} and F282^{7.32} of GALR1 and makes additional hydrogen bond with S281^{7.31} (**Fig. 2C**). Therefore, mutation of W2 in

galanin or F282^{7,32} results in significant loss of binding for the receptors (18, 31). F282^{7,32}A mutation in GALR1 almost abolished galanin potency (**Fig. 2G**), and F271^{7,32}A mutation in GALR2 reduced galanin potency by nearly 100-fold (**Fig. 2H**). A7E mutation that was identified as a cause of TLE disease likely causes a clash with nearby hydrophobic residues, accounting for reduced binding affinity for GALRs (11) (**Fig. 2C**). Y9 penetrates into the receptor core, about 10 Å above the toggle switch (*SI Appendix, Fig. S4B*), and is hydrogen-bonded by Q92^{2,61} in GALR1 or Q82^{2,61} in GALR2 (**Figs. 2 C and E**). Mutation of Q92^{2,61} or Q82^{2,61} to alanine reduced galanin potency by almost 100-fold (**Figs. 2 G and H**). Our structural observation is also consistent with previous studies showing that W2 and Y9 are vital for galanin binding to the receptors (31). However, because of distinct residues of ECL2 and ECL3 involved in binding galanin, the conformations of these regions are different between GALR1 and GALR2 (**Fig. 2F**). For instance, V274 in the ECL3 of GALR1 engages hydrophobic interaction with L11 (**Fig. 2C**), while the equivalent residue in GALR2, Q263 rotates away from galanin due to its longer side chain and hydrophilic nature, resulting in the conformational change of ECL3 (**Fig. 2F**). As a result, V274G mutation reduced agonist potency by about 370-fold, while Q263A mutation showed little effect (**Figs. 2 G and H**). ECL2 forms an antiparallel β-sheet, which is a characteristic of peptide receptors. It covers galanin as a lid-like structure and forms extensive hydrophobic interactions with L4, P13 and V16. The residues in ECL2 of GALR1 involved in binding galanin have bulkier aromatic side chains than that in GALR2 (**Fig. 2F**). Mutations of the equivalent residues W188^{ECL2} and H176^{ECL2} in GALR1 and GALR2 respectively had distinct effect on galanin potency (**Fig. 2 G and H**), suggesting that ECL2 in GALR1 and GALR2 differently contribute to galanin binding. An endogenous peptide spexin has A7M and G8L mutations in galanin and specifically activates GALR2 and GALR3 (*SI Appendix, Fig. S4A*). Structure of the spexin-bound GALR2-Gq complex reveals that L8 in spexin likely clashes with the bulkier residue W188^{ECL2} in GALR1 (*SI Appendix, Fig. S4C*), accounting for the specific binding of spexin for GALR2 and GALR3 (16). Owing to these conformational differences, R184^{5,35} in GALR2 but not K197^{5,35} in GALR1 makes hydrogen bonds with the backbone of galanin (**Fig. 2F**).

As a result, R184^{5.35} A mutation reduces galanin potency by about 10-fold, while K197^{5.35} A mutation shows little effect (**Figs. 2 G and H**). Taken together, these results suggest that mechanisms of galanin recognition by GALR1 and GALR2 are not identical, which allows the development of selective ligands targeting a specific subtype.

Activation mechanisms of GALR1 and GALR2

The inactive structures of GALR1 and GALR2 predicted by Alphafold may represent ligand-free structure, in which TM helices loosely pack against each other to allow the access of galanin (32) (**Fig. 3A**). Upon galanin binding, the orthosteric site undergoes significant conformational change, as indicated by the inward displacement of the extracellular portions of TM2 and TM6 and the outward shift of the TM1 and TM7 (**Figs. 3 A and B**). These conformation changes account for the outward motion of TM6 and inward motion of TM7 in the intracellular side (**Fig. 3C**). The conformational changes of the toggle switch W^{6.48} and P^{5.50}I/V^{3.40}F^{6.44} motif are common features of the class A GPCR activation. In contrast to the most class A GPCRs, where the orthosteric sites are in close proximity to the toggle switch, galanin binding site is distant from it. Hydrophobic interactions between F275 in ECL3 of GALR1 and, L10 and L11 in galanin result in the downward shift of F275, which propagates to the downward movement of W260^{6.48} via I266^{6.54} and H263^{6.51} (**Fig. 3B**). The downward shift of W^{6.48} is associated with the conformational change of the P^{5.50}I/V^{3.40}F^{6.44} motif, which allosterically disrupts the conserved ionic lock between R133^{3.50} and D132^{3.49}, leading to the outward displacement of TM6 (**Fig. 3C**). Inward displacement of TM7 in the intracellular side is observed in the active state of many other class A GPCRs, as indicated by the conformational change of NPXXY motif, in which Y303^{7.53} forms a water-mediated hydrogen bond with Y220^{5.58}. The inward displacement of TM7 is coupled by the outward shift of R285^{7.35} that arises from its interaction with galanin. Although the key residues involved in receptor activation are conserved between GALR1 and GALR2, their conformations vary significantly (**Fig. 3D**). This is because the hydrophobic interaction between V274 in ECL3 and L11 in galanin exists in GALR1, while this interaction is absent in

GALR2 due to the substitution of V274 in Q263, which leads to the upward shift of F264^{ECL3} as well as the toggle switch W^{6.48} and PIF motif in GALR2, compared to the equivalent residues in GALR1 (**Fig. 3D**). To further support the important role of F275^{ECL3}/F264^{ECL3} in GALR receptors activation, mutation of F275 in GALR1 or F264 in GALR2 reduced galanin potency by almost 100-fold (**Figs. 2 G and H**). The conformation differences of residues involved in receptor activation contribute to the structural variation in the cytoplasmic pocket of GALR1 and GALR2 and may play a role in G protein selectivity.

Zn²⁺ is a negative allosteric modulator (NAM) of GALR1

Previous studies have reported that Zn²⁺ can inhibit galanin binding to the receptors (18). To further investigate the functional role of zinc ion in galanin receptors signaling, we used the NanoBiT complementation-based assay to assess the effect of Zn²⁺ on activation of receptors by galanin in living cells. Mini-G proteins were used in the NanoBiT assay throughout this study, since they preserve appropriate coupling specificity, and can be recruited to the active GPCRs without further dissociation, which increases the signal-to-noise ratio in this assay. As expected, Zn²⁺ diminished the effect of 1 μ M galanin on GALR1 activation in a concentration dependent manner with an IC₅₀ value of 47.2 μ M (**Fig. 4A**). The diminished effect of Zn²⁺ is saturable, or has a “ceiling” level. In contrast, the diminished effect was observed in GALR2 when the concentration of Zn²⁺ reached millimolar range that is above the physiological concentration, indicating that Zn²⁺ had little effect on galanin-induced GALR2 activation. Our structures show that galanin receptors are enriched with histidine residues that may coordinate Zn²⁺ underneath the orthosteric binding pocket (**Fig. 4B**). Comparison of primary sequences of GALR1 and GALR2 from different species revealed that H267^{6.55} but not nearby histidine residues in GALR1 is mutated to Isoleucine in GALR2 (**SI Appendix, Fig. S5A**). As expected, the zinc effect was significantly abrogated in GALR1 when H267^{6.55} was mutated, while H112^{3.29}A, H263^{6.51}F or H264^{6.52}F mutation had little influence (**Fig. 4A and SI Appendix, Fig. S5B**). All these mutants of GALR1 can be activated by galanin, although the potency and efficacy of galanin for these mutants vary (**SI**

Appendix, Fig. S5C). We further tested the effect of Zn^{2+} on the concentration response curve of galanin. Zn^{2+} produced the concentration-dependent and saturable rightward shifts in the potency of galanin, and decreased the galanin maximum response as well (**Fig. 4C**). By contrast, Zn^{2+} had little effect on the galanin concentration-response curve of H267A mutant of GALR1 (**Fig. 4D**). H267 is located in the TM6 right below the galanin binding site. The extracellular part of TM6 near H267 moves inwards upon galanin binding, which leads to the receptor activation. As a result, when coordinated by H267 and other nearby residues, Zn^{2+} likely rigidifies the extracellular part of TM6 and restricts its conformational change, attenuating galanin-induced receptor activation. The exact coordination pattern of Zn^{2+} awaits further investigation. Nevertheless, these results indicate that Zn^{2+} is a NAM of GALR1.

Structural determinants of Gi/o and Gq/11 selectivity

A notable difference between structures of the GALR1-Go complex and the GALR2-Gq complex is the relative orientation of Go and Gq to the receptors (**Figs. 5 A and B**). When aligning the receptors, the $\alpha 5$ of Gao is rotated around the “wavy hook” of $\alpha 5$ by about 14° toward TM5, compared with G α q. This orientation difference was also observed in the structures of M1 and M2 muscarinic receptors (M1R and M2R) bound to G11 and Go respectively (33). In addition, because of the different interaction interface of the receptor and G protein, the flexibility of intracellular loops (ICL) in GALR1 and GALR2 differs (**Figs. 5A**). For instance, ICL1 is ordered in GALR2 owing to the hydrogen bond interaction between D312 in G β and the main chain carbonyl group of G53 in ICL1, whereas it is flexible in GALR1 due to the absence of this interaction (**SI Appendix, Fig. S6A**). The ICL2 of most Gi/o-coupled receptors forms an alpha helical structure, where hydrophobic residues at position 34.51 engage weak hydrophobic interactions with the hydrophobic pocket of Gai/o formed by V34 from the α N- β 1 loop, L195 from the β 2- β 3 loop and I343 and F336 from α 5 (**SI Appendix, Fig. S6B**). However, the ICL2 of GALR1 is disordered because of the substitution of the hydrophobic residue L1313^{4.51} in arginine and the absence of hydrophobic

interaction between the ICL2 of GALR1 and Gi (**Figs. 5 C and G**). By contrast, L131^{34.51} in the ICL2 of GALR2 is buried deep in the hydrophobic pocket of Gαq formed by L34 from the αN-β1 loop, V79 from the β2-β3 loop, and F228, K232 and I235 from α5, and engages strong hydrophobic interactions. In addition, P130^{34.50} at the junction of ICL1 and TM3 is stabilized through hydrophobic interactions with I235 and K232 in α5 of Gαq (**Fig. 5D**). Substitution of L131^{34.51} or P130^{34.50} in GALR2 in the equivalent residues in GALR1 impaired the ability of GALR2 to couple Gq (**SI Appendix, Fig. S7A**), accounting for the inability of GALR1 to couple Gq. However, substitutions of S140^{34.50}, R141^{34.51} and S149^{4.49} in ICL2 of GALR1 with the equivalent residues in GALR2 have little effect on coupling efficiency between GALR1 and Gαi (**SI Appendix, Fig. S7B**), suggesting that ICL2 in GALR1 is not involved in Gi coupling. Remarkably, GALR1 acquires the ability to couple Gq, as indicated by the NanoBiT assay as well as the IP1 assay (**Fig. 5H and SI Appendix, Fig. S7C**), when residues in ICL2 of GALR1 are substituted with that of GALR2, further supporting the important role of ICL2 in Gq coupling.

To understand the structural mechanism of the inability of GALR2 to couple Gi, we compared the interaction details between the GALR1-Gi and the GALR2-Gq complexes and mainly focused on residues of GALR1 involved in Go coupling that are not conserved in GALR2. R133^{3.50}, I137^{3.54}, L224^{5.62}, L227^{5.65}, L231^{5.69}, and T245^{6.33} in TM3, TM5 and TM6 of GALR1 engage extensive hydrophobic interactions with the extreme C-terminal part of α5 in Go (**Fig. 5E**). Most of these residues are conserved in GALR2 (**Fig. 5F**). Mutations of conserved residues in GALR1 and GALR2 impair the recruitment of Gi and Gq, respectively (**SI Appendix, Figs. S7D-7G**). Nevertheless, a notable difference between GALR1 and GALR2 are in ICL3. S238^{ICL3} and S235^{ICL3} in ICL3 of GALR1 make hydrogen bonds with D341 in Gαo, and K237^{ICL3} engages electrostatic interactions with residues in the GTPase domain of Gαo. All of three residues are mutated in GALR2, leading to loss of these interactions (**Fig. 5 E and F**). Mutations of S235^{ICL3} and K237^{ICL3} in GALR1 have modest effect on galanin potency, but dramatically reduced the maximum responses (**SI Appendix,**

Figs. S7D and S7E). Remarkably, GALR2 acquired the ability to bind Go, when ICL3 (214-225) of GALR2 including the three residues were replaced by the equivalent residues in GALR1 (**Fig. 5A**). Taken together, our data suggest that ICL2 in GALR2 and ICL3 in GALR1 are critical for determining the Gq and Go selectivity, respectively.

Structural determinants of Gs and Gq selectivity

Although it has been shown that interactions between the hydrophobic residue at position 34.51 of ICL2 and the hydrophobic pocket of $G\alpha$ are essential for the efficient coupling of Gq and Gs (34), it remains unclear how Gq and Gs are selectively recognized. Comparison of structures of D1 dopamine receptor (D1R)-Gs and GALR2-Gq revealed key structural elements in the receptors that determine Gq and Gs selectivity. In the GALR2-Gq complex, the conformation of ICL2 is stabilized by salt bridge interactions between $R^{34.57}$ ($M^{34.57}$ in D1R) and $D^{3.49}$ of the DRY motif as well as a hydrogen bond between $R^{34.57}$ and $Y(-4)$ in $G\alpha_q$, while ICL2 of D1R is stabilized by a hydrogen bond between $Y^{34.53}$ ($S^{34.53}$ in GALR2) and $D^{3.49}$, and a potential water-mediated hydrogen bond between $Y^{3.49}$ and $Y(-4)$ in $G\alpha_s$ (**Fig. 6A**). Notably, $Y^{34.53}M/V^{34.57}$ are prevalent in Gs-coupled receptors, while $R^{34.57}$ is enriched in Gq-coupled receptors (**Fig. 6D**). Mutations of YM in D1R and RS in GALR2 significantly reduced the potency of dopamine and galanin, respectively (**Figs. 6 E and F**). Moreover, $N(-3)$ (-1 indicates the last residue of $G\alpha$) in $G\alpha_q$ is inserted into a hydrophobic pocket formed by $N^{2.40}$, $F^{8.50}$ and other nearby residues, whereas $E(-3)$ flips outside this pocket, probably due to its longer side chain and negative-charge nature. Remarkably, when $E(-3)$ in $G\alpha_s$ but not the nearby residues $L(-1)$ and $Q(-5)$ was substituted with the equivalent residues in $G\alpha_q$, the coupling efficiency between GALR2 and Gs was significantly increased (**Fig. 6G**). These different interaction modes of GALR2-Gq and D1R-Gs account for the movement of $\alpha 5$ in Gs toward TM6 and the outward movement of TM6 in D1R, compared to that in GALR2 (**Fig. 6B**), explaining that most Gs-coupled receptors display a larger TM6 movement in the active state than Gq-coupled receptors. As a result of these conformational changes, Gs is closer to TM5 than Gq, highlighting the important role of TM5 in determining Gs selectivity. Indeed, TM5 in most Gs-coupled receptors have a C-terminal helical

extension, and previous studies have shown that the $AV^{5.65}Q^{5.68}\Phi^{5.69}$ (Φ represents hydrophobic residues) motif in TM5 is prevalent in receptors that exclusively couple to Gs and is critical for Gs coupling in D1R (35) (Companion paper). Residues at position 5.65 in Gs-coupled receptors prefer hydrophobic residues with small side chains such as alanine and valine because of its close distance from the hydrophobic pocket formed by L(-1), L(-2), and L(-7) in G α s (**Fig. 6C**). Mutation of A^{5.65} in leucine would cause a clash with this pocket, and impaired the Gs coupling (Companion paper). In contrast, leucine is dominant at position 5.65 in Gq-coupled receptors, due to its long distance from the hydrophobic pocket formed by V(-1), L(-2), and L(-7) in G α q (**Fig. 6C**). L^{5.65}A mutation in GALR2 weakens the interaction with this hydrophobic pocket and thus significantly decreased galanin potency (**Fig. 6F**). However, it is noteworthy that Gs- and Gq-coupled receptors show sequence preference at some positions of ICL2 and TM5, but also accommodate various residues at these positions (**Fig. 6D**), partly because of diverse receptor-G protein interfaces and promiscuous coupling of some GPCRs.

Discussion

Here, we report cryo-EM structures of the GALR1-Go and GALR2-Gq complex using the GPCR-G protein fusion strategy. The structures revealed distinct mechanisms of galanin recognition and receptor activation for GALR1 and GALR2, which contribute to structural variation in the cytoplasmic pocket of the receptors and may play an important role in determining the G protein selectivity. Moreover, we showed that Zn²⁺ is a negative allosteric modulator of GALR1 but not GALR2.

Zn²⁺, known as a neuromodulator, is widely distributed in the central nervous system (CNS), particularly enriched in the synaptic vesicles of glutamatergic neurons (36, 37). It is released to the synaptic cleft upon membrane depolarization, and modulates functions of ion channels and receptors on the pre- or post-synaptic membrane. It has been shown that zinc inhibits ionotropic glutamate AMPA and NMDAR receptors, fine-tuning synaptic transmission in the brain (38, 39). GALR1 is expressed on both glutamatergic and GABAergic

postsynaptic neurons. The spatial colocalization of zinc and GALR1 makes it possible for zinc to regulate the function of GALR1. Moreover, the IC₅₀ of zinc on GALR1 activation is 47.2 μ M, which is in the range of the physiological concentration of zinc (10 nM to 100 μ M) (40). Previous studies have also shown that zinc regulates endogenous ligand binding at several GPCRs including β 2 adrenergic receptors (β 2AR) (41), melanocortin receptors (42) and platelet-activating factor receptor (43). In this study, we showed that zinc attenuated GALR1 activation by galanin possibly through restricting the conformational change of TM6 that leads to receptor activation. Further studies are required to address whether zinc modulates a large number of GPCRs in the CNS and fine-tunes GPCR signaling, as does sodium (44).

Combining published structures of the GPCR-Gi complexes, we can roughly divide the class A Gi-coupled receptors into three classes based on the interaction features between ICL2 and G proteins: (i) receptors that exclusively couple to Gi and have a charge residue at the position 34.51 of ICL2 such as GALR1 and sphingosine-1-phosphate receptors (S1PR) (45); (ii) receptors that exclusively couple to Gi and have a hydrophobic residue at the position 34.51 such as D3 dopamine receptor, M2 muscarinic receptor (M2R) and μ opioid receptor (28, 33, 46); (iii) receptors that promiscuously couple to Gi and have a large hydrophobic residue at position 34.51 such as the neurotensin receptor 1 (NTSR1), β 2AR, and the cholecystokinin A receptor (47, 48). In the first class, when bound to the receptor, ICL2 is disordered, or forms a random coil structure. Since there is no hydrophobic interaction between 34.51 and G α i, ICL2 in receptors of this class plays a distinct role in determining Gi coupling efficiency (45) (**SI Appendix, Fig. S8A**). In GALR1, ICL2 is not involved in Gi coupling, whereas in S1PR, ICL2 is involved in hydrophilic interactions with Gi, and is important for Gi coupling; In the second class, ICL2 forms an alpha helical structure, and the residue 34.51 of ICL2 is located outside and distant from the hydrophobic pocket of G α i formed by residues from the α N- β 1 loop, the β 2- β 3 loop and α 5, and engages weak hydrophobic interactions (**SI Appendix, Figs. S8B and S8C**). Mutation of this residue had

little effect on the Gi coupling or GDP release from Gi/o (49, 50); In the third class, similar to Gs- and Gq-coupled receptors, the residue 34.51 is located close to the middle of the hydrophobic pocket of Gai and engages strong hydrophobic interaction (**SI Appendix, Figs. S8E and S8F**). In addition, some receptors in this class such as NTSR1 have the other conformation, where the residue 34.51 is located outside the hydrophobic pocket (**SI Appendix, Fig. S8D**). Previous studies have shown that F^{34.51}A mutant of β 2AR failed to activate Gi (50), suggesting that the hydrophobic interaction between ICL2 and Gi is very important for Gi coupling in the third class. Owing to the absence of or weak interaction between Gai and ICL2 in receptors that exclusively couple Gi, the cytoplasmic end of TM5 and TM6, and ICL3 have strong interactions with Gai, and are critical for determining Gi selectivity (**SI Appendix, Fig. S8**).

It has been recognized that the distal part of α 5 in G α plays a key role in determining G protein selectivity (51-53). We further identified a residue pair N/E(-3) in α 5 of Gq/Gs that contributes to structural differences and selective interactions between the D1R-Gs and GALR2-Gq. Substitution of this residue in Gs can promote coupling of GALR2 to noncognate Gs. Moreover, we revealed several signature residues in ICL2 and TM5 that dominate in Gs- and Gq-coupled receptors. Thus, our results provide novel insights into the molecular mechanisms of G protein selectivity by class A GPCRs.

Figure legends:

Fig. 1. Overall structures of galanin-bound GALR1-miniGo-scFv16 and GALR2-miniGq-scFv16 complexes.

(A) Schematic representation of GALR receptors signaling. GALR1 and GALR3 primarily couple to Gi/o, while GALR2 mainly signals through Gq.

(B) Cryo-EM structures of GALR1-miniGo-scFv16.

(C) Cryo-EM structures of GALR2-miniGq-scFv16.

Fig. 2. Mechanisms of galanin recognition by GALR1 and GALR2.

(A) EM density map for galanin from the structure of the GALR2-Gq complex.

(B) Electrostatic potential surface of GALR1 and ribbon representation of galanin (magenta) viewed from the extracellular side. Colors from red to blue represent negatively to positively charged regions.

(C) Detailed interaction between GALR1 and galanin.

(D) Electrostatic potential of the GALR2-galanin interface is distinct from that of the GALR1-galanin interface.

(E) Detailed interaction between GALR2 and galanin.

(F) Structural superposition of the GALR1-galanin and the GALR2-galanin complex. The

403 equivalent residues in GALR1 and GALR2 that play distinct roles in galanin binding are
404 shown. Arrows indicate the conformational changes.
405 (G) and (H) The effects of mutations in GALR1 and GALR2 on galanin potency as measured
406 by the cAMP inhibition assay and the IP1 accumulation assay respectively. The equivalent
407 residues that play distinct roles in GALR signaling are colored red. Data represent mean $\pm$
408 SEM of triplicate measurements in three independent experiments. Significance was
409 analyzed using one-way ANOVA, ****P<0.0001, ***P<0.001, **P<0.01.

410

411

412 **Fig. 3. Mechanisms of GALR1 and GALR2 activation.**

413 (A) Structural overlay of GALR1 in the active and inactive state (predicted by AlphaFold).

414 (B) Conformational changes of the P^{5.50}I^{3.40}F^{6.44} motif upon receptor activation.

415 (C) Conformational changes of residues in the cytoplasmic pocket including the
416 D^{3.49}R^{3.50}Y^{3.51} and the NPY motif upon receptor activation.

417 (D) Structural overlay of the active state of GALR1 and GALR2.

Fig. 4. Zinc is a NAM of GALR1.

(A) The effect of increasing concentration of zinc on receptor activation induced by 1 μM galanin, as evaluated by the NanoBiT assay, where the small fragment, and the large fragment are fused to the C-terminus of GALR1 and the N-terminus of mini-Go, respectively. The luminescence signals are normalized as percentages of the initial response of GALR1 to galanin without zinc treatment.

(B) Histidine residues are enriched underneath the galanin binding pocket of GALR1.

(C) and (D) The actions of increasing concentration of zinc on the galanin dose-response curve of WT (C) and H267A mutant (D) of GALR1 measured by the NanoBiT assay. The luminescence signals are normalized to the vehicle treatment as fold change.

Fig. 5. Mechanisms of Go and Gq selectivity in the GALR receptor family.

(A) and (B) Structural superposition of GALR1-Gao and GALR2-Gaq in two opposite views. Receptors are aligned.

(C) Interaction details between ICL2 of GALR1 and Gao.

(D) Interaction details between ICL2 of GALR2 and Gaq.

(E) Interaction details between TM5 and TM6 of GALR1 and Gao.

(F) Interaction details between TM5 and TM6 of GALR2 and Gaq.

(G) Sequence alignment of ICL2 from GALR receptor family.

(H) The IP1 accumulation assay evaluating the effects of ICL2 substitutions in GALR1 on GALR1-Gq coupling. All mutants are expressed at similar levels as WT.

(I) Substitution of ICL3 in GALR2 with that in GALR1 increases coupling efficiency of GALR2 and Go.

- 454 (F) Mutations of S^{34.53}R^{34.57} in the ICL2 of GALR2 almost abolish GALR2 and Gq coupling.
455 (G) The effects of mutations of the “wavy hook” in Gs on coupling efficiency of GALR2-Gαs,
456 as evaluated by the NanoBiT assay.

Materials and method

Cloning

The human GALR1 and GALR2 were cloned into pcDNA3.1(+) vector(Thermo Fisher Scientific) with an N-terminal hemagglutinin (HA) signal sequence and a FLAG epitope tag (DYKDDDDK). An engineered mini-Gαo was fused to the C-terminus of GALR1 (1-349) with three copies of 3C protease sites between them. GALR2 (1-314) was expressed as a fusion protein including two repeats of 3C protease site and a mini-Gαq sequence in the C-terminus of GALR2. ScFv16 was cloned into the pFastBac vector (Invitrogen) with an N-terminal GP64 signal sequence and a C-terminal 3C protease site, followed by an octa-histidine tag. His6-tagged Gβ1 and Gγ2 (C68S mutation) were cloned in the pFastBac Dual vector for insect cell expression.

Protein expression and purification

The plasmid expressing GALR1-mini-Gαo or GALR1-mini-Gαq was transiently expressed into Expi293F cells (Thermo Fisher Scientific) using polyethyleneimine (PEI, Polysciences). Cells were lysed in the lysis buffer (20 mM HEPES, pH 7.4) supplemented with protease inhibitor cocktail (Roche) using a glass dounce grinder and centrifuged at 1,000 x g for 3 minutes to remove the nucleus. The membrane fraction was pelleted by centrifugation at 65,000x g, at 4 °C for 1 hour, and homogenized in the solubilization buffer containing 20 mM HEPES pH 7.4, 150 mM NaCl, 1% Lauryl Maltose Neopentyl Glycol (LMNG), 0.2% Cholesteryl Hemisuccinate (CHS) and 60 nM galanin peptide (1-30). After centrifugation to remove debris, the supernatant containing solubilized GALRs-mini-Gα was supplemented with 2 mM CaCl₂ and loaded onto the M1 anti-FLAG antibody resin. The resin was washed with wash buffer containing 20 mM HEPES, pH 7.4, 300 mM KCl, 0.01% LMNG, 0.002% CHS, 2 mM CaCl₂, 10 mM MgCl₂, 2 mM ATP, 6 nM galanin and eluted with elution buffer containing 20 mM HEPES, pH 7.4, 150 mM NaCl, 0.01% LMNG, 0.002% CHS, 10 mM EDTA, 0.5 mg/ml 1x FLAG peptide and 60 nM galanin. Gβ₁γ₂ (C68S) and scFv16 were expressed and purified as previously described (21, 54).

Complex assembly

Purified GALR1-miniGo or GALR2-miniGq proteins, G β ₁ γ ₂ (C68S) and scFv16 were mixed with a molar ratio of 1:1.5:2 in 500 μ l of the equilibration buffer (20 mM HEPES, pH 7.4, 150 mM NaCl, 0.01% LMNG, 0.002% CHS, 60 nM galanin, 0.5 μ M TCEP) supplemented with 1 μ l PNGaseF and 0.5 μ l apyrase. The mixture was incubated on ice for 1 hour and further purified on a Superose 6 Increase 10/300 column pre-equilibrated with the equilibration buffer. The peak fractions containing the complex were supplemented with 60 μ M galanin and concentrated to about 6 mg/ml. For assembly of the GALR1-miniGao/G β ₁ γ ₂ (C68S) complex with 3C protease site cleaved, similar procedures were performed as above, except that 3C protease was added before purification on a Superose 6 Increase 10/300 column. For assembly of the spexin-bound GALR2 complexes, same procedures were performed, except that galanin was replaced by spexin during the purification process.

Cryo-EM sample preparation and data collection

300 mesh holey carbon grids (Quantifoil Au R1.2/1.3) were glow-charged, loaded into a Vitrobot MarkIV instrument chamber (Thermo Fisher Scientific) maintained at 8 °C and 100% humidity. 3.0 μ l of GALRs complex samples was applied onto the grid, blotted for 3.0-4.0 s with a blotting force of 4 before plunge freezing in liquid ethane. Cryo-EM movies were collected on a Titan Krios microscope equipped with a BioQuantum GIF/K3 direct electron detector (Gatan) under accelerating voltage of 300 kV at a nominal magnification of 64,000 x. Each movie stack was collected as 32 frames, with total dose of 50 e-/Å² for 2.56 s.

Cryo-EM Data processing

All movie stacks were collected and processed with MotionCor2 for motion correction (55), with 2x binned to a pixel size of 1.087 Å. Contrast Transfer Function (CTF) estimation was performed using patch-based CTF estimation in cryoSPARC_v3 (56). All processed images were then subjected to particle picking using Blob picker in cryoSPARC, followed by particle extraction. For the galanin-bound GALR1-mini-Go complex with 3C protease sites cleaved,

particles from 1,401 micrographs (Dataset A) were subjected to two rounds of 2D classification, generating 248,352 good particles. *Ab initio* reconstruction and non-uniform refinement were performed to get a reference map for GALR1. For the GALR1-mini-Go fusion protein complex, 886 micrographs (Dataset B) were collected, followed by particle picking using Blob picker and particle extraction. Particles from the two datasets were combined and subjected to two rounds of 2D classification, yielding 2,882,487 good particles. These particles were subjected to global 3D classification in RELION3.1 (57), followed by another round of 3D classification focused on the receptor. 426,045 particles from the best class were run through non-uniform refinement in cryoSPARC, resulting in a final 3.3 Å map.

For the galanin-bound GALR2-mini-Gq fusion protein complex, 1,337 micrographs were collected, and processing procedures were performed as above. In brief, two rounds of 2D classification using auto-picked particles resulted in 1,325,739 good particles, which were subjected to two rounds of 3D classification in RELION3.1 using the GALR1-Go complex map as initial model. 578,453 particles from three classes with clear secondary structure features were selected, and subjected to non-uniform refinement in cryoSPARC, resulting in a final 3.29 Å map. All 3D maps were post-processed with DeepEMhancer (58).

For the spexin-bound GALR2-mini-Gq complex, 1,139 movies were collected and processed as above. 1,015,461 good particles were selected from two rounds of 2D classification, and were subjected to heterogeneous refinement and non-uniform refinement in cryoSPARC. The final map is about 3.5 Å.

Model building

Homology models for GALR1 and GALR2 were generated using the structure of μ opioid receptor (PDB: 4DKL) in the SWISS-MODEL server. The homology model of GALR1 and the structure of mini-Gao/G β γ /scFv16 (PDB: 7D77) were fitted into the EM map in Chimera (59). The structure of mini-G α q/G β γ /scFv16 was extracted from the published structure

(PDB: 6WHA), and docked into the EM map together with the homology model of GALR2. All the models were manually built in COOT (60) and, are subjected to *real_space_refinement* in *Phenix* (61) using the reference structure and secondary structure restraints. The statistics for structure refinement are summarized in Table S1.

cAMP inhibition assay

Chinese hamster ovary (CHO) cells were seeded into six-well plates and cultured overnight until cell confluence reaches ~80%. Plasmids expressing GALR1 or mutants were transfected together with the GloSensor biosensor plasmid following a Lipofectamine cell transfection procedure (Invitrogen). Transfected cells were cultured for 1 day and re-seeded into 96-well plates by 3×10^4 cells per well. After 8 hours post seeding, the medium was exchanged to CO₂-independent medium (Gibco) supplemented with 500 µg/ml of D-luciferin. Cells were stimulated with various concentration gradients of galanin for 5 minutes, and then treated with 1 µM forskolin. The bioluminescence signal was constantly measured for 10 minutes, and the peak signal was acquired for the inhibitory dose curve fitting and IC₅₀ determination using GraphPad Prism 8 software. Significance analysis was performed using one-way analysis of variance method (one-way ANOVA in Prism 8).

IP1 accumulation assay

Gαq-mediated IP1 accumulation was measured using the IP-ONE Gq HTRF Kit from Cisbio. HEK-293T cells were seeded into 6-well plates, and 2 µg of GALR2 or mutant plasmids were transfected using PEI. After 2 days post-transfection, cells were suspended, washed one time with DPBS (Gibco), resuspended into HBSS buffer (Beyotime) and seeded into 384-well plates (Greiner) with ~7000 cells per well. Transfected cells were stimulated with various concentration gradients of galanin for 1 hour and subjected to IP1 accumulation detection following the assay protocol. Inhibitory dose curve was plotted and IC₅₀ was determined using GraphPad Prism 8 (dose-response-inhibitory, three parameters). Significance was analyzed using One-way ANOVA.

569

570 **NanoBiT assay**

571 To monitor the interaction between G proteins and GALR1 or GALR2 upon galanin
572 stimulation, a NanoLuc-based enzyme complementation system called NanoBiT assay (62)
573 was used (Promega). The C-terminus of GALR1 or GALR2 was fused with the small
574 fragment (smBit), and the large fragment (LgBit) element was fused to the N terminus of
575 mini-Gα proteins. HEK-293T cells were seeded into 6-well plates and transfected with 1 μg
576 of GPCR-smBit and 1 μg of LgBit-miniGα. After 2 days post transfection, cells were
577 suspended, washed with DPBS for one time and resuspended into the assay buffer
578 containing HBSS supplemented with 0.01% BSA (SIGMA), 10 mM HEPES (Beyotime) and
579 10 μM coelenterazine-h (YEASEN). The culture was equilibrated at room temperature (RT)
580 for 2 hours and subjected to stimulation with various concentration gradients of galanin and
581 instant bioluminescence measurement. The bioluminescence signal was acquired in the
582 time point when the signal went into the stationary phase, and the normalized signal (fold
583 change) was fitted to a three-parameter sigmoidal concentration-response curve in Prism 8
584 software.

585

586 **Zn²⁺ inhibition assay**

587 As zinc produced high background signal in the IP1 accumulation assay and cAMP inhibition
588 assay, the NanoBiT assay was used to measure the effect of Zn²⁺ effect on GALRs signaling.
589 The same constructs used in the NanoBiT assay were adopted. After 2 days
590 post-transfection, cells were resuspended and washed twice with the assay buffer (20 mM
591 HEPES, pH 7.3 and 150 mM NaCl), and resuspended into the assay buffer supplemented
592 with 10 μM coelenterazine-h and seeded into 96-well plates. After 30 minutes of incubation
593 at RT, cells were stimulated with various concentration gradients of galanin premixed with a
594 fixed concentration of ZnCl₂, or 1 μM of galanin pre-mixed with titrated concentration of
595 ZnCl₂. The bioluminescence signals in the stationary phase were acquired, and were
596 analyzed using three-parameter dose-response-stimulatory or dose-response-inhibitory

fitting methods in Prism 8 software.

Acknowledgements

We thank Dr. Xiangyu Liu at Tsinghua University for providing the plasmid expressing Gβ1γ2. We thank staff at Shuimu BioSciences for their help with cryo-EM data collection. All EM images were collected at Shuimu BioSciences. This work was supported by Chinese Ministry of Science and Technology, Beijing Municipal Science & Technology Commission (Z201100005320012) and Tsinghua University.

Author contributions

W.J. purified the protein complex, collected cryo-EM data, performed cryo-EM data processing and model building, performed cellular assay with input from S.Z. S.Z. and W.J. wrote the manuscript.

Competing interests

The authors declare no competing interests.

Data availability

The atomic structures have been deposited at the Protein Data Bank (PDB) under the accession codes XXX. The EM maps have been deposited at the Electron Microscopy Data Bank (EMDB) under the accession numbers XXX.

References

1. Tatemoto K, Rokaeus A, Jornvall H, McDonald TJ, & Mutt V (1983) Galanin - a novel biologically active peptide from porcine intestine. *FEBS Lett* 164(1):124-128.
2. Sipkova J, Kramarikova I, Hynie S, & Klenerova V (2017) The galanin and galanin receptor subtypes, its regulatory role in the biological and pathological functions. *Physiol Res* 66(5):729-740.
3. Lang R, *et al.* (2015) Physiology, signaling, and pharmacology of galanin peptides and receptors: three decades of emerging diversity. *Pharmacol Rev* 67(1):118-175.
4. Branchek TA, Smith KE, Gerald C, & Walker MW (2000) Galanin receptor subtypes. *Trends Pharmacol Sci* 21(3):109-117.
5. Wang S & Gustafson EL (1998) Galanin receptor subtypes. *Drug News Perspect* 11(8):458-468.
6. Lin EJ, *et al.* (2003) Recombinant AAV-mediated expression of galanin in rat hippocampus suppresses seizure development. *Eur J Neurosci* 18(7):2087-2092.
7. Haberman RP, Samulski RJ, & McCown TJ (2003) Attenuation of seizures and neuronal death by adeno-associated virus vector galanin expression and secretion. *Nat Med* 9(8):1076-1080.
8. Millon C, *et al.* (2019) Role of the galanin N-terminal fragment (1-15) in anhedonia: Involvement of the dopaminergic mesolimbic system. *J Psychopharmacol* 33(6):737-747.
9. Li SY, *et al.* (2017) Involvement of galanin and galanin receptor 1 in nociceptive

640 modulation in the central nucleus of amygdala in normal and neuropathic rats. *Sci*
641 *Rep* 7(1):15317.

642 10. Kokaia M, *et al.* (2001) Suppressed kindling epileptogenesis in mice with ectopic
643 overexpression of galanin. *Proc Natl Acad Sci U S A* 98(24):14006-14011.

644 11. Guipponi M, *et al.* (2015) Galanin pathogenic mutations in temporal lobe epilepsy.
645 *Hum Mol Genet* 24(11):3082-3091.

646 12. Xu XF, *et al.* (2016) Galanin and its receptor system promote the repair of injured
647 sciatic nerves in diabetic rats. *Neural Regen Res* 11(9):1517-1526.

648 13. Elliott-Hunt CR, Pope RJ, Vanderplank P, & Wynick D (2007) Activation of the galanin
649 receptor 2 (GalR2) protects the hippocampus from neuronal damage. *J Neurochem*
650 100(3):780-789.

651 14. Kroeger D, *et al.* (2018) Galanin neurons in the ventrolateral preoptic area promote
652 sleep and heat loss in mice. *Nat Commun* 9(1):4129.

653 15. Reichert S, Pavon Arocas O, & Rihel J (2019) The Neuropeptide Galanin Is Required
654 for Homeostatic Rebound Sleep following Increased Neuronal Activity. *Neuron*
655 104(2):370-384 e375.

656 16. Kim DK, *et al.* (2014) Coevolution of the spexin/galanin/kisspeptin family: Spexin
657 activates galanin receptor type II and III. *Endocrinology* 155(5):1864-1873.

658 17. Mills EG, Izzu-Engbeaya C, Abbara A, Comninou AN, & Dhillon WS (2021) Functions of
659 galanin, spexin and kisspeptin in metabolism, mood and behaviour. *Nat Rev*
660 *Endocrinol* 17(2):97-113.

- 661 18. Kask K, Berthold M, Kahl U, Nordvall G, & Bartfai T (1996) Delineation of the peptide
662 binding site of the human galanin receptor. *EMBO J* 15(2):236-244.
- 663 19. Church WB, Jones KA, Kuiper DA, Shine J, & Iismaa TP (2002) Molecular modelling
664 and site-directed mutagenesis of human GALR1 galanin receptor defines
665 determinants of receptor subtype specificity. *Protein Eng* 15(4):313-323.
- 666 20. Nehme R, *et al.* (2017) Mini-G proteins: Novel tools for studying GPCRs in their active
667 conformation. *PLoS One* 12(4):e0175642.
- 668 21. Maeda S, *et al.* (2018) Development of an antibody fragment that stabilizes
669 GPCR/G-protein complexes. *Nat Commun* 9(1):3712.
- 670 22. Smith KE, *et al.* (1997) Expression cloning of a rat hypothalamic galanin receptor
671 coupled to phosphoinositide turnover. *J Biol Chem* 272(39):24612-24616.
- 672 23. Fathi Z, *et al.* (1997) Cloning, pharmacological characterization and distribution of a
673 novel galanin receptor. *Brain Res Mol Brain Res* 51(1-2):49-59.
- 674 24. Carpenter KA, *et al.* (1999) The glycine residue in cyclic lactam analogues of
675 galanin(1-16)-NH₂ is important for stabilizing an N-terminal helix. *Biochemistry*
676 38(46):15295-15304.
- 677 25. Barany-Wallje E, Andersson A, Graslund A, & Maler L (2004) NMR solution structure
678 and position of transportan in neutral phospholipid bicelles. *FEBS Lett*
679 567(2-3):265-269.
- 680 26. Kruse AC, *et al.* (2013) Activation and allosteric modulation of a muscarinic
681 acetylcholine receptor. *Nature* 504(7478):101-106.

- 682 27. Shihoya W, *et al.* (2016) Activation mechanism of endothelin ETB receptor by
683 endothelin-1. *Nature* 537(7620):363-368.
- 684 28. Koehl A, *et al.* (2018) Structure of the micro-opioid receptor-Gi protein complex.
685 *Nature* 558(7711):547-552.
- 686 29. Hong C, *et al.* (2021) Structures of active-state orexin receptor 2 rationalize peptide
687 and small-molecule agonist recognition and receptor activation. *Nat Commun*
688 12(1):815.
- 689 30. Floren A, Land T, & Langel U (2000) Galanin receptor subtypes and ligand binding.
690 *Neuropeptides* 34(6):331-337.
- 691 31. Land T, *et al.* (1991) Linear and cyclic N-terminal galanin fragments and analogs as
692 ligands at the hypothalamic galanin receptor. *Int J Pept Protein Res* 38(3):267-272.
- 693 32. Jumper J, *et al.* (2021) Highly accurate protein structure prediction with AlphaFold.
694 *Nature* 596(7873):583-589.
- 695 33. Maeda S, Qu Q, Robertson MJ, Skiniotis G, & Kobilka BK (2019) Structures of the M1
696 and M2 muscarinic acetylcholine receptor/G-protein complexes. *Science*
697 364(6440):552-557.
- 698 34. Moro O, Lamah J, Hogger P, & Sadee W (1993) Hydrophobic amino acid in the i2
699 loop plays a key role in receptor-G protein coupling. *J Biol Chem*
700 268(30):22273-22276.
- 701 35. Xiao P, *et al.* (2021) Ligand recognition and allosteric regulation of DRD1-Gs
702 signaling complexes. *Cell* 184(4):943-956 e918.

- 703 36. Blakemore LJ & Trombley PQ (2017) Zinc as a Neuromodulator in the Central
704 Nervous System with a Focus on the Olfactory Bulb. *Front Cell Neurosci* 11:297.
- 705 37. Kay AR & Toth K (2008) Is zinc a neuromodulator? *Sci Signal* 1(19):re3.
- 706 38. Anderson CT, *et al.* (2015) Modulation of extrasynaptic NMDA receptors by synaptic
707 and tonic zinc. *Proc Natl Acad Sci U S A* 112(20):E2705-2714.
- 708 39. Kalappa BI, Anderson CT, Goldberg JM, Lippard SJ, & Tzounopoulos T (2015) AMPA
709 receptor inhibition by synaptically released zinc. *Proc Natl Acad Sci U S A*
710 112(51):15749-15754.
- 711 40. Vogt K, Mellor J, Tong G, & Nicoll R (2000) The actions of synaptically released zinc
712 at hippocampal mossy fiber synapses. *Neuron* 26(1):187-196.
- 713 41. Swaminath G, Steenhuis J, Kobilka B, & Lee TW (2002) Allosteric modulation of
714 beta2-adrenergic receptor by Zn(2+). *Mol Pharmacol* 61(1):65-72.
- 715 42. Holst B, Elling CE, & Schwartz TW (2002) Metal ion-mediated agonism and agonist
716 enhancement in melanocortin MC1 and MC4 receptors. *J Biol Chem*
717 277(49):47662-47670.
- 718 43. Nunez D, Kumar R, & Hanahan DJ (1989) Inhibition of [3H]platelet activating factor
719 (PAF) binding by Zn²⁺: a possible explanation for its specific PAF antiaggregating
720 effects in human platelets. *Arch Biochem Biophys* 272(2):466-475.
- 721 44. van der Westhuizen ET, Valant C, Sexton PM, & Christopoulos A (2015) Endogenous
722 allosteric modulators of G protein-coupled receptors. *J Pharmacol Exp Ther*
723 353(2):246-260.

- 724 45. Yuan Y, *et al.* (2021) Structures of signaling complexes of lipid receptors S1PR1 and
725 S1PR5 reveal mechanisms of activation and drug recognition. *Cell Res.*
- 726 46. Xu P, *et al.* (2021) Structures of the human dopamine D3 receptor-Gi complexes. *Mol*
727 *Cell*81(6):1147-1159 e1144.
- 728 47. Kato HE, *et al.* (2019) Conformational transitions of a neurotensin receptor 1-Gi1
729 complex. *Nature* 572(7767):80-85.
- 730 48. Liu Q, *et al.* (2021) Ligand recognition and G-protein coupling selectivity of
731 cholecystokinin A receptor. *Nat Chem Biol.*
- 732 49. Krishna Kumar K, *et al.* (2019) Structure of a Signaling Cannabinoid Receptor 1-G
733 Protein Complex. *Cell* 176(3):448-458 e412.
- 734 50. Kim HR, *et al.* (2020) Structural mechanism underlying primary and secondary
735 coupling between GPCRs and the Gi/o family. *Nat Commun* 11(1):3160.
- 736 51. Conklin BR, Farfel Z, Lustig KD, Julius D, & Bourne HR (1993) Substitution of three
737 amino acids switches receptor specificity of Gq alpha to that of Gi alpha. *Nature*
738 363(6426):274-276.
- 739 52. Conklin BR, *et al.* (1996) Carboxyl-terminal mutations of Gq alpha and Gs alpha that
740 alter the fidelity of receptor activation. *Mol Pharmacol* 50(4):885-890.
- 741 53. Semack A, Sandhu M, Malik RU, Vaidehi N, & Sivaramakrishnan S (2016) Structural
742 Elements in the Galphas and Galphaq C Termini That Mediate Selective G
743 Protein-coupled Receptor (GPCR) Signaling. *J Biol Chem* 291(34):17929-17940.
- 744 54. Zheng S, Abreu N, Levitz J, & Kruse AC (2019) Structural basis for KCTD-mediated

rapid desensitization of GABAB signalling. *Nature* 567(7746):127-131.

55. Zheng SQ, *et al.* (2017) MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nat Methods* 14(4):331-332.

56. Punjani A, Rubinstein JL, Fleet DJ, & Brubaker MA (2017) cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods* 14(3):290-296.

57. Scheres SH (2012) RELION: implementation of a Bayesian approach to cryo-EM structure determination. *J Struct Biol* 180(3):519-530.

58. Sanchez-Garcia R, *et al.* (2020) DeepEMhancer: a deep learning solution for cryo-EM volume post-processing. *BioRxiv*.

59. Pettersen EF, *et al.* (2004) UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem* 25(13):1605-1612.

60. Emsley P & Cowtan K (2004) Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol Crystallogr* 60(Pt 12 Pt 1):2126-2132.

61. Adams PD, *et al.* (2010) PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr* 66(Pt 2):213-221.

62. Inoue A, *et al.* (2019) Illuminating G-Protein-Coupling Selectivity of GPCRs. *Cell* 177(7):1933-1947 e1925.