1 **Factors affecting post-breeding endometritis, pregnancy rate and**
2 **embryonic/fetal death in sport mares in two French commercial stud farms:**
3 **special focus on age, parity and lactating status effects**

4 Emilie Derisoud^{a,b,+,*}, Bénédicte Grimard^{a,b,+}, Clothilde Gourtay^c, Pascale Chavatte-Palmer^{a,b,*}

5 ^aUniversité Paris-Saclay, UVSQ, INRAE, BREED, 78350, Jouy-en-Josas, France; ^bEcole Nationale
6 Vétérinaire d'Alfort, BREED, 94700, Maisons-Alfort, France ; ^cIFCE, Plateau technique du Pin, 61310,
7 Exmes, France

8 ⁺ Authors contributed equally to this work

9 ^{*}Corresponding author.

10 Emails:

11 emilie.derisoud@inrae.fr

12 benedicte.grimard@vet-alfort.fr

13 clothilde.gourtay@ifce.fr

14 pascale.chavatte-palmer@inrae.fr

15 Mail address:

16 Pascale Chavatte-Palmer

17 BREED (UMR1198) – Bâtiment 230

18 Biology de la reproduction, Environnement, Epigénétique et Développement

19 Domaine de Vilvert

20 78352 Jouy-en-Josas, FRANCE

21

22 **Abstract**

23 Fertility in mares depends on management and reproduction mode, but overall analyses have shown
 24 that there is an important and progressive decline in fertility in older mares. Age is often related to
 25 maternal parity, but the role of parity in the fertility decline in old mares is not known. Moreover, the
 26 positive or negative impacts of breeding mares yearly are a subject of controversy. The aim of this
 27 study was to identify whether these factors, as well as reproductive management, played a role on
 28 post-insemination endometritis and fertility in sports mares in commercial studs. Altogether, the
 29 breeding records of 277 mares (506 cycles) over one breeding season in two French commercial stud
 30 farms were analyzed. Age, parity, whether the mare was barren or suckling, follicular size before
 31 ovulation, the use of estrus and/or ovulation induction, artificial insemination (AI) protocol, post-
 32 breeding inflammation and treatment, Day 14 pregnancies and number of embryos, as well as
 33 subsequent foaling the next year were recorded. Data were analyzed using multivariate logistic
 34 regression models. The pregnancy rate by cycle was 41.9% and 76.5% of mares were pregnant at the
 35 end of the season. Mares older than 10 years, barren mares or mares inseminated in July or August
 36 were more likely to have inflammatory reaction after insemination ($p < 0.0001$, OR=3.29, 5.389 and
 37 3.329, respectively). Mares >10 years and mares inseminated with frozen semen were less likely to
 38 be pregnant on Day 14 vs younger mares and mares inseminated with fresh and refrigerated semen
 39 ($p < 0.05$, OR=0.622 and 0.582, respectively). As expected, mares with multiple ovulations were more
 40 likely to be pregnant on Day 14 compared to mares that had only one ovulation ($p < 0.05$, OR=1.791).
 41 Altogether, parity only tended to improve the likelihood of being pregnant on Day 14 ($p = 0.07$,
 42 OR=1.434 in parous vs nulliparous) but in older mares (age >10), parity increased pregnancy rates
 43 (44.09% vs 30.89% in parous vs nulliparous, $p < 0.05$). Post-breeding inflammation was not related to
 44 pregnancy rate on Day 14 (37.8% vs 44.2%, $p = 0.16$) but all mares with post-breeding inflammation
 45 received a treatment. No effect of suckling at insemination was observed. None of the studied factor

46 could be related to embryo/fetal mortality. In conclusion, maternal age seems to be more important
47 than parity and lactating status for reproductive success in mare. Cumulative effects on parity and
48 aging were observed on pregnancy rates.

49

50 **Keywords**

51 Horse, equine, reproduction, insemination, lactation, fertility

52

53

1. Introduction

Effectiveness of mare reproduction is an important economical factor in the horse industry. Breeding a mare is an expensive procedure with no guarantee: stallion fees, gynecological monitoring and mare maintenance are costly. Therefore, once mares begin their reproductive career, breeders generally aim to produce one foal per year per mare to remain profitable. This goal can only be achieved if mares remain fertile throughout their life span. Several factors may affect mare fertility such as their age, parity and lactation status during the breeding period.

The decline in mare fertility associated to age has been well documented in various breeds [1–18]. Moreover, Langlois et al. demonstrated that if fertility was the only factor for culling broodmares and if a 10% of fertility decline was chosen as cutoff value, then mares older than 8 should not be bred [2]. A survey sent to breeders participating in competing events, however, showed that mare's age was the least important consideration for the selection of mares for reproduction [19]. Moreover, since a mare's reproductive career often begins after their sporting career, it is common in sport horses to breed older mares. Indeed, in Canada, 37% of the standardbred mares were shown to be older than 11 years old at insemination [20]. These figures are even higher in Finland where 53% of Standardbred and 63% of Finnhorse mares exceed this age [3].

Mares can be bred shortly after foaling and breeders often aim to produce foals yearly, so that mares are often nursing at the time of breeding. Lactation affects mares' metabolism [21] but there is no consensus on whether it may induce effects on fertility and embryo/fetal loss [22–25].

As for lactation, the putative effects of mare's parity on fertility and embryo loss are controversial [2–5,9,10,12–16,18]. Horse production is unique in that mares may be both nulliparous and old at the same time, which is not the case in other animal production. As a matter of fact, in Finland, 20.5% of Finnhorse and 15.5% of Standardbred broodmares are both older than 10 and nulliparous [3].

77 Transient uterine inflammation is a normal physiological reaction in mares after breeding. It is
78 thought to be required for the elimination of bacteria and the excess of spermatozoa introduced in
79 the uterine lumen. For some mares, however, a persistent infection can develop and interfere with
80 fertility outcomes (for review [26]). Ultrasound images of uterine fluid accumulation are the most
81 commonly used diagnostic criterion for post-breeding endometritis. They are observed more
82 frequently in mares >8 years old compared to younger mares. In a study on 57 mares, Woodward et
83 al. observed that 45% (N=22) of the 8-16 years old mares and 88% of the older mares (≥17 years old,
84 N=26) had uterine fluid retention 48h after insemination with frozen semen, whereas none of the 9
85 younger mares (<8 years old) were affected [27]. To the authors' knowledge, the specific effects of
86 maternal parity and/or lactation on the post-breeding inflammation has not been explored.

87 The Thoroughbred studbook only allows hand breeding whereas the French Trotter studbook allows
88 insemination but only with fresh semen. There is no such limitation in sport horse breeds but studies
89 on fertility in sports mares are scarce and relatively ancient. There is no recent study considering at
90 the same time the method of reproduction and the effects of age, parity and lactation on mare
91 fertility.

92 The aim of this study was thus to identify effects of mare age, parity and nursing on post-breeding
93 endometritis, fertility and embryonic/fetal death in commercial stud conditions for sport mares.

94 **2. Material and methods**

95 **2.1. Ethics**

96 This is an observational study performed in commercial farms. All procedures were carried out
97 according to the European Union Directive 98/58/CE concerning farm animals protection and the
98 French legislation about animal care in rural codification.

99 **2.2. Animals**

Data were recorded in 2 French commercial stud farms, one located about 60km south west of Paris, the other 150 km south of Paris, both managed by one artificial insemination cooperative during one breeding season (2019).

From beginning of February to late August 2019, the study enrolled 277 sport mares over 506 estrus periods. For each mare, date of birth, breed, number of previous breedings and foalings were extracted from the national database (Infochevaux: <https://infochevaux.ifce.fr/fr/info-chevaux>). Mares were clustered according to age in two (≤ 10 years old vs > 10 years old) or four classes (≤ 5 years old,]5, 10],]10, 15], > 15 years old). They were also clustered in 2 classes according to previous fertility (number of previous foaling/number of previous breedings < 0.7 or > 0.7) and in 2 classes according to parity: nulliparous (never foaled before) vs parous (foaled at least once before) at the beginning of the breeding season.

2.3. Breeding management

2.3.1. Estrus monitoring

First, luteolysis was induced in mares that had a persistent corpus luteum or that were sent for breeding late in the season, using prostaglandin F₂ α analogs: Alfaprostol (Alfabedyl®) or Luprostol, (Prosolvin®, 1 to 1.5 mL IM according to mare size).

Estrus detection was performed using uterine horn firmness as estimated by rectal palpation together with ultrasound examination where the uterine oedema score (0-5) [28] and the largest follicle diameter were determined. Mares were monitored once or twice daily (according to breeding method, see below) from the time when the largest follicle diameter reached 35 mm and until ovulation was detected.

As ovulation approached (follicle diameter ≥ 35 mm and reduction of the uterine oedema score), ovulation was induced with hCG (Chorulon®, 1500 to 5000 IU IM related to mare's weight) or with a GnRH analog (Decapeptyl®, 0.1 mg triptorelin in 1 mL SC). The GnRH analog option was chosen when mares had a known resistance to hCG or when hCG had already been used before during the breeding season and was not effective to induce ovulation.

2.3.2. Insemination management

All mares were bred using artificial insemination (AI). Mares were inseminated with fresh (FAI), cooled transported (CTAI) or frozen semen (FZAI). Mare management differed according to the type of semen used.

Mares entitled to be inseminated with fresh semen were examined once daily during estrus. A first insemination was performed before ovulation (follicle diameter ≥ 35 mm and decrease in the uterine oedema score) and mares that had not ovulated 48h later were inseminated a second time. This was repeated until ovulation was observed.

Mares entitled to be inseminated with cooled transported semen were also examined once daily and inseminated before ovulation but subsequent inseminations were performed every 24h until ovulation.

Mares bred with frozen semen were examined every 12h if the number of available straws was >4 while examinations were performed every 6 h when the number of available straws was ≤ 4 . In this latter case, deep horn insemination was performed. The aim was to inseminate just before ovulation, as determined by ultrasound observation of preovulatory follicular wall thickening and follicle deformation, or just after ovulation.

2.3.3. Post-breeding endometritis and pregnancy diagnosis

All mares were examined by ultrasound the day after insemination. If uterine oedema and/or fluid accumulation were observed, mares were treated either with a single dose of oxytocin (Oxytocine S©, 10-20 UI in 1 to 1.5 mL IM or IV), or oxytocin in association with uterine lavage (one or two liters of warm, sterile saline solution), or in association with antibiotics or by uterine lavage alone. Treatment was decided by the veterinary surgeon and depended on the volume of fluid accumulated.

Pregnancy was assessed 14 days (D14) after AI by ultrasonography. In the case of twin pregnancy, squeezing was recommended to the breeder but not always performed. Pregnancy was confirmed on Day 30 if the mare was brought back to stud. In addition, some mares came back again in autumn for late pregnancy diagnosis.

2.3.4. Data recording

All data were recorded in the same dedicated software (Gynebase©, Equidéclit, France) by the veterinarians and technicians of the 2 studs.

Age, parity, lactational status at breeding (nursing vs non nursing), number of monitored cycles for each mare, estrus induction (yes/no), heat duration (number of days of observation by ultrasonography), size of the preovulatory follicle, induction of ovulation (yes/no), hormone used to induce ovulation (hCG or GnRH analog), ovulation observed (yes/no), number of follicles ovulated, AI mode (fresh semen, cooled transported or frozen semen), number of AI during estrus, date of insemination (last AI if more than one AI were performed during estrus), stallion identity, number of straws used (for frozen semen), uterine fluid accumulation after AI (yes/no), treatment used (oxytocin alone vs other), presence of single or twin embryos, squeezing when twins were diagnosed (yes/no), pregnancy diagnosis on D14 as well as on D30 or later, when possible, were recorded. The number of the monitored cycles needed to obtain a gestation was classified as 1 vs >1 when

pregnancy was achieved after more than one cycle. The number of straws used for frozen insemination was classified as either <8 vs ≥ 8 or ≤ 4 vs > 4 . Cutoff values of 4 and 8 were chosen according to previous (8 straws for 400 million mobile spermatozoa) and the current (4 straws of 50×10^6 mobile spermatozoa, 4 mL each) recommendation of the French Institute for Horses and Horse-riding (IFCE) for frozen semen.

Pregnancy rate was calculated as the ratio between the number of pregnant mares at the end of the breeding season and total number of mares bred (overall pregnancy rate at the end of the season) and as the ratio of number of positive diagnoses on D14 / total number of used cycles (pregnancy rate per cycle). Embryonic loss was estimated by the number of mares not pregnant on D30 after a positive pregnancy diagnosis on D14. Late embryonic loss was calculated by the number of mares not pregnant during the autumn after a positive pregnancy diagnosis on D14. Foal birth was checked the next year using the national database. Total embryonic/fetal loss was defined as the number of mares that did not foal after a positive pregnancy diagnosis on D14. Mares were only considered as non-foaling when abortion/stillbirth or new breeding without foaling were recorded in the national database in 2020.

2.4. Statistical analysis

Data were analyzed using SAS® Studio 3.8 (SAS® University Edition).

2.4.1. Univariate & multivariate analysis

Univariate analysis was used to evaluate the effects of qualitative variables on (1) the incidence of post-breeding endometritis, on (2) pregnancy rate and (3) embryonic/fetal loss. For (1) and (2), all

recorded cycles were used while for (3), only pregnant mares on D14 were considered. The statistical unit was, therefore, the cycle for (1) and (2) while it was the mare for (3).

For each criterion analyzed, results were compared between the different classes using a Chi-Squared test. Variables associated at $p < 0.20$ were included in the second step of the analysis.

As a second step, multivariate analysis was conducted using logistic regression (GLIMMIX procedure of SAS® Studio). Individual effects were considered by including mare identity in each model as a random effect. A backward stepwise elimination of non-associated ($p > 0.10$) variables was performed to develop multivariate models. Models presenting the lowest Akaike's Information Criterion were retained.

For these analyses, breeds and stallions could not be used as their numbers were too high without enough repeats to be considered in the statistical analysis.

For quantitative data, results are presented as mean $\pm$ standard error.

2.4.2. Analysis of age, parity and nursing status

The analyses were performed on data, using the estrous cycle as the reference unit. Young (≤ 10 years) vs old (> 10 years), nulliparous vs parous, nursing vs non nursing mares were compared using T test for quantitative variables and Xi Square for qualitative ones. Using Xi Square analysis, effects of mares' suckling status on post-breeding inflammation and effects of mares' parity on Day 14 pregnancy rates were assessed in each population of young or old mares.

2.4.3. Data availability statement

Data supporting the conclusions of this article are available in the Data INRAE repository, available at:

<https://doi.org/10.15454/4OIUKH>

3. Results

3.1. Descriptive analysis

3.1.1. Population of study

The 277 mares belonged to 32 breeds approved by the French Institute of the Horse (IFCE), with French saddlebred (Selle français) being the most represented breed (56% of mares included in this study). No other breed reached more than 15 individuals. Eight mares did not belong to any French approved breed. In addition, 13 mares belonged to racehorse breeds (3 English Thoroughbreds, 10 French Trotters). There were also 22 ponies and one draft horse. All mares were used for sport and leisure.

Characteristics of the 277 mares are detailed in Table 1. The average age was 12.7 years old (min 2, max 23, $\sigma = 4.8$). Altogether, 196 mares had been bred at least in one previous breeding season and the overall average of previous breeding seasons was 2.5 (min 0, max 17, $\sigma = 3.0$). For previously bred mares, mean fertility (number of foalings/number of breeding) was 0.71 (min 0, max 1, $\sigma = 0.31$). The interval between previous breeding and the 2019 reproductive season ranged from 1 to 13 years. Altogether, 66 (21.9%) mares were nulliparous and aged more than 10 years and 44 (15.8%) were suckling and older than 10 years.

Table 1: Characteristics of the 277 mares bred in 2019.

Variable	n	Percent
Age in 2 classes (Years)		
≤ 10	96	34.7

> 10	181	65.3
Age in 4 classes (Years)		
≤5	24	8.7
5-9	72	26.0
10-14	94	33.9
> 15	87	31.4
Parity		
Nulliparous	98	35.4
Parous	179	64.6
Lactation status in 2019		
Suckling	76	27.4
Non suckling	201	72.6
Fertility at previous breeding (n=196)		
< 0.7	82	41.8
≥ 0.7	114	58.2
Month of first AI		
March	36	13.0
April	70	25.3
May	86	31.0
June	67	24.2
July - August	18	6.5

232

233 **3.1.2. Pregnancy rate per mare**

Mares were bred with 154 different stallions. Altogether, 212 mares were diagnosed pregnant on D14 at the end of the season (76.5% overall fertility rate). These performances were obtained within an average of 1.8 cycle per mare (min 1, max 6). From the 154 mares that came back for pregnancy confirmation, 130 were still pregnant at Day 30, so that embryonic loss was 9.1%. Only forty-six mares came back later in the autumn, of which 44 mares were confirmed still pregnant (late embryonic loss: 4.3%).

Foaling information in 2020 were obtained for 206 pregnant mares on Day 14. Among them, 168 foalings were recorded indicating a total embryonic/fetal loss of 18.4%.

3.1.3. Breeding management

Data recorded per estrus period are presented on Table 2.

Among the 506 estrus recorded periods, 92 were induced by luteolysis of a previous corpus luteum using prostaglandins (18.2%). Data were recorded in average during 5.1 days per estrus period (min 1, max 16, $\sigma = 2.1$).

Of the 506 estrous periods, fresh semen was used for 90 cycles (FAI, 17.8%), cooled transported semen for 142 cycles (CTAI, 28.1%) and frozen semen for 274 (FZAI, 54.1%). Ovulation was induced in 360 estrous periods (71.2%). The induction of ovulation was performed in 40.0% of the FAI, 74.0% of the CTAI and 80.0% of the FZAI. Mostly hCG was used (n=322, 89.4%) with GnRH analog used in the other cases (n=38, 10.6%). Preovulatory dominant follicle mean diameter was 42.4 mm (min 28, max 60, $\sigma = 4.8$). Mares were inseminated in average 1.2 times per estrus (min 1, max 5, $\sigma = 0.6$). For FZAI, the mean number of straws used was 5.1 (min 1, max 12, $\sigma = 2.4$, recorded in 251 AI by 274).

Ovulation was observed by ultrasonography in 501 of the 506 estrous periods (99%). In most cases, only one ovulation was observed (n=425, 84.8%). Double or triple ovulations were observed in 75 (15.0%) and 1 estrus, respectively.

After insemination, uterine fluid accumulation was observed in 180 estrous periods (35.6%). Oxytocin was used for 173/180 treatments, mostly alone (139/180), sometimes associated with uterine lavage (27 cases). Antibiotics were used for local treatment (10/180) associated to oxytocin (n=3), to uterine lavage (n=4) or alone (n=3).

After AI, 212 cycles among the 506 led to a pregnancy on D14 (pregnancy rate per cycle: 41.9%). Twins were detected in 27 cases (5% of breeding and 12.7% of pregnancies). Squeezing was performed for 25/27 pregnancies. After squeezing, pregnancy was checked on Day 30 and 21/24 mares were still pregnant (87.5%).

Table 2: Characteristics of the 506 monitored estrus

Variable	n	Percent
Cycle number		
1	277	54.7
2	144	28.5
3	65	12.8
4	18	3.7
5	1	0.2
6	1	0.2
Luteolysis before AI		
No	414	81.8
Yes	92	18.2
Induction of ovulation		
No	146	28.8
Yes	360	71.2

Number of observed ovulations (n=501)		
1	425	84.8
2	75	15.0
3	1	0.2
Method of semen preservation		
Fresh semen (FAI)	90	17.8
Cooled transported (CTAI)	142	28.1
Frozen (FZAI)	274	54.1
Number of straws (n= 251)		
< 8	177	70.5
≥ 8	74	29.5
Number of straws (n= 251)		
≤ 4	126	50.2
> 4	125	49.8
Month of AI		
March	36	7.1
April	104	20.6
May	150	29.6
June	155	30.6
July-August	61	12.1
Number of AI by estrus period		
1	431	85.2
2	57	11.2
3	11	2.2
4	5	1.0
5	2	0.4

Observed ovulation

Yes	501	99.0
No	5	1.0

Uterine fluid accumulation/endometritis after AI

Yes	180	35.6
No	327	64.4

Uterine treatment (n=180)

Oxytocin alone	139	77.2
Other	41	22.8

268

269 3.2. Univariate & multivariate analysis

270 3.2.1. Factors associated with post-breeding inflammation

271 After univariate analysis, variables associated with post-breeding inflammation/endometritis at the
 272 threshold of 20% were: mare's age (using 10 years old as cutoff value), parity, lactation (suckling vs
 273 non suckling), induction of estrus by luteolysis, cycle number and month of AI (Figure 1). Post-
 274 breeding inflammation was not associated with semen conditioning (33.6, 40.2 and 37.9%
 275 respectively for FAI, CTAI and FZAI, $p=0.45$), number of AI per estrus (35.2% for 1 AI vs 40.3% for
 276 more than one AI per estrus, $p=0.40$), nor number of straws used (33.9 vs 33.8% for number of
 277 straws < 8 vs ≥ 8 , $p=0.98$ and 37.3 vs 30.4% for number of straws ≤ 4 vs > 4 , $p=0.25$).

278 After multivariate analysis, only 3 variables were significantly associated ($p<0.05$) with post-breeding
 279 inflammation: mare age, lactation and month of AI (Supplementary table 1). Inflammation was more
 280 frequent in mares older than 10 years than in younger mares (Figure 1). Non suckling mares at the
 281 time of insemination were also more affected in comparison with suckling ones (Figure 1). There was
 282 no interaction between age and nursing status ($p = 0.38$) but increased age amplified the number of

post-breeding inflammations in both non-suckling and suckling mares. Respectively, 22.9% and 8.2% of the non-suckling and suckling young mares were affected by post-breeding inflammation while 50.7% and 18.3% of mares older than 10 years old were affected.

The risk of inflammation was also increased in July and August compared to previous months (Figure 1).

3.2.2. Factors associated with pregnancy rate per cycle

After univariate analysis, the following factors were associated with pregnancy rate at the 20% threshold: age in 2 classes, parity, cycle number, AI modality, month of insemination, number of observed ovulations and post-breeding inflammation. All the associated variables were introduced in multivariate models. As significant effect of number of ovulations was observed after univariate analysis, only estrous cycle with ovulations that were observed by ultrasonography were considered (*i.e.*, 501 estruses among the 506 recorded).

Stepwise regression and backward elimination led to a model containing only 3 significant variables ($p < 0.05$): mare age, number of ovulations and semen conditioning (Supplementary Table 2). Data that significantly influenced pregnancy rates are summarized in Figure 2. Pregnancy rate was higher in mares younger vs older than 10 years mares. Pregnancy rate was increased when a multiple vs single ovulation was observed. FAI and CTAI resulted in more pregnancies on Day 14 than FZAI. Trends were observed ($p < 0.10$) for the effects of month of AI and parity. Pregnancy rate tended to be higher in April, May, July and August vs March and June. Parous mares tended also to have better pregnancy rates than nulliparous mares ($p = 0.07$, OR=1.434 in parous mares).

The study of the interaction between maternal age and parity showed that in mares aged of 10 years or less, being nulliparous or parous did not alter pregnancy rates (49.25 and 46.15% of pregnancy rate per cycle for young nulliparous and multiparous respectively, $n = 158$, $p = 0.70$). In mares older

than 10 years old, however, nulliparity accentuated the decrease in pregnancy rates. Indeed, the pregnancy rates per cycle was only 30.89% for old nulliparous mares vs 44.09% for old multiparous mares (OR = 1.96 in parous mares, $p = 0.016$).

After multivariate analysis, pregnancy rate was not significantly affected by post-breeding inflammation (37.8% of pregnancy rate after inflammation vs 44.2% in healthy mares, $p = 0.16$). In treated mares, treatment after post-breeding treatment did not affect pregnancy rate (37.4% of pregnancy for treatment with oxytocin alone, $n = 139$ vs 39.0% for other treatment, $n = 41$, $p = 0.85$).

3.2.3. Factors associated with embryonic/fetal loss between Day 14 and foaling

After the univariate analysis, age in four classes, induction of ovulation and month of insemination were associated with embryonic/fetal loss at the threshold of 20%. However, after multivariate analysis, none of these factors affected embryonic/fetal loss.

3.3. Focus on maternal age, parity and suckling status

3.3.1. Effect of maternal age

Mares 10 years old or younger were 7.2 ± 0.2 years old while mares older than 10 years were 15.6 ± 0.2 years old.

Neither parity, induction of the estrus cycle or of ovulation, month of AI, number of cycles required for a gestation at Day 14, number of straws in case of FZAI or the number of twin embryos at Day 14 were significantly related to maternal age (Supplementary Table 3).

Nevertheless, more young mares were pregnant at Day 14 and less had a post-breeding inflammation compared to old mares (Table 3). In addition, 31% of young were nursing while it represented only 20.7% of old mares ($p < 0.05$). Multiple ovulations occurred more frequently in old

330 mares compared to young mares. Young mares tended to be more inseminated using fresh and
331 frozen and less with refrigerated semen compared to older mares.

332 Estrus tended to be longer in old vs young mares (respectively, in average, 5.2 ± 0.1 vs 4.8 ± 0.2 days,
333 $p = 0.086$). The preovulatory follicle diameter tended to be smaller in old compared to young mares
334 (respectively, in average, 42.2 ± 0.2 vs 43.0 ± 0.4 mm, $p = 0.086$).

335 **Table 3: Characteristics in young (≤ 10 years old) vs old mares (> 10 years old)**

Variable	n	Young ≤ 10 years old n=158	Old > 10 years old n=343	p
Pregnancy rates on Day 14	501	47.5%	39.4%	0.021**
Post-breeding inflammation	501	18.3%	44.0%	$< 0.0001^{**}$
Lactation				
Non-nursing	381	69.0%	79.3%	0.012
Nursing	120	31.0%	20.7%	
Method of the semen preparation				
*				
FAI	87	19.0%	16.6%	0.093
CTAI	140	21.5%	30.9%	
FZAI	274	59.5%	52.5%	
Number of ovulations per cycle				
1				
> 1	425	90.5%	82.2%	0.016
	76	9.5%	17.8%	

* AI: Artificial insemination, FAI: insemination with fresh semen, CTAI: insemination with cooled transported semen, FZAI: insemination with frozen semen

** Results from multivariate analysis

3.3.2. Effect of maternal parity

Neither age, induction of the estrus cycle, semen preparation, month of AI, number of cycles or AI required for a gestation at Day 14, number of straws in case of FZAI nor the number of ovulation or twin embryos at Day 14 were significantly affected by maternal age (Supplementary Table 4).

Nulliparous mares tended to be less pregnant on Day 14 (Figure 2). Parity tended to be related to post-breeding inflammation. Maternal parity was different according to maternal age. More nulliparous mares were present in 5 or less and 10-15 years old group while less nulliparous were observed in 5-10- and 16 or more-years old groups. Nulliparous mares were mostly older than 10 years (64.7%, Supplementary Table 4) but nulliparous were in average younger than parous mares (Table 4) with a mean 2-year difference between nulliparous and parous mares (in average, respectively, 11.6 ± 0.3 vs 13.7 ± 0.3 years old, $p < 0.0001$). Moreover, suckling status was obviously related to parity and 38.6% of parous mares were suckling at insemination. Ovulation was induced more frequently in parous than in nulliparous mares. Parity did not modify estrus duration (5.1 ± 0.2 days for nulliparous vs 5.1 ± 0.1 days for parous mares, $p = 0.11$) but the size of the preovulatory follicle at ovulation was reduced in nulliparous compared to parous mares (respectively, 41.9 ± 0.3 and 42.8 ± 0.2 mm in diameter, $p=0.03$).

Table 4: Characteristics in nulliparous vs parous mares

Variable	n	Nulliparous	Parous	p
----------	---	-------------	--------	---

		n=190	n=311	
Pregnancy rates on D14	501	37.4%	44.7%	0.066**
Age				
≤ 5	40	17.4%	2.3%	< 0.0001
5 < Age ≤ 10	118	17.9%	27.0%	
10 < Age ≤ 15	185	43.2%	33.1%	
Age > 15	158	21.6%	37.6%	
Lactation				
Non nursing	381	100.0%	61.4%	< 0.0001
Nursing	120		38.6%	
Induction of ovulation				
No	145	35.8%	24.8%	0.008
Yes	356	64.2%	75.2%	

357 AI: Artificial insemination, FAI: insemination with fresh semen, CTAI: insemination with cooled

358 transported semen, FZAI: insemination with frozen semen

359 ** Results from multivariate analysis

360

361 3.3.3. Effect of nursing at insemination

362 Neither induction of the estrus cycle nor ovulation induction, semen preparation, number of AI per

363 estrus required for a gestation at Day 14, number of straws in case of FZAI nor the number of

364 ovulations or twin embryos at Day 14 were significantly affected by maternal suckling status

365 (Supplementary Table 5).

Significantly less suckling mares were affected by post-breeding infection than non-suckling ones (Table 5) but pregnancy rates were not affecting by lactational status (45.8% and 40.7% of pregnancy rate per cycle for, respectively suckling and non-suckling mares, $p = 0.32$). There were more suckling mares younger than 10 years of age and more non-suckling mares aged more than 10 years. Mares younger than 5 and mares aged between 10 to 15 years were less frequently nursing than mares aged 5-10. In the group of 5-10 years old, more suckling mares were present in comparison to non-suckling ones while the lactational status was similarly represented in 16 years old and older mares. In average, however, the age of suckling vs non-suckling mares was not different (respectively 13.0 ± 0.2 vs 12.6 ± 0.2 years old, $p = 0.39$). Less nursing mares were bred in March and April but more were bred in June than non-nursing ones (Table 5). Moreover, more nursing mares were pregnant at Day 14 within the first exploited cycle than non-nursing ones. Estrus was shorter (respectively, 4.7 ± 0.16 vs 5.2 ± 0.1 days, $p = 0.03$) and preovulatory follicle diameter was larger (respectively 43.8 ± 0.4 vs 42.0 ± 0.2 , $p < 0.0001$) in nursing vs non-nursing mares.

Table 5: Characteristics in non-suckling vs suckling mares

Variable	n	Non suckling n=381	Suckling n=120	p
Post-breeding inflammation	501	42.8%	14.2%	< 0.0001
Age				
≤ 10 years old	158	28.6%	40.8%	0.012
> 10 years old	343	71.4%	59.2%	
≤ 5	40	9.7%	2.5%	< 0.0001
5 < Age ≤ 10	118	18.9%	38.3%	

10 < Age ≤ 15	185	39.6%	28.3%	
Age > 15	158	31.8%	30.8%	
Month of AI				
March	35	8.1%	3.3%	0.005
April	103	22.8%	13.3%	
May	149	29.7%	30.0%	
June	153	26.7%	42.5%	
July - August	61	12.6%	10.8%	
Number of cycles for pregnancy				
1	276	52.5%	63.3%	0.037
> 1	225	47.5%	36.7%	

* AI: Artificial insemination, FAI: insemination with fresh semen, CTAI: insemination with cooled transported semen, FZAI: insemination with frozen semen

383

384 4. Discussion

385 Data presented here are summarized in Figure 3.

386 4.1. Post-breeding inflammation

387 4.1.1. Prevalence

388 In this study, more than a third of the monitored cycles were followed by post-breeding uterine fluid
389 accumulation. In a normal population of Thoroughbred mares, around 15% of mares are susceptible
390 to post-breeding inflammation that persists for several days [29]. In a more recent study in
391 Thoroughbreds in UK, however, post-breeding fluid accumulation occurred in 47.7% of analyzed

pregnancies [17]. The observed rate in the present study is therefore in the range of the literature. In comparison to other studies, however, the post-breeding inflammation rate was maybe overestimated. Indeed, as mares could be kept on the breeding stud for several cycles, the same mares could have had several inflammations over the entire breeding period, as this condition is generally persistent [30].

4.1.2. Effect of maternal age, parity and lactation status

In the studied population, there was twice more post-breeding endometritis in mares older than 10 years than in younger mares. These results agreed with literature. Indeed, one recent study in Thoroughbred stud farms showed that the percentage of mares with post-breeding intrauterine fluid accumulation [17], as well as the frequency of endometritis in other breeds, increase with mare age [27,31–35].

In addition, suckling at insemination appeared to have a protective effect against post-breeding endometritis. To the authors' knowledge, there is no study on the effect of nursing at insemination on post breeding endometritis. Since lactating mares are producing endogenous oxytocin, the well-known and widespread treatment against post-breeding inflammation [36] could explain the observed improvement in nursing mares. Indeed, nursing induces oxytocin release in the blood circulation, reaching a peak of around 10mIU/L in nursing pony and broodmares [37,38]. Plasma oxytocin concentrations of 10-7 mIU/L were reported to be present 20min after the beginning of the suckling period [37]. Since suckling bouts last around 20min during the first month after birth, and since foals nurse at hourly intervals, the release of oxytocin happens in a regular manner throughout the day [38].

4.1.3. Effect of the method of semen preservation and semen quantity

In this study, nor semen preservation method nor the volume determined by number of frozen straws, nor the number of inseminations were related to post-breeding endometritis. One study showed that a similar volume of uterine fluid accumulated in the uterus after insemination with cooled or frozen semen which is consistent with results obtained here [39]. One recent study, however, showed that the uterine inflammatory response was positively correlated to the number of spermatozoid used for the insemination with frozen semen [40]. This high inflammatory response was also faster to resume than the response for low doses of semen, which could explain the lack of influence of the number of straws in the present study.

4.1.4. Effect of month of insemination

Data showed that there were more inflammations/endometritis when insemination was performed late in the season, i.e., during July and August. Several hypotheses could explain this result. The first is that mares with good quality uteri needed less cycles to become pregnant and therefore that more mares with reproductive troubles were inseminated late in the season. Another hypothesis could be that there is more post-breeding inflammation from June/July onwards because of the increased environmental temperature. So far, there is no study on the effects of outdoor temperature on the incidence of endometritis in mares but in dairy cows, environmental heat, as evaluated by comparing the possibility of sheltering or not from the sun, increased rectal temperature by almost one degree [41] and reduced uterine blood flow [42]. The temperature increase, by acting directly on uterine temperature and/or on uterine blood flow could therefore promote inflammation in the mares' uterus.

4.1.5. Treatment and pregnancy rates

In the present study, when a post-breeding inflammation was detected, treatment was systematically applied. Most of the time, this treatment was limited to an injection of oxytocin. In a recent study about therapeutic practices in intensively managed Thoroughbred mares, almost half of the pregnant mares were treated with intrauterine antibiotics and the same proportion was treated with oxytocin. Oxytocin combined with intrauterine antibiotics are used prophylactically in Thoroughbreds in the UK to avoid uterine infections that breeders believe to be a cause of conception failure and embryo loss [17]. In Thoroughbreds, however, less than 10% of early pregnancy failures were associated with uterine inflammation [17,32]. Another study on 99 Thoroughbred mares showed that there was no association between post-breeding uterine fluid accumulation and pregnancy rates nor embryonic death [43], as observed here. In all studies, the systematic management to prevent/cure anormal inflammation could explain this lack of association. Therefore, today, the post-breeding inflammation seems to be well handled by stud farms.

In different breeds, the combination of antibiotics with oxytocin was shown to be more efficient than antibiotics or oxytocin alone to prevent decreased pregnancy rates after mating [44]. In another study on warmblood mares artificially inseminated, however, oxytocin alone was sufficient to increase pregnancy rates in comparison to no treatment [36]. In most of cases, here, oxytocin alone was applied and was sufficient to avoid altered pregnancy rate. This, therefore suggested that this treatment alone is efficient to avoid decreased pregnancy rates in artificially inseminated mares.

4.2. Pregnancy rates

4.2.1. Prevalence

In studies on different breeds in several countries using artificial insemination, pregnancy rates were between 40-80%, according to the semen preservation method [15,45–47]. In a comparable recent study on 328 sport mares artificially inseminated in the Netherlands, 46.6% of gestations were

obtained on Day 12 – 18 [48]. Here, pregnancy rates per cycles were similar with 41.9% of mares being pregnant by cycle regardless of the method used for semen preservation.

4.2.2. Effect of maternal age, parity and lactation status

Pregnancy rates were reduced when mares were older than 10 years old at the time of insemination. Several studies already showed a reduction of pregnancy rates with increased maternal age [1,7,10,31,49–51]. Most studies, however, observed differences in mares older than 14 years old and not as early as 10 years old. Here, the clustering in 4 classes (≤ 5 , 5-9, 10-14 and ≥ 15 years old) was not associated with alteration of pregnancy rates. This could be because pregnancy rates were only slightly reduced with age and due to the limited number of cycles studied, the present study was not able to demonstrate these changes.

Maternal parity tended to not reduce pregnancy rates in the overall population. Effect of maternal parity on pregnancy rates are controversial and related to mares' age. Indeed, when nulliparous mares were mostly older than 7 years, a detrimental effect of nulliparity was observed [2,15] but when nulliparous mares were younger, no deleterious or a favorable effect of parity was observed [2,14]. Here, more than 60% of nulliparous were older than 10 years and this could explain the observed tendency.

When considering only mares older than 10 years, D14 pregnancy rates were reduced in nulliparous mares while in the youngest group, the parity did not affect the incidence of pregnancy. Thus, the present data indicate a deleterious cumulative effect of nulliparity and aging on fertility in mares.

Lactation at the time of insemination did not influence pregnancy rates, 14 days post ovulation. Foaling mares, however, needed less cycles to become pregnant. In the literature, the effect of lactation on fertility is controversial as some studies observed that nursing mares are more fertile than non-nursing ones [2,8,31,50,51] while others do not observe any difference [4,7,10,14].

Insemination at foal heat was previously shown to reduce pregnancy rates [24,52] and to be associated with increased embryonic death [22,23,52]. Here, most foaling mares were bred on foal heat but this did not affect pregnancy rates.

4.2.3. Effect the number of ovulations

The more ovulations were observed, the higher the likelihood that the mare was to be pregnant at 14 days post ovulation. The result obtained here seem obvious as double ovulations increase the number of oocytes that could be fertilized.

4.2.4. Effect of the method of semen preservation and semen quantity

As previously observed [45,46,48,53], the modalities of insemination influenced the probability for a mare to be pregnant on Day 14. Indeed, pregnancy rate was higher after FAI or CTAI than after FZAI. It is well known that stallion semen quality decreases with cryopreservation and that a higher critical number of mobile spermatozoa per dose is required to reach same pregnancy rates than with fresh semen [53]. In this study, the number of straws was not related to pregnancy rate, as also shown by others [45]. The protocol involving ultrasonography every 6h and deep intra-uterine AI after observed ovulation seemed to be effective to reach acceptable pregnancy rates with 4 straws or less.

4.2.5. Effect of month of insemination

Month of insemination tended to influence the pregnancy rate, with reduced incidence of pregnancies in March and June. In another French study, March was one of the more prolific month in terms of foal productivity per mare but all breed were considered (Thoroughbred, Standardbred

and sport bred) [2]. In racehorse breeding, it is common to use light to advance the breeding season [54] since foals born early in the year have an advantage when they are sold as yearling. In sport horse breeding under European latitudes, however, March is often the beginning of the breeding season. At this time, mares enter the spring transitional period and less mares are bred, which could explain the reduced pregnancy rates on this month. The other French study also reported a fertility decrease when mares were bred in June and after [2]. The June effect, here, is more complicated to explain as July and August did not affect pregnancy rates. During their stay, all mares were housed in individual stable with no access to fresh grass. They were fed with hay. A change in food quality in June could not explain the present results. Nevertheless, increasing ambient temperature could explain the differences in pregnancy rates as temperatures reached a maximum of 35.8°C in one stud farm in June 2019. In July and August, temperatures were also high with more than 33°C as maximal temperature in the 2 studied stud farms. The lack of difference in July and August could be explained by the fact that only few mares were bred during this period and that technicians and veterinarians were maybe more caring.

4.3. Effects of age on other reproductive outcomes

4.3.1. Choice of semen preservation method

Maternal age did not influence the breeders' choice concerning ovulation management but influenced the modality of AI. Indeed, less frozen and fresh semen were chosen for the insemination of older mares. On one side, frozen semen is less efficient for producing a foal, as confirmed by literature [53] and this study. Moreover, reduced fertility has been observed in old mares, both in literature [for review see 47] and in the present study. The combination of both factors could explain that older mares tended to be inseminated less with frozen semen as the financial risk might have been too costly for breeders. Moreover, currently, equine breeders do not use more straws for old than for young mares, even if it has been shown that increasing straw number could improve fertility

[53]. On the other side, as more than half of the mares older than 10 years were parous in the present study, breeders could have been looking for stallions that are not available on the reproductive centers for fresh insemination to look for genetic diversity, thus explaining why cooled transported semen was preferred.

4.3.2. Length of estrus and ovulation outcomes

This retrospective study shows that the length of estrus tended to increase with mare age (only 0.4 day more in old mares), with a smaller preovulatory follicle. This was already observed as a prolonged interovulatory interval was associated with a prolonged follicular growth [56–63] and to a reduction in follicular growth and preovulatory follicular size in oldest mares [64–67].

In the studied population, more multiple ovulations were also observed in older mares. Several studies reported that the incidence of multiple ovulation continuously increases until the mare reaches 20 years of age [62,67–69]. More multiple ovulations in older mares, however, is not associated with more twinning at 14 days post ovulation. Studies have shown that even if fertilization appears to be equal between young and old mares, early embryo mortality is increased in older mares [49,70] and this could explain that there are no more twin embryos observed at 14 days in older mares than in younger ones.

4.4. Effects of parity on other reproductive outcomes

Nulliparous mares had smaller preovulatory follicles. As explained above, the monitoring of mares was not performed more than twice a day: since follicles continuously grow until breakdown, this observed difference should be considered with caution.

4.5. Effects of suckling at insemination on other reproductive outcomes

In sport horses, mares are often bred and subsequently foal during the spring. Foaling mares are most of the time pregnant at the beginning of the spring. It is, therefore, not surprising to observe a peak of breeding in May and June in foaling mares while barren mare breedings were more spread throughout the reproductive season. Moreover, as it appears that less estrus cycles were required to obtain a pregnancy in lactating mares, it is not surprising either to not observed many lactating mares in August.

Finally, the heat period was shortened and the preovulatory follicle was larger when mares were in lactation without changing the number of ovulations. To the author knowledge, there is no study on the effect of lactation at insemination time on estrus and follicle parameters.

5. Conclusion

In conclusion, maternal age appeared to be the most important factor affecting both post-breeding inflammation and pregnancy rates. Both could be explained by the degenerative changes that are observed in older mares. Breeders, should, therefore, be encouraged to pay more attention to the age of their broodmares and either, breed earlier in their life or culling them earlier to avoid excessive costs. Older mares are also of less interest concerning genetics point of view [71] that should also be considered by breeders. Moreover, as frozen semen was associated with decreased pregnancy rates, the use of fresh or cooled semen for the insemination of old mares should be advised.

In this study, it has also been shown that nulliparity as the same time to aging affected pregnancy rates. For their first gestation, mares should not be more than 10 years old to increase chance of

579 pregnancy. One advice to breeders could be to use the mares' time before competition to produce
580 their first foal as it will help for the later inseminations.

581 Suckling at insemination prevented post-breeding inflammation. The most probable hypothesis could
582 be that sucking provokes oxytocin release that is acting as a natural treatment of uterine fluid
583 accumulation.

584 Here, pregnancy rates were not affected by post-breeding inflammation, although uterine fluid
585 accumulation were more frequent than in other studies. This suggests that endometritis is well
586 handled and that oxytocin only as a treatment is efficient.

587

Declaration of competing interest

None.

Acknowledgements

The authors tanks V. Lehuraux, A Jugland, B. Brisset and P. Drevillon from the Stud of “Charmoy” (89) and Stud of “Les Bréviaires” (78) for the availability of the data and for their hospitality during the visits of their organizations. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- [1] Allen WR, Brown L, Wright M, Wilsher S. Reproductive efficiency of Flatrace and National Hunt Thoroughbred mares and stallions in England. *Equine Veterinary Journal* 2007;39:438–45. <https://doi.org/10.2746/042516407X1737581>.
- [2] Langlois B, Blouin C. Statistical analysis of some factors affecting the number of horse births in France. *Reprod Nutr Dev* 2004;44:583–95. <https://doi.org/10.1051/rnd:2004055>.
- [3] Katila T, Reilas T, Nivola K, Peltonen T, Virtala A-M. A 15-year survey of reproductive efficiency of Standardbred and Finnhorse trotters in Finland - descriptive results. *Acta Veterinaria Scandinavica* 2010:11.
- [4] Nath L, Anderson G, McKinnon A. Reproductive efficiency of Thoroughbred and Standardbred horses in north-east Victoria. *Australian Veterinary Journal* 2010;88:169–75. <https://doi.org/10.1111/j.1751-0813.2010.00565.x>.
- [5] Hanlon D, Stevenson M, Evans M, Firth E. Reproductive performance of Thoroughbred mares in the Waikato region of New Zealand: 1. Descriptive analyses. *New Zealand Veterinary Journal* 2012;60:329–34. <https://doi.org/10.1080/00480169.2012.693039>.
- [6] Haadem CS, Nødtvedt A, Farstad W, Thomassen R. A retrospective cohort study on fertility in the Norwegian Coldblooded trotter after artificial insemination with cooled, shipped versus fresh extended semen. *Acta Vet Scand* 2015;57:77. <https://doi.org/10.1186/s13028-015-0161-8>.
- [7] Bosh KA, Powell D, Neibergs JS, Shelton B, Zent W. Impact of reproductive efficiency over time and mare financial value on economic returns among Thoroughbred mares in central Kentucky. *Equine Veterinary Journal* 2009;41:889–94. <https://doi.org/10.2746/042516409X456059>.
- [8] Lane E, Bijnen M, Osborne M, More S, Henderson I, Duffy P, et al. Key Factors Affecting Reproductive Success of Thoroughbred Mares and Stallions on a Commercial Stud Farm. *Reprod Dom Anim* 2016;51:181–7. <https://doi.org/10.1111/rda.12655>.
- [9] Platt H. Aetiological aspects of abortion in the thoroughbred mare. *J Comp Pathol* 1973;83:199–205.
- [10] Morris LHA, Allen WR. Reproductive efficiency of intensively managed Thoroughbred mares in Newmarket. *Equine Veterinary Journal* 2002;34:51–60. <https://doi.org/10.2746/042516402776181222>.
- [11] Gibbs PG, Davison KE. A field study on reproductive efficiency of mares maintained predominately on native pasture. *Journal of Equine Veterinary Science* 1992;12:219–22. [https://doi.org/10.1016/S0737-0806\(06\)81449-8](https://doi.org/10.1016/S0737-0806(06)81449-8).
- [12] Baker CB, Little TV, McDOWELL KJ. The live foaling rate per cycle in mares. *Equine Veterinary Journal* 1993;25:28–30. <https://doi.org/10.1111/j.2042-3306.1993.tb04819.x>.
- [13] Hemberg E, Lundeheim N, Einarsson S. Reproductive Performance of Thoroughbred Mares in Sweden. *Reprod Domest Anim* 2004;39:81–5. <https://doi.org/10.1111/j.1439-0531.2004.00482.x>.
- [14] Brück I, Anderson G, Hyland J. Reproductive performance of Thoroughbred mares on six commercial stud farms. *Australian Vet J* 1993;70:299–303. <https://doi.org/10.1111/j.1751-0813.1993.tb07979.x>.
- [15] Squires E, Barbacini S, Matthews P, Byers W, Schwenzer K, Steiner J, et al. Retrospective study of factors affecting fertility of fresh, cooled and frozen semen. *Equine Veterinary Education* 2006;18:96–9. <https://doi.org/10.1111/j.2042-3292.2006.tb00425.x>.
- [16] Samper JC, Vidament M, Katila T, Newcombe JR, Estrada A, Sargeant J. Analysis of some factors associated with pregnancy rates of frozen semen: a multi-center study. *Theriogenology* 2002;58:647–50. [https://doi.org/10.1016/S0093-691X\(02\)00754-9](https://doi.org/10.1016/S0093-691X(02)00754-9).
- [17] Rose BV, Firth M, Morris B, Roach JM, Wathes DC, Verheyen KLP, et al. Descriptive study of current therapeutic practices, clinical reproductive findings and incidence of pregnancy loss in

- intensively managed thoroughbred mares. *Animal Reproduction Science* 2018;188:74–84.
<https://doi.org/10.1016/j.anireprosci.2017.11.011>.
- [18] Chevalier-Clément F. Pregnancy loss in the mare. *Animal Reproduction Science* 1989;20:231–44.
[https://doi.org/10.1016/0378-4320\(89\)90088-2](https://doi.org/10.1016/0378-4320(89)90088-2).
- [19] Whitaker TC, Brace C. Mare age as a selection consideration for sport horse broodmares; a breeder survey. *Proc BrSoc Anim Sci* 2007;2007:30–30.
<https://doi.org/10.1017/S1752756200019335>.
- [20] Physick-Sheard PW. Demographic analysis of the Canadian Standardbred broodmare herd. *Preventive Veterinary Medicine* 1995;24:285–99. [https://doi.org/10.1016/0167-5877\(95\)00492-F](https://doi.org/10.1016/0167-5877(95)00492-F).
- [21] Heidler B, Aurich JE, Pohl W, Aurich Chr. Body weight of mares and foals, estrous cycles and plasma glucose concentration in lactating and non-lactating Lipizzaner mares. *Theriogenology* 2004;61:883–93. [https://doi.org/10.1016/S0093-691X\(03\)00279-6](https://doi.org/10.1016/S0093-691X(03)00279-6).
- [22] Merkt H, Günzel A-R. A survey of Early Pregnancy Losses in West German Thoroughbred Mares. *Equine Veterinary Journal* 1979;11:256–8. <https://doi.org/10.1111/j.2042-3306.1979.tb01359.x>.
- [23] Blanchard TL, Thompson JA, Love CC, Brinsko SP, Ramsey J, O’Meara A, et al. Influence of day of postpartum breeding on pregnancy rate, pregnancy loss rate, and foaling rate in Thoroughbred mares. *Theriogenology* 2012;77:1290–6. <https://doi.org/10.1016/j.theriogenology.2011.10.034>.
- [24] Camillo F, Marmorini P, Romagnoli S, Vannozzi I, Bagliacca M. Fertility at the first post partum estrous compared with fertility at the following estrous cycles in foaling mares and with fertility in nonfoaling mares. *Journal of Equine Veterinary Science* 1997;17:612–6.
[https://doi.org/10.1016/S0737-0806\(97\)80189-X](https://doi.org/10.1016/S0737-0806(97)80189-X).
- [25] Camargo CE, Kozicki LE, Ruda PC, Breno Pedrosa V, Talini R, Weiss RR, et al. Reproductive efficiency in lactating mares inseminated early in the puerperium (<10 days post partum) versus non-lactating mares inseminated 180 days post partum: PHK 2017;33:458–64.
<https://doi.org/10.21836/PEM20170506>.
- [26] Christoffersen M, Troedsson M. Inflammation and fertility in the mare. *Reproduction in Domestic Animals* 2017;52:14–20. <https://doi.org/10.1111/rda.13013>.
- [27] Woodward EM, Christoffersen M, Campos J, Squires EL, Troedsson MHT. Susceptibility to persistent breeding-induced endometritis in the mare: Relationship to endometrial biopsy score and age, and variations between seasons. *Theriogenology* 2012;78:495–501.
<https://doi.org/10.1016/j.theriogenology.2012.02.028>.
- [28] Samper JC. Ultrasonographic Appearance and the Pattern of Uterine Edema to Time Ovulation in Mares. *Proceedings of the 43rd Annual Convention of the American Association of Equine Practitioners*, vol. 43, 1997, p. 189–91.
- [29] Zent WW, Troedsson MHT, Xue J-L. Postbreeding Uterine Fluid Accumulation in a Normal Population of Thoroughbred Mares: A Field Study. *Proc Am Assoc Equine Pract*, vol. 44, 1998, p. 64–5.
- [30] Canisso IF, Segabinazzi LGTM, Fedorka CE. Persistent Breeding-Induced Endometritis in Mares—A Multifaceted Challenge: From Clinical Aspects to Immunopathogenesis and Pathobiology. *IJMS* 2020;21:1432. <https://doi.org/10.3390/ijms21041432>.
- [31] Barbacini S, Marchi V, Zavaglia G. Equine frozen semen: results obtained in Italy during the 1994–1997 period. *Equine Veterinary Education* 1999;11:109–12. <https://doi.org/10.1111/j.2042-3292.1999.tb00930.x>.
- [32] Ricketts SW, Alonso S. The effect of age and parity on the development of equine chronic endometrial disease. *Equine Veterinary Journal* 1991;23:189–92. <https://doi.org/10.1111/j.2042-3306.1991.tb02752.x>.
- [33] Brinsko SP, Ball BA, Miller PG, Thomas PGA, Ellington JE. In vitro development of day 2 embryos obtained from young, fertile mares and aged, subfertile mares. *Reproduction* 1994;102:371–8.
<https://doi.org/10.1530/jrf.0.1020371>.
- [34] Carnevale EM, Ginther OJ. Relationships of age to uterine function and reproductive efficiency in mares. *Theriogenology* 1992;37:1101–15. [https://doi.org/10.1016/0093-691X\(92\)90108-4](https://doi.org/10.1016/0093-691X(92)90108-4).

- [35] Adams GP, Kastelic JP, Bergfelt DR, Ginther OJ. Effect of uterine inflammation and ultrasonically-detected uterine pathology on fertility in the mare. *J Reprod Fertil Suppl* 1987;35:445–54.
- [36] Rasch K, Schoon HA, Sieme H, Klug E. Histomorphological endometrial status and influence of oxytocin on the uterine drainage and pregnancy rate in mares. *Equine Veterinary Journal* 1996;28:455–60. <https://doi.org/10.1111/j.2042-3306.1996.tb01617.x>.
- [37] Sharma OP. RELEASE OF OXYTOCIN ELICITED BY SUCKLING STIMULUS IN MARES. *Reproduction* 1974;37:421–3. <https://doi.org/10.1530/jrf.0.0370421>.
- [38] Ellendorff F, Schams D. Characteristics of milk ejection, associated intramammary pressure changes and oxytocin release in the mare. *Journal of Endocrinology* 1988;119:219–27. <https://doi.org/10.1677/joe.0.1190219>.
- [39] Samper JC, Stanford MS, French HM, Chapwanya A. Post-breeding inflammation in mares after insemination with large and low doses of fresh or frozen semen: PHK 2016;32:24–6. <https://doi.org/10.21836/PEM20160103>.
- [40] Cazales N, Estradé MJ, Pereyra F, Fiala-Rechsteiner SM, Mattos RC. Sperm transport and endometrial inflammatory response in mares after artificial insemination with cryopreserved spermatozoa. *Theriogenology* 2020;158:180–7. <https://doi.org/10.1016/j.theriogenology.2020.09.021>.
- [41] Ealy AD, Drost M, Hansen PJ. Developmental Changes in Embryonic Resistance to Adverse Effects of Maternal Heat Stress in Cows¹. *Journal of Dairy Science* 1993;76:2899–905. [https://doi.org/10.3168/jds.S0022-0302\(93\)77629-8](https://doi.org/10.3168/jds.S0022-0302(93)77629-8).
- [42] Roman-Ponce H, Thatcher WW, Caton D, Barron DH, Wilcox CJ. Thermal Stress Effects on Uterine Blood Flow in Dairy Cows. *Journal of Animal Science* 1978;46:175–80. <https://doi.org/10.2527/jas1978.461175x>.
- [43] Malschitzky E, Schilela A, Mattos ALG, Garbade P, Gregory RM, Mattos RC. Intrauterine fluid accumulation during foal heat increases embryonic death. *Pferdeheilkunde* 2003;19:1–4.
- [44] Pycck JF, Newcombe JR. Assessment of the effect of three treatments to remove intrauterine fluid on pregnancy rate in the mare. *Veterinary Record* 1996;138:320–3. <https://doi.org/10.1136/vr.138.14.320>.
- [45] Sieme H, Bonk A, Hamann H, Klug E, Katila T. Effects of different artificial insemination techniques and sperm doses on fertility of normal mares and mares with abnormal reproductive history. *Theriogenology* 2004;62:915–28. <https://doi.org/10.1016/j.theriogenology.2003.12.011>.
- [46] Sieme H, Schäfer T, Stout TAE, Klug E, Waberski D. The effects of different insemination regimes on fertility in mares. *Theriogenology* 2003;60:1153–64. [https://doi.org/10.1016/S0093-691X\(03\)00113-4](https://doi.org/10.1016/S0093-691X(03)00113-4).
- [47] Watson ED, Barbacini S, Berrocal B, Sheerin O, Marchi V, Zavaglia G, et al. Effect of insemination time of frozen semen on incidence of uterine fluid in mares. *Theriogenology* 2001;56:123–31. [https://doi.org/10.1016/S0093-691X\(01\)00548-9](https://doi.org/10.1016/S0093-691X(01)00548-9).
- [48] Cuervo-Arango J, Claes AN, Stout TA. A retrospective comparison of the efficiency of different assisted reproductive techniques in the horse, emphasizing the impact of maternal age. *Theriogenology* 2019;132:36–44. <https://doi.org/10.1016/j.theriogenology.2019.04.010>.
- [49] Ball BA, Little TV, Hillman RB, Woods GL. Pregnancy rates at Days 2 and 14 and estimated embryonic loss rates prior to day 14 in normal and subfertile mares. *Theriogenology* 1986;26:611–9. [https://doi.org/10.1016/0093-691X\(86\)90168-8](https://doi.org/10.1016/0093-691X(86)90168-8).
- [50] Woods GL, Baker CB, Baldwin JL, Ball BA, Bilinski J, Cooper WL, et al. Early pregnancy loss in brood mares. *Journal of Reproduction and Fertility Supplement* 1987;35:455–9.
- [51] Sharma S, Dhaliwal GS, Dadarwal D. Reproductive efficiency of Thoroughbred mares under Indian subtropical conditions: A retrospective survey over 7 years. *Animal Reproduction Science* 2010;117:241–8. <https://doi.org/10.1016/j.anireprosci.2009.05.011>.
- [52] Malheiros de Souza JR, Dias Gonçalves PB, Bertolin K, Ferreira R, Sardinha Ribeiro AS, Ribeiro DB, et al. Age-dependent effect of foal heat breeding on pregnancy and embryo mortality rates in Thoroughbred mares. *Journal of Equine Veterinary Science* 2020:102982. <https://doi.org/10.1016/j.jevs.2020.102982>.

- [53] Vidament M, Dupere AM, Julienne P, Evain A, Noue P, Palmer E. Equine frozen semen: freezability and fertility field results. *THERIOGENOLOGY* 1997;48:907–17. [https://doi.org/10.1016/S0093-691X\(97\)00319-1](https://doi.org/10.1016/S0093-691X(97)00319-1).
- [54] Davies Morel MCG, Newcombe JR, Holland SJ. Factors affecting gestation length in the Thoroughbred mare. *Animal Reproduction Science* 2002;74:175–85. [https://doi.org/10.1016/S0378-4320\(02\)00171-9](https://doi.org/10.1016/S0378-4320(02)00171-9).
- [55] Derisoud E, Auclair-Ronzaud J, Palmer E, Robles M, Chavatte-Palmer P. Female age and parity in horses: how and why does it matter? *Reproduction, Fertility and Development* In press.
- [56] Ginther OJ, Carnevale EM, Bergfelt DR. DELAY IN EMERGENCE OF THE OVULATORY FOLLICULAR WAVE IN OLD MARES. *JOURNAL OF EQUINE VETERINARY SCIENCE* 1993;13:5.
- [57] Ginther OJ, Gastal MO, Gastal EL, Jacob JC, Siddiqui MAR, Beg MA. Effects of age on follicle and hormone dynamics during the oestrous cycle in mares. *Reprod Fertil Dev* 2008;20:955. <https://doi.org/10.1071/RD08121>.
- [58] Ginther OJ, Gastal MO, Gastal EL, Jacob JC, Beg MA. Age-related dynamics of follicles and hormones during an induced ovulatory follicular wave in mares. *Theriogenology* 2009;71:780–8. <https://doi.org/10.1016/j.theriogenology.2008.09.051>.
- [59] Carnevale EM, Bergfelt DR, Ginther OJ. Aging effects on follicular activity and concentrations of FSH, LH, and progesterone in mares. *Animal Reproduction Science* 1993;31:287–99. [https://doi.org/10.1016/0378-4320\(93\)90013-H](https://doi.org/10.1016/0378-4320(93)90013-H).
- [60] Vanderwall DK, Woods GL, Freeman DA, Weber JA, Rock RW, Tester DF. Ovarian follicles, ovulations and progesterone concentrations in aged versus young mares. *Theriogenology* 1993;40:21–32. [https://doi.org/10.1016/0093-691X\(93\)90338-6](https://doi.org/10.1016/0093-691X(93)90338-6).
- [61] Carnevale EM, Bergfelt DR, Ginther OJ. Follicular activity and concentrations of FSH and LH associated with senescence in mares. *Animal Reproduction Science* 1994;35:231–46. [https://doi.org/10.1016/0378-4320\(94\)90039-6](https://doi.org/10.1016/0378-4320(94)90039-6).
- [62] Marinone AI, Losinno L, Fumuso E, Rodríguez EM, Redolatti C, Cantatore S, et al. The effect of mare's age on multiple ovulation rate, embryo recovery, post-transfer pregnancy rate, and interovulatory interval in a commercial embryo transfer program in Argentina. *Animal Reproduction Science* 2015;158:53–9. <https://doi.org/10.1016/j.anireprosci.2015.04.007>.
- [63] Claes A, Ball BA, Scoggin KE, Roser JF, Woodward EM, Davolli GM, et al. The influence of age, antral follicle count and diestrous ovulations on estrous cycle characteristics of mares. *Theriogenology* 2017;97:34–40. <https://doi.org/10.1016/j.theriogenology.2017.04.019>.
- [64] Claes A, Ball BA, Scoggin KE, Esteller-Vico A, Kalmar JJ, Conley AJ, et al. The interrelationship between anti-Müllerian hormone, ovarian follicular populations and age in mares: Anti-Müllerian hormone and ovarian follicular populations. *Equine Vet J* 2015;47:537–41. <https://doi.org/10.1111/evj.12328>.
- [65] Uliani RC, Conley AJ, Corbin CJ, Friso AM, Maciel LFS, Alvarenga MA. Anti-Müllerian hormone and ovarian aging in mares. *Journal of Endocrinology* 2019;147–56. <https://doi.org/10.1530/JOE-18-0391>.
- [66] Korkmaz Ö, Emre B, Polat İM, Zonturlu AK, PiR Yağci İ, Pekcan M, et al. The Correlation Between Anti-Müllerian Hormone Concentrations and Reproductive Parameters in Different Age Groups in Purebred Arabian Mares. *Kafkas Univ Vet Fak Derg* 2020;26:53–7. <https://doi.org/10.9775/kvfd.2019.22227>.
- [67] Davies Morel MCG, Newcombe JR, Hayward K. Factors affecting pre-ovulatory follicle diameter in the mare: the effect of mare age, season and presence of other ovulatory follicles (multiple ovulation). *Theriogenology* 2010;74:1241–7. <https://doi.org/10.1016/j.theriogenology.2010.05.027>.
- [68] Morel MCGD, Newcombe JR, Swindlehurst JC. The effect of age on multiple ovulation rates, multiple pregnancy rates and embryonic vesicle diameter in the mare. *Theriogenology* 2005;63:2482–93. <https://doi.org/10.1016/j.theriogenology.2004.09.058>.
- [69] Panzani D, Rota A, Marmorini P, Vannozzi I, Camillo F. Retrospective study of factors affecting multiple ovulations, embryo recovery, quality, and diameter in a commercial equine embryo

801 transfer program. Theriogenology 2014;82:807–14.
802 <https://doi.org/10.1016/j.theriogenology.2014.06.020>.
803 [70] Ball BA, Little TV, Weber JA, Woods GL. Survival of Day-4 embryos from young, normal mares and
804 aged, subfertile mares after transfer to normal recipient mares. Reproduction 1989;85:187–94.
805 <https://doi.org/10.1530/jrf.0.0850187>.
806 [71] Palmer E, Chavatte-Palmer P. Contribution of reproduction management and technologies to
807 genetic progress in horse breeding. Journal of Equine Veterinary Science 2020;103016.
808 <https://doi.org/10.1016/j.jevs.2020.103016>.
809

Figure legends

Figure 1: Odds ratio for factors influencing the likelihood of post-breeding endometritis in sport mares

Figure 2: Odds ratio for factors influencing the likelihood of pregnancy in sport mares

Figure 3: Factors affecting post-breeding endometritis and pregnancy rates in the studied population of sport mares

Red arrows signifies that the factor increased the likelihood while blue arrows indicated that the factor decreased the likelihood.

Supplementary data legends

Supplementary table 1

Sup1_PostBreedingInfla_OR.xlsx

Odds ratio for factors that did not influence the likelihood of post-breeding endometritis in sport mares

Supplementary table 2

Sup2_DG14_OR.xlsx

Odds ratio for factors that did not influence the likelihood of pregnancy in sport mares

Supplementary table 3

Sup3_Age_NS.xlsx

Factors that were not associated with mares' age

Supplementary table 4

Sup4_parity_NS.xlsx

Factors that were not associated with mares' parity

Supplementary table 5

Sup4_nursing_NS.xlsx

Factors that were not associated with mares' nursing

Figures

845

846

847

848

849

850

851

852