K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

JOURNAL OF HERPETOLOGY

Characterizing Potential Tetrodotoxin Resistance in Domain IV of the Voltage-Gated Sodium Channel Nav1.4 of Pacific Chorus Frogs, *Pseudacris regilla*

Katherine O. Montana^{1,2,3}, Valeria Ramírez Castañeda¹, Rebecca D. Tarvin^{1,3}

¹*Museum of Vertebrate Zoology and Department of Integrative Biology, University of California, Berkeley, Berkeley, California, USA 94720*

²*Present address: Department of Biology, San Francisco State University, San Francisco, California, USA 94132 and Institute for Biodiversity Science and Sustainability, California Academy of Sciences, San Francisco, California, USA 94118*

³*Corresponding authors. kmontana@berkeley.edu, rdtarvin@berkeley.edu*

LRH: K.O. Montana et al.

RRH: Tetrodotoxin resistance in *Pseudacris regilla*

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

Abstract.—Animals that frequently encounter toxins often evolve mechanisms of toxin resistance. Both predators that consume toxic prey and organisms in physical contact with a toxin or pollutant in their environment may experience natural selection for resistance. Based on observations that Pacific Chorus Frogs (*Pseudacris regilla*) sometimes eat and mistakenly amplex tetrodotoxin (TTX)-defended *Taricha* newts, we hypothesized that *P. regilla* may possess resistance to TTX. We tested this prediction by comparing the amino acid sequences of the domain IV pore loop of the muscle voltage-gated sodium channel gene SCN4A (Nav1.4 protein) in populations of *P. regilla* that are sympatric and allopatric with *Taricha*. We identified a single substitution in the domain IV pore loop of Nav1.4 in *P. regilla* that has been hypothesized to provide TTX resistance based on computational modeling, though its role in resistance has been challenged. Nonetheless, both allopatric and sympatric *P. regilla* had this substitution, suggesting that it may be unrelated to TTX exposure from *Taricha*. Thus, there is no conclusive evidence that *P. regilla* has evolved TTX resistance encoded by amino acid substitutions in this domain. In addition, California occurrence data from the last 50 years indicate that *Taricha* activity peaks in January while the activity of *P. regilla* peaks in April. These relatively distinct activity patterns suggest that *P. regilla* may not be exposed to levels of TTX from *Taricha* that are high enough to require mutations in SCN4A. Nevertheless, other unidentified mechanisms of TTX resistance could be present in *P. regilla* and other species that are sympatric with toxic newts.

Key words: Toxin resistance; California; Environmental toxins; Target-site insensitivity; Salamanders; *Taricha*

37

38 Resumen.— Los animales que tienen contacto frecuente con toxinas suelen
39 desarrollar mecanismos de resistencia a las mismas. Tanto los depredadores que
40 consumen presas tóxicas como los organismos en contacto cercano con una toxina o
41 contaminante en su entorno pueden experimentar una presión de selección que los
42 lleva a evolucionar resistencia a toxinas. Basándonos en las observaciones de que las
43 ranas coro del Pacífico (*Pseudacris regilla*) a veces comen por error y/o amplexan las
44 salamandras del género *Taricha* que poseen tetrodotoxina (TTX), planteamos la
45 hipótesis de que *P. regilla* podría poseer resistencia a la TTX. Probamos esta
46 predicción comparando las secuencias de aminoácidos en el loop del poro del dominio
47 IV en el gen del canal de sodio voltaje dependiente muscular SCN4A (proteína Nav1.4)
48 en poblaciones de *P. regilla* que son simpátricas y alopátricas con *Taricha*.
49 Identificamos una única sustitución en el loop del poro del dominio IV de Nav1.4 en *P.*
50 *regilla* que se ha hipotetizado utilizando un modelo computacional que está relacionado
51 con la resistencia a TTX. Sin embargo, tanto el *P. regilla* alopátrico como el simpátrico
52 tienen esta sustitución, lo que sugiere que esta puede no estar relacionada con la
53 resistencia a la TTX proveniente de *Taricha*. Por lo tanto, no hay pruebas concluyentes
54 de que *P. regilla* haya evolucionado resistencia a la TTX causada sustituciones de
55 aminoácidos en este dominio de la proteína. Adicionalmente, los datos de ocurrencia en
56 California en los últimos 50 años indican una alternancia en la actividad de ambos
57 grupos, *Taricha* alcanza su punto máximo en enero, mientras que la de *P. regilla* lo
58 hace en abril. Estos patrones de actividad relativamente distintos sugieren que *P.*

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

regilla no está expuesta constantemente a niveles de TTX lo suficientemente altos como para requerir mutaciones en SCN4A. Sin embargo, otros mecanismos de resistencia a la TTX no identificados hasta ahora, podrían estar presentes en *P. regilla* y otras especies simpáticas a las salamandras tetrodotóxicas.

Palabras clave: Resistencia a las toxinas; California; Toxinas ambientales; Insensibilidad en el sitio de union; salamandras; Taricha

Many animals can synthesize or bioaccumulate toxic chemicals as an anti-predator defense. One common toxin is tetrodotoxin (TTX), which is employed as a defense by a variety of organisms including pufferfish and amphibians (e.g., dendrobatid frogs, *Atelopus*, *Polypedates*, *Brachycephalus*, salamandrid newts) (Daly et al., 1994; Tanu et al., 2001; Pires et al., 2002; Hanifin, 2010), as well as an offensive prey-capture technique by blue-ringed octopuses and flatworms (Ritson-Williams et al., 2006; Hwang et al., 2007; Williams, 2010). Bacterial symbionts of toxic newts and marine organisms produce TTX for their hosts, suggesting a bacterial origin in most if not all of these organisms (Magarlamov et al., 2017; Vaelli et al., 2020). TTX is a neurotoxin that blocks voltage-gated sodium channels by binding to highly conserved regions of the channel via hydrophobic forces and hydrogen bonds, fitting in the opening of the channel and blocking proper transport of Na⁺ ions (Kao and Levinson, 1986; Tikhonov and Zhorov, 2011). TTX binding prevents action potentials from propagating, resulting in cessation of nerve signals (Kao and Levinson, 1986; Tikhonov and Zhorov, 2011). In vertebrate animals lacking TTX resistance, ingested TTX can temporarily paralyze or, in some

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

cases, kill the organism. The poison usually becomes fatal when respiratory muscles fail, resulting in suffocation (Brodie, 1968).

While TTX can incapacitate or kill an animal that attempts to consume it, several species are able to come into contact with TTX and survive. Resistance to TTX intoxication usually evolves via one of three common mechanisms. In the first, amino acid substitutions in voltage-gated sodium channels (VGSCs), especially in domain IV (DIV) of Nav1.4, alter the shape of the pore loop (p-loop) and the surrounding area, which prevents TTX from binding (Geffeney et al., 2005; Tikhonov and Zhorov, 2005). Other regions of Nav1.4, such as domain III, and other voltage-gated sodium channels, such as Nav1.7, also harbor amino acid substitutions implicated in TTX resistance (Feldman et al., 2012; McGlothlin et al., 2016). Changes in the p-loops are particularly significant because they not only alter the sensitivity of the sodium channel to TTX but also to sodium ions; hence, there is often a tradeoff between TTX resistance and protein function (Chiamvimonvat et al., 1996; Geffeney et al., 2005; Lee et al., 2011; Tikhonov and Zhorov, 2011; Hague et al., 2018; but see Moniz et al., 2021). Thus, amino acid changes are usually confined to specific conserved regions that can provide resistance while allowing the organism to maintain sodium channel functionality (Feldman et al., 2012). The presence of amino acid substitutions that alter toxin binding is a mechanism known as target-site insensitivity and is the focus of this study.

Other mechanisms of TTX resistance include toxin-binding proteins, which have evolved in gastropods, pufferfish, horseshoe crabs, and shore crabs (Ho et al., 1994; Nagashima et al., 2002; Hwang et al., 2007; Hashiguchi et al. 2015), and diffusion

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

barriers, which are present in some invertebrates (Mebs et al., 2016). Some amphibians possess the toxin-binding, blood-plasma protein saxiphilin, but saxiphilin is incapable of binding TTX, at least in the animals that have been tested (Mahar et al., 1991; Llewellyn et al., 1997; Abderemane-Ali et al., 2021). Some species of mantids (*Tenodera sinensis*, *Sphodromantis viridis*, *Hierodula membranacea*, and *Miomantis caffra*) can survive tetrodotoxin ingestion but not toxin injection, suggesting the presence of a gut diffusion barrier (Mebs et al., 2016). Crayfish appear to have a similar mechanism (Wilson et al., 2014). Because we do not directly test for toxin-binding proteins or diffusion barriers, we cannot say whether *P. regilla* possesses these mechanisms of resistance.

Taricha newts (*Taricha torosa*, *T. granulosa*, *T. sierrae*, and *T. rivularis*, herein collectively referred to as *Taricha*) have TTX present on their skin (Brodie et al., 1974; Reimche et al., 2020) that likely functions as an anti-predator defense (Brodie et al. 2002). *Taricha* (and other salamandrids) possess amino acid changes in Na_v1.4 that help them resist their own TTX defenses (Hanifin and Gilly, 2015; Gendreau et al., 2021). The San Francisco Bay Area in California is one of two known hotspots of the arms race between newts and garter snake predators (Brodie et al., 2002). Here we explore the idea that high levels of TTX defenses in newts may select for resistance in other sympatric organisms outside the predator-prey dynamic, namely the Pacific Chorus Frog, *Pseudacris regilla*.

TTX from *Taricha* skin has been shown to enter waterways and affect the behavior of organisms present in the same aquatic systems. Larval *T. torosa* use TTX

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

as a cue to detect and avoid adult *T. torosa* that sometimes cannibalize larvae (Zimmer et al., 2006). TTX in water can also slow predatory patterns of dragonfly nymphs, which are macroinvertebrate prey of *T. torosa* (Bucciarelli and Kats, 2015). The New Zealand Mud Snail (*Potamopyrgus antipodarum*) moved out of its natural stream environment when encountering biologically relevant levels of TTX or chemical cues from *T. torosa* containing TTX (Ota et al., 2018). Water containing TTX also negatively increases newt trematode parasite mortality (Calhoun et al., 2017). Thus, naturally occurring levels of TTX produced by *T. torosa* can affect other organisms in the same aquatic environment.

Given that *P. regilla* have permeable skin, we hypothesized that they would be exposed to TTX through physical contact with newts or by occupying pond water that contains newts. We predicted that TTX exposure has selected for TTX resistance in *P. regilla* populations that are microsympatric with *Taricha*. We recognize that our hypothesis makes a simplifying assumption that the designation of *P. regilla* populations as sympatric or allopatric with *Taricha* is relevant and valid during the time in which selection may have occurred for TTX resistance, and that gene flow does not occur between sympatric and allopatric *P. regilla* populations. Although both of these assumptions may be violated by changes in climate and species distributions over time, evidence from the evolutionary arms race between garter snakes and TTX-bearing newts suggest that TTX potency and the degree of TTX resistance can change over a short period of time (Brodie et al., 2002), which aligns with our study that examines approximately the past 50 years. Occasional exposure to TTX may also be sufficient to select for target-site insensitivity (Durso et al., 2021).

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

We aimed to test whether the sympatry of *Taricha* newts with *Pseudacris regilla* has selected for target-site insensitivity in the skeletal muscle voltage-gated sodium channel of *P. regilla*. While *P. regilla* do not regularly prey upon *Taricha* (or vice versa), they likely come into contact with TTX in pond water shared with *Taricha* or via direct contact with *Taricha*, sometimes in mistaken amplexus behavior or even attempted predation (Fig. 1).

Our study centered on California populations, examining *Taricha* and *P. regilla* that do and do not overlap geographically in the present. We conducted demographic surveys of *Taricha granulosa*, *T. torosa*, and *P. regilla* at three San Francisco Bay Area ponds where the species co-occur to estimate the extent to which *P. regilla* may encounter TTX. Because these surveys were suspended after only a few months due to the COVID-19 pandemic, we then reviewed occurrence data available through the Global Biodiversity Information Facility (GBIF) to better estimate overlap of annual activity patterns. To determine whether *P. regilla* has structural modifications in the protein muscle voltage-gated sodium channel Nav1.4 that may provide TTX resistance, we sequenced and reviewed amino acid residues encoded by the SCN4A gene. Amino acid substitutions in this domain are known to confer resistance to TTX and have been found in several organisms that either have TTX defense or consume TTX-defended prey (Tikhonov and Zhorov, 2005; Venkatesh et al., 2005; Feldman et al., 2012; Hanifin and Gilly, 2015). We hypothesized that if the presence of *Taricha* selects for TTX resistance in *P. regilla*, populations sympatric with *Taricha* would have mutations in the p-loop of Nav1.4 while populations that are allopatric would not. Our work provides

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

insight into the potential cascading ecological consequences of chemical defenses, as we evaluate the unintended effects of animals wielding chemical defenses.

MATERIALS AND METHODS

Field Work.—We studied *Taricha* and *P. regilla* at four San Francisco Bay Area ponds. We conducted surveys at three of these sites where *Taricha granulosa*, *T. torosa*, and *P. regilla* co-occur: two in Briones Regional Park in Martinez, Contra Costa County, CA (Old Briones Road Trail 1 [OBRT 1] at 37.94571, -122.13395, and OBRT 2 at 37.94352, -122.14102, named after the trail closest to the sites) and the Japanese Pool (37.87440, -122.23760) at the University of California Botanical Garden (UCBG) in Berkeley, Alameda County, CA (Fig. 2). OBRT 1 and OBRT 2 are in an unshaded area of grazed open grassland that lies within an oak woodland habitat, and they connect to each other when water levels are high. The UCBG Japanese Pool is a man-made pool with a concrete floor and sides, located under tree cover within the visitor area of the UCBG (Fig. 2). We selected these three ponds because they were known to contain populations of *P. regilla* as well as *Taricha torosa* and *Taricha granulosa* (Katherine Montana, Rebecca Tarvin, pers. obs.). We also collected specimens from a fourth pond located in the Presidio of San Francisco (37.788496, -122.468605) that has historically contained *P. regilla* but no *Taricha* (records show absence of *Taricha* since at least 1966; pers. comm. J. Young, Presidio Trust). Mountain Lake is a natural lake surrounded by grass and receives run-off from the nearby Presidio Golf Course (Fig. 2). From October 2019 to February 2020, we conducted six surveys at the two Briones

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

ponds and five surveys at UCBG, with surveys at each site occurring three to four weeks apart from one another. We were unable to gather abundance data for more than this time period because the parks were closed from February until October 2020 during the early months of the COVID-19 pandemic.

Two researchers (usually K. Montana and V. Ramírez Castañeda) conducted surveys in the morning or early afternoon (between 8:00 AM and 1:30 PM). During each survey, surveyors split the pond perimeter into 7.62-m (25-feet) sections, and circled the pond on foot, counting how many individuals of each species were visible in the water from the shore within each transect. We note that surveyors were only able to count the number of visible newts; OBRT 1 and OBRT 2 were murky with plants and algae in which newts tended to hide and although the surface of the Japanese Pool at UCBG was glassy, the pool was filled with plants that could have obscured observations. Surveyors also counted the numbers of visible, individual adult *Pseudacris regilla*; *P. regilla* that were calling but not visible were not counted because surveyors could not ascertain if multiple calls were coming from a single or a few individuals. Though the TTX defenses of *T. granulosa* are thought to be more potent than that of *T. torosa* (Brodie et al., 1974), for the purposes of this study, we did not differentiate between the two species when we recorded their occurrences because they both contribute to the presence of TTX in pond water, and we could not easily identify individuals to species from the pond shores. We counted large (~5 cm or longer) *Taricha* larvae, adult *Taricha*, and *Taricha torosa* egg masses, which are conspicuous groups of eggs known to possess TTX via vertical transfer from the parent (Gall et al., 2012). The much smaller

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

T. granulosa eggs are typically hidden in the substrate and are more difficult to see; hence we did not include them in our survey. To simplify our survey protocol, we counted larval *Taricha*, which also have some amount of TTX (Gall et al., 2011), together with adults. We plotted survey results using R version 4.1.1 and the packages ggplot2 v.3.3.5 and ggpubr v.0.4.0 (Wickham, 2016; Kassambara, 2020; R Core Team, 2021) (Fig. 3). We compared survey results to GBIF occurrence data of *P. regilla* and *Taricha* from 1 January 1971 to 23 November 2021 by plotting the GBIF data using the following R packages: RColorBrewer v.1.1.2, ggplot2 v.3.3.5, chron v.2.3.56, cowplot v.1.1.1, and dplyr v.1.0.7 (Neuwirth, 2014; Wickham, 2016; James and Hornik, 2020; Wilke, 2020; Wickham et al., 2021).

After COVID-19 conditions permitted field work to continue in October 2020, we collected one *Pseudacris regilla* from the dry bed of OBRT 2 and eight *P. regilla* from the east arm of Mountain Lake in the Presidio of San Francisco. We surveyed OBRT 1 and UCBG at the same time but did not find any *P. regilla* at either location; OBRT 1 was dry.

Precipitation Data.—To better interpret amphibian abundance data from field surveys, we searched for precipitation data from NOAA for the weather stations closest to our field sites in Berkeley (UCBG) and Martinez (OBRT 1 and OBRT 2): Berkeley, CA USC00040693 and Martinez 2.2 SW, CA US1CACC0011 (NOAA, <https://www.ncdc.noaa.gov/cdo-web/search>). We measured pond size during each

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

survey event by adding the number of 7.62-m transects we walked during each survey to get the circumference and calculate the area, assuming the ponds were circular.

Notes on Taxonomy.—The taxonomy of *Pseudacris regilla* has changed several times over the last decades (Recuero et al., 2006; Fouquette and Dubois, 2014; Duellman et al., 2016). iNaturalist categorizes some *Pseudacris regilla* (e.g., in the San Francisco Bay Area) as *Pseudacris sierra*. However, these distinctions are not universally accepted by the herpetology community (Barrow et al., 2014; Crother et al., 2017). Here we consider *P. sierra*, *P. hypochondriaca*, and *Hyla regilla* synonyms of *Pseudacris regilla*. We downloaded occurrence data from GBIF for the *P. regilla* synonyms listed here and for all currently recognized species of *Taricha* (*T. granulosa*, *T. rivularis*, *T. sierrae*, *T. torosa*), as well as those no longer recognized (*T. lindoei*, *T. miocenica*, and *Palaeotaricha oligocenica*) and individuals identified only to the genus *Taricha*.

DNA Sequencing.—We selected 29 liver tissues of *Pseudacris regilla* from the Museum of Vertebrate Zoology at the University of California, Berkeley to broaden our geographic sampling (Table S1). We classified *P. regilla* samples as sympatric with *Taricha* if they were collected within 35 km or 500 m of elevation from the nearest *Taricha* occurrence documented on GBIF as of 23 November 2021; this included 11 individuals and the single *P. regilla* collected at OBRT 2. We classified allopatric *P. regilla* samples as those individuals collected at least 35 km or 500 m in elevation away

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

from the nearest *Taricha* observation; this included 18 individuals and the eight *P. regilla* from the Mountain Lake field site in the Presidio (Table S1). We predicted that Nav1.4 sequences from sympatric, but not allopatric, *P. regilla* would show amino acid substitutions potentially conferring TTX resistance. We utilized the R packages ggplot2 v.3.3.5 and tidyverse v.1.3.1 to map GBIF occurrences of *Taricha* throughout California, *P. regilla* museum samples that were sympatric with *Taricha*, *P. regilla* museum samples that were allopatric with *Taricha*, and our field sites (Wickham, 2016; Wickham et al., 2019; GBIF).

Our preliminary results suggested that a single amino acid substitution is present in *P. regilla*. Thus, to estimate when this substitution evolved, we also sequenced Nav1.4 from two close relatives of *P. regilla*, *Pseudacris cadaverina* and *Acris crepitans* (Table S1). The individuals of *P. cadaverina* and *A. crepitans* are from locations far from *Taricha* and, thus, were likely not exposed to TTX from *Taricha*. However, the range of *A. crepitans* overlaps with another TTX-bearing newt, *Notophthalmus viridescens* (Baird, 1854; Meham, 1967), so it may be exposed to TTX as well. Thus, if TTX from newts selects for mutations in Nav1.4 in sympatric frog species, we would expect to find amino acid substitutions in Nav1.4 of *A. crepitans* as well as in Nav1.4 of *P. regilla* sympatric with *Taricha*. In contrast, we expected to find no resistance-conferring amino acid substitutions in allopatric *P. regilla* or in *P. cadaverina*.

In the Evolutionary Genetics Lab (UC Berkeley), we extracted DNA from tissues according to the Miller et al. (1988) salt-extraction protocols, with modifications (see Supplementary Online Material for details). The primers used for *P. regilla* samples

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

targeted exon 24 (DIV segment 6 and p-loop) of the gene SCN4A: 2268F_SCN4A 5'-
TCTCCCGGCCCTCTTCAATA-3' and 2681R_SCN4A 5'-
TCGTCCTCGCATAAAGGCTC-3' (Tarvin et al., 2016). We added 0.5 µL of MgCl₂ to
each PCR sample. PCR protocols were: an initial denaturation for 2 min at 94 °C; 35
cycles of: 94 °C at 30 s for denaturation, an annealing temperature of 54 °C for 30 s,
and an elongation temperature of 72 °C for 45 s; and a final extension time of 7 min at
72 °C. PCR products were run on an agarose gel, purified with ExoSAP-IT PCR Product
Cleanup Reagent (ThermoFisher Scientific, 78201.1.ML) and sequenced using the
reverse primer 2681R_SCN4A at the UC Berkeley DNA Sequencing Facility. For PCR
and sequencing samples of *Pseudacris cadaverina* and *Acris crepitans*, we used the
same protocol and forward primer, but the reverse primer was redesigned based on the
DIV SCN4A *P. regilla* sequences we produced: P.regilla_SCN4AD4_R 5'-
ACTGCTTTCTTCTGTGGCCAC-3'.

We also obtained previously published SCN4A sequences from species that are
resistant to TTX, including *Erythrolamprus epinephelus*, which is potentially resistant to
TTX from *Atelopus*, as well as *Rhabdophis tigrinus*, *Thamnophis elegans*, *Thamnophis*
sirtalis, *Taricha torosa*, and *Taricha granulosa*, which are either confirmed or
hypothesized to possess target-site insensitivity to TTX via mutations in Nav1.4
(Feldman et al., 2009; Feldman et al., 2012; Hanifin and Gilly, 2015). We also include
sequences from *Nanorana parkeri* and *Rattus norvegicus*, which have not been
documented as being resistant to TTX (Table S2). We identified the location of the DIV
p-loop in Nav1.4 based on protein domains inferred by Tikhonov and Zhorov (2005). We

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

aligned sequences using Clustal Omega in Geneious (Sievers et al., 2011, Geneious Prime 2020.1.1). We reviewed the alignment by eye for amino acid differences between sequences, focusing on residues that have been shown to provide TTX resistance. We estimated a phylogeny of the species that we compared to *P. regilla* using 11 COI sequences (Table S2) and maximum likelihood on the IQTree web server, with two partitions (one for codon positions 1 and 2, another for codon position 3), the GTR + G substitution model, and support assessed using 1000 ultrafast bootstraps and the SH-aLRT test (Nguyen et al., 2015; Chernomor et al., 2015; Trifinopoulos et al., 2016; Hoang et al., 2018; IQ-Tree web server <http://iqtree.cibiv.univie.ac.at/>). We transformed the resulting phylogeny into an ultrametric chronogram using the function *chronos* in phytools v.0.7.90 and ape v.5.5 (Revell, 2012; Paradis and Schliep, 2019) (Fig. 6). We made all figures in R and Adobe Illustrator (Adobe, Inc., 2019; R Core Team, 2021).

RESULTS

Demographic surveys indicated that *P. regilla* may encounter TTX from *Taricha* in their environment. DNA sequencing identified one amino acid substitution in domain IV of Nav1.4 in *P. regilla* that could potentially be connected to TTX resistance, but the degree to which the mutation affects resistance is inconclusive.

Field Surveys.—While we never observed *P. regilla* at OBRT 2 or UCBG, their abundance at OBRT 1 peaked early in the observation period at 32 individuals on the first field trip date (20 October 2019). For OBRT field sites, *Taricha* abundance peaked

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

at the beginning of February; for the UCBG, abundance was low, and we may not have observed the seasonal peak (surveys were suspended at the end of February 2020 because of the COVID-19 pandemic). *Taricha torosa* egg masses were most abundant at the beginning of February 2020 at OBRT 1 and OBRT 2, while they were most abundant at UCBG at the end of February (though we cannot be certain these are true peaks because our surveys ended before March) (Fig. 3). California-wide GBIF observations for *P. regilla* and *Taricha* from 1 January 1971 to 23 November 2021 indicate two peaks of *P. regilla* abundance in April and August, while *Taricha* peaks in January (Fig. 4). Though the abundance peaks do not occur at precisely the same time, *P. regilla* and *Taricha* do overlap with one another and, thus, the species have opportunities to interact such that *Taricha* could expose *P. regilla* to TTX. These data are summarized across California and thus represent average activity across a range of habitat types. Nevertheless, when the data are constrained to a square roughly representing the San Francisco Bay Area (N 38.66226, S 37.09298, E -121.12701, W -123.23579), the results are similar, indicating a peak for *Taricha* in January and a peak for *P. regilla* in March (Supplementary Fig. 1).

When we surveyed the field sites in Briones Regional Park and the UC Botanical Garden from October 2019 to March 2020, California was experiencing drought. During this period, rain began in mid-November 2019 and stopped at the end of February 2020. The most rain recorded in one day during the six-month period in Berkeley, where UCBG is located, was 30.23 mm, the average recorded rainfall during that period was 1.76 mm/day, and the total number of recorded days with rain was 40 days (NOAA field

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

station Berkeley, CA USC00040693, elevation 94.49 m, 37.8744° N, 122.2605° W) (NOAA, 2022). The most rain recorded in one day during the six-month period in Martinez, where Briones is located, was 89.15 mm, the average recorded rainfall during that period was 1.61 mm/day, and the total number of recorded days with rain was 33 days (NOAA field station Martinez 2.2 SW, CA US1CACC0011, elevation 210.92 m, 37.9744° N, 122.1384° W) (NOAA, 2022). The area of the ponds fluctuated due to changes in precipitation and, in the case of the Japanese Pool, some drainage for maintenance (UCBG personnel, pers. com.). The area of OBRT 1 ranged between 319 and 912 m² with a survey perimeter of about 84-122 m. The area of OBRT 2 ranged between 1550 and 1733 m² with a perimeter of about 130-145 m. During our study period, OBRT 1 and OBRT 2 did not merge. The area of the Japanese Pool ranged between 114 and 137 m² with a perimeter of about 46-53 m.

Amino Acid Changes.— We found one amino acid change (I1519M) within the DIV p-loop of the sodium channel protein Nav1.4 in *Pseudacris regilla* compared to a sequence of *Rattus norvegicus* (a common species used in comparative TTX studies [e.g., Tikhonov and Zhorov, 2005; Tarvin et al. 2016] as it represents the mammalian TTX-sensitive Nav1.4 [Chahine et al., 1994]) (Fig. 6) (Tables S1 and S2). Importantly, there were no observed differences in amino acid sequences between *P. regilla* specimens that were sympatric and allopatric with *Taricha*. The I1519M substitution has been previously predicted by computational modeling to provide some amount of TTX resistance (Tikhonov and Zhorov, 2005; Tikhonov and Zhorov, 2011). The I1519M

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

substitution is also present in the snake *Rhabdophis tigrinus*, which consumes TTX-defended treefrogs, and is inferred to be associated with TTX resistance (Feldman et al., 2012). However, it has been suggested that the characterization of the I1519M substitution as contributing to TTX resistance is erroneous because the same substitution is found in TTX-sensitive voltage-gated sodium channels of other snakes (McGlothlin et al., 2016). Additionally, we found that the amino acid change I1519M occurs in the frog *Nanorana parkeri*, which is not known for being resistant to or in contact with tetrodotoxin; though, it does occur within the range of TTX-defended salamandrids such as *Tylostrotion verrucosus* (Che et al., 2010; Vitt and Caldwell, 2013; Wanninger et al., 2018). Close relatives of *P. regilla* (*P. cadaverina* or *A. crepitans*) did not possess the I1519M substitution, suggesting that the mutation arose within *P. regilla* or its ancestor.

DISCUSSION

Our results provide conflicting evidence regarding whether Nav1.4-based TTX resistance is present in *Pseudacris regilla*. In support of resistance is the presence of a single amino acid substitution in *P. regilla* Nav1.4 (Fig. 6) hypothesized to provide TTX resistance based on computational models (Tikhonov and Zhorov, 2005; Tikhonov and Zhorov, 2011), and evidence that *P. regilla* and *Taricha* overlap somewhat in annual activity patterns (Fig. 4). However, amino acid changes in important regions of the Nav1.4 protein do not always translate to a resistant phenotype (e.g., see Abderemane-Ali et al., 2021). Furthermore, both allopatric and sympatric *P. regilla* possessed the

substitution, but *A. crepitans* did not, although we expected that it might possess some resistance due to its overlap in range with *N. viridescens*. The presence of an identical substitution in *Nanorana parkeri* was unexpected and could either present evidence in support of resistance (the species does live near the TTX-defended salamandrid *Tylototriton verrucosus*), or against it, potentially indicative of functionally permissive amino acid diversity at that site in anurans. Below we review in more detail these lines of evidence.

Abundance data indicate that *P. regilla* and *Taricha* annual activity patterns overlap but that peak breeding times are not synchronized (Figs. 3, 4; Supplementary Fig. 1). Observations of mistaken amplexus and predation of newts by *P. regilla* indicate that the species do come into contact (Fig. 1). Still, TTX exposure may not be frequent or potent enough for *P. regilla* to undergo strong selection for resistance to TTX. In cases where environmental selective regimes are not predictable, more plastic responses like upregulation of detoxifying enzymes may be the primary way animals deal with toxin exposure (Zanger and Schwab, 2013). Nevertheless, even occasional exposure to TTX can result in maintenance of target-site insensitivity (Durso et al., 2021).

We hypothesized that we would find TTX-conferring amino acid substitutions in Na_v1.4 in *P. regilla* that are microsympatric with *Taricha* and no substitutions in *P. regilla* that are allopatric with *Taricha*, with a greater number of substitutions indicating a higher level of resistance as has been shown in salamanders and snakes (Geffeney et al., 2005; Lee et al., 2011; Hanifin and Gilly, 2015). Compared to *Rattus norvegicus* DIV,

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

we found one amino acid change at a site hypothesized to provide TTX resistance, I1519M, which is a TTX-sensing residue located in the p-loop (Tikhonov and Zhorov, 2005; Tikhonov and Zhorov, 2011; Feldman et al. 2012). This substitution was present in *P. regilla* both sympatric and allopatric with *Taricha* and in *Nanorana parkeri*. It is also found in *Rhabdophis tigrinus*, a species of snake that has been hypothesized to be resistant to TTX (Feldman et al., 2012), although no empirical tests have been undertaken and other authors have disputed the role of I1519M in TTX resistance (McGlothlin et al., 2016; noted as I1555M in these studies). The presence of this substitution in all *P. regilla* samples does not support our hypothesis that substitutions would only be found where *P. regilla* is exposed to TTX from sympatric *Taricha* (at least based on occurrence data from the last 50 years). Ultimately, our study suggests that the I1519M substitution evolved in the ancestor of *P. regilla*, which likely rules out contemporary selection from *Taricha*. The presence of the substitution in *N. parkeri* could also support that the mutation is unrelated to newt exposure. However, *N. parkeri* overlaps with other toxin-defended salamandrids like *T. verrucosus*, making it difficult to rule out resistance in *N. parkeri* or, consequently, in *P. regilla*. Similarly, *A. crepitans* co-occurs with the toxic newt *Notophthalmus viridescens*, but this species does not possess the I1519M substitution. Because there is disagreement whether the I1519M mutation provides resistance, we cannot say with certainty whether it provides TTX resistance in *P. regilla* without further experimental evidence.

While there is not strong evidence that *P. regilla* possesses target-site insensitivity in the p-loop of DIV in Na_v1.4, it is plausible that *P. regilla* may have other

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

ways of avoiding the effects of TTX, including amino acid mutations in other domains of Na_v1.4 or in other voltage-gated sodium channels (Feldman et al., 2012; McGlothlin et al., 2016). *Pseudacris regilla* could also possess the ability to smell TTX, as larval *Taricha* can, in order to avoid it (Zimmer et al., 2006). Alternatively, they could possess a diffusion barrier in the skin or the gut that would prevent high levels of TTX exposure, as occurs in mantids (Mebs et al., 2016). It is also possible that *P. regilla* possesses a binding protein that can scavenge TTX; other frogs have saxiphillin, a protein that binds saxitoxin, which is structurally similar to TTX (Mahar et al., 1991; Llewellyn et al., 1997). Similarly, little is known about the potential interactions between *Acris crepitans* and *Notophthalmus viridescens*, which could also select for TTX resistance in the frogs. To date, none of these alternative resistance mechanisms have been studied in *Pseudacris*, *Acris*, or *Notophthalmus*.

A number of reasons could explain why *P. regilla* would lack target-site insensitivity in Na_v1.4. Evolving TTX resistance could be costly to *P. regilla* due to possible alterations in the selectivity filter of sodium channels that frequently occurs when amino acid substitutions are present (Chiamvimonvat et al., 1996; Geffeney et al., 2005; Lee et al. 2011; Tikhonov and Zhorov, 2011; Moniz et al., 2021). Evolving costly mutations may require a stronger selective pressure than is currently occurring. Given the relative quantities of TTX on newt skin [up to 4.69 mg/newt (Hanifin et al., 2008)] compared to TTX levels in newt-containing water [$\sim 1 \times 10^{-7}$ mol/L (Zimmer et al., 2006)], the environmental toxicity of TTX is likely to cause a weaker selection for TTX resistance than exposure via chemical defenses from prey. The cost of exhibiting TTX

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

resistance combined with the low potential for *P. regilla* to interact with high concentrations of TTX in its environment may explain why we did not unequivocally identify target-site insensitivity in *P. regilla* DIV Nav1.4. Future work exploring the effects of TTX as an environmental toxin could include measuring TTX concentrations in newt-containing ponds, performing whole organism or *in vitro* assays of *P. regilla* to determine TTX sensitivity, and testing for other modes of TTX resistance in animals that live in sympatry with salamandrid newts.

Conclusions.—We do not find strong evidence for target-site insensitivity in DIV of the skeletal muscle voltage-gated sodium channel Nav1.4 of *Pseudacris regilla*. Although it is possible that the I1519M substitution in the p-loop of Nav1.4 may be related to TTX resistance, *P. regilla* may not encounter TTX from *Taricha* with high enough frequency or magnitude to select for TTX resistance. Thus, these animals may have behavioral or alternative physiological mechanisms to survive TTX exposure, or TTX exposure may not be severe enough to require adaptation by *P. regilla*. Investigating the environmental toxicity of TTX in the relationship between *P. regilla* and *Taricha* is an opportunity to explore the nexus of ecological interactions, neurotoxins, and resistance mechanisms. The likely absence of target-site insensitivity in *P. regilla* indicates that exposure to tetrodotoxin outside of trophic interactions may not strongly select for fixed mechanisms of toxin resistance.

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

Acknowledgments.—We affirm the sovereignty of the Karkin, Me-Wuk (Bay Miwok), Muwekma, Chochenyo, Ohlone, and Ramaytush people and the Confederated Villages of Lisjan; we thank them for stewarding the land where we worked and, thus, serving an integral role in the production of this study. We thank L. Smith in the Evolutionary Genetics Lab, as well as C. Spencer and the support of the Museum of Vertebrate Zoology collections staff. We also thank those who supported our work at field sites: UC Botanical Garden staff, including H. Forbes, J. Fong, E. Hupperts, and C. Loughran; East Bay Regional Parks District staff, including T. Lin and J. Miller; and J. Young with the Presidio Trust. We thank the other members of the Tarvin Lab for their editing prowess and general support. We thank A. Shabel for his advice in the early stages of this project, and K. Sanko, J. Toman, and L. Galindo who helped as field assistants. We appreciate the newts and frogs who were the focus of this project and who were sacrificed to offer the knowledge they hold. K. Montana thanks her family and ancestors for their guidance. R. Tarvin was supported by UC Berkeley start-up funds.

The East Bay Regional Parks District authorized the surveys at Briones under permit #19-1044, and staff at the UCBG gave written consent for surveys conducted on the premises. All animals were collected under approved collection permits and protocols: for the National Parks Service Scientific Research and Collecting Permit 2 number PRSF-2020-SCI-0002 within study PRSF-00035, with a written extension after expiration of the original permit, University of California IACUC AUP-2019-08-12457, and California Department of Fish and Wildlife Scientific Collection Permit S-190980001-19111-001.

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

Spanish translation was done using www.DeepL.com/Translator (free version),
which was then edited by Valeria Ramírez Castañeda.

LITERATURE CITED

(see attached document)

SUPPLEMENTARY ONLINE MATERIAL

Supporting Methods.—Specimens sequenced in this study, SCN4A exon 24
Table S1.—DNA Extraction Procedures, Adapted from Miller et al., 1988.
Table S2.— Nav1.4 (SCN4A exon 24) and COI sequences from GenBank
Supplementary Figure 1.—San Francisco Bay Area GBIF Observations of
Pseudacris regilla and *Taricha*, 1971-2021 (N 38.66226, S 37.09298, E -121.12701, W -
123.23579)

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

FIGURES

Fig. 1. Examples of *Pseudacris regilla* and *Taricha torosa* interactions: A. Amplexus in Briones Regional Park, Martinez, CA, 1 March 2019, photo credit K. Montana, user kmontana on iNaturalist (CC BY-NC 4.0; observation 100725148); B. Another instance of amplexus in Briones Regional Park, Martinez, CA, 21 February 2009, photo credit user ap2il on iNaturalist (CC BY-NC 4.0; observation 550227); C. Attempted predation on *Taricha torosa* by *Pseudacris regilla* in Mark West Springs, CA, 1 March 2014, photo credit user tiwane on iNaturalist (CC BY-NC 4.0; observation 2312). The frogs did not appear to be harmed in any of these scenarios, though the tongue posture in C suggests some numbing effects from newt exposure.

Fig. 2. Field sites: A. Old Briones Road Trail 1 (OBRT 1) pond, 37.94441, -122.13377; B. Old Briones Road Trail 2 (OBRT 2) pond, 37.94352, -122.14102; C. Japanese Pool at the UC Botanical Garden (UCBG), 37.87440, -122.23760; D. Presidio site, 37.788496, -122.468605; all photos taken by K. Montana.

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

Fig. 3. Numbers of A. *P. regilla* adults, B. *Taricha torosa* and *T. granulosa* adults and larvae, and C. *T. torosa* egg masses observed over field trips to the Old Briones Road Trail 1 (OBRT 1) (blue circle line), OBRT 2 (red triangle line), and U.C. Botanical Garden (yellow square line) ponds; 20 October 2019 to late February 2020.

Fig. 4. Percentage of total abundance of GBIF-recorded individuals of *Pseudacris regilla* (green) and *Taricha* (blue) in California by month, over the course of the last 50 years. *Taricha* includes *T. granulosa*, *T. rivularis*, *T. sierrae*, *T. torosa*, as well as those *Taricha* species no longer recognized (*T. lindoei*, *T. miocenica*, and *Palaeotaricha oligocenica*) plus individuals just identified to the genus *Taricha* (GBIF). Peak abundances for *P. regilla* occur around April and August each year, and peak abundances for *Taricha* occur in January each year. Samples included were from 1 January 1971 to 23 November 2021.

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

Fig. 5. Map of the state of California, USA showing location of *Pseudacris regilla* museum specimens sympatric with *Taricha* (green squares), *P. regilla* museum specimens allopatric with *Taricha* (orange triangles), and field sites (black squares) in relation to occurrences of *Taricha* (blue circles).

Fig. 6. Alignment of the p-loop of domain IV of the voltage-gated sodium channel Nav1.4 in several species including *Pseudacris regilla*, mapped onto a COI phylogeny inferred using IQTREE (support values are SH-aLRT/ultrafast bootstrap values from 1000 iterations each). Amino acid positions are numbered according to *Rattus norvegicus*, UniProt accession #P15390. Species marked with an asterisk* are known to be resistant to TTX, and the AA substitutions known or hypothesized to confer TTX resistance are shown in bold (Feldman et al., 2012; Hanifin and Gilly, 2015). The amino acid substitution I1519M is found in *P. regilla* as well as *N. parkeri* and *R. tigrinus*. *R. tigrinus* consumes *Polypedates leucomystax*, which wields tetrodotoxin and bufadienolides; the I1519M substitution is thought to be linked to TTX resistance (Hutchinson et al., 2007; Tikhonov and Zhorov, 2005; Tikhonov and Zhorov, 2011). *N. parkeri* has the I1519M substitution but is not known to consume or be resistant to any toxins. Close relatives of *P. regilla*, *P. cadaverina* and *A. crepitans*, do not have the I1519M substitution or any others. *E. epinephelus* has AA substitutions that are thought

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

569 to be connected to resistance to other toxins in their diets; they eat *Atelopus* toads that
 570 have guanidine alkaloids including tetrodotoxin as well as dendrobatid poison dart frogs
 571 that wield batrachotoxin and pumiliotoxin (Myers et al., 1978; Feldman et al., 2012;
 572 Ramírez Castañeda et al., 2017).

573

574

575

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

- 1 ABDEREMANE-ALI, F., N.D. ROSSEN, M.E. KOBIELA, R.A. CRAIG II, C.E. GARRISON, Z. CHEN,
- 2 C.M. COLLERAN, L.A. O'CONNELL, J. DU BOIS, J.P. DUMBACHER, AND D.L. MINOR, JR.
- 3 2021. Evidence that toxin resistance in poison birds and frogs is not rooted in
- 4 sodium channel mutations and may rely on 'toxin sponge' proteins." Journal of
- 5 General Physiology 153:e202112872. <https://doi.org/10.1085/jgp.202112872>.
- 6 Adobe Inc. 2019. Adobe Illustrator. Retrieved from
- 7 <https://adobe.com/products/illustrator>.
- 8 BAIRD, S.F. 1854. Descriptions of new genera and species of North American frogs.
- 9 Proceedings of the Academy of Natural Sciences of Philadelphia 7:59–62.
- 10 BARROW, L.N., H.F. RALICKI, S.A. EMME, AND E.M LEMMON. 2014. Species tree estimation
- 11 of North American chorus frogs (Hylidae: *Pseudacris*) with parallel tagged
- 12 amplicon sequencing. Molecular Phylogenetics and Evolution 75:78–90.
- 13 BRODIE, E.D., JR. 1968. Investigations on the skin toxin of the adult rough-skinned
- 14 newt, *Taricha granulosa*. Copeia 1968:307–313.
- 15 BRODIE, E.D., JR., J.L. HENSEL, AND J.A. JOHNSON. 1974. Toxicity of the urodele
- 16 amphibians *Taricha*, *Notophthalmus*, *Cynops* and *Paramesotriton*
- 17 (Salamandridae). *Copeia* 1974:506-511.
- 18 BRODIE, E.D., JR. B.J. RIDENHOUR, AND E. D. BRODIE III. 2002. The evolutionary
- 19 response of predators to dangerous prey: hotspots and coldspots in the
- 20 geographic mosaic of coevolution between garter snakes and newts.
- 21 Evolution 56:2067–2082.
- 22 BUCCIARELLI, G.M., AND L.B. KATS. 2015. Effects of newt chemical cues on the
- 23 distribution and foraging behavior of stream macroinvertebrates. Hydrobiologia

749:69–81.

CALHOUN, D.M., G.M. BUCCIARELLI, L.B. KATS, R.K. ZIMMER, AND P.T.J. JOHNSON. 2017. Noxious newts and their natural enemies: experimental effects of tetrodotoxin exposure on trematode parasites and aquatic macroinvertebrates. *Toxicon: Official Journal of the International Society on Toxinology* 137:120–127.

CHAHINE, M., P.B. BENNETT, A.L. GEORGE, AND R. HORN. 1994. Functional expression and properties of the human skeletal muscle sodium channel. *Pflugers Archive: European Journal of Physiology* 427:136–142.

CHE, J., W.W. ZHOU, J.S. HU, F. YAN, T.J. PAPENFUSS, D.B. WAKE, AND Y.P. ZHANG. 2010. Spiny frogs (*Paini*) illuminate the history of the Himalayan region and southeast Asia. *Proceedings of the National Academy of Sciences* 107:13765–13770.

CHERNOMOR, O., A. VON HAESELER, AND B.Q. MINH. 2016. Terrace aware data structure for phylogenomic inference from supermatrices. *Systematic Biology* 65:997–1008.

CHIAMVIMONVAT, N., M.T. PÉREZ-GARCÍA, G.F. TOMASELLI, AND E. MARBAN. 1996. Control of ion flux and selectivity by negatively charged residues in the outer mouth of rat sodium channels. *The Journal of Physiology* 491:51–59.

CROTHER, B.I., SOCIETY FOR THE STUDY OF AMPHIBIANS AND REPTILES, COMMITTEE ON STANDARD ENGLISH AND SCIENTIFIC NAMES, AMERICAN SOCIETY OF ICHTHYOLOGISTS AND HERPETOLOGISTS, CANADIAN ASSOCIATION OF HERPETOLOGISTS, PARTNERS IN AMPHIBIAN AND REPTILE CONSERVATION, SOCIETY FOR THE STUDY OF AMPHIBIANS AND REPTILES, AND HERPETOLOGISTS' LEAGUE. 2017. *Scientific and Standard*

English Names of Amphibians and Reptiles of North America North of Mexico, with Comments Regarding Confidence in Our Understanding, Eighth Edition.

DALY, J.W., F. GUSOVSKY, C.W. MYERS, M. YOTSU-YAMASHITA, AND T. YASUMOTO. 1994. First occurrence of tetrodotoxin in a dendrobatid frog (*Colostethus inguinalis*), with further reports for the bufonid genus *Atelopus*. *Toxicon* 32:279–285.

DUELLMAN, W.E., A.B. MARION, AND S. B. HEDGES. 2016. Phylogenetics, classification, and biogeography of the treefrogs (Amphibia: Anura: Arboranae). *Zootaxa* 4104:1–109.

DURSO, A.M., L.A. NEUMAN-LEE, G.R. HOPKINS, AND E.D. BRODIE, JR. 2021. Stable isotope analysis suggests that tetrodotoxin-resistant Common Gartersnakes (*Thamnophis sirtalis*) rarely feed on newts in the wild. *Canadian Journal of Zoology* 99:331–338.

FELDMAN, C.R., E.D. BRODIE, JR., E.D. BRODIE III, AND M.E. PFRENDER. 2009. The evolutionary origins of beneficial alleles during the repeated adaptation of garter snakes to deadly prey. *Proceedings of the National Academy of Sciences* 106: 13415-13420.

FELDMAN, C.R., E.D. BRODIE, JR., E.D. BRODIE III, AND PFRENDER, M.E. 2012. Constraint shapes convergence in tetrodotoxin-resistant sodium channels of snakes. *Proceedings of the National Academy of Sciences* 109: 4556-4561.

FOUQUETTE JR., M. J., AND A. DUBOIS. 2014. A checklist of North American amphibians and reptiles: the United States and Canada. Xlibris Corporation.

GALL, B.G., A.N. STOKES, S.S. FRENCH, E.A. SCHLEPPHORST, E.D. BRODIE III, AND E.D. BRODIE, JR. 2011. Tetrodotoxin levels in larval and metamorphosed newts

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

70 (*Taricha granulosa*) and palatability to predatory dragonflies. *Toxicon* 57:978–
 71 983.

72 GALL, B.G., A.N. STOKES, S.S. FRENCH, E.D. BRODIE III, AND E.D. BRODIE, JR. 2012.
 73 Predatory caddisfly larvae sequester tetrodotoxin from their prey, eggs of the
 74 rough-skinned newt (*Taricha granulosa*). *Journal of Chemical Ecology* 38: 1351–
 75 1357.

76 GBIF.org (23 November 2021) GBIF Occurrence Download
 77 <https://doi.org/10.15468/dl.qzrq4k>.

78 GBIF.org (23 November 2021) GBIF Occurrence Download
 79 <https://doi.org/10.15468/dl.z9tw6r>.

80 GBIF.org (30 November 2021) GBIF Occurrence Download
 81 <https://doi.org/10.15468/dl.dwz9yn>.

82 GBIF.org (30 November 2021) GBIF Occurrence Download
 83 <https://doi.org/10.15468/dl.ed2uqn>.

84 GEFFENEY, S.L., E. FUJIMOTO, E.D. BRODIE III, E.D. BRODIE, JR., AND P.C. RUBEN. 2005.
 85 Evolutionary diversification of TTX-resistant sodium channels in a predator–prey
 86 interaction. *Nature* 434: 759–763.

87 GENDREAU, K.L., A.D. HORNSBY, M.T.J. HAGUE, AND J.W. MCGLOTHLIN. 2021. Gene
 88 conversion facilitates the adaptive evolution of self-resistance in highly toxic
 89 newts. *Molecular Biology and Evolution*:4077–4094.

90 Geneious Prime 2020.1.1. <https://www.geneious.com>.

91 HAGUE, M.T.J., G. TOLEDO, SHANA L. GEFFENEY, C.T. HANIFIN, E.D. BRODIE, JR., AND E.D.
 92 BRODIE, III. 2018. Large-effect mutations generate trade-off between predatory

- 93 and locomotor ability during arms race coevolution with deadly prey. *Evolution*
- 94 *Letters* 2:406–416.
- 95 HANIFIN, C.T. 2010. The chemical and evolutionary ecology of tetrodotoxin (TTX) toxicity
- 96 in terrestrial vertebrates. *Marine Drugs* 8:577–593.
- 97 HANIFIN, C.T., E.D. BRODIE JR, AND E.D. BRODIE III. 2008. Phenotypic mismatches reveal
- 98 escape from arms-race coevolution. *PLOS Biology* 6:e60.
- 99 <https://doi.org/10.1371/journal.pbio.0060060>.
- 100 HANIFIN, C.T., AND W.F. GILLY. 2015. Evolutionary history of a complex adaptation:
- 101 tetrodotoxin resistance in salamanders. *Evolution; International Journal of*
- 102 *Organic Evolution* 69: 232–244.
- 103 HASHIGUCHI, Y., J. M. LEE, M. SHIRAIISHI, S. KOMATSU, S. MIKI, Y. SHIMASAKI, N. MOCHIOKA,
- 104 T. KUSAKABE, AND Y. OSHIMA. 2015. Characterization and evolutionary analysis of
- 105 tributyltin-binding protein and pufferfish saxitoxin and tetrodotoxin-binding protein
- 106 genes in toxic and nontoxic pufferfishes. *Journal of Evolutionary Biology*
- 107 28:1103–1118.
- 108 HO, B., D.S. YEO, AND J.L. DING. 1994. A tetrodotoxin neutralizing system in the
- 109 haemolymph of the horseshoe crab, *Carcinoscorpius rotundicauda*. *Toxicon:*
- 110 *Official Journal of the International Society on Toxinology* 32:755–762.
- 111 HOANG, D.T., O. CHERNOMOR, A. VON HAESELER, B.Q. MINH, AND L.S. VINH. 2018.
- 112 UFBoot2: improving the ultrafast bootstrap approximation. *Molecular Biology and*
- 113 *Evolution* 35:518–522.
- 114 HUTCHINSON, D.A., A. MORI, A.H. SAVITZKY, G.M. BURGHARDT, X. WU, J. MEINWALD, AND
- 115 F.C. SCHROEDER. 2007. Dietary sequestration of defensive steroids in nuchal

116 glands of the Asian snake *Rhabdophis tigrinus*. Proceedings of the National
117 Academy of Sciences 104:2265–2270.

118 HWANG, P.A., Y.H. TSAI, H.P. LIN, AND D.F. HWANG. 2007. Tetrodotoxin-binding proteins
119 isolated from five species of toxic gastropods. Food Chemistry 103:1153–1158.

120 JAMES, D., AND K. HORNIK. 2020. chron: chronological objects which can handle dates
121 and times. R package version 2.3-56.

122 KAO, C.Y., AND S.R. LEVINSON. 1986. Preface—tetrodotoxin, saxitoxin, and the
123 molecular biology of the sodium channel. Annals of the New York Academy of
124 Sciences 479:ix-x.

125 KASSAMBARA, A. 2020. ggpubr: 'ggplot2' Based Publication Ready Plots. R package
126 version 0.4.0. <https://CRAN.R-project.org/package=ggpubr>.

127 LEE, C.H., D.K. JONES, C. AHERN, M.F. SARHAN, AND P.C. RUBEN. 2011. Biophysical costs
128 associated with tetrodotoxin resistance in the sodium channel pore of the garter
129 snake, *Thamnophis sirtalis*. Journal of Comparative Physiology A 197:33–43.

130 LLEWELLYN, L.E., P.M. BELL, AND E.G. MOCZYDLOWSKI. 1997. Phylogenetic survey of
131 soluble saxitoxin-binding activity in pursuit of the function and molecular evolution
132 of saxiphilin, a relative of transferrin. Proceedings of the Royal Society B:
133 Biological Sciences 264:891–902.

134 MAGARLAMOV, T.Y., D.I. MELNIKOVA, AND A.V. CHERNYSHEV. 2017. Tetrodotoxin-
135 producing bacteria: detection, distribution and migration of the toxin in aquatic
136 systems. Toxins:166.

137 MAHAR, J., G.L. LUKÁCS, L. LI, S. HALL, AND E. MOCZYDLOWSKI. 1991. Pharmacological
138 and biochemical properties of saxiphilin, a soluble saxitoxin-binding protein from

- 139 the bullfrog (*Rana catesbeiana*). *Toxicon* 29:53–71.
- 140 MCGLOTHLIN, J.W., M.E. KOBIELA, C.R. FELDMAN, T.A. CASTOE, S.L. GEFFENEY, C.T.
- 141 HANIFIN, G. TOLEDO, F.J. VONK, M.K. RICHARDSON, E.D. BRODIE, JR., M.E.
- 142 PFRENDER, E.D. BRODIE, III. 2016. Historical contingency in a multigene family
- 143 facilitates adaptive evolution of toxin resistance. *Current Biology* 26:1616–1621.
- 144 MEBS, D., M. YOTSU-YAMASHITA, AND O. ARAKAWA. 2016. The praying mantis (*Mantodea*)
- 145 as predator of the poisonous red-spotted newt *Notophthalmus viridescens*
- 146 (Amphibia: Urodela: Salamandridae). *Chemoecology* 26:121–126.
- 147 MECHAM, J.S. 1967. *Notophthalmus viridescens*. Catalogue of American Amphibians
- 148 and Reptiles: 53.1-53.4.
- 149 MILLER, S.A., D.D. DYKES, AND H.F. POLESKY. 1988. A simple salting out procedure for
- 150 extracting DNA from human nucleated cells. *Nucleic Acids Research* 16:1215.
- 151 MONIZ, H.A., M.A. RICHARD, C.M. GIENGER, AND C.R. FELDMAN. 2021. Every breath you
- 152 take: assessing metabolic costs of toxin resistance in garter snakes
- 153 (*Thamnophis*). *Integrative Zoology*. <https://doi.org/10.1111/1749-4877.12574>.
- 154 MYERS, C.W., J.W. DALY, AND B. MALKIN. 1978. A dangerously toxic new frog
- 155 (*Phylllobates*) used by Emberá Indians of western Colombia, with discussion of
- 156 blowgun fabrication and dart poisoning. *Bulletin of the AMNH* 161:article 2.
- 157 NAGASHIMA, Y., K. YAMAMOTO, K. SHIMAKURA, AND K. SHIOMI. 2002. A tetrodotoxin-binding
- 158 protein in the hemolymph of shore crab *Hemigrapsus sanguineus*: purification
- 159 and properties. *Toxicon: Official Journal of the International Society on*
- 160 *Toxinology* 40:753–760.
- 161 NEUWIRTH, E. 2014. RColorBrewer: ColorBrewer palettes. R package version 1.1-2.

- 162 <https://CRAN.R-project.org/package=RColorBrewer>.
- 163 NGUYEN, L., H.A. SCHMIDT, A. VON HAESELER, AND B.Q. MINH. 2015. IQ-TREE: A fast and
- 164 effective stochastic algorithm for estimating maximum likelihood phylogenies.
- 165 Molecular Biology and Evolution 32:268-274.
- 166 NOAA (National Oceanic and Atmospheric Association). 2022. Climate data online
- 167 search—data tables for Berkeley, CA and Martinez CA.
- 168 <https://www.ncdc.noaa.gov/cdo-web/search>.
- 169 OTA, W.M., B. OLSEN, G.M. BUCCIARELLI, AND L.B. KATS. 2018. The effect of newt toxin
- 170 on an invasive snail. Hydrobiologia 817:341–348.
- 171 PARADIS E., AND K. SCHLIEP. 2019. ape 5.0: an environment for modern phylogenetics
- 172 and evolutionary analyses in R. Bioinformatics 35:526-528.
- 173 PIRES, O.R., A. SEBBEN, E.F. SCHWARTZ, S.W.R. LARGURA, C. BLOCH, R.A.V. MORALES,
- 174 AND C.A. SCHWARTZ. 2002. Occurrence of tetrodotoxin and its analogues in the
- 175 Brazilian frog *Brachycephalus ephippium* (Anura: Brachycephalidae). Toxicon:
- 176 Official Journal of the International Society on Toxinology 40:761–766.
- 177 RAMÍREZ-CASTAÑEDA, V. 2017. Origin of mutations in the NaV gene family associated
- 178 with resistance to neurotoxins in snake (*Erythrolamprus sp.*) predators of dart
- 179 poison frogs (Dendrobatidae). Master’s thesis. University of the Andes, Bogota,
- 180 Colombia.
- 181 R Core Team. 2021. R: A language and environment for statistical computing. R
- 182 Foundation for Statistical Computing, Vienna, Austria.
- 183 <https://www.R-project.org>.
- 184 RECUERO, E., Í. MARTÍNEZ-SOLANO, G. PARRA-OLEA, AND M. GARCÍA-PARÍS. 2006.

Phylogeography of *Pseudacris regilla* (Anura: Hylidae) in western North America, with a proposal for a new taxonomic rearrangement. *Molecular Phylogenetics and Evolution* 39:293–304.

RECUERO, E., Í. MARTÍNEZ-SOLANO, G. PARRA-OLEA, AND M. GARCÍA-PARÍS. 2006. Corrigendum to ‘Phylogeography of *Pseudacris regilla* (Anura: Hylidae) in western North America, with a proposal for a new taxonomic rearrangement’ [Mol. Phylogenet. Evol. 39 (2006) 293–304]. *Molecular Phylogenetics and Evolution* 41:511.

REIMCHE, J.S., E.D. BRODIE JR., A.N. STOKES, E.J. ELY, H.A. MONIZ, V.L. THILL, J.M. HALLAS, M.E. PFRENDER, E.D. BRODIE III, AND C.R. FELDMAN. 2020. The geographic mosaic in parallel: matching patterns of newt tetrodotoxin levels and snake resistance in multiple predator–prey pairs. *Journal of Animal Ecology* 89:1645–1657.

REVELL, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* 3:217-223.

RITSON-WILLIAMS, R., M. YOTSU-YAMASHITA, AND V.J. PAUL. 2006. Ecological functions of tetrodotoxin in a deadly polyclad flatworm. *Proceedings of the National Academy of Sciences* 103:3176–3179.

SIEVERS F., A. WILM, D.G. DINEEN, T.J. GIBSON, K. KARPLUS, W. LI, R. LOPEZ, H. MCWILLIAM, M. REMMERT, J. SÖDING, J.D. THOMPSON, D.G. HIGGINS. 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems Biology* 7:539.

TANU, M.B., Y. MAHMUD, K. TSURUDA, O. ARAKAWA, AND T. NOGUCHI. 2001. Occurrence of

208 tetrodotoxin in the skin of a rhacophoridid frog *Polypedates* sp. from Bangladesh.

209 Toxicon: Official Journal of the International Society on Toxinology 39:937–941.

210 TARVIN, R.D., J.C. SANTOS, L.A. O'CONNELL, H.H. ZAKON, AND D.C. CANNATELLA. 2016.

211 Convergent substitutions in a sodium channel suggest multiple origins of toxin

212 resistance in poison frogs. Molecular Biology and Evolution 37:607.

213 TIKHONOV, D.B., AND B.S. ZHOROV. 2005. Modeling p-loops domain of sodium channel:

214 homology with potassium channels and interaction with ligands. Biophysical

215 Journal 88:184–197.

216 TIKHONOV, D.B., AND B.S. ZHOROV. 2011. Possible roles of exceptionally conserved

217 residues around the selectivity filters of sodium and calcium channels. Journal of

218 Biological Chemistry 286:2998–3006.

219 TRIFINOPOULOS, J., L.T. NGUYEN, A. VON HAESELER, AND B.Q. MINH. 2016. W-IQ-TREE: a

220 fast online phylogenetic tool for maximum likelihood analysis. Nucleic Acids

221 Research 44:W232-W235.

222 VAELLI, P.M., K.R. THEIS, J.E. WILLIAMS, L.A. O'CONNELL, J.A. FOSTER, AND H.L. EISTHEN.

223 2020. The skin microbiome facilitates adaptive tetrodotoxin production in

224 poisonous newts. ELife 9:e53898. <https://doi.org/10.7554/eLife.53898>.

225 VENKATESH, B., S.Q. LU, N. DANDONA, S.L. SEE, S. BRENNER, AND T.W. SOONG. 2005.

226 Genetic basis of tetrodotoxin resistance in pufferfishes. Current Biology 15:2069–

227 2072.

228 VITT, L. J., AND J. P. CALDWELL. 2013. Herpetology: An Introductory Biology of

229 Amphibians and Reptiles. United Kingdom and United States of America.

230 WANNINGER, M., T. SCHWAHA, AND E. HEISS. 2018. Form and function of the skin glands

- 231 in the Himalayan newt *Tylototriton verrucosus*. Zoological Letters 4:15.
- 232 WICKHAM, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New
233 York. United States of America.
- 234 WICKHAM, H., M. AVERICK, J. BRYAN, W. CHANG, L.D. MCGOWAN, R. FRANÇOIS, G.
235 GROLEMUND, A. HAYES, L. HENRY, J. HESTER, M. KUHN, T.L. PEDERSEN, E. MILLER,
236 S.M. BACHE, K.MÜLLER, J. OOMS, D. ROBINSON, D.P. SEIDEL, V. SPINU, K.
237 TAKAHASHI, D. VAUGHAN, C. WILKE, K. WOO, AND H. YUTANI. 2019. Welcome to the
238 tidyverse. Journal of Open Source Software 4:1686.
239 <https://doi.org/10.21105/joss.01686>.
- 240 WICKHAM, H., R. FRANÇOIS, L. HENRY, AND K. MÜLLER. 2021. dplyr: a grammar of data
241 manipulation. R package version 1.0.7.
242 <https://CRAN.R-project.org/package=dplyr>.
- 243 WILKE, C.O. 2020. cowplot: streamlined plot theme and plot annotations for 'ggplot2'.
244 R package version 1.1.1. <https://CRAN.R-project.org/package=cowplot>.
- 245 WILLIAMS, B.L. 2010. Behavioral and chemical ecology of marine organisms with respect
246 to tetrodotoxin. Marine Drugs 8:381–398.
- 247 WILSON, N.J., A.N. STOKES, G.R. HOPKINS, E.D. BRODIE, JR., AND C.R. WILLIAMS. 2014.
248 Functional and physiological resistance of crayfish to amphibian toxins:
249 tetrodotoxin resistance in the white river crayfish (*Procambarus acutus*).
250 Canadian Journal of Zoology 92:939–945.
- 251 ZANGER, U.M., AND M. SCHWAB. 2013. Cytochrome P450 enzymes in drug metabolism:
252 regulation of gene expression, enzyme activities, and impact of genetic variation.

K.O. Montana et al.

Tetrodotoxin Resistance in *Pseudacris regilla*

253 Pharmacology & Therapeutics 138:103-41.
 254 <https://doi.org/10.1016/j.pharmthera.2012.12.007>
 255 ZIMMER, R.K., D.W. SCHAR, R.P. FERRER, P.J. KRUG, L.B. KATS, AND W.C. MICHEL. 2006.
 256 The scent of danger: tetrodotoxin (TTX) as an olfactory cue of predation risk.
 257 Ecological Monographs 76:585–600.
 258
 259
 260
 261
 262
 263

DIV S5-S6 p-loop

1519

FGNSIICLF EITTSAGWDGLLN P

SD

M

L V NV

S QS SD I

S QS SD I

M Q

M