1 **Meta-omics-aided isolation of an elusive anaerobic arsenic-methylating soil bacterium**

2

3 Karen Viacava^{1,2}, Jiangtao Qiao¹, Andrew Janowczyk³, Suresh Poudel⁴, Nicolas Jacquemin⁵,
4 Karin Lederballe Meibom¹, Him K. Shrestha^{4,6}, Matthew C. Reid⁷, Robert L. Hettich⁴, Rizlan
5 Bernier-Latmani^{1*}.

6

7 ¹ Environmental Microbiology Laboratory, School of Architecture, Civil and Environmental
8 Engineering, École Polytechnique Fédérale de Lausanne, Lausanne, Switzerland.

9 ² Soil Science Group, Institute of Geography, University of Bern, Bern, Switzerland.

10 ³ Bioinformatics Core Facility, Swiss Institute of Bioinformatics, Lausanne, Switzerland.

11 ⁴ BioSciences Division, Oak Ridge National Laboratory, Oak Ridge, TN, United States.

12 ⁵ Translational Bioinformatics and Statistics, Department of Oncology, Université de
13 Lausanne, Lausanne, Switzerland.

14 ⁶ Genome Science and Technology Graduate School, University of Tennessee, Knoxville,
15 TN, United States.

16 ⁷ School of Civil and Environmental Engineering, Cornell University, Ithaca, NY, United
17 States.

18 *Corresponding author. E-mail address rizlan.bernier-latmani@epfl.ch. Postal address EPFL
19 ENAC IIE EML CH A1 375 (Bâtiment CH) Station 6 CH-1015 Lausanne Switzerland. Tel.
20 +41 21 69 35001.

21 Competing Interests Statement: The authors declare there are no competing financial
22 interests in relation to the work described. The work was funded by the Swiss National
23 Science Foundation (SNSF) grant 310030_176146-1 and the SNSF NCCR Microbiomes
24 (grant # 51NF40_180575). The work at ORNL was conducted under the Plant-Microbe
25 Interface Science Focus Area, as supported by the U.S. Department of Energy, Office of
26 Science, Office of Biological and Environmental Research, Genomic Science Program.

27 Key words: metagenomics, metaproteomics, metatranscriptomics, *Paraclostridium*, arsenic
28 metabolism.

29

30 **Abstract**

31 Soil microbiomes harbour unparalleled functional and phylogenetic diversity. However,
32 extracting isolates with a targeted function from complex microbiomes is not straightforward,
33 particularly if the associated phenotype does not lend itself to high-throughput screening.
34 Here, we tackle the methylation of arsenic (As) in anoxic soils. As methylation was proposed
35 to be catalysed by sulphate-reducing bacteria. However, to date, there are no available
36 anaerobic isolates capable of As methylation, whether sulphate-reducing or otherwise. The
37 isolation of such a microorganism has been thwarted by the fact that the anaerobic bacteria
38 harbouring a functional arsenite S-adenosylmethionine methyltransferase (ArsM) tested to
39 date did not methylate As in pure culture. Additionally, fortuitous As methylation can result
40 from the release of non-specific methyltransferases upon lysis. Thus, we combined
41 metagenomics, metatranscriptomics, and metaproteomics to identify the microorganisms
42 actively methylating As in anoxic soil-derived microbial cultures. Based on the metagenome-
43 assembled genomes of microorganisms expressing ArsM, we isolated *Paraclostridium* sp.
44 strain EML, which was confirmed to actively methylate As anaerobically. This work is an
45 example of the application of meta-omics to the isolation of elusive microorganisms.

46

47 Introduction

48 Soil microbiomes represent a rich source of novel metabolisms and taxa [1–4]. However,
 49 isolating microorganisms from them to study specific functions can be challenging, and even
 50 more so in cases for which the phenotype is not identifiable with high-throughput methods
 51 [5, 6]. An example of challenging microorganisms to isolate are anaerobic As-methylating
 52 strains. Arsenic methylation, catalysed by arsenite (As(III)) S-adenosylmethionine
 53 methyltransferase (ArsM, in prokaryotes), entails the binding of one to three methyl group(s)
 54 to the As atom [7]. At present, there are no available microorganisms capable of anaerobic
 55 As methylation. This is because, adding to the constraints associated with maintaining an
 56 anoxic environment [8], there is no assay for ArsM activity that can be adapted for high-
 57 throughput assessment, despite recent endeavours [9]. Arsenic methylation occurs in
 58 anoxic, flooded rice paddy soils, is mediated by soil microorganisms [10], and results in the
 59 accumulation of methylated As in rice grains [11]. The bioaccumulation of methylated As in
 60 rice grains is considerably more efficient than that of inorganic As [12, 13].

61 The gene encoding ArsM (*arsM*) has been identified in phylogenetically diverse soil
 62 microorganisms [14–17]. Anaerobic As methylation is expected to produce a toxic trivalent
 63 monomethylated As species (MMAs(III)). The function of this transformation is hypothesized
 64 to be microbial warfare, by which the As-methylating organism inhibits microbial competitors
 65 via the production of MMAs(III) [18, 19]. If that is confirmed, it is conceivable that As
 66 methylation may not occur in pure cultures but only in microbial communities, triggered by
 67 metabolites produced by the microbiota. Alternatively, *arsM*-harbouring microorganisms that
 68 express As(III) efflux pump(s), the major pathway of As resistance within bacteria [20], may
 69 not methylate As due to the efficient removal of As(III) from the cytoplasm, which is the
 70 location of ArsM [21, 22]. This effect could be direct, i.e., insufficient substrate concentration,
 71 or indirect, i.e., the intracellular As(III) concentration is too low to induce *arsM* expression.
 72 Either occurrence (microbial warfare or rapid As(III) efflux) would render the isolation of pure
 73 cultures of As-methylating anaerobes very challenging using standard approaches. The

latter hypothesis is supported by recent work showing the lack of As methylation by anaerobic pure cultures harbouring functional ArsM enzymes [22].

An additional complexity is evidence for the fortuitous methylation of As upon cell lysis and the release of methyltransferases. This fortuitous activity was suggested for the methanogen *Methanosarcina mazei*, for which As methylation was initiated only when cell viability decreased [22], and by the *in vitro* methylation of various metals, including As, by MtaA, a methyltransferase involved in methanogenesis [23]. Thus, As methylation activity in cultures incubated beyond the exponential phase may simply be an experimental artefact [22]. Finally, the detection of methylated As requires relatively complex analytical tools (high pressure liquid chromatography coupled to inductively-coupled plasma mass spectrometry, HPLC-ICP-MS) that do not lend themselves readily to high-throughput screening of a large number of colonies [9]. As a result of these challenges, there are no anaerobic microorganisms available known to actively methylate As despite many efforts to identify them. In one instance, researchers had identified a Gram-positive sulphate-reducing bacterium (SRB) [24] that was reported to methylate As but this isolate is no longer available, precluding further investigation.

Thus, this study aimed to conclusively identify an active anaerobic As methylator in soil-derived microbial cultures using a multi-omics approach. The experimental strategy was to build Metagenome-Assembled Genomes (MAGs) from metagenomic data and to identify the subset of MAGs harbouring the gene *arsM* that also expressed the *arsM* RNA transcript (metatranscriptomics) and/or the enzyme ArsM (metaproteomics). Based on the genetic information from the target MAG, an isolation strategy was devised that allowed the recovery of a pure culture, later confirmed to be a novel anoxic As-methylating strain.

Materials and methods

Rice paddy soil microbiomes

The soil-derived cultures consisted of two anaerobic microbial enrichments derived from a Vietnamese rice paddy soil and described in Reid *et al.* [25]. The microbiota from the first

soil-derived microbiome was grown in ¼ strength tryptic soy broth (TSB) medium (7.5 g l⁻¹ TSB), used previously to enrich As-methylating microbes from a lake sediment [26], and henceforth referred to as the TSB culture. The medium for the second soil-derived microbiome, in addition to ¼ strength TSB, included electron acceptors and two additional carbon sources to simultaneously allow the growth of nitrate-, iron-, and sulphate-reducers, as well as fermenters and methanogens (EA medium: 5 mM NaNO₃, 5 mM Na₂SO₄, 5 mM ferric citrate, 0.2 g l⁻¹ yeast extract (Oxoid, Hampshire, UK) and 1 g l⁻¹ cellobiose, pH 7). This enrichment will be referred to as the EA culture. Both media were boiled, cooled down under 100% N₂ gas and 50 ml of medium were dispensed into 100-ml serum bottles. The bottle headspace was flushed with 100% N₂ gas prior to autoclaving. All culture manipulations were carried out using N₂-flushed syringes and needles. Cultures were grown at 30°C. Growth was quantified using optical density at 600 nm (OD₆₀₀).

Arsenic methylation assays

Pre-cultures from each enrichment were started from -80°C glycerol stocks. The EA culture started from the glycerol stock was transferred only after a dark precipitate, presumably iron sulphide resulting from sulphate reduction, was formed. The first experimental set-up consisted of bottles containing medium amended with As(III) as NaAsO₂ (+As condition) pre-inoculation or unamended (no-As control). For this set-up, cell pellets were sampled for DNA sequencing and proteome characterization during the stationary phase, and for RNA sequencing at the mid-exponential growth phase (see Figures S1, S2 and S3 in Supplementary Information (SI)). In a second experimental set-up, cultures were grown in unamended (no As(III) added) medium and As(III) was added at the mid-exponential growth phase. For this set-up, cell pellets were sampled before (no-As control) and 30 min after As amendment (+As condition) and were used solely for a second transcriptomic analysis. Triplicate biological experiments were performed for each condition (no-As, +As) and per soil-derived enrichment and were used for DNA and RNA sequencing and metaproteome characterization. Sampling for soluble As species, determination of As speciation, and total

As concentration are described in SI.

DNA sequencing and metagenomic analysis

DNA was extracted from the pellet (10 min, 4,500 g) of 4 ml of culture using the DNeasy Power Soil Kit (Qiagen, Hilden, Germany) homogenizing with a Precellys 24 Tissue Homogenizer (Bertin Instruments, Montigny-le-Bretonneux, France) (6,500 rpm for 10 s, repeated 3x with 10 s pause intervals). Metagenomic sequencing was performed by the Genomics Platform of the University of Geneva, Switzerland (iGE3) on a HiSeq 4000 (Illumina, San Diego, CA, US). Libraries were multiplexed and prepared using 100-base reads with paired ends according to the Nextera DNA Flex Library Preparation Kit protocol (Illumina). The quality of sequence reads was assessed with FastQC [27] and duplicated reads eliminated by FastUniq [28]. Reads from all biological replicates within the same experimental condition were assembled into contigs using MegaHit [29]. The contig abundance was determined by aligning the sequencing reads from each biological replicate back to the assembled contigs using Kallisto [30, 31]. The abundance for each gene was considered equivalent to the abundance of the contig in which it was encoded. Gene abundance is reported as 'transcripts per million' (TPM), referred to as TPM-DNA when used for gene abundance. TPM includes normalization for gene length and read sequencing depth [32]. Prodigal was used for the prediction of protein-coding genes [33], generating protein sequence libraries for each culture (EA, TSB) and condition (no-As control, +As condition). The annotation server GhostKOALA [34] was used to assign a KEGG Orthology (KO) database number to each protein-coding gene to identify its encoded function and taxonomic category. The 16S small subunit (SSU) rRNA sequences were identified in the contigs and their taxonomy assigned by Metaxa2 [35]. The relative abundance of the 16S SSU rRNA sequences identified in each of the four metagenomes was quantified using the Kallisto-calculated contig abundance. Contigs with length >2,000 bp were clustered into bins based on composition and coverage using CONCOCT [36], MetaBAT2 [37] and MaxBin 2.0 [38]. The final bin set was obtained by using the Bin_refinement module from MetaWRAP

[39]. Completeness, contamination, strain heterogeneity and community (%) in contigs for each bin were calculated using CheckM [40]. Matching bins between the no-As and +As metagenomes, and between the EA +As and TSB +As metagenomes were identified by pairwise comparison of the predicted genomes using dRep [41]. Bins with an average nucleotide identity >95% were considered identical genomes.

RNA sequencing and metatranscriptomic analysis

Each culture (5 ml) was harvested at mid-exponential phase for metatranscriptomic analysis. The cells were lysed and the RNA purified using the RNeasy Mini Kit following the manufacturer's instructions (RNAprotect Bacteria, Qiagen). The purified RNA was DNase-I treated (Promega, Madison, WI, US) (1h, 37 °C) and cleaned using the RNeasy Mini Kit a second time. Ribosomal RNA (rRNA) depletion (kit QIAseq FastSelect –5S/16S/23S, Qiagen), library preparation using single-end 100 bases reads (TrueSeq Stranded mRNA, Illumina) and RNA sequencing (on a HiSeq 4000) were performed by the iGE3 Platform. Reads were quality-assessed by FastQC, trimmed by Trimmomatic [42], post-sequencing rRNA-depleted by SortMeRNA [43] and aligned to their corresponding protein sequence library by Bowtie2 [44]. The program featureCounts [45] was employed to count the number of RNA reads aligned to the Prodigal-predicted protein-coding genes. The raw counts were used to calculate the TPM, referred as TPM-RNA when employed for transcript abundance. Finally, to assess RNA expression changes in the +As condition relative to the no-As condition, a differential abundance analysis was performed using the DESeq2 package [46] using the protein sequence libraries from the +As condition to align the RNA reads. A gene was considered to have a significant difference in transcription when the absolute log₂ fold change was ≥1 (i.e., 0.5 ≥fold change ≥2) and the adjusted *q* value ≤0.05.

Metaproteome characterization and metaproteomic analysis

The metaproteome analysis was performed at Oak Ridge National Laboratory (Oak Ridge, TN, US). Biomass pellets from 100 ml of culture were washed with 100 mM NH₄HCO₃ buffer

(ABC) (pH 8.0), re-suspended in lysis buffer (4% sodium dodecyl sulphate, 100 mM Tris-HCl, pH 8.0) and disrupted by bead-beating. Lysate proteins were reduced with 5 mM dithiothreitol (30 min, 37 °C), alkylated with 15 mM iodoacetamide (30 min in the dark, room temperature) and isolated by a chloroform-methanol extraction. Extracted proteins were solubilized in 4% sodium deoxycholate (SDC) in ABC and the concentration estimated with a Nanodrop (Thermo Fisher Scientific, Waltham, MA, US). Sequencing-grade trypsin (Promega) at a 1:75 enzyme:protein ratio (w/w) was used to digest the proteins and formic acid (1% final concentration) was used to precipitate the SDC and collect tryptic peptides. Aliquots of 12 µg of peptides were analysed by 2D LC-MS/MS consisting of a Vanquish UHPLC connected to a Q Exactive Plus MS (Thermo Fisher Scientific). Spectral data were collected using MudPIT (multidimensional protein identification technology) as described previously [47, 48]. Peptides were separated in three steps (35, 100, and 500 mM ammonium acetate eluent) with organic gradients after each step. Eluted peptides were measured and sequenced by data-dependent acquisition using previously described parameters [49].

Protein databases were created for the +As experimental condition (EA +As and TSB +As) from the corresponding protein sequence libraries generated by Prodigal. The MS/MS spectra raw files were processed in Proteome Discoverer version 2.4 (Thermo Fisher Scientific) with MS Amanda 2.0 [50] and Percolator [51]. Spectral data were searched against the protein database of the corresponding culture (i.e., EA or TSB). The following parameters were used in the search algorithm MS-Amanda 2.0 to derive tryptic peptides: MS1 tolerance = 5 ppm; MS2 tolerance = 0.02 Da; missed cleavages = 2; carbamidomethyl (C, + 57.021 Da) as static modification; and oxidation (M, + 15.995 Da) as dynamic modifications. The false discovery rate (FDR) threshold was set to 1% for strict FDR and 5% for relaxed FDR at the peptide-spectrum matched (PSM), peptide, and protein levels. FDR-controlled peptides were then quantified according to the chromatographic area-under-the-curve and mapped to their respective proteins. Areas were summed to estimate protein-level abundance.

For differential abundance analysis of proteins, the spectral data from the no-As control, EA no As and TSB no As, were searched against the EA +As and TSB +As protein databases, respectively. All the above-described parameters were maintained. The proteins with at least one peptide detected were exported from Proteome Discoverer. Protein data matrix from EA no As and EA +As were merged and TSB no As and TSB +As were merged. Protein abundance values were $\log_2$ transformed, LOESS-normalized among biological replicates and mean-centred across all conditions using the software InfernoRDN [52]. Stochastic sampling of the proteins was filtered by removing the proteins without abundance value in at least two of the biological triplicates in at least one condition (no-As control or +As condition). Remaining missing data were imputed by random numbers drawn from a normal distribution (width = 0.3 and downshift = 2.8 using the Perseus software <http://www.perseus-framework.org>) [53]. The differentially abundant proteins were identified by Student t-test method with adjusted q value ≤ 0.05 . Proteins were further filtered using the absolute $\log_2$ fold change ≥ 1 .

The isolation of the *Paraclostridium* sp. strain EML is described in SI.

Results

Arsenic methylation by soil-derived microbiomes

The first experimental set-up yielded samples for the metagenome, metaproteome and one of the metatranscriptomes (labeled metatranscriptome G for 'growth in the presence of As') (Figures S1, S2 and S3). The second set-up, assessing the microbiota's short-term response to As(III), provided sample for the second metatranscriptome (labeled metatranscriptome R for 'response to arsenic addition') (Figures S1-A and S2-A). Both EA and TSB cultures exhibited As methylation, reaching an efficiency of As(III) transformation of 27.7% and 19.5%, respectively (Figures S1 and S2).

Microbiota composition

The taxonomic classification of 16S SSU rRNA sequences show that, although eukaryotic

DNA was also identified, the main fraction of the communities was bacterial ($>89.0 \pm 0.8\%$ for EA cultures and $>98.5 \pm 0.3\%$ for TSB cultures, relative abundance) and was distributed amongst eight operational taxonomic units (OTUs) at the order level (Figure 1 and Tables S1-S4). Statistically significant changes (unpaired Student t-test and no significant difference considered when p value > 0.05) in the OTUs relative abundances, +As condition versus no-As control, are described in SI and summarized in Tables S5 and S6.

MAG selection

The contigs from the four metagenomes, EA (+As, no-As control) and TSB (+As, no-As control), were clustered separately into bins. High-quality ($\geq 90\%$ completeness and $\leq 5\%$ contamination) bins were designated as MAGs [54]. For the +As condition, the parsing process led to a total of 36 MAGs (Table 1). Additionally, matching bins were sought in the bins from the no-As control cultures (Tables S7 and S8). Only one of the 36 MAGs in the +As condition was left unpaired (TSB MAG 8).

For each MAG, a lineage was assigned by CheckM, based on lineage-specific marker genes [40]. The MAGs identified belonged to the phyla: *Firmicutes* (orders *Clostridiales*, *Selenomonadales* and *Lactobacillales*, and the genus *Clostridium*), *Proteobacteria* (*Enterobacteriaceae* family and *Deltaproteobacteria* class) and *Bacteroidetes* (order *Bacteroidales*). Fifteen MAGs presented non-zero strain heterogeneity (Table 1), an index of the phylogenetic relatedness of binned contigs based on the amino acid identity of the encoded proteins. For ten MAGs, the value is $\geq 50\%$, suggesting some phylogenetic relation with the contaminating strains. Five MAGs had heterogeneity values $\leq 33.33\%$, suggesting contamination with microorganisms that are not closely related. In the remaining 21 MAGs, the strain heterogeneity is 0%, i.e., no strain heterogeneity or no contamination (Tables S7 and S8).

Changes in the relative abundance of MAGs (no-As control vs. +As condition), relatedness of the +As EA and TSB microbial communities, along with the presence, transcription and translation of genes encoding key enzymes from major metabolic pathways of each MAG in

the +As condition are included in SI.

Arsenic resistance genes

The metagenomic libraries from the +As condition of the EA and TSB cultures were mined for arsenic resistance (*ars*) genes and their encoded proteins (the pipeline is described in SI). A total of 309 and 282 genes were annotated as *ars* genes in the EA and TSB +As metagenomic libraries, respectively (Tables S9 and S10). Of those, 255 and 226 were considered correctly annotated as *ars* genes based on BLAST and HMMER (refer to SI for pipeline), and 225 and 147 had above-threshold DNA abundances, respectively (Figure 2) (refer to SI for abundance threshold values). Individual abundance values of *ars* genes, transcripts and proteins in the +As condition and the no-As control and their transcript and protein relative abundance values in the +As condition vs. the no-As control for each MAG group from the EA and TSB cultures are available in Tables S11 and S12, respectively.

The *ars* genes encode proteins involved in the detoxification of As oxyanions: *arsB* and *acr3*, encoding As(III)-efflux systems; *arsA*, encoding the ATPase energizing the efflux of As(III) and As(III) chaperone; *arsD*, encoding a weak *ars* operon repressor [55]; *arsC1* and *arsC2*, encoding As(V) reductases coupling As reduction to the oxidation of glutaredoxin or thioredoxin, respectively; and *arsR* genes encoding As(III)-regulated repressors (ArsR1, ArsR2, and ArsR3) classified based on the location of the As(III)-binding cysteine residues [56–58].

The most common *ars* genes in EA and TSB culture metagenomes were *arsR*, *arsC*, and *arsP* (Figure 2). The first two genes are part of the canonical *ars* operon *arsRBC* [59], whilst *arsP*, encoding a recently discovered membrane transporter, has been found to be widely distributed in bacterial genomes [20]. Most of the surveyed *arsP* genes, 57% in EA and 50% in TSB, are encoded in putative *ars* operons, represented by *ars* genes contiguously encoded in the same contig (Tables S11 and S12), supporting their As-related function and correct annotation. The next most abundant genes were those responsible for As(III) efflux (*arsB*, *acr3*, and *arsA*), typically found in organisms living in reducing environments in

association with *arsC* [16, 60]. Finally, *arsM* and the two genes, *arsI* and *arsH*, encoding MMAs(III)-resistance mechanisms, were the least recurrent genes in the metagenomes. The results of gene and protein relative expression vs. the no-As control of the *ars* genes involved in the metabolism of inorganic As in the MAGs are described in SI.

Arsenic-methylating MAGs

The *arsM* gene can be expressed at similar, or slightly different levels in the absence or presence of As(III) in some organisms [61, 62], but expressed at significantly higher levels in the presence of As(III) in others [63–66]. Thus, we sought to identify *arsM* genes transcribed and ArsM proteins showing increased expression in the +As condition relative to the no-As control (Figure 3) but also those simply exhibiting expression, not necessarily increased relative to the control (Figure S4).

Sixteen phylogenetically distinct *arsM* genes were identified in the EA +As metagenome, but increased transcriptome reads or peptides (relative to the no-As control) were only detected for three genes (Figure 3). The first is an *arsM* in *Clostridiales* EA MAG 8 classified by GhostKOALA as belonging to *Paeniclostridium sordellii* (EA MAG 8, *arsM*-1, psor type strain, in Table S11). The second was found in *Clostridium* EA MAG 9, also detected in the metaproteome, and the GhostKOALA taxonomic classification of the corresponding gene (EA MAG 9, *arsM*-1 in Table S11) revealed that it was attributed to the unclassified species *Ruminococcaceae bacterium* CPB6 (Figure 3, Table S11) [67]. Finally, the third *arsM* was obtained from transcriptomic data but not clustered in any EA MAG (EA unbinned, *arsM*-5 in Table S11) and likewise classified as pertaining to *Paeniclostridium sordellii*.

In the TSB +As metagenome, nine distinct *arsM* genes were identified but none were detected in the metatranscriptome and only one exhibited increased expression in the metaproteome (Figure 3). It corresponds to an *arsM* gene from MAG 9 (TSB MAG 9, *arsM*-2 in Table S12). The expressed ArsM protein was assigned by GhostKOALA to a *Clostridiales* strain: *Clostridium botulinum* (*cby* type strain) (TSB MAG 9, *arsM*-2) (Figure 3, Table S12). Finally, there was one *arsM* expressed in the TSB +As metaproteome but with no increased

expression relative to the no-As control, it was classified as *Ruminococcaceae bacterium* CPB6 (TSB MAG 9, *arsM-1*) (Figure S4), the same organism identified in the EA culture (EA MAG 9, *arsM-1*).

In addition to evidence for active As methylation, there was evidence for active detoxification of methylated arsenic. Indeed, the metagenome included genes encoding proteins involved in the metabolism of methylated As such as *arsH*, *arsI*, *arsP*, and *arsR4* (Figures 2 and 3). These genes encode proteins involved in the detoxification of methylated arsenic like MMAs(III) and roxarsone: the oxidase ArSH, responsible for the oxidation of trivalent methylated As to the less toxic pentavalent form [68]; the demethylase ArSI that removes methyl groups from the As atom [69]; and the transmembrane transporter ArSP, thought to efflux methylated As [70]. The *arsR4* gene encodes an atypical MMAs(III)-responsive ArsR repressor, containing only two conserved cysteine residues [71]. The *Enterobacteriaceae* TSB MAG 14 exhibited activity of the oxygen-dependent ArSH protein [68] (Figure 3). An *arsR4*, shown to induce expression of *arsP* in the presence of MMAs(III) [71], had increased transcription along with an *arsP* encoded in the same contig in the *Selenomonadales* TSB MAG 19 (Figure 3, Table S12). Both gene transcripts were <5 TPM-RNA (Table S12) and thus, were not considered as transcribed in Figure S4. Finally, an ArSI protein, taxonomically related to class *Clostridia* ([*Eubacterium*] *rectale*), was expressed but encoded in an unbinned gene from the EA culture (Figure 3, Table S12).

Isolation of an arsenic-methylating anaerobic microorganism

Based on the analysis of the active metabolic activity from the EA MAG 8, expressing an ArsM (Figure S5), an appropriate selective medium was identified for its isolation. We utilized the fact that this MAG harbours and expresses the anaerobic assimilatory sulphite reductase encoded by the *asrABC* operon which is responsible for the NADH-dependent reduction of sulphite to sulphide [72–74] in sulphite-reducing *Clostridia* (SRC). From the nine *Clostridia* MAGs, only two expressed this capability in the EA microbiome (Figure S5). Thus, the isolation relied on growing the EA culture on agar medium selective for the SRC

phenotype. In TSC agar, designed for the enumeration of *Clostridium perfringens* in food [75], the colonies from SRC are black, as the ammonium ferric citrate forms iron sulphide during sulphite reduction. Additionally, D-cycloserine acts a selective agent for the isolation of *Clostridia* strains [76] while inhibiting facultative anaerobes [75]. Finally, the bromocresol purple contained in the agar allows the identification of sucrose fermenters, resulting in a change of colour from purple to yellow. As none of the genes involved in sucrose transport or hydrolysis were binned in EA *Clostridiales* MAG 8 (Figure S5), only non-sucrose fermenting black colonies were considered. Those colonies were selected and using a colony PCR screen specifically targeting the *arsM* gene of EA MAG 8, we isolated a *Clostridiales* strain encoding the gene of the expressed ArsM in the EA MAG 8 (protein id k119_30669_28, Table S11) (Figure S6).

The isolate consists of non-sucrose-fermenting, rod-shaped and spore-forming bacteria forming convex and circular black colonies on TSC agar (Figures S7 and S8). The BLAST (NCBI) search of the 16S rRNA sequence gives >99% identity to *Paraclostridium* strains (Table S13). On the basis of the 16S rRNA sequence, we assign the following name to the bacterium: “*Paraclostridium* species str. EML”. Strain EML was tested for As methylation under anaerobic conditions with 25 μ M As(III). The growth of strain EML was hindered by As(III) (Figure 4-A) and starting from ~4 hours, the isolate transformed As(III) to monomethylated soluble As representing $48.3 \pm 1.5\%$ of the soluble arsenic in the culture after 83 h (panels B and C from Figure 4). A fraction ($14.7 \pm 0.6 \mu$ M) of the arsenic was found associated with biomass almost exclusively as inorganic As (Figure 4-D).

Discussion

Our results demonstrate the successful translation of multi-omic information to a specific strategy for targeted microbial isolation. The metagenomes from the anaerobic soil-derived cultures identified the potential for As methylation in microorganisms from diverse taxa. While there were a large number of *ars* genes in the metagenomes, only a small proportion

was transcribed or translated in the presence of As when compared to the no-As control (Figure 2). This contrast was particularly evident for the gene responsible for As methylation, *arsM*. The post-genomic approaches of community gene and protein expression in TSB as in EA soil-derived microbiomes clearly pointed to the active As-methylating role of various fermenting bacteria from the order *Clostridiales*. This information paved the way for the identification of As-methylating microorganisms and the successful isolation of an anaerobic As methylator.

The TSB and EA media were chosen to selectively enrich for putative As methylators from the microbial soil community based on the study from Bright *et al.*, in which lake sediments enriched in TSB medium, either sulphate-amended or unamended, were shown to have greater As methylation rates than in iron- or manganese-reducing TSB cultures [26]. The selected media caused a great shift in the original soil microbial diversity [25] along with the loss of putative As-methylating microorganisms. Nonetheless, the As-methylating TSB and EA soil-derived cultures offered the opportunity to study active As methylation from paddy-soil microbiota in an environment that is less complex than soil but that remains environmentally relevant. In contrast to soil slurries, the absence of soil minerals in the soil-derived cultures facilitated the detection of soluble methylarsenicals and the extraction of DNA, RNA and proteins. The multi-omic approach made it possible to identify putative microorganisms driving As methylation and their metabolism. Targeting a specific *arsM* gene rather than the synthesis of methylarsenicals greatly accelerated colony screening, as colony PCR could be employed instead of analytical detection by HPLC-ICP-MS.

Had only the metagenomic approach been implemented, the data would have pointed to SRB MAGs as putative As methylators, as they harboured the most abundant *arsM* genes (Figure 5). Indeed, SRB have been proposed as drivers of As methylation in rice paddy soils based on the correlation in the abundance of *arsM* and dissimilatory sulphite reductase (*dsr*) genes [77] and RNA transcripts [78], and a decrease in As methylation by the addition chemical inhibitors of dissimilatory sulphate reduction (DSR) [77, 78]. Additionally, the use of degenerate primers for *arsM* amplification may underestimate *arsM* phylogenetic diversity, a

drawback overcome by metagenomic and metatranscriptomic sequencing. In the present findings, the SRB *Deltaproteobacteria* MAGs, although actively reducing sulphate (Figures S5 and S9), did not exhibit As-methylating activity as their *arsM* genes were neither transcribed nor translated (Figure 5). *Desulfovibrio* MAGs were metabolically active in both cultures, but amongst all their encoded *ars* genes, only an *arsR3* exhibited increased expression in the presence of As(III), providing strong evidence for their lack of involvement in As methylation in the TSB and EA cultures.

Previous work had identified another As-methylating *Clostridiales* strain, *Clostridium* sp. BXM [24], that performed fermentation and DSR but that is no longer available. The sole attribution of As-methylating activity to fermenting Firmicutes in that work, along with the isolation of the present sulphite-reducing fermenter, point to a key role for fermenting *Clostridiales* microorganisms harbouring sulphur-related metabolism in As methylation. Other studies have reported an increase in As methylation efficiency after the amendment of sulphate [79] or organic matter to soil [15, 80–82], or after the increase in dissolved organic carbon in soil [83]. The positive impact of sulphate amendment on As methylation was interpreted as pointing to the role of SRB in As methylation [79]. Here, we offer an alternative explanation, supported by examples of organic amendments enhancing As methylation [15, 80–82]. The sulphate amendment could have indirectly increased the availability of short-chain fatty acids through DSR, providing fermentable substrates. Thus, we propose that direct or indirect organic amendments result in the enrichment of fermenting communities, and consequently, in an increase in As methylation.

It was previously proposed that the As-methylating activity of anaerobic microorganisms may be limited by efficient efflux of intracellular As(III) [22], or that it may function as a defensive response against nutrient competition [18]. Indeed, the identification of MAGs exhibiting a detoxification response to methylarsenicals supports the hypothesis of the role of monomethylated As as an arsenic-bearing antibiotic. Although the expression of ArsI and ArsH, catalysing oxygen-dependent MMAs(III)-resistance mechanisms (Figure 3), is difficult to reconcile with anoxic conditions, it is conceivable that these proteins are capable of

additional functions in the absence of O₂. Up until now, the lack of available anaerobic microbial isolates able to methylate As *in vitro* [22] precluded the investigation of the hypotheses raised above. This work represents the first study applying a combination of three meta-omic techniques in order to characterize As metabolism in microbial communities and to perform meta-omics-aided isolation of a microorganism [84][85]. The successful isolation of *Paraclostridium* sp. EML is part of the “new era of omics information-guided microbial cultivation technology” described by Gutleben *et al.* [84] and represents a milestone to obtain novel targeted microbial isolates from the environment and to elucidate the controls on anaerobic As methylation.

Further work is needed to elucidate why ArsM expression was restricted to members of *Clostridiales* fermenters and did not occur in other organisms harbouring *arsM* genes. The availability of As-methylating anaerobes will allow investigation of why the *arsM* gene evolved under an anoxic atmosphere [86], of the controls on the production of toxic methylated As species in flooded rice paddies, and the development of microbially-mediated remediation technologies for As-contaminated soils via the synthesis of volatile methylarsenicals [87, 88].

Acknowledgements

The work was funded by the Swiss National Science Foundation (SNSF) grant 310030_176146-1 and the SNSF NCCR Microbiomes (grant # 51NF40_180575). The work at ORNL was conducted under the Plant-Microbe Interface Science Focus Area, as supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, Genomic Science Program.

Data availability

Metagenomic and metatranscriptomic raw sequencing reads are available at the National Centre for Biotechnology Information (NCBI) Sequence Read Archive (SRA), BioProject PRJNA714492 (post publication). Data from the meta-omic analyses and source data from

figures are available in Zenodo data repository (10.5281/zenodo.4605527, post publication).

Conflict of interest

The authors declare no conflict of interest.

Figures Legends

Figure 1. Operational taxonomic units (OTUs) at order and genus level (with > 1% relative abundance at genus level) identified from 16S SSU rRNA sequences from soil-derived cultures. Abbreviations: EA no As: EA culture no-As control EA +As: EA culture +As condition, TSB no As: TSB culture no-As control TSB +As: TSB culture +As condition. OTUs at the order level are indicated in bold in the legend. Plotted values are the average relative abundance and together with SD values and Student t test results are available in Tables S5 and S6.

Figure 2. Number of *ars* genes, encoded in the +As condition cultures and with increased expression in metatranscriptomes/metaproteome relative to no-As controls. Number of *ars* genes encoded in metagenomes and with increased expression in metatranscriptomes, R or G, or metaproteomes and the non-redundant overlap between genes with increased expression in metatranscriptomes and/or metaproteomes from +As condition EA (left panels) and TSB (right panels) cultures. Bar length and numbers on the right side of the bars correspond to the number of genes per *ars* gene category. Bold numbers on the lower left corner of each panel correspond to the sum of all *ars* genes per category.

Figure 3. Distribution of *ars* genes involved in methylated arsenic metabolism encoded in MAGs from the +As condition and differentially expressed in metatranscriptomes/metaproteome relative to the no-As EA control. Each numbered box represents an *ars* gene. The number in each box corresponds to the “Numbering”

column in Tables S11 and S12 where individual gene abundance and fold change values can be found. Com. (%): community (%) as defined in caption from Table 1.

Figure 4. Isolate “*Paraclostridium* sp. EML”: (A) growth as OD₆₀₀ with 25 µM As(III) and without, (B) proportion of soluble arsenic species in filtered medium containing 25 µM As(III), (C) concentration of arsenic species soluble in filtered medium containing 25 µM As(III) (solid lines) and biomass-bound (dashed lines) and (D) proportion of biomass-bound arsenic species. Data points and bars represent the mean value and error bars, plus and minus one standard deviation. Individual values for each measurement and biological replicate are available in Tables S23 and S24.

Figure 5. Gene abundance of *arsM* genes in MAGs from the +As condition cultures. Coloured bars correspond to *arsM* genes with increased expression in the metaproteome (blue-green) or in the metatranscriptome G (purple) from +As condition relative to the no-As control in EA (left panel) and TSB (right panel) cultures. The taxonomic classification shown on the right side of the error bars for selected *arsM* genes corresponds to the individual gene classification assigned by GhostKOALA - “Genus” column in Tables S11 and S12. Columns with matching symbols on the right side of the error bars, correspond to matching *arsM* genes between the cultures. Individual gene abundance can be found in Tables S11 and S12. Numbers inside bar and bar length represent mean and error bars one standard deviation.

Table Legend

Table 1. Metagenome assembled genomes (MAGs) from EA (upper Table A) and TSB (lower Table B) cultures in the +As condition. Marker lineage: taxonomic rank set by CheckM. Completeness and contamination (%): estimated completeness and contamination of genome as determined by CheckM from the presence/absence of single-copy marker genes and the expected colocalization of these genes. Strain heterogeneity: index between

0 and 100 where a value of 0 means no strain heterogeneity, high values suggest the majority of reported contamination is from closely related organisms (i.e., potentially the same species) and low values suggest the majority of contamination is from phylogenetically diverse sources. Proportion of binned proteins assigned to MAG (%): number of protein-coding genes assigned to the MAG divided by the total number of protein-coding genes binned. Community (%): sum of the number of reads mapped to the contigs in each MAG divided by the total number of reads mapped to all contigs including the unbinned contigs, and normalized to MAG size, assuming an average genome size for all unbinned populations.

References

1. Whitman WB, Coleman DC, Wiebe WJ. Prokaryotes: The unseen majority. *Proc Natl Acad Sci* 1998; **95**: 6578–6583.
2. Bardgett RD, Van Der Putten WH. Belowground biodiversity and ecosystem functioning. *Nature* 2014; **515**: 505–511.
3. Roesch LFW, Fulthorpe RR, Riva A, Casella G, Hadwin AKM, Kent AD, *et al.* Pyrosequencing enumerates and contrasts soil microbial diversity. *ISME J* 2007; **1**: 283–290.
4. Daniel R. The soil metagenome - A rich resource for the discovery of novel natural products. *Curr Opin Biotechnol* 2004; **15**: 199–204.
5. Bodor A, Bounedjoum N, Vincze GE, Erdeiné Kis Á, Laczi K, Bende G, *et al.* Challenges of unculturable bacteria: environmental perspectives. *Rev Environ Sci Biotechnol* 2020; **19**: 1–22.
6. Houle D, Govindaraju DR, Omholt S. Phenomics: The next challenge. *Nat Rev Genet* 2010; **11**: 855–866.
7. Ajees AA, Marapakala K, Packianathan C, Sankaran B, Rosen BP. Structure of an As(III) S-adenosylmethionine methyltransferase: Insights into the mechanism of arsenic biotransformation. *Biochemistry* 2012; **51**: 5476–5485.
8. Mori K, Kamagata Y. The challenges of studying the anaerobic microbial world. *Microbes Environ* 2014; **29**: 335–337.
9. Dong H, Xu W, Pillai JK, Packianathan C, Rosen BP. High-throughput screening-compatible assays of As(III) S-adenosylmethionine methyltransferase activity. *Anal Biochem* 2015; **480**: 67–73.
10. Lomax C, Liu WJ, Wu L, Xue K, Xiong J, Zhou J, *et al.* Methylated arsenic species in plants originate from soil microorganisms. *New Phytol* 2012; **193**: 665–672.
11. Zhao FJ, Zhu YG, Meharg AA. Methylated arsenic species in rice: Geographical variation, origin, and uptake mechanisms. *Environ Sci Technol* 2013; **47**: 3957–3966.
12. Zheng MZ, Li G, Sun GX, Shim H, Cai C. Differential toxicity and accumulation of

- 560 inorganic and methylated arsenic in rice. *Plant Soil* 2013; **365**: 227–238.
- 561 13. Abedin MJ, Feldmann J, Meharg AA. Uptake kinetics of arsenic species in rice plants.
562 *Plant Physiol* 2002; **128**: 1120–1128.
- 563 14. Dunivin TK, Yeh SY, Shade A. A global survey of arsenic-related genes in soil
564 microbiomes. *BMC Biol* 2019; **17**: 45.
- 565 15. Jia Y, Huang H, Zhong M, Wang F-H, Zhang L-M, Zhu Y-G. Microbial arsenic
566 methylation in soil and rice rhizosphere. *Environ Sci Technol* 2013; **47**: 3141–3148.
- 567 16. Xiao KQ, Li LG, Ma LP, Zhang SY, Bao P, Zhang T, *et al.* Metagenomic analysis
568 revealed highly diverse microbial arsenic metabolism genes in paddy soils with low-
569 arsenic contents. *Environ Pollut* 2016; **211**: 1–8.
- 570 17. Zhang SY, Su JQ, Sun GX, Yang Y, Zhao Y, Ding J, *et al.* Land scale biogeography
571 of arsenic biotransformation genes in estuarine wetland. *Environ Microbiol* 2017; **19**:
572 2468–2482.
- 573 18. Chen J, Rosen BP. The arsenic methylation cycle: How microbial communities
574 adapted methylarsenicals for use as weapons in the continuing war for dominance.
575 *Front Environ Sci* 2020; **8**: 43.
- 576 19. Ping Li Y, Ben Fekih I, Chi Fru E, Moraleda-Munoz A, Li X, Rosen BP, *et al.*
577 Antimicrobial activity of metals and metalloids. *Annu Rev Microbiol* 2021; **75**: 175–
578 197.
- 579 20. Yang Y, Wu S, Lilley RM, Zhang R. The diversity of membrane transporters encoded
580 in bacterial arsenic-resistance operons. *PeerJ* 2015; **3**: e943.
- 581 21. Yang P, Ke C, Zhao C, Kuang Q, Liu B, Xue X, *et al.* ArsM-mediated arsenite
582 volatilization is limited by efflux catalyzed by As efflux transporters. *Chemosphere*
583 2020; **239**: 124822.
- 584 22. Viacava K, Meibom KL, Ortega D, Dyer S, Gelb A, Falquet L, *et al.* Variability in
585 arsenic methylation efficiency across aerobic and anaerobic microorganisms. *Environ*
586 *Sci Technol* 2020; **54**: 14343–14351.
- 587 23. Thomas F, Diaz-Bone RA, Wuerfel O, Huber B, Weidenbach K, Schmitz RA, *et al.*

588 Connection between multimetal(loid) methylation in methanoarchaea and central
589 intermediates of methanogenesis. *Appl Environ Microbiol* 2011; **77**: 8669–8675.

590 24. Wang PP, Bao P, Sun GX. Identification and catalytic residues of the arsenite
591 methyltransferase from a sulfate-reducing bacterium, *Clostridium* sp. BXM. *FEMS*
592 *Microbiol Lett* 2015; **362**: 1–8.

593 25. Reid MC, Maillard J, Bagnoud A, Falquet L, Le Vo P, Bernier-Latmani R. Arsenic
594 methylation dynamics in a rice paddy soil anaerobic enrichment culture. *Environ Sci*
595 *Technol* 2017; **51**: 10546–10554.

596 26. Bright DA, Brock S, Reimer KJ, Cullen WR, Hewitt GM, Jafaar J. Methylation of
597 arsenic by anaerobic microbial consortia isolated from lake sediment. *Appl*
598 *Organomet Chem* 1994; **8**: 415–422.

599 27. Andrews S. FastQC: A quality control tool for high throughput sequence data [Online].
600 <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>. .

601 28. Xu H, Luo X, Qian J, Pang X, Song J, Qian G, *et al*. FastUniq: A fast *de novo*
602 duplicates removal tool for paired short reads. *PLoS One* 2012; **7**: e52249.

603 29. Li D, Liu CM, Luo R, Sadakane K, Lam TW. MEGAHIT: An ultra-fast single-node
604 solution for large and complex metagenomics assembly via succinct de Bruijn graph.
605 *Bioinformatics* 2015; **31**: 1674–1676.

606 30. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq
607 quantification. *Nat Biotechnol* 2016; **34**: 525–527.

608 31. Schaeffer L, Pimentel H, Bray N, Melsted P, Pachter L. Pseudoalignment for
609 metagenomic read assignment. *Bioinformatics* 2017; **33**: 2082–2088.

610 32. Wagner GP, Kin K, Lynch VJ. Measurement of mRNA abundance using RNA-seq
611 data: RPKM measure is inconsistent among samples. *Theory Biosci* 2012; **131**: 281–
612 285.

613 33. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal:
614 Prokaryotic gene recognition and translation initiation site identification. *BMC*
615 *Bioinformatics* 2010; **11**: 119.

- 616 34. Kanehisa M, Sato Y, Morishima K. BlastKOALA and GhostKOALA: KEGG tools for
617 functional characterization of genome and metagenome sequences. *J Mol Biol* 2016;
618 **428**: 726–731.
- 619 35. Bengtsson-Palme J, Hartmann M, Eriksson KM, Pal C, Thorell K, Larsson DGJ, *et al.*
620 Metaxa2: Improved identification and taxonomic classification of small and large
621 subunit rRNA in metagenomic data. *Mol Ecol Resour* 2015; **15**: 1403–1414.
- 622 36. Alneberg J, Bjarnason BS, De Bruijn I, Schirmer M, Quick J, Ijaz UZ, *et al.* Binning
623 metagenomic contigs by coverage and composition. *Nat Methods* 2014; **11**: 1144–
624 1146.
- 625 37. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, *et al.* MetaBAT 2: An adaptive
626 binning algorithm for robust and efficient genome reconstruction from metagenome
627 assemblies. *PeerJ* 2019; e7359.
- 628 38. Wu YW, Simmons BA, Singer SW. MaxBin 2.0: An automated binning algorithm to
629 recover genomes from multiple metagenomic datasets. *Bioinformatics* 2016; **32**: 605–
630 607.
- 631 39. Uritskiy G V., Diruggiero J, Taylor J. MetaWRAP - A flexible pipeline for genome-
632 resolved metagenomic data analysis. *Microbiome* 2018; **6**: 158.
- 633 40. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: Assessing
634 the quality of microbial genomes recovered from isolates, single cells, and
635 metagenomes. *Genome Res* 2015; **25**: 1043–1055.
- 636 41. Olm MR, Brown CT, Brooks B, Banfield JF. DRep: A tool for fast and accurate
637 genomic comparisons that enables improved genome recovery from metagenomes
638 through de-replication. *ISME J* 2017; **11**: 2864–2868.
- 639 42. Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina sequence
640 data. *Bioinformatics* 2014; **30**: 2114–2120.
- 641 43. Kopylova E, Noé L, Touzet H. SortMeRNA: Fast and accurate filtering of ribosomal
642 RNAs in metatranscriptomic data. *Bioinformatics* 2012; **28**: 3211–3217.
- 643 44. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods*

644 2012; **9**: 357–359.

645 45. Liao Y, Smyth GK, Shi W. featureCounts: An efficient general purpose program for
646 assigning sequence reads to genomic features. *Bioinformatics* 2014; **30**: 923–930.

647 46. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
648 RNA-seq data with DESeq2. *Genome Biol* 2014; **15**: 550.

649 47. Washburn MP, Wolters D, Yates JR. Large-scale analysis of the yeast proteome by
650 multidimensional protein identification technology. *Nat Biotechnol* 2001; **19**: 242–247.

651 48. McDonald WH, Ohi R, Miyamoto DT, Mitchison TJ, Yates JR. Comparison of three
652 directly coupled HPLC MS/MS strategies for identification of proteins from complex
653 mixtures: Single-dimension LC-MS/MS, 2-phase MudPIT, and 3-phase MudPIT. *Int J*
654 *Mass Spectrom* 2002; **219**: 245–251.

655 49. Clarkson SM, Giannone RJ, Kridelbaugh DM, Elkins JG, Guss AM, Michenera JK.
656 Construction and optimization of a heterologous pathway for protocatechuate
657 catabolism in Escherichia coli enables bioconversion of model aromatic compounds.
658 *Appl Environ Microbiol* 2017; **83**.

659 50. Dorfer V, Pichler P, Stranzl T, Stadlmann J, Taus T, Winkler S, *et al.* MS Amanda, a
660 universal identification algorithm optimized for high accuracy tandem mass spectra. *J*
661 *Proteome Res* 2014; **13**: 3679–3684.

662 51. Käll L, Canterbury JD, Weston J, Noble WS, MacCoss MJ. Semi-supervised learning
663 for peptide identification from shotgun proteomics datasets. *Nat Methods* 2007; **4**:
664 923–925.

665 52. Polpitiya AD, Qian WJ, Jaitly N, Petyuk VA, Adkins JN, Camp DG, *et al.* DAnTE: A
666 statistical tool for quantitative analysis of -omics data. *Bioinformatics* 2008; **24**: 1556–
667 1558.

668 53. Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, *et al.* The Perseus
669 computational platform for comprehensive analysis of (prote)omics data. *Nat Methods*
670 2016; **13**: 731–740.

671 54. Bowers RM, Kyripides NC, Stepanauskas R, Harmon-Smith M, Doud D, Reddy TBK,

- 672 *et al.* Minimum information about a single amplified genome (MISAG) and a
- 673 metagenome-assembled genome (MIMAG) of bacteria and archaea. *Nat Biotechnol*
- 674 2017; **35**: 725–731.
- 675 55. Lin Y-F, Walmsley AR, Rosen BP. An arsenic metallochaperone for an arsenic
- 676 detoxification pump. *PNAS* 2006; **103**: 15617–15622.
- 677 56. Shi W, Wu J, Rosen BP. Identification of a putative metal binding site in a new family
- 678 of metalloregulatory proteins. *J Biol Chem* 1994; **269**: 19826–19829.
- 679 57. Murphy JN, Saltikov CW. The ArsR repressor mediates arsenite-dependent regulation
- 680 of arsenate respiration and detoxification operons of *Shewanella sp.* strain ANA-3. *J*
- 681 *Bacteriol* 2009; **191**: 6722–6731.
- 682 58. Santha S, Pandaranayaka EPJ, Rosen BP, Thiyagarajan S. Purification,
- 683 crystallization and preliminary X-ray diffraction studies of the arsenic repressor ArsR
- 684 from *Corynebacterium glutamicum*. *Acta Crystallogr Sect F Struct Biol Cryst Commun*
- 685 2011; **67**: 1616–1618.
- 686 59. Fekih I Ben, Zhang C, Li YP, Zhao Y, Alwathnani HA, Saquib Q, *et al.* Distribution of
- 687 arsenic resistance genes in prokaryotes. *Front Microbiol* 2018; **9**: 2473.
- 688 60. Cai L, Yu K, Yang Y, Chen BW, Li XD, Zhang T. Metagenomic exploration reveals
- 689 high levels of microbial arsenic metabolism genes in activated sludge and coastal
- 690 sediments. *Appl Microbiol Biotechnol* 2013; **97**: 9579–9588.
- 691 61. Huang K, Xu Y, Packianathan C, Gao F, Chen C, Zhang J, *et al.* Arsenic methylation
- 692 by a novel ArsM As(III) S-adenosylmethionine methyltransferase that requires only
- 693 two conserved cysteine residues. *Mol Microbiol* 2018; **107**: 265–276.
- 694 62. Zhang J, Cao T, Tang Z, Shen Q, Rosen BP, Zhao FJ. Arsenic methylation and
- 695 volatilization by arsenite S-adenosylmethionine methyltransferase in *Pseudomonas*
- 696 *alcaligenes* NBRC14159. *Appl Environ Microbiol* 2015; **81**: 2852–2860.
- 697 63. Huang K, Chen C, Zhang J, Tang Z, Shen Q, Rosen BP, *et al.* Efficient arsenic
- 698 methylation and volatilization mediated by a novel bacterium from an arsenic-
- 699 contaminated paddy soil. *Environ Sci Technol* 2016; **50**: 6389–6396.

- 700 64. Yin XX, Chen J, Qin J, Sun GX, Rosen BP, Zhu YG. Biotransformation and
701 volatilization of arsenic by three photosynthetic cyanobacteria. *Plant Physiol* 2011;
702 **156**: 1631–1638.
- 703 65. Zhao C, Zhang Y, Chan Z, Chen S, Yang S. Insights into arsenic multi-operons
704 expression and resistance mechanisms in *Rhodopseudomonas palustris* CGA009.
705 *Front Microbiol* 2015; **6**: 986.
- 706 66. Wang G, Kennedy SP, Fasiludeen S, Rensing C, DasSarma S. Arsenic resistance in
707 Halobacterium sp. strain NRC-1 examined by using an improved gene knockout
708 system. *J Bacteriol* 2004; **186**: 3187–3194.
- 709 67. Tao Y, Zhu X, Wang H, Wang Y, Li X, Jin H, *et al.* Complete genome sequence of
710 *Ruminococcaceae* bacterium CPB6: A newly isolated culture for efficient n-caproic
711 acid production from lactate. *J Biotechnol* 2017; **259**: 91–94.
- 712 68. Chen J, Bhattacharjee H, Rosen BP. ArsH is an organoarsenical oxidase that confers
713 resistance to trivalent forms of the herbicide monosodium methylarsenate and the
714 poultry growth promoter roxarsone. *Mol Microbiol* 2015; **96**: 1042–1052.
- 715 69. Yoshinaga M, Rosen BP. A C-As lyase for degradation of environmental
716 organoarsenical herbicides and animal husbandry growth promoters. *PNAS* 2014;
717 **111**: 7701–7706.
- 718 70. Chen J, Madegowda M, Bhattacharjee H, Rosen BP. ArsP: A methylarsenite efflux
719 permease. *Mol Microbiol* 2015; **98**: 625–635.
- 720 71. Chen J, Nadar VS, Rosen BP. A novel MAs(III)-selective ArsR transcriptional
721 repressor. *Mol Microbiol* 2017; **106**: 469–478.
- 722 72. Doyle CJ, O'Toole PW, Cotter PD. Genomic characterization of sulphite reducing
723 bacteria isolated from the dairy production chain. *Front Microbiol* 2018; **9**: 1507.
- 724 73. Czyzewski BK, Wang DN. Identification and characterization of a bacterial
725 hydrosulphide ion channel. *Nature* 2012; **483**: 494–497.
- 726 74. Huang CJ, Barrett EL. Sequence analysis and expression of the *Salmonella*
727 *typhimurium* *asr* operon encoding production of hydrogen sulfide from sulfite. *J*

- 728 *Bacteriol* 1991; **173**: 1544–1553.
- 729 75. Harmon SM, Kautter DA, Peeler JT. Improved medium for enumeration of *Clostridium*
730 *perfringens*. *Appl Microbiol* 1971; **22**: 688–692.
- 731 76. George WL, Sutter VL, Finegold SM. Toxigenicity and antimicrobial susceptibility of
732 *Clostridium difficile*, a cause of antimicrobial agent-associated colitis. *Curr Microbiol*
733 1978; **1**: 55–58.
- 734 77. Wang M, Tang Z, Chen XP, Wang X, Zhou WX, Tang Z, *et al*. Water management
735 impacts the soil microbial communities and total arsenic and methylated arsenicals in
736 rice grains. *Environ Pollut* 2019; **247**: 736–744.
- 737 78. Chen C, Li L, Huang K, Zhang J, Xie WY, Lu Y, *et al*. Sulfate-reducing bacteria and
738 methanogens are involved in arsenic methylation and demethylation in paddy soils.
739 *ISME J* 2019; **13**: 2523–2535.
- 740 79. Zeng X, Jiang Y, Fan X, Chao S, Yang Y, Liu J, *et al*. Effects of sulfate application on
741 inhibiting accumulation and alleviating toxicity of arsenic in *Panax notoginseng* grown
742 in arsenic-polluted soil. *Water Air Soil Pollut* 2016; **227**.
- 743 80. Mestrot A, Feldmann J, Krupp EM, Hossain MS, Roman-Ross G, Meharg AA. Field
744 fluxes and speciation of arsines emanating from soils. *Environ Sci Technol* 2011; **45**:
745 1798–1804.
- 746 81. Huang H, Jia Y, Sun G-X, Zhu Y-G. Arsenic speciation and volatilization from flooded
747 paddy soils amended with different organic matters. *Environ Sci Technol* 2012; **46**:
748 2163–8.
- 749 82. Seyfferth AL, Morris AH, Gill R, Kearns KA, Mann JN, Paukett M, *et al*. Soil
750 incorporation of silica-rich rice husk decreases inorganic arsenic in rice grain. *J Agric*
751 *Food Chem* 2016; **64**: 3760–3766.
- 752 83. Zhao FJ, Harris E, Yan J, Ma J, Wu L, Liu W, *et al*. Arsenic methylation in soils and its
753 relationship with microbial *arsM* abundance and diversity, and As speciation in rice.
754 *Environ Sci Technol* 2013; 7147–7154.
- 755 84. Gutleben J, Chaib De Mares M, van Elsas JD, Smidt H, Overmann J, Sipkema D. The

756 multi-omics promise in context: From sequence to microbial isolate. *Crit Rev Microbiol*
757 2018; **44**: 212–229.

758 85. Chandran H, Meena M, Sharma K. Microbial biodiversity and bioremediation
759 assessment through omics approaches. *Front Environ Chem* 2020; **1**: 9.

760 86. Chen S-C, Sun G-X, Yan Y, Konstantinidis KT, Zhang S-Y, Deng Y, *et al.* The Great
761 Oxidation Event expanded the genetic repertoire of arsenic metabolism and cycling.
762 *PNAS* 2020; **117**: 10414–10421.

763 87. Liu S, Zhang F, Chen J, Sun G. Arsenic removal from contaminated soil via
764 biovolatilization by genetically engineered bacteria under laboratory conditions. *J*
765 *Environ Sci* 2011; **23**: 1544–1550.

766 88. Chen J, Sun GX, Wang XX, Lorenzo V De, Rosen BP, Zhu YG. Volatilization of
767 arsenic from polluted soil by *Pseudomonas putida* engineered for expression of the
768 *arsM* arsenic(III) S-adenosine methyltransferase gene. *Environ Sci Technol* 2014; **48**:
769 10337–10344.

770

EA +As

TSB +As

Encoded in metagenome

Increased expression in metatranscriptome R

Increased expression in metatranscriptome G

Increased expression in metaproteome

Increase expression in metatranscriptome G, metatranscriptome R and/or metaproteome

0

20

40

60

0

20

40

ars genes

Present in (+As condition):

Metagenome

Expression fold change (FC) vs. no-As control:

EA +As

TSB +As

Table 1. Metagenome assembled genomes (MAGs) from EA (upper Table A) and TSB (lower Table B) cultures in the +As condition. Marker lineage: taxonomic rank set by CheckM. Completeness and contamination (%): estimated completeness and contamination of genome as determined by CheckM from the presence/absence of single-copy marker genes and the expected colocalization of these genes. Strain heterogeneity: index between 0 and 100 where a value of 0 means no strain heterogeneity, high values suggest the majority of reported contamination is from closely related organisms (i.e., potentially the same species) and low values suggest the majority of contamination is from phylogenetically diverse sources. Proportion of binned proteins assigned to MAG (%): number of protein-coding genes assigned to the MAG divided by the total number of protein-coding genes binned. Community (%): sum of the number of reads mapped to the contigs in each MAG divided by the total number of reads mapped to all contigs including the unbinned contigs, and normalized to MAG size, assuming an average genome size for all unbinned populations.

A										
MAG	Bin	Marker lineage*	Completeness (%)	Contamination (%)	Strain heterogeneity (%)	Genome size (Mbp)	Community (%)	GC content	ArsM-encoding genes	Binner §
1	36	<i>Bacteroidales</i> (o)	98.5	0.4	0.0	3.8	7.26 ±0.32	39.1	0	A
2	15	<i>Clostridiales</i> (o)	98.7	0.0	0.0	2.2	0.52 ±0.06	58.4	0	B
3	21	<i>Clostridiales</i> (o)	95.2	0.0	0.0	4.3	5.73 ±0.24	28.5	0	C
4	4	<i>Clostridiales</i> (o)	90.7	0.0	0.0	2.1	0.30 ±0.03	57.5	0	A
5	24	<i>Clostridiales</i> (o)	97.8	0.3	0.0	2.0	25.21 ±0.23	43.2	0	A
6	9	<i>Clostridiales</i> (o)	100.0	1.3	50.0	3.2	12.68 ±0.68	54.9	0	A
7	35	<i>Clostridiales</i> (o)	98.0	3.3	55.6	5.3	0.84 ±0.21	44.0	0	A
8	31	<i>Clostridiales</i> (o)	97.9	3.5	0.0	3.8	1.48 ±0.12	28.2	1	A
9	20	<i>Clostridium</i> (g)	97.2	2.2	0.0	3.4	0.70 ±0.06	30.1	1	C
10	18	<i>Clostridium</i> (g)	96.5	2.9	16.7	4.0	0.27 ±0.05	30.0	0	A
11	11	<i>Deltaproteobacteria</i> (o)	99.2	0.7	100.0	3.4	0.63 ±0.05	57.4	2	BC
12	33	<i>Deltaproteobacteria</i> (o)	100.0	1.2	0.0	3.3	12.71 ±0.49	57.8	3	BC
13	28	<i>Firmicutes</i> (p)	99.9	0.0	0.0	2.5	1.02 ±0.10	47.2	0	B
14	27	<i>Firmicutes</i> (p)	91.9	3.3	92.3	3.1	0.84 ±0.07	49.2	1	BC
15	8	<i>Lactobacillales</i> (o)	99.6	0.0	0.0	2.7	3.30 ±0.61	36.8	0	A
16	1	<i>Lactobacillales</i> (o)	99.3	4.6	0.0	4.1	0.99 ±0.05	39.1	0	BC
17	16	<i>Selenomonadales</i> (o)	100.0	1.5	0.0	2.2	2.60 ±0.24	41.3	0	C

* (p) phylum, (o) order, or (g) genus.

§ A,B and C refer to MetaBAT 2, MaxBin 2.0 and CONCOCT respectively.

B										
MAG	Bin	Marker lineage*	Completeness (%)	Contamination (%)	Strain heterogeneity (%)	Genome size (Mbp)	Community (%)	GC content	N. of ArsM-encoding genes	Binner §
1	12	<i>Clostridiales</i> (o)	100.0	0.0	0.0	3.1	0.41 ±0.03	54.8	0	B
2	9	<i>Clostridiales</i> (o)	98.9	0.0	0.0	4.7	1.26 ±0.24	28.4	0	C
3	39	<i>Clostridiales</i> (o)	98.0	0.3	0.0	2.0	2.36 ±0.51	43.2	0	A
4	4	<i>Clostridiales</i> (o)	99.3	0.7	100.0	2.7	4.77 ±3.14	56.1	0	A
5	16	<i>Clostridiales</i> (o)	98.7	0.9	0.0	2.8	0.35 ±0.21	35.7	0	A
6	19	<i>Clostridiales</i> (o)	99.2	1.1	0.0	3.5	0.39 ±0.28	31.2	1	BC
7	1	<i>Clostridiales</i> (o)	98.7	1.3	50.0	2.6	0.25 ±0.07	56.1	1	A
8	15	<i>Clostridiales</i> (o)	97.3	2.5	16.7	2.7	0.17 ±0.02	60.5	0	C
9	28	<i>Clostridium</i> (g)	99.3	5.5	23.1	5.6	1.42 ±0.30	30.1	2	A
10	27	<i>Clostridium</i> (g)	98.6	6.9	0.0	4.6	2.10 ±0.60	32.3	1	A
11	32	<i>Deltaproteobacteria</i> (o)	94.8	0.0	0.0	3.1	0.21 ±0.06	59.3	0	BC
12	38	<i>Deltaproteobacteria</i> (o)	98.3	1.8	50.0	3.4	0.81 ±0.08	57.6	2	BC
13	10	<i>Enterobacteriaceae</i> (f)	96.6	0.7	33.3	4.3	0.39 ±0.08	52.8	0	B
14	42	<i>Enterobacteriaceae</i> (f)	95.7	2.1	12.5	5.1	6.77 ±0.35	56.3	0	BC
15	31	<i>Firmicutes</i> (p)	99.9	0.0	0.0	2.4	0.37 ±0.06	47.6	0	A
16	33	<i>Firmicutes</i> (p)	100.0	0.6	0.0	3.2	1.97 ±1.09	49.1	1	BC
17	7	<i>Lactobacillales</i> (o)	99.6	0.0	0.0	2.9	2.64 ±0.52	36.5	0	C
18	5	<i>Lactobacillales</i> (o)	98.9	4.2	0.0	4.1	1.94 ±0.81	39.1	0	AB
19	36	<i>Selenomonadales</i> (o)	100.0	1.5	0.0	2.3	0.81 ±0.11	41.1	0	A

* (p) phylum, (o) order, or (g) genus.

§ A,B and C refer to MetaBAT 2, MaxBin 2.0 and CONCOCT, respectively.