Recommendations for tissue homogenisation and extraction in DNA metabarcoding of Malaise trap samples

Vera MA Zizka^{1*}, Matthias F Geiger¹, Thomas Hörren², Ameli Kirse¹, Niklas W Noll¹, Livia Schäffler¹, Alice M Scherges¹, Martin Sorg²

¹ Leibniz Institute for the Analysis of Biodiversity Change (LIB), Zoological Research Museum Alexander Koenig (ZFMK), Centre for Biodiversity Monitoring and Conservation Science, Adenauerallee 160, 53113 Bonn, Germany

² Entomological Society Krefeld (EVK), Marktstraße 159, 47798 Krefeld, Germany

Keywords: arthropod metabarcoding, biodiversity, monitoring, natura 2000

* Corresponding author:

Vera MA Zizka

Leibniz Institute for the Analysis of Biodiversity Change (LIB), Zoological Research Museum Alexander Koenig, Adenauerallee 160, 53113 Bonn, Germany

Email: v.zizka@leibniz-zfmk.de

Phone: +49 (0) 2289122212

Abstract

With increased application of DNA metabarcoding in fast and high-resolution biodiversity assessment, various laboratory protocols have been optimised in recent years and their further evaluation is subject of current research. Homogenisation of bulk samples and subsequent DNA extraction from destructed tissue is one way of starting the metabarcoding process. This essential step in the protocol can either be conducted from wet sample material (e.g. bulk insect samples) soaked in fixative or from completely dried individuals. While the latter method appears to produce more consistent results, it is time consuming and more prone to cross-contamination. We tested both homogenisation approaches with regard to time efficiency and biodiversity assessment of complex arthropod bulk samples, in particular how the amount of processed tissue affects taxon recovery. Both approaches reveal similar taxa compositions and detect a similar total OTU diversity in a single extraction reaction. Increased amounts of tissue used in DNA extraction improved OTU diversity detection and recovered particularly specific low-biomass taxa, making this approach valuable for samples with high biomass and/or diversity. Due to less handling time and lower vulnerability for cross-contamination we recommend the processing of wet material when sample homogenisation is applied.

Introduction

For highly diverse groups as terrestrial arthropods and insects in particular, where morphological identification is difficult, slow and expensive, metabarcoding provides an efficient alternative (Bush et al., 2019; Evans et al., 2016; Morinière et al., 2019; van der Heyde et al., 2020; Yu et al., 2012). In recent years, a variety of studies evaluated and discussed promising sampling strategies (Gleason et al., 2020; Marquina et al., 2019; Pereira-da-Conceicao et al., 2020; Steinke et al., 2020), laboratory procedures (Elbrecht et al., 2019; Majaneva et al., 2018; Piñol et al., 2019; Zizka et al., 2019), bioinformatic analyses (Boyer et al., 2016; Frøslev et al., 2017; Porter and Hajibabaei, 2020; Turon et al., 2020) and ways of integration into existing biodiversity monitoring matrices (Buchner et al., 2019; Cordier et al., 2018; Mächler et al., 2020; Pawlowski et al., 2018). This resulted in a variety of different DNA metabarcoding protocols, whereas standardisation is still lacking even though it is a major prerequisite for inter-comparability and transferability of methods to applied concepts (Bush et al., 2019; McGee et al., 2019; Pawlowski et al., 2018).

Aside from eDNA (environmental DNA) metabarcoding, where free extracellular DNA is processed (e.g. from soil, water, faeces), DNA can be extracted from enclosed communities

(cDNA), more precisely, the sample's fixative ethanol (Batovska et al., 2021; Hajibabaei et al., 2012; Martins et al., 2019; Zizka et al., 2019) or propylene glycol (Martoni et al., 2021), from added lysis buffer (Giebner et al., 2020; Ji et al., 2013; Kirse et al., 2021) or from homogenised tissue of specimens (Hardulak et al., 2020; Mata et al., 2020; Zizka et al., 2020). While the latter approach is currently considered most effective to assess biodiversity pattern (Hardulak et al., 2020; Marquina et al., 2019; Persaud et al., 2021; Zenker et al., 2020; Zizka et al., 2019) it prevents subsequent morphological determinations (Nielsen et al., 2019). Homogenisation and tissue-based DNA extraction can be conducted from wet (Beentjes et al., 2019; Gibson et al., 2015; Porter et al., 2019) samples in ethanol or from dried tissue after ethanol evaporation (Elbrecht et al., 2019; Hardulak et al., 2020; Hausmann et al., 2020; Steinke et al., 2020). While powder homogenate of dried samples usually appears finer than that of wet material, handling is more prone to cross contamination and time consuming. Since most DNA extraction approaches tolerate only a limited amount of tissue per reaction, only a subsample of complete material is usually processed, ranging between 1-100 mg (Elbrecht et al., 2017; Hausmann et al., 2020; Majaneva et al., 2018; Marquina et al., 2019; Mata et al., 2020). Higher tissue volume during DNA extraction requires multiple reactions or more voluminous DNA extraction kits and is increases effort and material costs. However, DNA extraction from the subsample tissue assumes perfect homogenisation and equal distribution within storage tubes, and it remains uncertain to what extent variations in tissue composition affects the assessment of species contained in bulk samples. Insight on that question is essential to decide on how to optimise diversity detection and is thus a prerequisite for successful and reliable application of tissue-based DNA metabarcoding. While Buchner et al. 2021 analyse the overlap in detected taxa between subsamples only for wet homogenisation of Malaise traps, other studies cover aquatic samples or other trapping types with lower biomass and diversity (Beentjes et al., 2019; Elbrecht et al., 2017; Mata et al., 2020). In addition, if several subsamples are applied, taxa detection is not compared amongst them or further methodological investigations are applied (e.g. comparison of different extraction methods (Majaneva et al., 2018). The overlap between subsamples of homogenised tissue or increases in taxon recovery by use of larger quantities of homogenised tissue in Malaise trap metabarcoding has never been tested systematically but bears important information for high resolution metabarcoding of terrestrial insect biodiversity.

Here we use five time-interval Malaise trap samples collected in a protected area in Germany and investigate the effect of homogenisation strategy and tissue subsampling on biodiversity assessments. Based on our results, we formulate best-practice

recommendations for tissue-based DNA metabarcoding protocols esp. for Malaise trap samples, which ensure time and money efficiency, best quality of biodiversity assessment and also improved standardisation of DNA metabarcoding for biodiversity monitoring programs.

Material and Methods

Sampling

Samples were collected in the Nature reserve 'Latumer Bruch' near Krefeld in Western Germany. All samples originate from one Malaise trap (51.326701N, 6.632973E). Detailed information about samples taken between May and July are given in Table 1.

Tab. 1: Malaise trap samples analysed: duration: time collection bottle was installed on the Malaise trap, wet biomass: complete wet biomass (g) and biomass per day (g/day) over sampling interval

ID	duration	wet biomass (g)	wet biomass (g/day)
T 1	12.05 - 18.05.19	14.1	2.4
T 2	29.05 - 08.06.19	65.9	6.7
T 3	28.06 - 07.07.19	71.7	8
T 4	07.07 - 18.07.19	40	3.6
T 5	18.07 - 28.07.19	70.1	7

Malaise trap sampling was conducted in a standardised manner (for details see Ssymank et al., 2018). Samples were collected in 96% denatured ethanol (1% MEK). After collection, ethanol was replaced with new 96% undenatured ethanol and stored at -20°C for further processing.

Figure 1: Experimental setup. Malaise trap samples were sieved in ethanol and separated into size fraction **Large** (>4mm) and **Small** (<4mm). Wet tissue was homogenised and nine subsamples per size fraction with ~20 mg each were transferred to 1.5 µl Eppendorf tubes for separate DNA extraction. Homogenised tissue was dried and again homogenised in dry conditions. Again, nine subsamples per size fraction with ~20 mg each were transferred to 1.5 µl Eppendorf tubes. Subsequent DNA workflow was conducted as described in Material and Methods.

Laboratory work

Supernatant ethanol was removed and each sample was separated into two size classes by sieving of wet specimen through a 4 mm x 4 mm mesh with a wire diameter of 0.5 mm (untreated stainless steel). In the following, the size fractions will be referred to as either S (small, < 4mm) or L (large, > 4 mm). Depending on sample volume, individuals of both size classes were transferred to 30 ml tubes (Nalgene, wide-mouth bottle, polypropylene) or to 50 ml Falcon tubes, and approximately 20 g of grinding balls (5 mm diameter, stainless steel, Retsch GmbH) were added to each sample. Homogenisation of wet samples was conducted with a mixer mill MM 400 (Retsch GmbH) at a frequency of 30 s⁻¹ for five minutes. Nine

subsamples per sample and size class were transferred to 1.5 ml Eppendorf tubes, centrifuged for 1 minute at 10,000 rpm (Heraeus Fresco 21 Centrifuge, Thermo Scientific) and dried overnight in a shaking incubator (ILS6, VWR) at 50 °C (90 samples in total, 5 samples x 2 treatments x 9 subsamples). Subsamples were weighed (20 mg ± 6 mg) on a fine scale balance (Entris, Sartorius). Together with 6 negative controls (200 µl lysis buffer, no sample added) and DNA was extracted with the DNeasy 96 Blood and Tissue Kit (Quiagen, Hilden, Germany) following manufacturer instructions. Remaining sample tissue (in 30 ml tubes, not transferred to 1.5 Eppendorf tubes) was centrifuged for 1 minute at 4700 rpm (MegaStar 1.6, VWR Collection) and the supernatant discarded afterwards. Remaining tissue was left to dry in a shaking incubator at 50 °C for up to 3 days until complete ethanol evaporation. Dried tissue of samples containing less than 30 ml was again homogenised at a frequency of 30 s/sec for 5 minutes with the Retsch mixer mill (MM400). Samples consisting of more than 30 ml source material were homogenised for 3 minutes with a Turax mixer mill (Tube Mill 100 Control) at 25,000 rpm because dried material was clustered to a hard unit and could not be destructed with the former mill. Nine subsamples per sample were transferred to 1.5 Eppendorf tubes and weighed (23 mg ± 5 mg). To ensure processing of identical samples during this experiment, samples already homogenised under wet condition were dried and homogenised again under dry condition. Obviously, this includes an additional homogenisation step for those samples, which likely influences fineness of material. This will be picked up in the discussion. For simplicity, the former approach will hereinafter will be referred to as *wet homogenisation*, while the latter approach (wet with additional dry homogenisation) will be referred to as *dry homogenisation*. Together with 6 negative controls DNA was extracted with the DNeasy 96 Blood and Tissue Kit (Quiagen, Hilden, Germany) following manufacturer instructions for both plates. Extraction success and DNA quality was checked on a 1 % agarose gel.

A two-step PCR protocol was applied using standard Illumina Nextera primers for dual indexing of samples. The first PCR was performed with the PCR Multiplex Plus Kit (Qiagen, Hilden, Germany) using 12.5 µl master mix, 1 µl of DNA template, 0.2 µM of the fwH2 forward (Vamos et al., 2017) and Fol_degen_rev reverse (Yu et al., 2012) primers respectively. The primer pair targets a 313 bp long stretch of the COI DNA barcode region and was positively evaluated in a comparison on Malaise trap samples (Elbrecht et al., 2019). The PCR mix was filled up with 10.5 µl ddH₂O to a 25 µl reaction volume. The following PCR program was applied: initial denaturation at 95 °C for 5 min; 25 cycles of: 30 s at 95 °C, 30 s at 50 °C and 50 s at 72 °C; final extension of 5 min at 72 °C. First step PCR (PCR 1) product was used for the second PCR (PCR 2), also conducted with the PCR Multiplex Plus Kit (Qiagen, Hilden, Germany). Reaction included 1 µl DNA template from

PCR 1, 0.2 μ M of each tagging primer (Nextera, Illumina, San Diego, USA) and 12.5 μ l master mix filled up with 10.5 μ l H₂O. Primers included a nucleotide overhang as a binding site for primers in PCR 2, which was run with the same program as PCR 1 but with only 15 cycles. PCR success was evaluated on a 1 % agarose gel before PCR products were normalised using a SequalPrep Normalisation plate (Thermo Fisher Scientific, MA, USA) following the manufacturer's instructions with an end concentration of 25 ng per sample (100 μ l). Of each sample 10 μ l was pooled together and a left sided size selection was applied twice on the sample pool to remove primer residuals (ration 0.76x, SPRIselect Beckman Coulter). Library concentration was measured with a Quantus fluorometer (Promega, Madison, USA) and on a fragment analyser (Agilent Technologies, Santa Clara, CA, USA) and the pool was sent for sequencing on two Miseq runs (2 x 300 bp) to MacroGen Europe B.V., Netherlands.

Data analysis

The quality of sequences delivered by MacroGen was determined through the program Fastqc (Andrews et al., 2012). Subsequent data processing was conducted for all samples as implemented in JAMP v0.67 (<https://github.com/VascoElbrecht/JAMP>) using standard settings. Paired-end reads were merged (Edgar and Flyvbjerg, 2015) (module Merge_PE) with vsearch v2.15.0 (Rognes et al., 2016). Cutadapt 3.4 (Martin, 2017) was used to remove primers and to discard sequences of unexpected length so that only reads with a length of 303 - 323 bp were used for further analyses. The module Max_ee was used to discard all reads with an expected error >0.5 (Edgar and Flyvbjerg, 2015). Sequences were dereplicated, singletons were removed and sequences with $\geq 97\%$ similarity were clustered into Operational Taxonomic Units (OTUs) using uparse (Cluster_otus) (Edgar, 2013). OTUs with a minimal read abundance of 0.003% per sample were retained for further analysis and the program LULU was used for further qualitative filtering (Frøslev et al., 2017). Taxonomic assignment of molecular units was conducted by comparison with an Arthropoda reference database. The database was created by taxalogue (<https://github.com/nwnoll/taxalogue> commit: [62ce71819af40a6e605e9142f0ccd69318477596](https://github.com/nwnoll/taxalogue/commit/62ce71819af40a6e605e9142f0ccd69318477596)) with sequences from BOLD (<https://www.boldsystems.org>), NCBI GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>) and GBOL (<https://bolgermany.de/gbol1/ergebnisse/results>). Taxon names were normalised according to the NCBI Taxonomy (<https://www.ncbi.nlm.nih.gov/taxonomy>). Only those OTUs were kept that had at least 85% similarity with a sequence in a reference database (using vsearch version 2.14.1 (Rognes et al., 2016) with `-usearch_global` command (Edgar, 2010) and the following parameters: `-id 0.85`, `-dbmask none`, `-qmask none`, `-maxhits 1000`, `blast6out` and `-maxaccepts 0`). Custom scripts were used to extract the best hits and assign

each OTU a taxonomic name. For analysis of significant differences in OTU numbers between homogenisation methods and size fractions, a Kruskal-Wallis test was performed as integrated in the R package “dplyr” (Wickham et al., 2021). Dissimilarity indices were calculated with the package “vegan” (Oksanen et al., 2019). For species accumulation curves and associated calculations of extrapolated values the package “iNEXT” (Hsieh et al., 2016) was used. Figures were constructed with the R package ggplot (Wickham, 2016).

Results

On the two Miseq runs 13.3 and 14.7 million reads in forward and reverse direction were assigned to the specified index combinations. Raw data are uploaded at GenBank (accession number: XX). After quality filtering, on average 114,394 ($\pm 20,365$) reads were kept per sample. For both homogenisation approaches combined and all subsamples, in total 1,529 OTUs were clustered with the following order-level assignments: Coleoptera 158 (10.3%), Diptera 372 (24.3%), Hemiptera 134 (8.8%), Hymenoptera 689 (45.1%), Lepidoptera 52 (3.4%), other orders 99 (6.4%), no assignment to order level 25 (1.6%). Of those, 1088 OTUs were assigned to genus or species level. Detailed information about OTU assignments and read distribution on order level are summarised in Table 1 and Table S1. While the highest number of OTUs was assigned on average to dipterans (L: 41.6%, S: 37.5%) and hymenopterans (L: 38%, S: 32.3%), the main proportion of reads was related to dipterans (L: 60.8%, S: 77%) and < 10% to representatives of Hymenoptera (Table 1). This was most pronounced for size fraction S, where on average only 4.7% of the reads were assigned to this highly diverse order.

Table 1: Average proportion of OTUs and reads assigned per time interval and through all samples to the orders Coleoptera, Diptera, Hemiptera, Hymenoptera, Lepidoptera, other arthropod orders and OTUs not assigned to order level (see table S1) for both homogenization approaches and all subsamples combined

Order	Size	OTU number [%]	Read number [%]	Total OTU number
Coleoptera	L	5.7 $\pm$ 1.9	6 $\pm$ 4.9	158 (10.3%)
	S	11.1 $\pm$ 1	5.5 $\pm$ 2.8	
Diptera	L	41.6 $\pm$ 9.8	60.8 $\pm$ 13.3	372 (24.4%)
	S	37.5 $\pm$ 8.8	77 $\pm$ 11.9	
Hemiptera	L	3 $\pm$ 0.8	0.4 $\pm$ 0.3	134 (8.8%)
	S	11.2 $\pm$ 2.3	10.1 $\pm$ 10.7	

Hymenoptera	L	38 ± 5.2	18.7 ± 7.9	689 (45.1%)
	S	32.3 ± 7.5	4.7 ± 2.5	
Lepidoptera	L	7.6 ± 3.2	13.8 ± 5.4	52 (3.4%)
	S	2.9 ± 1.3	1.8 ± 1	
Other arthropods	L	2.8 ± 0.8	0.3 ± 0.2	98 (6.4%)
	S	3.6 ± 0.6	0.2 ± 0.1	
No assignment	L	1.2 ± 0.4	0.1 ± 0.2	25 (1.64%)
	S	1.3 ± 0.3	0.7 ± 0.5	

While different emptying dates and the different size classes per sample showed distinct community compositions (Fig. 2, $p < 0.002$), the condition during homogenisation (only wet or wet and additional dry homogenisation) had no effect on sample ordination in NMDS analysis ($p = 0.997$, Fig. 2). However, average Jaccard dissimilarity between subsamples homogenised in wet and additional dry condition (0.179 ± 0.06) was lower ($p < 0.001$) than dissimilarities between subsamples homogenised in only wet condition (0.207 ± 0.081) and when subsamples of one emptying date and size fraction were compared amongst homogenisation approaches (0.206 ± 0.089 , $p < 0.001$). Bray-Curtis dissimilarity was on average lower within wet and dry (0.11 ± 0.07) compared to only wet (0.11 ± 0.09) homogenised subsamples than comparing subsamples between methods (0.13 ± 0.08 , $p < 0.01$).

Figure 2: Non-metric multidimensional scaling based on A) Jaccard (presence/absence data) and B) Bray-Curtis (abundance data – regarding read numbers) dissimilarity matrices. Samples include the nine subsamples per emptying date (colour coding), size fraction (S and L, marked in figure) and condition during homogenisation (shape coding).

For samples processed under wet condition, on average $60.7\% \pm 8.5$ of calculated total diversity could be detected in size fraction S and $75.8\% \pm 7.9$ in size fraction L when only one subsample was processed in extraction (~ 20 mg of tissue, Fig. 3). With nine extraction subsamples ($9 \times \sim 20$ mg) $88.4\% \pm 3.2$ and $92.4\% \pm 4.2$ of calculated total richness were detected in fraction S and L respectively, while 95% of calculated species richness was assessed with 12 ± 5.4 and 17 ± 3.4 subsamples. For samples homogenised in wet and additional dry condition, a single extraction (~ 20 mg) of size class S revealed $64\% \pm 2.3$ of calculated species richness, while $79\% \pm 3.8$ could be detected in 20 mg of size class L (Fig. 3). On average $88.9\% \pm 1.6$ of total diversity was assessed with the nine applied subsamples for size fraction S and $93.1\% \pm 3.8$ for size fraction L, while 95% of total was calculated with 16 ± 2.1 (S, 320 mg) and 13 ± 7.1 (L, 260 mg) subsamples (~ 320 mg and ~ 260 mg, Fig. 3).

Detailed analysis of Hymenoptera in only wet homogenised tissue revealed on average $47.2\% \pm 13.3$ of extrapolated total species richness in size fraction S and $80.4\% \pm 10.4$ in size fraction L when a single subsample was processed (Fig.3). Increased to nine subsamples, $81.4\% \pm 7.6$ (S) and $96.2\% \pm 4.7$ (L) of calculated total diversity were detected. Extrapolations revealed, that 95% of total calculated diversity was achieved by processing 22 ± 5.2 (S, ~ 440 mg) and 7.4 ± 6 (L, ~ 140 mg) replicates. For wet and additional dry homogenisation, one extraction subsample revealed $54\% \pm 2.7$ (S) and $82.4\% \pm 9.6$ (L) of total species richness, while $86.2\% \pm 3.1$ and $95.1\% \pm 4.8$ could be assessed with nine extraction subsamples. Calculations revealed a detection of 95% from total hymenopteran species richness if 18 ± 4 (360 mg) and 9.8 ± 8 (200 mg) replicates were processed (Fig. 3).

For wet homogenisation, in detail analysis of dipteran representatives revealed $70.6\% \pm 5.6$ of calculated diversity in size fraction S and $77\% \pm 10.3$ in size fraction L, when only a single extraction subsample was processed. Detected richness increased to $92.1\% \pm 4.7$ and $91\% \pm 5.6$ when nine subsamples were processed (95% were reached with 14.2 ± 7.9 and 5 ± 1.9 subsamples). With additional dry homogenisation processing of one tissue subsample revealed $70.6\% \pm 2.7$ of extrapolated diversity for size fraction S and 81.1 ± 5.9 for size fraction L. With nine extraction subsamples, taxa detection increased to $91.2\% \pm 3.4$ (S) and $93.3\% \pm 5.5$ (L). Calculations revealed a detection of 95% from total Diptera if 15.6 ± 5.4

(~320 mg) and 14.2 ± 10.2 (~200 mg) replicates were processed. For detailed information about observed and calculated species richness see Figure 3.

Figure 3: Species accumulation curves. Increased number of OTUs with increased amount of processed tissue in extraction is illustrated as well as extrapolation up to 40 subsamples. All taxa homogenized in A) dry and B) wet condition; Diptera recovered in samples homogenised in C) dry and D) wet condition; Hymenoptera detected in samples homogenised in E) dry and F) wet condition. Size fraction is marked in the upper left corner of each subfigure (S = small, L = large).

296

297

298 **Figure 4:** Proportion of molecular units (OTUs) detected with additional DNA extraction
 299 subsamples. A) for samples of size fraction L (> 4mm) and B) for size fraction S (< 4 mm).
 300 The x-axis describes differences between processed subsamples (e.g. r1-r2 difference in
 301 relative number of OTUs detected with one subsamples (~20 mg) compared to two
 302 subsamples (~40 mg)).

303

304

305 Discussion

306 Homogenisation of bulk samples and subsequent DNA extraction from destructed tissue is
 307 widely applied when insect biodiversity is assessed through DNA metabarcoding (Beermann
 308 et al., 2021; Hardulak et al., 2020; Mata et al., 2020). It is up to now more efficient for bulk
 309 sample analysis than other extraction methods (Marquina et al., 2019; Persaud et al., 2021;
 310 Zenker et al., 2020). Here we set out to test different homogenisation protocols and how
 311 subsampling of homogenised tissue affects diversity estimates of highly diverse Malaise trap
 312 samples.

313

314 Comparison of different homogenisation approaches

315 The average dissimilarity between reactions homogenised under dry conditions was lower
 316 than dissimilarity between reactions processed under wet condition for presence/absence
 317 analysis, which indicates a higher homogeneity of tissue samples processed under dry
 318 condition. To ensure comparability and processing of identical sampling material (as stated

in Material and Methods section line 150-156), the dry homogenisation approach was based on material already destructured under wet condition. This additional step of 3 min homogenisation most probably influenced refinement of samples and similarity of subsamples. In addition, on average a lower tissue weight was processed subsequent to wet homogenisation due to a weight decrease after drying (difference 3 mg). However, Jaccard's dissimilarity indices of subsamples which were homogenised in wet condition was on average 0.21 ± 0.08 , mainly due to high inconsistencies between subsamples from June 8th with an average dissimilarity of 0.35 ± 0.02 compared to dissimilarities between subsamples of the other collection dates (0.18 ± 0.06 ; Fig. 2). The homogenisation of dried samples implements drying for approximately 48 h at temperatures around 50 °C to guarantee the complete evaporation of ethanol from the sample. In addition, the fine powder resulting from dry homogenisation is electrostatically charged and thus bears a high risk of cross contamination between samples and general lab contamination, and further increases the time for sample handling (Buchner et al., 2021; Elbrecht and Steinke, 2019). In comparison, the homogenisation of wet samples soaked in ethanol circumvents the drying step and is therefore more time efficient and suitable for large scale approaches as implemented in several studies on aquatic samples (Hajibabaei et al., 2019; Majaneva et al., 2018; Pereira-da-Conceicao et al., 2020). In addition, the handling of homogenised tissue in ethanol is simplified and reduces contamination risk (Elbrecht and Steinke, 2019). The minor differences we observed between the two applied methods and the above-mentioned experimental setup allows to recommend homogenisation of wet material for tissue-based DNA metabarcoding Malaise trap samples as it reduces processing time and contamination risk. Additionally, results indicate, that an additional homogenisation step of dried material as e.g. through bead-grinding should be integrated after material has been subsampled to increase the fineness of material as also conducted in Buchner et al., 2021.

Amount of tissue needed for homogenisation

A high, but incomplete proportion of the total insect diversity was assessed when processing a single tissue subsample in extraction, which opens up an additional perspective to previous studies highlighting wet homogenisation of samples (Buchner et al., 2021). This accounts for both tested variables, basis material for homogenisation (wet or dry) and size fractions (Fig. 3). For all samples at least 60% of calculated total arthropod diversity were detected with a single extraction from ~20 mg of tissue (Fig. 3 A+B), revealing strong alterations between size fractions time intervals and higher insect taxa. The processing of additional subsamples only moderately increase diversity estimates for size fraction L and samples of both size fractions from samples collected in May. While those samples constitute a comparatively low biodiversity, a strong increase was detected for remaining

samples of the small size fractions and collected in summer. Results indicate that extraction should be conducted from more tissue for highly diverse samples, either through increasing number of extraction subsamples, as it was also recommended in (Elbrecht and Steinke, 2019) or a higher tissue volume per DNA extraction. Additional rare material for extraction however comes with higher expenses but can increase taxa detection by 25-30% as an average over all taxa. Referring to the 3000 BINs detected within a year-long Malaise trap collection through single-specimen barcoding in Geiger et al., 2016 this can result in 600-900 additional molecular units for year-long application of a malaise trap.

The most pronounced increase of OTUs with additional extraction subsamples was observed for representatives of the order Hymenoptera, which is almost twice as high as for Diptera (Fig. 3, Tab. 2). Representatives of these two orders are the main targets in many Malaise trapping studies (Ssymank et al., 2018). However, as also indicated with previous studies, dipterans are present in much higher individual numbers, constituting a higher proportion of biomass in Malaise trap catches in Germany, while diversity of both groups are considered to be similar (Geiger et al., 2016). An underrepresentation of specific insect families, especially constituting taxa of low-biomass (Elbrecht et al., 2021) has been reported from previous metabarcoding studies (Elbrecht and Steinke, 2019; Krehenwinkel et al., 2017; Yu et al., 2012). This accounts e.g. for highly diverse parasitoid hymenopterans, which depict important ecosystem functions. While insufficient primer binding efficiency is discussed as the main reason for this phenomenon, our results indicate, that the extraction from an insufficient amount of raw tissue material could also bias detected diversity pattern. It also assumes that taxon biomass in complex bulk samples is affecting detection probability if a limited amount of tissue is processed in extraction but can be increased through insertion of higher tissue amounts. Again, this accounts especially for highly diverse samples as indicated here through the small size fraction. We demonstrate, that the use of higher amount of tissue of approx. 180 mg over nine subsamples increase the detection of low biomass taxa, even if general sequence coverage remains low (5% of reads assigned to Hymenoptera, Table 1). Additionally, the application of higher sequencing depth can increase taxa recovery and overlap between extraction subsamples. We here only tested a sequencing depth of on average $114,394 \pm 20,365$ reads per sample (after quality filtering) and detailed analysis to understand the linkage between higher sequencing depth and replication strategy is out of the scope of the present study. Further investigation could reveal an increase in sequencing depth as the most effective way to optimise taxon recovery under financial constraints.

Conclusion

We recommend homogenisation of wet material in tissue-based DNA metabarcoding of Malaise trap samples due to similar levels of taxon recovery from dry and wet tissue homogenisation combined with a lower time effort and contamination susceptibility of the latter approach. In both, dry and wet homogenisation, additional DNA extractions from more tissue results in a higher number of detected taxa, in particular those of low biomass. The amount of processed tissue and number of subsamples affects the resolution taxonomically. A decision on the more complete diversity detection associated with higher resources depends on the focused scientific goals of individual studies.

Data Accessibility

The raw sequence used in this study is available at NCBI SRA: XX

Competing Interests

The authors declare no competing interests

Acknowledgements

This work was supported by the Ministry for Environment, Agriculture, Conservation and Consumer Protection of the German State of North Rhine-Westphalia (No. III-1-620.08)

References

- Andrews, S., Krueger, F., Segonds-Pichon, A., Biggins, L., Krueger, C., Wingett, S., 2012. FastQC. Babraham, UK.
- Andújar, C., Creedy, T.J., Arribas, P., López, H., Salces-Castellano, A., Pérez-Delgado, A., Vogler, A.P., Emerson, B.C., 2020. Validated removal of nuclear pseudogenes and sequencing artefacts from mitochondrial metabarcode data. *bioRxiv* 2020.06.17.157347. <https://doi.org/10.1101/2020.06.17.157347>
- Arribas, P., Andújar, C., Bidartondo, M.I., Bohmann, K., Coissac, É., Creer, S., deWaard, J.R., Elbrecht, V., Ficetola, G.F., Goberna, M., Kennedy, S., Krehenwinkel, H., Leese, F., Novotny, V., Ronquist, F., Yu, D.W., Zinger, L., Creedy, T.J., Meramveliotakis, E., Noguerales, V., Overcast, I., Morlon, H., Vogler, A.P., Papadopoulou, A., Emerson, B.C., 2021. Connecting high-throughput biodiversity inventories: Opportunities for a site-based genomic framework for global integration and synthesis. *Mol. Ecol.* 30, 1120–1135. <https://doi.org/10.1111/mec.15797>
- Batovska, J., Lynch, S.E., Cogan, N.O.I., Brown, K., Darbro, J.M., Kho, E.A., Blacket, M.J., 2018. Effective mosquito and arbovirus surveillance using metabarcoding. *Mol. Ecol. Resour.* 18, 32–40. <https://doi.org/10.1111/1755-0998.12682>
- Batovska, J., Piper, A., Valenzuela, 2021. Developing a Non-destructive Metabarcoding Protocol for Detection of Pest Insects in Bulk Trap Catches.
- Beentjes, K.K., Speksnijder, A.G.C.L., Schilthuizen, M., Hoogeveen, M., Pastoor, R., van der Hoorn, B.B., 2019. Increased performance of DNA metabarcoding of

- macroinvertebrates by taxonomic sorting. PLOS ONE 14, e0226527.
<https://doi.org/10.1371/journal.pone.0226527>
- Boyer, F., Mercier, C., Bonin, A., Le Bras, Y., Taberlet, P., Coissac, E., 2016. obitools: a
unix-inspired software package for DNA metabarcoding. Mol. Ecol. Resour. 16, 176–
182. <https://doi.org/10.1111/1755-0998.12428>
- Buchner, D., Beermann, A.J., Laini, A., Rolaufts, P., Vitecek, S., Hering, D., Leese, F., 2019.
Analysis of 13,312 benthic invertebrate samples from German streams reveals minor
deviations in ecological status class between abundance and presence/absence
data. PLOS ONE 14, e0226547. <https://doi.org/10.1371/journal.pone.0226547>
- Buchner, D., Leese, F., 2020. BOLDigger – a Python package to identify and organise
sequences with the Barcode of Life Data systems. Metabarcoding Metagenomics 4,
e53535. <https://doi.org/10.3897/mbmg.4.53535>
- Bush, A., Compson, Z.G., Monk, W.A., Porter, T.M., Steeves, R., Emilson, E., Gagne, N.,
Hajibabaei, M., Roy, M., Baird, D.J., 2019. Studying Ecosystems With DNA
Metabarcoding: Lessons From Biomonitoring of Aquatic Macroinvertebrates. Front.
Ecol. Evol. 7, 434. <https://doi.org/10.3389/fevo.2019.00434>
- Bush, A., Monk, W.A., Compson, Z.G., Peters, D.L., Porter, T.M., Shokralla, S., Wright,
M.T.G., Hajibabaei, M., Baird, D.J., 2020. DNA metabarcoding reveals
metacommunity dynamics in a threatened boreal wetland wilderness. Proc. Natl.
Acad. Sci. 117, 8539. <https://doi.org/10.1073/pnas.1918741117>
- Cordier, T., Forster, D., Dufresne, Y., Martins, C.I.M., Stoeck, T., Pawlowski, J., 2018.
Supervised machine learning outperforms taxonomy-based environmental DNA
metabarcoding applied to biomonitoring. Mol. Ecol. Resour. 18, 1381–1391.
<https://doi.org/10.1111/1755-0998.12926>
- Elbrecht, V., Braukmann, T.W.A., Ivanova, N.V., Prosser, S.W.J., Hajibabaei, M., Wright, M.,
Zakharov, E.V., Hebert, P.D.N., Steinke, D., 2019. Validation of COI metabarcoding
primers for terrestrial arthropods. PeerJ 7, e7745. <https://doi.org/10.7717/peerj.7745>
- Elbrecht, V., Vamos, E.E., Meissner, K., Aroviita, J., Leese, F., 2017. Assessing strengths
and weaknesses of DNA metabarcoding-based macroinvertebrate identification for
routine stream monitoring. Methods Ecol. Evol. 8, 1265–1275.
<https://doi.org/10.1111/2041-210X.12789>
- Evans, D.M., Kitson, J.J.N., Lunt, D.H., Straw, N.A., Pocock, M.J.O., 2016. Merging DNA
metabarcoding and ecological network analysis to understand and build resilient
terrestrial ecosystems. Funct. Ecol. 30, 1904–1916. <https://doi.org/10.1111/1365-2435.12659>
- Frøslev, T.G., Kjølner, R., Bruun, H.H., Ejrnæs, R., Brunbjerg, A.K., Pietroni, C., Hansen,
A.J., 2017. Algorithm for post-clustering curation of DNA amplicon data yields reliable
biodiversity estimates. Nat. Commun. 8, 1188. <https://doi.org/10.1038/s41467-017-01312-x>
- Giebner, H., Langen, K., Bourlat, S.J., Kukowka, S., Mayer, C., Astrin, J.J., Misof, B.,
Fonseca, V.G., 2020. Comparing diversity levels in environmental samples: DNA
sequence capture and metabarcoding approaches using 18S and COI genes. Mol.
Ecol. Resour. 20, 1333–1345. <https://doi.org/10.1111/1755-0998.13201>
- Geiger, M. F., Moriniere, J., Hausmann, A., Haszprunar, G., Wägele, W., Hebert, P. D., &
Rulik, B. (2016). Testing the Global Malaise Trap Program—How well does the
current barcode reference library identify flying insects in Germany?. Biodiversity
data journal, (4).
- Gleason, J.E., Elbrecht, V., Braukmann, T.W.A., Hanner, R.H., Cottenie, K., 2020.
Assessment of stream macroinvertebrate communities with eDNA is not congruent
with tissue-based metabarcoding. Mol. Ecol. n/a. <https://doi.org/10.1111/mec.15597>
- Hajibabaei, M., Spall, J.L., Shokralla, S., van Konyenburg, S., 2012. Assessing biodiversity
of a freshwater benthic macroinvertebrate community through non-destructive
environmental barcoding of DNA from preservative ethanol. BMC Ecol. 12, 28.
<https://doi.org/10.1186/1472-6785-12-28>

- Hardulak, L.A., Moriniere, J., Hausmann, A., Hendrich, L., Schmidt, S., Doczkal, D., Müller, J., Hebert, P.D., Haszprunar, G., 2020. DNA metabarcoding for biodiversity monitoring in a national park: Screening for invasive and pest species. *Mol. Ecol. Resour.* 1–16. <https://doi.org/10.1111/1755-0998.13212>
- Hausmann, A., Segerer, A.H., Greifenstein, T., Knubben, J., Morinière, J., Bozicevic, V., Doczkal, D., Günter, A., Ulrich, W., Habel, J.C., 2020. Toward a standardized quantitative and qualitative insect monitoring scheme. *Ecol. Evol.* 10, 4009–4020. <https://doi.org/10.1002/ece3.6166>
- Ji, Y., Ashton, L., Pedley, S.M., Edwards, D.P., Tang, Y., Nakamura, A., Kitching, R., Dolman, P.M., Woodcock, P., Edwards, F.A., Larsen, T.H., Hsu, W.W., Benedick, S., Hamer, K.C., Wilcove, D.S., Bruce, C., Wang, X., Levi, T., Lott, M., Emerson, B.C., Yu, D.W., 2013. Reliable, verifiable and efficient monitoring of biodiversity via metabarcoding. *Ecol. Lett.* 16, 1245–1257. <https://doi.org/10.1111/ele.12162>
- Kirse, A., Bourlat, S.J., Lange, K., Fonseca, V.G., n.d. Arthropod diversity assessment from the ground up: Seasonal overlap between soil and Malaise traps. *Rev.*
- Mächler, E., Walser, J.-C., Altermatt, F., 2020. Decision-making and best practices for taxonomy-free environmental DNA metabarcoding in biomonitoring using Hill numbers. *Mol. Ecol.* n/a. <https://doi.org/10.1111/mec.15725>
- Majaneva, M., Diserud, O.H., Eagle, S.H.C., Hajibabaei, M., Ekrem, T., 2018. Choice of DNA extraction method affects DNA metabarcoding of unsorted invertebrate bulk samples. *Metabarcoding Metagenomics* 2, e26664. <https://doi.org/10.3897/mbmg.2.26664>
- Marquina, D., Esparza-Salas, R., Roslin, T., Ronquist, F., 2019. Establishing arthropod community composition using metabarcoding: Surprising inconsistencies between soil samples and preservative ethanol and homogenate from Malaise trap catches. *Mol. Ecol. Resour.* 19, 1516–1530. <https://doi.org/10.1111/1755-0998.13071>
- Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnetjournal* Vol 17 No 1 Gener. Seq. Data Anal. <https://doi.org/10.14806/ej.17.1.200>
- Martins, F.M.S., Galhardo, M., Filipe, A.F., Teixeira, A., Pinheiro, P., Paupério, J., Alves, P.C., Beja, P., 2019. Have the cake and eat it: Optimizing nondestructive DNA metabarcoding of macroinvertebrate samples for freshwater biomonitoring. *Mol. Ecol. Resour.* 19, 863–876. <https://doi.org/10.1111/1755-0998.13012>
- Martoni, F., Nogarotto, E., Piper, A.M., Mann, R., Valenzuela, I., Eow, L., Rako, L., Rodoni, B.C., Blacket, M.J., 2021. Propylene Glycol and Non-Destructive DNA Extractions Enable Preservation and Isolation of Insect and Hosted Bacterial DNA. *Agriculture* 11. <https://doi.org/10.3390/agriculture11010077>
- Mata, V.A., Ferreira, S., Campos, R.M., da Silva, L.P., Veríssimo, J., Corley, M.F.V., Beja, P., 2020. Efficient assessment of nocturnal flying insect communities by combining automatic light traps and DNA metabarcoding. *Environ. DNA* n/a. <https://doi.org/10.1002/edn3.125>
- McGee, K.M., Robinson, C.V., Hajibabaei, M., 2019. Gaps in DNA-Based Biomonitoring Across the Globe. *Front. Ecol. Evol.* 7, 337. <https://doi.org/10.3389/fevo.2019.00337>
- Morinière, J., Balke, M., Doczkal, D., Geiger, M.F., Hardulak, L.A., Haszprunar, G., Hausmann, A., Hendrich, L., Regalado, L., Rulik, B., Schmidt, S., Wägele, J.-W., Hebert, P.D.N., 2019. A DNA barcode library for 5,200 German flies and midges (Insecta: Diptera) and its implications for metabarcoding-based biomonitoring. *Mol. Ecol. Resour.* 19, 900–928. <https://doi.org/10.1111/1755-0998.13022>
- Oksanen, J., Blanchet, F.G., Friendl, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2019. *Vegan: community ecology package.*
- Pawlowski, J., Kelly-Quinn, M., Altermatt, F., Apothéoz-Perret-Gentil, L., Beja, P., Boggero, A., Borja, A., Bouchez, A., Cordier, T., Domaizon, I., Feio, M.J., Filipe, A.F., Fornaroli, R., Graf, W., Herder, J., van der Hoorn, B., Iwan Jones, J., Sagova-Mareckova, M., Moritz, C., Barquín, J., Piggott, J.J., Pinna, M., Rimet, F., Rinkevich, B., Sousa-Santos, C., Specchia, V., Trobajo, R., Vasselon, V., Vitecek, S.,

- Zimmerman, J., Weigand, A., Leese, F., Kahlert, M., 2018. The future of biotic indices in the ecogenomic era: Integrating (e)DNA metabarcoding in biological assessment of aquatic ecosystems. *Sci. Total Environ.* 637–638, 1295–1310. <https://doi.org/10.1016/j.scitotenv.2018.05.002>
- Pedro, P.M., Rodrigues de Sá, I.L., Rojas, M.V.R., Amorim, J.A., Ribeiro Galardo, A.K., Santos Neto, N.F., Furtado, N.V.R., Pires de Carvalho, D., Nabas Ribeiro, K.A., de Paiva, M., Pepe Razzolini, M.T., Sallum, M.A.M., 2020. Efficient Monitoring of Adult and Immature Mosquitoes Through Metabarcoding of Bulk Samples: A Case Study for Non-Model Culicids With Unique Ecologies. *J. Med. Entomol.* <https://doi.org/10.1093/jme/tjaa267>
- Pereira-da-Conceicao, L., Elbrecht, V., Hall, A., Briscoe, A., Barber-James, H., Price, B., 2020. Metabarcoding unsorted kick-samples facilitates macroinvertebrate-based biomonitoring with increased taxonomic resolution, while outperforming environmental DNA. *Environ. DNA* n/a. <https://doi.org/10.1002/edn3.116>
- Persaud, S.F., Cottenie, K., Gleason, J.E., 2021. Ethanol eDNA Reveals Unique Community Composition of Aquatic Macroinvertebrates Compared to Bulk Tissue Metabarcoding in a Biomonitoring Sampling Scheme. *Diversity* 13, 34.
- Piñol, J., Senar, M.A., Symondson, W.O.C., 2019. The choice of universal primers and the characteristics of the species mixture determine when DNA metabarcoding can be quantitative. *Mol. Ecol.* 28, 407–419. <https://doi.org/10.1111/mec.14776>
- Porter, T.M., Hajibabaei, M., 2020. Putting COI Metabarcoding in Context: The Utility of Exact Sequence Variants (ESVs) in Biodiversity Analysis. *Front. Ecol. Evol.* 8, 248. <https://doi.org/10.3389/fevo.2020.00248>
- Porter, T.M., Morris, D.M., Basiliko, N., Hajibabaei, M., Doucet, D., Bowman, S., Emilson, E.J.S., Emilson, C.E., Chartrand, D., Wainio-Keizer, K., Séguin, A., Venier, L., 2019. Variations in terrestrial arthropod DNA metabarcoding methods recovers robust beta diversity but variable richness and site indicators. *Sci. Rep.* 9, 18218. <https://doi.org/10.1038/s41598-019-54532-0>
- R Core Team, 2017. R: A language and environment for statistical computing.
- Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: a versatile open source tool for metagenomics. *PeerJ* 4, e2584. <https://doi.org/10.7717/peerj.2584>
- Ssymank, A., Sorg, M., Doczkal, D., Rulik, B., Merkel-Wallner, G., Vischer-Leopold, M., 2018. Praktische Hinweise und Empfehlungen zur Anwendung von Malaisefallen für Insekten in der Biodiversitätserfassung und im Monitoring. *Ser. Nat.* 1, 1–12.
- Steinke, D., Braukmann, T., Manerus, L., Woodhouse, A., Elbrecht, V., 2020. Effects of Malaise trap spacing on species richness and composition of terrestrial arthropod bulk samples. *bioRxiv* 2020.09.30.321430. <https://doi.org/10.1101/2020.09.30.321430>
- Turon, X., Antich, A., Palacín, C., Præbel, K., Wangenstein, O.S., 2020. From metabarcoding to metaphylogeography: separating the wheat from the chaff. *Ecol. Appl.* 30, e02036. <https://doi.org/10.1002/eap.2036>
- Vamos, E.E., Elbrecht, V., Leese, F., 2017. Short COI markers for freshwater macroinvertebrate metabarcoding. *Metabarcoding Metagenomics* 1, e14625. <https://doi.org/10.3897/mbmg.1.14625>
- van der Heyde, M., Bunce, M., Wardell-Johnson, G., Fernandes, K., White, N.E., Nevill, P., 2020. Testing multiple substrates for terrestrial biodiversity monitoring using environmental DNA metabarcoding. *Mol. Ecol. Resour.* 20, 732–745. <https://doi.org/10.1111/1755-0998.13148>
- Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.
- Wickham, H., Francois, R., Henry, L., Müller, K., 2021. dplyr: A Grammar of Data Manipulation.
- Yu, D.W., Ji, Y., Emerson, B.C., Wang, X., Ye, C., Yang, C., Ding, Z., 2012. Biodiversity soup: metabarcoding of arthropods for rapid biodiversity assessment and biomonitoring. *Methods Ecol. Evol.* 3, 613–623. <https://doi.org/10.1111/j.2041-210X.2012.00198.x>

- 598 Zenker, M.M., Specht, A., Fonseca, V.G., 2020. Assessing insect biodiversity with automatic
599 light traps in Brazil: Pearls and pitfalls of metabarcoding samples in preservative
600 ethanol. *Ecol. Evol.* 10, 2352–2366. <https://doi.org/10.1002/ece3.6042>
601 Zizka, V.M., Leese, F., Peinert, B., Geiger Matthias F, 2019. DNA metabarcoding from
602 sample fixative as a quick and voucher-preserving biodiversity assessment method.
603 *Genome* 62, 122–136. <https://doi.org/doi:10.1139/gen-2018-0048>
604 Zizka, V.M.A., Elbrecht, V., Macher, J.-N., Leese, F., 2019. Assessing the influence of
605 sample tagging and library preparation on DNA metabarcoding. *Mol. Ecol. Resour.*
606 19, 893–899. <https://doi.org/10.1111/1755-0998.13018>
607 Zizka, V.M.A., Geiger, M.F., Leese, F., 2020. DNA metabarcoding of stream invertebrates
608 reveals spatio-temporal variation but consistent status class assessments in a natural
609 and urban river. *Ecol. Indic.* 115, 106383.
610 <https://doi.org/10.1016/j.ecolind.2020.106383>
611

612

613