1 **Title page**

2

3 **SHORT TITLE:** Coexistence dynamics in squamates

4

5 **GEOGRAPHICAL AND ECOLOGICAL DRIVERS OF COEXISTENCE DYNAMICS IN**

6 **SQUAMATE REPTILES**

7

8 LAURA R. V. ALENCAR^{1,2*} & TIAGO B. QUENTAL¹

9

10 ¹Departamento de Ecologia, Instituto de Biociências, Universidade de São Paulo, São
11 Paulo, SP, 05508-900, Brazil

12 ²Current address: Department of Ecology & Evolutionary Biology, Yale University,
13 New Haven, CT, 06511, USA

14 *Corresponding Author

15

16 **ACKNOWLEDGEMENTS**

17 We thank all the curators and staff from the scientific collections visited: K. De
18 Queiroz, R. Wilson and A. Wynn (SNMNH), L. Scheinberg (CAS), F. Burbrink and D.
19 Kizirian (AMNH), H. Zaher and A. Benetti (MZUSP), D. Rabosky and G. Schneider
20 (UMMZ), J. Losos and J. Rosado (MCZ), J. McGuire and C. Spencer (MVZ), P.
21 Doughty (WAM), J. Rowley, S. Mahony, C. Portway, T. Cutajar (AM). We also thank
22 Lab MeMe members, especially G. Burin and D. Caetano, for their suggestions during
23 the development of this study. We thank A. Pigot for kindly providing R codes and J.R.
24 Hodge for proofreading. LRVA and TBQ thank Sao Paulo Research Foundation
25 (FAPESP) for grants #2016/14292-1 and 2018/05462-6. This study also received a

26 Visiting Collections Fellowship awarded by the Australian Museum Research Institute
27 to LRVA.

28

29 **CONFLICT OF INTEREST STATEMENT**

30 The authors declare no competing of interests.

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52 **RUNNING TITLE:** Coexistence dynamics in squamates

53

54 **GEOGRAPHICAL AND ECOLOGICAL DRIVERS OF COEXISTENCE DYNAMICS IN**

55 **SQUAMATE REPTILES**

56

57 **ABSTRACT**

58 **Aim:** Species richness varies widely across space. To understand the processes behind
59 these striking patterns, we must know what are the relevant drivers underlying species
60 coexistence. Several factors can potentially shape species coexistence such as the
61 speciation process, the time since divergence between lineages, environmental effects,
62 and intrinsic properties of the organisms. For the first time, we model the coexistence
63 dynamics of lizards and snakes across broad temporal and spatial scales, investigating
64 the role of species interactions, dispersal ability, and geographic area.

65 **Location:** Global

66 **Time period:** Last 20 million years

67 **Major taxa studied:** Squamata (lizards and snakes)

68 **Methods:** We used 448 closely related species pairs and their age since divergence
69 across 100 dated phylogenies. We categorized each pair as sympatric or allopatric and
70 as occurring on islands or continents. We measured morphological traits to quantify
71 niche divergence and used range and body size as proxies for dispersal ability. We
72 applied a model-comparison framework in lizards and snakes separately to evaluate
73 which factors best explained their coexistence dynamics.

74 **Results:** We found that distinct factors drive the coexistence dynamics in lizards and
75 snakes. In snakes, species pairs that coexist tend to occur on islands and are more
76 different in body size, suggesting that both geographical setting and species interactions

77 might be relevant factors. In contrast, we only found evidence that dispersal ability
78 shaped the coexistence of lizards, where species coexist when they have higher
79 dispersal abilities.

80 **Main conclusions:** Lizards and snakes greatly differ in coexistence dynamics. Higher
81 heterogeneity in coexistence dynamics within lizards and group-specific life-history
82 aspects might help to explain these findings. Our results emphasize that the interaction
83 between where organisms are and who they are, ultimately shapes biodiversity patterns.
84 We also highlight interesting avenues for further studies on species coexistence in deep
85 time.

86

87 **KEY WORDS** dispersal ability, competition, niche divergence, snakes, lizards, islands,
88 continents

89

90

91

92

93

94

95

96

97

98

99

100

101

102 INTRODUCTION

103

104 Biodiversity is heterogeneously distributed across the Earth, and to explain why
105 certain regions comprise more species than others has been one of the major challenges
106 in Ecology and Biogeography (Gaston, 2000; Ricklefs, 2004). At global and regional
107 scales, the number of species is determined by speciation, extinction and migration
108 (Wiens & Donoghue, 2004; Ricklefs, 2006). However, to fully understand such spatial
109 variation in species richness we also need to know the mechanisms that ultimately
110 promote species coexistence and generate the observed species richness patterns at finer
111 spatial scales (Weber & Strauss, 2016; Pigot et al., 2018). How and why the distribution
112 of species changes across space and time has been the focus of intense debate in the
113 literature (e.g. Jackson & Overpeck, 2000; Sexton et al., 2009; Louthan et al., 2015).

114 Several factors can potentially shape the distribution of organisms and
115 consequently shape species coexistence (Hutchinson, 1957; Vamosi et al., 2009; Wisz
116 et al., 2013). Distinct speciation modes, for example, are linked to distinct predictions
117 regarding the patterns of coexistence between two closely related species. Sympatric
118 speciation assumes that two species will coexist since their emergence (Grossenbacher
119 et al., 2014), whereas allopatric speciation will produce geographically isolated species
120 (Kozak & Wiens, 2006) that can eventually expand their distributions and subsequently
121 coexist (Pigot & Tobias, 2015; Weber & Strauss, 2016; Pigot et al., 2018). In this case,
122 the time since divergence between lineages can play an important role in determining
123 coexistence patterns.

124 Biotic interactions and the intrinsic properties of organisms, such as their
125 dispersal ability, can also affect the process of geographic expansion, either accelerating
126 or preventing species coexistence over time (Johnson & Stinchcombe, 2007; Vamosi et

127 al., 2009; Lowe & McPeck, 2014; Jönsson et al., 2016; Weber & Strauss, 2016). From
 128 an ecological perspective, the importance of competition in shaping coexistence is
 129 based on the “principle of competitive exclusion”, which suggests that species
 130 occupying similar ecological niches would not be able to coexist because they require
 131 the same limited resources to survive (Hutchinson, 1957; Hardin, 1960). The outcome
 132 of these ecological processes on an evolutionary time-scale could be character
 133 displacement among populations or a pair of species (Brown & Wilson, 1956; Schluter,
 134 2000), species sorting where only species with different ecological niches will coexist
 135 (Anderson & Weir, 2021), or even the extinction of an entire lineage (e.g. Silvestro et
 136 al., 2015). Indeed, the number of studies suggesting that competition is a relevant factor
 137 in shaping the distribution of organisms at global and macroevolutionary scales has
 138 rapidly increased (Gotelli et al., 2010; Pigot & Tobias, 2013; Silvestro et al., 2015;
 139 Pigot et al., 2018). Pigot and Tobias (2013) for example, found that the coexistence of
 140 closely related species of birds seems to be mediated by differences in their ecological
 141 niche. On the other hand, other factors might play a role, and these same authors later
 142 found that dispersal ability also determine how fast two species might expand their
 143 distributions and coexist (Pigot & Tobias, 2015; Pigot et al., 2018). Dispersal is indeed
 144 widely suggested to play a central role in the redistribution of organisms, contributing to
 145 the colonization of new areas and range shifts, representing a potentially important
 146 component driving coexistence dynamics (Lowe & McPeck, 2014; Jönsson et al., 2016;
 147 Weber & Strauss, 2016).

148 Although biotic interactions and dispersal ability have been frequently invoked to
 149 explain both spatial and temporal biodiversity patterns, the geographical scenario where
 150 species thrive is another important aspect to be considered. Islands are widely known
 151 for producing unique biodiversity patterns given their smaller size and geographic

isolation compared to continental settings (MacArthur & Wilson, 1967; Losos & Ricklefs, 2009; Baeckens & Van Damme, 2020). Changes in body size like gigantism and dwarfism are frequently reported in insular vertebrates ("island rule", Foster, 1964; Van Valen, 1973; see also Benítez-López et al., 2021 but see Meiri, 2007). Patterns like these are suggested to be the result of several mechanisms characterizing insular systems, such as reduced predation, relaxed inter-specific competition and resource limitation (Losos & Ricklefs, 2009; Baeckens & Van Damme, 2020; Benítez-López et al., 2021). However, the consequences of these insular particularities on the coexistence dynamics of species are less clear (Ricklefs, 2010; Pigot et al., 2018). On one hand, it is feasible to expect that coexistence might be achieved faster on islands compared to continents given that the first comprise smaller geographical limits and is usually characterized by a relaxation of biotic constraints. On the other hand, islands harbor populations that might be more likely to suffer stochastic extinctions or introgression possibly erasing any signs of sympatry (Pigot et al., 2018).

Several other ecological, physiological, and behavioral aspects can also influence species coexistence dynamics (Gaston, 2000; Buckley et al., 2012). Squamate reptiles (i.e. lizards and snakes) are ectothermic animals and their distributions will frequently be limited by their physiological requirements (Buckley et al., 2012; Vitt & Caldwell, 2014). As a result, their distributions can be strongly determined by climatic conditions (e.g. Buckley et al., 2012). Lizards and snakes, however, comprise more than 10.000 species ranging across a wide spectrum of latitudes and biomes (Roll et al., 2017; Uetz & Hosek, 2017). While the species richness of snakes shows the typical latitudinal gradient similar to endothermic organisms, lizards show a pattern distinct from any other tetrapod group, with a higher richness in Australia (see Roll et al., 2017). Furthermore, although closely related, snakes and lizards differ greatly in their

177 morphology and in several aspects of their natural history (Sites et al., 2011; Vitt &
178 Caldwell, 2014). Generally speaking, snakes have a slower metabolism and lower
179 population density compared to lizards (Vitt & Caldwell, 2014); lizards comprise more
180 diverse diet habits, including insectivorous, carnivorous and even herbivorous species
181 (Pianka & Vitt, 2003; Meiri, 2008), whereas snakes are strictly carnivorous (Greene,
182 1997). This impressive diversity is actually an indication that species distributions in
183 squamates might not be regulated by the same processes, and that aspects not related to
184 climate might also play an important role in shaping species distributions (e.g. Algar et
185 al., 2013; Cunningham et al., 2015). *Anolis* lizards, for example, comprise a classic
186 example of squamate reptiles among which species coexistence is strongly determined
187 by interspecific competition (Losos, 2011).

188 To reveal the relevant factors shaping the coexistence dynamics of a diverse group
189 such as squamates, and whether snakes and lizards share or not these factors, we need
190 investigations across broader taxonomic and spatial contexts. Here we take this
191 approach by investigating the role of biotic interactions, dispersal ability and different
192 geographical settings in shaping the coexistence dynamics across closely related species
193 of squamate reptiles. To do that, we use a model-comparison framework to (1) evaluate
194 the main speciation mode (i.e sympatric or allopatric) in lizards and snakes; (2) test if
195 the tendency to geographically overlap over time increases in species that are
196 ecologically distinct, (3) have high dispersal abilities, and/or (4) occur on islands. Our
197 study is the first to explicitly model the coexistence dynamics of squamates over time in
198 broad temporal and spatial scales. On top of that, we take into account the possible
199 effects of different ecological and geographical scenarios in driving coexistence. Our
200 results provide insights into the factors shaping species distributions and the dynamics
201 of coexistence, contributing to a better understanding of the processes shaping

202 biological communities and, at a broader scale, of those that modulate the evolution of
203 biodiversity.

204

205 **METHODS**

206

207 **Closely related species pairs**

208

209 To explore coexistence dynamics in squamates, we used one of the most
210 complete molecular phylogenies (Tonini et al., 2016) to define a pool of species pairs
211 ("sister" species), representing the most closely related species in the maximum-
212 likelihood topology. Our pool included only species from well sampled genera (70% or
213 more of all described species of a given genus had to be in the phylogeny) and for
214 which the phylogenetic placement of species pairs was highly supported (equal to or
215 higher than 0.95). To calculate genus sampling, we followed the taxonomy of the
216 Reptile Database up to January 2017 (Uetz et al., 2017). We were able to identify 538
217 species pairs (161 of snakes and 377 of lizards).

218

219 **Geographical ranges**

220

221 To determine if species comprising a given species pair coexist, we used species
222 polygon range maps mostly from Roll et al. (2017). For an additional 20 species, whose
223 maps were not available in Roll et al. (2017), we used those provided by the IUCN Red
224 List Assessment. We used the range map provided by Birskis-Barros et al. (2019) for
225 the South American rattlesnake *Crotalus durissus*, given the outdated distribution for
226 this species in the previous two databases. From the 538 species pairs, we could not

227 obtain range maps for four pairs given that at least one of the species did not have a map
228 available.

229 From the 534 species pairs for which we had range maps, we excluded species
230 that occur in marine or mangrove environments (10 pairs) and terrestrial pairs where
231 species are separated by marine barriers (species occurring on different landmasses or
232 islands, see Pigot & Tobias, 2015) (those represent 57 pairs). The dispersal dynamics
233 characterizing these species pairs might potentially differ from those where species are
234 terrestrial and occur on the same landmass and would potentially add noise to our
235 analyses.

236

237 **Age since divergence**

238

239 To model coexistence over time, we used the age since divergence of each
240 species pair estimated by Tonini et al. (2016) as our temporal measurement.
241 Given that estimates of age since divergence can be highly uncertain, we randomly
242 chose 100 different phylogenies from those generated by Tonini et al. (2016) to obtain a
243 range of possible age estimates for our species pairs. It is important to mention that
244 these 100 phylogenies were generated using the same backbone molecular phylogeny
245 and, therefore, the relationships between species pairs remain the same, but the age
246 estimates differ (see Tonini et al. 2016 for details). Therefore, we incorporated in our
247 analyses 100 different age since divergence estimates for each species pair (see below).

248 To avoid including species pairs for which evolutionary history might have been
249 strongly influenced by extinction we set a divergence time limit to our species pairs. We
250 first took the median of the 100 ages since divergence estimates for each pair and kept
251 those pairs comprising a median equal to or lower than 20 million years ago. After all

curatorial work, our final pool comprised 448 pairs of species (132 of snakes and 316 of lizards) that we used to investigate the coexistence dynamics in squamates (see Appendix I). These 448 pairs span a wide diversity of taxonomic groups and geographical areas representing the vast diversity observed in squamates (see Figure 1). We also ran additional analyses using only those pairs for which the median of the age since divergence was equal to or lower than 10 million years ago (124 of snakes and 243 of lizards), a more conservative way to include species pairs with respect to potential loss of history to extinction.

Geographical overlap

We considered a species pair as sympatric or allopatric based on the spatial overlap of their geographical ranges. We considered a pair as sympatric when more than 30% of the smaller species range overlapped with the range of the other species. We also ran additional analyses considering a given pair as sympatric when the smaller range overlapped more than 70%. These additional analyses allowed us to explore if our results on the coexistence dynamics of squamates would differ depending on how we define sympatry. We quantified geographic overlap using the Raster package in R (Hijmans, 2019). We should note that we choose a simpler geographical categorization for methodological purposes (see Statistical analyses), considering as “allopatric” species pairs that might have been originated under processes such as vicariance, parapatric speciation, or founder events (see Skeels & Cardillo, 2019).

Geographical and ecological traits

277 *Size of the geographical area*

278

279 To explore if the size of the geographical area affects the tendency by which
280 closely related species coexist over time, we categorized each species pair as occurring
281 on islands or continents. We considered as islands all geographically isolated land
282 masses smaller than Madagascar (the largest island considered). When the species
283 comprising a given pair occurred both on islands and continents, we considered the
284 geographical area as the one where both species occur. We did not have any cases
285 where both the species occurred on islands and continents.

286

287 *Ecological similarity*

288

289 To quantify how ecologically similar two species are and, thus, to explore the
290 importance of species interactions in affecting the tendency of closely related species to
291 coexist over time, we measured specimens deposited in several scientific collections
292 between the years of 2017 and 2019 (see supplementary methods). This massive data
293 collection allowed us to obtain measurements for more than 2000 specimens of lizards
294 and snakes comprising 300 and 152 species, respectively (150 and 76 sister pairs of
295 lizards and snakes; Appendix II and III). Therefore, we were able to obtain
296 morphological data for 47% and 57% of the pairs from our total pool (lizards and
297 snakes, respectively). On average, we were able to measure 6 specimens per species of
298 lizards and 5.7 specimens per species of snakes. For additional details on how we
299 determined the sex and the sexual maturity of the specimens see the supplementary
300 methods.

301 We focused on measuring morphological traits known to be associated with
 302 different axes of the ecological niches of squamates. Additionally, we aimed to include
 303 only males in our dataset as a way to decrease any noise caused by morphological
 304 differences resulting from sexual dimorphism. Measurements for lizards comprised
 305 snout-vent length (SVL), head length, width and height, jaw length, mid-body
 306 circumference, the length of the humerus, ulna, femur and tibia, and the distance
 307 between fore and hind limbs. Measurements for snakes comprised snout-vent length,
 308 head length, width and height, mid-body circumference, tail length and eye diameter.
 309 This dataset was combined with the morphological dataset for snakes of the family
 310 Viperidae generated by Alencar et al. (2017), and with a few measurements for snout-
 311 vent length taken from the literature (see Appendix II and III), all measurements from
 312 adult males only.

313 To explore the role of species interactions in driving the tendency of species to
 314 geographically overlap over time, we used two different metrics as proxies for
 315 ecological similarity of a given species pair. First, we calculated the absolute difference
 316 between the average of the log-transformed SVL of the species comprising each pair.
 317 The larger the difference between the SVL, the less ecologically similar species are
 318 expected to be (Wilson 1975; Ricklefs & Miles, 1994). We chose to use SVL as our
 319 metric for body size given it is traditionally used by herpetologists as a proxy for size
 320 (see Meiri, 2008). As a next step, we generated one morphospace for snakes and another
 321 for lizards but excluding legless lizards, and a second one for all lizards including
 322 legless lizards but excluding the distance between the fore and hind limbs measurement.
 323 To generate these morphospaces we first size-corrected the morphological variables
 324 (log-shape ratios) for each species by dividing each variable by the geometric mean
 325 calculated for each species (see below) and log-transforming the resulting value (Price

et al., 2019; Friedman et al., 2020). The geometric mean was calculated for lizards as the cube root of the product of species average SVL, mid-body circumference and length of the limbs, and for snakes as the cube root of the product of species' average SVL, tail length and mid-body circumference. We chose not to calculate the geometric mean across all morphological variables because most of them are inclusive measures of these dimensions of size (see Price et al., 2019). We chose to apply this size-correction method because it allows us to produce a metric that preserves the allometric component of shape (see Price et al., 2019). We performed phylogenetic principal component analyses using the R package phytools (Revell, 2009; Revell, 2012) to generate these morphospaces using the consensus phylogenetic tree from Tonini et al. (2016). Using these morphospaces, we calculated the Euclidean distance between species comprising each pair across all PhylPCA axes combined. The larger the euclidean distance, the less ecologically similar species are expected to be. In the end, we had two proxies for ecological similarity, (1) differences in SVL and (2) the Euclidean distance between species pairs within the morphospace.

To explore if the proxies for ecological similarity, that is, differences in SVL and differences in shape (Euclidean distance) were correlated, we performed PGLS analyses using ten phylogenetic trees from Tonini et al. (2016). These analyses suggested that despite being significantly correlated in lizards (R^2 0.59 – 0.77, $p < 0.05$), differences in the SVL and Euclidean distances are not correlated in snakes ($R^2 < 0.01$, $p > 0.1$). In the subsequent analyses, we decided to use only the SVL differences for lizards and keep both the differences in SVL and the Euclidean distance as proxies for ecological similarity in snakes.

Dispersal ability

351

352 Despite often being considered a species trait, here we assess dispersal ability as
 353 a trait characterizing a given species pair due to methodological reasons (see statistical
 354 analyses below). To do this, we used two different metrics. First, we used the log-
 355 transformed ratio between the average range size of each pair and the age since the
 356 divergence of that pair. Range size can be a good proxy for dispersal ability because
 357 larger ranges might reflect a higher ability of lineages to geographically expand (Brown
 358 et al., 1996; Pigot et al., 2018). We decided to take into account the age since
 359 divergence between species comprising each pair because older lineages might have
 360 larger geographical ranges simply because they had more time to expand geographically
 361 and not as a result of dispersion ability. Therefore, the ratio between range size and age
 362 of divergence reflects the rate at which a given species pair was able to geographically
 363 expand. We calculated this ratio 100 times for each pair using the different age
 364 estimates obtained previously. As a next step, we used the average SVL for each pair.
 365 Because larger animals might also have larger home ranges and/or territories (Brown et
 366 al., 1996; Bonner, 2011), we expect that larger squamates might have higher dispersal
 367 ability. It is important to note that the analyses using average SVL as a proxy for
 368 dispersal ability included less species pairs compared to our first proxy given we
 369 measured a smaller number of species than our total species pool (see above).

370 To explore if the two proxies were correlated with each other, we performed
 371 PGLS analyses on the average snout-vent length and the average range size/age ratio,
 372 using ten randomly sampled sets of the latter and the ten corresponding phylogenetic
 373 trees from Tonini et al. (2016). Interestingly, all these analyses suggested that despite
 374 being significantly correlated ($p < 0.05$), the relationships have either very low R^2 (0.06

– 0.08, for lizards) or moderate R^2 (0.14 – 0.18, for snakes). For this reason, we decided to keep both proxies in our subsequent analyses (see below).

Statistical analyses

We explored the coexistence dynamics of squamates over time by using different probabilistic models of species co-occurrence (Figure 2, Pigot et al., 2013, 2015). In this framework, the dynamic of coexistence is modelled as a constant rate Markov process, and maximum likelihood is used to perform model fitting and parameter estimates. In a general manner, the models allow us to calculate the probability that a species pair exists in its current geographical state (i.e. allopatry or sympatry) given their age since divergence and the parameters controlling the rates of transition from allopatry to sympatry (σ) and from sympatry to allopatry (ϵ) (Figure 2, see also Pigot & Tobias, 2015). Therefore, we were able to model how coexistence possibly changed over time by using a given set of species pairs. We fitted four different models to our dataset using the R package *msm* (Jackson, 2011): 1) allo-one-way, which assumes that lineages diverged in allopatry and then undergo a transition to sympatry ($\gamma = 1$, $\sigma > 0$, and $\epsilon = 0$); 2) symp-one-way, which assumes that lineages diverged in sympatry and then undergo a transition to allopatry ($\gamma = 0$, $\sigma = 0$, and $\epsilon > 0$); 3) allo-two-way, assumes that lineages diverged in allopatry, become sympatric but became allopatric again ($\gamma = 1$, $\sigma > 0$, $\epsilon > 0$); Symp-two-way, which assumes that lineages diverged in sympatry, undergo a transition to allopatry but come back to sympatry later ($\gamma = 0$, $\sigma > 0$, and $\epsilon > 0$).

A schematic workflow of the model comparisons we performed is presented in Figure S1. First, we fitted all four models 100 times across species pairs of lizards and

snakes separately using their age since divergence obtained from 100 different topologies (see above). We then evaluated how many times a given model was favored over the others using the Akaike Information Criterion (Burnham & Anderson, 2002) (see Table S1-S4). We considered the best model to be the one with the lowest AIC value and with a ΔAIC higher than two. To evaluate the effects of geographical position (island vs continent), ecological similarity (body size difference and euclidean distance) and dispersal ability (ratio of range area and age, average body size) in shaping the coexistence dynamics of squamates, we ran another set of analyses comparing all four models but also including the allo-one-way and the allo-two-way models with each candidate factor as a covariate of σ (the parameter describing the transition rate to sympatry). We decided not to include the Symp-two-way model with candidate factors as covariates in our model comparison because transitions to coexistence occur at a distinct stage in the symp-two-way model relative to those in the allopatric models. In summary, we performed model comparisons across the six models (allo-one-way, symp-one-way, allo-two-way, symp-two-way, allo-one-way with covariate, allo-two-way with covariate) five times for snakes and four times for lizards, including one distinct candidate driver in each of them (Table S1-S4, see summary in Figure S1). We performed model comparisons separately using each candidate factor as they frequently involved distinct species pairs datasets (e.g. morphology based proxies comprise fewer species pairs than range size, for example).

When candidate drivers significantly improved a model over the others, we quantified the hazard ratio to evaluate the direction of the relationship between transition rates to sympatry and the driver of coexistence (Table S5). To illustrate how transition rates to sympatry are affected by the drivers of coexistence, we extracted estimates of transition rates to sympatry under the best models selected and the

corresponding hazard ratios for different values of the drivers (Figure 3). We performed posterior predictive simulations to evaluate how well the best models could actually predict the empirical data (see supplementary methods).

RESULTS

For our main dataset that considered sympatric species pairs as those with at least 30% of the smallest distribution overlapping and included pairs that diverged 20 million years ago or less, 37% and 48% of the species pairs are sympatric in snakes and lizards, respectively (Table S6). Proportions were very similar when considering only the species pairs with a median age since divergence of less than 10 mya (see Table S6). However, as expected, the number of sympatric pairs decreased when we considered as sympatric only those pairs with more than 70% of the smallest distribution overlapping (Table S6). Regardless of the dataset, however, lizards consistently show a higher proportion of sympatric pairs compared to snakes (Table S6). In general, sympatric pairs tend to be older than allopatric pairs among snakes (Table S7). In lizards, however, sympatric and allopatric species pairs have very similar ages of divergence, with sympatric pairs still being slightly older across datasets comprising pairs that have diverged 20 mya or less (see Table S7).

Coexistence dynamics in squamates

Model selection without covariates

When comparing only the four models without adding any candidate traits that could potentially be driving coexistence dynamics, the “allo-two-way-model” was

450 considered the best model for about 75% of the age datasets analyzed for snakes either
 451 when considering species pairs that diverged less than 20 or 10 mya (Table S1-S2).
 452 These results suggest that snakes generate species mainly via allopatric speciation with
 453 transitions to sympatry over time and back to allopatry as lineages become older. On the
 454 other hand, by comparing the four models without adding the covariates, we could not
 455 recover a unique best model for lizards (Figure S1-S2). The symp-two-way-model and
 456 the allo-two-way-model are equally likely, which suggests that geographical overlap in
 457 lizards changed considerably through time. Additionally, no best model was detected
 458 when performing model comparison with an overlap cut-off of 70% either in lizards or
 459 snakes (see Table S3-S4).

460

461 *Model selection with the geographical setting as covariate*

462

463 Being on an island or continent seems to be an important driver of species
 464 coexistence in snakes but not necessarily in lizards. When considering the two
 465 additional models that include the geographical setting as a covariate on the transition
 466 rate to sympatry, model comparisons suggested that the allo-two-way model with the
 467 covariate was the best model across all age datasets in snakes, regardless of only
 468 including younger pairs or not (Table S1-S2). According to this model, transition rates
 469 to sympatry substantially increase in species pairs of snakes occurring on islands
 470 compared to those occurring on continents (Figure 3, Table S5). When performing
 471 model comparison under the 70% overlap cut-off, the allo-two-way model with the
 472 covariate was still recovered as the best model across 76% of the age datasets in snakes.
 473 However, no single best model was recovered when considering only the younger pairs
 474 (Table S3-S4), although the model with geographical setting as covariate was still

among the best models (it was tied with another model in 99 out of 100 comparisons). In contrast, no best model was selected for lizards and, therefore, we do not have evidence that being on an island or continent is relevant in driving their coexistence dynamics (Table S1-S4). It is also worth mentioning that, in general, the model that includes geographical setting as a covariate was frequently not among the best tied models in lizards, with the exception of when considering the 70% overlap threshold and species pairs that diverged less than 10 mya (33, 26, 56, or 97 out of 100 model comparisons depending on the dataset, see Tables S1-S4).

Model selection with ecological similarity as covariate

Ecological similarity seems to be an important driver of species coexistence in snakes but not in lizards. Model selection suggested that the allo-two-way model with body size difference as the covariate was the best model across all age datasets for snakes (all 100 model comparisons), regardless of whether the data were restricted to younger pairs (Table S1-S2). According to this model, transition rates to sympatry increase with increasing differences in body size between species pairs (Figure 3, Table S5). Despite always suggesting the same positive relationship, transition rates to sympatry and the hazard ratio have unreliably high values when estimated using ages of divergence extracted from some phylogenies (Figure 3, Table S5). This suggests that model fitting might have failed to converge in these datasets. However, simulations using parameter estimates under the best model are still able to recover proportions of sympatric and allopatric pairs similar to the empirical data (see below). When analyzing species pairs of snakes under the 70% overlap cut-off no model was preferred, but the model with body size difference as the covariate tied with another model about 60% of

the time (Table S3-S4). In contrast, we did not recover a best model when adding the differences in shape (measured by the Euclidean distance, see Methods) of species pairs as the covariate on transition rates to sympatry in snakes, although this model is frequently tied with other models (see Tables S1-S4). Additionally, we did not find strong evidence of ecological similarity being an important driver of coexistence in lizards (Table S1-S4). It is important to mention, however, that although we did not recover a best model for lizards, the model with body size difference as the covariate is among the best models in several of the model comparisons performed (Table S1-S4).

Model selection with dispersal ability as covariate

Contrary to the analyses above, dispersal ability seems to be a relevant driver of coexistence dynamics in lizards but not snakes (Table S1-S2). Model selection suggested that the allo-two-way model with the ratio of range size and age as the covariate was the best model across more than 80% of the age datasets of lizards when considering the overlap threshold of 30%, regardless of whether only younger pairs were included (Table S1-S2). According to this model, transition rates to sympatry increase with faster rates of geographic expansion (i.e. the larger the ratio between geographical range and time; Figure 3, Table S5). We did not recover a single best model when analyzing species pairs of lizards under the 70% overlap cut-off (Table S3-S4), but the model with dispersal as a covariate was usually among the tied models (above 80% of the comparisons). When performing model selection with mean body size (the other proxy for dispersion ability) as the covariate, we recovered the allo-two-way model that included the covariate as the best model for 57% of the age datasets but only in our main analyses (30% overlap and all species pairs, Table S1). Surprisingly,

525 however, according to this model transition rates to sympatry increase the smaller the
526 species (Figure 3, Table S5). We did not find a single best model for these model
527 comparisons using species pairs of snakes but models with dispersal proxies as
528 covariates usually tied with other models (Table S1-S4).

529

530 Posterior predictive simulations

531

532 For all best models with covariates selected in our main analyses (30% overlap
533 between species and including all species pairs), simulated proportions of sympatric
534 pairs frequently recovered the empirical proportions (Figure S2).

535

536 **DISCUSSION**

537

538 In lizards and snakes, neither time or sympatric speciation alone can explain the
539 coexistence patterns observed among extant species. On the other hand, by performing
540 extensive data collection and comparing statistical models, we found evidence for
541 completely distinct drivers underlying the coexistence dynamics of these two closely
542 related groups. In general, our results suggest that the species' geographical setting,
543 ecologically similarity, and dispersal ability might play different roles in shaping the
544 coexistence dynamics within each taxonomic group analyzed.

545 By taking a first look at the patterns of geographical overlap, we found that most
546 species pairs of snakes are allopatric. Allopatric speciation is indeed the rule across
547 vertebrates, with many vertebrate groups having more allopatric than sympatric sister
548 species (Barraclough & Vogler, 2000; Pigot & Tobias, 2015; García-Navas et al., 2020;
549 see review in Hernández-Hernández et al., 2021). In lizards, however, allopatric and

550 sympatric pairs had similar proportions. Allopatric pairs of snakes tend to be younger
 551 than sympatric ones, which conforms to a scenario where coexistence would be
 552 achieved with time. Indeed, without adding any covariates, the coexistence dynamics in
 553 snakes are best described by allopatric speciation with lineages eventually becoming
 554 sympatric as time goes by and becoming allopatric again as they get older (allo-two-
 555 way model, Table S1-S2). In lizards, on the other hand, sympatric species and allopatric
 556 pairs tend to have very similar ages, with allopatric pairs being older in some cases
 557 (Table S7). This result might suggest two scenarios: (1) sympatric speciation plays an
 558 important role in the diversification of lizards; (2) allopatric speciation is still the rule,
 559 but time alone does not explain the coexistence dynamics of lizards. In any case, we
 560 indeed could not recover a single best model in the first set of model comparisons for
 561 the group (Table S1-S4). It is worth noting that when a single model is chosen in
 562 lizards, the sympatric model is the one selected (19 vs 1, Table S1), and when two
 563 models are equally likely (80% of the time), the sympatric model is also among them
 564 (Table S1).

565 Taken together, these first results suggest that lizards might be more
 566 heterogeneous and have more dynamic distributions compared to snakes, either because
 567 their lineages are influenced by different processes and/or because their distributions
 568 change more through time. A wider array of processes acting simultaneously could
 569 potentially mask coexistence signals when we include all species in the same category
 570 “Lizards”. Indeed, lizards occurring in the same islands can show completely distinct
 571 patterns of diversification. At the Socotra Archipelago, for example, closely related
 572 species of the genera *Hemidactylus* and *Haemodracon* show sympatric distributions
 573 with marked differences in body size, whereas those of the genus *Pristurus* tend to be
 574 allopatric or parapatric with no apparent divergence in body size (see García-Porta et

575 al., 2016). Furthermore, species richness can indeed be achieved via distinct pathways
576 in distinct lizard clades and geographic regions. High species richness is associated with
577 low functional divergence in agamids and gekkoes but with high functional diversity in
578 varanids and scincids (Skeels et al., 2020). Exploring whether distinct clades are also
579 characterized by distinct coexistence dynamics might be an interesting next step.
580 However, we would need a much higher number of phylogenetically well-supported
581 species pairs within each of the distinct clades to explore group-specific coexistence
582 dynamics.

583 When adding covariates to the transition rates to sympatry, the allo-two-way
584 model is frequently recovered as the best model across the analyses for which we were
585 able to recover a single best model. In contrast to previous findings in birds (Pigot et al.,
586 2018), islands seem to promote species coexistence in snakes. Surprisingly, however,
587 this effect was not evident in lizards. In other words, the model with the geographical
588 setting as covariate does not have higher support compared to those without the
589 covariate in lizards. Indeed, in this set of analyses for lizards, ties were mainly between
590 the allo-two-way and symp-two-way models that lack covariates (Table S1). In general,
591 snakes have a slower life history and lower energetic demands compared to lizards
592 (Pough, 1973; Vitt & Caldwell, 2014). Therefore, resources on islands might appear
593 less scarce for snakes than lizards, potentially allowing snakes to more easily coexist in
594 insular environments given the relaxation of some biotic constraints (e.g. predation, and
595 perhaps less intense competition) and the smaller geographical limits of islands
596 compared with continental settings. Lizards, on the other hand, have high population
597 densities on islands (Novosolov et al., 2015) and the resultant scarcity of resources
598 might lead to much more intense competition. Hence, the smaller geographical limits

599 would not increase coexistence in insular lizards at least when compared with
600 continental settings.

601 Species interactions also underlie species coexistence in snakes, and divergence
602 in body size (but not shape) seems to be the most likely route to avoid competition and
603 promote coexistence, similar to what has been suggested for birds (Pigot et al., 2018).
604 On the other hand, we found no definitive or clear evidence that ecological similarity is
605 a relevant driver of species coexistence in lizards. For 99% of the time when we had
606 more than one best model (a tie) the model involving ecological divergence (complex-
607 allo-two-way) was among them for lizards (Table S1). The prediction that the likelihood
608 of coexistence increases with increasing the ecological difference between species,
609 assumes that some species pairs would accumulate more morphological differences
610 allopatrically allowing them to coexist faster than others (i.e. species sorting, Grant,
611 1972; Davies et al., 2007). Although we did not find definitive support for this
612 prediction in lizards, morphological divergence could still be driving the coexistence
613 dynamics at the local scale. If this is the case, we would expect that sympatric
614 populations of a given species pair would be more morphologically distinct compared to
615 allopatric populations of these same species (character displacement, Brown & Wilson,
616 1956; Davies et al., 2007). We might not have been able to capture these smaller-scale
617 differences given our large-scale approach and goals. A fruitful next step would be to
618 apply recently developed and promising frameworks to detangle the role of species
619 sorting and character displacement in speciation and species coexistence dynamics (see
620 Anderson & Weir, 2021), and to increase the intra-specific morphological sampling
621 allowing us to compare morphological divergence at the population-level.

622 Even though competition has widely been seen as an important factor shaping
623 community structure (e.g. Darwin, 1859; Webb et al., 2002; Cavender-Bares et al.,

2009), a lack of strong evidence of competitive interactions shaping trait-divergence has also been suggested (e.g. Meiri et al., 2011; Stuart & Losos, 2013; Shi et al., 2018, Slavenko et al., 2021). Slavenko et al. (2021) suggested that environmental filtering might be more important in driving morphological divergence in lizards than competitive interactions, which could potentially explain why we did not find clear evidence for morphological differences mediating coexistence dynamics in lizards. Another possibility is that morphological divergence could be more relevant in some geographical regions than others or within some groups of lizards (see Skeels et al., 2020). Competition might be more prevalent in climatically stable areas (Dobzhansky, 1950; MacArthur, 1969; Henriques-Silva et al., 2019) or regions with lower resource availability and, therefore, morphological divergence could be an important driver of coexistence at lower compared with higher latitudes or on islands compared with continents (e.g. García-Porta et al., 2016), respectively.

In contrast to what we found for competitive interactions, dispersal ability (measured by the ratio between range size and age of divergence) seems to be a relevant driver of species coexistence in lizards but not necessarily in snakes. It is worth mentioning that our dispersal metric that uses range size/age has its limitations and assumptions. For example, it is thought that species might both start and go extinct with smaller range sizes (Foote, 2007). Hence we might be pulling together species at very different stages which might mask the potential effect of dispersal and perhaps explain the lack of dispersal effects in snakes. By restricting the time window of analysis, this problem might be ameliorated but not explicitly taken into account. However, the lack of an association is still present in snakes when considering species pairs that diverged less than 10 Mya.

648 Lizards indeed have been suggested to have comparatively higher dispersal
649 abilities than snakes, at least regarding long-distance dispersal to oceanic islands (see
650 Pitta et al., 2013). It is important to note, however, that we are not able to disentangle
651 between the ability to move across geographical space and the ability to establish
652 populations in new geographical areas (see Jönsson et al., 2016), as we considered both
653 as being part of the “dispersal ability” of species. Simply being more mobile across
654 space would not necessarily mean that species would successfully establish populations
655 in new geographical areas, possibly allowing species coexistence. This scenario has the
656 potential to be especially true for snakes, for which our results suggest that competitive
657 interactions are relevant drivers of species coexistence. Therefore, in snakes, only those
658 species that are different enough might be able to rapidly coexist, even if others have
659 high dispersal abilities. The establishment of populations, and consequently coexistence
660 among species, might also be compromised if not enough time has elapsed for strong
661 reproductive barriers to emerge and prevent populations from fusing (Weir & Price,
662 2011). The interaction between dispersal and competition, as well as between
663 competition and geographical setting, in driving species coexistence deserves further
664 investigation (see Lowe & McPeck, 2014; Jönsson et al., 2016). Indeed, the statistical
665 framework to explore the interaction between coexistence drivers is already available
666 (Pigot et al., 2016, 2018). However, we would need a much higher number of
667 phylogenetically well-supported species pairs, as well as their morphological
668 information to be able to properly investigate these questions.

669 Despite being favored in some of the model comparisons, models with mean
670 body size as the covariate on the transition rates to sympatry suggested that the smaller
671 the lizard the higher these transition rates. This might look unexpected given that both
672 proxies for dispersal ability (ratio between range and age, and body size) were

673 positively, despite weakly, correlated (see Methods). However, the body size dataset
674 comprises a much smaller number of sister pairs compared to the range and age ratio
675 dataset, which could explain the differences in the coexistence dynamics depicted by the
676 models.

677 On the other hand, we cannot rule out the possibility that the preconception that
678 larger species would be better dispersers might not be totally true for lizards. Going
679 further, the reason for a negative relationship between body size and the transition rates
680 to sympatry could be interpreted in light of the well-known link between body size and
681 specific axes of the ecological niche in animals (Peters, 1986; Meiri, 2008; Clarke,
682 2021). Contrary to what has been suggested for several other organisms, Costa et al.
683 (2008) found that body size is negatively related to dietary niche-breadth in predatory
684 lizards and, therefore, the diversity of prey consumed decreases as body size increases.
685 Furthermore, several small lizard species are insectivorous (Pianka & Vitt, 2003) and
686 might face less severe pressures when dealing with the scarcity of food resources
687 compared to larger species that may rely on more limited food availability. This, in turn,
688 might enable smaller species to share the same resources and coexist. Although some
689 large bodied species of lizards are herbivorous, meaning that their food might be readily
690 available in some environments, the higher total metabolic rate of larger animals also
691 requires a greater caloric intake (Pough, 1973). Coupled with that, small insectivorous
692 lizards can be more mobile than larger herbivorous due to the higher energetic food
693 taken by the first (Pough, 1973). All of these factors could help to explain a higher
694 incidence of coexistence over time in smaller lizards.

695 Going further, the relative importance of each coexistence driver in distinct
696 stages of the coexistence process is an aspect that deserves to be explored in squamates.
697 As found by Pigot et al. (2018), patterns of narrow geographic overlap in birds seem to

698 be driven by dispersal abilities and age, whereas wider coexistence patterns are mainly
 699 driven by ecosystem productivity and niche-divergence. For several of the analyses
 700 performed here, especially when considering as sympatric the species pairs with a
 701 geographic overlap higher than 70% of the smallest distribution, model comparisons
 702 were not able to choose between models. This probably occurs because using the 70%
 703 cut-off inevitably decreases the number of sympatric pairs, probably affecting the
 704 ability of these analyses to discern between models. Therefore, the addition of more
 705 well-supported species pairs coupled with their morphological information is essential
 706 to detangle between the differential drivers of species coexistence in distinct moments
 707 in the past.

708

709 **Conclusions**

710 This study shows that lizards and snakes, although closely related, differ greatly
 711 in the drivers of species coexistence. Speciation seems to predominantly occur via
 712 allopatric speciation in snakes. Lizards, on the other hand, seem to be more
 713 heterogeneous and have more dynamic distributions, which likely prevented us from
 714 recovering a single best model without the addition of covariates (see Results). In
 715 snakes, species that occur on islands or have different body sizes are more likely to
 716 coexist. In contrast, lizard species are more likely to co-occur when they have higher
 717 dispersal abilities.

718 Indeed, it has been widely shown that lizards tend to exhibit unique diversity
 719 patterns that frequently do not follow the “rules” that usually apply to snakes or other
 720 vertebrate groups (Meiri, 2007; Powney et al., 2010; Roll et al., 2017; Novosolov et al.,
 721 2018). These differences might have profound consequences either for community
 722 structure and lineage diversification, and care should be taken when analyzing these two

723 taxonomic groups together (see also Slavenko et al., 2019; Whiting & Fox, 2020).
 724 Beyond this, our results emphasize that when analyzing biogeographical,
 725 macroecological or macroevolutionary patterns and processes, one should take into
 726 account not only the geographical scenario but also who these organisms are (see also
 727 White, 2016; Skeels et al., 2020). It is the interaction between where and who that will
 728 ultimately shape biodiversity patterns.

729

730 REFERENCES

731

- 732 Alencar, L. R. V., Martins, M., Burin, G., & Quental, T. B. (2017). Arboreality
 733 constrains morphological evolution but not species diversification in vipers.
 734 Proceedings of the Royal Society B, 284, 20171775.
 735
- 736 Algar, A. C., Mahler, D. L., Glor, R. E., & Losos, J. B. (2013). Niche incumbency,
 737 dispersal limitation and climate shape geographical distributions in a species-rich island
 738 adaptive radiation. Global Ecology Biogeography, 22, 391-402.
 739
- 740 Anderson, S. A. S., & Weir, J. T. (2021). Character displacement drives trait divergence
 741 in a continental fauna. Proceedings of the National Academy of Sciences of the United
 742 States of America, 118, e2021209118.
 743
- 744 Baeckens, S., & Van Damme, R. V. (2020). The island syndrome. Current Biology, 30,
 745 329-339.
 746
- 747 Barraclough, T. G., & Vogler, A. P. (2000). Detecting the Geographical Pattern of
 748 Speciation from Species-Level Phylogenies. The American Naturalist, 155, 419-434.
 749
- 750 Benítez-López, A., Santini, L., Gallego-Zamorano, J., Milá, B., Walkden, P.,
 751 Huijbregts, M. A. J., & Tobias, A. J. (2021). The island explains consistent pattern of
 752 body size evolution in terrestrial vertebrates. Nature Ecology & Evolution, 5, 768-786.
 753

754 Birskis-Barros, I., Alencar, L. R. V., Prado, P. I., Böhm, M., & Martins, M. (2019).
755 Ecological and conservation correlates of rarity in New World pitvipers. *Diversity*, 11,
756 1-15.
757
758 Bonner, J. T. (2011). *Why size matters: From bacteria to blue whales*. Princeton:
759 Princeton University Press.
760
761 Brown Jr, W. L., & Wilson, E. O. (1956). Character displacement. *Systematic Zoology*
762 5:49-64.
763
764 Brown, J. H., Stevens, G. C., & Kaufman, D. M. (1996). *Annual Review of Ecology*
765 *and Systematics*, 27, 597-623.
766
767 Buckley, L. B., Hurlbert, A. H., & Jetz, W. 2012. Broad-scale ecological implications of
768 ectothermy and endothermy in changing environments. *Global Ecology and*
769 *Biogeography*, 21, 873-885.
770
771 Burnham, K. P., & Anderson, D. R. (2002). *Model selection and mixed model*
772 *inference: a practical information-theoretic approach*. New York: Springer.
773
774 Cavender-Bares, J., Kozak, K. H., Fine, P. V. A., & Kembel, S. W. (2009). The
775 merging of community ecology and phylogenetic biology. *Ecology Letters*, 12, 693-
776 715.
777
778 Clarke, J. T. (2021). Evidence for general size-by-habitat rules in actinopterygian fishes
779 across nine scales of observation. *Ecology Letters*, 24, 1569-1581.
780
781 Costa, G. C., Vitt, L. J., Pianka, E. R., Mesquita, D. O., & Colli, G. R. (2008). Optimal
782 foraging constrains macroecological patterns: body size and dietary niche breadth in
783 lizards. *Global Ecology and Biogeography*, 17, 670-677.
784
785 Cunningham, H. R., Rissler, L. J., Buckley, L. B., & Urban, M. C. (2015). Abiotic and
786 biotic constraints across reptile and amphibian ranges. *Ecography*, 38, 001-008.
787

788 Darwin, C. (1859). *The Origin of Species by Means of Natural Selection*. London:
789 Murray.
790

791 Davies, T. J., Meiri, S., Barraclough, T. G., & Gittleman, J. L. (2007). Species co-
792 existence and character divergence across carnivores. *Ecology Letters*, 10, 146-152.
793

794 Dobzhansky, T. (1950). Evolution in the tropics. *American Scientist*, 38, 208-221.
795

796 Foote, M. (2007). Symmetric waxing and waning of marine invertebrate genera.
797 *Paleobiology*, 33, 517-529.
798

799 Foster, J. B. (1964). Evolution of mammals on islands. *Nature*, 202, 234-235.
800

801 Friedman, S. T., Price, S. A., Corn, K. A., Larouche, O., Martinez, C. M., &
802 Wainwright, P.C. (2020). Body shape diversification along the benthic-pelagic axis in
803 marine fishes. *Proceedings of the Royal Society B*, 287, 20201053.
804

805 García-Navas, V., Kear, B. P., & Westerman, M. (2020). The geography of speciation
806 in dasyurid marsupials. *Journal of Biogeography*, 47, 2042-2053.
807

808 García-Porta, J., Morales, H. E., Gómez-Díaz, E., Sindaco, R., & Carranza, S. (2016).
809 Patterns of diversification in islands: A comparative study across three gecko genera in
810 the Socotra Archipelago. *Molecular Phylogenetics and Evolution*, 98, 288-299.
811

812 Gaston, K. J. (2000). Global patterns in biodiversity. *Nature*, 405, 220-227.
813

814 Gotelli, N. J., Graves, G. R., & Rahbek, C. (2010). Macroecological signals of species
815 interactions in the Danish avifauna. *Proceedings of the National Academy of Sciences*
816 *of the United States of America*, 107, 5030-5035.
817

818 Grant, P. R. (1972). Convergent and divergent character displacement. *Biological*
819 *Journal of the Linnean Society*, 4, 39-68.
820

821 Greene, H. W. (1997). Snakes: The evolution of mystery in nature. Berkeley: University
822 of California Press.
823

824 Grossenbacher, D. L., Veloz, S. D., & Sexton, J. P. (2014). Niche and range size
825 patterns suggest that speciation begins in small, ecologically diverged populations in
826 North American Monkeyflowers (*Mimulus* spp.). *Evolution*, 68, 1270-1280.
827

828 Hardin, G. (1960). The competitive exclusion principle. *Science*, 131, 1292-1297.
829

830 Henriques-Silva, R., Kubisch, A., & Peres-Neto, P. R. (2019). Latitudinal-diversity
831 gradients can be shaped by biotic processes: new insights from an eco-evolutionary
832 model. *Ecography*, 42, 259-271.
833

834 Hernández-Hernández, T., Miller, E. C., Román-Palacios, & Wiens, J. J. (2021).
835 Speciation across the tree of life. *Biological Reviews*, 96, 1205-1242.
836

837 Hijmans, R. J. (2019). raster: Geographic Data Analysis and Modeling. R package
838 version 3.4-13. <https://CRAN.R-project.org/package=raster>
839

840 Hutchinson, G. E. (1957). Concluding remarks. Cold springs Harbor Symposia on
841 Quantitative Biology, 22, 415-427.
842

843 Jackson, C. H. (2011). Multi-State Models for Panel Data: The msm Package for R.
844 *Journal of Statistical Software*, 38, 1-29.
845

846 Jackson, S. T., & Overpeck, J. T. (2000). Responses of plant populations and
847 communities to environmental changes of the late quaternary. *Paleobiology*, 26, 194-
848 220.
849

850 Johnson, M. T. J., & Stinchcombe, J. R. (2007). An emerging synthesis between
851 community ecology and evolutionary biology. *Trends in Ecology and Evolution*, 22,
852 250-257.
853

854 Jønsson, K. A., Tøttrup, A. P., Borregaard, M. K., Keith, S.A., Rahbek, C., & Thorup,
855 K. 2016. Tracking animal dispersal: from individual movement to community assembly
856 and global range dynamics. *Trends in Ecology and Evolution*, 31, 204-214.
857

858 Kozak, K. H., & Wiens, J. J. (2006). Does niche conservatism promote speciation? A
859 case study in North American Salamanders. *Evolution*, 60, 2604-2621.
860

861 Losos, J. B. (2011). *Lizards in an evolutionary tree: ecology and adaptive radiation of*
862 *Anoles*. Berkeley: University of California Press.
863

864 Losos, J. B., & Ricklefs, R.E. (2009). Adaptation and diversification on islands. *Nature*,
865 457, 830-836.
866

867 Louthan, A.M., Doak, D.F., & Angert, A.L. (2015). Where and when do species
868 interactions set range limits? *Trends in Ecology and Evolution*, 30, 780-792.
869

870 Lowe, W. H., & McPeck, M. A. (2014). Is dispersal neutral? *Trends in Ecology and*
871 *Evolution*, 29, 444-450.
872

873 MacArthur, R. H. (1969). Patterns and communities in the tropics. *Biological Journal of*
874 *the Linnean Society*, 1, 19-30.
875

876 MacArthur, R. H., & Wilson, E. O. (1967). *The Theory of Island Biogeography*.
877 Princeton: Princeton University Press.
878

879 Meiri, S. (2007). Size evolution in island lizards. *Global Ecology and Biogeography*,
880 16, 702-708.
881

882 Meiri, S. (2008). Evolution and ecology of lizard body sizes. *Global Ecology and*
883 *Biogeography*, 17, 724-734.
884

885 Meiri, S., Simberloff, D., & Dayan, T. (2011). Community-wide character displacement
886 in the presence of clines: A test of Holarctic weasel guilds. *Journal of Animal Ecology*,
887 80, 824-834.

888

889 Novosolov, M., Rodda, G. H., Feldman, A., Kadison, A. E., Dor, R., & Meiri, S.

890 (2015). Power in numbers. Drivers of high population density in insular lizards. *Global*

891 *Ecology and Biogeography*, 25, 87-95.

892

893 Novosolov, M., Rodda, G. H., Gainsbury, A. M., & Meiri, S. (2018). Dietary niche

894 variation and its relationship to lizard population density. *Journal of Animal Ecology*,

895 87, 285-292.

896

897 Peters, R. H. (1986). The ecological implications of body size. Cambridge: Cambridge

898 University Press.

899

900 Pianka, E. R., & Vitt, L. J. (2003). *Lizards: Windows to the evolution of diversity*.

901 Berkeley: University of California Press.

902

903 Pigot, A. L., Jetz, W., Sheard, C., & Tobias, J. A. (2018). The macroecological

904 dynamics of species coexistence in birds. *Nature Ecology & Evolution*, 2, 1112-1119.

905

906 Pigot, A. L., & Tobias, J. A. (2013). Species interactions constrain geographic range

907 expansion over evolutionary time. *Ecology Letters*, 16, 330-338.

908

909 Pigot, A. L., & Tobias, J. A. (2015). Dispersal and the transition to sympatry in

910 vertebrates. *Proceedings of the Royal Society B*, 282, 20141929.

911

912 Pigot, A. L., Tobias, J. A., & Jetz, W. (2016). Energetic constraints on species

913 coexistence in birds. *Plos Biology*, 14, e1002407.

914

915 Pitta, E., Kassara, C., Giokas, S., & Sfenthourakis, S. (2013). Compositional

916 dissimilarity patterns of reptiles and amphibians in insular systems around the world.

917 *Ecological Research*, 28, 633-642.

918

919 Pough, F.H. (1973). Lizard energetics and diet. *Ecology*, 54, 837-844.

920

921 Powney, G. D., Grenyer, R., Orme, C. D. L., Owens, I. P. F., & Meiri, S. (2010). Hot,
922 dry and different: Australian lizard richness is unlike that of mammals, amphibians and
923 birds. *Global Ecology and Biogeography*, 19, 386-396.
924
925 Price, S. A., Friedman, S. T., Corn, K. A., Martinez, C. M., Larouche, O., &
926 Wainwright, P. C. (2019). Building a body shape morphospace of Teleostean fishes.
927 *Integrative and Comparative Biology*, 59, 716-730.
928
929 Revell, L. J. (2009). Size-correction and principal components for interspecific
930 comparative studies. *Evolution*, 63, 3258-3268.
931
932 Revell, L. J. (2012). Phytools: an R package for phylogenetic comparative biology (and
933 other things). *Methods in Ecology and Evolution*, 3, 217-223.
934
935 Ricklefs, R. E. (2004). A comprehensive framework for global patterns in biodiversity.
936 *Ecology Letters*, 7, 1-15.
937
938 Ricklefs, R. E. (2006). Evolutionary diversification and the origin of the diversity-
939 environment relationship. *Ecology*, 87, S3-S13.
940
941 Ricklefs, R. E. (2010). Host–pathogen coevolution, secondary sympatry and species
942 diversification. *Philosophical Transaction of the Royal Society of London B*, 365, 1139-
943 1147.
944
945 Ricklefs, R. E., & Miles, D. B. (1994). Ecological and evolutionary inferences from
946 morphology: an ecological perspective. In: Wainwright, P. C., Reilly, S. M. (Eds.),
947 *Ecological morphology: Integrative organismal biology* (pp. 13-41). Chicago:
948 University of Chicago Press.
949
950 Roll, U., Feldman, A., Novosolov, M., Allison, A., Bauer, A. M., Bernard, R., Böhm,
951 M., et al. (2017). The global distribution of tetrapods reveals a need for targeted reptile
952 conservation. *Nature Ecology & Evolution*, 1, 1677-1682.
953

954 Schluter, D. (2000). Ecological character displacement in adaptive radiations. *American*
955 *Naturalist*, 156, S4-S16.
956
957 Sexton, J. P., McIntyre, P. J., Angert, A. L., & Rice, K. J. (2009). Evolution and
958 ecology of species range limits. *Annual Review of Ecology, Evolution and Systematics*,
959 40, 415-436.
960
961 Shi, J. J., Westeen, E. P., Katlein, N. T., Dumont, E. R., & Rabosky, D. L. (2018).
962 Ecomorphological and phylogenetic controls on sympatry across extant bats. *Journal of*
963 *Biogeography*, 45, 1560-1570.
964
965 Silvestro, D., Antonelli, A., Salamin, N., & Quental, T. B. (2015). The role of clade
966 competition in the diversification of North American canids. *Proceedings of the*
967 *National Academy of Sciences of the United States of America*, 112, 8684-8689.
968
969 Sites, J. W., Reeder, T. W., & Wiens, J. J. 2011. Phylogenetic insights on evolutionary
970 novelties in lizards and snakes: sex, birth, bodies, niches and venom. *Annual Review of*
971 *Ecology, Evolution and Systematics*, 42, 227-244.
972
973 Skeels, A., & Cardillo, M. (2019). Reconstructing geography of speciation from
974 contemporary biodiversity data. *The American Naturalist*, 193, 240-255.
975
976 Skeels, A., Esquerré, D., & Cardillo, M. (2020). Alternative pathways to diversity
977 across ecologically distinct lizard radiations. *Global Ecology and Biogeography*, 29,
978 454-469.
979
980 Slavenko, A., Allison, A., & Meiri, S. (2021). Elevation is a stronger predictor of
981 morphological trait divergence than competition in a radiation of tropical lizards.
982 *Journal of Animal Ecology*, 90, 917-930.
983
984 Slavenko, A., Feldman, A., Allison, A., Bauer, A. M., Böhm, M., Chirio, L., Colli, G.
985 R. et al. (2019). Global patterns of body size evolution in squamate reptiles are not
986 driven by climate. *Global Ecology and Biogeography*, 28, 471-483.
987

988 Stuart, Y. E., & Losos, J. B. (2013). Ecological character displacement: glass half full or
989 half empty. *Trends in Ecology and Evolution*, 28, 402-408.
990

991 Tonini, J. F. R., Beard, K. H., Ferreira, R. B., Jetz, W., & Pyron, R. A. (2016). Fully-
992 sampled phylogenies of squamates reveal evolutionary patterns in threat status.
993 *Biological Conservation*, 204, 23-31.
994

995 Vamosi, S. M., Heard, S. B., Vamosi, J. C., & Webb, C. O. (2009). Emerging patterns
996 in the comparative analysis of phylogenetic community structure. *Molecular Ecology*,
997 18, 572-592.
998

999 Van Valen, L. (1973). A new evolutionary law. *Evolutionary Theory*, 1, 1-30.
1000

1001 Vitt, L. J., & Caldwell, J. P. (2014). *Herpetology*. Cambridge: Academic Press.
1002

1003 Uetz, P. & Hosek, J. (2017). The reptile database. <http://www.reptile-database.org>.
1004

1005 Webb, C. O., Ackerly, D. D., McPeck, M. A., & Donoghue, M. J. (2002). Phylogenies
1006 and community ecology. *Annual Review of Ecology and Systematics*, 33, 475-505.
1007

1008 Weber, M. G., & Strauss, S. Y. (2016). Coexistence in close relatives: beyond
1009 competition and reproductive isolation in sister taxa. *Annual Review of Ecology*,
1010 *Evolution and Systematics*, 47, 359-381.
1011

1012 Weir J. T., & Price T. D. (2011). Limits to Speciation Inferred from Times to Secondary
1013 Sympatry and Ages of Hybridizing Species along a Latitudinal Gradient. *The American*
1014 *Naturalist* 177:462-469.
1015

1016 White, A. E. (2016). Geographical Barriers and Dispersal Propensity Interact to Limit
1017 Range Expansions of Himalayan Birds. *The American Naturalist*, 188, 99-112.
1018

1019 Whiting, E. T., & Fox, D. L. (2020). Latitudinal and environmental patterns of species
1020 richness in lizards and snakes across continental North America. *Journal of*
1021 *Biogeography*, 48, 291-304.

1022

1023 Wiens, J. J., & Donoghue, M. J. (2004). Historical biogeography, ecology and species
1024 richness. *Trends in Ecology and Evolution*, 19, 639-644.

1025

1026 Wilson, D. S. (1975). The adequacy of body size as a niche difference. *The American*
1027 *Naturalist*, 109, 769-784.

1028

1029 Wisz, M. S., Pottier, J., Kissling, D.K., Pellisier, L., Lenoir, J., Damgaard, C. F., &
1030 Dormann, C. F. (2013). The role of biotic interactions in shaping distributions and
1031 realised assemblages of species: implications for species distribution modelling.
1032 *Biological Reviews*, 88, 15-30.

1033 **DATA AVAILABILITY**

1034 Data used in this study are part of the supplementary material or have been published
1035 elsewhere.

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

FIGURE LEGENDS

1059

1060 **Figure 1.** Species diversity analyzed in the present study across geographical regions (a,

1061 c) and the taxonomic diversity of the species pairs (b) of lizards and snakes included in

1062 our analyses.

1063

1064

1065 **Figure 2.** Summary of the statistical models compared in the present study and their
 1066 parameters. γ = relative frequency of allopatric speciation, σ = transition rate from
 1067 allopatry to sympatry, ϵ = transition rate from sympatry to allopatry. We included
 1068 ecological similarity (i.e. body size and shape differences), occurrence on islands or
 1069 continents, and the dispersion ability (i.e. range size/age ratio and mean body size)
 1070 separately as covariates on the transition rate to sympatry. Modified from Pigot &
 1071 Tobias (2015).

1072

1073

1074 **Figure 3.** Estimates of transitions rates to sympatry relative to (a) species pairs
1075 occurrence on islands or continents, (b) snout-vent length difference, and (c) dispersal
1076 ability (range size / age ratio and mean snout-vent length), under the best model selected
1077 for snakes (a, b) and lizards (c, d). Multiple dots and lines represent estimates when
1078 using distinct age datasets (see Methods). Dashed lines in panel (a) represent medians of
1079 transition rates taken across all estimates. Given the extremely high values of 12 of
1080 these estimates (1 continent and 11 island), we excluded these from the plot for
1081 visualization purposes.

1082

1083

1084