Compartment-specific energy requirements of photosynthetic carbon metabolism in *Camelina sativa* leaves

Thomas Wieloch^{a,*}, Thomas David Sharkey^{b,c,d}

^aDepartment of Medical Biochemistry and Biophysics, KB. H6, Umeå University, 901 87 Umeå, Sweden

^bDepartment of Energy-Plant Research Laboratory, Michigan State University, East Lansing, MI 48824, USA

^cDepartment of Biochemistry and Molecular Biology, Michigan State University, East Lansing, MI 48824, USA

^dPlant Resilience Institute, Michigan State University, East Lansing, MI 48824, USA

*Corresponding author: Thomas Wieloch, thomas.wieloch@umu.se, +46706460369

Author	ORCID
Thomas Wieloch	0000-0001-9162-2291
Thomas David Sharkey	0000-0002-4423-3223

Author Contributions: T.W., and T.D.S. designed the research. T.W. led the research. T.W. analysed and interpreted the data and wrote the manuscript with input from T.D.S.

Data Availability: All data generated or analysed during this study are included in this published article.

Acknowledgments: T.D.S. was supported by the Division of Chemical Sciences, Geosciences and Biosciences, Office of Basic Energy Sciences of the United States Department of Energy Grant DE-FG02-91ER20021. T.D.S. receives partial salary support from MSU AgBioResearch. However, the contents of this paper are solely the responsibility of the authors.

Main Conclusion

The oxidative pentose phosphate pathway provides cytosolic NADPH yet reduces carbon and energy use efficiency. Repressing this pathway and introducing cytosolic NADPH-dependent malate dehydrogenase may increase crop yields by $\approx 5\%$.

Abstract

Detailed knowledge about plant energy metabolism may aid crop improvements. Using published estimates of flux through central carbon metabolism, we phenotype energy metabolism in illuminated *Camelina sativa* leaves (grown at 22 °C, 500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) and report several findings. First, the oxidative pentose phosphate pathway (OPPP) transfers 3.3% of the NADPH consumed in the Calvin-Benson cycle to the cytosol. However, concomitantly respired CO_2 accounts for 4.8% of total rubisco activity. Hence, 4.8% of the flux through the Calvin-Benson cycle and photorespiration is spent on supplying cytosolic NADPH, a significant investment. Associated energy requirements exceed the energy output of the OPPP. Thus, autotrophic carbon metabolism is not simply optimised for flux into carbon sinks but sacrifices carbon and energy use efficiency to support cytosolic energy metabolism. To reduce these costs, we suggest bioengineering plants with a repressed cytosolic OPPP, and an inserted cytosolic NADPH-dependent malate dehydrogenase tuned to compensate for the loss in OPPP activity (if required). Second, sucrose cycling is a minor investment in overall leaf energy metabolism but a significant investment in cytosolic energy metabolism. Third, energy balancing strictly requires oxidative phosphorylation, cofactor export from chloroplasts, and peroxisomal NADH import. Fourth, mitochondria are energetically self-sufficient. Fifth, carbon metabolism has an ATP/NADPH demand ratio of 1.52 which is met if $\leq 21.7\%$ of whole electron flux is cyclic. Sixth, electron transport has a photon use efficiency of $\geq 62\%$. Lastly, we discuss interactions between the OPPP and the cytosolic oxidation-reduction cycle in supplying leaf cytosolic NADPH.

Keywords: ATP/NADPH ratio, bioenergetics, energy metabolism, glucose-6-phosphate shunt, oxidative pentose phosphate pathway, sucrose cycling

Introduction

Central carbon metabolism operates co-ordinately across several cell compartments and requires compartment-specific inputs of reductant and energy cofactors (NAD(P)H, Fd_{red}, ATP, UTP). These requirements vary with ambient and developmental conditions (Scheibe 2019). For instance, drought frequently causes stomatal closure and low intercellular CO₂ concentrations which promote photorespiratory NADH production in mitochondria and consumption in peroxisomes (see below). Since cofactor production is spatially restricted, transmembrane transport and conversion of cofactors are key to metabolic functioning.

In illuminated leaves, reductant and energy cofactors are primarily produced by the light reactions of photosynthesis inside chloroplasts. Additionally, the pyruvate dehydrogenase complex (PDC) in fatty acid biosynthesis supplies chloroplastic NADH (Fig. 1). In the cytosol, glucose-6-phosphate dehydrogenase (G6PD) and 6-phosphogluconate dehydrogenase (6PGD) in the oxidative pentose phosphate pathway (OPPP), non-phosphorylating glyceraldehyde-3-phosphate dehydrogenase (GAPN) in glycolysis, and isocitrate dehydrogenase (IDH) in amino acid biosynthesis supply NADPH while phosphorylating glyceraldehyde-3-phosphate dehydrogenase (GAPC), phosphoglycerate kinase (PGK), and pyruvate kinase (PK) in glycolysis supply NADH and ATP. In mitochondria, the photorespiratory glycine decarboxylase complex (GDC) and several enzymes of the tricarboxylic acid cycle including PDC, and IDH supply NADH.

On the other hand, numerous enzyme reactions spread across several cell compartments consume reductant and energy cofactors. In chloroplasts, PGK, phosphorylating glyceraldehyde-3-phosphate dehydrogenase (GAPDH), and phosphoribulokinase (PRK) in the Calvin-Benson cycle (CBC) consume NADPH, and ATP. Furthermore, glutamine α -ketoglutarate aminotransferase (GOGAT), and glycerate kinase (GK) in photorespiration, ADP-glucose pyrophosphorylase (AGPase) in starch biosynthesis, and acetyl-CoA carboxylase (ACC), 3-ketoacyl-ACP reductase (KAR), and 2,3-trans-enoyl-ACP reductase (ACPr) in fatty acid biosynthesis consume chloroplastic NAD(P)H, Fd_{red}, and ATP. In the cytosol, UDP-glucose pyrophosphorylase (UGPase) in sucrose biosynthesis, and hexokinase (HK), and fructokinase (FK) in sucrose cycling consume UTP, and ATP. In mitochondria, glutamine synthetase (GS) in photorespiration, and

glutamate dehydrogenase (GDH) in amino acid biosynthesis consume ATP, and NADH while photorespiratory hydroxypyruvate reductase (HPR) consumes NADH in peroxisomes.

Several mechanisms enable transmembrane transport of cofactors from sites of production to sites of consumption (Fig. 2). NAD⁺ carrier proteins enable direct transport of NAD(P)H across the inner membranes of chloroplasts and mitochondria but have low affinities for these reductants (Palmieri et al. 2009). By contrast, dicarboxylate transporters coupled with malate dehydrogenase enzymes (malate valves) enable indirect exchange of NAD(P)H among chloroplasts, peroxisomes, mitochondria, and the cytosol (Selinski and Scheibe 2019). While there is an NADPH-dependent malate dehydrogenase inside chloroplasts, most malate dehydrogenase provide NADH, not NADPH. By contrast, glyceraldehyde 3-phosphate/3-phosphoglycerate (GAP/3PGA) cycling enables NADPH transfer from chloroplasts to the cytosol (Kelly and Gibbs 1973b, a). In this cycle, chloroplastic PK and GAPDH reduce 3PGA to GAP consuming ATP and NADPH. Subsequently, GAP is exported to the cytosol where GAPN is reducing it to 3PGA producing NADPH. A similar cycle involving GAPC and PK (instead of GAPN) can provide cytosolic NADH and ATP at equimolar amounts (Stocking and Larson 1969). Additional cytosolic ATP can come from oxidative phosphorylation through the ADP/ATP carrier (Klingenberg 2008).

Furthermore, several mechanisms enable cofactor conversion. In the cytosol, equimolar amounts of ATP, and NADH can be converted to NADPH by the carbon-neutral cytosolic oxidation-reduction (COR) cycle (Figs. 1-2; Wieloch 2021). In its forward direction, cytosolic GAPN oxidises GAP to 3PGA producing NADPH. In its backward direction, cytosolic PGK and GAPC reduce 3PGA to GAP consuming ATP, and NADH. COR cycling may help balancing the cytosolic NADPH supply according to varying demands. For instance, we recently reported evidence for increased COR cycling under drought where increased synthesis of reactive oxygen species may require increased amounts of cytosolic NADPH to maintain the cellular redox balance (Wieloch et al. 2021). In mitochondria, NADH can be converted to ATP by oxidative phosphorylation. Similarly, cytosolic NAD(P)H can be converted to mitochondrial ATP by oxidative phosphorylation starting not at complex I but at type II NAD(P)H dehydrogenases or glycerol-3-phosphate dehydrogenase both located in the inner mitochondrial membrane facing the cytosol (Rasmusson et al. 2008).

In a recently published study, we estimated carbon fluxes across central metabolism in illuminated *Camelina sativa* leaves, including flux through the CBC, photorespiration, glycolysis, the tricarboxylic acid cycle, and starch, sucrose, fatty acid, and amino acid biosynthesis (Xu et al. in prep.). Importantly, our analyses explained the ^{13}C labelling lag of CBC metabolites (a longstanding conundrum in plant biochemistry) by reinjection of weakly labelled carbon from vacuolar and cytosolic sugar pools into the CBC. This process involves breakdown of sucrose into fructose and glucose, ATP-dependent re-phosphorylation to fructose 6-phosphate and glucose 6-phosphate, UTP-dependent sucrose resynthesis from fructose 6-phosphate and glucose 6-phosphate and reinjection of carbon from cytosolic glucose 6-phosphate into the CBC via the NADPH-producing OPPP (Fig. 1). Additionally, assimilation of the CO_2 released by the OPPP involves all cofactor-dependent reactions in the CBC and photorespiration (ATP, NAD(P)H, Fd_{red}). Therefore, we hypothesise sucrose cycling and OPPP-mediated reinjection of cytosolic carbon into the CBC significantly affect plant energy metabolism.

To test this hypothesis, we comprehensively characterise energy metabolism in illuminated *Camelina sativa* leaves. More specifically, we estimate compartment-specific cofactor demands of sucrose cycling and the OPPP and discuss associated effects on plant carbon and energy metabolism. Additionally, we estimate compartment-specific cofactor demands of central carbon metabolism and address the following questions (with respect to the conditions and set of reactions studied here):

- Is cofactor export from chloroplasts strictly required?
- Are mitochondria energetically self-sufficient?
- Does photorespiration have the same ATP/NADPH demand ratio as the CBC?
- How much cyclic electron flux is required to meet the cofactor demands of carbon metabolism?
- Where is the cytosolic ATP coming from?
- Is oxidative phosphorylation strictly required? (Note that oxidative phosphorylation does not strictly require tricarboxylic acid cycling.)
- Is mitochondrial NADH strictly required to balance peroxisomal NADH demands?

- Where is the cytosolic NADPH coming from?

Material and methods

Data

Previously, we estimated carbon fluxes across central metabolism in 4-week-old source leaves of wild-type *Camelina sativa* ecotype Suneson by ^{13}C isotopically non-stationary metabolic flux analysis (Xu et al. in prep.). These plants were grown at 22 °C, 50% relative humidity, and a light intensity of 500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ (8/16 h, day/night). Here, we utilise the carbon flux estimates to estimate associated cofactor fluxes (Fig. 1).

Cofactor demand by carbon metabolism

We resolved chloroplastic (.p), cytosolic (.c), mitochondrial (.m), and peroxisomal (.ox) fluxes. Within the Calvin-Benson cycle, we considered reactions catalysed by PGK.p (-1 ATP per turnover), GAPDH.p (-1 NAD(P)H per turnover), and PRK.p (-1 ATP per turnover). Within photorespiration, we considered reactions catalysed by GS.m (-1 ATP per turnover), GOGAT.p (-2 Fd_{red} per turnover), GDC.m (+1 NADH per turnover), HPR.ox (-1 NADH per turnover), and GK.p (-1 ATP per turnover). It is commonly believed that photorespiratory ammonium is recaptured by GS2. While this enzyme is located in both chloroplasts and mitochondria (Taira et al. 2004), we here only considered the latter isoform since it has been suggested to immediately recapture potentially harmful ammonium released by GDC.m (Buchanan et al. 2015). Furthermore, while there is NADPH- and Fd_{red} -dependent GOGAT.p, only the latter is essential for photorespiration and, therefore, considered here (Somerville and Ogren 1980). Within the OPPP, we considered reactions catalysed by G6PD.c (+1 NADPH per turnover), and 6PGD.c (+1 NADPH per turnover). Within starch and sucrose biosynthesis and sucrose cycling, we considered reactions catalysed by AGPase.p (-1 ATP per turnover), UGPase.c (-1 UTP per turnover), HK.c (-1 ATP per turnover), and FK.c (-1 ATP per turnover). Within glycolysis, we considered the reaction catalysed by PK.c (+1 ATP per turnover). Within the tricarboxylic acid cycle and amino acid biosynthesis, we only considered reactions catalysed by PDC.m (+1 NADH per turnover), IDH.m (+1 NADH per turnover), and GDH.m (-1 NADH per turnover) because other cofactor-dependent reactions reportedly carry negligible flux (Xu et al. in prep.). Within fatty acid biosynthesis, we considered reactions catalysed by PDC.p (+1 NADH per turnover), ACC.p (-1

ATP per turnover), KAR.p (-1 NADPH per turnover), and ACPr.p (-1 NADH per turnover). Our accounting makes three assumptions. First, all pyruvate entering fatty acid biosynthesis comes from the β -elimination of phosphate from a carbocation intermediate of ribulose 1,5-bisphosphate (RuBP) carboxylation by rubisco (Andrews and Kane 1991). At 25 °C, the ratio of rubisco pyruvate synthesis to carboxylation is 0.7% (Andrews and Kane 1991). Xu et al. (in prep.) reported carboxylation proceeds at a rate of 172.1 $\mu\text{mol RuBP g}^{-1} \text{FW h}^{-1}$ corresponding to $\approx 1.2 \mu\text{mol g}^{-1} \text{FW h}^{-1}$ pyruvate synthesis. Thus, pyruvate provided by rubisco significantly exceeds pyruvate requirements by fatty acid biosynthesis (0.4 $\mu\text{mol pyruvate g}^{-1} \text{FW h}^{-1}$). Second, since the major isoform of ACPr.p is NADH-dependent and a minor isoform works with both NADH and NADPH (Buchanan et al. 2015), our accounting only considers NADH as cofactor of ACPr.p. Third, only C₁₆ fatty acid are being produced. Initiation of C₁₆ fatty acid biosynthesis requires 2 molecules pyruvate, 1 ATP, and 1 NADPH, and yields 1 C₄ fatty acid, and 1 NADH. Elongation to a C₁₆ fatty acid requires another 6 molecules pyruvate, 6 ATP, and 6 NADPH. Overall, biosynthesis of a C₁₆ fatty acid requires 8 molecules pyruvate, 7 ATP, 7 NADPH and yields 1 NADH.

The cytosolic OPPP forms a shunt bypassing much of the CBC and releases CO₂ from metabolism. We have calculated rubisco carboxylation and oxygenation rates accounting for the fixation of this CO₂ as

$$v_c(\text{OPPP}) = \frac{v_{\text{OPPP}}}{1 - 0.5 v_o/v_c} \quad (1)$$

and

$$v_o(\text{OPPP}) = \frac{v_o}{v_c} * v_c(\text{OPPP}) \quad (2)$$

where v_{OPPP} denotes flux through the cytosolic OPPP, v_c and v_o denote overall rubisco carboxylation and oxygenation rates, respectively, and $v_c(\text{OPPP})$ and $v_o(\text{OPPP})$ denote rubisco carboxylation and oxygenation rates attributable to the CO₂ lost in the OPPP, respectively.

Results and discussion

Impacts of the cytosolic OPPP on plant carbon and energy metabolism

Flux through the cytosolic OPPP is likely a general feature of C_3 metabolism since it can explain labelling lags of CBC metabolites (Xu et al. in prep.) observed in several species fed $^{13}\text{CO}_2$ or $^{14}\text{CO}_2$ (Mahon et al. 1974; Canvin 1979; Hasunuma et al. 2010; Szecowka et al. 2013; Ma et al. 2014; Sharkey et al. 2020; Xu et al. 2021). In illuminated *Camelina sativa* leaves, the cytosolic OPPP supplies $14\ \mu\text{mol NADPH g}^{-1}\text{ FW h}^{-1}$ to the cytosol from the chloroplast, corresponding to 3.3% of the NAD(P)H consumed in the CBC (Tables 1-2). This NADPH may support cytosolic redox, signalling, and biosynthetic metabolism and mitochondrial ATP production (Rasmusson et al. 2008).

Cytosolic NADPH production by the OPPP is associated with the release of $7\ \mu\text{mol CO}_2\text{ g}^{-1}\text{ FW h}^{-1}$ from metabolism (Table S1). Fixation of this CO_2 accounts for 4.8% of the total activity of rubisco. That is, 4.8% of the total flux through the CBC and photorespiration is spent on providing cytosolic NADPH, a significant investment. Associated energy requirements ($20\ \mu\text{mol NADPH g}^{-1}\text{ FW h}^{-1}$, $2.4\ \mu\text{mol Fd}_{\text{red}}\text{ g}^{-1}\text{ FW h}^{-1}$, $33\ \mu\text{mol ATP g}^{-1}\text{ FW h}^{-1}$) exceed the energy output of the OPPP ($14\ \mu\text{mol NADPH g}^{-1}\text{ FW h}^{-1}$; Table 1). These carbon and energy costs would increase with increasing photorespiration. Thus, carbon metabolism in illuminated autotrophic tissue is not simply optimised for flux into carbon sinks but sacrifices both carbon and energy use efficiency to support cytosolic energy metabolism.

To reduce costs related to OPPP respiration, we suggest bioengineering plants with a repressed cytosolic OPPP, and a newly introduced cytosolic NADPH-dependent malate dehydrogenase finely tuned to compensate for the loss in OPPP activity. This latter addition will only be required under conditions where carbon-neutral GAP/3PGA cycles and the COR cycle fall short of compensating for the reduction in NADPH from the OPPP.

Impact of sucrose cycling on plant energy metabolism

Sucrose cycling involving breakdown of sucrose into fructose and glucose, ATP-dependent re-phosphorylation to fructose 6-phosphate and glucose 6-phosphate, and UTP-dependent sucrose resynthesis from fructose 6-phosphate and glucose 6-phosphate is another futile carbon cycle (Fig.

1). In the cytosol of illuminated *Camelina sativa* leaves, its operation requires $6.5 \mu\text{mol ATP g}^{-1} \text{FW h}^{-1}$ (Table 1; phosphorylation of 1 UDP to UTP consumes 1 ATP). This corresponds to 1% and 53% of the ATP requirement in chloroplasts and the cytosol, respectively (Table 2). Thus, sucrose cycling is a minor investment in overall leaf energy metabolism but a significant investment in cytosolic energy metabolism. However, it should be noted that several processes consuming cytosolic ATP are not considered here including cytosol-vacuole proton pumping (Hedrich et al. 2015).

Cofactor fluxes in central carbon metabolism

Table 2 lists cofactor consumption and production by central enzyme reactions in carbon metabolism of *Camelina sativa* leaves. Figure 2 summarises this information for specific cell compartments. Chloroplast metabolism consumes large amounts of ATP and NADPH and some Fd_{red} while producing small amounts of NADH. Cytosol metabolism consumes ATP and UTP while producing NADPH. If triose phosphates were the only cytosolic carbon substrates (i.e., if 3PGA was not exported) then there would be an additional supply of $2.15 (+0.19, -0.16) \mu\text{mol g}^{-1} \text{FW h}^{-1}$ of NADH and ATP by GAPC and PGK or NADPH by GAPN (Fig. 1, grey reactions). However, this scenario is not considered here. Peroxisome metabolism consumes NADH, and mitochondrion metabolism consumes ATP while producing NADH. Compared to other cell compartments, absolute energy cofactor fluxes (ATP, UTP) in chloroplast are 26.7-times higher while absolute reductant cofactor fluxes (NAD(P)H, Fd_{red}) are 17.9-times higher.

Is cofactor export from chloroplasts strictly required?

Cofactor export from chloroplasts is mediated by the malate valve and GAP/3PGA cycles (Fig. 2). To investigate whether these processes are strictly required, we will now illustrate how cofactors produced by extra-chloroplastic carbon metabolism are used most efficiently to meet extra-chloroplastic cofactor demands. In this theoretical scenario, the mitochondrial NADH surplus ($26.4 \mu\text{mol g}^{-1} \text{FW h}^{-1}$) is used to balance the peroxisomal NADH deficit ($25.4 \mu\text{mol g}^{-1} \text{FW h}^{-1}$) through NADH transmembrane transport by malate valves (Fig. 2). Assuming a P/O ratio of 2.5 (Hinkle 2005), remaining mitochondrial NADH ($1.03 \mu\text{mol g}^{-1} \text{FW h}^{-1}$) is converted to $2.6 \mu\text{mol ATP g}^{-1} \text{FW h}^{-1}$ via oxidative phosphorylation starting at complex I of the electron transport chain. Remaining mitochondrial and cytosolic ATP and UTP deficits equivalent to $34.1 \mu\text{mol ATP g}^{-1}$

FW h⁻¹ are met by transmitting electrons from cytosolic NADPH onto the mitochondrial electron transport chain via type II NAD(P)H dehydrogenases. This reduces ATP yields by about 30% compared to transmission via complex I (Rasmusson et al. 2008). That is, the cytosolic surplus of 14 μmol NADPH g⁻¹ FW h⁻¹ is converted to 24.4 μmol ATP g⁻¹ FW h⁻¹ in mitochondria which is partly exported to the cytosol via the ADP/ATP carrier (Klingenberg 2008). Overall, there remains an extra-chloroplastic deficit of 9.7 μmol ATP g⁻¹ FW h⁻¹. Thus, with respect to the conditions and set of reactions studied here, cofactor export from chloroplasts is strictly required to meet all extra-chloroplastic cofactor demands.

Are mitochondria energetically self-sufficient?

It may be most realistic to assume that the mitochondrial surplus of 26.4 μmol NADH g⁻¹ FW h⁻¹ is used first to balance the mitochondrial deficit of 25.5 μmol ATP g⁻¹ FW h⁻¹ rather than being directly shuttled out of mitochondria into peroxisomes (Fig. 2). To meet the mitochondrial ATP deficit by oxidative phosphorylation starting from complex I requires only 10.2 μmol NADH g⁻¹ FW h⁻¹. Thus, with respect to the conditions and set of reactions studied here, mitochondria are energetically self-sufficient.

Does photorespiration have the same ATP/NADPH demand ratio as the CBC?

Our cofactor accounting assumes that photorespiratory ammonium is recaptured by mitochondrial (not chloroplastic) GS. This has interesting consequences. Chloroplastic cofactor requirements upon RuBP carboxylation comprise 4 ATP by PGK, 4 NADPH by GAPDH, and 2 ATP by PRK (Fig. 1). Chloroplastic cofactor requirements upon RuBP oxygenation comprise 3 ATP by PGK, 3 NADPH by GAPDH, 2 ATP by PRK, 1 ATP by GK, and 2 Fd_{red} corresponding to 1 NADPH by GOGAT. Thus, overall, both processes have the same absolute chloroplastic cofactor requirements with an ATP/NADPH demand ratio of 1.5. Other processes, such as starch biosynthesis and serine withdrawal from photorespiration, will merely cause small changes of this ratio (see below).

How much cyclic electron flux is required to meet the cofactor demands of carbon metabolism?

Chloroplast carbon metabolism has much higher demands for reductant and energy cofactors than any other cell compartment (Fig. 2). These demands are readily satisfied by cofactor input mainly from the light reactions of photosynthesis without the need for transmembrane transport or

conversion of cofactors. Interestingly, chloroplast GAPDH accepts both NADPH and NADH (McGowan and Gibbs 1974). *In vivo*, however, NADPH supply from the light reactions is 7630-times higher than the NADH surplus from fatty acid biosynthesis (Table 2).

Overall, chloroplast metabolism requires $680 \mu\text{mol ATP g}^{-1} \text{FW h}^{-1}$ and $446 \mu\text{mol NAD(P)H g}^{-1} \text{FW h}^{-1}$ (Table 2). This accounts for Fd_{red} demands where 2 Fd_{red} correspond to 1 NADPH. Hence, the reaction network and conditions studied here have an ATP/NADPH demand ratio of 1.52. However, linear electron flux including Q cycling has an estimated ATP/NADPH supply ratio of 1.286 (assuming trans-membrane transport of 3 protons per electron and production of 3 ATP per 14 protons by ATP synthase) (Seelert et al. 2000). Thus, assuming neither the water-water cycle (affecting the ATP/NADPH supply ratio) nor GAP/3PGA cycles and the chloroplast malate valve (affecting the ATP/NADPH demand ratio) are operational, cyclic electron flux through photosystem I is required.

Production of $446 \mu\text{mol NADPH g}^{-1} \text{FW h}^{-1}$ requires linear electron flux at a rate of $892 \mu\text{mol g}^{-1} \text{FW h}^{-1}$ and yields $574 \mu\text{mol ATP g}^{-1} \text{FW h}^{-1}$. To meet the remaining deficit of $106 \mu\text{mol ATP g}^{-1} \text{FW h}^{-1}$, cyclic electron flux through photosystem I including Q cycling at a rate of $248 \mu\text{mol g}^{-1} \text{FW h}^{-1}$ is required (assuming trans-membrane transport of 2 protons per electron and production of 3 ATP per 14 protons by ATP synthase) (Seelert et al. 2000). Thus, with respect to the conditions and set of reactions studied here, cyclic electron flux accounts for 21.7% of whole electron flux and is a 27.8% fraction of linear electron flux.

Jointly, linear, and cyclic electron flux through photosystem I consume $2032 \mu\text{mol photons g}^{-1} \text{FW h}^{-1}$. Carbon flux in *Camelina sativa* leaves was measured at $500 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ corresponding to $3273 \mu\text{mol photons g}^{-1} \text{FW h}^{-1}$ (the fresh weight per leaf area for a typical *Camelina* leaf is $0.055 \pm 0.0041 \text{ g cm}^{-2}$ implying a leaf thickness of 0.55 mm). Thus, with respect to the conditions and set of reactions studied here, 62% of the photons falling on the leaf are absorbed and used for photosynthesis.

In our previous analysis (Xu et al. in prep.), carbon fluxes through GAP/3PGA cycles, the COR cycle (grey in Fig. 1), and the chloroplast malate valve were not estimated, i.e., associated cofactor

fluxes are unknown. These processes would, however, decrease the ATP/NADPH demand ratio of carbon metabolism and, thus, the requirement for cyclic electron flux. This is because GAP/3PGA cycles require chloroplastic ATP and NAD(P)H at a 1:1 ratio (Stocking and Larson 1969; Kelly and Gibbs 1973a), and the malate valve requires chloroplastic NAD(P)H only (Selinski and Scheibe 2019). COR cycling requires NADH and ATP in the cytosol which may be provided by combined action of the chloroplast malate valve and oxidative phosphorylation (Wieloch 2021). Furthermore, these processes would consume additional electrons and, thus, increase the photon use efficiency of electron transport. For example, no cyclic electron flux is required if $82.5 \mu\text{mol NAD(P)H g}^{-1} \text{FW h}^{-1}$ (corresponds to $\approx 60\%$ of net CO_2 assimilation) are exported by the chloroplast malate valve. This additional flux would result in 64.6% photon use efficiency of electron transport.

Where is the cytosolic ATP coming from? Is oxidative phosphorylation strictly required?

Overall, we estimated an extra-chloroplastic ATP deficit of $36.7 \mu\text{mol g}^{-1} \text{FW h}^{-1}$ (Fig. 2, Table 2) which may be balanced by the following mechanisms. First, cytosolic NADPH from the OPPP ($14 \mu\text{mol g}^{-1} \text{FW h}^{-1}$) may be converted to $24.4 \mu\text{mol ATP g}^{-1} \text{FW h}^{-1}$ via oxidative phosphorylation starting at type II NADPH dehydrogenase (yields 2.5×0.7 ATP per NADPH). This leaves an ATP deficit of $12.3 \mu\text{mol g}^{-1} \text{FW h}^{-1}$. Second, GAP/3PGA cycling involving cytosolic GAPN may supply cytosolic NADPH (Fig. 1). To balance the entire ATP deficit of $36.7 \mu\text{mol g}^{-1} \text{FW h}^{-1}$ via oxidative phosphorylation starting at type II NADPH dehydrogenases would require a flux of $21 \mu\text{mol GAP g}^{-1} \text{FW h}^{-1}$ through this pathway. Third, mitochondrial NADH from photorespiration ($26.4 \mu\text{mol g}^{-1} \text{FW h}^{-1}$) may be converted to up to $66 \mu\text{mol ATP g}^{-1} \text{FW h}^{-1}$ via oxidative phosphorylation starting at complex I of the mitochondrial electron transport chain (yields 2.5 ATP per NADH). Fourth, GAP/3PGA cycling involving cytosolic PGK may supply ATP (Fig. 1). This pathway provides ATP and NADH (from GAPC) at equimolar amounts. If the latter is imported into mitochondria by the malate valve and used for oxidative phosphorylation (Fig. 2), GAP/3PGA cycling at a rate of $10.5 \mu\text{mol g}^{-1} \text{FW h}^{-1}$ would suffice to meet the ATP deficit of $36.7 \mu\text{mol g}^{-1} \text{FW h}^{-1}$ (1 ATP from PGK, 2.5 ATP from NADH provided by GAPC). If NADH from GAPC enters oxidative phosphorylation via type II NADH dehydrogenase which reduces ATP yields by about 30% (see above), GAP/3PGA cycling at a rate of $13.3 \mu\text{mol g}^{-1} \text{FW h}^{-1}$ is required to meet the ATP deficit (1 ATP from PGK, 2.5×0.7 ATP from NADH). Fifth, GAP/3PGA cycling

involving cytosolic PGK may supply all extra-chloroplastic ATP required (Fig. 1). This would provide an additional $36.7 \mu\text{mol NADH g}^{-1} \text{FW h}^{-1}$ from GAPC which could be used to balance the peroxisomal NADH deficit ($25.4 \mu\text{mol g}^{-1} \text{FW h}^{-1}$). Remaining extra-chloroplastic NADH ($37.7 \mu\text{mol g}^{-1} \text{FW h}^{-1}$, includes the mitochondrial NADH surplus) would then most likely enter oxidative phosphorylation. Sixth, cytosolic NADH supplied by the chloroplast malate valve may be converted to ATP by mitochondrial NADH import (malate valve) and oxidative phosphorylation or oxidative phosphorylation via type II NADH dehydrogenase. Supply of $36.7 \mu\text{mol ATP g}^{-1} \text{FW h}^{-1}$ by these pathways requires net flux through the chloroplast malate valve of 14.7 or $21 \mu\text{mol NADH g}^{-1} \text{FW h}^{-1}$, respectively. Hence, numerous metabolic pathways can supply cytosolic ATP. All of them, however, involve oxidative phosphorylation. Thus, in illuminated *Camelina sativa* leaves under the conditions studied here, oxidative phosphorylation is strictly required. This is in agreement with results from a recent simulation study (Shameer et al. 2019).

Is mitochondrial NADH strictly required to balance peroxisomal NADH demands?

We estimated a peroxisomal NADH deficit of $25.4 \mu\text{mol g}^{-1} \text{FW h}^{-1}$ due to photorespiration (Fig. 2, Table 2). This deficit is often assumed to be balanced by the mitochondrial NADH surplus of $26.4 \mu\text{mol g}^{-1} \text{FW h}^{-1}$ coming mostly from photorespiration. However, there are other processes that may help to balance the peroxisomal NADH deficit. First, NAD(P)H from chloroplasts may be transferred to peroxisomes via malate valves. Second, combined action of GAP/3PGA cycling involving cytosolic GAPC and the peroxisomal malate valve may balance peroxisomal NADH demands. Concomitantly supplied ATP by PGK may help to balance extra-chloroplastic ATP deficits. Thus, to balance peroxisomal NADH demands, mitochondrial NADH is not strictly required. However, peroxisomal NADH import appears to be strictly required.

Where is the cytosolic NADPH coming from?

In the cytosol of illuminated *Camelina sativa* leaves, $14 \mu\text{mol NADPH g}^{-1} \text{FW h}^{-1}$ are supplied by the OPPP. Simultaneously, CO_2 accounting for 4.8% of total rubisco activity is released (see above). Relative carbon losses would further increase with decreasing carbon assimilation unless the OPPP is downregulated. However, at constant light, decreasing carbon assimilation (e.g., due to drought) promotes the synthesis of reactive oxygen species (Noctor et al. 2002) and their detoxification can be expected to increase the demand for cytosolic NADPH.

Primary flux control through the OPPP is exerted at the level of its first enzyme, G6PD. In the cytosol of Arabidopsis leaves, only G6PD5 was found (Wakao and Benning 2005). Reportedly, G6PD5 activity is not directly affected by redox agents (Wakao and Benning 2005). Correspondingly, cytosolic G6PD activity in *Solanum tuberosum* leaf discs was found to be constant in response to increasing H₂O₂ (Hauschild and von Schaewen 2003). By contrast, increases in activity occur with increasing sugar availability (sugars tested: mannose, glucose, fructose, sucrose) through *de novo* enzyme synthesis (Hauschild and von Schaewen 2003). As carbon assimilation decreases, so does sugar availability (Lawlor and Fock 1977; Sánchez-Rodríguez et al. 1999). This can be expected to result in decreasing cytosolic NADPH supply by G6PD when it is most needed (due to increasing oxidative load). Thus, especially (but not exclusively) under low carbon assimilation and high oxidative load, other sources of cytosolic NADPH may be required.

GAPN catalyses the irreversible oxidation of GAP to 3PGA and simultaneous reduction of NADP⁺ to NADPH (Fig. 1). Interestingly, a GAPN null mutant showed increased expression and activity of cytosolic G6PD5 (3.8-fold increase in At3g27300) and an increased oxidative load (Rius et al. 2006). This suggests that G6PD5 and GAPN interact in redox control and that GAPN is required in redox control since increased G6PD5 expression cannot fully compensate for the loss in GAPN function.

Just like GAPN, GAPC and PGK catalyse cytosolic GAP to 3PGA conversions. Intriguingly, the GAPN null mutant showed increased expression and activity of GAPC (GAPC1: 3.5-fold increase in At3g04120) (Rius et al. 2006). This was suggested to compensate for the lack of GAPN activity (Rius et al. 2006). However, GAPC and PGK supply NADH and ATP instead of NADPH which cannot be used in redox control. Alternatively, upregulation of GAPC may support COR cycling (Wieloch 2021). In contrast to the OPPP, COR cycling does not release CO₂ from metabolism. Each turn provides NADPH at GAPN and consumes ATP and NADH at PGK and GAPC (Fig. 1). Required NADH and ATP may be provided via the chloroplast malate valve and oxidative phosphorylation (Fig. 2). In this regard, a putative moonlighting function of GAPC is worth noting (Scheibe 2019). With increasing oxidative load (as observed in the GAPN null mutant), GAPC is

increasingly deactivated and moved to the nucleus (Piattoni et al. 2013; Hildebrandt et al. 2015). There, it can bind to the gene coding for chloroplast malate dehydrogenase, and expression of this gene was found to increase during the initial stages of oxidative stress (Hildebrandt et al. 2015). Furthermore, it is worth noting that the GAPN null mutant shows increased expression of cytosolic malate dehydrogenase (Rius et al. 2006). Taken together, this may result in increased malate valve capacities to support GAPN and PGK.

Recently, we reported isotope evidence for flux through the COR cycle and drought-induced upregulation (Wieloch et al. 2021). Still, more work is needed on its role in supplying cytosolic NADPH, its interaction with the cytosolic OPPP, and their environmental/developmental dependences. In this regard, wide-screening enzyme expression in a G6PD5 knockout mutant may be valuable.

Table 1 Metabolite and cofactor flux associated with sucrose cycling and carbon re-injection into the Calvin-Benson cycle by the cytosolic oxidative pentose phosphate pathway in illuminated *Camelina sativa* leaves [$\mu\text{mol g}^{-1} \text{FW h}^{-1}$].

Reaction	Metabolite flux	Cofactor	Cofactor flux
Calvin-Benson cycle			
PGK.p	20.00	ATP	-20.00
GAPDH.p	20.00	NAD(P)H	-20.00
PRK.p	10.62	ATP	-10.62
Photorespiration			
GS.m	1.21	ATP	-1.21
GOGAT.p	1.21	Fd _{red}	-2.43
GDC.m	1.21	NADH	1.21
HPR.ox	1.21	NADH	-1.21
GK.p	1.21	ATP	-1.21
Oxidative pentose phosphate pathway			
G6PD.c	6.98	NADPH	6.98
6PGD.c	6.98	NADPH	6.98
Sucrose cycling			
UGPase.c	2.16	UTP	-2.16
HK.c	2.16	ATP	-2.16
FK.c	2.16	ATP	-2.16

Metabolite flux as reported by Xu et al. (in prep.). Negative and positive cofactor fluxes denote cofactor consumption and production, respectively. Intracellular location of enzyme reaction: .p, chloroplast; .m, mitochondrion; .ox, peroxisome; .c, cytosol. Enzymes: 6PGD, 6-phosphogluconate dehydrogenase; FK, fructokinase; G6PD, glucose-6-phosphate dehydrogenase; GAPDH, phosphorylating glyceraldehyde-3-phosphate dehydrogenase; GDC, glycine decarboxylase complex; GK, glycerate kinase; GOGAT, glutamine α -ketoglutarate aminotransferase; GS, glutamine synthetase; HK, hexokinase; HPR, hydroxypyruvate reductase; PGK, phosphoglycerate kinase; PRK, phosphoribulokinase; UGPase, UDP-glucose pyrophosphorylase. Confidence intervals of flux estimates given in Table SI 2.

Table 2 Metabolite and cofactor flux in illuminated *Camelina sativa* leaves [$\mu\text{mol g}^{-1} \text{FW h}^{-1}$].

Reaction	Metabolite flux	Cofactor	Cofactor flux
Calvin-Benson cycle			
PGK.p	420.20	ATP	-420.20
GAPDH.p	420.20	NAD(P)H	-420.20
PRK.p	223.10	ATP	-223.10
Photorespiration			
GS.m	25.50	ATP	-25.50
GOGAT.p	25.50	Fd _{red}	-51.00
GDC.m	25.49	NADH	25.49
HPR.ox	25.38	NADH	-25.38
GK.p	25.38	ATP	-25.38
Oxidative pentose phosphate pathway			
G6PD.c	6.98	NADPH	6.98
6PGD.c	6.98	NADPH	6.98
Starch and sucrose biosynthesis, and sucrose cycling			
AGPase.p	10.51	ATP	-10.51
UGPase.c	7.86	UTP	-7.86
HK.c	2.16	ATP	-2.16
FK.c	2.16	ATP	-2.16
Glycolysis			
PK.c	0.99	ATP	0.99
Fatty acid biosynthesis			
PDC.p	0.44	NADH	0.44
ACC.p	0.44	ATP	-0.39
KAR.p	0.44	NADPH	-0.39
ACPr.p	0.44	NADH	-0.39
Tricarboxylic acid cycle			
PDC.m	0.92	NADH	0.92
IDH.m	0.92	NADH	0.92
Amino acid biosynthesis			
GDH.m	0.92	NADH	-0.92

Metabolite flux as reported by Xu et al. (in prep.). Negative and positive cofactor fluxes denote cofactor consumption and production, respectively. Intracellular location of enzyme reaction: .p, chloroplast; .m, mitochondrion; .ox, peroxisome; .c, cytosol. Enzymes: 6PGD, 6-phosphogluconate dehydrogenase; ACC, acetyl-CoA carboxylase; ACPr, 2,3-trans-enoyl-ACP reductase; AGPase, ADP-glucose pyrophosphorylase; FK, fructokinase; G6PD, glucose-6-phosphate dehydrogenase; GAPDH, phosphorylating glyceraldehyde-3-phosphate dehydrogenase; GDC, glycine decarboxylase complex; GDH, glutamate dehydrogenase; GK,

glycerate kinase; GOGAT, glutamine α -ketoglutarate aminotransferase; GS, glutamine synthetase; HK, hexokinase; HPR, hydroxypyruvate reductase; IDH, isocitrate dehydrogenase; KAR, 3-ketoacyl-ACP reductase; PDC, pyruvate dehydrogenase complex; PGK, phosphoglycerate kinase; PK, pyruvate kinase; PRK, phosphoribulokinase; UGPase, UDP-glucose pyrophosphorylase. Confidence intervals of flux estimates given in Table SI 3.

glyceraldehyde-3-phosphate dehydrogenase; GDC, glycine decarboxylase complex; GDH, glutamate dehydrogenase; GK, glycerate kinase; GOGAT, glutamine α -ketoglutarate aminotransferase; GS, glutamine synthetase; HK, hexokinase; HPR, hydroxypyruvate reductase; IDH, isocitrate dehydrogenase; KAR, 3-ketoacyl-ACP reductase; PDC, pyruvate dehydrogenase complex; PGK, phosphoglycerate kinase; PK, pyruvate kinase; PRK, phosphoribulokinase; UGPase, UDP-glucose pyrophosphorylase. Metabolites: 1,3BPG, 1,3-bisphosphoglycerate; 2PGA, 2-phosphoglycerate; 3PGA, 3-phosphoglycerate; 6PG, 6-phosphogluconate; 6PGL, 6-phosphogluconolactone; ACA, acetyl coenzyme A; ADPG, ADP-glucose; AKG, α -ketoglutarate; F6P, fructose 6-phosphate; FRU, fructose; G1P, glucose 1-phosphate; G6P, glucose 6-phosphate; GA, glycerate; GLC, glucose; GLN, glutamine; GLO, glyoxylate; GLU, glutamate; GLY, glycine; HP, hydroxypyruvate; ICT, isocitrate; PEP, phosphoenolpyruvate; PYR, pyruvate; Ru5P, ribulose 5-phosphate; RuBP, ribulose 1,5-bisphosphate; SER, serine; SUC, sucrose; TP, triose phosphate (glyceraldehyde 3-phosphate and dihydroxyacetone phosphate); UDPG, UDP-glucose.

Figure 2 Compartment specific cofactor consumption (negative values) and production (positive values) in 4-week-old photosynthesising *Camelina sativa* leaves. Error bars represent 95% confidence intervals. 1, malate valves exchange NADH through malate-oxaloacetate interconversion by malate dehydrogenase (Selinski and Scheibe 2019). A chloroplastic isoform accepts NADPH. 2 and 3, cofactor transfer through redox cycles proposed by Kelly and Gibbs (1973a) and Stocking and Larson (1969), respectively. 4, counter-exchange of cytosolic ADP for mitochondrial ATP by the ADP/ATP carrier (Klingenberg 2008). 5, conversion of ATP and NADH to NADPH by the cytosolic oxidation-reduction cycle (Wieloch 2021). 6, NADH to ATP conversion by oxidative phosphorylation. 7, conversion of cytosolic NAD(P)H to mitochondrial ATP by oxidative phosphorylation starting at type II NAD(P)H dehydrogenases and glycerol-3-phosphate dehydrogenase both located in the inner mitochondrial membrane facing the cytosol. Solid arrow, direct cofactor transport. Dotted arrow, indirect cofactor transport via redox reactions.

References

- Andrews TJ, Kane HJ (1991) Pyruvate is a by-product of catalysis by ribulosebisphosphate carboxylase/oxygenase. *Journal of Biological Chemistry* 266:9447–9452
- Buchanan BB, Gruissem W, Jones RL (eds) (2015) *Biochemistry and molecular biology of plants*, 2nd edn. John Wiley & Sons, Ltd, Chichester
- Canvin DT (1979) Photorespiration: Comparison between C₃ and C₄ plants. In: Gibbs M, Latzko E (eds) *Photosynthesis II: Photosynthetic Carbon Metabolism and Related Processes*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp 368–396
- Hasunuma T, Harada K, Miyazawa S-I, et al (2010) Metabolic turnover analysis by a combination of *in vivo* ¹³C-labelling from ¹³CO₂ and metabolic profiling with CE-MS/MS reveals rate-limiting steps of the C₃ photosynthetic pathway in *Nicotiana tabacum* leaves. *Journal of Experimental Botany* 61:1041–1051
- Hauschild R, von Schaewen A (2003) Differential regulation of glucose-6-phosphate dehydrogenase isoenzyme activities in potato. *Plant physiology* 133:47–62
- Hedrich R, Sauer N, Neuhaus HE (2015) Sugar transport across the plant vacuolar membrane: nature and regulation of carrier proteins. *Current Opinion in Plant Biology* 25:63–70
- Hildebrandt T, Knuesting J, Berndt C, et al (2015) Cytosolic thiol switches regulating basic cellular functions: GAPDH as an information hub? *Biological chemistry* 396:523–37
- Hinkle PC (2005) P/O ratios of mitochondrial oxidative phosphorylation. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* 1706:1–11
- Kelly GJ, Gibbs M (1973a) Nonreversible D-glyceraldehyde 3-phosphate dehydrogenase of plant tissues. *Plant Physiology* 52:111–118
- Kelly GJ, Gibbs M (1973b) A mechanism for the indirect transfer of photosynthetically reduced nicotinamide adenine dinucleotide phosphate from chloroplasts to the cytoplasm. *Plant Physiology* 52:674–676

- Klingenberg M (2008) The ADP and ATP transport in mitochondria and its carrier. *Biochimica et Biophysica Acta (BBA) - Biomembranes* 1778:1978–2021
- Lawlor DW, Fock H (1977) Water stress induced changes in the amounts of some photosynthetic assimilation products and respiratory metabolites of sunflower leaves. *Journal of Experimental Botany* 28:329–337
- Ma F, Jazmin LJ, Young JD, Allen DK (2014) Isotopically nonstationary ^{13}C flux analysis of changes in *Arabidopsis thaliana* leaf metabolism due to high light acclimation. *Proc Natl Acad Sci USA* 111:16967–16972
- Mahon JD, Fock H, Canvin DT (1974) Changes in specific radioactivity of sunflower leaf metabolites during photosynthesis in $^{14}\text{CO}_2$ and $^{12}\text{CO}_2$ at three concentrations of CO_2 . *Planta* 120:245–254
- McGowan RE, Gibbs M (1974) Comparative enzymology of the glyceraldehyde 3-phosphate dehydrogenases from *Pisum sativum*. *Plant Physiology* 54:312
- Noctor G, Veljovic-Jovanovic S, Driscoll S, et al (2002) Drought and oxidative load in the leaves of C_3 plants: a predominant role for photorespiration? *Annals of Botany* 89:841–850
- Palmieri F, Rieder B, Ventrella A, et al (2009) Molecular identification and functional characterization of *Arabidopsis thaliana* mitochondrial and chloroplastic NAD^+ carrier proteins. *The Journal of Biological Chemistry* 284:31249–31259
- Piattoni CV, Guerrero SA, Iglesias AA (2013) A differential redox regulation of the pathways metabolizing glyceraldehyde-3-phosphate tunes the production of reducing power in the cytosol of plant cells. *International Journal of Molecular Sciences* 14:8073–8092
- Rasmusson AG, Geisler DA, Moller IM (2008) The multiplicity of dehydrogenases in the electron transport chain of plant mitochondria. *Mitochondrion* 8:47–60
- Rius SP, Casati P, Iglesias AA, Gomez-Casati DF (2006) Characterization of an *Arabidopsis thaliana* mutant lacking a cytosolic non-phosphorylating glyceraldehyde-3-phosphate dehydrogenase. *Plant Molecular Biology* 61:945–957

- Sánchez-Rodríguez J, Pérez P, Martínez-Carrasco R (1999) Photosynthesis, carbohydrate levels and chlorophyll fluorescence-estimated intercellular CO₂ in water-stressed *Casuarina equisetifolia* Forst. & Forst. Plant, Cell & Environment 22:867–873
- Scheibe R (2019) Maintaining homeostasis by controlled alternatives for energy distribution in plant cells under changing conditions of supply and demand. Photosynthesis Research 139:81–91
- Seelert H, Poetsch A, Dencher NA, et al (2000) Proton-powered turbine of a plant motor. Nature 405:418–419
- Selinski J, Scheibe R (2019) Malate valves: old shuttles with new perspectives. Plant Biology 21:21–30
- Shameer S, Ratcliffe RG, Sweetlove LJ (2019) Leaf energy balance requires mitochondrial respiration and export of chloroplast NADPH in the light. Plant Physiol 180:1947–1961
- Sharkey TD, Preiser AL, Weraduwege SM, Gog L (2020) Source of ¹²C in Calvin-Benson cycle intermediates and isoprene emitted from plant leaves fed with ¹³CO₂. Biochem J 477:3237–3252
- Somerville CR, Ogren WL (1980) Inhibition of photosynthesis in Arabidopsis mutants lacking leaf glutamate synthase activity. Nature 286:257–259
- Stocking CR, Larson S (1969) A chloroplast cytoplasmic shuttle and the reduction of extraplastid NAD. Biochemical and Biophysical Research Communications 37:278–282
- Szeczowka M, Heise R, Tohge T, et al (2013) Metabolic fluxes in an illuminated Arabidopsis rosette. Plant Cell 25:694–714
- Taira M, Valtersson U, Burkhardt B, Ludwig RA (2004) *Arabidopsis thaliana* GLN2-encoded glutamine synthetase is dual targeted to leaf mitochondria and chloroplasts. Plant Cell 16:2048–2058

- Wakao S, Benning C (2005) Genome-wide analysis of glucose-6-phosphate dehydrogenases in Arabidopsis. *The Plant Journal* 41:243–256
- Wieloch T (2021) A cytosolic oxidation–reduction cycle in plant leaves. *J Exp Bot* 72:4186–4189
- Wieloch T, Werner RA, Schleucher J (2021) Carbon flux around leaf-cytosolic glyceraldehyde-3-phosphate dehydrogenase introduces a ^{13}C signal in plant glucose. *J Exp Bot* 72:7136–7144
- Xu Y, Fu X, Sharkey TD, et al (2021) The metabolic origins of non-photorespiratory CO_2 release during photosynthesis: A metabolic flux analysis. *Plant Physiology* 186:297–314
- Xu Y, Wieloch T, Shachar-Hill Y, et al (in prep.) Reimport of carbon from cytosolic and vacuolar sugar pools into the Calvin-Benson cycle explains photosynthesis labeling anomalies.