USP22 controls type III interferon signaling and SARS-CoV-2 infection through activation of STING

*Rebekka Karlowitz*¹, *Megan L. Stanifer*^{2,3}, *Jens Roedig*¹, *Geoffroy Andrieux*^{4, 5},
*Denisa Bojkova*⁶, *Sonja Smith*¹, *Lisa Kowald*¹, *Ralf Schubert*⁷, *Melanie Boerries*^{4, 5},
*Jindrich Cinatl Jr.*⁶, *Steeve Boulant*^{3,8}, *Sjoerd J. L. van Wijk*^{1,9,#}

¹ Institute for Experimental Cancer Research in Pediatrics, Goethe University Frankfurt, Komturstasse 3a, 60528 Frankfurt am Main, Germany

² Department of Infectious Diseases/Molecular Virology, Medical Faculty, Center for Integrative Infectious Diseases Research (CIID), University of Heidelberg, 69120 Heidelberg, Germany

³ Department of Molecular Genetics and Microbiology, University of Florida College of Medicine, Gainesville, Florida, USA

⁴ Institute of Medical Bioinformatics and Systems Medicine, Medical Center-University of Freiburg, Faculty of Medicine, University of Freiburg, 79110 Freiburg, Germany

⁵ German Cancer Consortium (DKTK) and German Cancer Research Center (DKFZ), partner site Freiburg, 79110 Freiburg, Germany

⁶ Institute of Medical Virology, University Hospital Frankfurt, Goethe University, 60596 Frankfurt am Main, Germany

⁷ Division for Allergy, Pneumology and Cystic Fibrosis, Department for Children and Adolescents, University Hospital Frankfurt, Goethe University, Theodor-Stern-Kai 7, 60590 Frankfurt am Main, Germany.

⁸ Department of Infectious Diseases, Virology, Medical Faculty, Center for Integrative Infectious Diseases Research (CIID), University of Heidelberg, 69120 Heidelberg, Germany

27

28 ⁹ German Cancer Consortium (DKTK) and German Cancer Research Center (DKFZ)
29 partner site Frankfurt/Mainz, Frankfurt am Main, Germany

30

31 **Running title:** USP22 controls type III IFN signaling and SARS-CoV-2 infection

32

33 # Corresponding author: Sjoerd J. L. van Wijk, Institute for Experimental Cancer
34 Research in Pediatrics, Goethe University Frankfurt, Komturstrasse 3a, 60528
35 Frankfurt am Main, Germany, Phone: +49 69 67866574, Fax: +49 69 6786659158,
36 Email: vanWijk@med.uni-frankfurt.de, s.wijk@kinderkrebsstiftung-frankfurt.de

37

38

39

Abstract

Pattern recognition receptors (PRRs) and interferons (IFNs) serve as essential antiviral defense against SARS-CoV-2, the causative agent of the COVID-19 pandemic. Type III IFN (IFN- λ) exhibit cell-type specific and long-lasting functions in autoinflammation, tumorigenesis and antiviral defense. Here, we identify the deubiquitinating enzyme USP22 as central regulator of basal IFN- λ secretion and SARS-CoV-2 infections in native human intestinal epithelial cells (hIECs). USP22-deficient hIECs strongly upregulate genes involved in IFN signaling and viral defense, including numerous IFN-stimulated genes (ISGs), with increased secretion of IFN- λ and enhanced STAT1 signaling, even in the absence of exogenous IFNs or viral infection. Interestingly, USP22 controls basal and 2'3'-cGAMP-induced STING activation and loss of STING reversed STAT activation and ISG and IFN- λ expression. Intriguingly, USP22-deficient hIECs are protected against SARS-CoV-2 infection, viral replication and the formation of *de novo* infectious particles, in a STING-dependent manner. These findings reveal USP22 as central host regulator of STING and type III IFN signaling, with important implications for SARS-CoV-2 infection and antiviral defense.

Key words: USP22, STING, cGAS, interferon, SARS-CoV-2, ISG, ubiquitin, deubiquitinating enzyme, pattern recognition receptors

Introduction

Sensing of “non-self” is a key feature of innate immunity and underlies the recognition of viruses, bacteria and fungi, but also plays important roles in cancer and autoimmune diseases^{1,2}. Pattern recognition receptors (PRRs), like Toll-like receptors (TLRs), Nucleotide-binding oligomerization domain (NOD)-like receptors (NLRs) and retinoic acid-inducible gene 1 protein (RIG-I)-like receptors (RLRs) are essential components of innate immune signaling and selectively recognize pathogen-associated molecular patterns (PAMPs). Dedicated PRRs, like TLR3, RIG-I, Melanoma differentiation-associated protein 5 (MDA5) and Cyclic GMP-AMP synthase (cGAS)-Stimulator of interferon genes protein (STING) recognize viral double stranded RNA (dsRNA) and dsDNA, and are important sensors for infections with RNA and DNA viruses, as well as infections with retroviruses¹⁻³. Whereas TLR3 recognizes dsRNA in endosomes, the prototypical RLRs, RIG-I and MDA5, sense cytosolic dsRNAs, while cGAS-STING detects viral dsDNA¹⁻⁴. STING is activated either directly via viral dsDNA, through the STING agonist 2’3’-cGAMP generated by the cyclic GMP-AMP synthase cGAS upon detection of viral dsDNA, or indirectly via RIG-1 and MDA5⁵. Activated STING interacts with TANK-binding kinase 1 (TBK1) and activates interferon regulatory factor (IRF) 1, -3, and -7 and Nuclear factor- κ B (NF- κ B), leading to the initiation of anti-viral and inflammatory transcriptional programs, including interferon-stimulated genes (ISGs) and interferons (IFN)⁵⁻⁸.

Interferons (IFNs) are secreted cytokines with important roles in immunity and anti-viral responses. IFN signaling relies on Janus kinase-Signal transducer and activator of transcription (JAK-STAT) activation, phosphorylation of STAT1/2 and the induction of ISG and IFN gene expression that influence viral replication^{9,10}. Although the vast majority of cell types can be triggered to express type I (IFN- α , - β , - ϵ , - κ and - ω) and

type III (IFN- λ 1, - λ 2, - λ 3 and - λ 4) IFNs, the expression of IFN-specific receptors is cell type restricted and determines IFN responses. For example, the type I IFN receptor (IFNAR) is ubiquitously expressed in many tissues, whereas expression of the type III IFN receptor IFNLR1 is mainly limited to epithelial cells, e.g. the gastro-intestinal and respiratory epithelium^{6-8,11,12}. Although type I and type III IFNs induce similar ISG signatures, type I IFNs generally trigger a more rapid increase and decay of ISG expression⁷. In addition, IFN- λ s have been described to be first-in-line against viral infections and might inhibit viral spread without triggering inflammatory responses, depending on IFN- λ receptor expression^{7,13,14}.

The novel severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is the causative agent of the pandemic Coronavirus disease 2019 (COVID-19) and belongs to the human coronaviruses (HCoV) that also includes SARS-CoV and MERS-CoV¹⁵. In many patients with severe COVID-19, SARS-CoV-2 infection induces the secretion of highly pro-inflammatory cytokines through cGAS-STING and NF- κ B-mediated signaling^{16,17}. Type I and III IFNs are important regulators of host viral defense against SARS-CoV-2^{6-8,11,12,18,19}, but at the same time, SARS-CoV-2 evades immune recognition via IFN and ISG suppression^{10,20}. Prolonged expression of low basal levels of type I and III IFNs might prime host responses against virus infection, including SARS-CoV-2²¹⁻²⁴. Although type III IFNs restrict SARS-CoV-2 infection in intestinal and airway epithelial cells^{19,25-29} and STING agonism reduces SARS-CoV-2 infection³⁰⁻³³, context-dependent damaging effects of type III IFNs on airway epithelia during viral infections have been described as well^{34,35}.

Innate immunity, PRRs and IFN signaling is closely regulated by ubiquitination, both by the host machinery as well as through viral E3 ligases and deubiquitinating enzymes

(DUBs) that hijack the host ubiquitin machinery ³⁶. STING, RIG-I, TLR3 and TBK1 are positively and negatively regulated by differential modification of polyubiquitin chains, including K11, K27, K48 and K63 linked chains ^{37,38}, by a variety of E3 ligases, such as TRIM56 ³⁹, TRIM32 ⁴⁰, MUL1 ⁴¹, AMFR ⁴², RNF5 ⁴³ and TRIM29 ⁴⁴ and RNF26 ⁴⁵. The interplay and functional consequences of ubiquitin modifications are complex and include proteasomal degradation as well as stabilization of protein-protein interactions. Importantly, IFN and anti-viral signaling are also heavily regulated by DUBs, like USP13 ⁴⁶, USP35 ⁴⁷ and CYLD ⁴⁸

Ubiquitin-specific peptidase 22 (USP22) is a DUB that is part of the deubiquitination module of the Spt-Ada-Gcn5-acetyltransferase (SAGA) complex, through which it regulates transcription via the control of histone H2A K119 and H2B K120 monoubiquitination (H2AK119ub1 and H2BK120ub1, respectively) ⁴⁹⁻⁵¹. Recently, additional USP22 substrates have emerged, with important roles in cell fate regulation and programmed cell death ⁵²⁻⁵⁴. Interestingly, USP22 has mostly been associated with IFN signaling and ISG expression upon infection with viruses ^{55,56}. However, up till now, the mechanisms how USP22 primes PRR and IFN signaling and prepares against anti-viral defense in native, uninfected settings remains unknown.

In light of the current COVID-19 pandemic and potential future pathogenic coronaviruses, identifying host factors that control SARS-CoV-2 infection is of extreme relevance. The roles of type III IFN in SARS-CoV-2 infections are only starting to emerge and are determined by tissue-specific factors as well. Here, we are the first to identify USP22 as a negative regulator of basal ISG expression, JAK/STAT activation and IFN signaling, even in the absence of exogenous IFNs or viral infection. Our findings elucidate USP22 as crucial host factor in shaping SARS-CoV-2 antiviral

defense by priming cellular anti-viral responsiveness prior to virus infection. Loss of USP22 in native, human intestinal epithelial cells (hIECs) triggers a strong upregulation of ISGs and, specifically, the type III IFN IFN- λ , mediated by STING. USP22 controls basal and 2'3'-cGAMP-induced STING ubiquitination, phosphorylation and activation, and combined loss of USP22 and STING rescues ISG expression, STAT signaling and IFN- λ production. Importantly, we found that USP22-deficient hIECs are prominently protected against SARS-CoV-2 infection, replication and the formation of novel infectious viral particles, which can be partially reversed by loss of STING expression.

Results

Profiling USP22-mediated gene expression in HT-29 hIECs

Substrate-specific deubiquitination is a central determinant of ubiquitin homeostasis and regulates receptor activation and internalization, proteasomal degradation and transcription. For the ubiquitin-specific protease USP22, both transcriptional and extranuclear targets have been identified. As part of the DUB module of the SAGA complex, USP22 regulates transcriptional elongation via H2AK119ub1 and H2BK120ub1⁴⁹⁻⁵¹. Up till now, the spectrum of target genes regulated by USP22 largely remains unclear, partially due to organism-, cell- and context-dependent redundancy in alternative DUBs that might compensate for loss of USP22⁵⁷. We previously reported that CRISPR/Cas9-mediated knockout (KO) of USP22 in human colon carcinoma cell line HT-29 affects RIPK3 ubiquitination during necroptosis, but without inducing major changes in RIPK1, RIPK3 and MLKL gene expression⁵⁴, suggesting gene-specific regulation of USP22. To identify the spectrum of USP22-regulated genes, we profiled USP22-dependent changes in gene expression in the human intestinal epithelial cell (hIEC) line HT-29. Quantification of alterations in gene expression in two independent HT-29 USP22 KO single cell clones revealed a marked alteration in gene expression, with 401 genes up-regulated and 182 down-regulated (**Figure 1A and Supplemental Figure 1A**). Loss of USP22 expression was accompanied with changes in H2Bub1, but not H2Aub1 (**Supplemental Figure 1B & C**). Among the top-50 differentially regulated genes, 30 were up- and 20 downregulated, with an adjusted P-value of < 0.05 (**Figure 1B**). Genes upregulated in both USP22 KO clones (#16 and #62) compared to control (non-human target: NHT) HT-29 cells include genes that encode for proteins involved in growth and differentiation, like Transforming Growth Factor β -1 (TGFB1), Tumor-associated calcium signal transducer 2 (TACSTD2) and Tyrosine-protein kinase Mer (MERTK)

and the cytosolic RNA- and DNA sensor DExD/H-Box Helicase 60 (DDX60). Downregulated genes include USP22, mitochondrial adenylate kinase 4 (AK4) that is involved in the regulation of mitochondrial function and ATP production ⁵⁸ and regenerating islet-derived protein 4 (REG4), a carbohydrate-binding lectin that has been identified as marker for deep crypt secretory cells (DSCs) that acts as niche for Lgr5-positive stem cells in the colon ⁵⁹. Differential regulation of gene expression, as well as loss of USP22 expression, was also demonstrated by independent qRT-PCR of the USP22-dependent upregulated genes TGFB1, SLFN5, TGM2 and DDX60, as well as downregulation of USP22, CXCR4 and AK4 (**Figure 1C**), confirming the quality of the microarray.

Loss of USP22 specifically enriches for genes involved in interferon signaling and response to viral infection

Next, gene-set enrichment analysis was performed on USP22-regulated genes to investigate if certain gene sets from gene ontology (GO) are specifically regulated by USP22. Interestingly, GO analysis revealed an enrichment of genes linked to type I and II interferon (IFN) signaling, as well as regulation of viral genome replication and several other viral processes, such as the regulation of viral genome replication, response to virus, response to IFN- γ , IFN- γ mediated signaling pathway in USP22 KO HT-29 cells as compared to control NHT HT-29 cells (**Figure 2A**). Interestingly, the GO terms of genes that are strongly downregulated are enriched in mitochondrial translation and gene expression, ribosomal and ribonucleoprotein complex biogenesis and the processing of tRNA, rRNA and ncRNA (**Figure 2A**).

Since previous studies suggest controversial roles of USP22 in IFN signaling ^{55,56,60,61}, we decided to further study USP22-dependent changes in genes involved in type I or

type II IFN signaling (**Figure 2B**). Loss of USP22 leads to the upregulation of many IFN stimulated genes (ISGs), some with important functions in viral defense, like OAS1, -2 and -3, MX1 and IFI27, suggesting a potential role of USP22 in the regulation of interferon signaling and viral responses (**Figure 2B**). Among the upregulated genes were components of the ISGylation machinery, like the ubiquitin-like modifier ISG15 and the ISG15-specific DUB USP18⁶². To validate the USP22-regulated changes in gene expression, qRT-PCR confirmed the increased expression of several ISGs, like BST2, PARP9, USP18, OAS3, IFIT1, IRF9, ISG15, OAS2, IFI27 and IFI6 in two independent HT-29 USP22 KO clones (**Figure 2C**). In addition, increased protein expression of MX1, IRF9, ISG56 and ISG20 could also be confirmed upon loss of USP22 (**Figure 2D**). These findings suggest that USP22 specifically controls the expression of genes involved in IFN signaling and virus defense, even in the absence of exogenous IFN stimulation or virus infection.

USP22 negatively regulates STAT1 signaling and IFN- λ 1 expression

The expression of ISGs is typically induced upon activation of IFN signaling pathways during pathogen invasion or autoinflammatory disease and serves to control inflammation and other defensive mechanisms⁹. Additionally, several IFNs are constitutively expressed at low levels as well⁶³ to prime and increase cellular responsiveness of IFN signaling upon activation by external stimuli²³. Interestingly, the expression levels of pan-IFN- α and IFN- β mRNA were upregulated upon loss of USP22, compared to non-human target control HT-29 (**Figure 3A**). This was accompanied by an increase in the expression of STAT1, an IFN-regulated ISG itself⁶⁴ as well as STAT1 phosphorylation, suggesting activation of IFN signaling pathways in USP22 KO HT-29 cells, compared to control cells (**Figure 3B**). Interestingly, in contrast to mRNA levels, in-depth analysis of USP22-mediated alterations in the

secretion of IFNs and IFN-related cytokines revealed only low basal levels of secreted IFN- α and IFN- β , suggesting that these cytokines might only weakly contribute to the observed ISG signature (**Figure 3C**). Surprisingly, the secretion and expression of IFN- λ 1, a type III IFN, was strongly upregulated in USP22 KO HT-29 cells compared to control cells (**Figure 3C & D**). In addition, loss of USP22 expression also induced elevated basal secretion of the pro-inflammatory cytokines CXCL10 and IL-8 and minor changes in the secretion of IFN- α 2 and GM-CSF, compared to controls. (**Figure 3C**). These findings suggest that USP22 negatively regulates IFN- λ 1 expression and ISG induction. Since type III IFN-induced target genes largely overlap with genes regulated by type I IFNs^{65,66}, type III IFN is likely the main IFN contributing to the USP22-dependent induction of ISG expression and STAT1 activation.

USP22 regulates type III IFN signaling via STING

Loss of USP22 expression specifically upregulates genes involved in IFN and viral responses. Within the context of viral infections, viral PAMPs, such as viral dsRNA and dsDNA are sensed by PRRs, like RIG-I, MDA5 and TLR3, and PRR activation mediates strong expression of IFNs and ISGs^{2,5,10}. Loss of USP22 leads to increased expression of RIG-I, MDA5 and TLR3 (**Figure 4A**). To investigate if these PRRs are functionally involved in USP22-mediated increased ISG signaling, the expression of RIG-I/DDX58, MDA5/IFIH1 and TLR3 was ablated with CRISPR/Cas9 in NHT and USP22 KO HT-29 cells (**Figure 4B - D**). Interestingly, despite efficient KO of the individual PRRs in both NHT and USP22 KO HT-29 cells, additional deletion of RIG-I, MDA5 or TLR3 did not decrease USP22-dependent STAT1 phosphorylation or ISG56 expression (**Figure 4B - D**). Interestingly, USP22-TLR3 double knockout (dKO) HT-29 cells even exhibit an increase in phosphorylated and total STAT1 levels as well as ISG56 expression, suggesting potential TLR3-specific effects of USP22 (**Figure 4D**).

An alternative source of IFN production might be from PRR-mediated detection of self-DNA (e.g. DNA damage and DNA double strand breaks), leading to the induction of IFN- α and IFN- λ via NF- κ B signaling ⁶⁷, as observed in several types of cancer. This is of special interest since USP22, apart from its role in transcriptional regulation, has also been associated with DNA damage responses ⁶⁸ and V(D)J recombination and CSR *in vivo* by facilitating c-NHEJ ⁶⁹. Since CRISPR/Cas9-mediated loss of USP22 did not lead to increased γ H2AX levels in hIECs compared to controls (**Supplemental Figure 2A**) despite increased NF- κ B signaling (**Supplemental Figure 2B**), it seems unlikely that DNA damage caused by loss of USP22 might contribute to IFN signaling.

Loss of RIG-I, MDA5 and TLR3 did not reverse the USP22-dependent IFN signature, suggesting that either functional redundancy between the selected PRRs could compensate for loss of individual PRRs or that additional PRRs are involved. Interestingly, expression of STING/TMEM173 was also increased in USP22 KO HT-29 cells (**Figure 4E and Supplemental Figure 2C**). STING can be activated via cGAS or indirectly via RIG-1 and MDA5, leading to complex formation with TBK1 and activation of IFN and NF- κ B signaling ⁵⁻⁸. To further investigate the potential PRR redundancy and the involvement of STING, NHT and USP22 KO HT-29 cells were stimulated with the TLR3-, RIG-I- and MDA5-agonist polyinosinic:polycytidylic acid (poly(I:C)), or the 45-bp non-CpG oligomer IFN-stimulating DNA (ISD) from *Listeria monocytogenes* that strongly activates the STING-TBK1-IRF3 axis ^{70,71}. Intriguingly, whereas poly(I:C) induced a prominent increase in the levels of total and phosphorylated STAT1 in both NHT and USP22 KO cells, ISD selectively induced increases in total and phosphorylated STAT1 levels in USP22 KO cells, but not in NHT control cells, which was also reflected in a prominent ISD-mediated induction of RIG-I

expression and STING activation (**Figure 4E**). In addition, ISD also induced strong expression of the representative ISGs OAS3 and IRF9 in USP22 KO cells compared to controls (**Figure 4F**). To confirm the role of STING in USP22-induced type III IFN signaling, USP22-STING dKO HT-29 cells were generated (**Figure 4G**). USP22-STING dKO cells exhibit strikingly reduced levels of basal and phosphorylated STAT1 protein compared to USP22 KO HT-29 cells (**Figure 4G**), suggesting a STING-dependent rescue of the USP22-dependent IFN signature. In line with this, USP22-induced ISG expression could be reversed as well in USP22-STING dKO HT-29 cells (**Figure 4H**). Additionally, USP22-mediated increases in IFN- λ expression could also largely be reduced upon USP22 STING dKO, whereas expression of IFN- α and IFN- β largely remains unaffected (**Figure 4I**). These findings reveal an important role of USP22 as negative regulator of STING-dependent type III IFN signaling in hIECs.

USP22 negatively regulates STING activation and ubiquitination

The differential response to ISD, but not poly(I:C), and the reversal of the IFN signature in USP22-STING dKO hIECs suggests an important role of USP22 in the control of STING-induced type III IFN signaling. However, until now, the mechanisms of how USP22 regulates STING function remain unclear. Therefore, we subjected HT-29 control and USP22 KO cells to the STING agonist 2'3'-cGAMP and observed a fast, strong and more prolonged activation and phosphorylation of STING, as well as increased TBK1 and IRF3 phosphorylation (**Figure 5A**). In addition, the analysis of 2'3'-cGAMP-treated USP22 KO HT-29 cells revealed a very prominent increase in IFNL1 expression in USP22 KO cells, accompanied by increased IFNA and IFNB expression as well, but to a much lesser extent (**Figure 5B**).

Since STING expression is controlled by IFNs, constitutive IFN-mediated priming upon USP22 deficiency might underly the upregulation of STING. To investigate the relevance of auto- and paracrine IFN signaling in the regulation of STING expression, control and USP22 KO HT-29 cells were incubated with the JAK/STAT inhibitor ruxolitinib. JAK/STAT inhibition increased STING protein and mRNA expression levels in USP22 KO cells, compared to controls, suggesting that IFN-dependent auto- or paracrine activation of STING expression is unlikely (**Figure 5C and Supplemental Figure 3A**). Of note, USP22-mediated increases in STAT1 phosphorylation could be reversed with ruxolitinib (**Figure 5C**).

STING is reported to be modified with several types of ubiquitin chains that mediate context dependent effects, ranging from proteasomal degradation to the stimulation of signaling functions. STING protein levels were slightly stabilized in cycloheximide (CHX)-treated USP22 KO HT-29 cells compared to controls (**Figure 5D**). In line with these observations, basal and 2'3'-cGAMP-induced STING ubiquitination was also increased in USP22 KO HT-29 cells, compared to NHT control cells (**Figure 5E**). Together, these findings suggest that USP22-mediated effects on type III IFN might be predominantly regulated by activating STING ubiquitination and lesser through auto- or paracrine IFN signaling.

Loss of USP22 protects against SARS-CoV-2 infection, replication and de novo infectious virus production in a STING-dependent manner

Previous studies revealed important, but highly context-dependent roles of STING and type III IFNs in the control of SARS-CoV-2 infection ^{19,30,32,33}. In addition, USP22 has been linked to viral signaling ⁵⁶. To investigate the significance of USP22 and the

resulting STING-mediated upregulation of type III IFN and ISG signaling for viral defense, the role of the USP22-STING axis was tested during SARS-CoV-2 infection. For this, we generated control and USP22 KO Caco-2 cells that express the virus receptors ACE-2 and TMPRSS2 and are susceptible to infection with SARS-CoV-2 virus¹⁹. Loss of USP22 expression in Caco-2 cells triggered phosphorylation of STAT1 and increased expression of STING, compared to wild-type (WT) and NHT CRISPR/Cas9 control Caco-2 cells (**Figure 6A**). Increased USP22-dependent upregulation of IFN signaling in Caco-2 cells was also reflected in the increased expression of the antiviral ISGs IRF9 and OAS3 (**Figure 6B**). Intriguingly, USP22-deficient Caco-2 cells also express higher levels of IFN- λ 1, compared to wild-type and CRISPR/Cas9 control non-human target cells, whereas IFN- α and IFN- β expression largely remained unaffected (**Supplemental Figure 4A**).

To test the functional relevance of the increased antiviral signaling upon loss of USP22 expression, WT, NHT and USP22 KO Caco-2 cells were subjected to infection with SARS-CoV-2 particles at a MOI of 1. Infected Caco-2 cells were fixed at 24 hours post infection (hpi) and subjected to quantification of SARS-CoV-2 replication via immunofluorescence with the NP antibody recognizing SARS nucleocapsid protein. Interestingly, USP22-deficient cells displayed a prominent decrease in SARS-CoV-2 infection compared to infected WT or NHT Caco-2 cells (**Figure 6C**), as determined by immunofluorescence of viral protein. In addition, 6 and 24 hpi, SARS-CoV-2-infected USP22-deficient Caco-2 cells had lower genome copy numbers, compared to WT and NHT control cells (**Figure 6D**). These findings agree with a decreased release of *de novo* infectious SARS-CoV-2 viral particles in supernatants of USP22 KO Caco-2 cells compared to WT and NHT Caco-2 cells (**Supplemental Figure 4B**). Intriguingly, USP22-STING dKO hIECs exhibit higher SARS-CoV-2 replication rates as well as the

353 formation of more *de novo* infectious viral particles compared to USP22 KO hIECs
354 confirming that the USP22-STING connection also affects antiviral defense against
355 SARS-CoV-2 infection (**Figure 6E & F and Supplemental Figure 4C**). Together,
356 these findings indicate that USP22 critically controls SARS-CoV-2 infection, replication
357 and the generation of novel infectious viral particles, partially through STING.

Discussion

Carefully controlled regulation of IFN secretion and signaling is essential for organizing innate immunity, inflammation and anti-viral defense and deregulation of IFNs occur in auto-inflammatory diseases and cancer ^{1,2}. Type I, II and III IFNs elicit complex and intertwined JAK/STAT-based signaling pathways that regulate the expression of IFN stimulated genes (ISG), IFNs, STATs and IRFs with important implications for anti-viral signaling ^{9,10}. Additionally, IFN responsiveness is heavily influenced by IFN receptor affinities, expression and assembly and positive and negative regulation via ISGs, often in cell- and organ-specific manners ⁹. IFN signaling, ISG function and PRR-mediated antiviral defense is carefully controlled by ubiquitination and multiple deubiquitinating enzymes (DUBs) have been linked to the regulation of IFN-specific JAK/STAT pathways and response to viral infection ³⁸.

Apart from studying USP22 functions on interferon signaling in mouse models ⁶⁰, previous findings exclusively investigate cellular functions of USP22 upon virus infection and applied overexpression models to investigate USP22 interactions and USP22-mediated ubiquitination ^{55,56}. Here, we are the first to study the basal functions of USP22 in the regulation of ISG expression and STAT signaling in native human intestinal epithelial cells (hIECs). We identify USP22 as negative regulator of type III IFN secretion in basal settings without the addition of exogenous IFNs or by viral infection. Our findings reveal that USP22 regulates both basal and 2'3'-cGAMP-induced STING ubiquitination and activation, even in the absence of ectopic IFNs or viral infection, and loss of STING expression reverses the effects of USP22 KO on IFN signaling. Finally, we test the functional relevance of basal USP22- and STING-mediated IFN and JAK/STAT priming on SARS-CoV-2 infection and identify a critical

role of USP22 in the control of SARS-CoV-2 infection, replication and *de novo* formation of infectious viral particles, in a STING-dependent manner.

Despite the finding that USP22 regulates ISG expression in hIECs, USP22 does not exclusively control ISG or IFN-related gene expression. Hence, a large fraction of IFN-unrelated genes is changed while the expression of other genes is not altered upon loss of USP22 expression. Until now, the basis for this selectivity remains unclear. In agreement with previous observations^{50,72}, loss of USP22 expression in hIECs indeed increased H2Bub1, a hallmark of transcriptionally active chromatin⁷³⁻⁷⁵. Interestingly, increased H2Bub1 could also be detected at nucleosomes at ISG-coding genes upon specific deletion of USP22 in the murine hematopoietic system, underlying the upregulation of ISG expression⁶⁰. This was accompanied by alterations in hematopoietic stem cells (HSCs), myelopoiesis, B cell development, T cell activation, the numbers of B and plasma cells, serum immunoglobulin levels and the appearance of autoantibodies, but not with an increased systemic secretion of IFNs⁶⁰. This is surprising since IFNs themselves are ISGs as well and IFN expression levels are often maintained at low basal levels to serve as priming signals that allow a fast and adequate increase of IFN responses upon virus infection. Indeed, a large fraction of USP22-regulated ISGs has been demonstrated to be involved as important anti-SARS-CoV-2 countermeasures⁷⁶. In addition, global and ISG-specific levels of H2Bub1 can be regulated by type I IFN signaling during infection with human adenovirus as well, in a manner depending on human Bre1/RNF20 and the viral E1A protein⁷⁷. Intriguingly, the RNF20/RNF40 E3 ligase complex, responsible for H2B ubiquitination⁷⁸, was shown to protect against SARS-CoV-2 infection, and RNF20 becomes cleaved and inactivated by the SARS-CoV-2 protease 3CLpro⁷⁹. At present, the functional role of 3CLpro-mediated inactivation of RNF20 for H2Bub1 still remains to be addressed.

409

410 Since loss of USP22 mostly affects type III IFN expression and secretion, USP22 likely
411 mediates ISG expression both via epigenetic regulatory mechanisms as well as
412 through long-term IFN-mediated priming effects in hIECs. In contrast to type I IFN, type
413 III IFN is mostly sensed in gastro-intestinal and airway epithelia and in the blood-brain
414 barrier ^{6-8,11,65}. IFN- λ mostly exhibits long-term signaling effects and plays important
415 roles in SARS-CoV-2 infection in airway epithelial and gastro-intestinal cells and
416 organoids and has been shown to critically control antiviral defense ^{19,25-29}.

417

418 The susceptibility towards SARS-CoV-2-infections is determined by USP22-mediated
419 regulation of STING. STING is described as mediator of IFN- λ 1 production in HT-29
420 cells, and during viral infection in primary human macrophages in a Ku70-dependent
421 manner ^{80,81}. We furthermore demonstrate for the first time that in the absence of viral
422 infections or exogenous IFN, loss of USP22 expression resulted in basal and 2'3'-
423 cGAMP-induced STING ubiquitination in hIECs. In addition, loss of STING expression
424 decreased the IFN/ISG signaling that occurred under USP22 deficiency, suggesting
425 that STING acts as a physical scaffold for USP22-dependent ubiquitin modifications.
426 STING ubiquitination serves different physiological roles, including determining protein
427 stability, mediating protein-protein interactions and cellular localization ³⁹⁻⁴⁸. Recently,
428 cGAS-STING activity has emerged as regulator of immunopathology in COVID-19,
429 highlighting the relevance of STING regulation ⁸². STING ubiquitination enables the
430 STING-TBK1 interaction upon cGAS-mediated recognition of cytosolic DNA and is
431 generally associated with activation of ISG expression ⁷². Until now, USP22-mediated
432 STING ubiquitination has only been described upon viral infection and upon ectopic
433 overexpression. For example, overexpressed USP22 modifies ectopically expressed
434 STING with HA-tagged K27 ubiquitin upon HSV-1 infection in HEK293T cells ⁵⁶.

USP22 controls nuclear accumulation of IRF3 and type I IFN signaling through KPNA2 deubiquitination only upon infection with SeV and HSV-1 and loss of USP22 expression decreased type I IFN responses upon virus infection, while USP22 deletion in uninfected cells did not trigger basal IFN signaling ⁵⁵. At present, the role of type III IFNs in SeV and HSV-1 infections remains unclear.

Taken together, our findings identify USP22 as central host factor that determines ISG expression and type III IFN production via STING, with important implications for SARS-CoV-2 infection and IFN priming.

Materials and Methods

Cell culture, reagents and chemicals

The human colon carcinoma cell line HT-29 was obtained from DSMZ (Braunschweig, Germany) and cultivated in McCoy's 5A Medium GlutaMAX™-I (Life Technologies, Inc., Eggenstein, Germany), supplemented with 10 % fetal calf serum (FCS) (Biochrom, Ltd., Berlin, Germany) and 1 % penicillin-streptomycin (Invitrogen). The human colorectal adenocarcinoma cell line Caco-2 was provided by Jindrich Cinatl Jr. (Frankfurt am Main, Germany) and maintained in MEM medium (Sigma), supplemented with 10 % FCS, 1 % penicillin-streptomycin, 2 % L-glutamine (Gibco). HEK293T cells were cultivated in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10 % FCS and 1 % penicillin/streptomycin. Vero E6 African green monkey kidney cells were obtained from ATCC and maintained in DMEM supplemented with 10 % FCS and 1 % penicillin/streptomycin. Cell lines were cultivated in humidified atmosphere at 37 °C with 5 % CO₂ and sub-culturing of cells was performed two or three times a week. All cell lines were regularly negatively tested for mycoplasma.

IFN-stimulating DNA (ISD), the cationic lipid-based transfection reagent LyoVec and cyclic [G(2',5')pA(3',5')p] (2'3'-cGAMP) were obtained from Invivogen (San Diego, USA) and Lipofectamine2000 was obtained from ThermoFisher Scientific (Dreieich, Germany). All other chemicals were obtained from Carl Roth (Karlsruhe, Germany) or Sigma, unless stated otherwise.

CRISPR/Cas9 gene editing

CRISPR/Cas9 KO cells were generated as described previously⁵⁴. Briefly, three independent guide RNAs (gRNAs), targeting USP22 (#1:

GCCATTGATCTGATGTACGG, #2: CCTCGAACTGCACCATAGGT and #3: ACCTGGTGTGGACCCACGCG), TMEM173 (#1: CATTACAACAACCTGCTACG, #2: GCTGGGACTGCTGTAAACG, #3: GCAGGCACTCAGCAGAACCA), DDX58 (#1: CATCTTAAAAAATTCCCACA, #2: GGAACAAGTTCAGTGAAGT, #3: TGCATGCTCACTGATAATGA), IFIH1 (#1: CTTGGACATAACAGCAACAT, #2: TGAGTTCCAAAATCTGACAT) or TLR3 (#1: ACGACTGATGCTCCGAAGGG, #2: ACTTACCTTCTGCTTGACAA, #3: GGAAATAAATGGGACCACCA) and control gRNAs (Addgene plasmid #51763, #51762 and #51760) were ligated into pLentiCRISPRv2 (Addgene plasmid # 52961) using restriction cloning. Plasmid fidelity was confirmed using Sanger sequencing. For the generation of viral particles, multiple gene-specific gRNAs were combined and co-transfected with pMD2.G (Addgene plasmid #12259) and psPAX2 (Addgene plasmid #12260) in HEK293T cells using FuGENE HD Transfection Reagent (Promega), according to the manufacturer's protocol. Viral supernatants were collected 48- and 72-hours post-transfection, pooled and used for transduction in the presence of Polybrene (Sigma-Aldrich), followed by selection with puromycin (Thermo Fischer Scientific). Knockout was confirmed with Western blotting. Where necessary, single-cell clones were selected using limited dilution. Double-knockout (dKO) cells were generated by transduction with USP22-targeting virus first, followed by transduction with viral particles with gRNAs against the appropriate secondary targets and puromycin selection.

RNA isolation, cDNA synthesis and quantitative real-time PCR

Appropriate cell lines were seeded in 6-well plates (Greiner) 48 hours prior to RNA isolation, treated as indicated or left untreated, followed by extraction of total RNA using the peqGOLD total RNA isolation kit (PepLab, Erlangen, Germany), according to the manufacturer's protocol. In brief, cells were lysed in RNA lysis buffer, centrifuged

at 12000 x g for 2 min., followed by the addition of an equal volume of 70 % ethanol to the flow-through, after which RNA was bound to RNA-binding columns by centrifugation at 10000 x g for 1 min. Upon washing with RNA Wash Buffer I and two additional wash steps with 80 % ethanol, the column was dried by centrifuging at 12000 x g for 2 min. RNA was eluted with nuclease free water at 12000 x g for 2 min after which 1 µg of RNA was transcribed into cDNA using the RevertAid H Minus First Strand Kit (ThermoFisher Scientific) and random primers, according to the manufacturer's protocol. Relative mRNA expression levels were quantified using SYBR green-based quantitative real-time PCR (Applied Biosystems, Darmstadt, Germany) using the 7900GR fast real-time PCR system (Applied Biosystems). Data were normalized to 28S housekeeping expression and the relative expression of target gene transcripts levels were calculated compared to the reference transcript using the $\Delta\Delta CT$ method⁸³. At least three independent experiments in duplicates are shown. All primers were purchased at Eurofins (Hamburg, Germany). Primer sequences are shown in **Supplementary Table 1.**

Gene expression profiling

To quantify global changes in gene expression, RNA was isolated as described above, followed by a DNase digest upon RNA binding using the peqGOLD DNase Digest Kit, according to the manufacturer's instructions. Samples were processed and gene expression was profiled at the DKFZ Genomics and Proteomics Core Facility (Heidelberg, Germany) using the Affymetrix human Clariom S array.

Gene expression profiling analysis

Raw .CEL files were processed with the oligo R package⁸⁴ and the normalized intensities were obtained after RMA normalization. Genes with differential expression

between NHT control and USP22 KO have been identified using the linear model-based approach limma R package ⁸⁵. An adjusted P-value < 0.05 was considered significant. Gene-set enrichment analysis was performed with gage R package ⁸⁶ using the MSigDB ⁸⁷ as gene set repository. An adjusted P-value < 0.05 was considered significant.

Multiplex quantification of cytokine secretion

Cells were seeded in 2 ml cell culture medium and supernatant was collected after 66 h, centrifuged at 300 x g, 4 °C for 5 min. and frozen in liquid nitrogen. Samples were analyzed using the LEGENDplex™ Human Anti-Virus Response Panel multiplex assay (BioLegend, San Diego, CA, USA) following the manufacturer's protocol. The analysis was performed with the BD FACSVerse™ flow cytometer (BD Biosciences, San Jose, CA, USA). At least 300 events were acquired per analyte. The data was analyzed with the LEGENDplex v.8 software (BioLegend).

Western Blot analysis

The indicated cell lines were seeded two days before lysis and treated as indicated, or left untreated. Lysis was done on ice using RIPA lysis buffer (50 mM Tris-HCl pH 8, 150 mM NaCl, 1 % Nonidet P-40 (NP-40), 150 mM MgCl₂, 0.5 % sodium deoxycholate), with phosphatase inhibitors (1 mM sodium orthovanadate, 1 mM β-glycerophosphate, 5 mM sodium fluoride), protease inhibitor cocktail (Roche, Grenzach, Germany), 0.1 % sodium dodecyl sulfate (SDS) and Pierce Universal Nuclease (Thermo Fisher Scientific) for 30 min, followed by centrifugation at 18000 x g for 25 min. at 4 °C. Protein concentrations of the cell lysates were measured using the BCA Protein Assay Kit from Pierce™, according to the manufacturer's instructions.

For Western blot detection, 20-40 µg of the lysates were boiled in Laemmli loading buffer (6x Laemmli: 360 mM Tris Base pH 6.8, 30 % glycerol, 120 mg/ml SDS, 93 mg/ml dithiothreitol (DTT), 12 mg/ml bromophenol blue) at 95 °C for 5 min, followed by Western blot analysis. The following antibodies are used: rabbit anti-STING (13647S, Cell Signaling Beverly, MA, USA), rabbit anti-phospho-STAT1 (9167L, Cell Signaling), mouse anti-STAT1 (9176S, Cell signaling), rabbit anti-USP22 (#ab195298, Abcam), mouse anti-glyceraldehyde 3-phosphate dehydrogenase (GAPDH) (5G4cc, HyTest, Turku, Finland), mouse anti-Vinculin (#V9131-100UL, Merck), rabbit anti-TBK1 (ab40676, Abcam), rabbit anti-phospho-TBK1 (ab109272, Abcam), rabbit anti-Histone H2B (#07-371, Merck), mouse anti-Ubiquitinyl-Histone H2B (#05-1312, Merck), rabbit anti-p65 (sc-372X, Santa Cruz Biotechnologies, Santa Cruz, CA, USA), rabbit anti-phospho-p65 (3033S, Cell Signaling), mouse anti-IRF3 (sc-33641, Santa Cruz), rabbit anti-phospho-IRF3 (4947S, Cell Signaling), rabbit anti-RIG-I (3743S, Cell Signaling), rabbit anti-MDA5 (5321S, Cell Signaling), rabbit anti-TLR3 (6961S, Cell Signaling), mouse anti-ISG56 (PA3-848, Thermo scientific), rabbit anti-MX1 (37849S, Cell Signaling), rabbit anti-IRF9 (76684S, Cell Signaling), rabbit anti-ISG20 (PA5-30073, Thermo scientific), rabbit anti-γ-H2AX (phospho Ser139) (NB100-384, Novus Biologicals) and mouse anti-NF-κB p52 (05-361, Millipore). Secondary antibodies labeled with horseradish peroxidase (HRP) were used for detection with enhanced chemiluminescence (Amersham Bioscience, Freiburg, Germany). HRP-conjugated goat anti-mouse IgG (ab6789, Abcam) was diluted 1:10000 and HRP-conjugated goat anti-rabbit IgG (ab6721, Abcam) was diluted 1:30000 in 5 % milk powder in PBS with 0.2 % Tween 20 (PBS-T). When necessary, membranes were stripped using 0.4 M NaOH for 10 min, followed by 1 h of blocking and incubation with a second primary antibody. Representative blots of at least two independent experiments are shown. When detected on separate membranes, only one representative loading control is

shown for clarity.

Stimulation of STING with 2'3'-cGAMP

The indicated cell lines were seeded 24 or 48 hours prior to stimulation in P/S-free cell culture medium. For stimulation, culture medium was removed and cell lines were permeabilized by incubation with digitonin buffer (50 mM HEPES, 100 mM KCl, 3 mM MgCl₂, 0.1 mM dithiothreitol, 85 mM sucrose, 0.2 % bovine serum albumin, 1 mM ATP, 5 µg/ml Digitonin) pH 7 in the presence or absence of 10 µg/ml 2'3'-cGAMP for 10 min. at 37 °C. After incubation, the permeabilization buffer was replaced with P/S-free cell culture medium and further incubated at 37 °C/5 % CO₂ for the indicated time points.

PRR stimulation with poly(I:C) and ISD

The indicated HT-29 cells were seeded 24 hours prior to treatment in sterile 6-well plates (Greiner). For each well, two µg of ISD (Invivogen) were pre-mixed with OptiMEM and, after 5 min. incubation at room temperature, mixed with premixed Lipofectamine2000-OptiMEM at a ratio of 3:1, according to the manufacturer's instructions. After incubation for 15 min. at room temperature, the indicated transfection mixes were added to the cells in P/S free medium. Cell lysis with RIPA or RNA lysis buffer was performed after 24 h. For stimulation with poly(I:C), the indicated HT-29 cells were seeded as described above and for each well, 2 µg of poly(I:C) was mixed with 20 µl LyoVec (Invivogen), incubated for 15 min. at room temperature to allow the formation of lipid-RNA complexes. The transfection mix was then added to the indicated HT-29 cells in P/S free medium at a 1:20 volume ratio and incubated for 24 h, after which cells were processed for Western blot or RNA isolation.

Tandem Ubiquitin Binding Entity (TUBE) pull-down analysis

Ubiquitinated proteins were enriched using GST-tagged tandem-repeated ubiquitin-binding entities (TUBEs)⁸⁸, as described before⁵⁴. Briefly, the indicated cell lines were seeded 48 hours prior to lysis and/or treatment, harvested in NP-40 lysis buffer (50 mM NaCl, 20 mM Tris pH 7.5, 1 % NP-40, 5 mM EDTA, 10 % Glycerol) supplemented with 25 mM NEM, 1 mM sodium orthovanadate, 1 mM sodium fluoride, 0.5 mM phenylmethylsulfonyl fluoride, protease inhibitor cocktail and Pierce Universal Nuclease on ice for 30 min. GST-TUBE beads were washed once with NP-40 buffer and incubated with 3 mg of protein lysate over night at 4 °C. Beads were washed four times with NP-40 buffer, followed by elution of ubiquitinated proteins by boiling in 2x Laemmli loading buffer at 96 °C for 6 min. Ubiquitinated proteins were analyzed using Western blot analysis.

SARS-CoV-2 infection

SARS-CoV-2 (strain BavPat1/2020) was obtained from the European Virology Archive and amplified in Vero E6 cells and used at passage 3. Virus titers were determined by TCID₅₀ assay. Caco-2 cells were infected using a MOI of 1 virus particle per cell. Medium was removed from Caco-2 cells and virus was added to cells for 1 h at 37°C. Viral supernatants were removed, infected cells were washed once with PBS and media was added back to the cells. Virus infection was monitored 24 h post-infection.

TCID₅₀ virus titration

Vero E6 cells were seeded (20000 per well) in 96-well plates 24 h prior to infection. A volume of 100 µl of viral supernatant from the indicated SARS-CoV-2-infected Caco-2 cells was added to the first well. Seven 1:10 dilutions were made (all samples were performed in triplicate). Infections were allowed to proceed for 24 h. At 24 h post infection (hpi), cells were fixed in 2 % paraformaldehyde (PFA) for 20 minutes at room

temperature. PFA was removed and cells were washed twice in PBS and permeabilized for 10 min. at room temperature in 0,5 % Triton-X/PBS. Cells were blocked in a 1:2 dilution of LI-COR blocking buffer (LI-COR, Lincoln, NE, USA) for 30 min at room temperature. Infected cells were stained with 1:1000 diluted anti-dsRNA (J2) for 1 h at room temperature, washed three times with 0.1 % PBT-T, followed by incubation with secondary antibody (anti-mouse CW800) and DNA dye Draq5 (Abcam, Cambridge, UK), diluted 1:10000 in blocking buffer and incubated for 1 h at room temperature. Cells were washed three times with 0.1 % PBS-T and imaged in PBS on a LI-COR imager.

Quantification of viral RNA

At 24 hpi, RNA was extracted from infected or mock-treated Caco-2 cells using the Qiagen RNAeasy Plus Extraction Kit (Qiagen, Hilden, Germany). For quantifying the SARS-CoV-2 genome abundance in mock and infected samples, cDNA was generated using 250 ng of RNA with the iSCRIPT reverse transcriptase (BioRad, Hercules, CA, USA), according to the manufacturer's instructions. qRT-PCR was performed using iTaq SYBR green (BioRad) following the instructions of the manufacturer and normalized on TBP. Primers were ordered at Eurofins, Luxemburg and are listed in **Supplementary Table 1.**

Indirect Immunofluorescence Assay

Cells seeded on iBIDI glass bottom 8-well chamber slides. At 24 post-infection, cells were fixed in 4% paraformaldehyde (PFA) for 20 mins at room temperature (RT). Cells were washed and permeabilized in 0.5% Triton-X for 15 mins at RT. Primary antibody SARS-CoV NP (Sino biologicals MM05) were diluted in phosphate-buffered saline (PBS) and incubated for 1h at RT. Cells were washed in 1X PBS three times and

incubated with secondary antibodies goat-anti mouse Alexa Fluor 568 and DAPI for 45 mins at RT. Cells were washed in 1X PBS three times and maintained in PBS. Cells were imaged by epifluorescence on a Nikon Eclipse Ti-S (Nikon).

Statistical analysis

Significance was assessed using Student's t-test (two-tailed distribution, two-sample, equal variance) using Microsoft Excel, unless indicated otherwise. *P*-values < 0.05 are considered significant (* *P* < 0.05; ** *P* < 0.01; *** *P* < 0.001, n.s.: not significant).

Resource availability

Further information and requests for resources and reagents should be directed to and will be fulfilled by the corresponding author, Sjoerd J. L. van Wijk (vanWijk@med.uni-frankfurt.de; s.wijk@kinderkrebsstiftung-frankfurt.de).

Materials availability

All unique reagents generated in this study are available from the corresponding author without restriction.

Data and code availability

Microarray data are available on Gene Expression Omnibus under the accession number GSE190036.

Acknowledgments

The authors thank the members of the van Wijk lab for advice, discussions and support during the study, Dr. M. Bewerunge-Hudler and her team from the Genomics and Proteomics Core Facility, German Cancer Research Center/DKFZ, Heidelberg, Germany for help and support with performing the microarray analysis and Christina Hugenberg for proofreading. S.J.L.v.W. is supported by the Deutsche Forschungsgemeinschaft (DFG) (WI 5171/1-1, FU 436/20-1 and project-ID 259130777 – SFB 1177), the Deutsche Krebshilfe (70113680), the Frankfurter Stiftung für krebskranke Kinder and the Dr. Eberhard and Hilde Rüdiger Foundation. M.B. is supported by the DFG – CRC 850 subprojects C9 and Z1, CRC1479 (Project ID: 441891347- S1), CRC 1160 (Project Z02), CRC1453 (Project ID 431984000 - S1) and TRR167 (Project Z01), the German Federal Ministry of Education and Research by MIRACUM within the Medical Informatics Funding Scheme (FKZ 01ZZ1801B). S.B. was supported by DFG project numbers 415089553 (Heisenberg program), 240245660 (SFB1129), 278001972 (TRR186), and 272983813 (TRR179), the state of Baden-Württemberg (AZ: 33.7533.-6-21/5/1), the BMBF (01KI20198A) and within the Network University Medicine - Organo-Strat COVID-19. M.L.S. was supported by the BMBF (01KI20239B) and DFG project 416072091.

Author contributions

R.K. performed experiments and analyzed data with help from J.R., S.S. and L.K., M.L.S and S.B. performed SARS-CoV-2 infections and accompanying experiments, gene expression analysis was performed by G.A. and M.B., R.S. provided access and support with the LEGENDplex analysis. D.B and J.C.Jr. provided the Caco-2 cell line

697 and expertise. R.K. and S.J.L.v.W. conceived the project and wrote the manuscript. All
698 authors have read, commented and agreed on the submitted version of the manuscript.

699

700 **Declaration of interest**

701 The authors declare no competing interests.

702

Figure legends

Figure 1: Profiling USP22-mediated gene expression in HT-29 hIECs. A. Volcano plot showing the differential gene expression patterns of two independent single-cell HT-29 USP22 CRISPR/Cas9 KO clones (#16 and #62) compared to CRISPR/Cas9 control (NHT) HT-29 cells. Color code represents the log2 foldchange compared to NHT. **B.** Heatmap of the top-50 differentially regulated genes between HT-29 USP22 KO single clones #16 and #62 and the NHT control. Color coding represents the row-wise scaled (Z-score) RNA intensities. Genes are sorted according to their log2 fold change, compared to NHT. **C.** Basal mRNA expression levels of the indicated genes were determined in control and two independent USP22 KO HT-29 single clones using qRT-PCR. Gene expression was normalized against 28S mRNA and is presented as x-fold mRNA expression compared to NHT. Mean and SD of three independent experiments in triplicate are shown. *P < 0.05; **P < 0.01, ***P < 0.001.

Figure 2: Loss of USP22 specifically enriches for genes involved in interferon signaling and response to viral infection. A. Bar plot showing the top-20 regulated Gene Ontology (GO) terms in two independent single-cell HT-29 USP22 CRISPR/Cas9 KO clones (#16 and #62) compared to control (non-human target: NHT) HT-29 cells. Color code represents the number of annotated genes within each gene set. **B.** Heatmap of differentially expressed genes contributing to the GO terms *response to type I interferon* (left) and *interferon gamma mediated signaling pathway* (right). Color code represents the log2 foldchange compared to NHT. **C.** Basal mRNA expression levels of GO enriched genes related to IFN signaling in control (NHT) and two independent USP22 KO HT-29 single clones using qRT-PCR. Gene expression was normalized against 28S mRNA and is presented as x-fold mRNA expression compared to NHT. Mean and SD of three independent experiments in triplicate are

shown. *P < 0.05; **P < 0.01, ***P < 0.001, n.s. not significant. **D.** Western blot analysis of basal MX1, IRF9, ISG56, ISG20 and USP22 expression levels in control and USP22 KO HT-29 cells (clone USP22 KO #62). GAPDH served as loading control. Representative blots of at least two different independent experiments are shown.

Figure 3: USP22 negatively regulates STAT1 signaling and IFN- λ 1 expression. A.

Basal mRNA expression levels of total IFNA (panIFNA) and IFNB1 in control (non-human target: NHT) and the CRISPR/Cas9-generated USP22 knock-out (KO) HT-29 single clone (USP22 KO #62). Gene expression was normalized against 28S mRNA and is presented as x-fold mRNA expression compared to NHT. Mean and SD of three independent experiments in triplicate are shown. *P < 0.05; **P < 0.01. **B.** Western blot analysis of basal phosphorylated and total levels of STAT1 and USP22 in control and USP22 KO HT-29 cells (USP22 KO #62). GAPDH served as loading control. Representative blots of at least two different independent experiments are shown. **C.** FACS-based analysis of the indicated basal secretion patterns of the viral defense cytokine panel in supernatants of control and USP22 KO HT-29 cells (USP22 KO #62). Data are presented as absolute levels of cytokines (in pg/ml). Samples below lower detection limit were set to zero, values above upper detection limit were set to detection limit. Mean and SD of three independent experiments in triplicate are shown. *P < 0.05; n.s. not significant. **D.** Basal mRNA expression levels of IFNL1 in control and USP22 KO HT-29 single clone (USP22 KO #62). Gene expression was normalized against 28S mRNA and is presented as x-fold mRNA expression compared to NHT. Mean and SD of three independent experiments in triplicate are shown. *P < 0.05.

Figure 4: USP22 regulates type III IFN signaling via STING. A. Western blot analysis of basal RIG-I, MDA5, TLR3 and USP22 expression levels in control (non-

human target: NHT) and the CRISPR/Cas9-generated USP22 knock-out (KO) HT-29 single clone (USP22 KO #62). GAPDH served as loading control. Representative blots of at least two different independent experiments are shown. **B.** Western blot analysis of basal RIG-I, phosphorylated and total STAT1, ISG56 and USP22 expression levels in control, USP22 KO HT-29 cells (USP22 KO #62) as well as two NHT-control and one USP22-DDX58 double KO (dKO) HT-29 single clones. GAPDH served as loading control. Representative blots of at least two different independent experiments are shown. **C.** Idem as **B.**, one MDA5/IFIH1 KO single clone instead of RIG-I/DDX58. **D.** Idem as **B.** three TLR3 KO single clones instead of RIG-I/DDX58. **E.** Western blot analysis of phosphorylated and total STAT1, RIG-I, STING and USP22 expression levels in control and USP22 KO HT-29 cells (USP22 KO #62) subjected to transfection with transfection reagent (control) alone or in the presence of ISD and poly(I:C) (2 µg/well) for 24 h. Vinculin served as loading control. Representative blots of at least two different independent experiments are shown. **F.** mRNA expression levels of OAS3 (*left*) and IRF9 (*right*) in control and USP22 knock-out (KO) HT-29 cells (USP22 KO #62) subjected to transfection with ISD and poly(I:C) (each 2 µg/well) for 24 h. Gene expression was normalized against 28S mRNA and is presented as x-fold mRNA expression compared to NHT. Mean and SD of three independent experiments in triplicate are shown. *P < 0.05; **P < 0.01; n.s. not significant. **G.** Western blot analysis of phosphorylated and total STAT1, STING and USP22 expression levels in control, USP22 KO HT-29 cells (USP22 KO #62) as well as in the indicated NHT, USP22, control and STING dKO cells. GAPDH served as loading control. Representative blots of at least two different independent experiments are shown. **H.** Basal mRNA expression levels of the indicated genes in control USP22 KO HT-29 cells (USP22 KO #62) as well as in the indicated NHT, USP22, control and STING dKO cells. Gene expression was normalized against 28S mRNA and is presented as x-fold mRNA

expression compared to NHT. Mean and SD of three independent experiments in triplicate are shown. $**P < 0.01$; $***P < 0.001$. **I.** Basal mRNA expression levels of IFNA, IFNB and IFNL1 in control USP22 KO HT-29 cells (USP22 KO #62) as well as in the indicated NHT, USP22, control and STING dKO cells. Gene expression was normalized against 28S mRNA and is presented as x-fold mRNA expression compared to NHT. Mean and SD of three independent experiments in triplicate are shown. $**P < 0.01$; $***P < 0.001$.

Figure 5: USP22 negatively regulates STING activation and ubiquitination. A.

Western blot analysis of STING, phosphorylated and total TBK1, phosphorylated and total IRF3 and USP22 expression levels in control (non-human target: NHT) and CRISPR/Cas9-generated USP22 knock-out (KO) HT-29 single clone (USP22 KO #62) subjected to 2'3'-cGAMP (10 μ g/ml) for the indicated timepoints. GAPDH served as loading control. Representative blots of at least two different independent experiments are shown. **B.** mRNA expression levels of IFNA, IFNB and IFNL1 in control and USP22 KO HT-29 cells (USP22 KO #62) subjected to 2'3'-cGAMP (10 μ g/ml) for 3 h. Gene expression was normalized against 28S mRNA and is presented as x-fold mRNA expression compared to NHT. Mean and SD of three independent experiments in triplicate are shown. $*P < 0.05$; $**P < 0.01$. **C.** Western blot analysis of STING, phosphorylated and total STAT1 and USP22 expression levels in control and USP22 KO HT-29 cells (USP22 KO #62) subjected to the JAK/STAT inhibitor ruxolitinib (5 μ M) for the indicated timepoints. GAPDH served as loading control. Representative blots of at least two different independent experiments are shown. **D.** Western blot analysis of STING and USP22 expression levels in control and USP22 KO HT-29 cells (USP22 KO #62) subjected to cycloheximide (CHX) (100 μ g/ml) for the indicated timepoints. Vinculin served as loading control. Representative blots of at least two different

independent experiments are shown. **E.** Western blot analysis of Tandem Ubiquitin Binding Entity (TUBE)-enriched ubiquitin-modified STING from control and USP22 KO HT-29 cells (USP22 KO #62) subjected to 2'3'-cGAMP (10 µg/ml) for 24 h. GAPDH served as loading control and Ponceau S staining confirms equal loading of GST-TUBE beads. Representative blots of at least two different independent experiments are shown.

Figure 6: Loss of USP22 protects against SARS-CoV-2 infection, replication and de novo infectious virus production in a STING-dependent manner. A. Western blot analysis of phosphorylated and total STAT1, STING and USP22 expression levels in wild-type (WT), control (non-human target: NHT) and two CRISPR/Cas9-generated USP22 knock-out (KO) Caco-2 single clones (USP22 KO #1 and #6). GAPDH served as loading control. Representative blots of at least two different independent experiments are shown. **B.** Basal mRNA expression levels of IRF9 and OAS3 in WT, control and USP22 KO Caco-2 cells (USP22 KO #1 and #6). Gene expression was normalized against 28S mRNA and is presented as x-fold mRNA expression compared to NHT. Mean and SD of three independent experiments in triplicate are shown. *P < 0.05; **P < 0.01, n.s. not significant. **C.** Quantification of immunofluorescence-stained SARS-CoV-2-infected cells, normalized against non-infected cells. WT, control and USP22 KO Caco-2 cells (USP22 KO #1 and #6) were stained with anti-dsRNA (J2) at 24 hpi. Mean and SD of three independent experiments in triplicate are shown. ***P < 0.001. **D.** Quantification of relative SARS-CoV-2 genome expression of SARS-CoV-2-infected WT, control and USP22 KO Caco-2 cells (USP22 KO #1 and #6) at 6 hpi (left) and 24 hpi (right). Data are normalized against non-infected cells. Mean and SD of three independent experiments in triplicate are shown. ***P < 0.001. **E.** Western blot analysis of STING and USP22 expression levels in control-NHT, control-USP22 KO

833 #1 and #6, STING-NHT and STING-USP22 KO #1 and #6 double KO (dKO) HT-29
 834 cells. Vinculin served as loading control. Representative blots of at least two different
 835 independent experiments are shown. **F.** Quantification of relative SARS-CoV-2
 836 genome expression of SARS-CoV-2-infected control-NHT, control-USP22 KO #1 and
 837 #6, STING-NHT and STING-USP22 KO #1 and #6 dKO HT-29 cells at 24 hpi. Mean
 838 and SD of three independent experiments in triplicate are shown. **P < 0.005.
 839

References

- 1 Takeuchi, O. & Akira, S. Pattern recognition receptors and inflammation. *Cell* **140**, 805-820, doi:10.1016/j.cell.2010.01.022 (2010).
- 2 Li, D. & Wu, M. Pattern recognition receptors in health and diseases. *Signal Transduct Target Ther* **6**, 291, doi:10.1038/s41392-021-00687-0 (2021).
- 3 Pichlmair, A. & Reis e Sousa, C. Innate recognition of viruses. *Immunity* **27**, 370-383, doi:10.1016/j.immuni.2007.08.012 (2007).
- 4 Kato, H. *et al.* Length-dependent recognition of double-stranded ribonucleic acids by retinoic acid-inducible gene-I and melanoma differentiation-associated gene 5. *The Journal of experimental medicine* **205**, 1601-1610, doi:10.1084/jem.20080091 (2008).
- 5 Decout, A., Katz, J. D., Venkatraman, S. & Ablasser, A. The cGAS-STING pathway as a therapeutic target in inflammatory diseases. *Nat Rev Immunol* **21**, 548-569, doi:10.1038/s41577-021-00524-z (2021).
- 6 Mesev, E. V., LeDesma, R. A. & Ploss, A. Decoding type I and III interferon signalling during viral infection. *Nature Microbiology* **4**, 914-924, doi:10.1038/s41564-019-0421-x (2019).
- 7 Lazear, H. M., Schoggins, J. W. & Diamond, M. S. Shared and Distinct Functions of Type I and Type III Interferons. *Immunity* **50**, 907-923, doi:10.1016/j.immuni.2019.03.025 (2019).
- 8 Kotenko, S. V., Rivera, A., Parker, D. & Durbin, J. E. Type III IFNs: Beyond antiviral protection. *Semin Immunol* **43**, 101303, doi:10.1016/j.smim.2019.101303 (2019).
- 9 Schoggins, J. W. & Rice, C. M. Interferon-stimulated genes and their antiviral effector functions. *Curr Opin Virol* **1**, 519-525, doi:10.1016/j.coviro.2011.10.008 (2011).

- 10 Park, A. & Iwasaki, A. Type I and Type III Interferons - Induction, Signaling, Evasion, and Application to Combat COVID-19. *Cell Host Microbe* **27**, 870-878, doi:10.1016/j.chom.2020.05.008 (2020).
- 11 Ye, L., Schnepf, D. & Staeheli, P. Interferon-lambda orchestrates innate and adaptive mucosal immune responses. *Nat Rev Immunol* **19**, 614-625, doi:10.1038/s41577-019-0182-z (2019).
- 12 Mesev, E. V., LeDesma, R. A. & Ploss, A. Decoding type I and III interferon signalling during viral infection. *Nat Microbiol* **4**, 914-924, doi:10.1038/s41564-019-0421-x (2019).
- 13 Forero, A. *et al.* Differential Activation of the Transcription Factor IRF1 Underlies the Distinct Immune Responses Elicited by Type I and Type III Interferons. *Immunity* **51**, 451-464 e456, doi:10.1016/j.immuni.2019.07.007 (2019).
- 14 Galani, I. E. *et al.* Interferon-lambda Mediates Non-redundant Front-Line Antiviral Protection against Influenza Virus Infection without Compromising Host Fitness. *Immunity* **46**, 875-890 e876, doi:10.1016/j.immuni.2017.04.025 (2017).
- 15 Fung, T. S. & Liu, D. X. Human Coronavirus: Host-Pathogen Interaction. *Annu Rev Microbiol* **73**, 529-557, doi:10.1146/annurev-micro-020518-115759 (2019).
- 16 Neufeldt, C. J. *et al.* SARS-CoV-2 infection induces a pro-inflammatory cytokine response through cGAS-STING and NF- κ B. *bioRxiv*, 2020.2007.2021.212639, doi:10.1101/2020.07.21.212639 (2020).
- 17 Tay, M. Z., Poh, C. M., Rénia, L., MacAry, P. A. & Ng, L. F. P. The trinity of COVID-19: immunity, inflammation and intervention. *Nature Reviews Immunology* **20**, 363-374, doi:10.1038/s41577-020-0311-8 (2020).
- 18 Ivashkiv, L. B. & Donlin, L. T. Regulation of type I interferon responses. *Nat Rev Immunol* **14**, 36-49, doi:10.1038/nri3581 (2014).

- 19 Stanifer, M. L. *et al.* Critical Role of Type III Interferon in Controlling SARS-CoV-2 Infection in Human Intestinal Epithelial Cells. *Cell reports* **32**, 107863, doi:10.1016/j.celrep.2020.107863 (2020).
- 20 Triana, S. *et al.* Single-cell analyses reveal SARS-CoV-2 interference with intrinsic immune response in the human gut. *Molecular Systems Biology* **17**, e10232, doi:<https://doi.org/10.15252/msb.202110232> (2021).
- 21 Erlandsson, L. *et al.* Interferon-beta is required for interferon-alpha production in mouse fibroblasts. *Curr Biol* **8**, 223-226, doi:10.1016/s0960-9822(98)70086-7 (1998).
- 22 Phipps-Yonas, H., Seto, J., Sealfon, S. C., Moran, T. M. & Fernandez-Sesma, A. Interferon-beta pretreatment of conventional and plasmacytoid human dendritic cells enhances their activation by influenza virus. *PLoS Pathog* **4**, e1000193, doi:10.1371/journal.ppat.1000193 (2008).
- 23 Stewart, W. E., 2nd, Gosser, L. B. & Lockart, R. Z., Jr. Priming: a nonantiviral function of interferon. *J Virol* **7**, 792-801, doi:10.1128/JVI.7.6.792-801.1971 (1971).
- 24 Kuri, T. *et al.* Interferon priming enables cells to partially overturn the SARS coronavirus-induced block in innate immune activation. *J Gen Virol* **90**, 2686-2694, doi:10.1099/vir.0.013599-0 (2009).
- 25 Felgenhauer, U. *et al.* Inhibition of SARS-CoV-2 by type I and type III interferons. *J Biol Chem* **295**, 13958-13964, doi:10.1074/jbc.AC120.013788 (2020).
- 26 Stanifer, M. L., Guo, C., Doldan, P. & Boulant, S. Importance of Type I and III Interferons at Respiratory and Intestinal Barrier Surfaces. *Frontiers in immunology* **11**, 608645, doi:10.3389/fimmu.2020.608645 (2020).

- 27 Vanderheiden, A. *et al.* Type I and Type III Interferons Restrict SARS-CoV-2 Infection of Human Airway Epithelial Cultures. *J Virol* **94**, doi:10.1128/JVI.00985-20 (2020).
- 28 Busnadiego, I. *et al.* Antiviral Activity of Type I, II, and III Interferons Counterbalances ACE2 Inducibility and Restricts SARS-CoV-2. *mBio* **11**, doi:10.1128/mBio.01928-20 (2020).
- 29 Lamers, M. M. *et al.* An organoid-derived bronchioalveolar model for SARS-CoV-2 infection of human alveolar type II-like cells. *EMBO J* **40**, e105912, doi:10.15252/emboj.2020105912 (2021).
- 30 Zhu, Q. *et al.* Inhibition of coronavirus infection by a synthetic STING agonist in primary human airway system. *Antiviral Research* **187**, 105015, doi:<https://doi.org/10.1016/j.antiviral.2021.105015> (2021).
- 31 Liu, W. *et al.* Activation of STING Signaling Pathway Effectively Blocks Human Coronavirus Infection. *J Virol* **95**, doi:10.1128/jvi.00490-21 (2021).
- 32 Li, M. *et al.* Pharmacological activation of STING blocks SARS-CoV-2 infection. *Science Immunology* **6**, eabi9007, doi:10.1126/sciimmunol.abi9007 (2021).
- 33 Humphries, F. *et al.* A diamidobenzimidazole STING agonist protects against SARS-CoV-2 infection. *Science Immunology* **6**, eabi9002, doi:10.1126/sciimmunol.abi9002 (2021).
- 34 Broggi, A. *et al.* Type III interferons disrupt the lung epithelial barrier upon viral recognition. *Science* **369**, 706-712, doi:10.1126/science.abc3545 (2020).
- 35 Major, J. *et al.* Type I and III interferons disrupt lung epithelial repair during recovery from viral infection. *Science* **369**, 712-717, doi:10.1126/science.abc2061 (2020).

- 36 Isaacson, M. K. & Ploegh, H. L. Ubiquitination, ubiquitin-like modifiers, and
deubiquitination in viral infection. *Cell Host Microbe* **5**, 559-570,
doi:10.1016/j.chom.2009.05.012 (2009).
- 37 van Huizen, M. & Kikkert, M. The Role of Atypical Ubiquitin Chains in the
Regulation of the Antiviral Innate Immune Response. *Front Cell Dev Biol* **7**, 392,
doi:10.3389/fcell.2019.00392 (2019).
- 38 Heaton, S. M., Borg, N. A. & Dixit, V. M. Ubiquitin in the activation and
attenuation of innate antiviral immunity. *The Journal of experimental medicine*
213, 1-13, doi:10.1084/jem.20151531 (2016).
- 39 Tsuchida, T. *et al.* The ubiquitin ligase TRIM56 regulates innate immune
responses to intracellular double-stranded DNA. *Immunity* **33**, 765-776,
doi:10.1016/j.immuni.2010.10.013 (2010).
- 40 Zhang, J., Hu, M. M., Wang, Y. Y. & Shu, H. B. TRIM32 protein modulates type
I interferon induction and cellular antiviral response by targeting MITA/STING
protein for K63-linked ubiquitination. *J Biol Chem* **287**, 28646-28655,
doi:10.1074/jbc.M112.362608 (2012).
- 41 Ni, G., Konno, H. & Barber, G. N. Ubiquitination of STING at lysine 224 controls
IRF3 activation. *Sci Immunol* **2**, doi:10.1126/sciimmunol.aah7119 (2017).
- 42 Wang, Q. *et al.* The E3 ubiquitin ligase AMFR and INSIG1 bridge the activation
of TBK1 kinase by modifying the adaptor STING. *Immunity* **41**, 919-933,
doi:10.1016/j.immuni.2014.11.011 (2014).
- 43 Zhong, B. *et al.* The ubiquitin ligase RNF5 regulates antiviral responses by
mediating degradation of the adaptor protein MITA. *Immunity* **30**, 397-407,
doi:10.1016/j.immuni.2009.01.008 (2009).

- 44 Xing, J. *et al.* TRIM29 promotes DNA virus infections by inhibiting innate immune response. *Nature Communications* **8**, 945, doi:10.1038/s41467-017-00101-w (2017).
- 45 Fenech, E. J. *et al.* Interaction mapping of endoplasmic reticulum ubiquitin ligases identifies modulators of innate immune signalling. *eLife* **9**, doi:10.7554/eLife.57306 (2020).
- 46 Sun, H. *et al.* USP13 negatively regulates antiviral responses by deubiquitinating STING. *Nature Communications* **8**, 15534, doi:10.1038/ncomms15534 (2017).
- 47 Zhang, J. *et al.* Deubiquitinase USP35 restrains STING-mediated interferon signaling in ovarian cancer. *Cell Death & Differentiation* **28**, 139-155, doi:10.1038/s41418-020-0588-y (2021).
- 48 Zhang, L. *et al.* The deubiquitinase CYLD is a specific checkpoint of the STING antiviral signaling pathway. *PLOS Pathogens* **14**, e1007435, doi:10.1371/journal.ppat.1007435 (2018).
- 49 Zhang, X. Y. *et al.* The putative cancer stem cell marker USP22 is a subunit of the human SAGA complex required for activated transcription and cell-cycle progression. *Mol Cell* **29**, 102-111, doi:10.1016/j.molcel.2007.12.015 (2008).
- 50 Zhao, Y. *et al.* A TFTC/STAGA module mediates histone H2A and H2B deubiquitination, coactivates nuclear receptors, and counteracts heterochromatin silencing. *Molecular cell* **29**, 92-101, doi:10.1016/j.molcel.2007.12.011 (2008).
- 51 Zhang, X. Y., Pfeiffer, H. K., Thorne, A. W. & McMahon, S. B. USP22, an hSAGA subunit and potential cancer stem cell marker, reverses the polycomb-catalyzed ubiquitylation of histone H2A. *Cell Cycle* **7**, 1522-1524, doi:10.4161/cc.7.11.5962 (2008).

- 52 Kosinsky, R. L. *et al.* USP22-dependent HSP90AB1 expression promotes resistance to HSP90 inhibition in mammary and colorectal cancer. *Cell Death Dis* **10**, 911, doi:10.1038/s41419-019-2141-9 (2019).
- 53 Kosinsky, R. L. *et al.* USP22 exerts tumor-suppressive functions in colorectal cancer by decreasing mTOR activity. *Cell Death Differ*, doi:10.1038/s41418-019-0420-8 (2019).
- 54 Roedig, J. *et al.* USP22 controls necroptosis by regulating receptor-interacting protein kinase 3 ubiquitination. *EMBO Rep* **22**, e50163, doi:10.15252/embr.202050163 (2021).
- 55 Cai, Z. *et al.* USP22 promotes IRF3 nuclear translocation and antiviral responses by deubiquitinating the importin protein KPNA2. *The Journal of experimental medicine* **217**, doi:10.1084/jem.20191174 (2020).
- 56 Liu, Q. *et al.* Broad and diverse mechanisms used by deubiquitinase family members in regulating the type I interferon signaling pathway during antiviral responses. *Sci Adv* **4**, eaar2824, doi:10.1126/sciadv.aar2824 (2018).
- 57 Atanassov, B. S. *et al.* ATXN7L3 and ENY2 Coordinate Activity of Multiple H2B Deubiquitinases Important for Cellular Proliferation and Tumor Growth. *Mol Cell* **62**, 558-571, doi:10.1016/j.molcel.2016.03.030 (2016).
- 58 Lanning, N. J. *et al.* A mitochondrial RNAi screen defines cellular bioenergetic determinants and identifies an adenylate kinase as a key regulator of ATP levels. *Cell Rep* **7**, 907-917, doi:10.1016/j.celrep.2014.03.065 (2014).
- 59 Sasaki, N. *et al.* Reg4⁺ deep crypt secretory cells function as epithelial niche for Lgr5⁺ stem cells in colon. *Proc Natl Acad Sci U S A* **113**, E5399-5407, doi:10.1073/pnas.1607327113 (2016).

1014 60 Dietlein, N. *et al.* Loss of Usp22 enhances histone H2B monoubiquitination and
1015 stimulates intracellular and systemic interferon immunity. *bioRxiv*,
1016 2021.2004.2009.439190, doi:10.1101/2021.04.09.439190 (2021).

1017 61 Hong, A., Lee, J. E. & Chung, K. C. Ubiquitin-Specific Protease 22 (USP22)
1018 Positively Regulates RCAN1 Protein Levels Through RCAN1 De-Ubiquitination.
1019 *J Cell Physiol* **230**, 1651-1660, doi:10.1002/jcp.24917 (2015).

1020 62 Basters, A., Knobloch, K. P. & Fritz, G. USP18 - a multifunctional component
1021 in the interferon response. *Biosci Rep* **38**, doi:10.1042/BSR20180250 (2018).

1022 63 Gough, D. J., Messina, N. L., Clarke, C. J. P., Johnstone, R. W. & Levy, D. E.
1023 Constitutive type I interferon modulates homeostatic balance through tonic
1024 signaling. *Immunity* **36**, 166-174, doi:10.1016/j.immuni.2012.01.011 (2012).

1025 64 Cheon, H. & Stark, G. R. Unphosphorylated STAT1 prolongs the expression of
1026 interferon-induced immune regulatory genes. *Proc Natl Acad Sci U S A* **106**,
1027 9373-9378, doi:10.1073/pnas.0903487106 (2009).

1028 65 Kotenko, S. V. *et al.* IFN-lambdas mediate antiviral protection through a distinct
1029 class II cytokine receptor complex. *Nat Immunol* **4**, 69-77, doi:10.1038/ni875
1030 (2003).

1031 66 Sheppard, P. *et al.* IL-28, IL-29 and their class II cytokine receptor IL-28R. *Nat*
1032 *Immunol* **4**, 63-68, doi:10.1038/ni873 (2003).

1033 67 Brzostek-Racine, S., Gordon, C., Van Scoy, S. & Reich, N. C. The DNA damage
1034 response induces IFN. *Journal of immunology* **187**, 5336-5345,
1035 doi:10.4049/jimmunol.1100040 (2011).

1036 68 Ramachandran, S. *et al.* The SAGA Deubiquitination Module Promotes DNA
1037 Repair and Class Switch Recombination through ATM and DNAPK-Mediated
1038 gammaH2AX Formation. *Cell Rep* **15**, 1554-1565,
1039 doi:10.1016/j.celrep.2016.04.041 (2016).

1040 69 Li, C. *et al.* The H2B deubiquitinase Usp22 promotes antibody class switch
1041 recombination by facilitating non-homologous end joining. *Nat Commun* **9**,
1042 1006, doi:10.1038/s41467-018-03455-x (2018).

1043 70 Stetson, D. B. & Medzhitov, R. Recognition of cytosolic DNA activates an IRF3-
1044 dependent innate immune response. *Immunity* **24**, 93-103,
1045 doi:10.1016/j.immuni.2005.12.003 (2006).

1046 71 Ishikawa, H., Ma, Z. & Barber, G. N. STING regulates intracellular DNA-
1047 mediated, type I interferon-dependent innate immunity. *Nature* **461**, 788-792,
1048 doi:10.1038/nature08476 (2009).

1049 72 Wang, Z., Zhu, L., Guo, T., Wang, Y. & Yang, J. Decreased H2B
1050 monoubiquitination and overexpression of ubiquitin-specific protease enzyme
1051 22 in malignant colon carcinoma. *Hum Pathol* **46**, 1006-1014,
1052 doi:10.1016/j.humpath.2015.04.001 (2015).

1053 73 Lee, J. S. *et al.* Histone crosstalk between H2B monoubiquitination and H3
1054 methylation mediated by COMPASS. *Cell* **131**, 1084-1096,
1055 doi:10.1016/j.cell.2007.09.046 (2007).

1056 74 Xiao, T. *et al.* Histone H2B ubiquitylation is associated with elongating RNA
1057 polymerase II. *Mol Cell Biol* **25**, 637-651, doi:10.1128/MCB.25.2.637-651.2005
1058 (2005).

1059 75 Minsky, N. *et al.* Monoubiquitinated H2B is associated with the transcribed
1060 region of highly expressed genes in human cells. *Nat Cell Biol* **10**, 483-488,
1061 doi:10.1038/ncb1712 (2008).

1062 76 Martin-Sancho, L. *et al.* Functional landscape of SARS-CoV-2 cellular
1063 restriction. *Mol Cell* **81**, 2656-2668 e2658, doi:10.1016/j.molcel.2021.04.008
1064 (2021).

1065 77 Fonseca, G. J. *et al.* Adenovirus evasion of interferon-mediated innate immunity
1066 by direct antagonism of a cellular histone posttranslational modification. *Cell*
1067 *Host Microbe* **11**, 597-606, doi:10.1016/j.chom.2012.05.005 (2012).

1068 78 Zhu, B. *et al.* Monoubiquitination of human histone H2B: the factors involved
1069 and their roles in HOX gene regulation. *Mol Cell* **20**, 601-611,
1070 doi:10.1016/j.molcel.2005.09.025 (2005).

1071 79 Zhang, S., Wang, J. & Cheng, G. Protease cleavage of RNF20 facilitates
1072 coronavirus replication via stabilization of SREBP1. *Proc Natl Acad Sci U S A*
1073 **118**, doi:10.1073/pnas.2107108118 (2021).

1074 80 Sui, H. *et al.* STING is an essential mediator of the Ku70-mediated production
1075 of IFN- λ 1 in response to exogenous DNA. *Science signaling* **10**,
1076 doi:10.1126/scisignal.aah5054 (2017).

1077 81 Chen, J. *et al.* STING-Dependent Interferon- λ 1 Induction in HT29 Cells, a
1078 Human Colorectal Cancer Cell Line, After Gamma-Radiation. *International*
1079 *journal of radiation oncology, biology, physics* **101**, 97-106,
1080 doi:10.1016/j.ijrobp.2018.01.091 (2018).

1081 82 Di Domizio, J. *et al.* The cGAS-STING pathway drives type I IFN
1082 immunopathology in COVID-19. *Nature*, doi:10.1038/s41586-022-04421-w
1083 (2022).

1084 83 Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using
1085 real-time quantitative PCR and the 2^{(-Delta Delta C(T))} Method. *Methods* **25**,
1086 402-408, doi:10.1006/meth.2001.1262 (2001).

1087 84 Carvalho, B. S. & Irizarry, R. A. A framework for oligonucleotide microarray
1088 preprocessing. *Bioinformatics (Oxford, England)* **26**, 2363-2367,
1089 doi:10.1093/bioinformatics/btq431 (2010).

1090 85 Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-
1091 sequencing and microarray studies. *Nucleic Acids Res* **43**, e47,
1092 doi:10.1093/nar/gkv007 (2015).

1093 86 Luo, W., Friedman, M. S., Shedden, K., Hankenson, K. D. & Woolf, P. J. GAGE:
1094 generally applicable gene set enrichment for pathway analysis. *BMC*
1095 *Bioinformatics* **10**, 161, doi:10.1186/1471-2105-10-161 (2009).

1096 87 Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based
1097 approach for interpreting genome-wide expression profiles. *Proceedings of the*
1098 *National Academy of Sciences* **102**, 15545-15550,
1099 doi:10.1073/pnas.0506580102 (2005).

1100 88 Hjerpe, R. *et al.* Efficient protection and isolation of ubiquitylated proteins using
1101 tandem ubiquitin-binding entities. *EMBO Rep* **10**, 1250-1258,
1102 doi:10.1038/embo.2009.192 (2009).

1103

A

B

C

A

B

C

D

Karlowitz et al Figure 2

A **B** **C** **D** **E**

Supplementary Information

USP22 controls type III interferon signaling and SARS-CoV-2 infection through activation of STING

Rebekka Karlowitz, Megan L. Stanifer, Jens Roedig, Geoffroy Andrieux, Denisa Bojkova, Sonja Smith, Lisa Kowald, Ralf Schubert, Melanie Boerries, Jindrich Cinatl Jr., Steeve Boulant, Sjoerd J. L. van Wijk[#]

[#] Corresponding author: Sjoerd J. L. van Wijk, Institute for Experimental Cancer Research in Pediatrics, Goethe University Frankfurt, Komturstrasse 3a, 60528 Frankfurt am Main, Germany, Phone: +49 69 67866574, Fax: +49 69 6786659158, Email: vanWijk@med.uni-frankfurt.de, s.wijk@kinderkrebsstiftung-frankfurt.de

Summary: Four Supplementary Figures including Supplementary Figure legends and a Supplementary Table

Supplemental Figure 1 (related to Figure 1). A. Scatter plot demonstrating the changes in gene expression of CRISPR/Cas9 control (NHT) HT-29 cells with two independent single-cell HT-29 USP22 KO clones (#16 and #62). Color code represents the average log2 foldchange. **B, C.** Western blot analysis of mono-ubiquitinated (H2Bub1) and total levels of Histone 2B (H2B) (**B**) and Histone 2A (H2A) (**C**) as well as USP22 in control and USP22 KO HT-29 cells (USP22 KO #62). Vinculin served as loading control. Representative blots of at least two different independent experiments are shown.

Supplemental Figure 2 (related to Figure 3). **A.** Western blot analysis of phosphorylated γ -H2AX (p. γ -H2AX) and USP22 expression levels in control (non-human target: NHT) and CRISPR/Cas9-generated USP22 knock-out (KO) HT-29 cells (USP22 KO) subjected to vehicle or etoposide (100 μ M) for 2 h. GAPDH served as loading control. Representative blots of at least two different independent experiments are shown. **B.** Western blot analysis of phosphorylated and total p65, p100/p52 and USP22 expression levels in control and USP22 KO HT-29 cells (USP22 KO #62). GAPDH served as loading control. Representative blots of at least two different independent experiments are shown. **C.** Basal mRNA expression levels of TMEM173/STING in control and USP22 KO HT-29 cells (USP22 KO #62) using qRT-PCR. Gene expression was normalized against 28S mRNA and is presented as x-fold mRNA expression compared to NHT. Mean and SD of three independent experiments in triplicate are shown. *P < 0.05.

A

Supplemental Figure 3 (related to Figure 5). A. mRNA expression levels of TMEM173/STING in control (non-human target: NHT) and CRISPR/Cas9-generated USP22 knock-out (KO) HT-29 cells (USP22 KO) using qRT-PCR. Cells were treated with ruxolitinib (5 μ M) for the indicated timepoints. Gene expression was normalized against 28S mRNA and is presented as x-fold mRNA expression compared to NHT. Mean and SD of three independent experiments in triplicate are shown. *** $P < 0.001$, n.s. not significant.

Supplemental Figure 4 (related to Figure 6).

A. Basal mRNA expression levels of IFNA, IFNB and IFNL1 in wild-type (WT), control (non-human target: NHT) and CRISPR/Cas9-generated USP22 knock-out (KO) Caco-2 single clones (USP22 KO #1 and #6). Gene expression was normalized against 28S mRNA and is presented as x-fold mRNA expression compared to NHT. Mean and SD of four (IFNA, IFNB) or three (IFNL1) independent experiments in triplicate are shown. *P < 0.05. **B.** TCID50/mL, determined via titration of supernatant from SARS-CoV-2-infected WT, control and USP22 KO Caco-2 cells (USP22 KO #1 and #6) 6 and 24 hpi on Vero cells. Mean and SD of three independent experiments in triplicate are shown.

C. Idem, 24 hpi, supernatant additionally of NHT-, USP22 KO #1- and #6-STING dKO
Caco-2 cells.

Supplementary Table 1: List of qRT-PCR primers used in this study

	<i>Forward primer</i>	<i>Reverse primer</i>
TGFB1	ACTACTACGCCAAGGAGGTCAC	TGCTTGAACCTTGTCATAGATTTTCG
SLFN5	AGCAAGCCTGTGTGCATTC	ACCACTCTGTCTGAAAATACTGGA
TGM2	GGCACCAAGTACCTGCTCA	AGAGGATGCAAAGAGGAACG
DDX60	AATCCCACAGGACTGCACA	TCGACCAAATACCTTCTGCAA
USP22	GAAGATCACCACGTATGTGTCC	CATTCATCCTGCTCTCTTTGC
AK4	CACTGGTGAACCGTTAGTCCA	AGCACTCCTCGGCTCTTGT
CXCR4	GGCCCTCAAGACCACAGTCA	TTAGCTGGAGTGAAAACCTGAAG
BST2	CCACCTGCAACCACACTG	CCTGAAGCTTATGGTTTAATGTAGTG
PARP9	CTGTCTGCACCGAGGAGAG	GCGCTTCAAAGCATAGACTGT
USP18	TCCCGACGTGGAACCTCAG	CAGGCACGATGGAATCTCTC
OAS3	TCCCATCAAAGTGATCAAGGT	ACGAGGTCGGCATCTGAG
IFIT1	CTTGTGGGTAATACAGTGGAGATG	GCTCCAGACTATCCTTGACCTG
IRF9	AGCCTGGACAGCAACTCAG	GAAACTGCCCACTCTCCACT
ISG15	GAGGCAGCGAACTCATCTTT	AGCATCTTCACCGTCAGGTC
OAS2	TGCAGGGAGTGGCCATAG	TCTGATCCTGGAATTGTTTTAAGTC
IFI27	GTGGCCAAAGTGGTCAGG	CCAATCACAACCTGTAGCAATCC
IFI6	AACCGTTTACTCGCTGCTGT	GGGCTCCGTCACCTAGACCT
panIFNA	TCCATGAGVTGATBCAGCAGA	ATTTCTGCTCTGACAACCTCCC
IFNB1	ATGACCAACAAGTGCTCCTCC	GGAATCCAAGCAAGTTGTAGCTC
IFNL1	GGACGCCTTGGAAGAGTCAC	AGCTGGGAGAGGATGTGGT
COV1	GCCTCTTCTGTTCTCATCAC	AGACAGCATCACCGCCATTG
TBP	CCACTCACAGACTCTCACAAC	CCACTCACAGACTCTCACAAC
DDX58	TGTGGGCAATGTCATCAAAA	GAAGCACTTGCTACCTCTTGC
TMEM173	ACATTCGCTTCCTGGATAAACT	CTGCTGTCTATCTGCAGGTTT
IFIH1	TTTTGCAGATTCTTCTGTAGTTTCA	TGCTGTTATGTCCAAGACTTTCA