Title: Viviparity is sufficient for the evolution of genomic imprinting

Authors: Sean A. Montgomery^{1,2}, Tetsuya Hisanaga¹, Nan Wang³, Elin Axelsson¹, Svetlana Akimcheva¹, Milos Sramek¹, Chang Liu³, Frédéric Berger^{1*}

Affiliations:

¹ Gregor Mendel Institute, Austrian Academy of Sciences, Vienna BioCenter; Dr. Bohr-Gasse 3, 1030 Vienna, Austria.

² Vienna BioCenter PhD Program, Doctoral School of the University of Vienna and Medical University of Vienna, A-1030, Vienna, Austria

³ Institute of Biology, University of Hohenheim, Garbenstrasse 30, 70599 Stuttgart, Germany

*Corresponding author. Email: frederic.berger@gmi.oeaw.ac.at

Abstract:

Genomic imprinting is an epigenetic phenomenon whereby one parental allele is preferentially expressed. Despite the assumption that extraembryonic annexes are necessary for the innovation of genomic imprinting, the evolutionary conditions under which embryonic genomic imprinting can evolve are unclear. We use the viviparous model *Marchantia polymorpha* to assess whether imprinting can occur in embryos that develop inside the mother but lack complex extraembryonic tissues. Contrasting with the limited number of imprinted loci in flowering plants and mammals, the paternal genome is globally repressed in *Marchantia* embryos. Silencing is mediated by Polycomb-deposited methylation of histone H3 at lysine 27 after gametogenesis but before karyogamy. Regulation of this new form of imprinting is essential for embryonic development. This report provides the proof of principle that imprinting can regulate embryogenesis in a larger range of viviparous organisms and extends our view on the diverse forms which imprinting can take.

Main text:

In multicellular eukaryotes, gene expression in early developmental stages needs to be tightly controlled. A specific epigenetic control of embryogenesis, defined as “genomic imprinting”, was identified in organisms with embryos dependent on maternal nutrition provided through placental or placenta-like tissues¹⁻⁴. Genomic imprinting results in the asymmetric, unequal expression of parental alleles. In both plants and mammals, a couple of hundred imprinted loci have been described that are either paternally or maternally expressed. A large majority of imprinted genes are controlled by DNA methylation over specific cis-control elements¹⁻⁴, although H3K27me3 has been shown to control a few loci in mouse^{5,6} and some paternally expressed loci in *Arabidopsis*^{7,8}. Another form of imprinting has been described in mammals. Females inactivate one of the X sex chromosomes to achieve a balanced dosage of gene expression. X chromosome inactivation (XCI) in marsupials⁹ and in extraembryonic lineages¹⁰ and early embryo^{11,12} in mouse is a complex mechanism that involves long non-coding RNAs, DNA methylation and the deposition of H3K9me3 and H3K27me3¹³.

Parental genomic imprinting has only been described in therian mammals and flowering plants, and many imprinted genes are primarily expressed in the nutritive extraembryonic tissues that define these species, namely the placenta and endosperm, respectively^{4,14-16}. Due to the large evolutionary distance between animals and plants, genomic imprinting is thought to have evolved independently, and its association with complex placental tissues has led to the assumption that the innovation of genomic imprinting is dependent on the presence of these tissues^{17,18}. The occurrence of genomic imprinting in mammals and plants is also tightly linked with viviparity, the development of embryos within mothers. Thus, it remains unknown whether placentation or viviparity is sufficient for the evolution of genomic imprinting. Imprinting has not been found in viviparous animals without complex placentas, but only a limited number of loci were examined in these

studies^{18–20}. One of the two major land plant lineages, bryophytes, also evolved viviparity without extraembryonic tissues comparable with the endosperm of flowering plants. While it has been theorized that bryophytes may have imprinting, these predictions have not been tested experimentally^{21–25}.

To determine whether genomic imprinting occurs in species with viviparous embryos but lacking placental tissues, we used the model bryophyte *Marchantia polymorpha* (hereafter referred to as *Marchantia*). *Marchantia* is suitable to test these predictions because haploid vegetative individuals are either male or female (Fig. 1a) and the genome has been fully sequenced^{26,27}, thus allowing for a genome-wide investigation into imprinting.

Embryonic transcription is maternally biased

To assess whether imprinting may occur in *Marchantia*, we first looked for evidence of allele-specific expression in diploid embryos. We performed crosses between two wild-type natural accessions, Cam-2 as the mother and Tak-1 as the father, and obtained transcriptomes from embryos thirteen days after fertilization (daf). Each transcriptome was prepared from single hand-dissected embryos that were washed several times to remove potential contaminating RNA from the surrounding maternal tissue (Extended Data Fig. 1, Supplementary Video 1)²⁸. We utilized single-nucleotide polymorphisms (SNPs) between male and female accessions to calculate the ratio of reads originating from maternal alleles versus paternal alleles (maternal ratio, p_m ; ascribed a value between 0 and 1, ranging from 0 if only paternal reads were detected to 1 if only maternal reads were detected). We only considered genes with at least fifty reads containing informative SNPs. Transcription was overall maternally biased for 98% of resolved genes, with 73% of genes maternally biased (as defined in²⁹; $0.65 \leq p_m < 0.95$) and 25% of genes only expressed from maternal alleles ($p_m \geq 0.95$) (Fig. 1b, c). The exact

reciprocal cross was not possible because inbred genetically near-identical pairs of males and females do not exist, but to confirm that the maternal bias did not result from the pair of natural accessions used, we analyzed published RNA-seq data from a cross of different accessions, Tak-2 and Tak-1³⁰. The published transcriptomes were generated from samples collected by laser-capture microdissection, an orthogonal sample collection method that offered equally high sample purity²⁸. A similarly strong maternal bias in transcription was observed, with 99% of genes maternally biased or expressed only from maternal alleles (Extended Data Fig. 2a, b). Thus, these data ruled out that the observed allele-specific gene expression originated from natural variation amongst wild-type parents or from maternal contamination during sample collection. This was also evident from the comparison amongst transcriptomes of isolated embryos, vegetative tissue, and sexual organs which revealed the distinctness of isolated embryos from other tissues (Fig. 1d). Additionally, we tested whether the genes resolved by our analyses formed a sample representative of all genes. We found no correlation between the maternal ratio and expression level of a gene (Extended Data Fig. 2c), nor did the transcription maternal ratio vary significantly along the length of each autosome (Extended Data Fig. 2d, e). Thus, we infer that the genes we were able to resolve with SNPs were representative of a genome-wide maternal bias in transcription. We conclude that in *Marchantia* embryos, genes are primarily or exclusively expressed from their maternal allele. The lack of paternal allele expression suggests the presence of a repressive chromatin modification specifically on the paternal genome.

H3K27me3 is paternally biased

To better understand what chromatin-related mechanisms may be driving the maternal bias in embryonic transcription, we examined differentially expressed genes between vegetative parents and embryos. In total, 3879 genes were upregulated in embryos relative to both

mothers and fathers (Extended Data Fig. 3a), while 3466 genes were downregulated (Extended Data Fig. 3b). Upregulated genes were more expressed from the maternal genome than downregulated genes (Extended Data Fig. 3c, effect size (Cohen's d) = 0.276) highlighting a maternal control over the embryonic transcriptome. Since imprinting is an epigenetic process, we focused further on chromatin-related genes. Of the 215 chromatin-related genes in the *Marchantia* genome²⁷, 151 were upregulated and 7 downregulated (Extended Data Fig. 3d, Supplementary Table 1). Of these, 20 genes were specifically expressed in the embryonic stage (Supplementary Table 1). Two noteworthy genes were paralogs of the catalytic subunit of the Polycomb Repressive Complex 2 (PRC2), *E(z)2* and *E(z)3* (Extended Data Fig. 3e). PRC2 is a conserved multi-subunit complex that deposits H3K27me3 and is associated with gene silencing³¹. The other three subunits of PRC2, *FIE*, *Su(z)12*, and *MSI1*, were expressed in all tissues (Extended Data Fig. 3e). Therefore, we hypothesized that H3K27me3 might be present on silenced paternal alleles in the *Marchantia* embryo.

We first set out to determine whether H3K27me3 was enriched on paternal alleles in *Marchantia* embryos. We profiled chromatin modifications using CUT&RUN^{32,33} on sorted nuclei from *Marchantia* embryos. We used SNPs between male and female accessions to distinguish the parental allele of origin for CUT&RUN reads to calculate a maternal ratio (p_m) for genomic regions of interest. Enrichment in H3K27me3 was paternally biased for 88% of genes resolved ($0.05 < p_m \leq 0.35$) (Fig. 2a, b) and genes with paternally biased H3K27me3 had maternally biased transcription (Fig. 2c). Genes paternally marked with H3K27me3 were located across all autosomes (Extended Data Fig. 4a, b), indicating the broad, pervasive nature of the phenomenon. In contrast, a paternal bias was not observed in profiles of H3K9me1, H3, and H3K36me3 (5%, 2%, and 2% of genes with $0.05 < p_m \leq 0.35$, respectively) (Extended Data Fig. 4c-e). We conclude that levels of H3K27me3 enrichment

anticorrelate with maternally biased transcription and spreads over most paternal alleles in *Marchantia* embryos. These findings suggested that H3K27me3 covers the entire genome of paternal origin.

The paternal genome is partitioned into dense H3K27me3 compartments

To test if the paternal genome was coated with H3K27me3, we performed immunofluorescence experiments to observe the localization of this modification within embryonic nuclei. As a control, nuclei of parental adult vegetative cells showed evenly distributed speckles of heterochromatin marked by H3K27me3 (Fig. 3a)²⁶. In stark contrast, one to three large heterochromatic compartments, as defined by dense DNA staining, covered 10% of the area in embryonic nuclei (Fig. 3a, Extended Data Fig. 5a, b). A strong correlation between heterochromatic foci and H3K27me3 was apparent (Fig. 3a). 44% of the H3K27me3 signal was contained within these compartments (Extended Data Fig. 5c), whereas 70% of the area of the compartments was contained within H3K27me3 domains (Extended Data Fig. 5d). In contrast, only 20% and 10% of H3K9me1 and H3K36me3 signal, respectively, were contained within heterochromatic compartments (Extended Data Fig. 5c). We conclude that the portion of the genome marked by H3K27me3 represents the largest fraction of heterochromatin organized as a couple of dense compartments in nuclei of the embryo.

All paternal autosomes are coated with H3K27me3

The *Marchantia* life cycle has a vegetative haploid phase with a diploid phase of embryogenesis (Fig. 1a). In *Marchantia* diploid embryonic cells, the genome is packaged in $2n = 18$ chromosomes including two sex chromosomes and sixteen autosomes. The large size of the heterochromatic compartments marked by H3K27me3 suggested that they contained

entire chromosomes. Accordingly, immunofluorescence of mitotic cells revealed that eight of the sixteen autosomes were densely coated with H3K27me₃, whereas the other half had no detectable H3K27me₃ (Fig. 3b). In contrast, mitotic vegetative haploid cells showed an uneven speckled pattern of H3K27me₃ over the eight autosomes (Fig. 3b). This observation mirrored the strong paternal bias of H3K27me₃ enrichment, and we conclude that the paternal genome is covered by H3K27me₃ and partitioned into heterochromatic compartments within embryonic nuclei.

In addition to the eight autosomes, each parent carries a small sex chromosome (U in females and V in males³⁴). Sex chromosomes detected using FISH were not associated with H3K27me₃ heterochromatic foci in *Marchantia* embryos (Fig. 3c, Extended Data Fig. 5e). Instead, we observed that both U and V sex chromosomes rather associated with H3K9me₁ heterochromatic foci (Fig. 3d, Extended Data Fig. 5e). We conclude that the sex chromosomes are excluded from H3K27me₃ heterochromatic compartments and form small constitutive heterochromatic foci in both embryonic and vegetative nuclei.

H3K27me₃ is deposited in paternal pronuclei

Like in most animals, in the *Marchantia* zygote the male pronucleus contributed by the sperm remains separated from the female pronucleus contributed by the egg, thus providing an opportunity for the deposition of an epigenetic mark on the genome of one parent^{35,36}. As *Marchantia* sperm chromatin is comprised of protamines and is devoid of histones³⁷, we hypothesized that paternal H3K27me₃ is deposited on paternal alleles sometime after fertilization. Male and female pronuclei remain separate until four daf (Fig. 4a, b)³⁵, thus we examined if H3K27me₃ was deposited at 3 daf, before pronuclear fusion. As we could not isolate pronuclei for chromatin profiling, we instead performed immunofluorescence

experiments. At 3 daf, we observed both H3K27me3 and H3 in the paternal pronucleus (Fig. 4c, Extended Data Fig. 6), demonstrating that H3K27me3 is deposited in the paternal pronucleus before its fusion with the maternal pronucleus. Therefore, paternal alleles become imprinted by H3K27me3 while they are spatially segregated from maternal alleles prior to the fusion of pronuclei. The conservative restoration of H3K27me3 after DNA replication³⁸ provides a mechanism to propagate the initial paternal “coat” of H3K27me3 to all autosomes and silence the paternal genome throughout embryonic development.

PRC2 deposits H3K27me3 and silences the paternal genome

To directly test the effect of embryo-specific PRC2 subunits on paternal H3K27me3 imprinting, we knocked-out both embryo-specific paralogs of the catalytic subunit, *E(z)2* and *E(z)3* (Extended Data Fig. 7a). These mutants did not display any aberrant phenotype prior to fertilization (Extended Data Fig. 7b, c). We crossed mutant females (Cam-2 *e(z)2/e(z)3*) to wild type males and observed a loss or dispersion of large heterochromatic foci that overlapped with H3K27me3 foci in embryos compared to wild-type embryos (Fig. 5a, Extended Data Fig. 8a). In these embryos, enrichment of H3K27me3 as measured by CUT&RUN was significantly decreased over most genes (Extended Data Fig. 8b, one sample t-test $p < 0.0001$). Thus, maternal inheritance of both *e(z)2* and *e(z)3* altered patterns of H3K27me3 associated heterochromatic foci and reduced H3K27me3 deposition. The paralog *E(z)1* is expressed in embryos (Extended Data Fig. 3e) and likely accounted for the remaining detected H3K27me3, but it proved impossible to test this hypothesis due to lethality at the haploid vegetative stage in knockdowns of *E(z)1*³⁹. To determine whether paternal alleles were the source of H3K27me3 loss, we distinguished the parental genome of origin of CUT&RUN sequencing reads and calculated maternal ratios. The paternally biased enrichment of H3K27me3 was significantly reduced (Fig. 5b, one sample t-test $p < 0.0001$,

effect size = 1.03) and only 51% of genes were categorized as paternally biased ($0.05 < p_m \leq 0.35$) (Extended Data Fig. 7c), down from 88% in wild type (Fig. 2a). Furthermore, there was a negative correlation between H3K27me3 enrichment and maternal ratio (Fig. 5c), indicating that loci that lost H3K27me3 had predominantly lost paternal H3K27me3 in embryos that lost the maternal alleles of *E(z)2* and *E(z)3*. We conclude that the deposition of H3K27me3 on most paternal loci in *Marchantia* embryos depends on maternally supplied PRC2 activity.

If H3K27me3 did indeed repress paternal alleles, we expected expression from paternal alleles at loci that lost paternal H3K27me3 in *e(z)2/e(z)3* mutants. To test this idea, we generated transcriptomes from mutant embryos and examined maternal ratios of gene expression. Overall, transcription became more biallelic in mutants than in wild-type embryos (Fig. 5b, one sample t-test $p < 0.0001$, effect size = -1.03). Only 47% of genes were maternally biased ($0.65 \leq p_m < 0.95$) and 15% of genes completely expressed from maternal alleles ($p_m \geq 0.95$), a stark deviation from wild type values of 73% and 25% (Extended Data Fig. 9a and compare with Fig. 1b). Comparing both mutant maternal ratios of transcription with patterns of H3K27me3, we observed that the H3K27me3 maternal ratio negatively correlated with the maternal ratio of transcription (Extended Data Fig. 9b, c) and that H3K27me3 enrichment positively correlated with the maternal ratio of transcription (Extended Data Fig. 9b, d), indicating that loci with less paternal H3K27me3 in mutants were more transcribed from paternal alleles. High paternal H3K27me3 and maternal transcription did not correlate with the level of gene expression (Extended Data Fig. 9b, e-g), suggesting that absolute gene expression levels did not influence paternal allele repression. Therefore, paternal alleles regain expression in the absence of maternal PRC2 and upon the loss of paternal H3K27me3. We conclude that H3K27me3 deposited by embryo-specific PRC2 subunits is required for the collective repression of paternal alleles.

Considering that in the land plant lineage, the first diploid embryos evolved in bryophytes, the maintenance of a virtual haploid state by imprinting of paternal autosomes is surprising. To assess the physiological relevance of the loss of paternal allele repression, we quantified the growth and survival of mutant embryos. Embryonic growth was significantly slower in mutants than in wild type, as measured by total size (Extended Data Fig. 10a). Only 20% of mutant embryos survived to maturity versus 95% of wild type (Extended Data Fig. 10), and only 5% of all mutant embryos produced spore-bearing structures, compared to 77% of wild type (Extended Data Fig. 10c). Of those mutants that produced spores, none produced viable spores, thus rendering all mutants unable to continue their life cycle (Fig. 5d). We conclude that our results are consistent with the model that PRC2-mediated paternal allele repression is essential for viability and fecundity in *Marchantia* (Fig. 6).

Discussion

In conclusion, we report genomic imprinting in the embryos of the bryophyte *Marchantia polymorpha*. The repressive mark H3K27me3 is first deposited over the whole paternal pronucleus, in contrast to mammals where H3K27me3-mediated imprinting impacts only a handful of loci and is deposited in the female gamete⁴⁰. The formation of heterochromatic foci associated with H3K27me3 is reminiscent of the compaction and compartmentalization of the X chromosome during XCI in mammals^{41–43}. However, since half of the genome is marked exclusively by H3K27me3, its partition results in large compartments, the compaction of which depends on PRC2 activity. Interference of maternal PRC2-mediated H3K27me3 deposition ultimately halts embryonic development. Thus, PRC2 initiates and maintains silencing of the entire paternal genome, which is essential for the development of the diploid embryo of *Marchantia*.

We believe that genomic imprinting in *Marchantia* represents a new form of imprinting that we name “imprinted autosomal inactivation”. It is distinct from parental genomic imprinting described in flowering plants and therian mammals for three main reasons: the epigenetic mark is deposited after fertilization; the entire genome of one parent is silenced by the epigenetic mark; and imprinting imposes a global maternal control of embryogenesis. The outcome of imprinting in *Marchantia* also differs from the elimination of the paternal genome following heterochromatin formation in several insect species^{44,45}. In insects, heterochromatinized paternal chromosomes are eliminated by still unknown mechanisms, though the timing of elimination varies amongst species^{46,47}. In contrast, the paternal genome in *Marchantia* embryos subjected to “imprinted autosomal inactivation” is still passed on to the next generation. Additionally, imprinting in *Marchantia* differs from XCI in mammals because the autosomes and not the sex chromosomes are imprinted and H3K9me is not involved⁴⁸. Similar mechanisms may be at play in mediating whole-chromosome compaction and repression in mammals, insects and *Marchantia*. The molecular mechanisms underlying the establishment and abolition of “imprinted autosomal inactivation” are not addressed by our model (Fig. 6), however their elucidation will be of interest to make cross-kingdom comparisons with other instances of genomic imprinting.

As early land plants evolved with separate sexes³⁴, it is likely that “imprinted autosomal inactivation” first evolved in an ancestor with a life cycle like that of *Marchantia* earlier than 440 million years ago. The presence of genomic imprinting in bryophytes supports that the primary causes for the evolution of imprinting are viviparity and its collateral maternal support of embryo development. Imprinting has not been discovered in non-therian viviparous animals, but only orthologous loci imprinted in mammals were investigated, thus a genome-wide search may yield new insights^{18–20}. Our findings support the idea that viviparity is sufficient for the evolution of imprinting and provide a new

framework to explore the evolution of imprinting in a much more diverse range of organisms than previously considered across eukaryotes.

Main references:

1. McGrath, J. & Solter, D. Maternal Thp lethality in the mouse is a nuclear, not cytoplasmic, defect. *Nature* **308**, 550–551 (1984).
2. Barton, S. C., Surani, M. A. & Norris, M. L. Role of paternal and maternal genomes in mouse development. *Nature* **311**, 374–6
3. Feil, R. & Berger, F. Convergent evolution of genomic imprinting in plants and mammals. *Trends Genet.* **23**, 192–199 (2007).
4. Pires, N. D. & Grossniklaus, U. Different yet similar: evolution of imprinting in flowering plants and mammals. *F1000Prime Rep.* **6**, 63 (2014).
5. Mei, H. *et al.* H2AK119ub1 guides maternal inheritance and zygotic deposition of H3K27me3 in mouse embryos. *Nat. Genet.* **53**, 539–550 (2021).
6. Inoue, A., Chen, Z., Yin, Q. & Zhang, Y. Maternal Eed knockout causes loss of H3K27me3 imprinting and random X inactivation in the extraembryonic cells. *Genes Dev* **32**, 1525–1536 (2018).
7. Borg, M. *et al.* Targeted reprogramming of H3K27me3 resets epigenetic memory in plant paternal chromatin. *Nat. Cell Biol.* **22**, 621–629 (2020).
8. Moreno-Romero, J., Del Toro-De León, G., Yadav, V. K., Santos-González, J. & Köhler, C. Epigenetic signatures associated with imprinted paternally expressed genes in the Arabidopsis endosperm. *Genome Biol* **20**, 41 (2019).
9. Sharman, G. B. Late DNA Replication in the Paternally Derived X Chromosome of Female Kangaroos. *Nature* **230**, 231–232 (1971).
10. Takagi, N. & Sasaki, M. Preferential inactivation of the paternally derived X chromosome in the extraembryonic membranes of the mouse. *Nature* **256**, 640–642 (1975).
11. Mak, W. *et al.* Reactivation of the Paternal X Chromosome in Early Mouse Embryos.

- Science* **303**, 666–669 (2004).
12. Okamoto, I., Otte, A. P., Allis, C. D., Reinberg, D. & Heard, E. Epigenetic dynamics of imprinted X inactivation during early mouse development. *Science* **303**, 644–649 (2004).
 13. Żylicz, J. J. & Heard, E. Molecular Mechanisms of Facultative Heterochromatin Formation: An X-Chromosome Perspective. *Annu. Rev. Biochem.* **89**, 255–282 (2020).
 14. Barlow, D. P. & Bartolomei, M. S. Genomic Imprinting in Mammals. *Cold Spring Harb. Perspect. Biol.* **6**, a018382–a018382 (2014).
 15. Tucci, V., Isles, A. R., Kelsey, G., Ferguson-Smith, A. C. & Erice Imprinting Group. Genomic Imprinting and Physiological Processes in Mammals. *Cell* **176**, 952–965 (2019).
 16. Gehring, M. & Satyaki, P. R. Endosperm and Imprinting, Inextricably Linked. *Plant Physiol.* **173**, 143–154 (2017).
 17. Patten, M. M., Cowley, M., Oakey, R. J. & Feil, R. Regulatory links between imprinted genes: evolutionary predictions and consequences. *Proceedings. Biol. Sci.* **283**, (2016).
 18. Renfree, M. B., Suzuki, S. & Kaneko-Ishino, T. The origin and evolution of genomic imprinting and viviparity in mammals. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **368**, 20120151 (2013).
 19. Griffith, O. W., Brandley, M. C., Belov, K. & Thompson, M. B. Allelic expression of mammalian imprinted genes in a matrotrophic lizard, *Pseudemoia entrecasteauxii*. *Dev. Genes Evol.* **226**, 79–85 (2016).
 20. Lawton, B. R. *et al.* Allelic expression of IGF2 in live-bearing, matrotrophic fishes. *Dev. Genes Evol.* **215**, 207–212 (2005).
 21. Haig, D. Filial mistletoes: the functional morphology of moss sporophytes. *Ann Bot*

- 111, 337–345 (2013).
22. Haig, D. & Wilczek, A. Sexual conflict and the alternation of haploid and diploid generations. *Philos Trans R Soc L. B Biol Sci* **361**, 335–343 (2006).
23. Montgomery, S. A. & Berger, F. The evolution of imprinting in plants: beyond the seed. *Plant Reprod.* (2021). doi:10.1007/s00497-021-00410-7
24. Shaw, A. J., Szovenyi, P. & Shaw, B. Bryophyte diversity and evolution: windows into the early evolution of land plants. *Am J Bot* **98**, 352–369 (2011).
25. Carey, S. B., Kollar, L. M. & McDaniel, S. F. Does degeneration or genetic conflict shape gene content on UV sex chromosomes? *Bryophyt. Divers. Evol.* **43**, 133–149 (2021).
26. Montgomery, S. A. *et al.* Chromatin Organization in Early Land Plants Reveals an Ancestral Association between H3K27me3, Transposons, and Constitutive Heterochromatin. *Curr. Biol.* **30**, 573–588.e7 (2020).
27. Bowman, J. L. *et al.* Insights into Land Plant Evolution Garnered from the *Marchantia polymorpha* Genome. *Cell* **171**, 287–304 e15 (2017).
28. Schon, M. A. & Nodine, M. D. Widespread Contamination of Arabidopsis Embryo and Endosperm Transcriptome Data Sets. *Plant Cell* **29**, 608–617 (2017).
29. Wang, X. & Clark, A. G. Using next-generation RNA sequencing to identify imprinted genes. *Heredity (Edinb)*. **113**, 156–166 (2014).
30. Frank, M. H. & Scanlon, M. J. Transcriptomic evidence for the evolution of shoot meristem function in sporophyte-dominant land plants through concerted selection of ancestral gametophytic and sporophytic genetic programs. *Mol. Biol. Evol.* **32**, 355–367 (2015).
31. Margueron, R. & Reinberg, D. The Polycomb complex PRC2 and its mark in life. *Nature* **469**, 343–349 (2011).

32. Skene, P. J. & Henikoff, S. An efficient targeted nuclease strategy for high-resolution mapping of DNA binding sites. (2016).
33. Zheng, X. & Gehring, M. Low-input chromatin profiling in Arabidopsis endosperm using CUT&RUN. *Plant Reprod.* **32**, 63–75 (2019).
34. Iwasaki, M. *et al.* Identification of the sex-determining factor in the liverwort *Marchantia polymorpha* reveals unique evolution of sex chromosomes in a haploid system. *Curr. Biol.* (2021). doi:10.1016/j.cub.2021.10.023
35. Hisanaga, T. *et al.* Deep evolutionary origin of gamete-directed zygote activation by KNOX/BELL transcription factors in green plants. *Elife* **10**, (2021).
36. Hisanaga, T. *et al.* Building new insights in plant gametogenesis from an evolutionary perspective. *Nat. plants* **5**, 663–669 (2019).
37. Reynolds, W. F. & Wolfe, S. L. Changes in basic proteins during sperm maturation in a plant, *Marchantia polymorpha*. *Exp. Cell Res.* **116**, 269–273 (1978).
38. Jiang, D. & Berger, F. DNA replication–coupled histone modification maintains Polycomb gene silencing in plants. *Science* **357**, 1146–1149 (2017).
39. Flores-Sandoval, E., Dierschke, T., Fisher, T. J. & Bowman, J. L. Efficient and inducible use of artificial MicroRNAs in *Marchantia polymorpha*. *Plant Cell Physiol.* **57**, 281–290 (2016).
40. Inoue, A., Jiang, L., Lu, F., Suzuki, T. & Zhang, Y. Maternal H3K27me3 controls DNA methylation-independent imprinting. *Nature* **547**, 419–424 (2017).
41. Galupa, R. & Heard, E. X-Chromosome Inactivation: A Crossroads Between Chromosome Architecture and Gene Regulation. *Annu. Rev. Genet.* **52**, 535–566 (2018).
42. Nozawa, R.-S. *et al.* Human inactive X chromosome is compacted through a PRC2-independent SMCHD1-HBiX1 pathway. *Nat. Struct. Mol. Biol.* **20**, 566–573 (2013).

43. Plath, K. *et al.* Role of Histone H3 Lysine 27 Methylation in X Inactivation. *Science* **300**, 131–135 (2003).
44. de la Filia, A. G. *et al.* Males That Silence Their Father’s Genes: Genomic Imprinting of a Complete Haploid Genome. *Mol. Biol. Evol.* **38**, 2566–2581 (2021).
45. Crouse, H. V. The Controlling Element in Sex Chromosome Behavior in *Sciara*. *Genetics* **45**, 1429–1443 (1960).
46. Bain, S. A. *et al.* Sex-specific expression and DNA methylation in a species with extreme sexual dimorphism and paternal genome elimination. *Mol. Ecol.* **30**, 5687–5703 (2021).
47. Morse, G. E. & Normark, B. B. A molecular phylogenetic study of armoured scale insects (Hemiptera: Diaspididae). *Syst. Entomol.* **31**, 338–349 (2005).
48. Heard, E. *et al.* Methylation of Histone H3 at Lys-9 Is an Early Mark on the X Chromosome during X Inactivation. *Cell* **107**, 727–738 (2001).
49. Cohen, J. Statistical Power Analysis. *Curr. Dir. Psychol. Sci.* **1**, 98–101 (1992).
50. Sugano, S. S. *et al.* Efficient CRISPR/Cas9-based genome editing and its application to conditional genetic analysis in *Marchantia polymorpha*. *PLoS One* **13**, e0205117 (2018).
51. Tsuboyama, S., Nonaka, S., Ezura, H. & Kodama, Y. Improved G-AgarTrap: A highly efficient transformation method for intact gemmalings of the liverwort *Marchantia polymorpha*. *Sci. Rep.* **8**, 10800 (2018).
52. Kao, P. & Nodine, M. D. Profiling Transcriptomes of Manually Dissected *Arabidopsis* Embryos. in 113–126 (2020). doi:10.1007/978-1-0716-0342-0_9
53. Picelli, S. *et al.* Full-length RNA-seq from single cells using Smart-seq2. *Nat. Protoc.* **9**, 171–181 (2014).
54. Higo, A. *et al.* Transcriptional Framework of Male Gametogenesis in the Liverwort

- Marchantia polymorpha L. *Plant Cell Physiol.* **57**, 325–338 (2016).
55. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).
 56. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841–842 (2010).
 57. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).
 58. Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).
 59. R Core Team. R: A Language and Environment for Statistical Computing. (2018).
 60. Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences. *F1000Research* **4**, 1521 (2015).
 61. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).
 62. Torchiano, M. effsize: Efficient Effect Size Computation. (2020).
doi:10.5281/zenodo.1480624
 63. Kolde, R. pheatmap: Pretty Heatmaps. (2019).
 64. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).
 65. McKenna, A. *et al.* The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).
 66. Krueger, F. & Andrews, S. R. SNPsplite: Allele-specific splitting of alignments between genomes with known SNP genotypes. *F1000Research* **5**, 1479 (2016).
 67. Borg, M., Buendía, D. & Berger, F. A simple and robust protocol for immunostaining

- Arabidopsis pollen nuclei. *Plant Reprod.* **32**, 39–43 (2019).
68. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682 (2012).
 69. Wang, N. & Liu, C. Study of Cell-Type-Specific Chromatin Organization: In Situ Hi-C Library Preparation for Low-Input Plant Materials. in 115–127 (2020).
doi:10.1007/978-1-0716-0179-2_9
 70. Bi, X. *et al.* Nonrandom domain organization of the Arabidopsis genome at the nuclear periphery. *Genome Res.* **27**, 1162–1173 (2017).
 71. Murray, M. G. & Thompson, W. F. Rapid isolation of high molecular weight plant DNA. *Nucleic Acids Res.* **8**, 4321–4326 (1980).
 72. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag New York, 2016).
 73. Kassambara, A. ggpubr: ‘ggplot2’ Based Publication Ready Plots. (2020).

Figure legends:

Figure 1 | Maternally biased transcription in embryos. **a**, Life cycle of *Marchantia polymorpha*. Haploid (1n) vegetative males and females produce male and female reproductive structures, which subsequently produce sperm and egg. The diploid (2n) embryo persists for around 20 days before meiosis and the production of haploid spores. Ploidy of each stage is indicated. **b**, Percentage of measured genes within each category of maternal ratio (p_m) of transcription in wild-type embryos. Segments are for paternal ($p_m < 0.05$), paternal bias ($0.05 < p_m \leq 0.35$), no bias ($0.35 < p_m < 0.65$), maternal bias ($0.65 \leq p_m < 0.95$), and maternal ($0.95 \leq p_m$) expression of genes, with the number of genes indicated above each bar. **c**, Histogram of the maternal ratio (p_m) of transcription per gene in wild-type (Cam-2 x Tak-1) embryos. Each bin is 0.01 units wide. **d**, Principal component analysis of transcriptomes from wild-type embryos (Cam-2 x Tak-1), vegetative tissues from female (Cam-2) and male (Tak-1) parents, and female and male sexual organs. The first two principal components are plotted, and the percentage of variance explained is indicated.

Figure 2 | Paternally biased H3K27me3 in embryos. **a**, Percentage of measured genes within each category of maternal ratio (p_m) of H3K27me3 in wild-type embryos. Segments are for paternal ($p_m < 0.05$), paternal bias ($0.05 < p_m \leq 0.35$), no bias ($0.35 < p_m < 0.65$), maternal bias ($0.65 \leq p_m < 0.95$), and maternal ($0.95 \leq p_m$) H3K27me3 of genes, with the number of genes indicated above each bar. **b**, Histogram of the maternal ratio (p_m) of H3K27me3 per gene in wild-type (Cam-2 x Tak-1) embryos. Each bin is 0.01 units wide. **c**, Scatterplot of maternal ratios of H3K27me3 and transcription per resolved gene.

Figure 3 | H3K27me3 heterochromatic foci in embryos. **a**, Immunofluorescence of H3K27me3 in interphase wild-type embryonic and vegetative nuclei. DNA is stained with DAPI. **b**, Immunofluorescence of H3K27me3 in mitotic wild-type embryonic and vegetative cells. DNA is stained with DAPI. Contrast was enhanced for the DAPI channel of vegetative nuclei for illustration purposes. Outlines of the H3K27me3-coated chromosomes are indicated with dashed yellow lines. **c**, Immuno-FISH for sex chromosomes and H3K27me3 in interphase wild-type embryonic nuclei. **d**, Immuno-FISH for sex chromosomes and H3K9me1 in interphase wild-type embryonic nuclei. The female sex chromosome is chrU and the male sex chromosome is chrV. Scale bars are as indicated for all images.

Figure 4 | H3K27me3 is deposited in paternal pronuclei. **a**, Annotated confocal image of a *Marchantia* zygote 3 days after fertilization (daf) and surrounding vegetative mother tissue. The paternal pronucleus is visible in the vicinity of the maternal pronucleus. Nuclei are stained with DAPI. Indicated are the fertilized zygotic cell (dashed yellow circle), maternal pronucleus (pink circle), vegetative mother tissue (green lines) surrounding the zygote, and paternal pronucleus (cyan circle). Scale bar as indicated. **b**, Composite maximum intensity projection confocal image of a *Marchantia* zygote expressing *SUN-GFP* at 3 daf plus surrounding vegetative mother tissue. Nuclear membranes are marked by localization of

SUN-GFP, shown in green. The paternal pronucleus is smaller than and adjacent to the maternal pronucleus. Autofluorescence from chloroplasts in vegetative mother cells is shown in red, and both channels are overlayed on a transmitted light image. Scale bar as indicated. **c**, Immunofluorescence image 3 daf of a *Marchantia* zygote. Both maternal and paternal pronuclei are indicated in pink and cyan, respectively. The inset depicts a zoomed in view of the paternal pronucleus with separate images for H3K27me3 (red), H3 (green), DAPI (blue), and the merged image. Contrast is enhanced for each image and channel independently for visualization purposes. Scale bars as indicated.

Figure 5 | Embryonic PRC2 represses the paternal genome. **a**, Immunofluorescence of H3K27me3 in interphase wild-type and mutant embryonic nuclei. DNA is stained with DAPI. **b**, Violin plots of maternal ratios for wild-type (WT) and mutant H3K27me3 and transcription. Cohen's *d* effect size values are indicated for pairwise comparisons of mutant to WT H3K27me3 maternal ratio and mutant to WT transcription maternal ratio, where $|d| > 0.8$ is a large effect, as previously reported⁴⁹. **c**, Scatterplot of H3K27me3 enrichment versus H3K27me3 maternal ratio per gene in WT and mutant embryos. Genes with an arcsinh-transformed H3K27me3 enrichment greater than 2 are displayed as triangles at the upper boundary of the plot. **d**, Spore germination assay for spores resulting from WT and mutant embryos. A serial dilution of a suspension of spores from a single embryo is shown.

Figure 6 | Model of genomic imprinting in *Marchantia*. Model of H3K27me3 deposition in WT paternal pronuclei and subsequent propagation throughout embryogenesis. Closed lollipops depict H3K27me3 on genes. Pink and blue circles depict maternal and paternal (pro)nuclei, respectively. Pink and striped ovals depict whole embryos and the parental genome from which transcription is occurring. Yellow circles depict mature spores. The lack of H3K27me3 on paternal pronuclei in mutant zygotes allows for the transcription of paternal alleles in the embryo, ultimately leading to the lack of viable spore production.

Methods:

Plant lines and growth conditions

Wild-type male Tak-1, female Cam-2, and female Tak-2 accessions of *Marchantia polymorpha* ssp. *ruderalis* were used in this study. Cam-2 and Tak-2 *e(z)2/e(z)3* mutants were generated as described below.

Female plants for crosses were grown at room temperature on Grodan Vital (Grodan) supplemented with liquid Hyponex fertilizer (Hyponex) under constant white light. Male plants for crosses were grown at 22 °C on Neuhaus N3 substrate soil (Humko) under 16 hours of far-red light and 80% humidity. Plants grown for the collection or observation of vegetative tissues were grown under axenic conditions on half-strength Gamborg B5 media without vitamins (Duchefa Biochemie) and 1% (w/v) agar under constant white light.

Crosses were performed by collecting mature antheridiophore discs in water and pipetting the water containing released sperm onto archegoniophores.

Generation of *e(z)2/e(z)3* mutants

To construct a plasmid to disrupt *E(z)2* and *E(z)3* simultaneously, two oligonucleotide pairs (TH219: ctcgAAATAGAAAGTGGCGCCT/TH220: aaacAGGCGCCACTTTCTATTT for *E(z)2*; TH223: ctcgATCATATACCCTCGGCTC /TH224: aaacGAGCCGAGGGTATATGAT for *E(z)3*) were annealed and cloned into the BsaI sites of pMpGE_En04 and pBC-GE14 to yield pMpGE_En04-MpEz2-sg1 and pBC-GE14-MpEz3-sg1, respectively. These two plasmids were assembled via BglII restriction sites and ligated to yield pMpGE_En04-MpEz2-sg1-MpEz3-sg1. The resulting DNA fragment containing two MpU6promoter-gRNA cassettes was transferred into pMpGE010 (cat. no. 71536, Addgene)⁵⁰ using the Gateway LR reaction (Thermo Fisher Scientific) to yield pMpGE010_MpEz2-sg1-MpEz3-sg1. This construct was introduced into Cam-2 gemmae using the G-AgarTrap

method⁵¹. Transformants were selected for on 0.5 Gamborg B5 plates without vitamins (Duchefa Biochemie) supplemented with hygromycin and genotyped using the following primer pairs: TH300: TACGCCCTCTCCCATTGAAC/TH301: GATACGAAGAGAACGAACCTGC for *E(z)2* and TH306: TGAGCTACATGGCTACTCTCAACC/TH307: AGCTTGGAACACGGATCTCCTG for *E(z)3*.

Expression analysis

Vegetative samples from Cam-2 and Tak-1 were collected from 100mg of apical notches from 14 day old plants grown from gemmae. The tissue was frozen in liquid nitrogen in Precellys tubes (Bertin Instruments) with 2.8mm Stainless steel beads (Bertin Corp.) and disrupted with a Precellys Evolution tissue homogenizer (Bertin Instruments) using the following settings: 7200RPM 10s, 5s pause, repeated thrice. RNA was extracted using a Spectrum Plant Total RNA kit (Sigma-Aldrich).

Embryo samples were collected by hand dissection, with one embryo per replicate (fig. S1). Embryos and the surrounding maternal calyptra tissue were dissected out of the archegoniophore into 10% RNALater (Qiagen) on Microscope slides with cavities (Marienfeld Superior) and the embryo was further dissected out of the surrounding maternal tissue. Each embryo was washed four times in a series of wells containing 150μL 10% RNALater to remove any maternal RNAs, as previously described for the pure isolation of plant embryos⁵². Each embryo was then placed in 30μL 100% RNALater on ice until sample collection was completed. The solution was diluted to 10% RNALater by the addition of 270μL RNase-free water (Zymo), vortexed gently, and the solution removed. Samples were either resuspended in 30μL 100% RNALater and stored at -70°C or in 100μL TRI reagent (Zymo). Samples were crushed with a micropestle and RNA was extracted using a Direct-zol

RNA MicroPrep kit (Zymo). All RNA samples were treated to remove DNA using a DNase Treatment and Removal kit (Invitrogen).

RNA-seq libraries were generated from total RNA following the Smart-seq2 protocol⁵³. cDNA synthesis was performed on 1µL of total RNA. 1µL of 10µM 5'-Bio-anchored oligo dT ([Btn]AAGCAGTGGTATCAACGCAGAGTACTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTT TVN) and 1µL 10mM dNTPs were added to each sample and incubated at 72°C for 3 minutes and immediately placed on ice. 7µL of a mastermix containing 0.5µL SuperScript IV (Invitrogen), 0.25µL RiboLock (ThermoFisher Scientific), 2µL Superscript IV buffer (Invitrogen), 1µL 100mM DTT (Invitrogen), 2µL 5M Betaine (Sigma-Aldrich), 0.9µL MgCl₂, 0.1µL nuclease-free water (Invitrogen), and 1µL 10µM 5'-Bio TSO ([Btn]AAGCAGTGGTATCAACGCAGAGTACrGrG+G, Exiqon) was added to each sample and the cDNA synthesis reaction took place under the following thermocycling conditions: 42°C 90min, <50°C 2min, 42°C 2min>x10, 70°C 15min. 40µL of a mastermix containing 14.5µL nuclease-free water, 25µL Q5 Hot Start 2x MasterMix (New England Biolabs), and 0.5µL 10µM 5' Bio-ISPCR oligo ([Btn]AAGCAGTGGTATCAACGCAGAGT) was added to each sample and the PCR pre-amplification took place under the following thermocycling conditions: 98°C 3min, <98°C 15s, 67°C 20s, 72°C 6min>x12, 72°C 5 minutes. Samples were cleaned up by bead purification using 1x volume of MBSPure beads (Molecular Biology Services, IMP) and samples were eluted in 15µL of 10mM Tris-HCl. 5-50ng of each sample was used for the tagmentation reaction, containing 2.5µL of 4x TAPS-DMF buffer and 1µL of Tn5 (Molecular Biology Services, IMP), which was 3 minutes at 55°C, after which samples were immediately placed on ice. Samples were purified using a DNA Clean and Concentrator kit (Zymo) using manufacturer's instructions and eluted in 10µL of 10mM

Tris-HCl. Tagmented samples were amplified by the addition of 2.5µL each of 10µM barcoded forward and reverse primers (see ref⁵³) and 15µL Q5 2x HiFi MasterMix (New England Biolabs) using the following thermocycling conditions: 72°C 3 minutes, 98°C 20s, <98°C 10s, 63°C 30s, 72°C 3 minutes>x5. Amplified samples were cleaned up by bead purification using 1x volume of MSBPure beads (Molecular Biology Services, IMP). Samples were sequenced on an Illumina NovaSeq to generate 50bp paired-end reads. Three biological replicates each of male (Tak-1) and female (Cam-2) vegetative tissue, 11 of wild-type (Cam-2 x Tak-1) embryos, and 17 of mutant (Cam-2 *e(z)2/e(z)3* x Tak-1) embryos were used for subsequent analyses.

Published transcriptomes from male and female reproductive tissues, antheridiophores and archegoniophores, respectively⁵⁴ and wild-type Tak-2 x Tak-1 embryos³⁰ were downloaded from the SRA database.

Reads were mapped to the Takv6 genome³⁴ wherein all SNP positions between Tak-1 and Cam-2 or between Tak-1 and Tak-2 were replaced with N's, depending on the genotype of the sample (refer to SNP analysis section below). Reads were preprocessed with SAMtools v1.9⁵⁵ and BEDTools v2.27.1⁵⁶, trimmed with Trim Galore (<https://github.com/FelixKrueger/TrimGalore>) and mapped with STAR v2.7.1⁵⁷. Transcripts per Million (TPM) values were calculated by RSEM v1.3.2⁵⁸. Data from RSEM were imported into R v3.5.1⁵⁹ using the tximport package v1.10.1⁶⁰. Differential gene analysis was performed using DeSeq2 v1.22.2⁶¹. Principal component analysis was performed in R v3.5.1⁵⁹. Effect size (Cohen's *d*) was calculated in R using effsize v0.7.6⁶² where $|d| < 0.2$ is no effect, $0.2 < |d| < 0.5$ is a small effect, $0.5 < |d| < 0.8$ is a medium effect, and $|d| > 0.8$ is a large effect, as previously reported⁴⁹. Heatmaps were generated in R using the pheatmap v1.0.12 package⁶³.

Chromatin profiling by CUT&RUN and analysis

Embryos and the surrounding calyptra were hand-dissected from archegoniophores and placed in Galbraith buffer (45mM MgCl₂-6H₂O, 30mM Trisodium citrate, 20mM MOPS) pH 7.0 plus 0.1% Triton X-100 and 1x cOmplete Protease Inhibitor Cocktail (Roche) on ice. Samples were crushed using a mortar and pestle on ice to release nuclei and were filtered through a 40µm filter (VWR) before staining with 2µg/mL DAPI. Nuclei were sorted on a BD FACSARIA III (BD Biosciences) to discriminate diploid embryonic nuclei from haploid maternal nuclei. Samples were sorted into 100µL of Wash buffer (20mM HEPES pH 7.5, 150mM NaCl, 0.5mM Spermidine, 1x cOmplete Protease Inhibitor Cocktail (Roche)) with 40,000 nuclei per replicate. Bio-Mag Plus Concanavalin A coated beads (Polysciences, Inc.) were activated by mixing 10µL per sample of ConA beads in 1.5mL Binding buffer (20mM HEPES-KOH pH 7.9, 10mM KCl, 1mM CaCl₂, 1mM MnCl₂). The beads were placed on a magnet, liquid was removed, and the beads were resuspended in 1.5mL Binding buffer. Liquid was again removed from the beads on a magnet and beads were resuspended in 10µL Binding buffer per sample. 10µL of the activated beads were added to each sorted nuclei sample and incubated at room temperature for 10min on a rotator. Liquid was removed from the bead-bound nuclei on a magnet and samples were resuspended in 50µL Antibody buffer (Wash buffer plus 2mM EDTA). 0.5µL of each antibody (H3K27me3 Millipore #07-449; H3K36me3 Abcam ab9050; H3K9me1 Abcam ab9045; H3 Abcam ab1791) used was added to samples while gently vortexing and samples were incubated overnight at 4°C on a shaker. Liquid was removed from the samples on a magnet and washed twice in 1mL Wash buffer before resuspending in 50µL Wash buffer. 1.16µL of 30µg/mL pAG-MNase (Molecular Biology Service, IMP) was added to each sample with gently vortexing and placed on a shaker for 10min at room temperature. Liquid was removed from the samples on a magnet and washed twice in 1mL Wash buffer before resuspending in 150µL Wash buffer. 3µL

100mM CaCl₂ was added to ice-cold samples while gently vortexing and shaken at 4°C for two hours. 100μL STOP buffer (340mM NaCl, 20mM EDTA, 4mM EGTA, 50μg/mL RNase A (ThermoFisher Scientific), 50μg/mL glycogen, 10pg/mL heterologous HEK293 DNA) was added to stop the reaction. Samples were incubated at 37°C for 10min at 500RPM then spun at 4°C for 5min at 16,000G. Samples were placed on a magnet and the liquid containing released DNA fragment was transferred to a new tube. 2.5μL 10% SDS and 2.5μL 20mg/mL Proteinase K (ThermoFisher Scientific) was added to each sample, mixed by inversion, and incubated for 1hr at 50°C. 250μL buffered phenol-chloroform-isoamyl solution (25:24:1) was added to each sample, followed by vortexing and transfer to MaXtract tubes (Qiagen). Samples were spun for 5min at 16,000G. 250μL chloroform was added and samples were spun for 5min at 16,000G. The top aqueous phase was transferred to a fresh tube containing 2μL 2mg/mL glycogen. 625μL 100% EtOH was added before vortexing and chilling at -20°C overnight. DNA extraction continued with spinning for 10min at 4°C at 20,000G. The supernatant was poured off and 1mL 100% EtOH was added to the samples before spinning again for 1min at 4°C at 16,000G. Supernatant was discarded and samples air-dried before dissolving in 50μL 0.1x TE. A NEBNext Ultra II DNA library prep kit for Illumina (New England Biolabs) was used according to the manufacturer's instructions for sample library preparation. Samples were sequenced on either an Illumina HiSeqv4 or NovaSeq to generate 50bp paired-end reads. Two biological replicates were used for each sample for H3K27me₃, H3K36me₃, H3K9me₁ and H3 in wild-type (Cam-2 x Tak-1) embryos and mutant (Cam-2 *e(z)2/e(z)3* x Tak-1) embryos.

Reads were mapped to the Takv6 genome³⁴ wherein all SNP positions between Tak-1 and Cam-2 were replaced with N's (refer to SNP analysis section below). File processing and mapping parameters were performed as previously published²⁶. Chromatin enrichment per

gene was calculated by counting the number of reads and normalizing to 1x genome coverage.

SNP analysis

Whole genome sequencing of Cam-2 was done as previously described³⁴. 5g of 14 day old Cam-2 plants grown from gemmae were collected and frozen in liquid nitrogen. Samples were crushed using a mortar and pestle on ice and ground further in 25mL PVPP buffer (50mM Tris-HCl pH 9.5, 10mM EDTA, 4M NaCl, 1% CTAB, 0.5% PVPP, 1% beta-mercaptoethanol). The mixture was divided into two 50mL Falcon tubes and incubated at 80°C for 30 minutes in a water bath. Samples were cooled to room temperature and 7.5mL chloroform was added to each tube, followed by 5mL TE-saturated phenol after mixing. Samples were spun at 20,000G for 5 minutes at room temperature and the upper aqueous phase was transferred to a new 50mL tube. 1x volume of water and 4x volume of 100% EtOH were added and mixed, and samples were frozen at -70°C. Tubes were thawed and spun at 20,000G at 4°C for 15 minutes. The supernatant was poured off, samples were spun again briefly, and the remaining supernatant pipetted off. 2mL of 1x TE was added to each tube and incubated at 60°C for 10 minutes without mixing. The supernatant was transferred to another tube and incubated at 60°C for 10 minutes without mixing. 2μL of RNaseA (ThermoFisher Scientific) was added and samples incubated at 37°C for 5 minutes. 500μL was split into 2mL tubes and 50μL of 3M sodium acetate pH 5.2 and 1mL 100% EtOH were added. Samples were incubated at -20°C for 30min and spun at 13,000RPM for 15min. After removing the supernatant, pellets were rinsed twice with 1mL 70% EtOH and spun at 13,000RPM for 5min. Pellets were dried for 90s at 65°C and resuspended in 1mL 1xTE.

Library preparation was done by tagmentation. Briefly, 1μL gDNA was mixed with 2.5μL 4x TAPS-DMF buffer and 5μL activated Tn5 (Molecular Biology Services, IMP).

Tagmentation proceeded for 5min at 55°C before cooling on ice. Samples were purified with a Zymo DNA Clean and Concentrator kit (Zymo) according to manufacturer instructions and eluted in 10µL 10mM Tris-HCl. PCR amplification was done by adding 2.5µL each of 10µM forward and reverse primers, plus 15µL NEBNext 2x HiFi PCR MasterMix (New England Biolabs) and thermocycling with the following conditions: 72°C 3min, 98°C 30s, <98°C 10s, 63°C 30s, 72°C 3min>x5. Samples were cleaned up by bead purification and sequenced on an Illumina NextSeq550 to generate 75bp paired-end reads.

Reads were preprocessed with SAMtools v1.9⁵⁵, BEDTools v2.27.1⁵⁶ and Picard v2.18.27 (<http://broadinstitute.github.io/picard/>) before mapping to the Tak-1 genome with bwa v0.7.17⁶⁴. SNPs were called using gatk v4.0.1.2 and the reference genome with all SNPs replaced with N's was created⁶⁵.

Mapped reads from CUT&RUN and RNA-Seq experiments were assigned to paternal or maternal genomes using SNPSplit v0.3.4⁶⁶. Counts for the number of reads originating from either genome were calculated per sample using SAMtools v1.9⁵⁵ and BEDTools v2.27.1⁵⁶. The maternal ratio was determined by dividing the number of maternal reads by total reads per gene. For CUT&RUN data, only data from genes with more than ten total reads in each replicate were retained. For RNA-Seq data, only data from genes with more than fifty reads in total across all replicates were retained. Additionally, data from genes that were completely maternally biased in male Tak-1 RNA-Seq data or were completely paternally biased in female Cam-2 RNA-Seq data were excluded from further maternal ratio analyses.

Interphase nuclei immunofluorescence slide preparation

Sporophytes were hand-dissected from archegoniophores and placed in Galbraith buffer (45mM MgCl₂-6H₂O, 30mM Trisodium citrate, 20mM MOPS) pH 7.0 plus 0.1% Triton X-

100 and 1x cOmplete Protease Inhibitor Cocktail (Roche) on ice. Samples were crushed in a mortar and pestle on ice and filtered through a 40µm filter (VWR). 16% paraformaldehyde was added to reach a final concentration of 4% PFA and incubated on ice for 20min. Glycine was added to a final concentration of 125mM and nuclei were spotted onto glass slides and dried at room temperature for 20min.

Mitotic cells immunofluorescence slide preparation

Sporophytes were hand-dissected from archegoniophores and placed in 1x PBS with 0.1% Triton-X 100 (PBST) and 4% paraformaldehyde on ice. Samples were fixed by applying a vacuum for 15min followed by 45min at 4°C. Samples were washed thrice for 10min each with PBST at 4°C with gentle shaking. Cell walls were digested by incubating samples in PBST plus 1% cellulase (Duchefa Biochemie) at 37°C for 10min in a damp chamber. Samples were washed thrice for 10min each with PBST at 4°C with gentle shaking. Intact tissues were placed in 10µL PBST on a glass slide and squashed with a cover slip. Slides were dipped in liquid nitrogen and the cover slip was removed with a razor blade.

Zygotic cells immunofluorescence slide preparation

To obtain archegonia holding synchronized zygote, fertilization timing was synchronized using an in vitro fertilization method described previously³⁵. At 3 daf, archegoniophores were dissected under a Lynx EVO stereomicroscope (Vision Engineering) and clusters of archegonia were collected into Fixative buffer (4% PFA, 1xPBS). Fixed tissues were then dehydrated and embedded in paraffin using the Donatello tissue processor (Diapath). Paraffin sectioning was done with a HM355S microtome (Microm) with 4µm thickness. Slides were deparaffinized and rehydrated with a Gemini autostainer (Fisher Scientific) with the following protocol: Xylene 5min, Xylene 5min, EtOH 100% 5min, EtOH 100% 5min, EtOH

95% 5min, EtOH 70% 5min, EtOH 30% 5min, Running tap water 1min, Water. Antigen retrieval was performed by boiling slides in Sodium Citrate buffer pH 6.0.

Immunostaining of slides

Immunostaining of slides was done by an InsituPro VSi staining system (Intavis) as previously described, with minor modifications⁶⁷. Slides were washed for 10min with TBS with 0.1% Tween-20 (TBST) 5 times then blocked with Blocking buffer (1x TBS, 0.1% Tween-20, 2% bovine serum albumin (BSA), 5% normal goat serum (NGS)) twice for 30min each. One antibody per slide (H3K27me3 Millipore #07-449; H3K36me3 Abcam ab9050; H3K9me1 Abcam ab9045) was diluted 1:100 and slides were incubated for 6hrs. After washing with TBST six times for 10min each, slides were incubated with a 1:500 dilution of secondary antibody (Goat Anti-Rabbit IgG H&L, Alexa Fluor® 488, ab150077; Goat Anti-Mouse IgG H&L, Alexa Fluor® 568, ab175473; Goat Anti-Rabbit IgG H&L, Alexa Fluor® 647, preadsorbed, ab150083) and slides incubated for 2hrs. After eight 10min washes with TBST, slides were dried and counterstained with 1.5µg/mL 4',6-diamidino-2-phenylindole (DAPI) and mounted in Vectashield antifade mounting medium with DAPI (Vector Laboratories) before being sealed with a coverslip and nail varnish.

Immunofluorescence image acquisition

Images were acquired with a LSM 780 scanning laser confocal microscope (Zeiss).

Interphase nuclei image deconvolution

Immunofluorescence images of interphase nuclei were deconvolved with Huygens Professional v21.04 (Scientific Volume Imaging B.V.) using the CMLE algorithm with 40 iterations and SNR values as follows: 6 for WT H3K27me3 samples H3K27me3 channel, 8

for WT H3K27me3 samples DAPI channel, 2 for WT H3K9me1 samples H3K9me1 channel, 4 for WT H3K9me1 samples DAPI channel, 4 for WT H3K36me3 samples H3K36me3 channel, 5 for WT H3K36me3 samples DAPI channel, 2 or 4 for mutant H3K27me3 samples H3K27me3 channel, 3 for mutant H3K27me3 samples DAPI channel.

Interphase nuclei image nuclei segmentation

Nuclei were identified from DAPI signal marking DNA in each immunofluorescence image.

An adaptive thresholding technique was used, based on the creation of a sequence of 20 threshold values spanning a range from a clearly too low threshold to a clearly too high threshold. A sequence of masks was thus obtained for each 3-dimensional image by thresholding it using these values. Subsequently, a maximum intensity projection of each mask was computed and size of each mask projection was evaluated. Typically, starting from the lowest threshold, such sequence first decreased rapidly, followed by a wide plateau, and ending by a decreasing tail near the highest threshold value (fig. S10A). The nearly constant plateau was detected by thresholding absolute value of neighbor size differences. Again, a sequence of thresholds was used, starting from a minimum equal to 1/10 of average of the differences, until the length of such plateau was larger than 6. Finally, the segmentation in the middle of the plateau was taken as the final one. In the last step eventual holes in the 3-dimensional mask were filled by a hole-filling operation and eventual thin gaps in the mask were filled by binary closing.

Nucleus masks thus obtained were subsequently visually inspected. Except for a few cases, the nuclei were labeled correctly. The incorrect ones were manually adjusted by means of drawing functions in FIJI v2.0.0⁶⁸.

Interphase nuclei image foci segmentation

Foci within the nucleus area were detected separately for the DAPI and the immunofluorescence channels by k-means classification in 2, 3 and 4 classes, which in our case of single-valued data specified one, two or three threshold values. The foci mask was then computed by thresholding the input data using the highest of these thresholds. The results were visually inspected and classification in 3 classes was then taken for further processing and evaluation. Only the largest foci with size bigger than 20% of the nucleus size were considered (fig. S10B).

In a limited number of cases, one or two foci were missing, or foci were too large. In the first case there were two possibilities how to identify more foci: either by decreasing the expected foci size or by decreasing the threshold value. Thus, in a loop we multiplied both these values by a coefficient until the desired number of foci was reached. In the second case with too large foci, classification in 4 classes was used, which increased the highest threshold and simultaneously decreased the size of the foci.

Combined Fluorescence *in situ* hybridization (FISH) and immunostaining method

Tissue fixation, nuclei isolation and flow cytometry were performed as described⁶⁹.

A circular barrier was made with an ImmEdge™ Hydrophobic Barrier PAP Pen (Vector Laboratories) on the charged adhesion slide of a glass slide (ThermoFisher Scientific). Size of the circle was ~ 0.7 cm diameter and ≥ 0.5 cm line thickness. Slides were dried for 30 min. 20µl of the nuclei suspension was transferred into PCR tubes and nuclei were incubated at 65°C for 30 min within a PCR Thermal Cycler. Heat shock treated nuclei were immediately transferred to ice for 5 min. 5 or 10µl of 0.1mg/ml RNase A (in 2x SSC buffer) was spotted into the circle drawn on the slide and mixed with 10µl (containing at least 1×10^4 nuclei) of heat shock treated nuclei. The solution was spread within the circle barrier.

Slides were incubated at 37°C in a slide hybridizer (ThermoBrite) for 1 hour under a humid environment. At the end of this incubation, a very thin layer of solution remained on the glass slide. After incubation, the slides were treated for about 1min each by dipping up and down till the streaks go away in an ethanol series (100%, 95%, 90%, 80%, 60%, 30% EtOH). The slides were then treated in antigen retrieval buffer (10 mM sodium citrate pH 6.0) at room temperature for 5 min and then the antigen retrieval was started by boiling the slides for 10-12 min in a microwave at 700W. Slides were post-fixed in 4% formaldehyde solution 10 min after the slides cooled down to room temperature. After post-fixation, the slides were treated for about 1 min each by dipping up and down in ethanol series (30%, 60%, 80%, 90%, 95%, 100% EtOH). Slides were dried at room temperature for 1 hour.

The subsequent probe denaturation, hybridization, washing, and detection steps were performed according to ref⁷⁰ with minor changes. Anti-Histone H3 (mono methyl K9) antibody (Abcam, ab9045) or Anti-trimethyl-Histone H3 (Lys27) Antibody (Millipore, #07-449) was diluted 1:500 in antibody buffer (5% BSA, 4x SSC, 0.2% Tween 20). 10µl of the antibody mixture was pipetted onto the slides. The slides were incubated in a humid box at 37°C for 1 hour. After 1 hour of antibody binding, slides were washed for 5 min in a solution of 4x SSC with 0.2% Tween 20 in a foil-wrapped jar at room temperature on the shaker 3 times. 100µl 1:150 Anti-rabbit Alexa Fluor 546-conjugated goat antibody (Invitrogen, AB_2534093) was dropped onto the slides. The slides were incubated at 37°C for 1 hour followed by 3 times (5 mins each) washing steps. Then, the slides were mounted with 5µl SlowFade™ Diamond Antifade Mountant (Invitrogen). Slides were covered with a coverslip and sealed with nail polish. Images were acquired with a LSM 710 scanning laser confocal microscope (Zeiss).

Probe labeling for FISH

Probes were labeled according to the Nick Translation-based DNA Probe Labeling method (Roche). Tak-1 and Tak-2 genomic DNA was extracted by CTAB method⁷¹. For U chromosome probe and U chromosome competition probe, the Tak-2 gDNA was used as template. For V chromosome probe and V chromosome competition probe the Tak-1 gDNA was used as template. Fluoroprobe labelling mix: For V/U chromosome probe (dATP, dCTP, dGTP, dTTP, Dig-dUTP); For V/U chromosome competition probe (dATP, dCTP, dGTP, dTTP). For U chromosome FISH, U-chromosome probe and 5 times V-chromosome competition probe were loaded. For V chromosome FISH, V-chromosome probe and 5 times U-chromosome competition probe were loaded.

Tissue clearing and DAPI staining of zygotes

Tissue clearing and DAPI staining for 3 daf zygotes were done as described previously³⁵. Stained samples were mounted in Vectashield antifade mounting medium with DAPI (Vector Laboratories). Images were taken by with a LSM780 confocal microscope (Zeiss).

Nuclear envelope visualization

To observe the nuclear envelope of 3 daf zygotes, ECpro:SUN-GFP³⁵ females were fertilized with wild-type sperm and zygotes were excised under a Lynx EVO stereomicroscope (Vision Engineering) and mounted in half-strength Gamborg B5 media without vitamins (Duchefa Biochemie) liquid medium. Samples were observed under a Nikon C2 confocal laser-scanning microscope (Nikon Instech).

Mutant fitness analyses

Four gemmae from Cam-2 and Cam-2 $e(z)2/e(z)3$ plants were grown together. Images of each gemmaling was taken at four, seven and ten days after planting using a Lynx EVO stereomicroscope (Vision Engineering). The area of each gemmaling was calculated using FIJI v2.0.0⁶⁸ and plotted as a smoothed curve using the loess function and formula $y \sim x$ in R v3.5.1⁵⁹ with the ggplot2 v3.3.5 package⁷².

Gemmae from Cam-2 and Cam-2 $e(z)2/e(z)3$ were planted on Grodan and monitored until the first archegoniophores were visible. Pictures were taken after all replicates had produced archegoniophores to illustrate the synchronicity of archegoniophore developmental stage.

Images of fully dissected embryos (see Expression analysis sample collection section above for details) were taken with Lynx EVO stereomicroscope (Vision Engineering). The height and width of each embryo was calculated in FIJI v2.0.0⁶⁸ using images of a calibration slide as reference. The sample area was calculated by multiplying height and width.

Aborted embryos can be identified by a browning of tissue, collapse of tissue within the calyptra, and the outgrowth of the perianth without growth of the embryo within. Embryo survival was calculated as the number of green, non-collapsed embryos per archegoniophore divided by the number of perianths with or without live embryos.

Mature embryos can be identified by the yellowing of tissue due to the production of spores within. The percentage of embryos producing spores was calculated as the number of mature yellow embryos per archegoniophore divided by the number of perianths with or without live embryos.

Spore germination was assessed by counting the number of sporelings growing out from spots of serially diluted spore solutions from single sporophytes. Mature embryos were dissected from archegoniophores, dried for one week, and frozen at -70°C. Frozen embryos

were thawed and ruptured in 100µL sterile water using a sterile pipette tip. 80µL of the spore suspension was transferred to a tube containing 420µL sterile water. 500µL of 0.1% NaDCC (Sigma-Aldrich) was added to each sample and tubes were inverted and spun at 13,000RPM for 1 minute. The supernatant was removed, and spores were resuspended in 100µL sterile water. 20µL of spore suspension was spotted onto plates of half-strength Gamborg B5 media without vitamins (Duchefa Biochemie) and 1% agar. 20µL of spore suspension was carried to a tube containing 60µL sterile water. The process was repeated until dilutions of 1:1, 1:4, 1:16 and 1:64 were spotted. Images of sporeling germination and growth were taken at 11 days after planting.

Statistical analyses

Statistical comparisons means of FISH immunostaining images and mutant growth analyses were performed with Wilcoxon tests in R v3.5.1⁵⁹ with the ggpubr v0.4.0 package⁷³. Spearman correlations were calculated in R and with the ggpubr package.

Data availability:

The CUT&RUN and RNA-seq sequencing datasets generated for the current study will be made available in the Gene Expression Omnibus (GEO) upon publication. Whole-genome sequencing data are deposited under BioProject accession number PRJNA795113. Publicly available datasets can be accessed under the DDBJ Sequence Read Archive accession numbers DRR050346-DRR050348 and DRR050351-DRR050353 and the NCBI Sequence Read Archive accession numbers SRR1553297-SRR1553299. Source data are provided with this paper. Original images will be deposited online upon publication.

Code availability:

The code used in this paper will be deposited online upon publication.

Acknowledgements:

We thank A. Pauli, A. Burga, and I. Patten for suggestions and critical reading of the manuscript. F.B. acknowledges support from the PlantS, next-generation sequencing and histopathology facilities at the Vienna BioCenter Core Facilities (VBCF), and the BioOptics facility and Molecular Biology Services from the Institute for Molecular Pathology (IMP). This work was funded by FWF grants P26887, P28320, P32054, P33380 to F.B., FWF doctoral school DK W1238 to S.A.M., and European Research Council under the European Union's Horizon 2020 research and innovation programme 757600 to C.L.

Author contributions:

S.A.M. and F.B. conceived and designed the experiments. S.A.M. performed the whole genome sequencing, RNA-seq, CUT&RUN, embryonic nuclei immunostaining, and fitness measurements. T.H. performed zygotic nuclei immunostaining. T.H. and S.A. generated material used in this study. N.W. performed immuno-FISH experiments. E.A. and S.A.M. performed statistical analyses and curated data. M.S. and S.A.M. analyzed image data. F.B. and C.L. supervised the study. S.A.M. and F.B. wrote the manuscript with input from C.L. and T.H.

Competing interest declaration:

The authors declare no competing interests.

Correspondence and requests for materials should be addressed to Frédéric Berger

Supplementary Video 1.

Movie of the dissection of a representative Marchantia embryo from surrounding calyptra of maternal origin.

Supplementary Table 1.

List of Marchantia chromatin-related genes and their expression status in embryos relative to other tissues.

Extended data figure legends:

Extended Data Fig. 1 | Maternally biased transcription in embryos. **a**, Images of hand-dissected embryos after removal of perianth and calyptra of maternal origin. Solid single arrows indicated isolated embryos. Double arrows indicate the removed calyptra. Scale bar as indicated. **b**, Percentage of measured genes within each category of maternal ratio (p_m) of transcription in wild-type (Tak-2 x Tak-1) embryos. Segments are for paternal ($p_m < 0.05$), paternal bias ($0.05 < p_m \leq 0.35$), no bias ($0.35 < p_m < 0.65$), maternal bias ($0.65 \leq p_m < 0.95$), and maternal ($0.95 \leq p_m$) expression of genes, with the number of genes indicated above each bar. **c**, Histogram of the maternal ratio (p_m) of transcription per gene in wild-type (Tak-2 X Tak-1) embryos. Each bin is 0.01 units wide. **d**, Scatterplot of gene expression (arcsinh transformed Transcripts per Million (TPM)) versus transcription maternal ratio per gene in wild-type (Cam-2 X Tak-1) embryos. Spearman correlation is indicated. **e**, Transcription maternal ratio per gene along the length of each chromosome. Vertical lines indicate the end of each chromosome. **f**, Violin plots of transcription maternal ratio of genes per chromosome. Sex chromosomes are excluded as alleles could not be resolved.

Extended Data Fig. 2 | Differential gene expression analysis of embryos to parents. **a**, Venn diagram of upregulated genes in wild-type embryos compared to male (Tak-1) and female (Cam-2) parents. **b**, Venn diagram of downregulated genes in wild-type embryos compared to male (Tak-1) and female (Cam-2) parents. **c**, Violin plots of maternal ratio of transcription for shared differentially expressed genes in embryos versus wild-type vegetative tissue from parents. Cohen's d effect size values are indicated for pairwise comparisons of Down to None, Up to None, and Up to Down where $|d| < 0.2$ is no effect and $0.2 < |d| < 0.5$ is a small effect, as previously reported (32). **d**, Volcano plot of a differential gene expression analysis between wild-type embryos and the male parent. The negative log transformed p-value per gene is plotted against the log2 fold-change in expression. Vertical dashed lines indicate a log2 fold-change of -1 and 1. The horizontal dashed line indicates a p-value of 0.05. Dots not in grey indicate chromatin-related genes, blue for significantly downregulated genes, red for significantly upregulated genes, black for genes not significantly downregulated nor upregulated. **e**, Heatmap of gene expression of Polycomb Repressive Complex 2 subunits across Marchantia development. Vegetative male (Tak-1) and female (Cam-2) tissues give rise to male and female sexual organs (antheridiophores and archegoniophores, respectively; data from (37)). Wild-type embryos are from Cam-2 x Tak-1 crosses. Values shown are arcsinh transformed Transcript per Million values.

Extended Data Fig. 3 | Paternally biased H3K27me3 in embryos. **a**, H3K27me3 maternal ratio per gene along the length of each chromosome. Vertical lines indicate the end of each chromosome. **b**, Violin plots of H3K27me3 maternal ratio of genes per chromosome in wild-type embryos. Sex chromosomes are excluded as alleles could not be resolved. **c-f**, Percentage of measured genes within each category of maternal ratio (p_m) of **c**, H3K9me1, **d**, H3, and **e**, H3K36me3 in wild-type embryos. Segments are for full paternal ($p_m < 0.05$), paternal bias ($0.05 < p_m \leq 0.35$), no bias ($0.35 < p_m < 0.65$), maternal bias ($0.65 \leq p_m < 0.95$), and full maternal ($0.95 \leq p_m$) chromatin enrichment of genes, with the number of genes indicated above each bar.

Extended Data Fig. 4 | Quantification of immunofluorescence experiments. **a**, Number of heterochromatic foci per nucleus in wild-type embryos. **b**, Percentage of nuclear area covered by heterochromatic foci in wild-type embryos. **c**, Percentage of immunofluorescence signal located within heterochromatic foci in wild-type embryos. **d**, Percentage of heterochromatic foci area overlapping with H3K27me3 foci in wild-type embryos. **e**, Quantification of sex chromosome FISH signal located within H3K27me3 or H3K9me1 heterochromatic foci. *P* values are indicated, unpaired two-tailed Wilcoxon test.

Extended Data Fig. 5 | Immunofluorescence of pronuclei. Images of wild-type zygotes at 3 days after fertilization immunostained against H3K27me3 (red) and H3 (green) and counterstained with DAPI (blue). Paternal pronuclei are indicated with a cyan box in each image. Contrast is enhanced for each image and channel independently for visualization purposes. Scale bars as indicated.

Extended Data Fig. 6 | Vegetative phenotypes of $e(z)2/e(z)3$ mutants. **a**, Structure of $E(z)2$ and $E(z)3$ genes and the mutations generated in each. **b**, Growth of Cam-2 $e(z)2/e(z)3$ mutants in the vegetative stage relative to Cam-2 wild type. Area of vegetative haploid plant growth as a function of the number of days after planting gemmae (propagules). **c**, Timing of the appearance of sexual organs in Cam-2 $e(z)2/e(z)3$ mutants relative to Cam-2 wild type as a function of the number of days after planting. *P* value is indicated, unpaired two-tailed Wilcoxon test.

Extended Data Fig. 7 | Chromatin organization of mutant embryos. **a**, Representative set of wild-type and mutant embryo immunofluorescence images. Images are maximum intensity projections. Scale bars are as indicated. **b**, Log2 ratio of H3K27me3 enrichment between mutant and wild type. Brown scale indicates greater H3K27me3 enrichment in the mutant, whereas the blue scale indicates greater H3K27me3 enrichment in wild type. **c**, Percentage of measured genes within each category of maternal ratio (p_m) of H3K27me3 in mutant embryos. Segments are for full paternal ($p_m < 0.05$), paternal bias ($0.05 < p_m \leq 0.35$), no bias ($0.35 < p_m < 0.65$), maternal bias ($0.65 \leq p_m < 0.95$), and full maternal ($0.95 \leq p_m$) H3K27me3 of genes, with the number of genes indicated above each bar.

Extended Data Fig. 8 | Transcription in mutant embryos. **a**, Percentage of measured genes within each category of maternal ratio (p_m) of transcription in mutant embryos. Segments are for full paternal ($p_m < 0.05$), paternal bias ($0.05 < p_m \leq 0.35$), no bias ($0.35 < p_m < 0.65$), maternal bias ($0.65 \leq p_m < 0.95$), and full maternal ($0.95 \leq p_m$) expression of genes, with the number of genes indicated above each bar. **b**, Heatmap of Spearman correlations of gene features in mutant embryos. **c**, Scatterplot of transcription maternal ratio versus H3K27me3 maternal ratio per gene in mutant embryos. **d**, Scatterplot of arcsinh transformed H3K27me3 enrichment versus transcription maternal ratio per gene in mutant embryos. **e**, Scatterplot of arcsinh transformed Transcript per Million (TPM) values versus transcription maternal ratio per gene in mutant embryos. **f**, Scatterplot of arcsinh transformed TPM values versus H3K27me3 enrichment per gene in mutant embryos. **g**, Scatterplot of arcsinh transformed TPM values versus H3K27me3 maternal ratio per gene in mutant embryos. Spearman correlations are indicated for each scatterplot.

Extended Data Fig. 9 | Phenotypic defects of mutant embryos. **a**, Embryo size of wild-type and mutant embryos 13 and 16 days after fertilization (daf) measured by the area of a bounding box. **b**, Percentage of wild-type and mutant embryos per female sex organ that survive to maturity at 25 and 32 daf. **c**, Percentage of wild-type and mutant embryos per female sex organ that have produced spores at 25 and 32 daf. *P* values are indicated, unpaired two-tailed Wilcoxon test.

Extended Data Fig. 10 | Immunofluorescence image analysis. **a**, Nucleus segmentation. Top panel: A sequence of 20 segmentations created by thresholding. Bottom panel: Differences (*d*) between nucleus mask sizes (blue curve). Plateau identified by estimating a threshold value d_{thr} (orange curve). The selected segmentation is in the center of the plateau (green arrow). **b**, Foci segmentation. Left: H3K27me3 and DAPI image bands with the overlaid nucleus border detected from the DAPI band. Right: Masks of the nucleus and detected foci.

a **b** **c**

a **b**