Running Head: Can livestock be functional substitutes of native herbivores?

Functional substitutability of native herbivores by livestock for long-term soil carbon sequestration depends on microbial decomposers.

Shamik Roy and Sumanta Bagchi*

Centre for Ecological Sciences, Indian Institute of Science, Bangalore – 560012, India

*Email: sbagchi@iisc.ac.in

ORCID: SR, 0000-0002-9916-5651; SB, 0000-0002-4841-6748.

Author contributions

S.R. and S.B. designed research; S.R. performed research; S.R. and S.B. analysed data; and S.R. and S.B. wrote the paper.

Conflict of Interest: The authors declare that they have no conflict of interest.

Keywords: Carbon use efficiency, Land-use, Grassland, 4-per-mille, Microbiome rewilding, Structural equation modelling (SEM)

ABSTRACT

The influence of native mammalian herbivores on ecosystem functions is declining as domestic livestock have displaced them to become the dominant grazers across the world. Alongside rewilding, restoration, and conservation, it is now important to address whether livestock can assume the functional roles of displaced native herbivores in grasslands, savannas, and steppes. If yes, this has implications for the future of biodiversity, ecosystem functions and services, food-security, and other social and environmental dimensions. We test competing alternative hypotheses with a replicated, long-term, landscape-level experiment to ask whether livestock in the Trans-Himalayan ecosystem can match decadal-scale (2006-16) soil-C sequestration under native herbivores. Despite comparable settings, livestock were imperfect substitutes and could only attain three-fourths of soil-C stocks under native herbivores. We evaluate multiple lines of evidence surrounding this difference with a grazing-exclusion experiment (nutrient-cycling, microbial metabolism, and e-DNA), and assess their inter-relationships. Livestock and native herbivores differed in their effects on several soil microbial variables. Structural equation models supported the hypotheses that livestock alter soil microbial communities towards lower fungal abundance, which is detrimental for microbial carbon use efficiency (CUE) and ultimately for soil-C storage. Lags in soil-C storage can be addressed through microbial restoration and rewilding of soil. Therefore, the idea of functional substitutability must be leveraged alongside microbial rewilding to restore and conserve ecosystem functions. Together they can reconcile conflicting demands from food-security and ecosystem services in ways that have eluded conventional approaches to become an integral part of nature-based climate solutions.

INTRODUCTION

Large mammalian herbivores have undergone widespread declines in their abundance and distribution across the world's grasslands, steppes, and savannas. Pre-historic humans might have hunted many megafauna to extinction. More recently, humans have progressively replaced them with livestock across the world. Relatively intact assemblages of native herbivores are now increasingly confined to parks and reserves; elsewhere, livestock have not only become the most abundant herbivores, but they also represent the most geographically expansive land-use today (Asner et al. 2004, Bar-On et al. 2018). Further displacement of native herbivores seems inevitable due to steadily rising demands for livestock-products to accommodate evolving human-diets (Thornton 2010, Leclère et al. 2020). This severely restricts and alters the erstwhile influence of native herbivores on ecosystem structure and functions, their broad influence on global biogeochemical cycles, and their influence on soil-C in particular (Brodie et al. 2021, Roy and Bagchi 2021).

Increasingly, it is important to ask whether livestock can maintain the functional roles of the displaced native herbivores (Malhi et al. 2016, Cromsigt et al. 2018, Lundgren et al. 2020). If livestock can maintain ecological functions and services of their native counterparts, then such functional substitutability can have implications for the future of biodiversity, ecosystem functions and services, food-security, and other social and environmental dimensions (Sayer et al. 2013, Veblen et al. 2016, Willemsen et al. 2020, Western et al. 2020, Delabre et al. 2021). The extent to which livestock can maintain ecosystem functions and services, particularly soil-C sequestration, becomes important to inform policies over sustainable use of grazing ecosystems and leverage their potential as natural climate solutions (Smith 2014, Cromsigt et al. 2018, Bossio et al. 2020).

Opportunities to restore and maintain ecosystem processes using livestock as surrogates of displaced native herbivores arise from the mutual similarity in their functional

traits (Malhi et al. 2016, Lundgren et al. 2020, Schowanek et al. 2021). Indeed, traits are better predictors of ecosystem functions than taxonomic identity of species (Hooper et al. 2005). Analyses of key traits, such as body size, diet-choice, and fermentation type, hypothesize that livestock can be approximate surrogates for native herbivores, i.e., nearest neighbors in trait-space (Malhi et al. 2016, Lundgren et al. 2020, Schowanek et al. 2021). However, empirical evidence on ecosystem functions and services is needed before these ideas can be implemented toward ecosystem management, and such evidence is lacking (Reinhart et al. 2021).

From this functional-substitutability hypothesis one can ask (Betts et al. 2021) – if there is sufficient overlap in trait-space between livestock and native herbivores, then to what extent can they act as surrogates? It is challenging to address this question because many modern livestock-production systems no longer have native counterparts within comparable ecological settings that can allow a fair comparison. Unlike intensive livestock production systems that are often associated with various forms of environmental degradation, traditional pastoral production systems where both livestock and native herbivores occur in multiple-use landscapes use (Asner et al. 2004, du Toit et al. 2009, Veblen et al. 2016, Western et al. 2020) can be testbeds to assess whether livestock are good substitutes, or not. While this functional premise does not expect livestock to be perfect replacements for the displaced native herbivores (Bagchi and Ritchie 2010, Sitters et al. 2020), it is important to estimate the degree to which their ecological roles are matched to identify gaps and shortfalls as well as the scope for improved livestock production systems (Smith 2014, Western et al. 2020, Reinhart et al. 2021, Delabre et al. 2021).

Here we address impacts of livestock and native herbivores on soil-C sequestration as grazing by large mammals is an important component of the global carbon cycle (Schmitz et al. 2014, Roy and Bagchi 2021). We compare their ecological impacts in Spiti region of

northern India, which is part of the greater Trans-Himalayan ecosystem spread across the Tibetan plateau and surrounding mountains. Soil-C sequestration in the Trans-Himalaya is particularly important under human land-use for livestock production and under ongoing climate change (Murthy and Bagchi 2018, Naidu and Bagchi 2021). We investigate long-term (decadal-scale) net soil-C sequestration under livestock and native herbivores, and their influence on key microbial decomposer functions related to the C-cycle (Falkowski et al. 2008, Sinsabaugh et al. 2016, Roy and Bagchi 2021). We take advantage of several key features in the Trans-Himalaya: (1) the livestock are a multi-species assemblage that show considerable trait-overlap with the native herbivores (Fig. 1), (2) livestock and native herbivores occupy distinct but juxtaposed watersheds with comparable environmental settings (Fig. S1-S4), (3) their abundances are comparable so that grazing pressure on vegetation is also similar (see Methods, Fig. S5). Absent major confounding influences, the livestock and native assemblages in Trans-Himalaya can help evaluate their functional substitutability (Malhi et al. 2016, Cromsigt et al. 2018, Lundgren et al. 2020, Schowanek et al. 2021). Specifically, we investigate (1) whether, and by how much, does grazing by livestock and native herbivores differ in its influence on long-term (decadal-scale) soil-C sequestration, and (2) whether any differences can be attributed to how each assemblage influences microbial decomposer functions in soil.

MATERIALS AND METHODS

STUDY AREA

Traditional pastoral livestock production is the major land-use in Spiti region of the Trans-Himalayan grazing ecosystem. Watersheds near human settlements are used by a multi-species livestock assemblage, and these are juxtaposed with watersheds that retain native herbivores (Fig. 1, Fig. S1). Natural terrain and barriers (canyons, rivers, high ridges) that

restrict frequent animal movement, maintain replicate watersheds under two alternative types of herbivore assemblages (Fig. S1, S2). Historical local extinctions of native herbivores and their replacement by livestock over the past decades remains a major conservation challenge in this region (Mishra et al. 2002, Namgail et al. 2013), and around the world (Veblen et al. 2016, Western et al. 2020). The extant native herbivores are ibex (*Capra sibirica*), bharal (*Pseudois nayaur*), and yak (*Bos grunniens grunniens*) that are related to wild yak (*Bos grunniens mutus*; also known as *B. grunniens* and *B. mutus*, respectively; Leslie and Schaller 2009). Currently, kiang (*Equus kiang*) are known from the fringes of the study area, and there are sporadic reports of argali (*Ovis ammon*). But Tibetan antelope (*Pantholops hodgsonii*) and Tibetan gazelle (*Procapra picticaudata*) are no longer found in the region. The livestock consist of goat, sheep, donkey, horse, cattle and yak-cattle hybrids (Fig. 1). The livestock and native herbivores (after including kiang and argali) show sufficiently high overlap in key traits (Lundgren et al. 2020), such as body mass, dietary guild, fermentation type, sexual-dimorphism, graminoid, limb morphology (Fig. 1). In recent decades, total animal biomass has remained comparable (6.9×10^4 kg for native and 7.2×10^4 kg for livestock; Bagchi and Ritchie 2010, Singh et al. 2015), such that grazing pressure on vegetation has also remained similar for sufficiently long periods (Fig. S5).

The two types of watersheds are comparable in their abiotic conditions (Fig. S3–S4). Climate in Spiti is cold and semi-arid with temperature ranging from 25 °C in summer to below -30 °C in winters (Fig. S6). Precipitation occurs as winter-snow ($100\text{--}200$ cm yr⁻¹) and summer-rain ($150\text{--}300$ mm yr⁻¹), and vegetation growth-season is short (May–August, Fig. S6). Vegetation in this treeless landscape consists of grasses (*Poa*, *Elymus*, *Festuca*), sedges (*Carex*, *Kobresia*), forbs and shrubs (*Artemisia*, *Lindelofia*, *Caragana*). Both types of watersheds have an overlapping pool of plant species with spatial variation in vegetation composition. Livestock watersheds have high abundance of sedges and grasses, have higher

ground cover as well as higher aboveground plant production than grass-and-forb vegetation in the native watersheds (Fig. S5, Bagchi and Ritchie 2010).

SOIL CARBON

Between 2006 and 2016, we measured soil-C at regular intervals in four watersheds with native herbivores, and another four watersheds with livestock (Fig. S1, S2). In each watershed, we set up 3-4 replicated permanent plots ($10 \times 10 \text{ m}^2$), for a total of $n=15$ plots under native herbivores, and another $n=15$ plots under livestock. We collected soils with a 5 cm diameter and 20 cm deep cores, and measured carbon content (TruSpec, Leco, USA, and Elementar Analysensysteme, GmbH, Germany) to compare long-term soil-C stocks (mg C cm^{-3}) under livestock and native herbivores.

GRAZING-EXCLUSION EXPERIMENT

We distinguish whether any observed difference in soil-C between the watersheds is merely an artefact of pre-existing site-specific conditions, or it can be attributed to grazing by the alternative herbivore assemblages. For this, starting in 2005, we set up experimental grazer-exclusion with a paired-adjacent fence at each location (also $10 \times 10 \text{ m}^2$, each). In 2017, after up to 12 years of herbivore-exclusion, we estimated grazer-effects on key soil microbial functions in the paired-adjacent fenced and control plots.

SOIL MICROBIAL FUNCTIONS

We estimated the response of key soil microbial functions to grazing, that are related to the carbon cycle and net soil-C sequestration: (1) Basal respiration (BR), (2) Potential, or substrate-induced respiration (SIR), (3) Soil microbial community composition, broadly as fungal and bacterial fractions, (4) Microbial biomass C and N (MBC and MBN), (5) labile

Carbon (LC) and labile Nitrogen (LN) fractions of soil organic matter, (6) Metabolic quotient (qCO_2), and (7) Microbial carbon use efficiency (CUE). Since these microbial processes have seasonal dynamics (Bagchi et al. 2017), we measured them at five time-points during the growing season, between May and September. We followed standard laboratory protocols to measure these microbial variables (Robertson et al. 1999, Bagchi et al. 2017, Roy and Bagchi 2021). Briefly, for BR we used an alkali-trap where we incubated 4 g soil at 60% water holding capacity and measured respired CO_2 ($mg\ C\ g^{-1}\ soil\ day^{-1}$). Similarly, we measured potential microbial respiration ($mg\ C\ g^{-1}\ soil\ day^{-1}$) using substrate-induced respiration (SIR) after adding 0.5% (w/v) of glucose to soil.

We estimated broad microbial community composition as fungal and bacterial contributions to potential respiration under selective inhibition using Streptomycin (anti-bacterial) and Cycloheximide (anti-fungal). We used chloroform fumigation-extraction to estimate MBC ($mg\ C\ g^{-1}\ soil$), and MBN ($mg\ N\ g^{-1}\ soil$). We measured SOM from weight loss on ignition at 450 °C after 4 h. We measured labile fractions ($mg\ g^{-1}\ soil$) of SOM using two-step hydrolysis with H_2SO_4 . We estimated metabolic quotient as the ratio of BR and MBC (qCO_2 , respiration per unit biomass, hr^{-1}). We estimated CUE from established relationships between key microbial extracellular enzymes (carbon-acquiring Beta-glucosidase, alongside nitrogen-acquiring Leucine aminopeptidase and *N*-acetylglucosaminidase), MBC and MBN, and labile fractions of C and N in SOM (Sinsabaugh et al. 2013, 2016). We measured activity of the three enzymes using their specific fluorogenic substrates (4-MUB beta-D-glucopyranoside for Beta-glucosidase, L-Leucine-7-amido-4-methylcoumarin hydrochloride for Leucine aminopeptidase, and 4-MUB N-acetyl-beta-D-glucosaminide for N-acetylglucosaminidase) (German et al. 2011) in soils from growth season of 2017. These parameters influence microbial metabolism to determine their investment in growth (G) and respiration (R), so that $CUE = \frac{G}{G+R}$ (Sinsabaugh et al. 2013, 2016).

Theoretically CUE can range between 0 and 1, i.e., nil to perfect efficiency, with higher values indicating greater potential for C-sequestration. But, CUE in nature generally does not exceed 0.6 (Sinsabaugh et al. 2013, 2016).

SOIL MICROBIAL DNA

Based on patterns seen in fungal and bacterial contribution to soil respiration, we evaluated community composition in more detail using their DNA markers in 2019 (Fierer et al. 2005, Maestre et al. 2015). We collected soils from ten paired-plots at four time points during the growing season and extracted microbial DNA using MOBIO Power Soil DNA Extraction Kit following manufacturer's protocols. Next, we quantified the Internal Transcribed Spacer (ITS-1 and ITS-2) regions of rRNA for fungal abundance, and region of 16s rRNA for bacterial abundance with qPCR (Fierer et al. 2005, Maestre et al. 2015). We used these genetic data to confirm whether DNA-based estimates align with respiration-based estimates of fungal and bacterial abundance. Additional details of laboratory analyses are in Appendix.

DATA ANALYSIS

We evaluated competing hypotheses (Tredennick et al. 2021, Betts et al. 2021) to determine whether the observed variation in the data merely represent site-specific and time-specific heterogeneity between replicates, or it is necessary to invoke herbivore assemblage-type (livestock or native) as an explanatory variable. We used a linear mixed-effects model to evaluate long term trends in soil-C and grazer-exclusion effects on each microbial variable. For soil-C, we used a null-model (M_0^{soilC}) with time (years, 2006-16) and plot identity as random-effects to account for background variation due to sampling locations and times. This model was evaluated against the alternative competing model (M_A^{soilC}) which included assemblage-type as a fixed-effect, alongside both random-effects. We compared both models

based on parsimony, goodness-of-fit, and likelihood-ratio test (i.e., AIC, RMSE, and LR).

Next, we determined whether there was gain or loss of soil-C with a separate model where time (years) was a fixed-effect and plot identity as random effect.

We estimated grazer-effect on each microbial variable as the log-ratio of values in the fenced and paired-adjacent control plots i.e., $\Delta = \ln \left(\frac{\text{Grazing}_i}{\text{Fenced}_i} \right)$ for the i^{th} pair, and calculated the mean and 95% CI of the log-ratio (Roy and Bagchi 2021). As earlier, we first evaluated a null-model to assess whether the observed variation in grazer-effects can simply be attributed to heterogeneity across sampling locations and times (months, May-September). As the null-model for the j^{th} microbial variable (M_0^j), we used plot identity and month as random-effects. In the competing alternative model for each variable (M_A^j), we included herbivore assemblage-type as a fixed-effect alongside the random-effects. As before, we used AIC, RMSE and LR for model comparison to judge whether herbivore assemblage-type is a necessary explanatory variable for the data (Tredennick et al. 2021). We assessed whether the data met statistical assumptions from observed and theoretical quantiles of residuals ; the variables required no further transformation (Δ is already in ln-scale). We used ‘nlme’ library in R (Pinheiro et al. 2020) to perform these analyses.

We assessed the inter-relationships between grazer-effects (Δ) on different microbial variables with structural equations models (SEM). SEM help evaluate hypothesized pathways over how change in one variable can influence another (Grace 2006). We identified SEM paths from *a-priori* examples in the literature (Fig. S9). To avoid multicollinearity in SEM arising from redundancy among variables, we removed those variables which were either highly correlated with each other ($|r| > 0.49$), or were derived from each other (Fig. S10). So, ΔSIR , ΔB , ΔMBN , and ΔqCO_2 were not included in this step, and we built SEM using ΔMBC , ΔLN , ΔLC and ΔF . We did not use genetic data from ITS and 16s rRNA in the SEM because these were collected from a subset of plots (10 fenced and control pairs in each

watershed), were also not from the same year as the other microbial variables, and because they broadly matched respiration-based estimates of community composition. As earlier, we evaluated two competing SEMs – with and without herbivore assemblage-type as an explanatory variable, i.e., SEM₀ and SEM_A. We incorporated SEM paths as mixed-effects models for each variable as piece-wise structural equations (Lefcheck 2016). We assessed goodness of fit with Fisher’s C statistic (Shipley 2009, Roy and Bagchi 2021) and report standardised path coefficients alongside their statistical significance ($\alpha=0.05$). When SEM paths are supported by data, then they offer a plausible explanation for the underlying process. When SEM paths are not supported, the hypothesized process may not give rise to the observed data.

RESULTS

LONG-TERM TREND IN SOIL-CARBON

Model comparison showed that spatio-temporal heterogeneity was not sufficient to explain long-term variation in soil-C, and it could instead be attributed to herbivore assemblage-type. There was stronger support for M_A^{soilC} than for M_0^{soilC} (difference in AIC=-25.3, and difference in RMSE=-0.61, and LR=27.3, $P<0.001$). Soils under native herbivores stored 7.7 ± 1.5 SE more carbon (mg C cm^{-3}) than under livestock ($F_{1,176}=26.6$, $P<0.001$). Effectively, areas under livestock could realize up to 71% of the soil-C found under native herbivores (Fig. 2).

SOIL MICROBIAL RESPONSES TO GRAZING

Analysis of grazer-effects revealed that the observed difference in soil-C (Fig. 2) was related to how soil microbes respond to grazing by livestock or native herbivores (Fig. 3). Livestock and native herbivores had comparable effects on B, 16s rRNA abundance, MBC, MBN, LC,

LN, and qCO₂ (Fig. 3, Table S1). For these seven variables, the competing alternative models were not supported, and their variability could be attributed to spatio-temporal heterogeneity rather than to assemblage-type (Fig. 3, Table S1). But livestock and native herbivores differed in their influence on BR, SIR, F, ITS abundance, and CUE. For these five variables, there was stronger support for the competing models than the null models (Table S1), as assemblage-type was an important explanatory variable over and above background spatio-temporal heterogeneity (Fig. 3). Grazing by livestock increased both basal and potential microbial respiration (BR and SIR), compared to grazing by native herbivores (Fig. 3). But grazing by livestock reduced both the fungal fraction of microbial respiration and abundance of fungal genetic markers (F and ITS). Respiration-based estimates of fungal and bacterial fractions were consistent with DNA-based estimates (Fig. 3). Grazing by livestock reduced CUE compared to native herbivores (Fig. 3).

INTER-RELATIONSHIPS BETWEEN SOIL MICROBIAL RESPONSES

To assess how differences in grazer-effects on one microbial variable may influence other variables, we evaluated two competing SEM models. The null model was supported by the data (SEM₀: Fisher's C = 3.34, $P=0.503$, $df = 4$, AIC = 69.34). This showed, regardless of herbivore assemblage-type, grazer-effects on microbial community composition (fungal fraction) affects CUE, and grazers also influence the labile-C pool via their effects on microbial biomass and labile-N (Fig. 4). The alternative competing model was also supported by the data (SEM_A: Fisher's C = 2.108, $P=0.716$, $df = 4$, AIC = 88.11). It showed that change in assemblage-type can influence CUE via its effect on microbial community composition (Fig. 4). Specifically, change in herbivore-assemblage from native to livestock has negative influence on fungal abundance, which in turn is detrimental for CUE (Fig. 4). Both SEM models indicate that direction and magnitude of grazer-effects are consistent with the premise

that higher CUE under native herbivores improves net soil-C storage, compared to livestock (Fig. 2, Fig. 4).

DISCUSSION

Previous studies from around the world have emphasized that grazing by large mammalian herbivores leads to net soil-C storage (Conant and Paustian 2002, Witt et al. 2011, Abdalla et al. 2018, Derner et al. 2019, Liu et al. 2020), and grazing ecosystems should be integral to nature-based climate solutions (Smith 2014, Crooms et al. 2018, Willemen et al. 2020, Bossio et al. 2020, Roy and Bagchi 2021). But it has remained unclear whether this service can be maintained when natural ecosystems are repurposed for livestock production (Reinhart et al. 2021). And if so, then to what extent?

Our results address several aspects of functional substitutability of native herbivores by livestock. We find that a multi-species assemblage of livestock can have remarkable functional similarities with the native assemblages they displace (Fig. 1). And, at comparable abundances, both assemblages gain soil-C (Fig. 2) at approximately 1‰ to reiterate the role for grazing ecosystems in nature-based climate solutions (Smith 2014, Crooms et al. 2018, Willemen et al. 2020, Naidu and Bagchi 2021, Roy and Bagchi 2021). But livestock do not emerge as perfect substitutes since they achieve only about three-fourths of the soil-C stock, compared to the native herbivores (Fig. 2). Nevertheless, consistent with the functional substitutability hypothesis, a key ecosystem service provided by native herbivores, soil-C sequestration, can also be partly realized in livestock-production systems. This can have implications for how we manage environmental impacts of food (Sayer et al. 2013, Smith 2014, Veblen et al. 2016, Willemen et al. 2020, Western et al. 2020, Delabre et al. 2021), and meet the rising demand for livestock-products in the coming decades (Thornton 2010, Leclère et al. 2020).

To determine the scope of improved management, it is important to identify the potential drivers and mechanisms that explain why livestock lag native herbivores in soil-C sequestration. Prevailing management practices have largely focused on two aspects—overgrazing and vegetation community change—which result in altered C-input to soil from plants (Bagchi and Ritchie 2010, Reinhart et al. 2021). However, this prevailing focus on C-input alone can be simplistic and inadequate since it does not account for subsequent metabolism of soil organic matter by microbes (Fontaine et al. 2004, Six et al. 2006, Falkowski et al. 2008, Sinsabaugh et al. 2013, Roy and Bagchi 2021). Our results from the grazer-exclusion experiment identify soil microbial responses as the proximate explanation for why livestock are not perfect substitutes for native herbivores (Fig. 3). Tests of the competing hypotheses address whether the observed patterns in soil-C reflects idiosyncratic and/or pre-existing differences between watersheds, or they are linked to the identity of the herbivores. We attribute this missing soil-C to how microbial processes respond to grazing when native herbivores are replaced by livestock (Fig. 3-4). However, this still leaves the possibility that livestock watersheds may have been historically overgrazed, and their soils have not yet recovered. Even though vegetation may recover from historical overgrazing, it may take much longer for soil microbial functions to recover (López Zieher et al. 2020). Seen together, these highlight the importance of soil microbial restoration in tandem with management of grazing intensity and stocking rates. CUE is a fundamental microbial trait that controls residence time of carbon in the soil, and it determines the flow of carbon through ecosystems (Six et al. 2006, Falkowski et al. 2008, Sinsabaugh et al. 2013). The observed grazer-effects on fungi (Fig. 3) are also known from other ecosystems (Bardgett et al. 1997, Zhang and Fu 2021). Direction and magnitude of difference in the microbial community under livestock reduces CUE (Fig. 3-4), and this is consistent with lower soil-C

stocks (Fig. 2) because fungi are known to have higher CUE than bacteria (Fontaine et al. 2004, Six et al. 2006).

Several overlapping factors can alter microbial communities towards lower fungi, and their individual roles may not be easily disentangled from each other. First, variation in vegetation composition can result in differences in the quality of litter-input to soil, and this in turn can influence the rate and efficiency with which microbes metabolize soil organic matter (Fontaine et al. 2004, Six et al. 2006, Sinsabaugh et al. 2013). Second, as discussed above, microbial communities may have not yet recovered from any historical overgrazing in the past (López Zieher et al. 2020), even though vegetation biomass does not indicate overgrazing is currently a concern (Fig. S5). Third, veterinary care offered to the livestock leads to inadvertent release of antibiotics and other drugs to the soil from their dung and urine, and this can alter microbial communities towards lower CUE (Wepking et al. 2019). Such overlapping factors need to be addressed together to improve soil-C sequestration, rather than in isolation.

Our results from alternative competing SEMs offer insights into how net soil-C sequestration can be improved (Fig. 4). Since replacing native herbivores with livestock is detrimental for CUE, promoting wildlife conservation over livestock-production would favor soil-C (SEM_A, Fig. 4). This would compromise livelihoods and food-security, and it circles back to the inconsistencies and conflicts between the goals of conservation and human land-use. This conventional and exclusionary approach may no longer be tenable and is also increasingly difficult to justify (Adams and Mulligan 2002); it is timely to assess alternative ideas (Smith 2014). The null model (SEM₀, Fig. 4) can offer insights into ways to resolve these conflicting demands on land-use. It suggests altered microbial communities compromise CUE regardless of whether the community shift is a fallout of land-use, or not. This opens opportunities for restoration of microbial communities to sustain ecosystem

services under human land-use (Paluch et al. 2013, Wubs et al. 2016, Mills et al. 2017).
Admittedly, there are many gaps in our current understanding of structure-function
relationships for hyper-diverse microbial communities (Bier et al. 2015). Yet, there are
encouraging signs that effective restoration can be achieved even with incomplete taxonomic
knowledge (Paluch et al. 2013, Wubs et al. 2016, Mills et al. 2017). Since livestock already
deliver 71% of the ecosystem service provided by native herbivores, further incorporating
microbial rewilding of soils into livestock-production may offset the 29% lag against native
herbivores—in the process, making livestock-production more environmentally responsible
(Smith 2014, Cromsigt et al. 2018, Willemsen et al. 2020, Bossio et al. 2020).

In conclusion, our results highlight: (1) multi-species and functionally diverse
assemblages of livestock can partly approximate the ecological roles of native herbivores,
and (2) restoration and rewilding of soil microbial communities may enable livestock to
emulate the services provided by native herbivores. Alongside continued management of
stocking rates, functional substitutability and microbial restoration in-tandem may help
reconcile rising demands for livestock-products with conservation of ecosystem functions
and services.

Acknowledgements

CSIR-India provided graduate fellowship to SR; SB was supported by MHRD-India. DST-
SERB, DST-FIST, DBT-IISc, and MoEFCC provided funding support for fieldwork and lab-
analyses. Govt. of Himachal Pradesh has supported our long-term research in Spiti. We
received generous support from Abhilash Vijay Nair, Ananya Jana, Atish Roy Chowdhury,
Debarshi Dasgupta, Deepak Shetti, Dilip Naidu, Dipshikha Chakravorty, Dorje Chhering,
Dorje Chhewang, Ishika Pramanick, Kavita Isvaran, Praveen Karanth, Pronoy Baidya, Raju

Roy, and Ritika Chatterjee. Discussions with Ben Bond-Lamberty and Mayank Kohli helped us prepare the manuscript.

References:

- Abdalla, M., A. Hastings, D. R. Chadwick, D. L. Jones, C. D. Evans, M. B. Jones, R. M. Rees, and P. Smith. 2018. Critical review of the impacts of grazing intensity on soil organic carbon storage and other soil quality indicators in extensively managed grasslands. *Agriculture, Ecosystems and Environment* 253:62–81.
- Adams, W. M., and M. Mulligan. 2002. Decolonizing nature: Strategies for conservation in a post-colonial era. Page (W. Adams and M. Mulligan, Eds.) *Decolonizing Nature: Strategies for Conservation in a Post-colonial Era*. 1st edition. Routledge, London.
- Asner, G. P., A. J. Elmore, L. P. Olander, R. E. Martin, and A. T. Harris. 2004. Grazing systems, ecosystem responses, and global change. *Annual Review of Environment and Resources* 29:261–299.
- Bagchi, S., and M. E. Ritchie. 2010. Introduced grazers can restrict potential soil carbon sequestration through impacts on plant community composition. *Ecology Letters* 13:959–968.
- Bagchi, S., S. Roy, A. Maitra, and R. S. Sran. 2017. Herbivores suppress soil microbes to influence carbon sequestration in the grazing ecosystem of the Trans-Himalaya. *Agriculture, Ecosystems and Environment* 239:199–206.
- Bardgett, R. D., D. K. Leemans, R. Cook, and P. J. Hobbs. 1997. Seasonality of the soil biota of grazed and ungrazed hill grasslands. *Soil Biology and Biochemistry* 29:1285–1294.
- Bar-On, Y. M., R. Phillips, and R. Milo. 2018. The biomass distribution on Earth. *Proceedings of the National Academy of Sciences* 115:6506–6511.
- Betts, M. G., A. S. Hadley, D. W. Frey, S. J. K. Frey, D. Gannon, S. H. Harris, H. Kim, U. G. Kormann, K. Leimberger, K. Moriarty, J. M. Northrup, B. Phalan, J. S. Rousseau, T. D. Stokely, J. J. Valente, C. Wolf, and D. Zárrete-Charry. 2021. When are hypotheses useful in ecology and evolution? *Ecology and Evolution* 11:5762–5776.
- Bier, R. L., E. S. Bernhardt, C. M. Boot, E. B. Graham, E. K. Hall, J. T. Lennon, D. R. Nemergut, B. B. Osborne, C. Ruiz-González, J. P. Schimel, M. P. Waldrop, and M. D. Wallenstein. 2015. Linking microbial community structure and microbial processes: an empirical and conceptual overview. *FEMS Microbiology Ecology* 91:fiv113.
- Bossio, D. A., S. C. Cook-Patton, P. W. Ellis, J. Fargione, J. Sanderman, P. Smith, S. Wood, R. J. Zomer, M. von Unger, I. M. Emmer, and B. W. Griscom. 2020. The role of soil carbon in natural climate solutions. *Nature Sustainability* 2020 3:5 3:391–398.
- Brodie, J. F., S. Williams, and B. Garner. 2021. The decline of mammal functional and evolutionary diversity worldwide. *Proceedings of the National Academy of Sciences* 118:e1921849118.
- Conant, R. T., and K. Paustian. 2002. Potential soil carbon sequestration in overgrazed grassland ecosystems. *Global Biogeochemical Cycles* 16:90-1-90–9.
- Cromsigt, J. P. G. M., M. te Beest, G. I. H. Kerley, M. Landman, E. le Roux, and F. A. Smith. 2018. Trophic rewilding as a climate change mitigation strategy? *Philosophical Transactions of the Royal Society B: Biological Sciences* 373.

- Delabre, I., L. O. Rodriguez, J. M. Smallwood, J. P. W. Scharlemann, J. Alcamo, A. S. Antonarakis, P. Rowhani, R. J. Hazell, D. L. Aksnes, P. Balvanera, C. J. Lundquist, C. Gresham, A. E. Alexander, and N. C. Stenseth. 2021. Actions on sustainable food production and consumption for the post-2020 global biodiversity framework. *Science Advances* 7:eabc8259.
- Derner, J. D., D. J. Augustine, and D. A. Frank. 2019. Does Grazing Matter for Soil Organic Carbon Sequestration in the Western North American Great Plains? *Ecosystems* 22:1088–1094.
- Falkowski, P. G., T. Fenchel, and E. F. Delong. 2008. The Microbial Engines That Drive Earth's Biogeochemical Cycles. *Science* 320:1034–1039.
- Fierer, N., J. A. Jackson, R. Vilgalys, and R. B. Jackson. 2005. Assessment of Soil Microbial Community Structure by Use of Taxon-Specific Quantitative PCR Assays. *Applied and Environmental Microbiology* 71:4117–4120.
- Fontaine, S., G. Bardoux, L. Abbadie, and A. Mariotti. 2004. Carbon input to soil may decrease soil carbon content. *Ecology Letters* 7:314–320.
- German, D. P., M. N. Weintraub, A. S. Grandy, C. L. Lauber, Z. L. Rinkes, and S. D. Allison. 2011. Optimization of hydrolytic and oxidative enzyme methods for ecosystem studies. *Soil Biology and Biochemistry* 43:1387–1397.
- Grace, J. B. 2006. *Structural Equation Modeling and Natural Systems*. Cambridge University Press.
- Hooper, D. U., F. S. Chapin, J. J. Ewel, A. Hector, P. Inchausti, S. Lavorel, J. H. Lawton, D. M. Lodge, M. Loreau, S. Naeem, B. Schmid, H. Setälä, A. J. Symstad, J. Vandermeer, and D. A. Wardle. 2005. Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecological Monographs* 75:3–35.
- Leclère, D., M. Obersteiner, M. Barrett, S. H. M. Butchart, A. Chaudhary, A. De Palma, F. A. J. DeClerck, M. Di Marco, J. C. Doelman, M. Dürauer, R. Freeman, M. Harfoot, T. Hasegawa, S. Hellweg, J. P. Hilbers, S. L. L. Hill, F. Humpenöder, N. Jennings, T. Krisztin, G. M. Mace, H. Ohashi, A. Popp, A. Purvis, A. M. Schipper, A. Tabeau, H. Valin, H. van Meijl, W. J. van Zeist, P. Visconti, R. Alkemade, R. Almond, G. Bunting, N. D. Burgess, S. E. Cornell, F. Di Fulvio, S. Ferrier, S. Fritz, S. Fujimori, M. Grooten, T. Harwood, P. Havlík, M. Herrero, A. J. Hoskins, M. Jung, T. Kram, H. Lotze-Campen, T. Matsui, C. Meyer, D. Nel, T. Newbold, G. Schmidt-Traub, E. Stehfest, B. B. N. Strassburg, D. P. van Vuuren, C. Ware, J. E. M. Watson, W. Wu, and L. Young. 2020. Bending the curve of terrestrial biodiversity needs an integrated strategy. *Nature* 585:551–556.
- Lefcheck, J. S. 2016. piecewiseSEM: Piecewise structural equation modelling in r for ecology, evolution, and systematics. *Methods in Ecology and Evolution* 7:573–579.
- Leslie, D. M., and G. B. Schaller. 2009. *Bos grunniens* and *Bos mutus* (Artiodactyla: Bovidae). *Mammalian Species* 836:1–17.
- Liu, X., H. Sheng, Z. Wang, Z. Ma, X. Huang, and L. Li. 2020. Does Grazing Exclusion Improve Soil Carbon and Nitrogen Stocks in Alpine Grasslands on the Qinghai-Tibetan Plateau? A Meta-Analysis. *Sustainability* 12:977.
- López Zieher, X. M., L. Vivanco, and L. Yahdjian. 2020. Soil bacterial communities remain altered after 30 years of agriculture abandonment in Pampa grasslands. *Oecologia* 193:959–968.
- Lundgren, E. J., D. Ramp, J. Rowan, O. Middleton, S. D. Schowanek, O. Sanisidro, S. P. Carroll, M. Davis, C. J. Sandom, J.-C. Svenning, and A. D. Wallach. 2020. Introduced herbivores restore Late Pleistocene ecological functions. *Proceedings of the National Academy of Sciences* 117:7871–7878.

- Maestre, F. T., M. Delgado-Baquerizo, T. C. Jeffries, D. J. Eldridge, V. Ochoa, B. Gozalo, J. L. Quero, M. García-Gómez, A. Gallardo, W. Ulrich, M. A. Bowker, T. Arredondo, C. Barraza-Zepeda, D. Bran, A. Florentino, J. Gaitán, J. R. Gutiérrez, E. Huber-Sannwald, M. Jankju, R. L. Mau, M. Miriti, K. Naseri, A. Ospina, I. Stavi, D. Wang, N. N. Woods, X. Yuan, E. Zaady, and B. K. Singh. 2015. Increasing aridity reduces soil microbial diversity and abundance in global drylands. *Proceedings of the National Academy of Sciences of the United States of America* 112:15684–15689.
- Malhi, Y., C. E. Doughty, M. Galetti, F. A. Smith, J.-C. Svenning, and J. W. Terborgh. 2016. Megafauna and ecosystem function from the Pleistocene to the Anthropocene. *Proceedings of the National Academy of Sciences* 113:838–846.
- Mills, J. G., P. Weinstein, N. J. C. Gellie, L. S. Weyrich, A. J. Lowe, and M. F. Breed. 2017. Urban habitat restoration provides a human health benefit through microbiome rewilding: the Microbiome Rewilding Hypothesis. *Restoration Ecology* 25:866–872.
- Mishra, C., S. E. Wieren, I. M. A. Heitkönig, and H. H. T. Prins. 2002. A theoretical analysis of competitive exclusion in a Trans-Himalayan large-herbivore assemblage. *Animal Conservation* 5:251–258.
- Murthy, K., and S. Bagchi. 2018. Spatial patterns of long-term vegetation greening and browning are consistent across multiple scales: Implications for monitoring land degradation. *Land Degradation & Development* 29:2485–2495.
- Naidu, D. G. T., and S. Bagchi. 2021. Greening of the earth does not compensate for rising soil heterotrophic respiration under climate change. *Global Change Biology* 27:2029–2038.
- Namgail, T., S. E. van Wieren, and H. H. T. Prins. 2013. Distributional congruence of mammalian herbivores in the Trans-Himalayan Mountains. *Current Zoology* 59:116–124.
- Paluch, E. C., M. A. Thomsen, and T. J. Volk. 2013. Effects of Resident Soil Fungi and Land Use History Outweigh Those of Commercial Mycorrhizal Inocula: Testing a Restoration Strategy in Unsterilized Soil. *Restoration Ecology* 21:380–389.
- Pinheiro, J., D. Bates, S. DebRoy, D. Sarkar, and R. R Development Core Team. 2020. nlme: Linear and Nonlinear Mixed Effects Models, R package version 3.1-148.
- Reinhart, K. O., H. S. Sanni Worogo, and M. J. Rinella. 2021. Ruminating on the science of carbon ranching. *Journal of Applied Ecology*.
- Robertson, G. P., D. C. Coleman, C. S. Bledsoe, and P. Sollins. 1999. Standard soil methods for long-term ecological research. Oxford University Press, New York, NY.
- Roy, S., and S. Bagchi. 2021. Large Mammalian Herbivores and the Paradox of Soil Carbon in Grazing Ecosystems: Role of Microbial Decomposers and Their Enzymes. *Ecosystems* In press:1–13.
- Sayer, J., T. Sunderland, J. Ghazoul, J.-L. Pfund, D. Sheil, E. Meijaard, M. Venter, A. K. Boedhihartono, M. Day, C. Garcia, C. van Oosten, and L. E. Buck. 2013. Ten principles for a landscape approach to reconciling agriculture, conservation, and other competing land uses. *Proceedings of the National Academy of Sciences* 110:8349–8356.
- Schmitz, O. J., P. A. Raymond, J. A. Estes, W. A. Kurz, G. W. Holtgrieve, M. E. Ritchie, D. E. Schindler, A. C. Spivak, R. W. Wilson, M. A. Bradford, V. Christensen, L. Deegan, V. Smetacek, M. J. Vanni, and C. C. Wilmers. 2014. Animating the Carbon Cycle. *Ecosystems* 17:344–359.
- Schowanek, S. D., M. Davis, E. J. Lundgren, O. Middleton, J. Rowan, R. Ø. Pedersen, D. Ramp, C. J. Sandom, and J. Svenning. 2021. Reintroducing extirpated herbivores could partially reverse the late Quaternary decline of large and grazing species. *Global Ecology and Biogeography* 30:896–908.

- Shipley, B. 2009. Confirmatory path analysis in a generalized multilevel context. *Ecology* 90:363–368.
- Singh, R., R. K. Sharma, and S. Babu. 2015. Pastoralism in transition: Livestock abundance and herd composition in Spiti, Trans-Himalaya. *Human Ecology* 43:799–810.
- Sinsabaugh, R. L., S. Manzoni, D. L. Moorhead, and A. Richter. 2013. Carbon use efficiency of microbial communities: stoichiometry, methodology and modelling. *Ecology Letters* 16:930–939.
- Sinsabaugh, R. L., B. L. Turner, J. M. Talbot, B. G. Waring, J. S. Powers, C. R. Kuske, D. L. Moorhead, and J. J. Follstad. Shah. 2016. Stoichiometry of microbial carbon use efficiency in soils. *Ecological Monographs* 86:172–189.
- Sitters, J., D. M. Kimuyu, T. P. Young, P. Claeys, and H. Olde Venterink. 2020. Negative effects of cattle on soil carbon and nutrient pools reversed by megaherbivores. *Nature Sustainability* 3:360–366.
- Six, J., S. D. Frey, R. K. Thiet, and K. M. Batten. 2006. Bacterial and Fungal Contributions to Carbon Sequestration in Agroecosystems. *Soil Science Society of America Journal* 70:555–569.
- Smith, P. 2014. Do grasslands act as a perpetual sink for carbon? *Global Change Biology* 20:2708–2711.
- Thornton, P. K. 2010. Livestock production: recent trends, future prospects. *Philosophical Transactions of the Royal Society B: Biological Sciences* 365:2853–2867.
- du Toit, J. T., R. Kock, and J. C. Deutsch. 2009. *Wild Rangelands: Conserving Wildlife While Maintaining Livestock in Semi-Arid Ecosystems*. John Wiley and Sons Ltd.
- Tredennick, A. T., G. Hooker, S. P. Ellner, and P. B. Adler. 2021. A practical guide to selecting models for exploration, inference, and prediction in ecology. *Ecology* 102:e03336.
- Veblen, K. E., L. M. Porensky, C. Riginos, and T. P. Young. 2016. Are cattle surrogate wildlife? Savanna plant community composition explained by total herbivory more than herbivore type. *Ecological Applications* 26:1610–1623.
- Wepking, C., B. Badgley, J. E. Barrett, K. F. Knowlton, J. M. Lucas, K. J. Minick, P. P. Ray, S. E. Shawver, and M. S. Strickland. 2019. Prolonged exposure to manure from livestock-administered antibiotics decreases ecosystem carbon-use efficiency and alters nitrogen cycling. *Ecology Letters* 22:2067–2076.
- Western, D., P. Tyrrell, P. Brehony, S. Russell, G. Western, and J. Kamanga. 2020. Conservation from the inside-out: Winning space and a place for wildlife in working landscapes. *People and Nature* 2:279–291.
- Willemen, L., N. N. Barger, B. ten Brink, M. Cantele, B. F. N. Erasmus, J. L. Fisher, T. Gardner, T. G. Holland, F. Kohler, J. S. Kotiaho, G. P. von Maltitz, G. Nangendo, R. Pandit, J. A. Parrotta, M. D. Potts, S. D. Prince, M. Sankaran, A. Brainich, L. Montanarella, and R. Scholes. 2020. How to halt the global decline of lands. *Nature Sustainability* 3:164–166.
- Witt, G. B., M. V. Noël, M. I. Bird, R. J. S. Beeton, and N. W. Menzies. 2011. Carbon sequestration and biodiversity restoration potential of semi-arid mulga lands of Australia interpreted from long-term grazing exclosures. *Agriculture, Ecosystems and Environment* 141:108–118.
- Wubs, E. R. J., W. H. Van Der Putten, M. Bosch, and T. M. Bezemer. 2016. Soil inoculation steers restoration of terrestrial Ecosystems. *Nature Plants* 2:1–5.
- Zhang, H., and G. Fu. 2021. Responses of plant, soil bacterial and fungal communities to grazing vary with pasture seasons and grassland types, northern Tibet. *Land Degradation & Development* 32:1821–1832.

Figures

Fig. 1

Figure 1. Trait space of livestock and native herbivores in Spiti, India. We find there is considerable overlap in multidimensional trait space between native herbivores (green) and livestock (blue). Native herbivores in the study area are ibex, bharal, yak; asterisk (kiang, argali) indicates that these animals are found in nearby regions. The livestock consist of goat, sheep, donkey, horse, cattle and yak-cattle hybrids. Six traits were used to construct the trait-space: body mass, dietary guild, fermentation type, sexual-dimorphism, graminoid, and limb morphology (Lundgren et al. 2020) with Principal Coordinates Analysis on the Gower's distance matrix.

599 Fig. 2

600

601

602 Figure 2. Long-term change in soil-C (mean $\pm$ 95CI) under grazing by native herbivores and
603 livestock in Spiti, India. Watersheds with livestock had overall lower soil-C than those with
604 native herbivores, but both gained soil-C during the decade at approximately 1%.

605

606

Fig. 3

Figure 3. Grazer-effect ($\Delta = \ln\left(\frac{\text{grazed}}{\text{fenced}}\right)$), as mean $\pm$ 95CI of livestock and native herbivores on twelve soil microbial variables in Spiti, India. These variables are basal microbial respiration (BR), potential microbial respiration (SIR), Fungal fraction (F), ITS abundance (ITS), Bacterial fraction (B), 16s gene abundance (16s), Microbial Biomass Carbon (MBC), Microbial Biomass Nitrogen (MBN), Labile-C (LC), Labile-N (LN), Metabolic quotient ($q\text{CO}_2$), and microbial carbon use efficiency (CUE). Direction and magnitude of differences in BR, SIR, F, ITS abundance, and CUE varied with herbivore-assemblage type. But the remaining variables were not influenced by herbivore-assemblage type.

Fig. 4

Figure 4. Mixed-effects structural equation models to evaluate two competing hypotheses – with and without herbivore assemblage-type as an explanatory variable, i.e., SEM_A and SEM₀. For each variable, grazer-effect ($\Delta = \ln\left(\frac{\text{grazed}}{\text{fenced}}\right)$) was calculated from grazed and ungrazed paired plots. The variables are Δ BR: basal respiration, Δ MBC: microbial biomass-C, Δ LC: labile-C, Δ LN: labile-N, Δ F: fungal biomass, Δ CUE: carbon use efficiency. In SEM_A, as herbivore-assemblage goes from native to livestock, a decrease in Δ F leads to a decrease in Δ CUE. Alternatively, in SEM₀, increase in Δ F increases Δ CUE, independent of herbivore-assemblage type. Green and red arrows represent significant positive and negative standardized path coefficients respectively, while grey arrows represent non-significant paths ($\alpha = 0.05$). Marginal and conditional R^2 are reported, and asterix represent statistical significance (***) $P \leq 0.001$, ** $P \leq 0.01$, and * $P \leq 0.05$.