Optimal resistance management for mixtures of high-risk fungicides: robustness to the initial frequency of resistance and pathogen sexual reproduction

Nick P Taylor^{*1} and Nik J Cunniffe¹

*For correspondence:

nt409@cam.ac.uk (NPT)

¹Department of Plant Sciences, University of Cambridge, Cambridge, UK

Abstract

There is a strong consensus that selection for fungicide resistant pathogen strains can be most effectively limited by using applications of mixtures of fungicides designed to balance disease control against selection. However, how to do this in practice is not entirely characterised. Previous work indicates optimal mixtures of pairs of fungicides which are both at a high risk of resistance can be constructed using pairs of doses which select equally for both single resistant strains in the first year of application. What has not been addressed thus far is the important real-world case in which the initial levels of resistance to each fungicide differ, for example because the chemicals have been available for different lengths of time. We show how recommendations based on equal selection in the first year can be sub-optimal in this case. We introduce a simple alternative approach, based on equalising the frequencies of single resistant strains in the year that achieving acceptable levels of control is predicted to become impossible. We show that this strategy is robust to changes in parameters controlling pathogen epidemiology and fungicide efficacy. We develop our recommendation using a pre-existing, parameterised

23 model of *Zymoseptoria tritici* (the pathogen causing Septoria leaf blotch on wheat), which
 24 exemplifies the range of plant pathogens which predominantly spread clonally, but for which
 25 sexual reproduction forms an important component of the life cycle. We show that pathogen
 26 sexual reproduction can influence the rate at which fungicide resistance develops, but does not
 27 qualitatively affect our optimal resistance management recommendation.

28

29 Introduction

30 World food security faces multiple threats, including the growing global population (*Godfray et al.,*
 31 *2010*), climate change (*Tai et al., 2014*) and plant disease (*Strange and Scott, 2005*). However, it is
 32 estimated that food production will need to increase by 60% by 2050 (*Ristaino et al., 2021*). Despite
 33 annual spending of roughly 16 billion US dollars on fungicides globally, estimated crop losses due to
 34 disease stand at 20% (*Jorgensen et al., 2017*). Fungicide resistance challenges our ability to maintain
 35 control of fungal pathogens, but effective resistance management strategies prolong control of
 36 these yield-limiting crop diseases and have been studied for decades (*Staub, 1991*). We explore the
 37 optimal management of fungicide resistance in crop pathogens using mixtures containing pairs
 38 of fungicides which are ‘high-risk’ for resistance. In particular, we explore the effects of sexual
 39 pathogen reproduction and variation in initial levels of resistance.

40 We use Septoria leaf blotch (*Zymoseptoria tritici*), the most prevalent disease of wheat world-
 41 wide (*Suffert et al., 2011*), as our case study. An estimated 70% (≈€1bn) of the European cereal
 42 fungicide market is primarily targeted towards the management of *Z. tritici* of winter wheat (*Tor-*
 43 *riani et al., 2015*). Most fungicide modelling studies focus on Septoria, so there are existing pa-
 44 rameterised models available (e.g. *Hobbelen et al. (2013); Elderfield et al. (2018)*). Further, it is a
 45 heterothallic fungus (*Suffert et al., 2016*) capable of both sexual and asexual reproduction (*Suffert*
 46 *et al., 2011; Eriksen et al., 2001; Singh et al., 2021*). It therefore exemplifies the large number of
 47 plant pathogens for which sexual reproduction is potentially important in the epidemiology (*Agrios,*
 48 *2004*) and evolution (*McDonald et al., 1996*). For Septoria, reported proportions of sexual repro-
 49 duction differ widely between experiments (*Chen and McDonald, 1996; Zhan et al., 1998*), but it is
 50 known that ascospores produced via sexual reproduction initiate Septoria epidemics within a field
 51 (*Shaw and Royle, 1989*). Although ascospores are quantitatively the most significant form of pri-

mary inoculum (*Suffert et al., 2011*), most fungicide resistance modelling studies do not consider pathogen sexual reproduction. Here we seek to understand the effect of the inclusion of pathogen sexual reproduction on the resulting resistance management recommendation. We neglect the effect of ascospores within the growing season, due to previous studies suggesting they have a small effect on the severity of an epidemic (*Eriksen et al., 2001*), in part caused by the longer latent period of the sexual pseudothecia compared to the clonal pycnidia. Optimal management principles for *Septoria* may transfer to other fungal and oomycete crop pathogens which reproduce sexually, e.g. *Phytophthora infestans*, cause of the potato disease late blight (*Fones et al., 2020*).

A common resistance management strategy is to use fungicide mixtures with more than one mode of action present in the mixture. These fungicides are often categorised as ‘low-risk’ or ‘high-risk’ for resistance, depending on whether resistant pathogen strains exist in the population and the likelihood of developing resistance, amongst other factors (*Brent and Hollomon, 2007*). Often fungicide mixtures contain two fungicides that are high-risk for development of resistance. These mixtures are of increasing relevance since there are few low-risk fungicides available and the high risk options are typically of higher efficacy (*van den Bosch et al., 2014b*). Further, low-risk (multi-site) fungicides are increasingly rare; for example chlorothalonil has been banned for use by the EU due to environmental concerns since 2019 (*Murray, 2019*). Previous modelling studies have found fungicide mixtures to be more effective as a resistance management strategy than alternating use of fungicides (*Elderfield et al., 2018*) or spatially concurrent applications (*Hobbelen et al., 2013*), where different fields receive treatments from different modes of action. That mixtures outperformed alternations or concurrent use was robust to fitness costs, partial resistance, changes in fungicide parameters and the initial frequency of the double resistant strain. For this reason we concentrate exclusively here on the optimal strategy for mixtures of two high-risk fungicides and seek to test how to optimally construct high-risk fungicide mixtures if current levels of resistance to the two mixing partners differ, or if between-season pathogen sexual reproduction is considered.

It was reported by *van den Bosch et al. (2014b)* that, across 17 publications, mixtures of high-risk fungicides resulted in a reduction in selection for resistance in 20 out of 24 pathogen-crop fungicide combinations. There is ongoing debate about how high-risk mixtures (i.e. mixtures of high risk fungicides) should be constructed. Although the so-called ‘governing principles’ (*van den Bosch et al., 2014a*) suggest that increased fungicide dose increases selection for a given mode of action, increased dose of a mixing partner can reduce selection for the other mode of action in the

83 mixture (*van den Bosch et al., 2014b*). Modelling work shows that the optimal way to mix a low-
84 risk and a high-risk fungicide is to use the maximum dose of the low-risk chemical and the minimal
85 viable dose of the high-risk chemical (*Hobbelen et al., 2011a; Elderfield et al., 2018*). However,
86 maximising the dose of either fungicide when the mixture contains two high-risk chemicals could
87 lead to excessive selection pressure on that fungicide, so different recommendations are required.

88 *Hobbelen et al. (2013)* use modelling to address the case where the fungicide mixture contains
89 two high-risk chemicals. They consider four pathogen strains: one that is resistant to both chemi-
90 cals; one that is sensitive to both; and two more that are sensitive to one fungicide but resistant to
91 the other. Their results suggest that the choice of doses used is crucial to the resulting durability of
92 the strategy. *Hobbelen et al. (2013)* suggest the optimal fungicide mixture has a dose pairing that
93 is as weak as possible whilst achieving sufficient yield, and selects equally for both single resistant
94 strains. These authors addressed the case where both single resistant strains are initially at the
95 same frequency, and explored what happens for different amounts of the double resistant strain.
96 However, that study did not address the common real-world scenario where the initial levels of
97 resistance to the two chemicals differ. The optimal strategy in this case is not yet described and –
98 as well as the effect of sexual reproduction – is the focus of this paper.

99 Initial levels of resistance commonly differ for different fungicides due to differing natural inci-
100 dences of resistant strains, or because one fungicide was introduced to market much earlier than
101 its mixing partner. For instance, resistance to benzimidazoles developed rapidly in the mid-1980s
102 (*Blake et al., 2018*), but resistance to Quinone outside Inhibitors (QoI) fungicides was not detected
103 in the UK until 2001 (*Cheval et al., 2017*). Benzimidazoles were introduced in the 1960s, but QoIs
104 were not introduced until the 1990s (*Leadbeater, 2014*), suggesting levels of resistance and rate of
105 increase of resistance differed greatly between these fungicide classes. Fungicide sensitivity has
106 been reported to differ for high-risk methyl benzimidazole carbamate (MBC) fungicides depending
107 on year and region, with sensitive proportions of *Oculimacula acuformis* (eyespot disease of cere-
108 als) comprising 92% in Germany in 1985, but only 4% and 16% in France and the UK respectively in
109 the same year (*Parnell et al., 2008*). Differing resistance frequencies in *Botrytis cinerea* (gray mould
110 of raspberries) to seven different fungicides from a variety of fungicide classes were reported in
111 Northern Germany (*Rupp et al., 2017*). Further, initial resistance frequencies may be influenced by
112 mutation-selection balance (*van den Bosch and Gilligan, 2008*) which depends on the fitness costs
113 of resistance, which will depend on the mutation, and hence the fungicide.

Fungicides can be described by ‘dose-response curves’ which are measures of their efficacy. These curves differ depending on the mode of action and effectiveness of each chemical. We seek to show how different dose-response parameters influence the optimal strategy even when levels of resistance to the two fungicides vary and/or pathogen sexual reproduction is present. Understanding the effect of different dose response curves on the optimal strategy is crucial to decision making when constructing mixtures containing pairs of existing fungicides and/or new chemicals that come on to the market. These decisions are made by agronomists and growers but typically informed by recommendations from the Fungicide Resistance Action Committee (FRAC).

In this paper we address the following questions:

1. What is the effect of varying initial levels of resistance on optimal resistance management strategies for mixtures of pairs of high risk fungicides?
2. When do existing strategy recommendations fail, and how can we improve upon them?
3. How robust is our new recommendation to alterations in parameters controlling pathogen epidemiology and fungicide efficacy?
4. What is the effect of the balance of between-season sexual and asexual reproduction on the model and the strategy recommendation?

Methods

The model is an adapted version of one presented by *Hobbelen et al. (2013)*, which addresses two high-risk fungicides used together to control Septoria. The model is compartment-based and measures different categories of leaf tissue (Table 1). After infection, healthy (susceptible) tissue (S) transitions to exposed tissue (E) (infected but not infectious) and then to infectious tissue (I), before removal (R). The initial infection is given by a primary inoculum (P). The model also includes growth and senescence of living tissue. The subscripts s and r denote whether the pathogen strain is sensitive or resistant to fungicide application (Figure 2). The model also tracks the active concentration of both fungicides in the mixture over time. We split the modelled year into two distinct time periods – within and between growing seasons.

We generalise the model presented by *Hobbelen et al. (2013)* by introducing between-season sexual reproduction, because Septoria’s sexual ascospores are reported to contribute to a large proportion of the primary inoculum that initialises each epidemic (*Eriksen et al., 2001; Suffert et al., 2011*). We denote the proportion p_B , and initially set $p_B = 0$ in line with *Hobbelen et al. (2013)*

144 before exploring the effect of between-season sexual reproduction by scanning over a range of
145 values. We neglect within-season reproduction both for simplicity and because previous research
146 suggests its effect on epidemic severity is small (*Eriksen et al., 2001*). Another change to the model
147 presented by *Hobbelen et al. (2013)* is that we consider one field instead of two, since we omit the
148 less effective ‘concurrent field’ strategy.

149 Pathogen strains

150 When studying fungicide resistance evolution, modellers often consider an emergence and a se-
151 lection phase separately (*van den Bosch and Gilligan, 2008; Milgroom, 1990; van den Bosch et al.,*
152 *2011*). The former concerns the initial stochastic phase where new resistant strains appear through
153 random mutation and invasion. We focus on the selection phase, which is where the pathogen
154 population changes once fungicide treatments are used and a corresponding selection pressure
155 applied.

156 We label the two fungicides *A* and *B*. It is assumed that there are four pathogen strains; the dou-
157 ble sensitive strain, two single resistant strains and the double resistant strain. These are denoted
158 by *ss*, *sr*, *rs* and *rr* respectively, where *r* indicates resistant and *s* indicates sensitive to fungicide *A* or
159 *B* (for example the labelling *rs* would correspond to a pathogen strain that is resistant to fungicide
160 *A* but sensitive to fungicide *B*).

161 Within-season

162 Primary infection

163 The modelled season starts at T_{emerge} (Table 1), which corresponds to the emergence of ‘leaf five’
164 (*Elderfield et al., 2018; van den Berg et al., 2013*), rather than the start of the growing season.
165 The dynamics between the start of the growing season and T_{emerge} are approximated by the initial
166 conditions which are used to subsume both the primary infection and the initial dynamics at the
167 start of the season (*Elderfield et al., 2018*).

168 The initial infection comes from a primary inoculum P_{mn} for strain $mn \in \{rr, rs, sr, ss\}$. This inocu-
169 lum is assumed to decay exponentially, at the same rate for all strains. In fact, aside from the effect
170 of the fungicide application, the strains are treated as identical. In particular, this assumes no fit-
171 ness cost to presence of fungicide resistance. Letting t be the time since the start of the season

(measured in degree-days) and $P_{mn,0}$ be the initial amount of inoculum for strain $mn \in \{rr, rs, sr, ss\}$:

$$P_{mn} = P_{mn,0} \exp(-\nu t). \quad (1)$$

Host growth

We define the total amount of tissue

$$A = S + E_{ss} + E_{rs} + E_{sr} + E_{rr} + I_{ss} + I_{rs} + I_{sr} + I_{rr} + R. \quad (2)$$

The growth of the wheat crop is given by the following function:

$$g(A) = r(1 - A). \quad (3)$$

The growth is monomolecular (*Cunniffe and Gilligan, 2010*), and includes density dependence such that the rate of production of host tissue decreases as the total amount of tissue (A) increases. We non-dimensionalised the tissue quantities, so that after growth finishes, each quantity represents the proportion (out of 1) of tissue of that particular type. This differs from the scale used by *Hobbelen et al. (2013)*, where tissue quantities are measured out of 4.2, the maximum leaf area index (due to vertically stacked leaves). The growth function is scaled by a growth rate r .

Senescence

We use the following senescence function Γ :

$$\Gamma(t) = \begin{cases} 0.005 \left(\frac{t - T_{GS61}}{T_{GS87} - T_{GS61}} \right) + 0.1 e^{-0.02(T_{GS87} - t)}, & \text{if } t \geq T_{GS61}, \\ 0, & \text{if } t < T_{GS61}. \end{cases} \quad (4)$$

This function is inherited from the models by *Hobbelen et al. (2013)* and *Elderfield et al. (2018)*, and represents the rapid increase in the senescence of healthy tissue towards the end of the season. Senescence begins at growth stage 61, $t = T_{GS61}$ (where we use Zadok's growth scale for the growth of wheat (*Zadoks et al., 1974*)). Senescence is assumed to only affect tissue from the S and E compartments, but the disease is assumed to disrupt this process meaning that there is no senescence of tissue in the I compartment. By harvesting time at $t = T_{GS87}$ we get (almost) complete senescence of the healthy tissue.

Effect of fungicides

We denote the response of a pathogen strain m to the application of a fungicide F by $\delta_{F,m}(t)$. The time dependency (Figure 1D) arises because the chemical concentration changes depending on

the time since the crop was sprayed. We assume fungicide applications occur instantaneously and that the concentration of fungicide decays exponentially (Figure 1C). Two-treatment fungicide programs were found to balance effective control with resistance management in *van den Berg et al. (2016)*, so we focus on strategies which have two spray applications per year. We assume the two fungicide treatments are applied at T_{GS32} and T_{GS39} each year (Table 1).

The fungicides decrease the rate of the transition of tissue from healthy to exposed (β), and from exposed to infected (γ), corresponding to both protectant and eradican activity. Both transition rates are assumed to decrease by the same amount, as in *Hobbelen et al. (2013)*. The fungicide response $\delta_{F,s}$ lies in the interval $[0, 1]$, and multiplies each respective transition rate (Table 2).

For a dose C_F of a fungicide F , we use dose responses to sensitive strains (Figure 1A,B) of the type

$$\delta_{F,s}(C_F) = 1 - \epsilon_F(C_F), \quad (5)$$

where $\delta_{F,s}$ is the effect on any strains sensitive to it, and

$$\epsilon_F(C_F) = \omega_F \left(1 - e^{-\theta_F C_F} \right). \quad (6)$$

Here ω is the maximum effect of the fungicide and θ is a curvature parameter. Higher efficacy fungicides have higher values of ω and/or θ . We assume that resistant strains are completely unaffected by an application of fungicide, in the same way as *Elderfield et al. (2018)*. This means that $\delta_{F,r} = 1$ for any fungicide F .

Figure 1. Fungicide applications limit pathogen growth rates before concentrations decay. Dose response curves for pyraclostrobin (A) and a weaker but slower decaying alternative fungicide (B). We predominantly focus on dose responses within a similar parameter range to pyraclostrobin, a high risk strobilurin fungicide. This particular dose response curve is parameterised with a maximum efficacy $\omega = 1$, and a curvature of $\theta = 9.6$ (Hobbelen et al., 2013). We plot $\delta_{F,s}$, the multiplier on the growth rate for sensitive strains, so that the growth rate is unchanged ($\delta_{F,s} = 1$) when the chemical concentration is 0, but the growth rate approaches 0 as the concentration of pyraclostrobin increases (A). However, the weaker fungicide has a maximum efficacy $\omega = 0.8$ (B), meaning that the growth multiplier approaches 0.2 (dotted line) at high concentrations. The curvature parameter θ characterises how steeply the curve drops, i.e. how sharply the effect on the pathogen depends on the concentration of chemical. The concentration depends on time since application (C), starting at 0 at the beginning of the modelled season (1212 degree days as in Elderfield et al. (2018) and increasing instantaneously at $t = 1456$ and $t = 1700$ when the doses are applied, before exponentially decaying with rates that differ here for the two fungicides. The effect of the fungicides correspondingly vary with time since the first spray (D). We model scenarios in which two sprays are applied, and here a full dose of both fungicides is applied. **Parameter values:** pyraclostrobin: $(\omega, \theta, \Lambda) = (1, 9.6, 1.11 \times 10^{-2})$; weaker alternative fungicide: $(\omega, \theta, \Lambda) = (0.8, 7, 8 \times 10^{-3})$. Pyraclostrobin parameterisation is as in Hobbelen et al. (2013).

210 Primary and secondary infection

211 The net rate of infection for any strain mn is given by the sum of the rates of primary infections
212 and secondary infections. The rate of primary infections is given by $\delta_{A,m}\delta_{B,n}P_{mn}$, and the rate of
213 secondary infection is given by $\delta_{A,m}\delta_{B,n}I_{mn}$, so the rate of new infections of strain mn is given by:

$$G_{mn} = \delta_{A,m}\delta_{B,n}(I_{mn} + P_{mn}). \quad (7)$$

214 Initially primary infection P_{mn} contributes highly to G_{mn} , but since the primary inoculum decays
215 away exponentially it becomes less important relative to secondary infections as the season pro-
216 gresses.

217 Within-season model equations

218 We summarise the within-season dynamics as follows:

$$\frac{dS}{dt} = g - \Gamma S - \frac{\beta S}{A} \sum_{mn \in \{rr, sr, rs, ss\}} G_{mn}, \quad (8)$$

$$\frac{dE_{mn}}{dt} = \frac{\beta S}{A} G_{mn} - \Gamma E_{mn} - \gamma \delta_{A,m}\delta_{B,n}E_{mn} \quad \text{for } mn \in \{rr, sr, rs, ss\}, \quad (9)$$

$$\frac{dI_{mn}}{dt} = \gamma \delta_{A,m}\delta_{B,n}E_{mn} - \mu I_{mn} \quad \text{for } mn \in \{rr, sr, rs, ss\}, \quad (10)$$

$$\frac{dR}{dt} = \Gamma \left[S + \sum_{mn \in \{rr, sr, rs, ss\}} E_{mn} \right] + \mu \sum_{mn \in \{rr, sr, rs, ss\}} I_{mn}. \quad (11)$$

219 Note that the notation $mn \in \{rr, sr, rs, ss\}$ means that we actually have 4 equations for E_{mn} and
220 for I_{mn} , each corresponding to one of the pathogen strains rr, sr, rs, ss .

221 Between-season dynamics

222 Any remaining primary inoculum from the previous winter is assumed to have decayed entirely
223 by the end of the growing season. We assume a constant total initial amount of inoculum in each
224 season (denoted ψ_0), as is used by *Hobbelen et al. (2013)*; *Elderfield et al. (2018)*.

225 We keep the proportion of between-season sexual reproduction as a free parameter (p_B). Then
226 the remaining proportion $1 - p_B$ of the initial population is assumed to be clonal offspring. Initially
227 we consider $p_B = 0$ as in *Hobbelen et al. (2013)*; *Elderfield et al. (2018)*. We later scan over all
228 possible values of p_B to demonstrate the effect of alternative parameter choices.

229 We denote the proportion offspring of strain mn as X_{mn} , Y_{mn} for the asexual and sexual cases
230 respectively. The calculation of these quantities is described below. The levels of primary inoculum

at the start of the next season are given by:

$$P_{rr} = \psi_0 \left((1 - p_B) X_{rr} + p_B Y_{rr} \right), \quad (12)$$

$$P_{rs} = \psi_0 \left((1 - p_B) X_{rs} + p_B Y_{rs} \right), \quad (13)$$

$$P_{sr} = \psi_0 \left((1 - p_B) X_{sr} + p_B Y_{sr} \right), \quad (14)$$

$$P_{ss} = \psi_0 \left((1 - p_B) X_{ss} + p_B Y_{ss} \right). \quad (15)$$

Let I_{mn}^* be the level of infection for strain mn at the end of the previous modelled season. We assume that the fractions of each pathogen strain at the start of a season are calculated based only on the fractions of infectious tissue infected by each strain at the previous season's end, as in *Hobbelen et al. (2011a)*. We define the sum of all disease strains:

$$I_{TOT}^* = \sum_{m,n} I_{mn}^*. \quad (16)$$

We may also define F_{mn} , the frequency of strain mn as a proportion of total disease:

$$F_{rr} = \frac{I_{rr}^*}{I_{TOT}^*}; \quad F_{rs} = \frac{I_{rs}^*}{I_{TOT}^*}; \quad F_{sr} = \frac{I_{sr}^*}{I_{TOT}^*}; \quad F_{ss} = \frac{I_{ss}^*}{I_{TOT}^*}. \quad (17)$$

The proportions of asexual offspring, X_{mn} , are simply given by:

$$X_{rr} = F_{rr}; \quad X_{rs} = F_{rs}; \quad X_{sr} = F_{sr}; \quad X_{ss} = F_{ss}. \quad (18)$$

For the sexual offspring Y_{mn} , we assume perfect random mating of unlinked resistance genes in a haploid population, which leads to frequencies as in *Felsenstein (1965)*. The unlinked assumption is valid unless the loci involved are close together on the same chromosome. A full derivation of these frequencies is given in Appendix 1.

Defining the linkage disequilibrium

$$D = F_{ss} F_{rr} - F_{sr} F_{rs}, \quad (19)$$

we obtain:

$$Y_{rr} = F_{rr} - \frac{1}{2} D, \quad (20)$$

$$Y_{rs} = F_{rs} + \frac{1}{2} D, \quad (21)$$

$$Y_{sr} = F_{sr} + \frac{1}{2} D, \quad (22)$$

$$Y_{ss} = F_{ss} - \frac{1}{2} D. \quad (23)$$

Combining the results for X_{mn} and Y_{mn} leads to the following result:

$$P_{rr} = \psi_0 \left(F_{rr} - \frac{1}{2} p_B D \right), \quad (24)$$

$$P_{rs} = \psi_0 \left(F_{rs} + \frac{1}{2} p_B D \right), \quad (25)$$

$$P_{sr} = \psi_0 \left(F_{sr} + \frac{1}{2} p_B D \right), \quad (26)$$

$$P_{ss} = \psi_0 \left(F_{ss} - \frac{1}{2} p_B D \right). \quad (27)$$

Note that the sum of these values gives ψ_0 , the total initial amount of inoculum.

Measures of strategy performance

Effective life

The term *effective life* is used by *Hobbelen et al. (2013, 2011a)* to denote the number of years for which a fungicide or fungicide application strategy is effective, meaning growers achieve yields above a certain threshold, which we set at 95% of the disease-free yield (*Hobbelen et al., 2013*). We assume that a grower requires a yield above this threshold for economic reasons. We will use effective life to assess the effectiveness of the fungicide mixtures we test.

Selection ratio

Hobbelen et al. (2011b) use the *selection ratio* as a measure of how strongly a particular strategy results in selection for the resistant strain. They define the selection ratio in terms of the frequency of the resistant strain before the first spray and at the end of the growing season. We generalise this form to apply to any of our four pathogen strains, so that in any year N we use:

$$SR_{mn,N} = \Phi_{mn,N,end} / \Phi_{mn,N,start}. \quad (28)$$

Here Φ_{mn} represents the density of strain mn as a proportion of total disease, at the start or end of the season. If the selection ratio for a fungicide is greater than 1 then the pathogen strain has increased in frequency in year N .

Results

During the season, susceptible host tissue increases in density in the season before natural senescence later on (Figure 2B,C). The four pathogen strains are affected by the fungicide mixture to different extents (Figure 2D,E). This leads to selection for resistant strains and a gradual loss in yield over successive seasons due to reduced control of the pathogen (Figure 2F). The proportions

266 of the pathogen population carrying resistance to either fungicide increase over successive sea-
 267 sons (Figure 2G). This increase in resistance frequency for either fungicide (e.g. the strains rs and
 268 rr are resistant to fungicide A) is approximately logistic and is a result of the selection pressure
 269 applied. This is characterised by an initially gradual change in resistance frequency before a rapid
 270 increase.

Figure 2. Selection for resistant strains within each season leads to yield losses over time. The diagram (A) depicts the within-season dynamics. The solid lines represent transitions, the dashed lines effects (for example the amount of infectious tissue affects the rate of infection) and the dotted line represents the instantaneous arrival of primary inoculum (P) at the beginning of each season in the model. The four pathogen strains considered are denoted ss , rs , sr and rr , representing the double sensitive, two single-resistant, and the double resistant strain. S , E , I , R represent tissue categories: susceptible; exposed (latently infected); infectious (symptomatically infected); and removed. The model includes host growth, senescence, and density dependent pathogen growth (B,C,D,E). 'Leaf proportion' represents the proportion of the maximum possible leaf area index (area of leaf tissue) (Hobbelen et al., 2013; Elderfield et al., 2018). The modelled season begins at 1212 degree-days (emergence of leaf 5), as in Hobbelen et al. (2013). Note how the fungicide applications at $t = 1456$ and $t = 1700$ cause the exposed tissue (of sensitive strains) to stop increasing (D) and the infectious tissue (of sensitive strains) to decay (E). Over successive seasons, the yield declines slowly at first before a drop from years 6-8 takes it below the 95% 'unacceptable yield' threshold (F) as the resistance frequencies increase (G). The resistance frequencies for a fungicide are given by the proportion of total disease taken up by strains resistant to that fungicide (e.g. for fungicide A we consider strains rs and rr). The levels are initially very low but their rapid increase causes the loss in disease control and yield breakdown.

Parameter values: default values as used by Hobbelen et al. (2013), doses: (1, 0.5). Fungicides A and B parameterised to match efficacy of pyraclostrobin. Initial resistance frequencies: (rr , rs , sr) = (10^{-8} , 10^{-3} , 10^{-6}) – note that these are slightly higher than the default values used in later figures. The disease progress curves are for the 5th growing season. Choosing higher resistance frequencies and a later growing season means the resistant strains are sufficiently high in density to be visible on the same scale as the sensitive strain.

271 Finding an optimal high-risk mixture strategy

272 We seek the optimal dose-pairing to minimise selection for resistant strains and prolong the effec-
273 tiveness of the mixture while providing sufficient disease control to be economic. We restrict our
274 search to fixed-dose strategies; those strategies for which each year the same dose choices are
275 used, although the dose of fungicide *A* and of fungicide *B* may differ to each other in the mixture.

276 Previous work on high-risk fungicide mixtures has neglected the scenario where levels of the
277 single resistant strains differ initially. We seek to determine the optimal strategy depending on
278 these initial conditions, as well as fungicide parameterisation (dose response and decay rate) and
279 the balance of asexual and sexual pathogen reproduction. Initially we assume that there is no
280 sexual reproduction in the pathogen population, but we relax this assumption later.

281 Dose space

282 We define 'dose space' as the set of pairs of fungicide doses which take values between 0 and 1,
283 since we assume a "full dose" of 1 is the maximum dose that would be permitted (Figure 3A,B,C).
284 This allows us to describe all permissible dose-pairs for our fixed fungicide strategy. We seek the
285 optimal region within dose space, which is the region containing dose combinations which lead to
286 the longest effective life for the mixture. This is a region rather than a single pair of doses, since
287 there may be multiple dose combinations which break down in the same (optimal) year (Figures
288 3A, 4A-D).

289 The optimal region differs for different initial resistance frequencies, different fungicide param-
290 eters or different proportions of between-season sexual reproduction. To find the optimal region,
291 we use a grid of dose choices which discretises both concentrations and considers pairs of doses of
292 each chemical (labelled fungicide *A* and *B* respectively). This 'brute-force' method allows us to sim-
293 ply read off which dose combinations are best for a given model parameterisation. This method
294 was previously used by *Hobbelen et al. (2013)*, although we use a finer grid of 51 as opposed to
295 11 (i.e. for each parameterisation we consider $51^2 = 2601$ rather than $11^2 = 121$ pairs of doses).
296 We also seek a more biologically-motivated and general method to characterise the optimal dose
297 region, depending on characterisations of the pathogen and fungicides (initial levels of resistance,
298 mode of reproduction, decay rates and dose-response curves).

299 Strategies to test

300 We will compare the following candidate strategies:

- 301 • Equal selection in the first year (ESFY),
- 302 • Equal resistance frequencies at breakdown (ERFB).

303 Equal selection in the first year

304 When strains are completely resistant to fungicide *A* and *B*, **Hobbelen et al. (2013)** assert that the
 305 most durable strategy exerts an approximately equal selection pressure on both single-resistant
 306 strains. They considered situations where the frequencies of the single resistant strains were equal
 307 initially and show how equal selection in the first growing season gives the best outcome. Therefore
 308 we choose this as our first candidate strategy.

309 To explore this strategy we define a metric Ω_{SFY} :

$$\Omega_{SFY} = \frac{SR_{rs,A}}{SR_{rs,A} + SR_{sr,B}}, \quad (29)$$

310 where $SR_{mn,1}$ is the selection ratio for strain *mn* in year 1, as defined in Equation 28. Then $\Omega_{SFY} =$
 311 0.5 means there is equal selection for both single resistant strains in the first year of chemical
 312 application (Figure 3B), whereas $\Omega_{SFY} > 0.5$ means there is greater selection for fungicide *A*, and
 313 $\Omega_{SFY} < 0.5$ means there is greater selection for fungicide *B*. There is a curve in dose space defined
 314 by the doses satisfying $\Omega_{SFY} = 0.5$. In the equal dose-response, equal initial resistant frequency
 315 case, this contour is the line $y = x$ (for sufficiently large pairs of doses that we get an acceptable
 316 yield).

317 Minimal doses (that still achieve sufficient disease control) along the $\Omega_{SFY} = 0.5$ contour are
 318 equivalent to what was recommended previously (**Hobbelen et al., 2013**). We will explore this
 319 strategy (equal selection in first year, low dose), and generalise it slightly to additionally explore a
 320 strategy which takes any dose along this contour rather than only considering minimal doses (equal
 321 selection in first year). Considering all doses on the contour gives an increased chance of finding
 322 good dose combinations, and we are also interested in the relative performance of higher dose
 323 mixtures relative to lower dose mixtures. We define ‘low doses’ by considering the minimum viable
 324 and maximum permitted so-called ‘dose sum’ along this contour (we define dose sum as dose of
 325 fungicide *A* + dose of fungicide *B*). Low doses are within the lowest third (arbitrarily chosen) of
 326 dose sums which give acceptable control, a range which depends on the parameterisation.

327 If the initial frequencies of the resistant strains differ – for example for pairs of fungicides that
328 were introduced to market at different times – the ESFY recommendation sometimes fails to give
329 the optimal outcome (Figures 3A, 4B,D).

330 Equal resistance frequencies at breakdown

331 We propose an alternative strategy – one that ensures resistance frequencies are equal in the first
332 year where the yield becomes unacceptable (the ‘breakdown year’). This takes into account the
333 effect of the strategy over its entire course rather than only its first year, and it more effectively
334 accounts for differing initial levels of resistance to the two fungicides. This strategy performs as
335 well or better when compared to the ESFY strategy, using the default parameterisation and a range
336 of initial conditions (Figure 3A,C,D).

337 To explore this strategy, we will define another quantity: Δ_{RFB} . This is defined in terms of the
338 difference between the (logits of the) single resistance frequencies at breakdown:

$$\Delta_{RFB} = \text{logit}(I_{rs}) - \text{logit}(I_{sr}), \quad (30)$$

339 where

$$\text{logit}(x) = \log_{10}\left(\frac{x}{1-x}\right), \quad (31)$$

340 and I_{rs} , I_{sr} are the densities of single resistant strains to fungicides *A* and *B* respectively, measured
341 at the end of the breakdown year. This quantity informs us about the state of the system in the
342 breakdown year, and whether our strategy led to a greater degree of resistance to one fungicide
343 compared to its mixing partner. The quantities Ω_{SFY} and Δ_{RFB} are conceptually similar, focusing
344 on the effect of the strategy on the single resistant strains. However Δ_{RFB} relates to *resistance*
345 *frequency* in the *breakdown* year as opposed to *selection* in the *first* year (Ω_{SFY}). We use a different
346 function involving $\text{logit}(I_{mn})$ for Δ_{RFB} because the densities of the single resistant strains can be
347 as small as 10^{-10} , so logarithmic scales are more appropriate, whereas because the selection is
348 calculated in terms of a ratio, it typically lies between 1 and 10.

349 If the resistant frequencies are equal in the breakdown year then $\Delta_{RFB} = 0$. If there is more
350 resistance to fungicide *A* than *B*, Δ_{RFB} is positive and if there is more resistance to fungicide *B*
351 than *A* then Δ_{RFB} is negative. There is a contour in dose space described by $\Delta_{RFB} = 0$ (Figure 3A,C).
352 We will refer to this as the Δ_{RFB} contour.

353 For the identical fungicide pair model parameterisation, some points along the Δ_{RFB} contour lie
354 in the optimal region (Figure 3A). For these points, resistance frequencies in the breakdown year

are equal and our strategy is optimal. Note that there are also points along the Δ_{RFB} contour that are not optimal – this is because the strength of the mixture must be carefully chosen according to the fungicide/pathogen parameters. This process is examined in the wider parameter scan below. Note that in Figure 3D the optimal mixture strength is slightly higher than the weakest possible mixture strength. This is because the weakest possible mixture strength gives a yield in the first year of 95%. Any small increase in resistance then causes strategy failure. A slightly higher mixture strength achieves sufficiently good control to withstand these small changes in resistance in the early years, but much higher mixture strengths lead to stronger selection for resistant strains and hence much poorer resistance management. This trade-off causes the non-monotonic shape of the ERFB line in Figure 3D, which shows how effective life varies with mixture strength (parameterised as dose sum). Although *Hobbelen et al. (2013)* suggest that the optimal choice of dose is just high enough to provide effective control, this result suggests doses slightly higher than minimal are optimal. This result may have been obscured by the coarser 11 by 11 grid in dose space in that work.

If initial resistance frequencies and fungicide parameterisations are the same, this recommendation is equivalent to equal selection in the first year as found in *Hobbelen et al. (2013)* (Figure 4A). However, the ESFY recommendation may not work if the initial resistance frequencies and fungicide parameterisations are not identical. If the initial resistance levels differ then ERFB may outperform ESFY (Figure 4B,D). However, we note that if the fungicide parameterisations differ but the initial resistance levels are the same, then the two strategies are very closely aligned (Figure 4C). This is because if selection is equal every year then resistance frequencies will be equal at breakdown. Density dependent effects mean that equal selection in the first year does not guarantee equal selection in subsequent years, but this effect is small enough to be ignored in many cases.

Figure 3. The equal single resistant frequencies at breakdown strategy is optimal in some cases where equal selection in first year is sub-optimal. A, B, C: 'Dose space' for the scenario where the two fungicides are assumed to act with equal efficacy, but the initial levels of resistance are unequal, with greater levels of resistance to fungicide B initially. The points in dose space represent all of the possible choices of chemical concentrations that a grower could make. Different choices lead to different effective lives (the number of years for which this strategy gives acceptable yields). There is a grey region for which the mixture isn't sufficiently strong to give acceptable yields even in the first year. The optimal region is the green region with the highest effective life of 12 years (A). The position and size of the optimal region depend on the initial resistance frequencies and the fungicide parameters. Examples of the best doses along each contour are denoted by stars, and the highest permissible doses (strongest mixture) are denoted by circles. The dotted blue line represents those doses for which selection is equal after the first year of treatment (the ESFY strategy). The contour falls approximately along the line $y = x$ for the model parameterisation shown here, because the chemicals have equal efficacy and in the first year the initial resistance frequencies are so low that density dependent effects have a negligible effect on selection. There are no doses along this contour which lie in the optimal region of dose space. This is an example of a case in which the recommendation given by *Hobbelen et al. (2013)* fails. The values of Ω_{SFY} in dose space are shown in (B). The dashed black line represents those doses which lead to equal resistance frequencies for the two fungicides in the breakdown year (the ERFB strategy). Note that there are doses along this contour which fall in the optimal region (A). The Δ_{RFB} contour in dose space is shown in (C). This strategy reacts much more to differing initial levels of resistance. The curve is smooth within each effective life region, but jumps as it moves into regions with a different effective life (so different breakdown year). This is because the values of Δ_{RFB} are found using the relative amounts of single resistant strains in the breakdown year; in each different region the breakdown year differs and the relative amounts do not align exactly in different seasons. The effective lives are shown against different values of 'dose sum' which is the sum of the doses of fungicide A and B, as we move along the $\Omega_{SFY} = 0$ and $\Delta_{RFB} = 0$ contours (D). We find that low doses (just above minimal) along the Δ_{RFB} contour are optimal. **Parameter values:** both fungicides with dose response curves and decay rate parameter as per pyraclostrobin (*Hobbelen et al., 2013*): $(\theta, \omega, \Lambda) = (9.6, 1, 0.0111)$, initial resistance frequencies: $(I_{rs}, I_{sr}, I_{rr}) = (10^{-7}, 10^{-3}, 1.0010009487970706 \times 10^{-10})$. *Hobbelen et al. (2013)* use default initial values of 10^{-5} for the single resistant strains - we chose values 10^2 bigger/smaller than these values.

Figure 4. When does equal resistance at breakdown outperform equal selection in the first year? Dose space in four different scenarios. When the fungicides have identical dose responses, and the initial (single) resistance frequencies are the same, the ERFB and ESFY strategies are equivalent, and both are optimal (A). When the initial frequencies differ, but the fungicide parameterisations are the same, the strategies differ (B). This is demonstrated by the difference in location of the ERFB and ESFY contours. In this case ERFB outperforms ESFY; the ERFB contour passes through the optimal region but the ESFY contour does not. When the fungicide parameterisations differ, but initial (single) resistance frequencies are the same, the ERFB and ESFY strategies are very closely aligned, and both optimal (C). When both fungicide parameters and initial (single) resistance frequencies differ, again the strategies differ, but ERFB outperforms ESFY (D).

Parameter values:

A,B: Both fungicides with dose response curves and decay rate parameter as per pyraclostrobin (Hobbelen et al., 2013): $(\theta, \omega, \Lambda) = (9.6, 1, 0.0111)$.

C: fungicide A as per pyraclostrobin, fungicide B: $(\theta, \omega, \Lambda) = (7.5, 0.95, 0.0111)$.

D: fungicide A as per pyraclostrobin, fungicide B: $(\theta, \omega, \Lambda) = (7, 0.8, 0.0111)$.

A,C: Equal initial single RFs; $(I_{rs}, I_{sr}, I_{rr}) = (10^{-7}, 10^{-7}, 9.992007221626409 \times 10^{-15})$.

B,D: Different initial single RFs; $(I_{rs}, I_{sr}, I_{rr}) = (10^{-7}, 10^{-3}, 1.0010009487970706 \times 10^{-10})$.

378 The effect of sexual reproduction

379 We explored the effect of sexual reproduction for a range of initial frequencies and fungicide pa-
 380 rameterisations, which were selected randomly and independently (Table 3). We tested the maxi-
 381 mum effective life from a grid of 21 by 21 doses for different values of p_B for each set of parameters.
 382 Three such examples which demonstrate the possible behaviours are shown in Figure 5.

383 Initially we explored the initial value of the double resistant strain we expect at linkage equilib-
 384 rium; meaning the initial value satisfied $I_{rr}I_{ss} = I_{sr}I_{rs}$. These are the frequencies we would find in
 385 the absence of selection and if frequencies were combined at random. We also demonstrate the
 386 effect of increasing p_B when I_{rr} was not at linkage equilibrium, which could occur due to stochas-
 387 ticity or due to selection within a season, in the scenario where between-season reproduction was
 388 not entirely sexual. We tested two alternative scenarios: (i) low double resistant initial density (ii)
 389 high double resistant initial density. These were set to be 10^{-4} times lower/ 10^4 times higher than
 390 linkage equilibrium respectively. Although these are exaggerated values compared to the varia-
 391 tion from linkage equilibrium which might be expected in practice, they demonstrate the way the
 392 between-season reproduction influences the model output (Figure 5).

393 When the initial proportion of double resistant was equal or greater than the value expected
 394 from linkage equilibrium, higher proportions of sexual reproduction slowed the increase in the
 395 double resistant strain (Figure 5B,C). This is because the sexual reproduction step reduces the
 396 amount of the double resistant if there is an increased proportion of double resistant strain relative
 397 to single resistant strains. This is clear from the equations for between-season reproduction:

$$P_{rr} = F_{rr} - \frac{1}{2}p_B(F_{rr}F_{ss} - F_{sr}F_{rs}) = F_{rr} - \frac{1}{2}p_B D, \quad (32)$$

398 which shows that if $D = F_{rr}F_{ss} - F_{sr}F_{rs} > 0$, then the sexual reproduction step will suppress the
 399 double resistant strain.

400 When there is no effect on effective life as p_B varies (Figure 5A,B), it is because the suppression
 401 of the double resistant strain is offset by an increase in one or both of the single resistant strains
 402 (Appendix 2). Whether the single resistant strains are sufficient to lead to strategy failure depends
 403 on whether the fungicides in the mixture are low efficacy, so that each mixing partner cannot
 404 adequately control the single resistant strain which is sensitive to it.

405 Interestingly, for some cases where the double resistant strain was much lower than linkage
 406 equilibrium, there was a non-monotonic response in the maximum effective life as p_B increased

407 (Figure 5A). This is because any amount of sexual reproduction greatly increases the level of the
408 double resistant strain after the first between-season sexual reproduction step. After this, higher
409 levels of sexual reproduction tend to slow the increase in the double resistant strain that was
410 caused by selection, as above. This behaviour is explored in Appendix 2.

Figure 5. Effect of sexual reproduction proportion depends on the initial level of the double resistant strain. We examine three randomly generated scenarios (Table 3). For each scenario, we consider three possibilities: that the double resistant (rr) strain is lower than expected, relative to the equilibrium value of a sexually reproducing population in the absence of selection (**A**); that the rr strain is as expected (**B**); that the rr strain is higher than expected (**C**). We compare the maximum effective life from a 21 by 21 grid of doses for 11 different values of p_B between 0 and 1. To avoid the lines overlapping we shifted each scenario vertically by a small amount, but each effective life should be an integer. For low initial incidences of the double resistant (rr) strain, between-season sexual reproduction sometimes acts to decrease the effective life (e.g. scenarios 2 and 3). In scenario 2 there is a non-monotonic response. For moderate/expected initial incidences of the rr strain, there is often a monotonic increase in effective life with increasing proportions of sexual reproduction (e.g. scenarios 2, 3). However, sometimes there is no change in the maximum effective life in the grid with and without sexual reproduction (e.g. scenario 1). For high initial incidences, the effective life often increases when sexual reproduction is present (as in all scenarios here). This is because sexual reproduction acts to decrease the relative amount of the double resistant strain (see Appendix 2). As the level of double resistant strain increases, the effective life observed tends to decrease. For example in Scenario 2 (green) it decreases from 16 (**A**) to 8 (**C**) when $p_B = 0$.

Parameter values: random fungicide parameter values and single resistant frequencies, generated from distributions described in Table 3. **A:** initial double resistant strain (rr) density 10^{-4} times lower than value from linkage equilibrium (VLE); **B:** initial rr density at VLE; **C:** initial rr density 10^4 times higher than VLE. Values: ($p_{rr}, p_{rs}, p_{sr}, \Lambda_A, \Lambda_B, \omega_A, \omega_B, \theta_A, \theta_B$) = **Scenario 1:** ($7.049916 \times [10^{-19}/10^{-15}/10^{-11}]$, 5.545997×10^{-7} , 1.269711×10^{-8} , 0.009758 , 0.022030 , 0.729797 , 0.661193 , 7.362942 , 6.642679); **Scenario 2:** ($1.085892 \times [10^{-15}/10^{-11}/10^{-7}]$, 1.596583×10^{-6} , 6.801290×10^{-6} , 0.007417 , 0.009834 , 0.574543 , 0.706497 , 11.143576 , 11.170345); **Scenario 3:** ($2.445955 \times [10^{-15}/10^{-11}/10^{-7}]$, 7.155682×10^{-5} , 3.417953×10^{-7} , 0.005803 , 0.006279 , 0.875035 , 0.717337 , 8.544356 , 11.404773).

411 Previously mixtures were found to outperform alternations in the absence of sexual reproduc-
412 tion (Elderfield et al., 2018; Hobbelen et al., 2013). When between-season reproduction is included,
413 mixtures still outperformed alternations in every case across 100 tested randomly generated sce-
414 narios (Appendix 3).

415 **Generalising to different pathogen and fungicide parameterisations**

416 We tested the robustness of the observation that there are optimal dose combinations along the
 417 Δ_{RFB} contour, and that the ERFB strategy outperforms the ESFY, by testing a wide range of possible
 418 parameter values. We performed a randomisation scan test across different pathogen and fungi-
 419 cide parameter values to check whether this recommendation applies in different scenarios, includ-
 420 ing varying levels of between-season sexual reproduction, initial levels of resistance, and fungicide
 421 dose-responses and decay rates (Table 4). We have seen that the introduction of between-season
 422 reproduction can have an effect on the maximum effective life (Figure 5), and seek to show that
 423 the ERFB strategy is still optimal despite this effect and is robust to different initial conditions and
 424 fungicide parameter values.

425 **Pathogen parameters**

426 The model uses a proportion of between-season sexual reproduction p_B . In this section we allow
 427 p_B to vary between 0 and 1, since there is no agreed-upon experimental value that we could use.

428 To characterise the pathogen population with respect to the two fungicides, we allow the initial
 429 level of resistant strains to also vary. We independently choose values for the proportion of the
 430 population taken up by the two single resistant strains and for the double resistant strain. The
 431 remaining proportion of the population is the double sensitive strain.

432 **Fungicide parameters**

433 The fungicides in the model are characterised by three parameters: the decay rate of the chemical;
 434 the curvature parameter; and the asymptote of the dose-response curve. Fungicides with a higher
 435 curvature or asymptote value are more efficacious. Longer decay rates also increase the efficacy
 436 of a fungicide. We vary all three parameters for each fungicide in the scan.

437 **Parameter scan process**

438 For each simulation in the ensemble, we sample randomly from independent distributions for
 439 each parameter (Table 4). We repeated 500 times, giving 500 scenarios with different fungicide
 440 and pathogen characteristics as described above.

441 For each parameter combination, we ran the model over a 51×51 grid of dose pairings. We
 442 found the ERFB and ESFY contours in dose space, and compared the optimal effective life from
 443 doses on these contours to the optimal value from the grid. The ERFB and ESFY contours are not

444 constrained to lie on the set of pairs of doses we used for our grid, and using only these values
 445 would not necessarily lead to a smooth curve in dose space. Instead we characterised points on
 446 each contour in terms of the total dose applied, which approximately relates to the strength of the
 447 mixture. In particular, we identified a set of lines in dose space along which $C_B = C_{DS} - C_A$, where
 448 C_A, C_B are the concentrations of fungicide *A* and *B*, and C_{DS} is the so-called ‘dose sum’ (i.e. dose
 449 of fungicide *A* + dose of fungicide *B*). For each such line we used a numerical optimiser to find
 450 the single pair of doses which minimised the squared deviation from the relevant value (0 or 0.5
 451 for Δ_{RFB} and Ω_{SFY} respectively). The optimiser used was ‘minimize’ from the open-source Python
 452 package ‘scipy.optimize’.

453 This process finds this point precisely rather than to the ± 0.02 that would be obtained using the
 454 grid. Repeating the process for a total of 100 dose sums C_{DS} identified a smoothly varying contour
 455 of pairs of doses precisely identifying the contour and avoiding the rounding error that would have
 456 been associated with ‘snapping’ to the grid of points. This means that when effective lifetimes were
 457 compared, it was possible for the tested strategies to find a higher, equal or lower optimal value
 458 compared to the grid.

459 Restriction on parameter space

460 We restricted the parameter scan to only those fungicides for which equal selection in first year and
 461 equal resistance at breakdown was possible. For certain particularly weak fungicides, the minimum
 462 dose of the mixing partner (required to attain acceptable yield) was sufficiently large that equal
 463 selection was not possible, or if initial levels of resistance to the mixing partner were high then the
 464 mixing partner would always break down first.

465 We excluded any of these problematic parameter combinations by restricting our scan to those
 466 for which a full dose of either chemical on its own could give sufficient control ($> 95\%$) at least in
 467 the first year of application. This restriction corresponds to excluding fungicides which were not
 468 strong enough to provide adequate control if used as a solo product in the first year, and means
 469 that the corners (1,0) and (0,1) of dose space correspond to doses that give acceptable control. The
 470 restriction was sufficient to ensure that both strategies were viable in each scenario tested.

471 Parameter scan results

472 The ERFB strategy is optimal in 99% of cases (i.e. 495 out of 500). This means there is at least
 473 one point along the Δ_{RFB} contour which gives an effective life at least as good as the optimum

effective life from the grid. The ESFY strategy is less effective, performing optimally in 72% of cases. These results are summarised in Table 5. There were only 5 cases where ERFB was sub-optimal (by one year only). These cases are described in Appendix 5. Although there are many cases where the ESFY strategy performs worse than the ERFB contour, it is usually only one year away from the optimum, and is at most 5 years from the optimum. There is a single case in which the ESFY strategy outperformed the ERFB strategy by one year. This case is described in Appendix 5.

There was a relationship between the difference in initial frequencies (on a log scale) and improvement in performance from using ERFB instead of ESFY (Figure 6A). This is as expected -- the strategy which takes differences in initial frequencies into account (ERFB) more effectively performs better relative to the other strategy (ESFY) when these differences are larger. There is less of a clear pattern when looking at the initial values of the double resistant strain, although notably for very large initial values (greater than 10^{-5}) there is no difference in strategy performance, since the double resistant strain dominates and causes the loss of control (Figure 6B). This is because the improvement of ERFB on ESFY is usually caused by changes in the relative amounts of fungicide *A* and *B* in the mixture (due to the different positions of the contours). Both fungicide *A* and *B* are ineffective against the double resistant strain, so changes in the relative amounts of each does not increase the effective life if the double resistant strain is the dominant cause of loss of control. There is no clear pattern in the performance when looking at differences in the asymptote fungicide parameter (Figure 6C), which again suggests that the initial frequencies are the most important feature in terms of whether ERFB outperforms ESFY.

These results mean we can use the Δ_{RFB} contour to conceptually reduce the problem of dose selection from a two-dimensional problem (choice of two doses) to a one-dimensional problem: 'how far along the Δ_{RFB} contour is best?' In other words, we have established that single resistance frequencies should be equal in the final year, and the only remaining question is: 'how strong should the mixture be?'

On analysing the parameter scan, we find that whenever the ERFB strategy is successful, low doses are optimal (Figure 6D/Table 6). So the optimal strategy when initial resistance frequencies vary is to pick low doses along the equal resistance in breakdown year contour. These doses are often very close to minimal, but usually fractionally stronger than weakest possible acceptable mixtures. This ensures that small losses of control due to small increases in resistance are insufficient to push the strategy below the 95% threshold, as we saw in Figure 3D.

Table 1. Parameters and state variables used in the HRHR model. Sources: *Hobbelen et al. (2013)*; *Elderfield et al. (2018)*. Although the default value for the proportion of between-season sexual reproduction (p_B) is 0, we test different values to explore the effect of this parameter on the results of the model. The value of ψ_0 is changed by the non-dimensionalisation process, and the function g used by *Hobbelen et al. (2013)* was $r(k - A)$ for $k = 4.2$, not $r(1 - A)$. All other parameter values are unchanged by this process. This explains why the tissue state variables (and some of the other parameters) have no units in our version of the underlying model.

Symbol	Meaning	Type	Default value/range/equation	Units
S	Susceptible tissue	Variable	[0, 1]	-
E_{mn}	Latently infected (exposed) tissue, strain mn	Variable	[0, 1]	-
I_{mn}	Infectious tissue, strain mn	Variable	[0, 1]	-
R	Removed tissue	Variable	[0, 1]	-
A	Total tissue ($= S + R + \sum_{m,n} [E_{mn} + I_{mn}]$)	Variable	[0, 1]	-
P_{mn}	Primary inoculum, strain mn	Variable	$[0, \psi_0]$	-
C_i	Fungicide i concentration	Variable	≥ 0	Label dose (fraction of)
t	Time	Variable	≥ 0	degree-days
$g(t)$	Production of host leaf tissue	Function	Equation 3	-
$\Gamma(t)$	Senescence	Function	Equation 4	-
$\delta_{F,m}(t)$	Fungicide F growth rate factor, strain m	Function	Equation 5	-
r	Host growth rate	Parameter	1.26×10^{-2}	degree-days ⁻¹
β	Infection rate	Parameter	1.56×10^{-2}	degree-days ⁻¹
γ^{-1}	Latent period	Parameter	266	degree-days
μ^{-1}	Infectious period	Parameter	456	degree-days
ψ_0	Initial inoculum amount	Parameter	$(1.09/4.2) \times 10^{-2}$	-
ν	Inoculum decay rate	Parameter	8.5×10^{-3}	degree-days ⁻¹
p_B	Between-season sexual reproduction proportion	Parameter	0	-
Λ_F	Fungicide F decay rate	Parameter	1.11×10^{-2}	degree-days ⁻¹
θ_F	Fungicide F curvature	Parameter	9.6	Label dose ⁻¹
ω_F	Fungicide F asymptote	Parameter	1	-
T_{emerge}	Start of modelled season	Parameter	1212	degree-days
T_{GS32}	First treatment time	Parameter	1456	degree-days
T_{GS39}	Second treatment time	Parameter	1700	degree-days
T_{GS61}	Senescence begins	Parameter	2066	degree-days
T_{GS87}	End of season	Parameter	2900	degree-days

Table 2. Effect of fungicides on the pathogen strains in the model. Each value in the interval $[0, 1]$ multiplies the rate of the transition of tissue from susceptible to exposed (β) and from exposed to infectious (γ). This means that the rr strain is unaffected by the presence of fungicide (multiplied by 1), whereas the ss strain has transition rates reduced by a factor of $\delta_{A,s}\delta_{B,s} < 1$. Note that ‘fungicide effect’ and ‘realised effect’ are equivalent, since $\delta_{A,r} = \delta_{B,r} = 1$, i.e. resistant strains are assumed to be completely unaffected by an application of fungicide.

Strain	Fungicide effect	Realised effect
rr	$\delta_{A,r}\delta_{B,r}$	1
sr	$\delta_{A,s}\delta_{B,r}$	$\delta_{A,s}$
rs	$\delta_{A,r}\delta_{B,s}$	$\delta_{B,s}$
ss	$\delta_{A,s}\delta_{B,s}$	$\delta_{A,s}\delta_{B,s}$

Table 3. Ranges/values taken by parameters in the scan over values of the between-season sexual reproduction proportion. The range of fungicide decay rates depends on Λ_0 , set as 1.11×10^{-2} degree days $^{-1}$ as in Table 1. The value of the double resistant strain is fixed such that $D = 0$ initially, and then we explore what happens when the double resistant strain is at a density much lower or higher (10^{-5} or 10^5 times lower/higher) in each case (see Figure 5). Parameters were independently generated and parameter sets were accepted as long as the yield in the first year at full dose was at least 95% of the disease free yield, so that there was a valid strategy possible.

Parameter	Symbol	Range/Value	Distribution
Pathogen between-season sex proportion	p_B	$[0, 1]$	Uniform
Single resistant strain initial frequencies	$I_{mn}(0)$	$[10^{-8}, 10^{-4}]$	Log-uniform
Double resistant strain initial frequencies	$I_{rr}(0)$	Varied	Fixed by sr , rs strains
Fungicide decay rates	Λ_i	$\left[\frac{1}{3}\Lambda_0, 3\Lambda_0\right]$	Uniform
Fungicide asymptotes	ω_i	$[0.4, 1]$	Uniform
Fungicide curvatures	θ_i	$[4, 12]$	-

Table 4. This table shows the ranges/values taken by parameters in the ERFB robustness scan. Table 1 describes the other default parameter values, their sources and their units. The range of fungicide decay rates depends on Λ_0 , set as 1.11×10^{-2} degree days $^{-1}$ as in Table 1. Certain parameter combinations were excluded to ensure that both ERFB and ESFY were possible, as described in the main text.

Parameter	Symbol	Range/Value	Distribution
Pathogen between-season sex proportion	p_B	[0, 1]	Uniform
Single resistant pathogen strain initial frequencies	$I_{mn}(0)$	$[10^{-10}, 10^{-3}]$	Log-uniform
Double resistant pathogen strain initial frequencies	$I_{rr}(0)$	$[10^{-15}, 10^{-4}]$	Log-uniform
Fungicide decay rates	Λ_i	$\left[\frac{1}{3}\Lambda_0, 3\Lambda_0\right]$	Uniform
Fungicide asymptotes	ω_i	[0, 1]	Uniform
Fungicide curvatures	θ_i	[0, 12]	-

Table 5. This table shows the results of the ERFB robustness parameter scan, which tested 500 different randomly generated scenarios. ‘Cases worked’ is the number/percentage of runs in which the strategy was at least as durable as the optimal strategy from the 51×51 grid. ERFB worked in all but 5 exceptional cases which were within one year of the optimum, whereas ESFY (low dose) and ESFY (all doses) both failed to give optimal results in 28% of cases. ERFB worked every time to within one year whereas both ESFY tactics were at least two years from the optimum in over 10% of cases (12.4% for all doses, 12.6% for low doses only). The cases where ERFB was sub-optimal are addressed in Appendix 5.

Strategy	Cases worked	Cases worked (%)
Equal resistance frequency at breakdown	495	99%
Equal selection in the first year	360	72%
Equal selection in the first year, low dose	360	72%

Table 6. We split the Δ_{RFB} contour into 3 depending on the dose sum, splitting evenly into thirds between the minimum and maximum viable values on the contour. Minimum values are determined by the lowest doses to give acceptable yields, and maximum are determined by the largest dose sum such that both doses are less than or equal to 1. Low strength mixtures were optimal in 99% of cases, meaning their effective life was at least as good as the maximum value obtained on the 51×51 grid. There are 5 (i.e. 1% of cases) in which there were ERFB was not optimal (as explained in Appendix 5), so low strength mixtures were optimal in every case in which ERFB was optimal. Note that in some cases there were optimal values in more than one segment of the contour, meaning that these percentages do not sum to 100. In only 3.8% of cases did high strength mixtures perform optimally (and in these cases low doses were also optimal).

Strategy	Optimal runs	Optimal runs (%)
Low strength	495	99%
Medium strength	90	18%
High strength	19	3.8%

Figure 6. Initial resistance frequencies drive the improvement in the ERFB strategy over ESFY. **A,B,C:** we plot the difference between the effective life of the ERFB strategy and the ESFY strategy for each of the 500 scenarios in the parameter scan (Table 4). The equal resistance at breakdown strategy always produces an equal (black) or better (blue) effective life when compared to equal selection in first year (all doses) strategy. Greater differences in initial single resistance frequencies tend to correspond to an increased chance of the ERFB strategy outperforming the ESFY strategy by a greater amount (**A**). The quantity on the *x*-axis of subplot **A** is the log difference in the single resistance frequencies $|\log(I_{rs,0}) - \log(I_{sr,0})|$. If the double resistant frequency is particularly large, the strategies perform equally well, since the double resistant strain that causes strategy failure (**B**). For smaller levels of the double resistant frequency, larger differences in performance are sometimes observed. Changing the relative strengths of the fungicides does not lead to a clear pattern about the relative performance of the two strategies (**C**). The quantity on the *x*-axis, 'asymptote metric', of subplot **C** is $\omega_A/(\omega_A + \omega_B)$, where ω_F is the asymptote of the dose-response curve for a fungicide *F* (Table 1/Figure 1). There is a single case where ESFY outperforms ERFB (red), so that ERFB EL - ESFY EL is negative – this case is explored in Appendix 5.

In the examples tested in the parameter scan, low doses were equal (green) or better (purple) than high doses (**D**). The quantity on the *y*-axis of subplot **D**, 'high/low dose metric' is $EL_{low}/(EL_{low} + EL_{high})$, which is the maximum effective life offered by doses on the low or high sections of the ERFB contour, which we (arbitrarily) split into 3 depending on the dose sum between the minimum and maximum viable (acceptable yields and doses ≤ 1) values on the contour. There are a small number of doses for which high doses perform equally well, but for the vast majority of points lower doses are preferable (and in no case are high doses better).

Discussion

Optimising deployment of fungicides in order to delay spread of fungicide resistant strains remains a major challenge (*Cunniffe et al., 2015a*). Previous work has indicated that optimal mixtures of pairs of fungicides which are both at a high risk of resistance can be constructed by using pairs of doses which select equally for both single resistant strains in the first year of application (*Hobbelen et al., 2013*). However, we have shown that this recommendation can give sub-optimal results in the common real-world case in which the initial levels of resistance to the chemicals are not equal. We have presented an alternative strategy which gave optimal results essentially all of the time across a broad scan. This was an improvement when compared to the existing strategy recommendation which was optimal 72% of the time, where we tested a range of epidemiological and fungicide efficacy parameters, as well as different initial conditions for the proportion of single- and double-resistant pathogen strains. When the initial single resistant frequencies differed by at least 10^{-4} , the mean difference in effective life for the strategies was 1.01 years with a range of 0 to 4 years and the mean effective life of the equal resistance frequencies at breakdown (ERFB) strategy was 10.75 years. The average improvement in this case (as a percentage of the optimal ERFB effective life) was 7.61% with a range of 0 to 28.57%.

The strategy which consistently gave optimal results required doses which make the single resistance frequencies equal in the breakdown year. That means that the levels of resistance to each fungicide in the mixture are equal in the year that the yield becomes unacceptable due to pathogen evolution. This concept was shown to work even when fungicide parameters (asymptote, curvature and decay rate) and pathogen parameters (initial frequencies and between-season sexual reproduction proportion) were varied. The equal resistance frequency at breakdown strategy can be framed in biological terms rather than the purely mathematical framework relied on by a naive grid-search method. This leads to the following general and practically applicable recommendations:

- reduce doses of fungicides which have higher levels of existing resistance;
- lower doses are preferable to higher;
- relatively lower doses of the higher efficacy fungicide in the mixture are preferable;
- higher efficacy could be in terms of: the decay rate (slower decaying); the curvature (the effectiveness increases more rapidly with dose); or the asymptote (the maximum level of

535 control for that chemical).

536 The first point means that equal selection in the first year (ESFY) is only optimal if the initial
537 resistance levels are sufficiently close, or equal as in *Hobbelen et al. (2013)*. If they differ, the
538 mixture should be adjusted such that selection is weaker for the fungicide with an increased level of
539 existing resistance, such that the two single-resistant strains reach equal frequency as the mixture
540 becomes unable to achieve acceptable levels of control. Conversely, in the limit as initial resistance
541 to one fungicide decreases, its optimal dose increases towards full dose and the dose of the mixing
542 partner decreases to minimal levels required to give control. This matches the recommendation
543 given by *Elderfield et al. (2018)* about mixtures of low-risk and high-risk fungicides. Although the
544 idea of low doses and relatively lower doses of higher efficacy fungicides have been presented
545 before (*Hobbelen et al., 2013*), they have only previously been tested in the absence of pathogen
546 sexual reproduction. We have confirmed that these recommendations also apply when there is
547 between-season sexual reproduction.

548 The requirement that single resistance frequencies were equal at breakdown defines a curve of
549 possible dose pairings in dose space (the ERFB contour). This contour starts from a minimum viable
550 mixture strength and runs to a maximum viable strength in dose space due to legal limitations
551 on doses i.e. the point on the contour at which the dose of either fungicide first reaches 1. The
552 parameter scan showed that low doses along this contour were optimal in virtually every case.
553 The 5 exceptions are described in Appendix 5 and were based on an edge effect caused by the
554 95% yield threshold - meaning that in these cases greater control was achieved in the final year by
555 higher doses of either fungicide *A* or *B* despite slightly worse resistance management.

556 The parameter scan results can be extended to a wider range of fungicides than explicitly tested
557 in the scan. This is because there is a mathematical link between the curvature and dose param-
558 eters which means two fungicide/dose combinations are equivalent if the curvature multiplied by
559 chemical concentration is constant. This means that two fungicides with the same asymptote and
560 decay rate parameters, but different curvatures can be treated as identical with an appropriate
561 scaling of dose – see also Appendix 4). This means that the ERFB strategy is optimal across many
562 more choices of the curvature parameters than were explicitly tested in the scan.

563 Although sexual reproduction has a major role in the population genetics of *Septoria* (*Chen*
564 *and McDonald, 1996; Singh et al., 2021*), it has been omitted from previous models of fungicide

resistance (*Hobbelen et al., 2013; Elderfield et al., 2018*), justified because Septoria spreads largely clonally within season. We use the simplest model of sexual reproduction, that mating is fully random between unlinked loci and only occurs between seasons. Although loci may be linked for genes which are close together on the same chromosome, resistance of Septoria to some high-risk fungicide classes is conferred by mutations to genes on different chromosomes, leading to independent assortment. For example, resistance to methyl benzimidazoles are caused by mutations to chromosome 1 whereas resistance to succinate dehydrogenase inhibitors can be caused by mutations to chromosome 4, 7 or 8 (*Hartmann et al., 2020*). The proportion of sexual reproduction between seasons is assumed to be constant, despite evidence that sexual reproduction is influenced by density of infection (*Suffert et al., 2018; Eriksen et al., 2001*). However, we showed that the inclusion of sexual reproduction did not affect the optimal strategy recommendation for a variety of different values of the proportion of between-season sexual reproduction. Further, the model already normalises for density each year in the constant inoculum assumption which was used in previous models (*Elderfield et al., 2018; Hobbelen et al., 2013*). Although the model neglects within-season reproduction, we explored scenarios where all between-season reproduction was sexual, meaning that we have explored scenarios involving extensive pathogen sexual reproduction. Our results apply to haploid pathogens, although the situation may be more nuanced in the diploid case. The insecticide literature suggests that higher doses of insecticide mixtures can be preferable in tackling resistance in sexually reproducing, diploid pest species, particularly in combination with refugia (*Andow and Zwahlen, 2006*).

The question of mixtures vs alternations is still under debate (*Corkley et al., 2021*), but we showed that high-risk mixtures outperform alternations when between-season reproduction occurs (Appendix 3), which has previously only been shown in the case without pathogen sexual reproduction (*Hobbelen et al., 2013*).

Although these results apply to the wheat-Septoria pathosystem, they could potentially transfer to other pathosystems that are managed by mixtures of high risk fungicides. Future work may consider extending these ideas to such pathosystems e.g. grapevine powdery mildew (*Elderfield et al., 2018*) or potato blight (*Pacilly et al., 2018; Cohen and Rubin, 2020*). However, we also note that past research on mixtures containing a high risk fungicide and a low risk fungicide (*Elderfield et al., 2018*) showed a remarkable consistency of optimal recommendation across pathosystems and model structures, and we have no reason to believe results would differ for other pathosys-

596 tems.

597 The same ideas should generalise to mixtures of three or more fungicides - all modes of action
598 in the mixture would need balancing in a similar manner. However, further work would be required
599 to confirm this and to explore other strategies possible with more fungicides such as alternations
600 with three chemicals or varying pairs of two chemicals. Although growers use three-way fungicide
601 mixtures (*Phelan, 2017*), to the best of our knowledge, no modelling study exists of a mixture of
602 three or more fungicides as applied to Septoria. Three-way fungicide mixtures were also tested
603 for their control of *Phytophthora infestans* on tomato or potato and *Plasmopara viticola* on grape
604 (*Samoucha, 1987*). Combining our results with those of *Elderfield et al. (2018)* would suggest that
605 a full dose of a low-risk chemical combined with minimal doses along the ERFB contour is likely to
606 produce the longest effective life for a three-way mixture containing two high-risk fungicides and
607 one low-risk fungicide (applied to Septoria).

608 There are various possible criticisms of the model used to generate these results, such as the
609 lack of consideration of spatial effects (*Shaw, 2000; Parnell et al., 2005, 2006*), which could explore
610 how resistance might develop on a regional scale. Management decisions made by one grower
611 affect the entire community due to dispersal of aerial inoculum (*Laranjeira et al., 2020*). A spatially
612 explicit model would also allow reactive disease management to be optimised (*Cunniffe et al.,*
613 *2015b, 2016*), which would correspond to levels of fungicide treatment depending on local disease
614 intensity, or even the results of a decision support type system (*Carisse, 2010; Jorgensen et al.,*
615 *2017; Lázaro et al., 2021*).

616 Emergence of new resistant strains is not considered (*Hobbelen et al., 2014*), nor is partial re-
617 sistance (*Mikaberidze et al., 2017; Hobbelen et al., 2013*), cross-resistance (*Rehfus et al., 2018*) or
618 fitness costs (*Mikaberidze et al., 2014; Mikaberidze and McDonald, 2015*). There is no stochasticity
619 in the model, so variation within a growing season and between different growing seasons is ig-
620 nored (*te Beest et al., 2008*), as is precise timing of fungicide applications (*van den Berg et al., 2013,*
621 *2016*). Effects of host plant resistance (*Mikaberidze and McDonald, 2020; Carolan et al., 2017*) and
622 of multiple pathogens (where the crop is infected by e.g. brown rust as well as Septoria (*Garin*
623 *et al., 2018*)) are neglected. These various factors potentially have effects on the strength of mix-
624 ture required, although different relative dosages of the two fungicides could still be used to give
625 greater protection to the chemical at most urgent threat.

626 Future work might focus on exploring total lifetime yield (*Elderfield et al., 2018*) or economic

output (*te Beest et al., 2013; van den Bosch et al., 2020*) rather than effective life. The latter would involve incorporating the price of grain into the calculations, as well as the cost of each fungicide application and other costs from each year's farming. Using economics rather than effective life would likely incentivise lower doses in general, and relatively higher doses of cheaper fungicides if the difference in price relative to efficacy was significant. For instance, if a cheap high efficacy fungicide were used, lower doses might be best for effective life, but higher doses might lead to cost savings within an individual year. In a system where grain prices are high, each additional year of relatively high yield is very valuable, so resistance management becomes the most important factor. This means economics would probably only change where is optimal within the small region of dose space which contains the maximum effective life. In a system where grain prices were relatively low, then minimising doses of expensive fungicides may become a higher priority than choosing a strategy which is optimal for resistance management. These factors are important when considering policy changes to incentivise sustainable use of fungicide mixtures. Use of modelling to inform policy change on sustainable agriculture should be informed by agronomic, environmental, economic, and social considerations (*Mouratiadou et al., 2021*).

The breakdown in the final year recommendation may seem less practical than a first year recommendation, since it may be difficult to predict the future evolution of the pathogen population, leading to greater uncertainty in the recommendation. However, points in dose space that are close to the optimal ones usually also have a long effective life (Figure 4). This means that even if the estimates of the initial resistance frequencies or other parameters are imperfect, a good decision that is close to the optimal can be made. The model could be used to find the best estimate for the optimal dose-pairing given imperfect information about the levels of resistance or fungicide parameters. It can be difficult to estimate the proportion of resistance strains particularly when their incidence is very low (*van den Bosch et al., 2014a*) and there is regional variation (*Garnault et al., 2019*). However, as the resistance frequencies increase and become easier to reliably estimate, the model output could be updated with an improved estimate for the corresponding optimal dose combination. As is invariably the case, more complex disease management recommendations (such as the equal resistance at breakdown recommendation) require good prior knowledge for good outcomes (*Hyatt-Twynam et al., 2017*).

There are more practical criticisms which can be made. Growers typically apply doses in multiples of a quarter of a full dose. This means that a very precise theoretical prescription for an

optimal dose may not be used under current practices. However, our results show that pairs of fungicides should be used in such a way as to avoid resistance developing too rapidly to either component of the mixture. This may be achievable to an extent even if we restrict our attention to multiples of quarter-doses. Further, if modelling shows that using more precise doses would result in dramatic increases in the durability of a particular fungicide mixture then they should be recommended.

An interesting area for future work would be to consider time varying disease management strategies. Given that the resistant frequencies vary each year, it may be possible to prolong the effective lifetime by increasing dose as the level of resistance increases (and the reliability of estimates resistance frequency improves). Further, it would allow other strategies like alternating the use of mixtures that favour fungicide *A* or *B*, which could outperform strategies that are static in time. We hope to address these questions in future work. One approach to this would be to use optimal control theory (Bussell *et al.*, 2019; Bussell and Cuniffe, 2020, 2022), another would be to use dynamic programming (Bellman, 1952; Onstad and Rabbinge, 1985).

Availability of code online

An implementation of the model in the freely-available programming language Python (Python Software Foundation, available at <http://www.python.org>) is online at <https://github.com/nt409/HRHR>.

Acknowledgements

NPT acknowledges the Biotechnology and Biological Sciences Research Council of the United Kingdom (BBSRC) for support via a University of Cambridge DTP PhD studentship (Project Reference 2119688). The authors also thank Alexey Mikaberidze, Christopher Gilligan and Julia Davies for useful discussions.

References

- Agrios, G. (2004). *Plant pathology: fifth edition*, volume 9780080473789. Academic Press.
- Andow, D. A. and Zwahlen, C. (2006). Assessing environmental risks of transgenic plants. *Ecology Letters*, 9:196–214.
- Bellman, R. (1952). *On the Theory of Dynamic Programming*, volume 38. National Academy of Sciences.

685 Blake, J. J., Gosling, P., Fraaije, B. A., Burnett, F. J., Knight, S. M., Kildea, S., and Paveley, N. D. (2018). Changes
686 in field dose–response curves for demethylation inhibitor (DMI) and quinone outside inhibitor (QoI) fungi-
687 cides against *Zymoseptoria tritici*, related to laboratory sensitivity phenotyping and genotyping assays. *Pest*
688 *Management Science*, 74:302–313.

689 Brent, K. J. and Hollomon, D. W. (2007). *Fungicide resistance in crop pathogens: how can it be managed?* Fungicide
690 Resistance Action Committee.

691 Bussell, E. H. and Cunniffe, N. J. (2020). Applying optimal control theory to a spatial simulation model of sudden
692 oak death: ongoing surveillance protects tanoak while conserving biodiversity. *Journal of the Royal Society*
693 *Interface*, 17.

694 Bussell, E. H. and Cunniffe, N. J. (2022). Optimal strategies to protect a sub-population at risk due to an estab-
695 lished epidemic. *Journal of the Royal Society Interface*, 19.

696 Bussell, E. H., Dangerfield, C. E., Gilligan, C. A., and Cunniffe, N. J. (2019). Applying optimal control theory to
697 complex epidemiological models to inform real-world disease management. *Philosophical Transactions of*
698 *the Royal Society B*, 374.

699 Carisse, O. (2010). *Fungicides*. InTech.

700 Carolan, K., Helps, J., Van Den Berg, F., Bain, R., Paveley, N., and Van Den Bosch, F. (2017). Extending the
701 durability of cultivar resistance by limiting epidemic growth rates. *Proceedings of the Royal Society B: Biological*
702 *Sciences*, 284.

703 Chen, R. S. and McDonald, B. A. (1996). Sexual reproduction plays a major role in the genetic structure of
704 populations of the fungus *Mycosphaerella graminicola*. *Genetics*, 142:1119–1127.

705 Cheval, P., Siah, A., Bomble, M., Popper, A. D., Reignault, P., and Halama, P. (2017). Evolution of QoI resistance
706 of the wheat pathogen *Zymoseptoria tritici* in Northern France. *Crop Protection*, 92:131–133.

707 Cohen, Y. and Rubin, A. E. (2020). A new strategy for durable control of late blight in potato by a single soil
708 application of an oxathiapiprolin mixture in early season. *PLOS ONE*, 15:e0238148.

709 Corkley, I., Fraaije, B., and Hawkins, N. (2021). Fungicide resistance management: maximizing the effective life
710 of plant protection products. *Plant Pathology*.

711 Cunniffe, N. J., Cobb, R. C., Meentemeyer, R. K., Rizzo, D. M., and Gilligan, C. A. (2016). Modeling when, where,
712 and how to manage a forest epidemic, motivated by sudden oak death in California. *Proceedings of the*
713 *National Academy of Sciences of the United States of America*, 113:5640–5645.

714 Cunniffe, N. J. and Gilligan, C. A. (2010). Invasion, persistence and control in epidemic models for plant
715 pathogens: the effect of host demography. *Journal of The Royal Society Interface*, 7:439–451.

716 Cunliffe, N. J., Koskella, B., E. Metcalf, C. J., Parnell, S., Gottwald, T. R., and Gilligan, C. A. (2015a). Thirteen
717 challenges in modelling plant diseases. *Epidemics*, 10:6–10.

718 Cunliffe, N. J., Stutt, R. O., DeSimone, R. E., Gottwald, T. R., and Gilligan, C. A. (2015b). Optimising and commu-
719 nicating options for the control of invasive plant disease when there is epidemiological uncertainty. *PLOS*
720 *Computational Biology*, 11:e1004211.

721 Elderfield, J. A. D., Lopez-Ruiz, F. J., van den Bosch, F., and Cunliffe, N. J. (2018). Using epidemiological principles
722 to explain fungicide resistance management tactics: why do mixtures outperform alternations? *Phytopathol-*
723 *ogy*, 108:803–817.

724 Eriksen, L., Shaw, M. W., and Østergård, H. (2001). A model of the effect of pseudothecia on genetic recombina-
725 tion and epidemic development in populations of *Mycosphaerella graminicola*. *Phytopathology*, 91:240–248.

726 Felsenstein, J. (1965). The effect of linkage on directional selection. *Genetics*, 52:349.

727 Fones, H. N., Bebbler, D. P., Chaloner, T. M., Kay, W. T., Steinberg, G., and Gurr, S. J. (2020). Threats to global
728 food security from emerging fungal and oomycete crop pathogens. *Nature Food* 2020 1:6, 1:332–342.

729 Garin, G., Pradal, C., Fournier, C., Claessen, D., Houlès, V., and Robert, C. (2018). Modelling interaction dynamics
730 between two foliar pathogens in wheat: a multi-scale approach. *Annals of Botany*, 121:927–940.

731 Garnault, M., Duplaix, C., Leroux, P., Couleaud, G., Carpentier, F., David, O., and Walker, A.-S. (2019). Spatiotem-
732 poral dynamics of fungicide resistance in the wheat pathogen *Zymoseptoria tritici* in France. *Pest Management*
733 *Science*, 75:1794–1807.

734 Godfray, H. C. J., Crute, I. R., Haddad, L., Muir, J. F., Nisbett, N., Lawrence, D., Pretty, J., Robinson, S., Toulmin, C.,
735 and Whiteley, R. (2010). The future of the global food system.

736 Hartmann, F. E., Vonlanthen, T., Singh, N. K., McDonald, M. C., Milgate, A., and Croll, D. (2020). The complex
737 genomic basis of rapid convergent adaptation to pesticides across continents in a fungal plant pathogen.
738 *Molecular Ecology*.

739 Hobbelen, P. H., Paveley, N. D., Oliver, R. P., and van den Bosch, F. (2013). The usefulness of fungicide mixtures
740 and alternation for delaying the selection for resistance in populations of *Mycosphaerella graminicola* on
741 winter wheat: a modeling analysis. *Phytopathology*, 103:690–707.

742 Hobbelen, P. H., Paveley, N. D., and van den Bosch, F. (2011a). Delaying selection for fungicide insensitivity
743 by mixing fungicides at a low and high risk of resistance development: a modeling analysis. *Phytopathology*,
744 101:1224–1233.

745 Hobbelen, P. H., Paveley, N. D., and van den Bosch, F. (2014). The emergence of resistance to fungicides. *PLoS*
746 *ONE*, 9:91910.

747 Hobbelen, P. H. F., Paveley, N. D., Fraaije, B. A., Lucas, J. A., and van den Bosch, F. (2011b). Derivation and testing
748 of a model to predict selection for fungicide resistance. *Plant Pathology*, 60:304–313.

749 Hyatt-Twynam, S. R., Parnell, S., Stutt, R. O. J. H., Gottwald, T. R., Gilligan, C. A., and Cuniffe, N. J. (2017). Risk-
750 based management of invading plant disease. *New Phytologist*, 214:1317–1329.

751 Jorgensen, L. N., van den Bosch, F., Oliver, R. P., Heick, T. M., and Paveley, N. D. (2017). Targeting fungicide
752 inputs according to need. *Annual Review of Phytopathology*, 55:181–203.

753 Laranjeira, F. F., Silva, S. X., Murray-Watson, R. E., Soares, A. C., Santos-Filho, H. P., and Cuniffe, N. J. (2020).
754 Spatiotemporal dynamics and modelling support the case for area-wide management of citrus greasy spot
755 in a Brazilian smallholder farming region. *Plant Pathology*, 69:467–483.

756 Lázaro, E., Makowski, D., and Vicent, A. (2021). Decision support systems halve fungicide use compared to
757 calendar-based strategies without increasing disease risk. *Communications Earth & Environment* 2021 2:1,
758 2:1–10.

759 Leadbeater, A. J. (2014). Plant health management: fungicides and antibiotics. *Encyclopedia of Agriculture and*
760 *Food Systems*, pages 408–424.

761 McDonald, B. A., Mundt, C. C., and Chen, R.-S. (1996). The role of selection on the genetic structure of pathogen
762 populations: evidence from field experiments with *Mycosphaerella graminicola* on wheat. *Euphytica* 1996 92:1,
763 92:73–80.

764 Mikaberidze, A. and McDonald, B. A. (2015). Fitness cost of resistance: impact on management. *Fungicide*
765 *Resistance in Plant Pathogens*, pages 77–89.

766 Mikaberidze, A. and McDonald, B. A. (2020). A tradeoff between tolerance and resistance to a major fungal
767 pathogen in elite wheat cultivars. *New Phytologist*, 226:879–890.

768 Mikaberidze, A., McDonald, B. A., and Bonhoeffer, S. (2014). Can high-risk fungicides be used in mixtures
769 without selecting for fungicide resistance? *Phytopathology*, 104:324–331.

770 Mikaberidze, A., Paveley, N., Bonhoeffer, S., and Van Den Bosch, F. (2017). Emergence of resistance to fungi-
771 cides: the role of fungicide dose. *Phytopathology*, 107:545.

772 Milgroom, M. G. (1990). A stochastic model for the initial occurrence and development of fungicide resistance
773 in plant pathogen populations. *Phytopathology*, 80:410.

774 Mouratiadou, I., Latka, C., van der Hilst, F., Müller, C., Berges, R., Bodirsky, B. L., Ewert, F., Faye, B., Heckeley,
775 T., Hoffmann, M., Lehtonen, H., Lorite, I. J., Nendel, C., Palosuo, T., Rodríguez, A., Rötter, R. P., Ruiz-Ramos,
776 M., Stella, T., Webber, H., and Wicke, B. (2021). Quantifying sustainable intensification of agriculture: the
777 contribution of metrics and modelling. *Ecological Indicators*, 129:107870.

778 Murray, J. (2019). What is the chlorothalonil ban imposed by the EU?

Onstad, D. W. and Rabbinge, R. (1985). Dynamic programming and the computation of economic injury levels for crop disease control. *Agricultural Systems*, 18:207–226.

Pacilly, F. C., Hofstede, G. J., Lammerts van Bueren, E. T., Kessel, G. J., and Groot, J. C. (2018). Simulating crop-disease interactions in agricultural landscapes to analyse the effectiveness of host resistance in disease control: the case of potato late blight. *Ecological Modelling*, 378:1–12.

Parnell, S., Gilligan, C. A., Lucas, J. A., Bock, C. H., and Bosch, F. V. D. (2008). Changes in fungicide sensitivity and relative species abundance in *Oculimacula yallundae* and *O. acuformis* populations (eyespot disease of cereals) in Western Europe. *Plant Pathology*, 57:509–517.

Parnell, S., Gilligan, C. A., and van den Bosch, F. (2005). Small-scale fungicide spray heterogeneity and the coexistence of resistant and sensitive pathogen strains. *Analytical and Theoretical Plant Pathology*, 95:632–639.

Parnell, S., van den Bosch, F., and Gilligan, C. A. (2006). Large-scale fungicide spray heterogeneity and the regional spread of resistant pathogen strains. *Analytical and Theoretical Plant Pathology*, 96:549–555.

Phelan, P. (2017). Timing is everything with fungicides - Farming Independent.

Rehfus, A., Strobel, D., Bryson, R., and Stammler, G. (2018). Mutations in *sdh* genes in field isolates of *Zymoseptoria tritici* and impact on the sensitivity to various succinate dehydrogenase inhibitors. *Plant Pathology*, 67:175–180.

Ristaino, J. B., Anderson, P. K., Bebber, D. P., Brauman, K. A., Cunniffe, N. J., Fedoroff, N. V., Finegold, C., Garrett, K. A., Gilligan, C. A., Jones, C. M., Martin, M. D., MacDonald, G. K., Neenan, P., Records, A., Schmale, D. G., Tateosian, L., and Wei, Q. (2021). The persistent threat of emerging plant disease pandemics to global food security. *Proceedings of the National Academy of Sciences*, 118.

Rupp, S., Weber, R. W. S., Rieger, D., Detzel, P., and Hahn, M. (2017). Spread of *Botrytis cinerea* strains with multiple fungicide resistance in German horticulture. *Frontiers in Microbiology*, 0:2075.

Samoucha, Y. (1987). Use of two- and three-way mixtures to prevent buildup of resistance to phenylamide fungicides in *Phytophthora* and *Plasmopara*. *Phytopathology*, 77:1405.

Shaw, M. W. (2000). Models of the effects of dose heterogeneity and escape on selection pressure for pesticide resistance. *Analytical and Theoretical Plant Pathology*, 90:333–339.

Shaw, M. W. and Royle, D. J. (1989). Airborne inoculum as a major source of *Septoria tritici* (*Mycosphaerella graminicola*) infections in winter wheat crops in the UK. *Plant Pathology*, 38:35–43.

Singh, N. K., Karisto, P., and Croll, D. (2021). Population-level deep sequencing reveals the interplay of clonal and sexual reproduction in the fungal wheat pathogen *Zymoseptoria tritici*. *Microbial Genomics*, 7:000678.

810 Staub, T. (1991). Fungicide resistance: practical experience with antiresistance strategies and the role of inte-
811 grated use. *Annual Review of Phytopathology*, pages 421–442.

812 Strange, R. N. and Scott, P. R. (2005). Plant disease: A threat to global food security. *Annual Review of Phy-*
813 *topathology*, 43:83–116.

814 Suffert, F., Delestre, G., Carpentier, F., Gazeau, G., Walker, A. S., Gélisse, S., and Duplaix, C. (2016). Fashionably
815 late partners have more fruitful encounters: impact of the timing of co-infection and pathogenicity on sexual
816 reproduction in *Zymoseptoria tritici*. *Fungal Genetics and Biology*, 92:40–49.

817 Suffert, F., Delestre, G., and Gélisse, S. (2018). Sexual reproduction in the fungal foliar pathogen *Zymoseptoria*
818 *tritici* is driven by antagonistic density dependence mechanisms. *Microbial Ecology* 2018 77:1, 77:110–123.

819 Suffert, F., Sache, I., and Lannou, C. (2011). Early stages of septoria tritici blotch epidemics of winter wheat:
820 build-up, overseasoning, and release of primary inoculum. *Plant Pathology*, 60:166–177.

821 Tai, A. P., Martin, M. V., and Heald, C. L. (2014). Threat to future global food security from climate change and
822 ozone air pollution. *Nature Climate Change*, 4:817–821.

823 te Beest, D., Paveley, N., Shaw, M., and van den Bosch, F. (2008). Disease-weather relationships for powdery
824 mildew and yellow rust on winter wheat. *Phytopathology*, 98:609–617.

825 te Beest, D. E., Paveley, N. D., Shaw, M. W., and van den Bosch, F. (2013). Accounting for the economic risk caused
826 by variation in disease severity in fungicide dose decisions, exemplified for *Mycosphaerella graminicola* on
827 winter wheat. *Analytical and Theoretical Plant Pathology*, 103:666–672.

828 Torriani, S. F., Melichar, J. P., Mills, C., Pain, N., Sierotzki, H., and Courbot, M. (2015). *Zymoseptoria tritici*: A major
829 threat to wheat production, integrated approaches to control. *Fungal Genetics and Biology*, 79:8–12.

830 van den Berg, F., Paveley, N. D., and van den Bosch, F. (2016). Dose and number of applications that maxi-
831 mize fungicide effective life exemplified by *Zymoseptoria tritici* on wheat – a model analysis. *Plant Pathology*,
832 65:1380–1389.

833 van den Berg, F., van den Bosch, F., and Paveley, N. D. (2013). Optimal fungicide application timings for disease
834 control are also an effective anti-resistance strategy: A case study for *Zymoseptoria tritici* (*Mycosphaerella*
835 *graminicola*) on wheat. *Phytopathology*, 103:1209–1219.

836 van den Bosch, F., Blake, J., Gosling, P., Helps, J. C., and Paveley, N. (2020). Identifying when it is financially
837 beneficial to increase or decrease fungicide dose as resistance develops: An evaluation from long-term field
838 experiments. *Plant Pathology*, 69:631–641.

839 van den Bosch, F. and Gilligan, C. A. (2008). Models of fungicide resistance dynamics. *Annual Review of Phy-*
840 *topathology*, 46:123–147.

- 841 van den Bosch, F., Oliver, R., van den Berg, F., and Paveley, N. (2014a). Governing principles can guide fungicide-
842 resistance management tactics. *Annual Review of Phytopathology*, 52:175–195.
- 843 van den Bosch, F., Paveley, N., Shaw, M., Hobbelen, P., and Oliver, R. (2011). The dose rate debate: Does the
844 risk of fungicide resistance increase or decrease with dose? *Plant Pathology*, 60:597–606.
- 845 van den Bosch, F., Paveley, N., van den Berg, F., Hobbelen, P., and Oliver, R. (2014b). Mixtures as a fungicide
846 resistance management tactic. *Phytopathology*, 104:1264–1273.
- 847 Zadoks, J. C., Chang, T. T., and Konzak, C. F. (1974). A decimal code for the growth stages of cereals. *Weed*
848 *Research*, 14:415–421.
- 849 Zhan, J., Mundt, C. C., and McDonald, B. A. (1998). Measuring immigration and sexual reproduction in field
850 populations of *Mycosphaerella graminicola*. *Phytopathology*, 88:1330–1337.

Appendix 1

Between-season sexual reproduction – derivation

For between-season sexual reproduction, we assume that the recombination is random, so that each (haploid) offspring has an equal probability of inheriting each allele from either (haploid) parent (Appendix 1 Table 1). We have two unlinked (assumed) loci, each with two alleles: resistant or sensitive.

Pathogen strains	<i>ss</i>	<i>sr</i>	<i>rs</i>	<i>rr</i>
<i>ss</i>	<i>ss</i>	<i>ss/sr</i>	<i>ss/rs</i>	<i>ss/sr/rs/rr</i>
<i>sr</i>	<i>ss/sr</i>	<i>sr</i>	<i>ss/sr/rs/rr</i>	<i>rr/sr</i>
<i>rs</i>	<i>ss/rs</i>	<i>ss/sr/rs/rr</i>	<i>rs</i>	<i>rr/rs</i>
<i>rr</i>	<i>ss/sr/rs/rr</i>	<i>rr/sr</i>	<i>rr/rs</i>	<i>rr</i>

Appendix 1 Table 1. The possible offspring of sexual reproduction between parents given by the left column and top row are shown. For those entries which contain four strains, the probability of each is 1/4, so each number is accordingly scaled. Similarly those entries containing two strains each have probability 1/2. Note that both parents and offspring are haploid; for example *sr* corresponds to a haploid which carries the allele for sensitivity (*s*) to fungicide *A* on one chromosome, and the allele for resistance (*r*) to fungicide *B* on another.

Let F_{mn} denote the frequency (a proportion $\in [0, 1]$ of total disease) of strain *mn* immediately before the single, instantaneous round of between-season sexual reproduction which we model. Let Y_{mn} represent the frequency of strain *mn* immediately afterwards. Then collecting together all the terms, we find:

$$\begin{aligned}
 Y_{ss} &= F_{ss}^2 + \frac{1}{2}(F_{ss}F_{sr} + F_{sr}F_{ss}) + \frac{1}{2}(F_{ss}F_{rs} + F_{rs}F_{ss}) + \frac{1}{4}(F_{ss}F_{rr} + F_{rr}F_{ss}) + \frac{1}{4}(F_{sr}F_{rs} + F_{rs}F_{sr}) \\
 &= F_{ss}^2 + F_{ss}F_{sr} + F_{ss}F_{rs} + \frac{1}{2}F_{ss}F_{rr} + \frac{1}{2}F_{sr}F_{rs},
 \end{aligned} \tag{33}$$

and similarly:

$$Y_{sr} = F_{sr}^2 + F_{sr}F_{ss} + F_{sr}F_{rr} + \frac{1}{2}F_{ss}F_{rr} + \frac{1}{2}F_{sr}F_{rs}, \tag{34}$$

$$Y_{rs} = F_{rs}^2 + F_{rs}F_{ss} + F_{rs}F_{rr} + \frac{1}{2}F_{ss}F_{rr} + \frac{1}{2}F_{sr}F_{rs}, \tag{35}$$

$$Y_{rr} = F_{rr}^2 + F_{rr}F_{sr} + F_{rr}F_{rs} + \frac{1}{2}F_{ss}F_{rr} + \frac{1}{2}F_{sr}F_{rs}. \tag{36}$$

This can be simplified by noting that, $F_{ss} + F_{sr} + F_{rs} + F_{rr} = 1$, and that, using the *ss* strain

879

880

881

882

883

884

885

886

887

888

889

890

891

892

893

as an for example:

$$\begin{aligned} Y_{ss} &= F_{ss}(F_{ss} + F_{sr} + F_{rs} + F_{rr}) - \frac{1}{2}F_{ss}F_{rr} + \frac{1}{2}F_{sr}F_{rs} \\ &= F_{ss} - \frac{1}{2}F_{ss}F_{rr} + \frac{1}{2}F_{sr}F_{rs}. \end{aligned} \quad (37)$$

Now we introduce the linkage disequilibrium:

$$D = F_{ss}F_{rr} - F_{sr}F_{rs}. \quad (38)$$

Then we find:

$$Y_{ss} = F_{ss} - \frac{1}{2}D; \quad Y_{sr} = F_{sr} + \frac{1}{2}D; \quad Y_{rs} = F_{rs} + \frac{1}{2}D; \quad Y_{rr} = F_{rr} - \frac{1}{2}D. \quad (39)$$

This form is equivalent to that of **Felsenstein (1965)** for a haploid population, with two alleles at two loci and discrete generations (we use F_{mi} in place of their x_i). We assume that the recombination fraction $r = 1/2$, corresponding to the two loci being unlinked. We also assume that the probability of surviving from meiosis to fertilisation is equal for all strains, in line with our assumption that the strains behave identically in the absence of fungicide.

Appendix 2

Between-season sexual reproduction – effect on the model

Depending on initial resistance frequencies and fungicide parameterisations, we can observe:

- non-monotonic response in maximum effective life as p_B increases (Appendix 2 Figure 1A);
- no effect on maximum effective life as p_B increases (Appendix 2 Figure 1B);
- monotone increase in maximum effective life as p_B increases (Appendix 2 Figure 1C).

Appendix 2 Figure 1. Three effects are possible with sexual reproduction. This figure is exactly as in Figure 5 (main text), but highlighting the three qualitative behaviours we see as the proportion of sexual reproduction (p_B) increases. These are: (A) non-monotonicity; (B) no change in effective life; and (C) monotone increase in effective life as p_B increases. **Parameter values:** as in Figure 5 (main text); random fungicide parameter values and single resistant frequencies, generated from distributions described in Table 3. **A:** initial double resistant strain (rr) density 10^{-4} times lower than value from linkage equilibrium (VLE); **B:** initial rr density at VLE; **C:** initial rr density 10^4 times higher than VLE. Values: ($p_{rr}, p_{rs}, p_{sr}, \Lambda_A, \Lambda_B, \omega_A, \omega_B, \theta_A, \theta_B$) = **Scenario 1:** ($7.049916 \times [10^{-19}/10^{-15}/10^{-11}], 5.545997 \times 10^{-7}, 1.269711 \times 10^{-8}, 0.009758, 0.022030, 0.729797, 0.661193, 7.362942, 6.642679$); **Scenario 2:** ($1.085892 \times [10^{-15}/10^{-11}/10^{-7}], 1.596583 \times 10^{-6}, 6.801290 \times 10^{-6}, 0.007417, 0.009834, 0.574543, 0.706497, 11.143576, 11.170345$); **Scenario 3:** ($2.445955 \times [10^{-15}/10^{-11}/10^{-7}], 7.155682 \times 10^{-5}, 3.417953 \times 10^{-7}, 0.005803, 0.006279, 0.875035, 0.717337, 8.544356, 11.404773$).

For medium or high initial levels of the double resistant, we typically only see the latter two behaviours (no change or monotone increase). This is because sexual reproduction acts to suppress the double resistant strain, as shown by the between-season equation for

the rr strain:

$$P_{rr} = F_{rr} - \frac{1}{2}p_B D. \quad (40)$$

Therefore if $D = F_{rr}F_{ss} - F_{sr}F_{rs} > 0$, we expect a decrease in the level of double resistant after the sexual reproduction step (for any $p_B > 0$, i.e. any non-zero amount of sexual reproduction). Note that $D > 0$ is equivalent to $F_{rr} > F_{sr}F_{rs}/F_{ss}$. Initially $F_{ss} \approx 1$, so the threshold $F_{sr}F_{rs}/F_{ss} \approx F_{sr}F_{rs}$. The effect of sexual reproduction decreasing the rate of the double resistant strain increasing is particularly clear in Appendix 2 Figure 2H, and is present but to a lesser extent in Appendix 2 Figure 2D.

When the effective life is the same across different levels of sexual reproduction, this is caused by increases in the single-resistant strains relative to the asexual case (Appendix 2 Figure 2E,F,I,J). This offsets the decrease in the double-resistant strain. Whether the single or double resistant strain plays the major role in yield losses depends on the efficacy of the fungicides – very high efficacy fungicides are capable of controlling the single resistant strains since one component of the mixture remains effective. For example, relatively good control of the single resistant strains in Appendix 2 Figure 2A,B,C means that the effective life is quite long, since it takes a long time for the double resistant to reach sufficiently high frequencies to reduce yields below 95%. In contrast, poor control from fungicide A and good control from fungicide B in Appendix 2 Figure 2D,E,F means that the strains resistant to fungicide B (strains sr , rr) rapidly increase in density and lead to loss of control whilst strain rs declines.

If the level of double resistant is very low ($F_{rr} < F_{sr}F_{rs}/F_{ss}$), then $D < 0$ and the double resistant increases in density (Appendix 2 Figure 2A). This increase is only seen in the first year, after which point the double resistant remains at a level at least as high as $F_{sr}F_{rs}/F_{ss}$ due to the greater selection for the rr strain in comparison to the other strains every year. The non-monotonic effect is due to even low levels of sexual reproduction leading to an increase in the double resistant after the first off-season, but higher levels of sexual reproduction again suppressing the double resistant strain in subsequent years. This is shown by the decreased gradient of sexual proportion $p_B = 1$ vs sexual proportion $p_B = 0.5$ in Appendix 2 Figure 2A.

Appendix 2 Figure 2. Higher between-season sexual reproduction slows the double resistant strain. Three different scenarios, demonstrating the three qualitative behaviours from Appendix 2 Figure 1: non-monotonicity in effective life as p_B increases; no change in effective life as p_B increases; monotonic increase as p_B increases. The dotted line is the effective life of the asexual reproduction case. Note that here we compare full dose of both fungicides in each case, whereas in Appendix 2 Figure 1 and main text, Figure 5, we compare the optimal dose from the 21 by 21 grid of doses in each case. The randomly generated scenarios that we show here are different to any shown in Appendix 2 Figure 1, but here we are comparing something different (full dose rather than optimal dose). **NB** the subplots in each column are for the same scenario, but the scenarios are completely different between columns (not just a change in initial resistance frequencies).

Left column: the 'non-monotonicity' case. In this example the double resistant strain is lower than the value at linkage equilibrium (VLE) by 10^{-4} . Here high values of p_B , or $p_B = 0$ give longer effective lives than intermediate values. **Middle column:** the 'no effect' case. In this example the double resistant strain is at VLE. Here there is a faster increase in the double resistant strain when $p_B = 0$ (D) but this is offset by reduced growth of the single resistant strains (E,F). **Right column:** the 'monotonic increase' case. In this example the double resistant strain is higher than the VLE by 10^4 . Here the faster growth of the double resistant strain when $p_B = 0$ causes a shorter effective life than we see in the case where p_B takes larger values.

Parameter values. Generated at random: (I_{rr} , I_{sr} , I_{rs} , ω_A , ω_B , θ_A , θ_B , Λ_A , Λ_B) = **left column:** (1.085892×10^{-15} , $6.8012899586885546 \times 10^{-6}$, $1.596583402894029 \times 10^{-6}$, 0.574543 , 0.706497 , 11.143576 , 11.170345 , 0.007417 , 0.009834); **middle column:** (1.140463×10^{-10} , $1.545001824789278 \times 10^{-6}$, $7.38107054353044 \times 10^{-5}$, 0.983611 , 0.82889 , 9.581831 , 5.728716 , 0.032598 , 0.003884); **right column:** (1.121325×10^{-10} , $1.0174155443038875 \times 10^{-6}$, $1.1002632709098869 \times 10^{-8}$, 0.697464 , 0.480298 , 5.136889 , 5.748469 , 0.016088 , 0.011044).

978 **Appendix 3**

979 **Between-season sexual reproduction – mixtures vs alternations**

980 We now explore the effect of varying the between-season sexual reproduction p_B on the
981 model output and the optimal strategy. To check that mixtures outperform alternations
982 when between-season sexual reproduction was present, we ran a simple scan where between-
983 season reproduction was entirely sexual. The case where reproduction was entirely asex-
984 ual has previously been examined in *Hobbelen et al. (2013)*; *Elderfield et al. (2018)*, and
985 mixtures were found to outperform alternations.

986 We tested 100 randomly generated scenarios (Table 1). We varied the asymptote, curva-
987 ture and decay rate parameters of both fungicides, as well as the initial resistance frequen-
988 cies, ignoring any parameter sets where either of the alternation strategies had an effective
989 life of 0 (i.e. could never achieve a yield $\geq 95\%$). Using a grid of 51×51 doses, we checked
990 whether the maximum effective life from the mixture tactic in each case was greater than
991 or equal to the maximum effective life from either alternation tactic (applying fungicide *A*
992 first each year or applying fungicide *B* first each year). In every case, the mixture tactic was
993 at least as good as either alternation tactic (Appendix 3 Figure 1).

Variable	Range/value	Distribution
Initial proportion, strain rs, sr	$[10^{-10}, 10^{-4}]$	Log-uniform
Initial proportion, strain rr	$[10^{-15}, 10^{-6}]$	Log-uniform
Fungicide asymptote	$[0.4, 1]$	Uniform
Fungicide curvature	$[4, 12]$	Uniform
Fungicide decay rate A	$[1/3\Lambda_0, 3\Lambda_0]$	Uniform

995 **Appendix 3 Table 1.** Bounds for the parameter scan used to confirm that mixtures outperform
996 alternations. For this scan we used a grid of 51×51 doses, and checked that the maximum effective
997 life from the mixture grid was greater than or equal to the maximum from the alternation grid for
998 each example. We tested the case where between-season reproduction was entirely sexual. Unless
999 stated above, all other parameters take their default values (see Table 1). Note the fungicide decay
1000 rate range depends on Λ_0 , which is 1.11×10^{-2} degree days $^{-1}$ as in Table 1.

Appendix 3 Figure 1. Mixtures outperform alternation even with between-season sexual reproduction. Here are the optimal effective lives from 100 randomly generated scenarios. The scenarios are arranged from smallest to largest effective life of the mixture strategy. Here we consider the optimal dose from a 51 by 51 grid. **Parameter values:** randomly generated, see Appendix 3 Table 1.

Appendix 4

Parameter scan – link between curvature and dose

The following argument will show the link between curvature and dose which allows us to generalise the parameter scan results (Tables 5, 6 and Figure 6) to a wider range of fungicides. In particular, we show that any pair of fungicides with identical decay rate and asymptote but differing curvature can be considered to behave identically with an appropriate change of dose. To meet legal dose requirements (doses less than 1), we note that this applies to any higher efficacy fungicides (increased curvature) than those tested, and some lower efficacy ones (see below).

For a dose C of a fungicide F , we use dose responses of the type

$$\delta_{F,s}(C) = 1 - \epsilon(C), \quad (41)$$

where $\delta_{F,s} \leq 1$ is the factor by which the rates of the transition of tissue from healthy to latent (β), and from latent to infected (γ) are reduced, and ϵ is the effect on sensitive strains:

$$\epsilon(C) = \omega(1 - e^{-\theta C}). \quad (42)$$

We also assume fungicide concentrations decay exponentially, so that t units after a dose application the concentration will be:

$$C(t) = (C_0 + D)\exp(-\Delta t), \quad (43)$$

where Δ is the decay rate, C_0 was the concentration immediately before the application, and D was the applied dose.

Consider a pair of fungicides 1 and 2 which differ only in their curvature, i.e. have parameters $(\omega, \theta_1, \Delta)$ and $(\omega, \theta_2, \Delta)$. Then if we apply a dose of fungicide 2 such that

$$C_2 = \frac{C_1 \theta_1}{\theta_2}, \quad (44)$$

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

then we find:

$$\epsilon_1(C_1) = \omega \left(1 - \exp \left(-\theta_1 (C_0 + D) \exp(-\Delta t) \right) \right), \quad (45)$$

$$\begin{aligned} \epsilon_2(C_2) &= \omega \left(1 - \exp \left(-K \theta_2 \frac{(C_0 + D) \theta_1}{\theta_2} \exp(-\Delta t) \right) \right) \\ &= \omega \left(1 - \exp \left(-\theta_1 (C_0 + D) \exp(-\Delta t) \right) \right) = \epsilon_1(C_1). \end{aligned} \quad (46)$$

That is, if fungicide 1 has a curvature that is a factor of K bigger than fungicide 2, we can apply doses that are a factor of K smaller than fungicide 2. Then we end up with an identical form for ϵ_1 and ϵ_2 , and so an identical form for $\delta_{1,s}$ and $\delta_{2,s}$, even when accounting for the decay in concentration over time. This means that the two fungicides behave identically if these doses are applied (Appendix 4 Figure 1).

1049

1050

1051

1052

1053

1054

1055

1056

1058

Appendix 4 Figure 1. Fungicides with different curvatures can behave identically with appropriate dose choice. One fungicide has a curvature $\theta_h = 10$ and the other has a curvature of $\theta_l = 5$. Both have the same asymptote and decay rate, which is required for this effect. This parameterisation results in the former fungicide having a higher efficacy (A, B). Applying appropriately scaled doses: $D_h = D_l \theta_l / \theta_h$, so that $D_l = 1$, $D_h = 0.5$, (C), the effect is identical (D). Since the lower efficacy fungicide has half the curvature, it requires double the dose. **Parameter values:** higher efficacy fungicide $\theta_h = 10$. Lower efficacy fungicide: $\theta_l = 5$. Both have asymptote $\omega = 0.8$ and decay rate $\Lambda = 1.11 \times 10^{-2}$. Doses applied in B: ($D_h, D_l = 0.5, 1$).

1061

1062

1059

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075

Legal dose caveat

Note that to ensure that we remain within legal requirements for doses ($C < 1$), we require that

$$C_2 = \frac{C_1 \theta_1}{\theta_2} \leq 1, \quad (47)$$

so the parameter scan analysis can only directly be transferred to fungicides with curvatures satisfying

$$\theta_2 \geq \theta_1 C_{1,optimal}. \quad (48)$$

Since minimal doses tend to be optimal, this means our results may be generalised for a range of fungicides that are lower or equal efficacy than those in the scan ($\theta_1 C_{1,optimal} \leq \theta_2 \leq \theta_1$), and any fungicides that are higher efficacy (higher curvature than θ_1).

For instance, if a particular fungicide with curvature 8 had an optimal dose of 0.25 in a mixture, then we could apply our results to any fungicide with curvature $\theta \geq 8 \times 0.25 = 2$, using optimal dose $2/\theta \leq 1$.

1076 Appendix 5

1077 Parameter scan – sub-optimal runs

1078 Runs in which equal resistance frequencies at breakdown was sub-optimal

1079 There were initially 8 runs of the 500 for which equal resistance frequencies at breakdown
1080 (ERFB) appeared to not give optimal results. However, when re-running so that a much
1081 larger number of doses along the contour were chosen (500 and in a narrower range around
1082 the optimum), 3 of these cases were found to be optimal, leaving only 5 runs that were sub-
1083 optimal after the more computationally intensive search. The remaining 5 runs (Appendix
1084 5 Figure 1) the results remained sub-optimal by one year even when the denser grid was
1085 used. Parameter values for these runs are in Appendix 5 Table 1.

1086 In each of these cases, the optimal region is very small. There is a much larger region
1087 which achieves an effective life within one year of the optimal value. In this larger region,
1088 the resistance management offered is relatively good. The doses in the optimal region offer
1089 better control in that year than the surrounding region. The location of this region depends
1090 on the efficacy of the two fungicides, and in these 5 cases the ERFB contour does not pass
1091 through the optimal region. We examine one of these runs (scenario B) in Appendix 5 Figure
1092 2.

1093 Some dose-pairs near the optimal region are sub-optimal effective lives because they of-
1094 fer slightly poorer control, despite good resistance management. Other higher dose choices
1095 near the optimal region are sub-optimal because the poorer quality resistance management
1096 is bad enough to not offset the greater control caused by higher doses in the final year.

Appendix 5 Figure 1. All sub-optimal runs. All 5 sub-optimal runs are shown, here on a 101 by 101 grid. In each case the optimal region is very small and gives a one year improvement relative to the best dose along the ERFB contour. In each case the contour narrowly misses the optimal region. **Parameter values:** see Appendix 5 Table 1.

		A	B	C	D	E
	ω_A	0.492317	0.812490	0.815263	0.814526	0.924196
	ω_B	0.637730	0.692780	0.675103	0.806968	0.930859
	θ_A	6.816520	2.951951	10.259259	7.407919	4.753716
1103	θ_B	9.333866	7.009989	3.628018	9.627502	10.083944
1104	Λ_A	0.003854	0.004001	0.013152	0.011560	0.012227
	Λ_B	0.009849	0.006771	0.005065	0.004593	0.004933
	p_{rr}	7.540441×10^{-8}	1.165713×10^{-9}	1.679725×10^{-9}	3.379948×10^{-9}	2.632505×10^{-10}
	p_{rs}	2.332734×10^{-5}	5.311738×10^{-10}	2.214624×10^{-9}	8.008551×10^{-9}	1.153372×10^{-9}
	p_{sr}	2.697823×10^{-6}	1.312246×10^{-10}	5.736566×10^{-4}	2.206591×10^{-5}	5.557970×10^{-8}
	p_B	0.6347437	0.6778327	0.488108	0.282625	0.108196
1106	Appendix 5 Table 1. Parameter values for the sub-optimal runs shown in Appendix 5 Figure 1.

Appendix 5 Figure 2. Final year control causes the small number of exceptions where the ERFB strategy is sub-optimal. The optimal strategy achieves a yield of 95.003% in year 12, compared to the ERFB strategy which achieves 94.92% in this year (B). The optimal strategy fails in year 13, whereas the ERFB strategy fails in year 12 (A). Despite lower incidences of all resistant strains under the ERFB strategy, (F, G, H), the control offered by the doses is lower in year 12, since the dose of fungicide B is much lower. Each resistant strain increases faster under the optimal strategy, meaning that the differences in frequency increase over time (C, D, E).

Parameter values: see Appendix 5 Table 1, scenario B. Doses: optimal = 0.18, 0.2, ERFB = 0.185, 0.175.

Taking the first of the sub-optimal examples (Appendix 5 Figure 1A), we see that for a small perturbation in initial resistance frequencies this effect disappears and ERFB is optimal (Appendix 5 Figure 3). This supports the conclusion that the effect is caused by control in the final year, which is why every sub-optimal run of the 500 was only worse than the optimal by a single year. By the failure year, usually resistance frequencies are increasing rapidly and yield is falling steeply (main text, Figure 2F, 2G). This is why it is rare to have such

1122

1123

1124

a small optimal region; more commonly all doses within a larger optimal region are further above 95% in the penultimate year and further below 95% in the breakdown year.

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

Appendix 5 Figure 3. ERFB is optimal under small perturbation of initial conditions of sub-optimal run. This is the sub-optimal run with a grid size of 201 by 201. The optimal region is very small and outperforms ERFB because the increased control is offered by a higher dose of fungicide *B* (A). However, a small perturbation away from the initial conditions removes this effect, and we find that the ERFB strategy is optimal again (B). Here the green region no longer exists and the optimum effective life is now 9 (yellow).

If we gradually decrease resistance frequencies, the scenario can change from something analogous to panel B to something analogous to panel A. This is because the doses which achieve yields fractionally above 95% in the final year may not be the same as those which are best for longer term resistance management, since control and resistance management may not align in any given year. In practice there are very few cases where this phenomenon is observable, as shown by the fact it does not impact 99% of cases.

Parameter values:

Pathogen parameters, A: see Appendix 5 Table 1, scenario A.

Pathogen parameters, B: $(I_{rr}^B, I_{rs}^B, I_{sr}^B, I_{ss}^B) = (1.1 \times I_{rr}^A, 0.9 \times I_{rs}^A, 1.1 \times I_{sr}^A, 1 - I_{rr}^B - I_{rs}^B - I_{sr}^B)$.

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

Run in which equal selection in the first year outperformed equal resistance frequencies at breakdown

There was a single case out of the 500 in which equal selection in the first year (ESFY) gave an improved output relative to ERFB. Here the best ERFB effective life matched the optimal value from the 51 by 51 grid, but the best ESFY effective life was one year greater than this optimal grid value (Appendix 5 Figure 4). Note that Appendix 5 Figure 4 uses a denser grid of 201 by 201 to adequately illustrate this effect. See also Appendix 5 Figure 5. Note that the comparison in the parameter scan was between the 51 by 51 grid, ESFY and ERFB; it is possible that there are other cases where a finer grid might find small regions which achieve one year greater than the ERFB optimum and the 51 by 51 grid optimum.

1152
1153
1154
1155
1156
1157
1158
1159
1160

Appendix 5 Figure 4. The single case where ESFY outperformed ERFB. In this case the ESFY contour passes through the optimal region, but the ERFB contour does not. This was the only case in which this happened. Although ERFB is sub-optimal on the 201 by 201 grid, it was optimal on the coarser 51 by 51 grid.

Parameter values: Initial frequencies: $I_{rr} = 1.3122930696727086 \times 10^{-10}$, $I_{rs} = 1.3687346286628245 \times 10^{-7}$, $I_{sr} = 9.216054320447214 \times 10^{-7}$, $I_{ss} = 0.9999989413898756$, $p_B = 0.0334369453801207$. Fungicide parameters: $\omega_A = 0.5100900734551262$, $\theta_A = 5.336781378462673$, $\Lambda_A = 0.003895561458581$, $\omega_B = 0.8667335745106877$, $\theta_B = 4.618154403228932$, $\Lambda_B = 0.0080007828566827$.

Appendix 5 Figure 5. The single case where ESFY outperformed ERFB - explained. The optimal strategy achieves a yield of 95.002% in year 11, compared to the ERFB strategy which achieves 94.99% in this year (B). The optimal strategy fails in year 12, whereas the ERFB strategy fails in year 11 (A). In this case the ERFB strategy selects less strongly for the r_s and r_r strains but more strongly for the s_r strain. **Parameter values:** Initial frequencies: $I_{rr} = 1.3122930696727086 \times 10^{-10}$, $I_{rs} = 1.3687346286628245 \times 10^{-7}$, $I_{sr} = 9.216054320447214 \times 10^{-7}$, $I_{ss} = 0.9999989413898756$, $p_B = 0.0334369453801207$. Fungicide parameters: $\omega_A = 0.5100900734551262$, $\theta_A = 5.336781378462673$, $\Lambda_A = 0.003895561458581$, $\omega_B = 0.8667335745106877$, $\theta_B = 4.618154403228932$, $\Lambda_B = 0.0080007828566827$, Doses: ESFY = 0.19, 0.26, ERFB = 0.215, 0.23.