1 Improved fluorescent proteins for dual-color post-embedding CLEM

2 Dingming Peng^{1, 4, 8}, Na Li^{1, 2, 3, 5, 8}, Wenting He¹, Kim Ryun Drasbek⁵, Tao Xu^{4, 6*}, Mingshu Zhang^{1, 4*}
3 and Pingyong Xu^{1, 4, 6, 7*}

4 ¹ Key Laboratory of RNA Biology, Institute of Biophysics, Chinese Academy of Sciences, Beijing, China.

5 ² Sino-Danish College, University of Chinese Academy of Sciences, Beijing, China.

6 ³ Sino-Danish Center for Education and Research, Beijing, China.

7 ⁴ Bioland Laboratory (Guangzhou Regenerative Medicine and Health Guangdong Laboratory), 510005, Guangzhou, China.

8 ⁵ Center of Functionally Integrative Neuroscience, Dept of Clinical Medicine, Aarhus University, Aarhus, Denmark.

9 ⁶ National Laboratory of Biomacromolecules, CAS Center for Excellence in Biomacromolecules, Institute of Biophysics, Chinese
10 Academy of Sciences, Beijing, China.

11 ⁷ College of Life Sciences, University of Chinese Academy of Sciences, Beijing, China.

12 ⁸ These authors contributed equally.

13 *e-mail: pyxu@ibp.ac.cn; zhangmingshu@ibp.ac.cn; xutao@ibp.ac.cn

15 Abstract

16 Post-embedding correlative light and electron microscopy (CLEM) has the advantage of high-precision
17 registration and enables light and electron microscopy imaging of the same slice. However, its broad
18 application has been hampered by the limited available fluorescent proteins (FPs) and low signal-to-
19 background ratio (SBR). Here, we developed a green photoswitchable FP, mEosEM-E with substantially
20 high on/off contrast in EM samples embedded in Epon resin which maximally preserves cellular
21 structures but quenches the fluorescence of FPs. Taking advantage of the photoswitching property of
22 mEosEM-E, the autofluorescence background from the resin was significantly reduced by a subtraction-
23 based CLEM (sCLEM) method. Meanwhile, we identified a red fluorescent protein (RFP) mScharlet-H
24 that exhibited higher brightness and SBR in resin than previously reported RFPs. With mEosEM-E and
25 mScharlet-H, dual-color post-Epon-embedding CLEM images with high SBR and no cross-talk signal
26 were successfully performed to reveal the organization of nucleolar proteins. Moreover, a dissection of
27 the influences of different EM sample preparation steps on the fluorescence preservation for several RFPs
28 provides useful guidance for further probe development.

30 **Keywords:** Dual-color CLEM; high SBR; RSFP; probe development

32 Introduction

33 Light microscopy (LM) highlights and discriminates one or a few biomolecules using fluorescent
34 labeling. Electron microscopy (EM), on the other hand, reveals the ultrastructural context of the cell
35 where the biomolecules reside. Correlative light and electron microscopy (CLEM) [1] integrates the
36 information of both LM and EM in a complementary way, and thus provides localization, structural and
37 functional insights into the biomolecular machines, and shows great application prospects in cellular
38 physiology [2], virology [3] and neuroscience [4]. In CLEM imaging, LM can not only indicate the
39 location of the target protein, but also be used as a guide to quickly find slices containing target cells or
40 proteins in a large number of EM slices, and to quickly locate the cell on the target slice for EM imaging.
41 The advantage of fluorescence navigation is unique and is not possessed by EM labeling techniques,

such as immunogold labeling, APEX-gold [5], APEX2 [6], miniSOG [7], and other metal [8] or chemical [9] tags.

Same as any other imaging technique, the practical performance of CLEM is largely determined by the labeling probe. However, most fluorescent proteins (FPs) experienced severe fluorescence loss during standard EM sample preparation, making it difficult to perform LM and EM on the same section that can be correlated with high precision. To circumvent this problem, several strategies have been used. Pre-embedding CLEM [10, 11] obtains fluorescence microscopy (FM) image ahead of EM sample preparation, avoiding fluorescence quenching caused by a series of chemical treatments. However, because of the morphology distortion during EM sample preparation and sectioning, it suffers from poor registration between FM and EM images. Several modified EM sample preparation protocols with no or reduced OsO₄ concentration have been reported to preserve sufficient fluorescence signal for post-embedding CLEM [12, 13], however, deteriorated EM images were often observed.

In 2015, Paez-Segala et al. engineered the first FP, mEos4b, to retain fluorescence after 1% OsO₄ treatment [14]. However, the hydrophilic GMA resin was used for mEos4b-labeled samples. Compared to the GMA resins, the hydrophobic Epon resin has advantages in maintaining cellular ultrastructure as well as sectioning quality [3, 15] due to its higher toughness and hardness. Nevertheless, Epon resin undermines fluorescence more severely. To solve this problem, we have previously developed mEosEM [16], an FP that survives 1% OsO₄ fixation and Epon embedding and enables super-resolution CLEM (SR-CLEM) due to the preserved photomodulable property. Recently, Tanida et al. found that mKate2 [17], mCherry2, mWasabi, and GoGFP-v0 [18] could also preserve fluorescence after standard EM sample preparation and dual-color CLEM imaging was achieved using mWasabi and mcherry2. However, both the green and the red channel images showed very low signal-to-background ratio (SBR), making it difficult to distinguish the real fluorescence signals from that of the background or noises.

Low SBR can be attributed to three aspects. First, most of the fluorescence signals of the FPs mentioned above are quenched after TEM sample preparation, and only a small amount of the remaining fluorescence signal is used for CLEM imaging. Second, the thickness of ultrathin sections is generally only about 100 nm, and the number of FP-labeled molecules in the sections is limited. Third, the Epon resin has strong autofluorescence, especially in the green channel (Figure S1). Therefore, developing FPs with high in-resin SBR and repressing the autofluorescence background are effective ways to solve the problem. For dual-color CLEM, another phenomenon worth noting is that red fluorescent proteins (RFPs) on Epon-embedded slices emitted green fluorescence when illuminated with 488-nm laser (Figure S8), which interfered the signal in the green channel. Therefore, FPs and imaging methods that can solve these problems are desirable.

Reversibly photoswitchable FP (RSFP) can be utilized to suppress the unmodulatable fluorescent background and enhance the signal contrast by means of optical lock-in detection (OLID) [19], synchronously amplified fluorescence image recovery (SAFIRE) [20, 21] and out-of-phase imaging after optical modulation (OPIOM) [22]. We speculated that similar strategies could be applied with an OsO₄- and Epon-resistant RSFP to eliminate the resin background and the RFP crosstalk signal. In this study, we developed a fluorescence background-reduced CLEM method (sCLEM) using a simple subtraction of images of RSFPs at fluorescent on- and off-states and efficiently extracted the fluorescence signals of the FPs from that of the background. The higher on/off contrast of the RSFP, the better SBR of the final image. Therefore, we evolved a mEosEM variant termed mEosEM-E, with high on-state brightness and on/off contrast ratio after standard EM sample preparation, and demonstrated its utility in sCLEM imaging.

On the other hand, despite a few FPs were reported for post-Epon-embedding CLEM [14, 16-18], all these probes were discovered by OsO₄ resistance assay, the results of which are sometimes inconsistent with the final performance of the probe in CLEM imaging. As a matter of fact, nearly every step during EM sample preparation will quench the fluorescence of FPs. However, a comparative study that dissects the influence of each step on fluorescence signal reduction is lacking. In order to develop an optimal RFP that can be coupled with mEosEM-E for dual-color CLEM imaging, we assessed the fluorescence preservation of nine commonly used RFPs after each key step of EM sample preparation, including pre-fixation, OsO₄ fixation, ethanol dehydration, and Epon embedding. The results showed that the OsO₄ resistance assay is not the optimal criterion for CLEM probe development, while the performance of a probe should be evaluated in the final sample section. mScarlet-H, which preserved the highest on-section fluorescence and SBR among the nine RFPs tested, is the best RFP reported to date for the same section post-Epon-embedding CLEM. Finally, high SBR and accurate dual-color CLEM imaging of nucleolar proteins was successfully achieved for the first time using mEosEM-E and mScarlet-H double labeling.

Results

Green mEosEM-E with high SBR for sCLEM

To obtain a CLEM probe with high on/off contrast, we chose mEosEM as the template. mEosEM is a photoconvertible FP (PCFP) that can be converted to an RFP upon 405-nm illumination, but it also has photoswitching property at the green state like an RSFP. After the TEM sample preparation, mEosEM lost the characteristic of light conversion while retaining the photoswitching property[16]. However, this photoswitching property was not used for background removal in CLEM imaging, and the on/off contrast was not optimized for this use.

Our previous study revealed that the first amino acid of the chromophore tripeptide (XYG) of the Eos FP family not only determines the photomodulable type (RSFP or PCFP) of the FP but also greatly affect its photoswitching property [23]. We reasoned that because the first amino acid of the chromophore is located inside the barrel structure of the FP, mutagenesis at this site would have no negative effect on the resistance to the EM sample preparation, while largely tuning the photoswitching contrast. As expected, saturation mutagenesis at His63 of mEosEM produced a series of RSFP mutants displaying a variety of switching contrast and kinetics (Figure S2). We found that among several improved mutants, mEosEM-E (mEosEM H63E) displayed substantially improved contrast, in other words, largely reduced normalized residual fluorescence as compared to mEosEM after TEM sample preparation (Figure 1A). The on/off contrast ratio of mEosEM-E is ~2.7-fold higher than that of mEosEM on average (Figure 1B).

To demonstrate the superiority of mEosEM-E in CLEM, we imaged HEK 293T cell sections expressing mitochondria-targeting mEosEM-E or mEosEM. Fluorescence image sequences were acquired under the illumination of the 488-nm laser alone followed by the 488- and 405-nm lasers together. We termed the images acquired without the illumination of 405-nm laser as “OFF images” and those with 405-nm laser as “ON images”. Every 20 frames of the OFF and subsequent ON images from the image stack were averaged respectively. Then, the subtraction was performed pixel by pixel between the averaged ON and OFF images to acquire the final “ON – OFF image” (Figure S3). We named the subtraction-based LM as sLM, and accordingly CLEM as sCLEM. It can be clearly seen that the autofluorescent background in the OFF image was almost completely removed in the ON – OFF image of mEosEM-E-labeled cells (Figure 1C). For both mEosEM and mEosEM-E, the SBR in the ON – OFF image was significantly increased compared to the ON image (Figure 1D & E). Notably, the SBR of mEosEM-E fluorescence is 3.46-fold higher than that of mEosEM in the ON – OFF image (Figure 1F). Further exploration revealed that the substantial higher SBR in the ON – OFF image of mEosEM-E is mainly attributed to its higher SBR in the ON image (Figure S4).

Next, same section post-Epon-embedding sCLEM was demonstrated in HEK 293T cells, of which the nuclear lamina (Figure 1G-I) and the mitochondria matrix (Figure 1J-L) were labeled by mEosEM-E, separately. The results showed that high SBR and nearly background-free fluorescence images were obtained and well aligned with the EM images.

Altogether, mEosEM-E proved to be a better CLEM probe for high SBR imaging and is potentially useful for investigating proteins of low expression level.

Figure 1. mEosEM-E is an improved green RSFP for high SBR sCLEM imaging. (A) Representative normalized photoswitching curves of Epon-embedded HEK 293T cell sections (100 nm) expressing mitochondria-targeted mEosEM (red) and mEosEM-E (blue). Samples were excited and switched off with the 488-nm laser (0.41 kW/cm²), and switched on with the 405-nm (0.21 kW/cm²) laser for 1 s. (B) Statistics photoswitching contrast ratio of mEosEM (red) and mEosEM-E (blue). Data are summarized in Table S1. Bars represent mean $\pm$ SD. P values were determined with two-tailed t-test (n = 9). **** indicates $p < 0.0001$. (C) Representative OFF (left), ON (middle), and ON — OFF (right) images of Epon-embedded HEK 293T cell sections (100 nm) expressing mitochondria targeting mEosEM-E. Scale bar, 5 μ m. (D, E) SBR of mEosEM- (D) and mEosEM-E- (E) labeled cell samples in OFF (purple), ON

(green), and ON — OFF (orange) images. Data are summarized in Table S2 & 3. P values were determined with two-tailed t-tests ($n = 9$). *** indicates $p < 0.001$, **** indicates $p < 0.0001$. (F) Statistics SBR of mEosEM (red) and mEosEM-E (blue) in ON — OFF images. Two-tailed t-tests were performed ($n = 9$). **** indicates $p < 0.0001$. Data are summarized in Table S4. (G-L) Post-Epon-embedding CLEM of mEosEM-E. FM, EM, and CLEM images of nuclear envelope (G-I) and mitochondria (J-L) labeled by mEosEM-E (100 nm slices). Scale bars, 2 μm .

Identification of mScarlet-H as a high-performance red CLEM probe

To find the optimal red probe for dual-color CLEM, nine commonly used RFPs were chosen for investigation: mScarlet, mScarlet-H, mScarlet-I, FusionRed-MQV, mRuby3, mApple, tdTomato, mKate2 [17], mCherry2 [18], of which the latter two were previously demonstrated feasible for post-Epon-embedding CLEM. Instead of performing only the OsO_4 -resistant assay reported in the previous papers [16-18], we thoroughly analyzed the fluorescence preservation after each key step of the EM sample preparation procedure, including aldehyde fixation, OsO_4 fixation, ethanol dehydration, Epon embedding, and high-temperature polymerization.

A direct comparison of the absolute fluorescence intensity in fixed cells, which is a representative of the practical performance of the probe, showed that pre-fixation with 2% formaldehyde and 2.5% glutaraldehyde already significantly reduced fluorescence intensities of several RFPs, up to 50% of their initial value. While a few others, such as mScarlet-H and FusionRed-MQV were much less affected by pre-fixation, indicating a varied response among different RFPs (Figure S5). The fluorescence intensities of all RFPs were markedly reduced after 1% OsO_4 fixation, nevertheless, mScarlet-I, mRuby3, mScarlet, and mScarlet-H are the top 4 FPs retaining high residue fluorescence, which are all substantially higher than that of the previously reported mKate2 and mCherry2 (Figure 2A). However, the subsequent dehydration step using 100% ethyl alcohol changed the ordering of RFPs, making only mRuby3 standout, while others diminished to a similar level (Figure 2B). We speculated that the advantage of mRuby3 at this stage is due to the insensitivity of mRuby3 to the dehydration treatment, which was later confirmed by the results from dehydration treatment without OsO_4 fixation (Figure S6). Notably, the robustness of mRuby3 to the dehydration treatment following 1% OsO_4 fixation is not as strong as that to a single dehydration treatment, indicating a superimposed effect of the two treatments.

Next, we performed an *in vitro* thermostability test at 60 $^{\circ}\text{C}$ to mimic the effect of high-temperature on fluorescence preservation during Epon resin polymerization. Among nine selected RFPs, mScarlet-H showed the best thermostability that is significantly higher compared to the others (Figure 2C). After complete EM sample preparation, mScarlet-H showed the highest residue fluorescence, followed by mScarlet-I, which are both significantly higher than the previously reported mKate2 and mCherry2, while the residue fluorescence of mRuby3 was surprisingly low (Figure 2D). Statistic results showed that mScarlet-H has the highest SBR among tested RFPs (Figure S7). We speculated that mRuby3 might be super sensitive to the Epon embedding procedure. In addition, even though mApple and tdTomato show comparable fluorescence residue to mKate2 and mCherry2 after OsO_4 fixation and dehydration treatment, and even better thermostability than mCherry2 and mKate2, they totally lose their fluorescence after complete EM sample preparation (data not shown).

Additionally, we also measured the photostability of these RFPs on standard EM sample sections under the illumination condition of CLEM imaging. mScarlet-H had higher photostability than other RFPs except mRuby3 (Figure 2E, F).

Taken together, we recommend mScarlet-H as the best red probe for standard CLEM as it has the highest in-resin fluorescence and a preferable photostability among RFPs after complete TEM sample preparation.

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

Figure 2. Identification of optimal RFPs for CLEM imaging. (A) Residual fluorescence intensity of RFPs after 1% OsO₄ post-fixation. Red stars represent comparisons to mCherry2, black stars represent comparisons to mKate2. (B) Residual fluorescence intensity of RFPs after 1% OsO₄ post-fixation followed by absolute ethanol dehydration. (C) Thermostability of RFPs at 60 °C. (D) Fluorescence intensity of RFPs on Epon-embedded sections (100 nm). (E) Normalized photobleaching curves of RFPs on Epon-embedded sections (100 nm). Enlarged view of the boxed area is shown on top. (F) Statistics of the photobleaching time when the fluorescence intensity of RFPs reduced to 1/e of their initials. Bars represent mean ± SD. P values were determined with two-tailed t-tests in (A-C) (n = 3) and (F) (n = 5), Mann-Whitney U test was performed in (D) (n = 106). n.s. indicates p > 0.05, * indicates p < 0.05, ** indicates p < 0.01, *** indicates p < 0.001, **** indicates p < 0.0001. Data are summarized in Table S5 & 9.

224

mScarlet-H is a generalizable red probe for post-Epon-embedding CLEM

225

226

227

228

229

230

231

232

233

234

235

236

To exemplify the utility of mScarlet-H in Epon-embedded same-section CLEM, we constructed mScarlet-H fusions with the mitochondria targeting peptide, the endoplasmic reticulum membrane protein Sec61 β , and histone H2B, individually. Cell sections transiently expressing mScarlet-H fusions were prepared according to the standard TEM sample preparation procedure. The fluorescence signal of mScarlet-H from thin sections (100 nm) was consecutively recorded for multiple frames. The first 100 frames were averaged to smooth the background noise and increase the signal-to-noise ratio. Mitochondria, endoplasmic reticulum, and Histone H2B were correctly targeted and clearly identified under fluorescence microscopy (FM) (Figure 3A, D, G). Next, to obtain the corresponding ultrastructure of these cell components, the same sections were subsequently imaged using TEM (Figure 3B, E, H). All three structures were successfully aligned with high accuracy (Figure 3C, F, I). These results demonstrate that mScarlet-H is a high-performance red probe that can be generalized for same-section CLEM imaging of various cellular targets.

Figure 3. Post-Epon-embedding CLEM imaging by mScarlet-H. FM, EM, and CLEM images of the mitochondrial matrix (A-C), the endoplasmic reticulum sec61β protein (D-F), and the nucleosome H2B protein (G-I) labeled by mScarlet-H (100 nm slices). Scale bars, 5 μm.

Dual-color post-Epon-embedding same-section CLEM using mEosEM-E and mScarlet-H

mScarlet-H and mEosEM-E were then combined to perform dual-color same-section CLEM imaging. The nuclear envelope (laminA) and mitochondria matrix in HEK 293T cells were labeled by mScarlet-H and mEosEM-E, respectively. After standard EM sample preparation, dual-color wide-field FM images of cell sections (100 nm) were sequentially acquired with a 561-nm laser for 100 frames to image mScarlet-H, followed by 20 frames with a 488-nm laser, and 20 frames with 488- plus 405-nm lasers to image mEosEM-E. For the red channel, images were averaged to obtain the final image; while for the green channel, sLM was applied as mentioned above. As shown in Figure 4A, the mScarlet-H-labeled laminA in the red channel exhibited a good nuclear envelope structure with a high SBR. However, in the green channel, regardless of whether mEosEM-E was in the on- (Figure 4B) or off- state (Figure 4C), the SBR of the fluorescence signal was very low. Furthermore, in addition to mEosEM-E-labeled mitochondrial structure, nuclear envelope structure due to mScarlet-H crosstalk could also be observed as indicated by the merged image of mEosEM-E (on-state) and mScarlet-H (Figure 4D, box). Notably, using the photoswitching property of mEosEM-E and the sCLEM method, the structure of mitochondria could be obtained with a very high SBR (Figure 4E). Moreover, the mScarlet-H crosstalk signal was effectively eliminated (Figure 4F) by the sCLEM imaging scheme, as the green signal of mScarlet-H does not possess the photoswitching property. A further investigation showed that the crosstalk phenomenon is universal for all the tested RFPs that survived standard EM sample preparation (Figure S8). A very likely reason is that under high-energy illumination, severe chromophore photoreduction happens that leads to obvious red-to-green photoconversion of the RFPs, a mechanism that has been reported before [24-26].

Next, we applied dual-color CLEM to investigate how nucleolar proteins are organized in the nucleolus. The nucleolus of mammalian cells has three morphologically distinct components: the fibrillar center (FC), the dense fibrillar component (DFC), and the granular component (GC), which perform different functions in ribosome biogenesis [27]. We labeled the GC marker B23 with mEosEM-E and labeled Nopp140, whose localization has been controversially reported to be at the FC or DFC [28, 29], with mScarlet-H. The results showed that in the green channel, typical GC structures were clearly visible in FM images after sLM, again exemplifying the utility of the subtraction strategy (Figure 4G-I). Merging of green and red channels revealed that the red signal was encircled by the green signal (Figure 4J-L).

Further correlative FM and EM images (Figure 4M-O) showed that the fluorescence signal of Nopp140 was closely matched to the solid high electron-dense regions (Figure 4N, arrowheads) in the EM image, indicating a DFC localization, which is consistent with the fact that it's a functional different area other than GC (Figure 4M, O). Additionally, we also double-labeled Nop52 and Nopp140 by mEosEM-E and mScarlet-H, respectively. Unexpectedly, while Nop52 was often used as a marker of GC [30], Nopp140 was not located in the vacant holes devoid of the Nop52 signals, but partially overlapped with Nop52 (Figure S9). These results indicated that Nop52 localized both at the GC and the DFC.

Taken together, these results strongly demonstrate that mEosEM-E and mScarlet-H are optimal probes for high-quality dual-color post-Epon-embedding CLEM imaging, and sCLEM with mEosEM-E offers a great advantage in reducing the green background of the Epon resin as well as in eliminating the signal crosstalk of RFPs.

Figure 4. Dual-color post-Epon-embedding CLEM using mEosEM-E and mScarlet-H. (A-F) Dual-color imaging of mEosEM-E-labeled mitochondria and mScarlet-H-labeled LaminA protein in HEK 293T cell sections (100 nm). Representative red channel (A), green channel ON (B), and OFF (C) FM images. (D) Merged image of (A) and (B). White boxes indicate areas that showed crosstalk signals from mScarlet-

H. (E) Green channel ON — OFF image. (F) Merged image of (A) and (E). White boxes indicate the same areas in (D). Scale bar, 5 μ m. (G-O) Dual-color CLEM imaging of nucleolar proteins. OFF (G), ON (H), and ON — OFF (I) images of mEosEM-E labeled B23 in HEK 293T cells sections (100 nm). White arrowheads indicate areas that were not visible in (G) and (H). Scale bar, 2 μ m. Green channel FM (J), Red channel FM (K), merged channel FM (L), and CLEM (M) images of HEK 293T cell sections expressing mEosEM-E labeled B23 and mScarlet-H labeled Nopp140. Gamma value: 1.6 for both channels. Scale bars, 2 μ m. (N-O) Enlarged EM (N) and CLEM (O) images of boxed area in (M), white arrowheads indicate the localization of Nopp140 that showed higher EM contrast compared to B23. Gamma value: 1.6 for red channel. Scale bar, 1 μ m.

Discussion

In CLEM imaging, there is always a trade-off between the performance of the FM and the EM. Thus, it is challenging to obtain good FM and EM images of the same ultrathin section at the same time. Developing CLEM probes with resistance to the treatment of EM sample preparation is an efficient way to solve this problem, thus, making CLEM with both high-quality FM and EM possible.

Post-Epon-embedding CLEM was enabled by several FPs reported very recently, however, when these probes were used in practical CLEM imaging, the SBR of FM is very low. The problem is more prominent for imaging low-abundance proteins. In the current study, we utilized the optical switching properties of RSFPs and proposed an sCLEM method, by which the FP signal can be easily distinguished from that of the unmodulated background. sCLEM works more efficiently than the OLID technique in our case, which may be due to the phenomenon that the background fluorescence also showed minor on and off responses to 405- and 488-nm excitation. Compared to the single-frame subtraction (Max-Min), subtraction between the multi-frame averages of the on and off states ($\text{Sum}_{\text{AVG}} - \text{Sum}_{\text{AVG}}$) has an additional benefit of largely attenuated noise. Notably, the sCLEM method hits two birds with one stone. In addition to effectively removing the background signal of the Epon resin and improving the SBR of the FM image, it also removes the signal crosstalk of the RFP in the green channel during dual-color CLEM. To further improve the SBR of FM image, we used mEosEM as the template and developed a variant mEosEM-E with enhanced on-state brightness and on/off contrast that is beneficial for fluorescent background elimination and weak signal extraction. We also identified mScarlet-H as the best RFP for post-Epon-embedding CLEM. This was validated by actual imaging applications, where we proved that mEosEM-E/mScarlet-H is an excellent FP pair for same-section dual-color CLEM of post-Epon-embedded samples. Compared with mEos4b, mEosEM retained higher fluorescence intensity after osmic acid treatment [16]. Considering that the fluorescence intensity retained in GMA-embedded samples is mainly dependent on the osmic acid-resistant properties of the FP, we suspected that mEosEM-E/mScarlet-H would also be excellent choices for the hydrophilic GMA resin.

Besides the development of an advanced green CLEM probe, we also performed a meticulous study that thoroughly dissected the influence of key chemical treatments during TEM sample preparation on the fluorescence preservation of RFPs. Interestingly, we found that the responses of RFPs to these treatments vary greatly. Some RFPs (mRuby3, tdTomato) showed good OsO_4 resistance, however, their superiorities diminished after complete EM sample preparation, suggesting that they are more sensitive to the subsequent sample treatments. Therefore, we reasoned that the OsO_4 resistance assay is not a suitable gold standard for CLEM probe screening, even though OsO_4 fixation seems to be the uppermost factor for fluorescence reduction. Development of an optimal probe requires investigation of the probe's final performance after complete sample preparation, even though this will significantly increase the workload of probe development, and add challenges for high-throughput screening. Furthermore, this study also provided useful information that may assist the rational design of new types of probes. For example,

mRuby3 exhibited superior robustness towards dehydration, which may provide clues for developing FPs that can work in a hydrophobic environment.

Before this study, nucleolar proteins have never been inspected by the same section CLEM. Our dual-color CLEM results revealed that although both have been considered as a GC marker, Nop52 and B23 have different sub-nucleolar localizations. The partial co-localization of Nop52 with Nopp140 suggests that B23 would be a better marker to represent GC. Moreover, we confirmed a DFC localization of Nopp140 in our experimental conditions.

Further directions for the development of post-Epon-embedding CLEM probes would be: 1) the establishment of a high-throughput automatic on-section screening system; 2) a red RSFP CLEM probe for background-reduced and/or SR dual-color CLEM, for which mScarlet-H may be a good starting template; 3) both green and red CLEM probes with even higher on-section brightness, SBR, and photostability.

Materials and methods

Development of mEosEM-E

Saturation mutagenesis of mEosEM at His63 was performed in the pEGFP-N1-mito-mEosEM plasmid with Q5 high fidelity polymerase (New England Biolabs). The amplified fragments containing homologous arms and mutation sites were transformed into the Top10 competent cells (Tsingke) and sequenced (Tsingke). HEK 293T cells were transfected with sequencing validated mutants and were prepared following the standard EM sample preparation procedure. Cell slices with a thickness of 100 nm were imaged under a homemade wide field fluorescence microscope (Olympus IX71 body and Olympus PLAN APO 100×, 1.49 NA oil objective) to characterize the photoswitching properties of different mutants. mEosEM-E was identified for its highest photoswitching contrast ratio.

Plasmid construction

For prokaryotic expression plasmids pRsetA-RFPs (mScarlet, mScarlet-I, mScarlet-H, mKate2, FusionRed-MQV, mCherry2, mRuby3, mApple, tdTomato), RFPs fragments were PCR amplified and digested with BamHI and EcoRI restriction enzymes. Then fragments were ligated into the pRsetA vector digested with the same enzymes. For eukaryotic expression plasmids, pmEosEM-N1-mito was digested with AgeI and NotI restriction enzymes to replace mEosEM with mEosEM-E or RFPs fragments with the same restriction enzyme sticky ends. pEGFP-C1-Sec61, pmEos3.2-N1-H2B, and pmEosEM -C1-LaminA plasmids were digested with AgeI/BglII, XhoI/NotI, and NheI/BglII restriction enzymes respectively to replace EGFP, mEos3.2 and mEosEM with Scarlett-H fragment. For the construction of pmEosEM-C1-B23, pmEosEM-C1-Nop52, and pmScarlet-H-C1-Nopp140 plasmids, the full-length cDNA of B23, Nop52 and Nopp140 were PCR amplified from the HEK 293T cDNA library, digested with EcoRI/SalI, HindIII/SalI, and BglII/SalI restriction enzymes and inserted into pEGFP-C1-mEosEM-LaminA and pEGFP-C1-mScarlet-H-LaminA plasmids to replace LaminA. Q5 polymerase and T4 ligase were purchased from New England Biolabs. All restriction enzymes were purchased from Thermo Fisher Scientific.

Cell culture and transfection

HEK 293T and U-2 OS cells were cultured in Dulbecco's Modified Eagle Medium (Gibco) and McCoy's 5A Modified Medium (Gibco), respectively, supplemented with 10% FBS (Gibco) and 1% penicillin-streptomycin (TransGene Biotech). Cells were maintained at 37°C in an incubator supplied with 5% CO₂ (vol/vol). Transfections were performed with purified plasmids using lipofectamine 2000 (Invitrogen)

(with a ratio of 1 µg DNA: 3 µl lipofectamine) in 6 cm pertri dishes or 12-well cell culture plates following the manufacturer's instructions.

EM sample preparation

Successfully transfected cells were trypsinized and then harvested by centrifugation. Cells were fixed with 2% paraformaldehyde (Electron Microscopy Sciences) and 2.5% glutaraldehyde (Electron Microscopy Sciences) in 100 mM PBS at 4°C overnight. Pre-fixed cells were washed 2 times with PBS buffer and 2 times with Milli-Q water on ice. Then post-fixation with 1% OsO₄ was performed on ice for 1 h. After 4 washes with Milli-Q water, cells were gradually dehydrated with a series of concentration gradients of ethanol (30% 50% 70% 90% 100%) for 6 min each and followed by dehydration with 100% acetone for 6 min. Cells were then infiltrated step by step in mixtures of Epon (50% Epon812, 30.5% NMA, 18% DDSA, 1.5% DMP-30) and acetone with gradient concentrations (50% Epon for 2 h, 75% Epon for 3 h, 100% Epon for 12 h, 100% Epon for 12 h). Finally, cells were embedded in 100% Epon at 60°C for 16 h and then sectioned into 100-nm slices for subsequent experiments.

Photoswitching property analysis

HEK 293T cells expressing mitochondria targeted mEosEM and mutants were used for photoswitching property analysis. For the analysis before EM sample preparation, cells were excited by continuous 488-nm laser (0.41 kW/cm²), while every 10 s, a 405-nm laser (0.21 kW/cm²) pulse of 0.1 s were added to turn on the FPs. Six cycles were recorded for all FPs. For analysis after EM sample preparation, 100-nm cell sections were imaged under a continuous 488-nm laser for 50 frames, after which the 405-nm laser was added for 1 s to record the fluorescence signal of the FPs at the on-state. The contrast ratio was calculated as follow:

$$\text{Contrast ratio} = (\text{Max} - \text{Mean}) / \text{Mean}$$

Max represents the maximum value of the signal in all acquired images, Mean represents the averaged signal value of the first 20 frames acquired before the application of 405-nm laser.

High-content analysis of pre-fixation, post-fixation, and dehydration resistance of RFPs

U-2 OS cells expressing pRFPs-mito were seeded into 96-well optical polymer base microplates and fixed with 2% paraformaldehyde (Electron Microscopy Sciences) and 2.5% glutaraldehyde (Electron Microscopy Sciences) in 100 mM PBS at 37°C for 15min. The pre-fixed cells were washed 4 times with PBS and then post fixed with 1% OsO₄. After 10 min incubation at 4°C, OsO₄ was removed and cells were washed 5 times with PBS and stored in PBS. For dehydration, PBS buffer was replaced with absolute ethanol for 20 min without washing. Fluorescence images were acquired by the Opera PhenixTM High Content Screening System (PerkinElmer) using a 20×, 0.4 NA water objective with an excitation laser of 568 nm. Data quantification and analysis were performed using Harmony 4.9 software (PerkinElmer).

Protein purification and thermostability measurement

BL21(DE3) competent cells (Tsingke) were transformed with prokaryotic expression plasmids pRsetA-RFPs and single clones were cultured in liquid LB medium to the logarithmic growth phase. Then 0.8 mM IPTG (Isopropyl β- d-1-thiogalactopyranoside) was added to induce the expression of RFPs. After induction at 16 °C for 24 h, cells were harvested by centrifugation. The pelleted bacteria were resuspended in binding buffer including 10 mM imidazole and lysed by ultrasonication. Protein was purified through affinity chromatography (Ni-NTA His-Bind resin, Qiagen), followed by gel filtration chromatography (Superdex 200 Increase 10/300 GL, GE Healthcare). Purified protein was diluted in PBS buffer (pH = 7.2) and the fluorescence intensity was recorded at 60 °C in the Rotor-Gene 6600 real-

time PCR cycler (Qiagen) for 18 h. The thermostability was defined as the ratio between the initial and the final fluorescence intensities.

Photostability measurement

HEK 293T cell samples expressing mitochondria targeted RFPs were prepared as described above for EM and sectioned into 100 nm slices. Sample slices were illuminated with a 561-nm laser and the fluorescence signal was acquired by time-lapse imaging. Photostability was defined as the time when fluorescence intensity reached 1/e of its initial.

CLEM imaging

Cleaning and coating of the coverslips with pioloform (Ted Pella) were processed as previously reported [31]. Sectioned slices were placed on well prepared coverslips and submerged with mowiol buffer to recover the fluorescence. After a 30 min incubation, slices were imaged using a widefield fluorescence microscope (Olympus IX71) equipped with a 100×, 1.49 NA oil objective (Olympus PLAN APO). Images were acquired using an electron-multiplying charge-coupled device (EMCCD) camera (Andor iXon DV-897 BV). For the red channel, 100 frames were acquired for averaging during excitation by a 561-nm laser. For the green channel, 20 frames were first acquired using 488-nm laser excitation alone, then another 20 frames were acquired while a 405-nm laser pulse of 1 s was added to switch on the mEosEM-E molecules. For both channels, the exposure time was 50 ms. After fluorescence signal recording, DIC images of 100×, 16×, and 10× magnifications were sequentially collected to assist the target retrieving during subsequent EM imaging. After LM imaging, a rectangle on the pioloform film with a slice on it was scored by a knife. 12% hydrofluoric acid was dropped on the periphery of the rectangle to detach the pioloform film from the glass coverslip. When the coverslip was submerged under water, the detached pioloform film would float on the surface and was captured by an uncoated slot grid. Next, the section slice was stained with 2% UA and 1% Sato's triple lead. Finally, the sample was imaged under TECNAI SPIRIT TEM (FEI). Gold nanoparticles were used as the fiducial marker. FM and EM images were correlated by eC-CLEM following the previous protocol [16].

Registration of green and red channel FM images

TetraSpeck™ Microspheres (ThermoFisher Scientific) were diluted in PBS buffer and spotted on a coverslip (Fisher Scientific). Dual-color fluorescence signals were recorded simultaneously under 488- and 561-nm lasers. The registration was performed with the Fiji plugin "Multi Registration" in ImageJ.

Funding

The project was funded by National Key Research and Development Program of China (2017YFA0505300); National Natural Science Foundation of China (32027901, 31870857, 21778069, 21927813); The Strategic Priority Research Program of Chinese Academy of Sciences (XDB37040301).

Acknowledgments

We are grateful to Ya Wang from Research Platform for Protein Sciences at Institute of Biophysics for her support of Opera Phenix™ High Content Screening system data analysis. We thank Center for Biological Imaging at Institute of Biophysics for EM imaging support.

Author contributions

P.X., M.Z., D.P., and N.L. designed the experiments. D.P., N.L., and W.H. performed the experiments. P.X., M.Z., D.P., and N.L. analyzed the data. P.X., M.Z., and T.X. supervised the project and wrote the manuscript. K.D provided guidance on the project.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Wells, B. and L.L. Cour, *A technique for studying one and the same section of a cell in sequence with the light and electron microscope*. Journal of microscopy, 1971. **93**(1): p. 43-48.
- Hoffman, D.P., et al., *Correlative three-dimensional super-resolution and block-face electron microscopy of whole vitreously frozen cells*. Science, 2020. **367**(6475).
- Bykov, Y.S., et al., *Correlative light and electron microscopy methods for the study of virus–cell interactions*. FEBS letters, 2016. **590**(13): p. 1877-1895.
- Fang, T., et al., *Nanobody immunostaining for correlated light and electron microscopy with preservation of ultrastructure*. Nature methods, 2018. **15**(12): p. 1029-1032.
- Rae, J., et al., *A robust method for particulate detection of a genetic tag for 3D electron microscopy*. Elife, 2021. **10**: p. e64630.
- Lam, S.S., et al., *Directed evolution of APEX2 for electron microscopy and proximity labeling*. Nature methods, 2015. **12**(1): p. 51-54.
- Shu, X., et al., *A genetically encoded tag for correlated light and electron microscopy of intact cells, tissues, and organisms*. PLoS biology, 2011. **9**(4): p. e1001041.
- Risco, C., et al., *Specific, sensitive, high-resolution detection of protein molecules in eukaryotic cells using metal-tagging transmission electron microscopy*. Structure, 2012. **20**(5): p. 759-766.
- Tabata, S., et al., *Electron Microscopic Detection of Single Membrane Proteins by a Specific Chemical Labeling*. Iscience, 2019. **22**: p. 256-268.
- Polishchuk, R.S., E.V. Polishchuk, and A. Luini, *Visualizing live dynamics and ultrastructure of intracellular organelles with preembedding correlative light-electron microscopy*. Methods in cell biology, 2012. **111**: p. 21-35.
- Polishchuk, R.S., et al., *Correlative light-electron microscopy reveals the tubular-saccular ultrastructure of carriers operating between Golgi apparatus and plasma membrane*. The Journal of cell biology, 2000. **148**(1): p. 45-58.
- Kukulski, W., et al., *Correlated fluorescence and 3D electron microscopy with high sensitivity and spatial precision*. Journal of Cell Biology, 2011. **192**(1): p. 111-119.
- Watanabe, S., et al., *Protein localization in electron micrographs using fluorescence nanoscopy*. Nature methods, 2011. **8**(1): p. 80-84.
- Paez-Segala, M.G., et al., *Fixation-resistant photoactivatable fluorescent proteins for CLEM*. Nature methods, 2015. **12**(3): p. 215-218.
- Luft, J.H., *Improvements in epoxy resin embedding methods*. The Journal of biophysical and biochemical cytology, 1961. **9**(2): p. 409-414.
- Fu, Z., et al., *mEosEM withstands osmium staining and Epon embedding for super-resolution CLEM*. Nature methods, 2020. **17**(1): p. 55-58.
- Tanida, I., et al., *Visualization of cytoplasmic organelles via in-resin CLEM using an osmium-*

- 519 *resistant far-red protein*. Scientific reports, 2020. **10**(1): p. 1-7.
- 520 18. Tanida, I., et al., *Two-color in-resin CLEM of Epon-embedded cells using osmium resistant green*
521 *and red fluorescent proteins*. Scientific reports, 2020. **10**(1): p. 1-8.
- 522 19. Marriott, G., et al., *Optical lock-in detection imaging microscopy for contrast-enhanced imaging*
523 *in living cells*. Proceedings of the National Academy of Sciences, 2008. **105**(46): p. 17789-17794.
- 524 20. Chen, Y.-C., et al., *Optically modulated photoswitchable fluorescent proteins yield improved*
525 *biological imaging sensitivity*. Journal of the American Chemical Society, 2015. **137**(40): p.
526 12764-12767.
- 527 21. Chen, Y.-C., et al., *Facile autofluorescence suppression enabling tracking of single viruses in live*
528 *cells*. Journal of Biological Chemistry, 2019. **294**(50): p. 19111-19118.
- 529 22. Querard, J., et al., *Photoswitching Kinetics and Phase - Sensitive Detection Add Discriminative*
530 *Dimensions for Selective Fluorescence Imaging*. Angewandte Chemie International Edition,
531 2015. **54**(9): p. 2633-2637.
- 532 23. Chang, H., et al., *A unique series of reversibly switchable fluorescent proteins with beneficial*
533 *properties for various applications*. Proceedings of the National Academy of Sciences, 2012.
534 **109**(12): p. 4455-4460.
- 535 24. Helmerich, D.A., et al., *Photobluening of organic dyes can cause artifacts in super-resolution*
536 *microscopy*. Nature Methods, 2021. **18**(3): p. 253-257.
- 537 25. Kremers, G.-J., et al., *Photoconversion in orange and red fluorescent proteins*. Nature methods,
538 2009. **6**(5): p. 355-358.
- 539 26. Protasova, E.A., et al., *Chromophore reduction plus reversible photobleaching: how the mKate2*
540 *"photoconversion" works*. Photochemical & Photobiological Sciences, 2021: p. 1-13.
- 541 27. Farley, K.I., et al., *Determinants of mammalian nucleolar architecture*. Chromosoma, 2015.
542 **124**(3): p. 323-331.
- 543 28. Chen, H.-K., et al., *Human Nopp140, which interacts with RNA polymerase I: implications for*
544 *rRNA gene transcription and nucleolar structural organization*. Molecular and cellular biology,
545 1999. **19**(12): p. 8536-8546.
- 546 29. Okuwaki, M., *The structure and functions of NPM1/Nucleophsmin/B23, a multifunctional*
547 *nucleolar acidic protein*. Journal of biochemistry, 2008. **143**(4): p. 441-448.
- 548 30. Hernandez-Verdun, D., *Nucleolus: from structure to dynamics*. Histochemistry and cell biology,
549 2006. **125**(1): p. 127-137.
- 550 31. Xu, P., et al., *A protocol for Epon-embedding-based correlative super-resolution light and*
551 *electron microscopy*. 2019.

552

553

554 Supplementary Materials

Figure S1	Autofluorescence of Epon resin slice
Figure S2	Photoswitching properties of mEosEM and mutants before EM sample preparation
Figure S3	Scheme of sLM imaging of EM sample
Figure S4	SBR comparison of mEosEM and mEosEM-E in OFF- and ON-images
Figure S5	Pre-fixation influence on RFPs
Figure S6	Fluorescence intensity of RFPs in pre-fixed cells followed by dehydration
Figure S7	SBR comparison of RFPs after EM sample preparation
Figure S8	Green fluorescence emission properties of RFPs
Figure S9	Dual-color post-Epon-embedding CLEM imaging of Nopp52 and Nopp140
Table S1	Statistics for contrast ratio comparison between mEosEM and mEosEM-E
Table S2	Statistics for SBR of mEosEM in OFF, ON, and ON — OFF images
Table S3	Statistics for SBR of mEosEM-E in OFF, ON, and ON — OFF images
Table S4	Statistics for SBR comparison between mEosEM and mEosEM-E in ON — OFF images
Table S5	Statistics for fluorescence intensity comparison of RFPs after 1% OsO ₄ post-fixation
Table S6	Statistics for fluorescence intensity comparison of RFPs after 1% OsO ₄ post-fixation and dehydration treatment
Table S7	Statistics for thermostability comparison of RFPs
Table S8	Statistics for fluorescence intensity comparison of RFPs after EM sample preparation
Table S9	Statistics for photostability comparison of RFPs
Table S10	Statistics for SBR comparison between mEosEM and mEosEM-E in OFF images
Table S11	Statistics for SBR comparison between mEosEM and mEosEM-E in ON images
Table S12	Statistics for fluorescence intensity comparison of RFPs in live cells
Table S13	Statistics for fluorescence intensity comparison of RFPs after pre-fixation
Table S14	Statistics for fluorescence intensity comparison of RFPs after dehydration treatment
Table S15	Statistics for SBR comparison of RFPs after EM sample preparation

Figure S1. Autofluorescence of Epon resin slice. (A) A corner of Epon slice was imaged under 488-nm laser (0.41 kW/cm²). (B) Fluorescence intensity profile of the yellow line in (A). Fluorescence intensity was plotted against distance.

Figure S2. Photoswitching properties of mEosEM and mutants before EM sample preparation. (A-Q) Normalized fluorescence intensity of mEosEM and mutants was plotted against time. Cells expressing different fluorescent proteins were continuously illuminated with a 488-nm laser (8 W/cm²), while every 10 s, a 405-nm laser pulse (0.1 s, 9 W/cm²) were applied to switch on the FPs. Exposure time, 50 ms.

Figure S3. Scheme of sLM imaging of EM sample. 100-nm cell sections were imaged under a continuous 488-nm laser for 50 frames, after which the 405-nm laser was added for 20 frames to record the fluorescence signal of the FPs at the on-state. The 20 frames immediately before 405-nm illumination were averaged to produce OFF images, and the subsequent 20 frames with 405-nm illumination were averaged to produce ON images. The sLM images (ON — OFF images) were generated by pixel-by-pixel subtraction using adjacent ON and OFF images. Scale bar, 5 μ m.

Figure S4. SBR comparison of mEosEM and mEosEM-E in OFF- and ON-images. **(A)** Statistics of SBR in OFF-images between mEosEM and mEosEM-E. **(B)** Statistics of SBR in ON-images between mEosEM and mEosEM-E. Bars represent mean ± SD. P-value were determined with two-tailed t-test in **(A-B)** (n = 9). n.s. indicates p>0.05, **** indicates p<0.0001. Data are summarized in Table S10 & 11.

Figure S5. Pre-fixation influence on RFPs. **(A)** Fluorescence intensity of RFPs in live cells. Cells expressing RFPs were seeded in 96-well plates and imaged. **(B)** Fluorescence intensity of RFPs in pre-fixed cells. Cells expressing RFPs were seeded in 96-well plates and treated with fixation buffer (2.5% Glutaraldehyde and 2% Paraformaldehyde in PBS) for 15 min then washed 3 times with PBS buffer. Data were recorded using a high-content screening system. Excitation wavelength, 561 nm. Exposure time, 40 ms. Bars represent mean $\pm$ SD (n = 3). Data are summarized in Table S12 & 13.

Figure S6. Fluorescence intensity of RFPs in pre-fixed cells followed by dehydration. Cells expressing RFPs were seeded in 96-well plates and fixed with fixation buffer (2.5% Glutaraldehyde and 2% Paraformaldehyde in PBS buffer) for 15 min, and then treated with absolute ethanol for 20 min. Data were recorded using a high-content screening system. Excitation wavelength, 561 nm. Exposure time, 40 ms. Bars represent mean $\pm$ SD. P-value were determined with two-tailed t-tests ($n = 3$). * indicates $p < 0.05$. ** indicates $p < 0.01$. *** indicates $p < 0.001$. **** indicates $p < 0.0001$. Data are summarized in Table S14.

Figure S7. SBR comparison of RFPs after EM sample preparation. HEK 293T cells expressing RFPs were prepared under standard EM sample preparation and sectioned into 100 nm slices. Mann-Whitney U test was performed (n = 106). n.s. indicates $p > 0.05$, **** indicates $p < 0.0001$. Data are summarized in Table S15.

Figure S8. Green fluorescence emission properties of RFPs. HEK 293T cells expressing RFPs were prepared under standard EM sample preparation and sectioned into 100 nm slices. Sample slices were imaged by wide field fluorescence microscopy with sequential illumination of 561- and 488-nm laser. (A), (C), (E), (G), (I), (K) and (M) Slices were illuminated under 488-nm laser (0.92 kW/cm²) and signals were recorded from the green channel. (B), (D), (F), (H), (J), (L) and (N) Slices were illuminated under 561-nm laser (0.57 kW/cm²) and signals were recorded from the red channel. Scale bars, 10 μ m.

Figure S9. Dual-color post-Epon-embedding CLEM imaging of Nopp52 and Nopp140. (A-D) Dual-color CLEM imaging of mEosEM-E labeled Nopp52 and mScarlet-H labeled Nopp140 protein in HEK 293T cell sections (100 nm). Green channel FM (A), red channel FM (B), EM (C), and CLEM (D) images. Scale bar, 2 μ m. (E-F) Enlarged EM (E) and red channel CLEM (F) images of boxed area in (D). Gamma value: 1.6 for both channels. Scale bar, 1 μ m.

	mEosEM	mEosEM-E
Contrast Ratio	2.147	4.660
	2.513	7.366
	2.216	7.561
	2.256	7.351
	2.132	4.959
	2.233	4.190
	2.404	4.832
	2.289	6.791
	1.597	6.082
mean	2.199	5.977
s.d.	0.256	1.336
P value	2.13E-05	

Table S1. Statistics for contrast ratio comparison between mEosEM and mEosEM-E. Two-tailed t-test was performed between mEosEM and mEosEM-E, n = 9.

	mEosEM (OFF)	mEosEM (ON)	mEosEM (ON-OFF)
Signal-to-Background Ratio	2.130	3.684	12.289
	2.812	5.321	12.680
	1.921	3.336	11.894
	1.716	2.604	6.375
	1.416	2.035	5.226
	1.746	2.792	9.019
	1.478	2.304	9.097
	1.524	2.251	6.370
	1.593	2.039	3.676
mean	1.815	2.930	8.514
s.d.	0.437	1.062	3.298
P value			0.000762

Table S2. Statistics for SBR of mEosEM in OFF, ON, and ON — OFF images. Two-tailed t-test was performed between mEosEM (ON) and mEosEM (ON — OFF), n = 9.

	mEosEM-E		mEosEM-E
	(OFF)	mEosEM-E (ON)	(ON-OFF)
Signal-to-Background Ratio	2.086	5.785	28.925
	1.651	4.731	16.408
	1.931	7.015	34.881
	2.020	7.132	27.124
	1.899	5.406	38.488
	2.978	8.000	22.849
	2.263	6.488	28.004
	1.966	6.808	37.050
	2.061	6.655	31.652
mean	2.095	6.447	29.487
s.d.	0.370	0.990	7.029
P value			7.87E-06

628 **Table S3.** Statistics for SBR of mEosEM-E in OFF, ON, and ON — OFF images. Two-tailed t-test was
629 performed between mEosEM-E (ON) and mEosEM-E (ON — OFF), n = 9.
630

	mEosEM	mEosEM-E
Signal-to-Background Ratio	12.289	28.925
	12.680	16.408
	11.894	34.881
	6.375	27.124
	5.226	38.488
	9.019	22.849
	9.097	28.004
	6.370	37.050
	3.676	31.652
mean	8.514	29.487
s.d.	3.298	7.029
P value	4.70E-06	

Table S4. Statistics for SBR comparison between mEosEM and mEosEM-E in ON — OFF images. Two-tailed t-test was performed between mEosEM and mEosEM-E, n = 9.

634

	mScarlet	mScarlet-I	mScarlet-H	mKate2	FusionRed-MQV	mCherry2	mRuby3	mApple	tdTomato
Fluorescence									
Intensity	741.800	791.900	609.200	322.500	445.800	552.600	743.900	311.400	482.700
	712.100	764.000	592.400	320.800	440.100	505.400	738.000	304.400	495.900
	724.400	833.300	625.800	330.300	438.100	519.300	776.600	310.600	477.000
Mean	726.100	796.400	609.133	324.533	441.333	525.767	752.833	308.800	485.200
s.d.	14.923	34.868	16.700	5.066	3.995	24.255	20.793	3.831	9.695
P value									
(compared to									
mKate2)	0.000127	0.00150	0.000478				0.000423		
P value									
(compared to									
mCherry2)	0.000714	0.000695	0.0108				0.000286		

635 **Table S5.** Statistics for fluorescence intensity comparison of RFPs after 1% OsO₄ post-fixation. Two-
636 tailed t-tests were performed between mKate2 and mScarlet, mScarlet-I, mScarlet-H, mRuby3,
637 respectively, n = 3. Two-tailed t-tests were performed between mCherry2 and mScarlet, mScarlet-I,
638 mScarlet-H, mRuby3, respectively, n = 3.
639

	mScarlet	mScarlet-I	mScarlet-H	mKate2	FusionRed-MQV	mCherry2	mRuby3	mApple	tdTomato
Fluorescence									
Intensity	321.400	277.300	314.200	298.100	314.400	332.200	395.900	317.100	325.000
	296.800	281.100	319.100	297.800	272.400	308.000	385.100	312.300	325.200
	333.600	285.700	309.500	305.300	283.900	297.300	407.500	309.300	295.700
Mean	317.267	281.367	314.267	300.400	290.233	312.500	396.167	312.900	315.300
s.d.	18.745	4.206	4.800	4.246	21.704	17.880	11.202	3.934	16.974
P value	0.00637	0.00114	0.00216	0.00177	0.00492	0.00435		0.00282	0.00388

Table S6. Statistics for fluorescence intensity comparison of RFPs after 1% OsO₄ post-fixation and dehydration treatment. Two-tailed t-tests were performed between mRuby3 and other RFPs, n = 3.

	mScarlet	mScarlet-L	mScarlet-H	mKate2	FusionRed-MQV	mCherry2	mRuby3	mApple	tdTomato
Thermostability									
Ratio	0.840	0.682	0.994	0.071	0.022	0.214	0.852	0.550	0.794
	0.842	0.742	0.959	0.071	0.021	0.206	0.856	0.575	0.728
	0.856	0.750	0.975	0.066	0.021	0.205	0.822	0.603	0.761
Mean	0.846	0.725	0.976	0.069	0.021	0.208	0.843	0.576	0.761
s.d.	0.009	0.037	0.017	0.003	0.001	0.005	0.018	0.026	0.033
P value	0.00151	0.00223		8.67E-05	0.000109	5.71E-05	0.000840	7.76E-05	0.00199

643 **Table S7.** Statistics for thermostability comparison of RFPs. Two-tailed t-tests were performed
644 between mScarlet-H and other RFPs, n = 3.
645

	mScarlet	mScarlet-I	mScarlet-H	mKate2	FusionRed-MQV	mCherry2	mRuby3
Mean	238.979	283.083	327.015	210.963	153.836	136.175	158.454
s.d.	135.404	123.250	178.850	107.039	58.967	48.306	77.657
P value	0.000005	0.144		1.78E-09	3.23E-21	1.89E-25	1.19E-18

646 **Table S8.** Statistics for fluorescence intensity comparison of RFPs after EM sample preparation. Mann-
647 Whitney U tests were performed between mScarlet-H and each RFP, n = 106.
648

	mScarlet	mScarlet-I	mScarlet-H	mKate2	FusionRed-MQV	mCherry2	mRuby3
t1/e	341.040	359.090	476.280	366.780	269.560	307.030	636.990
	377.370	389.060	389.790	424.040	345.240	307.270	471.980
	351.770	353.860	452.630	426.160	273.080	327.990	527.020
	300.120	310.990	436.420	371.070	275.120	389.660	521.720
	380.550	433.362	440.220	290.670	253.760	381.610	482.540
Mean	344.147	366.071	443.090	362.633	267.320	366.420	510.427
s.d.	32.610	45.379	31.644	55.259	35.600	40.197	65.429
P value	0.0470	0.162		0.176	6.37E-05	0.0508	0.0299

649 **Table S9.** Statistics for photostability comparison of RFPs. Two-tailed t-tests were performed between
650 mScarlet-H and other RFPs, n = 3.
651

652

	mEosEM-OFF	mEosEM-E-OFF
Contrast Ratio	2.130	2.086
	2.812	1.651
	1.921	1.931
	1.716	2.020
	1.416	1.899
	1.746	2.978
	1.478	2.263
	1.524	1.966
	1.593	2.061
mean	1.815	2.095
s.d.	0.436	0.370
P value	0.162	

653 **Table S10.** Statistics for SBR comparison between mEosEM and mEosEM-E in OFF images. Two-
654 tailed t-test was performed between mEosEM and mEosEM-E, n = 9.

655

	mEosEM-ON	mEosEM-E-ON
Contrast Ratio	3.684	5.785
	5.321	4.731
	3.336	7.015
	2.604	7.132
	2.035	5.406
	2.792	8.000
	2.304	6.488
	2.251	6.808
	2.039	6.655
mean	2.930	6.447
s.d.	1.062	0.990
P value	1.94E-06	

Table S11. Statistics for SBR comparison between mEosEM and mEosEM-E in ON images. Two-tailed t-test was performed between mEosEM and mEosEM-E, n = 9.

	mScarlet	mScarlet-l	mScarlet-H	mKate2	FusionRed-MQV	mCherry2	mRuby3	mApple	tdTomato
Fluorescence									
Intensity	2928	3256	1453	1406	1568	1752	3011	3364	4674
	2906	3269	1457	1344	1763	1715	3100	3423	4637
	2887	3449	1348	1401	1719	1781	3104	3465	4643
Mean	2907	3324.667	1419.333	1383.667	1683.333	1749.333	3071.667	3417.333	4651.333
s.d.	20.51828	107.8718	61.80885	34.44319	102.2758	33.08071	52.57693	50.73789	19.85783

Table S12. Statistics for fluorescence intensity comparison of RFPs in live cells. n = 3.

659
660

	mScarlet	mScarlet-l	mScarlet-H	mKate2	FusionRed-MQV	mCherry2	mRuby3	mApple	tdTomato
Fluorescence									
Intensity	1490	1819	955.6	651.3	1117	974.2	1995	703.9	1860
	1483	1903	966.4	639.8	1228	1019	2198	715.2	2041
	1450	1841	924.8	642.8	1178	1041	2000	803.6	1745
Mean	1474.333	1854.333	948.9333	644.6333	1174.333	1011.4	2064.333	740.9	1882
s.d.	21.36196	43.55839	21.58642	5.965177	55.59077	34.04233	115.7857	54.59295	149.2213

Table S13. Statistics for fluorescence intensity comparison of RFPs after pre-fixation. n = 3.

661
662

	mScarlet	mScarlet-l	mScarlet-H	mKate2	FusionRed-MQV	mCherry2	mRuby3	mApple	tdTomato
Fluorescence									
Intensity	429.100	583.500	648.900	685.600	518.800	938.800	1553.000	1153.000	680.100
	442.600	491.500	604.900	681.800	487.400	878.500	1615.000	917.600	650.000
	440.500	780.200	639.500	665.700	483.300	823.900	1525.000	865.900	549.900
Mean	437.400	618.400	631.100	677.700	496.500	880.400	1564.333	978.833	626.667
s.d.	7.264	147.480	23.172	10.565	19.421	57.474	46.058	153.032	68.164
P value	0.000420	0.00461	8.06E-05	0.000526	0.000102	0.000119		0.0157	0.000101

Table S14. Statistics for fluorescence intensity comparison of RFPs after dehydration treatment. Two-tailed t-tests were performed between mRuby3 and other RFPs, n = 3.

	mScarlet	mScarlet-I	mScarlet-H	mKate2	FusionRed-MQV	mCherry2	mRuby3
Mean	1.791	1.910	2.034	1.706	1.524	1.456	1.541
s.d.	0.385	0.335	0.495	0.307	0.181	0.147	0.237
P value	0.000009	0.182		6.14E-09	8.95E-22	6.34E-27	1.64E-18

Table S15. Statistics for SBR comparison of RFPs after EM sample preparation. Mann-Whitney U tests were performed between mScarlet-H and each RFP, n = 106.