Locally adaptive temperature response of vegetative growth in *Arabidopsis thaliana*

Pieter Clauw¹✉, Envel Kerdaffrec², Joanna Gunis¹, Ilka Reichardt³, Viktoria Nizhynska¹, Stefanie Koemeda⁴, Jakub Jez⁴, and Magnus Nordborg¹✉

¹Gregor Mendel Institute of Molecular Plant Biology, Austrian Academy of Sciences, Vienna BioCenter (VBC), 1030 Vienna, Austria

²Department of Biology, University of Fribourg, CH-1700 Fribourg, Switzerland

³Max Planck Institute of Molecular Cell Biology and Genetics, Dresden, 01307 Dresden, Germany

⁴Plant Sciences Facility, Vienna BioCenter Core Facilities GmbH (VBCF), 1030 Vienna, Austria

We investigated early vegetative growth of natural *Arabidopsis thaliana* accessions in cold, non-freezing temperatures, similar to temperatures these plants naturally encounter in fall at northern latitudes. We found that accessions from northern latitudes produced larger seedlings than accessions from southern latitudes, partly as a result of larger seed size. However, their subsequent vegetative growth when exposed to colder temperatures was slower. The difference was too large to be explained by random population differentiation, and is thus suggestive of local adaptation, a notion that is further supported by substantial transcriptome and metabolome changes in northern accessions. We hypothesize that the reduced growth of northern accessions is an adaptive response, and a consequence of reallocating resources towards cold acclimation and winter survival.

Correspondence:

pieter.clauw@gmi.oeaw.ac.at

magnus.nordborg@gmi.oeaw.ac.at

Introduction

Local adaptation studies in *Arabidopsis thaliana* have found important roles for life history traits such as seed dormancy and flowering time (1). Temperature is a main regulator of these traits and local populations are adapted to their local temperatures (2–4). Also plant growth is affected by temperature and previous studies have detected genetic variation underlying rosette sizes and growth rates (5, 6) as well as signals of polygenic adaptation (7). How vegetative growth is adapted to local temperatures remains unclear, however.

Growth can be seen as the end-sum of a vast number of physiological processes. All of these are genetically determined but can also be heavily influenced by environmental conditions. Growth is therefore not only genetically a complex trait but also enormously plastic. The most straightforward environmental effect is when conditions are so adverse that growth reaches a physiological limit, making it impossible for the plant to grow any further. This is so-called “passive plasticity” (8, 9). Yet, when survival is at stake, it may also be in the interest of the plant to actively inhibit growth upon deteriorating environmental conditions (10) so-called “active plasticity” (8, 9). Since vegetative growth ultimately determines photosynthetic surface and thus energy input that can be invested in the next generation, it is in direct trade-off with survival. Allocation of resources towards either growth or survival is thus an important balance to keep, and plants are expected to be adapted to constantly perceiving and respond-

ing to specific environmental changes as cues for upcoming, potentially, adverse conditions.

Cold acclimation is a well-studied example of plants sensing cold temperatures as a cue for upcoming winter and consequently preparing for freezing temperatures. The increased freezing tolerance upon cold acclimations is accomplished by changing membrane composition, producing cryoprotective polypeptides such as COR15A (11, 12) and accumulating compatible solutes with cryoprotective properties such as raffinose, sucrose and proline (13–15). Main regulators of cold acclimation are CBF1/DREB1b, CBF2/DREB1c and CBF3/DREB1a, three AP2/ERF transcription factors, for which allelic variation in CBF2 has been linked to natural variation in freezing tolerance (16–18).

Here we investigated the role of growth in adaptation to cold temperatures by comparing vegetative growth of 249 accessions (Figure 1) grown in 16°C and 6°C for a period of 3 weeks following seedling establishment (Figure 2). Rosette growth of each plant was measured twice a day during temperature treatments, using automated phenotyping. The experiment generated rosette growth estimates at a high temporal resolution in two ecologically realistic temperature conditions in a wide set of accessions, allowing us to look for patterns of local adaptation.

Results

Estimating plant growth parameters.

Our experiment yielded dense (two measurements per day) time series growth data for over 7,000 individual plants (249 accessions X 2 treatments X 5 replicate accessions X 3 repeated experiments). These data were used to model plant growth and estimate growth parameters for further analysis. Unlimited growth should be exponential, but plant growth is known to slow down with increasing size, and therefore a power-law function, $\frac{dM}{dt} = rM^\beta$, with $\beta < 1$ is typically a better fit than a pure exponential function (for which $\beta = 1$ — in the equation, M is the size, r is the growth rate, and β is a scaling factor that allows rate of size increase to change with size). Growth according to a power-law function describes early stages of plant growth especially well (20) for the rosette size measurements in this study. To calculate the rosette size from a power-law function at a given time point only three parameters are required: the initial size (M_0),

Fig. 1. Geographic origin of the 249 accessions. Map color shows winter temperature (mean temperature of coldest quarter). Accessions are colored according to sub-population (19).

Fig. 2. Timeline of the experiments. Upon vernalisation, seeds germinated and seedlings established over 14 days. After 14 days plants were exposed to either 16°C treatment or 6°C treatment. Insert shows the 24 hour temperature profile for the 16°C (red) and 6°C (green) treatments, with light period indicated in yellow.

Fig. 3. Variation among accessions of initial size (M_0), growth rate (r) and the temperature response of the growth rate.

growth rate (r), and β . Note that M_0 is the rosette size at the start of the temperature treatment 14 days after stratification (Figure 2) and is thus not affected by the temperature treatment. We used a non-linear mixed model to obtain estimates for the initial size, growth rates and beta. Accession was added as fixed effect for initial size and growth rate, temperature and accession X temperature interactions were added as fixed effects for growth rate only. β was considered to be constant over accessions and temperatures. The “temperature response” of the growth rate was calculated for each accession as the slope between the growth rate at 16°C and 6°C. As expected, all accessions grew faster when it was warmer. The observed phenotypic variation (Figure 3) is to a large extent explained by genetic variation; broad-sense heritabilities are 0.41 for initial size, and 0.57 and 0.32 for growth rate at 16°C and 6°C respectively.

Growth parameters correlate with the environment of origin.

If growth rates are locally adaptive, they may reflect the envi-

ronment of origin of each accession. To investigate this, we correlated our estimated growth rates with climate data. The climate variables showing the strongest correlations with the different growth parameters were linked to temperature during winter months (Figure S1), also when correlations were corrected for population structure (Figure S2). In particular, the mean temperature during the coldest quarter (henceforth referred to as “winter temperature”) was strongly correlated with our parameter estimates, and we focus on it in what follows.

Initial size.

Accessions from colder climates generally had higher initial rosette size (M_0), 2 weeks after germination, than accessions from warmer climates, but then grew more slowly during the temperature experiment — regardless of temperature regime (Figure 4).

At least part of the explanation for this pattern is likely to be differences in seed size between accessions. Using seed size measurements from previous experiments, we found that seed size is positively correlated with initial size, and also with winter temperature (Figure S3), at least for the subset of 123 Swedish accessions. Winter temperature is still significantly associated with initial size when corrected for seed

Fig. 4. Variation among accessions of initial size (M_0), growth rate (r) and the temperature response of the growth rate.

size by adding it as a random effect. Seed size alone can therefore not explain the geographic pattern we observe for initial size and there must be a role for variation in growth rate during the very initial phases of seedling growth.

Growth rates.

While the initial sizes correlate negatively with winter temperature, we observed the opposite relation for the growth rates. Despite being larger initially, accessions from colder climates grew more slowly during both the 16°C and 6°C treatments (Figure 4). Since resources are not limiting, this suggests that the northern lines are actively inhibiting their growth, and that growing slower may be beneficial in colder climates, perhaps in preparation for winter. Accessions from colder climates were also less sensitive to the temperature experiment in the sense that the temperature response of the growth rate increased with winter temperature of origin (Figure 4).

Cold acclimation response

In agreement with the variation in growth rates, metabolite measurements taken at the final day of our experiment showed clear differences between accessions from cold and warm regions, and many of these differences involved metabolites with a known role in cold acclimation (21). Since the transcriptomic component of cold acclimation is well studied, we analyzed the expression profiles of 251 known cold acclimation genes in 8 accessions chosen to be representative in terms of their growth and metabolome profiles (Figure S4). The selected genes are known to be differentially expressed upon exposure to cold, and their expression is under control of at least one of the known transcription factors regulating cold acclimation: CBF1, CBF2, CBF3, HSFC1 (22) or ZAT12 (23). In our experiment, expression of these

Fig. 5. Expression of known cold acclimation genes. Expression is shown as the gene-wide z-scores of the normalized counts. The z-scores allow for grouping genes with a similar expression behavior over the different accessions in both temperatures. The top bar indicates winter temperature (°C) for each accession's origin. Both dendrograms along y-axis and x-axis respectively show hierarchical clustering of genes, and of accessions in both temperatures.

genes is likewise more affected by temperature than expected by chance (Figure 5; χ^2 -test: p-value < 0.001), and separates the two accessions from the coldest region (northern Sweden) from the rest in the 16°C treatment, and the three accessions from the warmest regions (Spain and Azerbaijan) from the rest in the 6°C treatment. Expression of different subsets of the selected cold-acclimation genes show clear correlations with winter temperature of origin (Figure S5). In particular, the genes that were previously found to be up-regulated upon cold exposure showed higher expression in accessions from cold climates (Figure S6). Since the expression of these cold acclimation genes is linked to the strength of cold acclimation, these accessions likely differ in their ability to cope with freezing temperatures upon cold treatment.

Growth is polygenic and shows signs of local adaptation

We used genome-wide association to investigate the genetic architecture underlying variation for the different growth parameters (Figure S7). The most significant association was found for overall growth rate in 16°C (Figure 6). Inflated significance levels after correcting for population structure are consistent with what we would expect from a polygenic trait (Figure 6) and were also observed for the other traits, except for growth rate in 6°C (Figure S7). Plausible candidates within 10kb of the most significant SNP (chr5:23,334,281) include *CIPK21* and *MYB36*. *CIPK21* encodes a CBL-

interacting protein kinase that is upregulated in cold conditions and is involved in the salt and osmotic stress response (24). MYB36 is a key regulator of root endodermal differentiation (25). Slightly more distant, 22kb away, is *COL5*, a transcription factor that is part of the gene network that is regulated by AN3, a regulator of cell proliferation in leaf growth (26).

To test for potential polygenic adaptation, we compared the phenotypic divergence to the expected neutral genome-wide genetic divergence. This can be done using a so-called $Q_{ST} - F_{ST}$ test (27–29), however this test is not well suited for species with complex population structure, and so we used a variation that was designed to detect adaptive differentiation for traits measured in structured GWAS panels (30). Instead of looking at divergences between predefined populations, this method uses principal components (PCs) of the genetic relatedness matrix as axes of potential adaptive differentiation. Adaptive differentiation is then detected as a correlation between the focal phenotype and any of these relatedness PCs that is significantly different than expected under neutrality.

Adaptive differentiation was detected for initial size and for growth rate in 16°C and its temperature response. Both traits show adaptive differentiation along different genetic axes (Figure 7). Initial size shows significant adaptive differentiation along PC6 (p-value < 0.05), whereas growth rate in 16°C and its temperature response showed significant adaptive differentiation along PC5 (p-values < 0.05). The adaptive differentiation for initial size along PC6 seems to stem from higher initial sizes in Swedish accessions, compared to central European accessions. The adaptive differentiation along PC5 seems to be driven by the lower growth rate temperature responses in Asian and Northern Swedish accessions in contrast to higher growth rates in a subset of southern Swedish accessions. The accessions in our set that come from Northern Sweden and Asia hail from the coldest climates. Thus these results suggest adaptive differentiation driven by adaptation to cold winters.

Discussion

This study explores the natural variation of rosette growth in non-freezing temperatures. Despite high plasticity, we detect genetic variation for the different growth parameters, and environmental correlations suggest local adaptation. GWAS analyses reveal, not surprisingly, a polygenic trait architecture. Indeed, across the genome we detect adaptive differentiation for certain growth parameters. We speculated that the slower growth measured in accessions from colder climates reflect relocation of resources from growth towards cold acclimation. Both metabolome and gene expression data are consistent with accessions from colder climates preparing for a harsh winter. In our temperature experiment, we see that the growth of northern lines is affected less than southern lines by switching from 16°C to 6°C.

Our conclusion that slower growth is likely adaptive in populations facing fiercer winters is in line with recent results of Wieters et al. (7), who concluded that the reduced growth

in Northern lines was adaptive and not a consequence of an accumulation of deleterious mutations at the species border. If slower growth were indeed a consequence of accumulated deleterious mutations we would expect to see slower growth also during the initial seedling establishment. On the contrary, we saw a fast seedling establishment for accessions from colder regions. We speculate that the fast seedling establishment is a potential adaptation for short growth seasons, which often coincide with colder climates (high latitude or high altitude). This fast seedling establishment seems to be partly supported by larger seeds. These larger seeds may provide more nutrients to initiate faster seedling establishment, while this is of less importance in warmer climates with longer growth seasons. Further work is needed to disentangle initial growth from seed size effects and confirm that there is a causal relationship between seed size and fast seedling establishment, whether this is due to seed nutrient storage, and whether it is adaptive.

The adaptation of growth to local climates is likely to be influenced by a trade-off with cold acclimation. General growth-survival trade-offs have long been observed and are described in general ecological strategy schemes such as Grime's C-S-R triangle (31) and the leaf-height-seed scheme (32). Specific trade-offs between growth and cold/frost survival were observed for wheat (33, 34), alfalfa (35), *Dactylis glomerata* (36), and multiple tree species (37–40). Here we observed higher expression of genes involved in cold acclimation in accessions from colder regions. Even though this is based on a limited set of 8 accessions, metabolome measurements in all 249 accessions lead to the same conclusion. Metabolites involved in cold acclimation such as raffinose, sucrose and proline were found in higher concentrations in accessions from colder climates (21). We believe that accessions from colder environments are relocating more energy and resources from growth towards preparations for upcoming freezing temperatures. This likely results in stronger cold acclimation and consequently increased freezing tolerance in the accessions from colder regions. Indeed, accessions originating from colder environments show increased freezing tolerance upon cold acclimation (41–44).

There is strong evidence from QTL mapping that genetic variation in the *CBF2* gene is one of the drivers for adaptation to freezing stress (16, 45). Here we looked at growth phenotypes and did not detect associations with the CBF loci. In the transcriptome analysis we did pick-up a role for CBF and other known cold-acclimation genes. The most significant locus detected in our GWAS analysis (for growth rate in 16°C) lies in the vicinity of *COL5*, a gene that is part of a leaf growth regulatory network (26) and whose expression is induced by both cold treatment and *CBF1*, *CBF2* and *CBF3* overexpression (22). In our transcriptome data, the *COL5* gene was found in cluster D, showing higher expression in accessions coming from warmer regions. It is however unclear what its exact regulatory role in growth in cold conditions might be.

In summary, we detected adaptive differentiation for growth between accessions from warm and cold climates. Our tran-

Fig. 6. GWAS results for the growth rate in 16°C. (A) Manhattan plot showing the significance of the association between the phenotype and each of the tested SNPs (MAF > 10%). The Bonferroni-corrected threshold is shown with a dashed red line. (B) Q-Q-plot showing the relation between observed and expected $-\log_{10}(\text{p-value})$ distributions. Red line shows the observed relationship. The gray line and band show the expected relationship under the null hypothesis of no differentiation between both distributions.

Fig. 7. Adaptive differentiation of initial size, growth rate at 16°C, and the temperature response of growth rate along different axes of genetic differentiation. Accessions are coloured according to their respective admixture groups, as specified in (19). The gray ribbon represents the expected correlation between phenotype and axis of genetic differentiation under neutrality. The blue line represents the observed correlation between phenotype and axis of genetic differentiation.

scriptome data and previous metabolome data suggest that resources are relocated from growth to cold acclimation in accessions from colder regions. This allows these accessions to be fully prepared for the coming of winter.

Materials and Methods

Plant growth and phenotyping.

Seeds of 249 natural accessions (Suppl. Data 1) of *Arabidopsis thaliana* described in the 1001 genomes project (19) were sown on sieved (6 mm) substrate (Einheitserde ED63). Pots were filled with 71.5 g $\pm$ 1.5 g of soil to assure homogenous packing. The prepared pots were all covered with blue mats (46) to enable a robust performance of the high-throughput image analysis algorithm. Seeds were stratified (4 days at 4°C in darkness) after which they germinated and left to grow for 2 weeks at 21°C (relative humidity: 55%; light intensity: 160 $\mu\text{mol m}^{-2} \text{s}^{-1}$; 14 h light). The temperature treatments were started by transferring the seedlings to either 6 °C or 16 °C. To simulate natural conditions temperatures fluctuated diurnally between 16-21 °C, 0.5-6 °C and 8-16 °C for the 21 °C initial growth conditions and the 6 °C and 16 °C treatments, respectively (Figure 2). Light intensity was kept constant at 160 $\mu\text{mol m}^{-2} \text{s}^{-1}$ throughout the experiment. Relative humidity was set at 55% but in colder temperatures it rose uncontrollably to maximum 95%. Daylength was 9h during the 16°C and 6°C treatments. Each temperature treatment was repeated in three independent experiments. Five replicate plants were grown for every genotype per experiment. Plants were randomly distributed across the growth

chamber with an independent randomisation pattern for each experiment. During the temperature treatments (14 DAS – 35 DAS), plants were photographed twice a day (1 hour after/before lights switched on/off), using an RGB camera (IDS uEye UI-548xRE-C; 5MP) mounted to a robotic arm. At 35 DAS, whole rosettes were harvested, immediately frozen in liquid nitrogen and stored at -80°C until further analysis. Rosette areas were extracted from the plant images using Lemnatec OS (LemnaTec GmbH, Aachen, Germany) software.

Non-linear modeling.

Non-linear modeling was used to describe plant growth in a minimum number of parameters. In a first step we constructed a simple non-linear model with plant size being explained by either the exponential (equation 1) or the power-law function (equation 2), with individual plant as a random effect for each of the model parameters; M_0 , r and. With being only present in the power-law model. Models were constructed using the nlsList and nlme functions from the nlme package for R. Exponential and power-law SelfStart functions were used from Paine et al. (20). Based on Akaike Information criterion and likelihood ratio test generated by the anova function, we decided to use the power-law model for further analyses.

$$\frac{dM}{dt} = rM \quad (1a)$$

$$M_t = M_0 e^{rt} \quad (1b)$$

$$\frac{dM}{dt} = rM^\beta \quad (2a)$$

$$M_t = (M_0^{1-\beta} + rt(1-\beta))^{1/1-\beta} \quad (2b)$$

In a second step, we constructed a model with fixed effects for the different power-law parameters. For initial size (M_0) we added accession as fixed effect. Temperature treatment only started from the initial time point onwards, and thus could not have an effect on the initial plant size. The growth rate, on the other hand, should be affected by temperature, therefore we included accession, temperature and their interaction as fixed effects for growth rate (r). No fixed effects were added for β . The idea here is that is an adjustment factor for decreasing growth rates (when $\beta < 1$) with increasing plant sizes, which is general for plant growth, or at least for our data in this case. Individuals, nested with experiment, were added as random effects for each of the model parameters. The correlation structure intrinsic to measuring the same individuals over time was accounted for by adding the first order continuous autoregressive correlation structure (corCAR1). The estimated fixed effects of this model were then used to obtain initial size estimates for each accession and growth rate estimates for each accession in both temperatures. These estimates were used for all further analyses. Exceptions to this are the broad-sense heritabilities, which were based on initial size and growth rate estimates that were obtained by including random effects, in order to get estimates for each individual plant.

Climate correlations.

The different phenotypes were correlated with each of the different (bio)climate variables downloaded from www.worldclim.org (47). Correlations were calculated as Pearson's correlations using the cor function in R (48). Population structure may confound the correlation between phenotype and climate. Therefore we included a population structure corrected phenotype-climate correlation (Figure S2). To this end we used a mixed-effects model as implemented in the lme4 function from the coxme package with phenotype as dependent variable, climate variable as fixed effects and the kinship matrix as random effect. Phenotype and climate variables were standardized, so that regression coefficients were comparable to correlation coefficients. Even though the strength and significance of the correlations weaken upon population structure correction, the growth parameters still demonstrate the same pattern of most strongly correlating with winter temperatures.

Seed size correlations.

We used the seeds produced by (49) and limited our measurements to the set of 123 Swedish accessions that overlapped with our growth dataset. After seed stratification for four days at 4°C in darkness, mother plants were grown for 8 weeks at 4°C under long-day conditions (16h light; 8h dark) to ensure proper vernalization. Temperature was raised to 21°C (light) and 16°C (dark) for flowering and seed ripen-

ing. Seeds were kept in darkness at 16°C and 30% relative humidity, from the harvest until seed size measurements. For each genotype three replicates were pooled and about 200-300 seeds were sprinkled on 12 x 12 cm square, transparent Petri dishes. Image acquisition was performed as described in (50) by scanning dishes on a cluster of eight Epson V600 scanners. The resulting 1200 dpi .tiff images were analyzed in the ImageJ software (2.1.0/1.53c). Images were converted to 8-bit binary images and thresholded with the setAutoThreshold("Defaultdark") command, and seed area was measured in squared mm by running the Analyse Particles command (inclusion parameters: size=0.04-0.25 circularity=0.70-1.00). All scripts used for image processing are available at https://github.com/vevel/seed_size.

Transcriptome profiling.

35 days after stratification, rosette tissue of all plants were harvested and flash frozen in liquid nitrogen. Random samples in each replicate and temperature were taken for 8 accessions to profile the transcriptome with RNA-sequencing. Total RNA was extracted using the KingFisher Duo Prime System (Thermo Fisher Scientific) together with a high performance RNA bead isolation kit (Molecular Biology Service, VBC Core Facilities, Vienna). To determine the quantity of RNA samples we used Fluorometer Qubit 4 (Invitrogen) and Qubit RNA BR Kit (Invitrogen). For each sample, 1 µg of total RNA was treated with the poly(A) RNA Selection Kit (Lexogen) and eluted in 12 µl of Nuclease-Free Water. Libraries were prepared according to the manufacturer's protocol in NEBNext Ultra II Directional RNA Library Prep Kit (New England BioLabs) and individually indexed with NEBNext Multiplex Oligos for Illumina (New England BioLabs). The quantity and quality of each amplified library were analyzed by using Fragment Analyzer (Agilent) and HS NGS Fragment Kit (Agilent). Libraries were sequenced with an Illumina HiSeq2500 in paired-end mode with read-length of 125bp. Sequencing was performed by the Next Generation Sequencing Facility at Vienna BioCenter Core Facilities (VBCF), member of the Vienna BioCenter (VBC), Austria. Gene expression was quantified by using quasi-mapping in salmon, version 1.2.1 (51). The salmon indices were built separately for each accession, as we incorporated the SNP variation from the (19) into the reference transcriptome.

Broad-sense heritabilities.

Broad-sense heritability (H^2) for initial size was calculated using the estimates for each individual plant, over all experiments. H^2 for growth rate was calculated separately for each temperature, again using estimates for each individual plant. A mixed model with genotype as random effect and experiment as fixed effects was used to estimate the variance explained by genotype (V_g). Residual variance was taken as an environmental variance (V_e). H^2 was then calculated as the ratio between V_g and the sum of V_g and V_e . The mixed model was constructed with the lmer function in the lme4 R package.

Genome-wide association mapping.

Genome-wide association mapping was done for each of the growth parameters, in both temperatures and also the temperature response for the growth rate. We used a mixed model with phenotype as dependent variable, genotype as fixed effect and genetic relatedness as random factor. Genotypes non-imputed SNPs obtained from the 1001 genomes consortium (19). This model was run in GEMMA, version 0.98.3 (52).

Testing for adaptive differentiation.

Adaptive differentiation was tested with the method described by Josephs et al. (30) and the accompanying R package quaint (<https://github.com/emjosephs/quaint>). Kinship matrix was calculated using the make_k function in the quaint package in R. Genetic principal components were then calculated from the eigen decomposition of the kinship matrix. Adaptive differentiation of each phenotype along the first 10 principal components was tested with the calcQpc function in the quaint R package. Principal components 11–248 were used to build the expected phenotypic differentiation under neutrality.

ACKNOWLEDGEMENTS

We want to thank past and current members of the Nordborg group for their help in setting up and harvesting these experiments. Thanks also to Daniele Filiault for her valuable comments on the manuscript.

Supplementary Information

Scripts can be found in https://github.com/picla/growth_16C_6C/. Scripts for seedsize analysis can be found in https://github.com/vevel/seed_size.

Supplemental dataset 1. List of all 249 accessions. Supplemental dataset 2. Cold acclimation genes and their expression cluster membership as shown in Figure 5. All RNA-sequecing were uploaded to SRA under <http://www.ncbi.nlm.nih.gov/bioproject/807069> All generated phenotyping data are filed under 10.5281/zenodo.6076948

Bibliography

- Margarita Takou, Benedict Wieters, Stanislav Kopriva, George Coupland, Anja Linstädter, and Juliette De Meaux. Linking genes with ecological strategies in arabidopsis thaliana. *J. Exp. Bot.*, 70(4):1141–1151, February 2019.
- Alejandra Martínez-Berdejá, Michelle C Stitzer, Mark A Taylor, Miki Okada, Exequiel Ezcurra, Daniel E Runcie, and Johanna Schmitt. Functional variants of DOG1 control seed chilling responses and variation in seasonal life-history strategies in arabidopsis thaliana. *Proc. Natl. Acad. Sci. U. S. A.*, 117(5):2526–2534, February 2020.
- Gordon G Simpson and Caroline Dean. Arabidopsis, the rosetta stone of flowering time? *Science*, 296(5566):285–289, April 2002.
- Jo Hepworth, Rea L Antoniou-Kourounioti, Rebecca H Bloomer, Catja Selga, Kristina Berggren, Deborah Cox, Barley R Collier Harris, Judith A Irwin, Svante Holm, Torbjörn Säll, Martin Howard, and Caroline Dean. Absence of warmth permits epigenetic memory of winter in arabidopsis. *Nat. Commun.*, 9(1):639, February 2018.
- Johanna A Bac-Molenaar, Dick Vreugdenhil, Christine Granier, and Joost J B Keurentjes. Genome-wide association mapping of growth dynamics detects time-specific and general quantitative trait loci. *J. Exp. Bot.*, 66(18):5567–5580, September 2015.
- Elodie Marchadier, Mathieu Hanemian, Sébastien Tisné, Lién Bach, Christos Bazakos, Elodie Gilbault, Parham Haddadi, Laetitia Virlovet, and Olivier Loudet. The complex genetic architecture of shoot growth natural variation in arabidopsis thaliana. *PLoS Genet.*, 15(4):e1007954, April 2019.
- Benedict Wieters, Kim A Steige, Fei He, Evan M Koch, Sebastián E Ramos-Onsins, Hongya Gu, Ya-Long Guo, Shamil Sunyaev, and Juliette de Meaux. Polygenic adaptation of rosette growth in arabidopsis thaliana. *PLoS Genet.*, 17(1):e1008748, January 2021.

- A Forsman. Rethinking phenotypic plasticity and its consequences for individuals, populations and species. *Heredity*, 115(4):276–284, October 2015.
- Mark Van Kleunen and Markus Fischer. Constraints on the evolution of adaptive phenotypic plasticity in plants, 2005.
- Hannes Claeys and Dirk Inzé. The agony of choice: how plants balance growth and survival under water-limiting conditions. *Plant Physiol.*, 162(4):1768–1779, August 2013.
- N N Artus, M Uemura, P L Steponkus, S J Gilmour, C Lin, and M F Thomashow. Constitutive expression of the cold-regulated arabidopsis thaliana COR15a gene affects both chloroplast and protoplast freezing tolerance. *Proc. Natl. Acad. Sci. U. S. A.*, 93(23):13404–13409, November 1996.
- P L Steponkus, M Uemura, R A Joseph, S J Gilmour, and M F Thomashow. Mode of action of the COR15a gene on the freezing tolerance of arabidopsis thaliana. *Proc. Natl. Acad. Sci. U. S. A.*, 95(24):14570–14575, November 1998.
- T Nanjo, M Kobayashi, Y Yoshida, Y Kakubari, K Yamaguchi-Shinozaki, and K Shinozaki. Antisense suppression of proline degradation improves tolerance to freezing and salinity in arabidopsis thaliana. *FEBS Lett.*, 461(3):205–210, November 1999.
- Sarah J Gilmour, Audrey M Sebolt, Maite P Salazar, John D Everard, and Michael F Thomashow. Overexpression of the arabidopsis CBF3 transcriptional activator mimics multiple biochemical changes associated with cold acclimation. *Plant Physiol.*, 124(4):1854–1865, December 2000.
- Teruaki Tajima, Chieko Ohsumi, Satoshi Iuchi, Motoaki Seki, Mie Kasuga, Masatomo Kobayashi, Kazuko Yamaguchi-Shinozaki, and Kazuo Shinozaki. Important roles of drought- and cold-inducible genes for galactinol synthase in stress tolerance in arabidopsis thaliana. *Plant J.*, 29(4):417–426, February 2002.
- C G Oakley, J Ågren, R A Atchison, and others. QTL mapping of freezing tolerance: links to fitness and adaptive trade-offs. *Molecular*, 2014.
- Sunchung Park, Sarah J Gilmour, Rebecca Grumet, and Michael F Thomashow. CBF-dependent and CBF-independent regulatory pathways contribute to the differences in freezing tolerance and cold-regulated gene expression of two arabidopsis ecotypes locally adapted to sites in sweden and Italy. *PLoS One*, 13(12):e0207723, December 2018.
- Carlos Alonso-Blanco, Concepción Gómez-Mena, Francisco Llorente, Maarten Koornneef, Julio Salinas, and José M Martínez-Zapater. Genetic and molecular analyses of natural variation indicate CBF2 as a candidate gene for underlying a freezing tolerance quantitative trait locus in arabidopsis. *Plant Physiol.*, 139(3):1304–1312, November 2005.
- 1001 Genomes Consortium. 1,135 genomes reveal the global pattern of polymorphism in arabidopsis thaliana. *Cell*, 166(2):481–491, June 2016.
- C E Timothy Paine, Toby R Mathews, Deborah R Vogt, Drew Purves, Mark Rees, Andy Hector, and Lindsay A Turnbull. How to fit nonlinear plant growth models and calculate growth rates: an update for ecologists. *Methods Ecol. Evol.*, 3(2):245–256, April 2012.
- Jakob Weiszmänn, Pieter Clauw, Joanna Jagoda, Ilka Reichardt-Gómez, Stefanie Koemeda, Jakub Jez, Magnus Nordborg, Dirk Walthers, Thomas Nägele, and Wolfram Weckwerth. Plasticity of the primary metabolome in 241 cold grown arabidopsis thaliana accessions and its relation to natural habitat temperature. preprint on BioRxiv, September 2020.
- Sunchung Park, Chin-Mei Lee, Colleen J Doherty, Sarah J Gilmour, Yongsig Kim, and Michael F Thomashow. Regulation of the arabidopsis CBF regulon by a complex low-temperature regulatory network, 2015.
- Jonathan T Vogel, Daniel G Zarka, Heather A Van Buskirk, Sarah G Fowler, and Michael F Thomashow. Roles of the CBF2 and ZAT12 transcription factors in configuring the low temperature transcriptome of arabidopsis. *Plant J.*, 41(2):195–211, January 2005.
- Girdhar K Pandey, Poonam Kanwar, Amarjeet Singh, Leonie Steinhilber, Amita Pandey, Akhilesh K Yadav, Indu Tokas, Sibaji K Sanyal, Beom-Gi Kim, Sung-Chul Lee, Yong-Hwa Cheong, Jörg Kudla, and Sheng Luan. Calcineurin B-Like Protein-Interacting protein kinase CIPK21 regulates osmotic and salt stress responses in arabidopsis. *Plant Physiol.*, 169(1):780–792, September 2015.
- Louisa M Liberman, Erin E Sparks, Miguel A Moreno-Risueno, Jalean J Petricka, and Philip N Benfey. MYB36 regulates the transition from proliferation to differentiation in the arabidopsis root. *Proc. Natl. Acad. Sci. U. S. A.*, 112(39):12099–12104, September 2015.
- Liesbeth Vercruyssen, Aurine Verkest, Nathalie Gonzalez, Ken S Heyndrickx, Dominique Eeckhout, Soon-Ki Han, Teddy Jégou, Rafal Archacki, Jelle Van Leene, Megan Andriankaja, Stefanie De Bodt, Thomas Abeel, Frederik Coppens, Stijn Dhondt, Liesbeth De Milde, Matthias Vermeersch, Katrien Maleux, Kris Gevaert, Andrzej Jerzmanowski, Moussa Benhamed, Doris Wagner, Klaas Vandepoel, Geert De Jaeger, and Dirk Inzé. ANGUSTIFOLIA3 binds to SWI/SNF chromatin remodeling complexes to regulate transcription during arabidopsis leaf development. *Plant Cell*, 26(1):210–229, January 2014.
- T Prout and J S Barker. F statistics in drosophila buzzatii: selection, population size and inbreeding. *Genetics*, 134(1):369–375, May 1993.
- Michael C Whitlock. Evolutionary inference from QST. *Mol. Ecol.*, 17(8):1885–1896, April 2008.
- K Spitz. Population structure in daphnia obtusa: quantitative genetic and allozymic variation. *Genetics*, 135(2):367–374, October 1993.
- Emily B Josephs, Jeremy J Berg, Jeffrey Ross-Ibarra, and Graham Coop. Detecting adaptive differentiation in structured populations with genomic data and common gardens. *Genetics*, 211(3):989–1004, March 2019.
- J P Grime. Primary strategies in plants. *Transactions of the Botanical Society of Edinburgh*, 43(2):151–160, November 1979.
- Mark Westoby. A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant Soil*, 199(2):213–227, February 1998.
- H K Hayes and O S Aamodt. Inheritance of winter hardiness and growth habit in crosses of marquis with minhardi and minturki wheats. *J. Agric. Res.*, 35(223):236, 1927.
- Karl S Quisenberry. *Inheritance of Winter Hardiness, Growth Habit, and Stem-rust Reaction in Crosses Between Minhardi Winter and H-44 Spring Wheats*. U.S. Department of Agriculture, 1931.
- Yves Castonguay, Serge Laberge, E Charles Brummer, and Jeffrey J Volenec. Alfalfa winter hardiness: A research retrospective and integrated perspective. In *Advances in Agronomy*, volume 90, pages 203–265. Academic Press, January 2006.
- Pauline Bristiel, Lauren Gillespie, Liv Østrem, Jennifer Balachowski, Cyrille Violle, and Flo-

rence Volaire. Experimental evaluation of the robustness of the growth–stress tolerance trade-off within the perennial grass *dactylis glomerata*. *Funct. Ecol.*, 32(8):1944–1958, August 2018.

37. Kari Koehler, Alyson Center, and Jeannine Cavender-Bares. Evidence for a freezing tolerance-growth rate trade-off in the live oaks (*quercus* series *virentes*) across the tropical-temperate divide. *New Phytol.*, 193(3):730–744, February 2012.

38. Craig Loehle. Height growth rate tradeoffs determine northern and southern range limits for trees. *J. Biogeogr.*, 25(4):735–742, July 1998.

39. Marco A Molina-Montenegro, Jorge Gallardo-Cerda, T S M Flores, and Cristian Atala. The trade-off between cold resistance and growth determines the *nothofagus pumilio* treeline. *Plant Ecol.*, 213(1):133–142, January 2012.

40. Jessica A Savage and Jeannine Cavender-Bares. Phenological cues drive an apparent trade-off between freezing tolerance and growth in the family *salicaceae*. *Ecology*, 94(8): 1708–1717, August 2013.

41. Ying Zhen, Preeti Dhakal, and Mark C Ungerer. Fitness benefits and costs of cold acclimation in *arabidopsis thaliana*. *Am. Nat.*, 178(1):44–52, July 2011.

42. Ellen Zuther, Elisa Schulz, Liam H Childs, and Dirk K Hincha. Clinal variation in the non-acclimated and cold-acclimated freezing tolerance of *arabidopsis thaliana* accessions. *Plant Cell Environ.*, 35(10):1860–1878, October 2012.

43. Matthew A Hannah, Dana Wiese, Susanne Freund, Oliver Fiehn, Arnd G Heyer, and Dirk K Hincha. Natural genetic variation of freezing tolerance in *arabidopsis*. *Plant Physiol.*, 142(1):98–112, September 2006.

44. Matthew W Horton, Glenda Willems, Eriko Sasaki, Maarten Koornneef, and Magnus Nordborg. The genetic architecture of freezing tolerance varies across the range of *arabidopsis thaliana*. *Plant Cell Environ.*, 39(11):2570–2579, 2016.

45. Malia A Gehan, Sunchung Park, Sarah J Gilmour, Chuanfu An, Chin-Mei Lee, and Michael F Thomashow. Natural variation in the c-repeat binding factor cold response pathway correlates with local adaptation of *arabidopsis* ecotypes. *Plant J.*, 84(4):682–693, November 2015.

46. Astrid Junker, Moses M Muraya, Kathleen Weigelt-Fischer, Fernando Arana-Ceballos, Christian Klukas, Albrecht E Melchinger, Rhonda C Meyer, David Riewe, and Thomas Altmann. Optimizing experimental procedures for quantitative evaluation of crop plant performance in high throughput phenotyping systems. *Front. Plant Sci.*, 5:770, 2014.

47. Stephen E Fick and Robert J Hijmans. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *Int. J. Climatol.*, 37(12):4302–4315, October 2017.

48. R Core Team. R: a language and environment for statistical computing, 2017.

49. Envel Kerdaffrec, Danièle L Filiault, Arthur Korte, Eriko Sasaki, Viktoria Nizhynska, Ümit Seren, and Magnus Nordborg. Multiple alleles at a single locus control seed dormancy in swedish *arabidopsis*. *Elife*, 5, December 2016.

50. Moises Exposito-Alonso, Claude Becker, Verena J Schuenemann, Ella Reiter, Claudia Setzer, Radka Slovak, Benjamin Brachi, Jörg Hagmann, Dominik G Grimm, Jiahui Chen, Wolfgang Busch, Joy Bergelson, Rob W Ness, Johannes Krause, Hernán A Burbano, and Detlef Weigel. The rate and potential relevance of new mutations in a colonizing plant lineage. *PLoS Genet.*, 14(2):e1007155, February 2018.

51. Rob Patro, Geet Duggal, Michael I Love, Rafael A Irizarry, and Carl Kingsford. Salmon provides fast and bias-aware quantification of transcript expression. *Nat. Methods*, 14(4): 417–419, April 2017.

52. Xiang Zhou and Matthew Stephens. Genome-wide efficient mixed-model analysis for association studies. *Nat. Genet.*, 44(7):821–824, June 2012.

652 Supplemental data

Fig. S1. Correlations between growth parameters and (bio)climate variables. (Bio)climate variables taken from the worldclim database (<https://www.worldclim.org>). Colors are correlation coefficients, correlations with FDR corrected p-values lower than 0.05 are indicated with a star. The order of the climate variables and phenotypes is based on hierarchical clustering.

Fig. S2. Population structure corrected correlations between growth parameters and (bio)climate variables. (Bio)climate variables taken from the worldclim database (<https://www.worldclim.org>). Colors are the coefficients for the climate variable in the mixed model with phenotype as dependent variable and population structure as random factor. Phenotypes and climate variables were standardized, making regression coefficients comparable to correlation coefficients. Correlations with FDR corrected p-values lower than 0.05 are indicated with a star. The order of the climate variables and phenotypes is based on hierarchical clustering.

Fig. S3. Correlation between initial size and seed size (A) and between seed size and winter temperature (B) for a subset of 123 Swedish accessions.

Fig. S4. Growth parameters and metabolic distance of rna-sequenced accessions in relation to local mean temperature of coldest quarter. Initial size (A), growth rate at 16°C (B) and 6°C (C), and metabolic distance (D), as a measure of temperature response over all 37 measured primary metabolites (21). Accessions selected for RNA-sequencing are depicted in black, remaining accessions are shown in gray.

Fig. S5. Cluster-specific expression in relation to winter temperature. Gene-wide standardized expression in 16°C (red) and 6°C (blue) values are plotted for each gene in clusters 1-7 (A-G), as defined in Figure 5. Expression values of each gene are connected with thin lines. Thick lines represent the correlation of the cluster's expression with the accession's winter temperature.

Fig. S6. Gene expression correlations with winter temperature. Correlation coefficients of each gene's correlations with winter temperature are grouped by the experimental temperature (16°C and 6°C) and by the expression direction upon cold exposure as measured by Park et al. and Vogel et al. (22) (23).

Fig. S7. GWAS results for the initial size, growth rate at 6°C and the temperature response of the growth rate. Left: Manhattan plots showing the significance of the association between the initial size, growth rate in 6°C and the growth rate's temperature response, and each of the tested SNPs. The Bonferroni-corrected threshold is visualized with the dashed red line. Right: QQ-plots showing the relation between observed and expected $-\log_{10}(\text{p-value})$ distributions for each of the respective GWAS. Red line shows the observed relationship. Gray line and band show the expected relationship under the null hypothesis of no differentiation between both distributions.