1 **Genome-wide analyses of histone modifications and chromatin accessibility**
2 **reveal the distinct genomic compartments in the Irish potato famine pathogen**
3 *Phytophthora infestans*

4

5

6

7

8

9 Han Chen^{1#}, Haidong Shu^{1#}, Yufeng Fang², Wenrui Song¹, Zhi Li¹, Yujie Fang¹,
10 Yuanchao Wang¹ and Suomeng Dong^{1*}

11

12

13

14 ¹Department of Plant Pathology and The Key Laboratory of Plant Immunity, Nanjing
15 Agricultural University, Nanjing, China.

16 ²GreenLight Biosciences Inc, Research Triangle Park, North Carolina, United States of
17 America

18

19

20

21

22

23

24

25 #These authors contributed equally to this work

26 *Corresponding author, E-mail: smdong@njau.edu.cn

Abstract

Phytophthora infestans, the causal agent of potato late blight, is a devastating plant disease that leads to Irish potato famine and threatens world-wide food security. Despite the genome of *P. infestans* has provided fundamental resource for studying the aggressiveness of this pandemic pathogen, the epigenomes remain poorly understood. Here, utilizing liquid chromatography-tandem mass spectrometry (LC-MS/MS), we demonstrate post-translational modifications (PTM) at *P. infestans* core histone H3. The PTMs not only include these prevalent modifications in eukaryotes, and also some novel marks, such as H3K53me2 and H3K122me3. We focused on the trimethylations of H3K4, H3K9 and H3K27 and H3K36, and profiled *P. infestans* epigenomes employing Native Chromatin Immunoprecipitation followed by sequencing (N-ChIP-seq). In parallel, we mapped *P. infestans* chromatin accessibility by Assay for Transposase-Accessible Chromatin with high-throughput sequencing (ATAC-seq). We found that adaptive genomic compartments display significantly higher levels of H3K9me3 and H3K27me3, and are generally in condense chromatin. Interestingly, we observed that genes encoding virulence factors, such as effectors, are enriched in open chromatin regions that barely have the four histone modifications. With a combination of genomic, epigenomic, transcriptomic strategies, our study illustrates the epigenetic states in *P. infestans*, which will help to study genomic functions and regulations in this pathogen.

48 Keywords: *Phytophthora infestans*, epigenetics, adaptive genome, histone modifications,

49 chromatin accessibility, gene expression

50

Author summary

Epigenetics play an important role in various biological processes of eukaryotes, including pathogenicity of plant pathogens. However, the epigenetic landscapes are marginally known in oomycetes that are fungal-like organisms and comprise lots of destructive plant pathogens. In this study, using the Irish potato famine pathogen *Phytophthora infestans* as a model, we conducted genome-wide studies of histone post modifications and chromatin accessibility, and demonstrate the relationship of gene expression and evolution with the epigenetic marks. We found that one of the most important classes of virulence proteins, effectors, are enriched in open chromatin regions that barely have eu- and hetero-chromatic marks. This study provides an overview of the oomycete epigenetic atlas, and advances our understanding of the regulation of virulence factors in plant pathogens.

Introduction

Late blight, a devastating disease is caused by *Phytophthora infestans* that strikes tomatoes and potatoes. The disease is notorious for triggering the 1840s Irish potato famine that resulted in the death of roughly one million people and the displacement of another million [1-3]. Today, *P. infestans* still remains a big threat to global plant health, causing ~US\$6.5 billion losses annually [4-6]. Physiologically and morphologically *P. infestans* resembles filamentous fungi; however, it belongs to oomycete in the Stramenopila kingdom [7, 8]. A large number of oomycetes are destructive plant pathogens whose virulence extensively rely on virulence factors including extracellular toxins, hydrolytic enzymes and inhibitors, and effectors that can enter the cytoplasm of plant cells [9]. In particular, effectors are one of the most important factors that alter host physiology via initiating and allowing an infection to develop [10].

P. infestans has a large and complex genomes (~240 Mb), approximately 74% of the genome is composed of repeats, such as transposable elements [11]. Due to fast bursts of the transposable element activities, repeat-rich genomic regions are highly dynamic and prone to evolutionary changes at accelerated rates compared with the rest of the genome, which shapes the *P. infestans* genome “two-speed” [9, 12, 13]. Interestingly, these repeat-rich regions tend to harbor genes that contribute to virulence such as effector genes, which is critical for host adaptation [9, 12, 13]. To date, studies

have shown functions and evolutionary trajectory of the effector genes (reviewed in [9, 12, 14]), but how these genes are regulated is marginally known.

Epigenetic regulation is crucial for gene expression in the eukaryotes, which is generally influenced by chromatin states, non-coding RNA (ncRNA) and associated machinery, and modifications of DNA or histones, etc. [15, 16]. Importantly, pathogenicity traits, such as virulence [17], sexual reproduction [18], growth [19, 20], and drug resistance [21] have been shown to correlate with epigenetic regulation in eukaryotic pathogens. For instance, in the fungal kingdom, histone modifications, ncRNA mediated silencing and chromosome alterations have been shown to regulate the expression of virulence factors, antifungal drug resistance, and interaction with hosts during infections [21-26]. Similarly, in parasites, histone methylations were reported to control growth and virulence [17, 27]. In oomycetes, epigenetic studies are limited, but studies indicated that epigenetic profiles of oomycetes are distinctive comparing to other organisms, and that ncRNA, histone modifications and chromosome states contributed to virulence. Strikingly, the prevalent 5-cytosine methylation (5mC) is missing in *Phytophthora* species, instead 6-adenine methylation (6mA) is widely distributed across the genomes [28]. This unusual DNA methylation profile makes *Phytophthora* a great model to study the evolution of DNA methylation. Recently, ChIP-seq based on the H3 variant CenH3 (CENP-A) uncovered the centromeres in *P. sojae*, which lack H3K4me2 but embed within the heterochromatin marks H3K9me3 and H3K27me3 [29]. *In silico*, several histone

modification enzymes were identified in *Phytophthora*, including histone acetyltransferases (HATs), deacetylases (HDACs) [30], some of which were involved in metabolic and biosynthetic process, sexual reproduction and virulence [31, 32]. In addition, gene silencing has been shown to be mediated by multiple different mechanisms, and be an effective way for oomycete pathogens to modulate virulence factors [33-35]. In *P. sojae*, transcription of the effector *Avr3a* was regulated by sRNA, which led to transgenerational silencing [33], while silencing of another effector gene *Avr1b* was correlated with H3K27me3 deposition [36]. In comparison, transgene-induced silencing of the elicitor gene *INF1* involves chromatin alteration in *P. infestans*, *INF1*-silenced strains harbor distinctive chromatin accessibility in the *INF1* loci [37, 38].

Despite evidence indicated that epigenetic processes affect growth, reproduction, and virulence, we lack an overview of the epigenetic states especially the histone modifications and chromosome states in oomycetes. In this study, we used *P. infestans* as a model to systemically investigate the modifications of histone H3 by LC-MS/MS. We focused on four important H3 modifications, H3K4me3, H3K9me3, H3K27me3 and H3K36me3 that are hallmarks of eu-and hetero- chromatins, and profiled their distributions across the genome. We also assessed the chromatin accessibility via ATAC-seq. We demonstrate that the histone H3 methylations and chromatin accessibility reflect *Phytophthora* genome structure, evolution and gene expression, and correlate closely with virulence factors. These findings provide a new insight of oomycete

123 epigenome structures, and advance our understanding of oomycete genome

124 architectures, which sheds the light on future studies aimed at the epigenetic regulation

125 of plant pathogens.

126

127

Results

Detection of histone H3 post-translational modifications (PTMs) in *P. infestans*

We are interested in dissecting how *P. infestans* histone H3 are decorated. A previous study has cloned *P. sojae* H3 (PsH3), and demonstrated its predominant nuclear localization in the *P. sojae* transformants by fusing to GFP [39]. To identify histone H3 in *P. infestans*, we performed blast searches against *P. infestans* T30-4 genome employing the PsH3 ortholog. We found five H3 homologs (PITG_03551, PITG_05675, PITG_20725, PITG_06953 and PITG_13828) present in *P. infestans*. Closer examination of each H3 homologs revealed that PITG_13828 is CENP-A (CenH3), because of its sequence and structure similarity to the *P. sojae* CENP-A [29] (data not shown). Intriguingly, PITG_03551 and PITG_05675 (PiH3-1) have an identical amino acid sequence, but different nucleotide sequences, so did for PITG_20725 and PITG_06953 (PiH3-2) (S1 Fig., S2 Fig.).

To detect histone H3 post-translational modifications (PTMs) in *P. infestans*, we conducted high-performance liquid chromatography-mass spectrometry (HPLC-MS) employing trypsin digested histones. In total, we detected 23 PTM forms in *P. infestans* (Fig. 1A). Most reported H3 PTMs can be found over the *P. infestans* H3 tails, such as the conventional mono-, di-, and tri- methylation of lysine [K], acetylation of lysine, tyrosine and serine [K/T/S], and recently discovered butyrylation [K], 2-hydroxyisobutyrylation [K], crotonylation [K], hydroxylation [Y], and malonylation [K]. Of

note, we did not detect the acetylation of K27 that was a common acetylation form in other organisms. On the other hand, we found some unique PMTs, such as di-methylation at K53, tri-methylation at K122 (S3 Fig., S4 Fig.).

Further examination revealed distinct distributions of PTMs over the two H3 orthologues. Five PTM variations occurred across 10 amino acid substitutions between two types of H3 (S3 Fig.). Acetylation and phosphorylation at T32 were detected in PiH3-1. Besides, acetylation at T31, di-methylation at K53 and phosphorylation at Y78 were detected in PiH3-2. It suggests different biological functions of H3 variants. Taken together, these findings indicate that despite most of the histone PTMs are conserved in *P. infestans* H3, several distinctive modification patterns are detected over several residues.

Profiling the genome-wide distribution of four histone H3 methylations and chromatin accessibility

To study the function of H3 PTMs, we focused on four histone methylations H3K4me3, H3K36me3, H3K9me3 and H3K27me3, which are the hallmarks of transcriptionally active euchromatin and transcriptionally silent heterochromatin. To validate the expression of the four histone H3 methylations in *P. infestans*, we carried out western blot using antibodies against these marks. As shown in S5 Fig., all the four histone methylations were detected in *P. infestans*, but not in the *Escherichia coli* expressed H3

protein, as it lacks post histone modifications.

To study genome-wide distribution of H3K4me3, H3K36me3, H3K9me3, and H3K27me3, we performed native chromatin immunoprecipitation (N-ChIP) employing the four antibodies that were tested in western blot, followed by high-throughput Illumina DNA sequencing. We generated high-quality map of each histone methylation with an average of 30 million reads that were uniquely mapped to the *P. infestans* reference genome (S6 Fig.). The four histone methylations displayed distinct distributions: For the euchromatic marks, the majority of H3K4me3 peaks (62.6%) were distributed in gene bodies, while 9.86% were in upstream of genes and 25.25% in intergenic regions (Fig. 2A). Moreover, we found that H3K4me3 was highly enriched in transcription start sites (TSS) (Fig. 2B). In comparison, H3K36me3 signals were scarce in upstream regions, but were highly enriched in gene bodies (41.46%) and intergenic regions (54.02%) (Fig. 2A and B). With respect to the heterochromatic marks H3K9me3 and H3K27me3, they generally exhibited a similar distribution pattern: 95.44% H3K9me3 peaks and 96.01% H3K27me3 peaks were localized in the intergenic regions (Fig. 2A, B and C).

One remarkable feature of heterochromatic marks is its association with repetitive sequences [40]. In *P. sojæ*, repetitive sequences, in particular, transposable elements (TEs) were highly enriched with H3K27me3 and H3K9me3 [41]. Similarly, we found that TEs in *P. infestans*, were highly enriched with H3K9me3 and H3K27me3, while lacked H3K4me3 and H3K36me3 (Fig. 2C and S7A Fig.). Further investigation of different TE

families revealed that 55% H3K9me3 and 66% H3K27me3 signals were associated with LTR retrotransposons that is the major TE family is the *P. infestans* genome (S7B Fig). Interestingly, the two heterochromatic marks exhibited quite different distribution over the DNA transposons, with 16% H3K9me3 and 5% H3K27me3 (S7B, Fig).

To further investigate the chromatin state over the *P. infestans* genome, we profiled the chromatin accessibility based on Assay for Transposase-Accessible Chromatin with high-throughput sequencing (ATAC-seq). A high-quality chromatin accessibility map was generated by aligning 100 million ATAC-seq reads uniquely to the reference genome (S6 Fig). We found that almost half of the ATAC-seq reads were mapped to the intergenic regions, while 29.05% were in gene bodies and 12.59% were in upstream of genes (Fig. 2A). To examine the relationship between chromatin accessibility and the four histone modifications, we implemented correlation analysis by PCA and heatmap. We found that heterochromatin was anticorrelated with chromatin accessibility (Pearson co-efficiency ranging from -0.15 to -0.23), while half euchromatin were somewhat correlated with chromatin accessibility (Pearson co-efficiency ranging from 0.12 to 0.20) (S8 Fig, S9 Fig). Comparing to H3K4me3, we found that the major ATAC-seq signals approximal gene bodies were around 100 bp upstream of TSS, indicating that the open chromatin state of promoter regions (Fig. 2B). It suggests the unique role of chromatin accessibility in chromatin state defining compared with those four PTMs.

Chromatin states strongly correlate with gene expression

We next sought to examine the correlation of H3 modifications and chromatin accessibility with gene expression. To determine expression levels of genes from the same growth stage that was used for profiling chromatin states, we performed RNA-seq from the *P. infestans* mycelial stage. We classified all *P. infestans* genes into six categories from “Silent” to “Top” expression, based on their expression levels, (Fig. 2D). We found that genes of higher expression were generally located in chromatin regions that are more accessible, and that H3K4me3 and H3K36me3 were positively correlated with gene expression (Fig. 2D) suggesting open chromatin and these euchromatic histone marks are associated with active genes. In contrast, H3K9 and H3K27 trimethylations exhibited an anticorrelation with expression level, indicating these two histone marks are associated with silencing (Fig. 2D). Genome browser views of individual genomic locations also suggest a strong association of transcription activity with chromatin accessibility and the four histone marks (Fig. 2C).

Collectively, these results illustrate that methylation of H3K4 and H3K36 is associated with expressed genes, which are correlated with accessible chromatin; while the methylation of H3K9 and H3K27 is enriched in areas of the genome that are transcriptionally silent and are generally consisted of TEs, which tend to be closed chromatin conformation.

Chromatin states reflect the bipartite genome structure

The genomes of *Phytophthora* species show a bipartite “two-speed” architecture that is consisted of gene-dense regions (GDRs), and gene-spare regions (GSRs) [11, 13, 42]. The GSR compartments are associated with accelerated gene evolution, serving as a cradle for adaptive evolution [11, 13, 42]. To investigate the relationship between chromatin state and the genome architecture, we measured the average level of each epigenetic state for genes located in GDRs and GSRs. We discovered that GSRs were enriched with the heterochromatic marks H3K9me3 and H3K27me3, and had condensed chromatin, but lacked the euchromatic marks H3K4me3 and H3K36me3 (Fig. 3). In contrast, GDRs were enriched with the euchromatic marks H3K4me3 and H3K36me3, and were associated with more accessible chromatin (Fig. 3). Thus, the epigenetic states display close correlation with the “two-speed” genome architecture.

Chromatin states are associated with the conservation of protein-coding genes

To address the connection between epigenetic state and protein evolution, we first sought to identify genes that were evolutionarily conserved in *P. infestans*. We selected 17 eukaryotic species including oomycetes, fungi, plants and animals, and measured the ratios of non-synonymous/synonymous codon substitution (dN/dS) for protein-coding genes. We found that the most conserved genes across eukaryotes had lowest dN/dS ratios, while the genes specific to *P. infestans* had highest dN/dS ratios, suggesting

species-specific genes are evolved later (S10A, B Fig.). In agreement with this observation, high dN/dS density was also found in GSRs, further implying that the genes in these regions were under higher evolutionary pressure (S10C Fig.).

Based on the gene evolution analysis, we examined the epigenetic states over each protein-coding gene in *P. infestans*. We found that *P. infestans* specific genes had higher level of H3K9me3 and H3K27me3, and were generally associated with condensed chromatin. In contrast, conserved genes had higher level of H3K4me3 and H3K36me3, and were preferably associated with more open chromatin regions (Fig. 4A). We found that genes that were under positive selection accumulated more H3K9me3 and H3K27me3 signals, whereas the genes under purifying selection accumulated more H3K4me3 and H3K36me3 (Fig. 4B). These findings are consistent with the chromatin accessibility and methylation pattern of the “two speed” genome architecture that we found above, namely the conserved genes are usually distributed in accessible GDR, and thus are marked by H3K4me3 and H3K36me, whereas the rapidly evolved TEs are enriched in the condensed GSR and thus associated with H3K9me3 and H3K27me3.

Effector genes are enriched in highly accessible chromatin region with less histone methylation level

To characterize the functions of genes that were undergone histone methylation, we performed gene ontology (GO) analysis. We found that histone methylations widely

contributed to biological processes (S11 Fig.). Interestingly, while the processes regulated by the euchromatic marks H3K4me3 and H3K36me3 were mostly different, these regulated by H3K9me3 and H3K27me3 were largely coincident (S11 Fig.), further indicating cross-talks may occur between the two heterochromatic marks for regulating gene expression. Notably, GO analysis indicated that histone modifications broadly contribute to pathogenesis related functions, such as pectate lyase activity and extracellular region, indicating the epigenetic regulation are critical for virulence. This prompted us to investigate the relationship between histone methylations and secreted proteins (secretome), in particular RxLR effectors [43, 44]. We found that comparing to other protein-coding genes, genes encoding secretome and RxLR effectors had overall higher H3K9me3 and H3K27me3 and lower H3K4me3 and H3K36me3 densities (S12A, and S12B Fig.). It is possibly because the majority genes encoding RxLR effectors were repressed in mycelia stage.

To further investigate the role of chromatin state in regulating virulence, we generated a multivariate Hidden Markov Model based on the distribution of the four aforementioned histone marks and chromatin accessibility by chromHMM. *P. infestans* genome was divided into five distinct states including open chromatin (OC) that harbored ATAC-seq signals but lacked the four histone marks; strong transcription (ST) region that had abundant H3K4me3, H3K36me3, and ATAC-seq signals; H3K9me3 dominant repression region (H3K9DR); H3K27me3 dominant repression region (H3K27DR); and

quiescent (Quies) state that were absent of all the tested marks (Fig 5A). We found that 53.7% (868/1616) genes encoding secreted virulence factors, such as carbohydrate-active enzymes (CAZyme) and various effector families such as Crinkler effectors (CRN), necrosis- and ethylene-inducing-like proteins (NLP), small cysteine-rich effector proteins (SCR) and RxLR, were associated with the OC and ST states (Fig. 5D, E). Interestingly, despite OC state accounts for the smallest percentage (5.29% of the genome), 44% RxLR effector-coding genes were associated with this state (Fig. 5E). A further analysis showed CAZyme, NLP and RxLR gene families significantly enriched in OC state (Fig. 5F). GO analysis of genes from different chromatin state revealed, the GO items like extracellular region, cell wall organization and pectin catabolic process were the most enriched ones in OC state associated genes (S13 Fig.), suggesting the role of OC state in *P. infestans* pathogenicity gene regulation.

To address how the expression profiles of the virulence genes correlated with the OC state. We examined the transcriptomes of the *P. infestans* T30-4 strain at different life/infection stages, such as mycelium (MY), 2 dpi (days post incubation) and 3 dpi. We found that about half of the OC state-associated secreted virulence genes (CAZyme, CRN, NLP, SCR and RxLR) displayed infection-induced pattern, for instance secreted pectin monooxygenase encoding gene *PiAA17C* [45], and RxLR effector genes *Avrvnt1*, *AvrSmira1* and *Pi02860* [46-48] which are key virulence genes. In contrast, most ST state-associated secreted virulence genes were relative highly expressed, although

308 some of them were infection-induced like *Avr3a* (Fig. 5G). To further explain the
 309 expression difference, we summarized the upregulation fold change of OC state and ST
 310 state-associated secreted virulence genes. Among 251 secreted virulence genes in OC
 311 state, 41.03% genes were 5-fold upregulated, and 35.45% genes were 10-fold
 312 unregulated at least in one infection stage. Otherwise, the ratio is lower in ST state (Fig.
 313 5H). It suggests that OC state-associated virulence genes could be highly induced in
 314 infection stages. Altogether, our analyses implied the importance of OC state in virulence
 315 gene regulation.

Discussion

In this study, we identified the histone H3 PTMs in *P. infestans* by HPLC/MS, and generated an epigenetic atlas on basis of histone H3 methylations and chromatin accessibility. We found that most H3 PTMs were conserved in *P. infestans*. By examining the tri-methylation profiles of H3K4, H3K36, H3K9 and H3K27 and chromatin accessibility, we concluded that these chromatin states were closely associated with gene expression, genome structure, protein evolution, and virulence factors regulation. Our results provide evidence that epigenetic scenario is associated with pathogen genome in oomycete, and highlight chromatin state of virulence factors.

We found *Phytophthora* H3 proteins and H3 PTMs are slightly different from these reported in animal, plants and Protista, indicating dynamics of these highly conserved protein. With blast search, we found two H3 orthologues in *P. infestans*; however, with sequence composition and phylogenetic analyses, we cannot distinguish the canonical orthologue H3.1 and the variant H3.3 in *Phytophthora* (S2 Fig.). On the other hand, HPLC/MS revealed differences of PTM between the two *P. infestans* H3 orthologues (S3 Fig). This suggests that oomycete histone H3 may have a distinct evolutionary trajectory, additional experiment such as gene expression pattern examination in and outside of the S phase, and nucleosome assembly will help to clarify the two H3 variants in *Phytophthora*.

We found some unique modifications present in *P. infestans*, such as methylations at residues H3K53 and H3K122. H3K53me maybe species-specific, as in several organisms the residue 53 in H3 not lysine but arginine (S2B Fig.). In *Arabidopsis*, the H3K53 site was also reported to possess a modification; however, instead of methylation, it was 2-hydroxyisobutyrylation (Khib). Intriguingly, this unusual Khib modification contributed to the plant adaption to stress [49], suggesting H3K53me2 may also participate in some biological activities in *P. infestans*. Despite studies have shown that H3K122ac is a critical transcriptional regulator that defines genome-wide genetic elements and chromatin features associated with active transcription in mammalian cells [50], there is no report of H3K122 methylation, thus it will be of interest to examine its function in oomycetes. While most histone H3 PTMs reported to date can be found in *P. infestans*, we found acetylation of H3K27 and H3K36 are missing in the pathogen. Interestingly, histone acetyltransferase families that represent for H3K27ac and H3K36ac were identified [30], and a recent paper found both H3K27ac and H3K36ac in *P. infestans* [51]. It is possible that this modification level was too low to be detected in our mass spectrometry. Overall, we confirmed that large number of PTMs were presented in *P. infestans*, meanwhile, some unique PTMs were detected.

In *P. sojae*, it has been shown that H3K9me3 and H3K27me3 are generally coincident over the genome [29]. We generally observe a similar distribution pattern in *P. infestans*. Surprisingly, closer examination of TE demonstrated that the two

heterochromatic marks were separated over the DNA transposons (S7 Fig.), implying a different mode may be used for silencing the types of TE. Another interesting features of epigenetic marks we found in *P. infestans* is that H3K36me3 distributed over the gene bodies of active genes (Fig. 2B), which is distinctive from strong methylation peak at the 3' end of the gene body in human and mouse, and strong peak at the 5' end of the gene body *Arabidopsis* and rice, but is similar to the pattern in the brown alga *Ectocarpus siliculosus*, yeast, and *C. elegans* [52, 53], suggesting divergent regulators and mechanisms for establishing H3K36 methylation among species.

Genome-wide profiling of chromatin states provides insights into the unique genomic compartments of plant pathogens. Principally, here we directly connected the chromatin accessibility, histone methylation status and two-speed genome feature, and greatly enriched the dimension of two-speed genome (Fig. 3). Similar to our results, high enrichment of the heterochromatic marks H3K9me3 and H3K27me3, rather than the euchromatic mark H3K4me2 was found in the rapidly evolved accessory chromosomes in the wheat fungal pathogen *Zymoseptoria tritici* [54]. Recent research in the verticillium wilt fungus *Verticillium dahliae* revealed adaptive lineage-specific genomic regions contain many heterochromatin features, but more accessible than true heterochromatin [55]. The heterochromatin features of adaptive genomic compartments in plant pathogens remains a question that the formation and maintenance of these regions. Mechanically, experimental evolution evidence proved that loss of H3K27me3 by

knockout methyltransferases Kmt6 drastically reduces the loss of accessory chromosomes [56]. It explained why these regions are more susceptible to genetic diversity. Further experimental research in *P. infestans* is needed to understand the unique genomic compartments regulation. Besides, open chromatin sites reflect the potential of recombination [57, 58], and *A. thaliana* showed that mutational rates are significantly predicted by some epigenetic modifications [59]. It suggests that epigenetic status is associated with evolutionary hotspots. So, it's reasonable to infer that chromatin state details directly reflect sequence mutation and epimutation ratio in specific strains. Higher resolution of two-speed genome and the molecular basis of genome architectural reshaping are fascinating points to us.

Dissecting the chromatin state of genes encoding secreted virulence protein is one of the key steps to illuminate *P. infestans* pathogenicity regulation. Interestingly, the GO analysis indicated that gene related to extracellular region and pectin catabolic process are enriched in the OC state (Fig. 5, S13 Fig.), meanwhile, those items was also enriched in H3K9me3 and H3K27me3 methylated genes (S11 Fig.). We inferred that those virulence genes involved in these GO items were not overlap and in different regulation model. Firstly, most virulence genes in OC state had the potential to be induced in infection stages, this kind of genes performed strong induction pattern due to transcriptional releasing in infection stages (Fig. 5G, 5H). Secondly, the other virulence genes were strongly repressed by methylation of histone lysine residues. Those kinds of

genes might be epigenetically regulated in *Phytophthora* [36], which was also observed in other filamentous pathogens [60, 61]. Epigenetic silencing of TE by heterochromatin marks to prevent TE proliferation in the TE-rich region is quite common in filamentous pathogen genomes [62], and we indeed found adjacent TEs around *RxLR* gene cluster (S12 Fig.). Moreover, it has been accepted that epigenetic marks are tissue- and stage-specific [63-65], revealing epigenomes of different stages and strains can dissect holistic virulence gene regulation model. Here, we propose that functional regulatory elements that contributed to establish and maintain normal chromatin state could be drug targets to defeat diseases. In conclusion, our findings reported herein provide clues for virulence gene regulation mechanism investigation and pest management.

Materials and methods

Oomycete and fungal cultures

P. infestans strain T30-4, and *P. sojae* strain P6497 were used in the study. *P. infestans* was routinely maintained on RSA/V8 medium at 18°C in dark, and *P. sojae* on 10% V8 medium at 25°C in dark [28]. *M. oryzae* strain Guy11 was regularly cultured in CM medium at 28°C [66].

LC-MS/MS analysis

Core histones were extracted from *P. infestans* mycelia employing EpiQuik Total Histone Extraction Kit (EpiGentek, OP-0006-100), according to the manufacture instruction. After being separated by 12% SDS-PAGE, proteins with ~ 17 kDa size were extracted and analyzed by mass spectrum at Thermo Fisher. Given that H3 proteins have multiple trypsin digestion site (lysine and arginine), a portion of sample was treated with propionic anhydride (PA) to block trypsin digestion at the lysine sites to get maximal sequencing read coverage. The trypsin digested peptides (derivatized and non-derivatized) were injected to HLC/MS/MS (C18 HPLC column and Q Exactive HF-X Mass Spectrometer), and the acquired MS/MS data was analyzed by the pFind3.0 [67] with default parameters.

Protein extraction and western blot

To extract oomycete and fungal proteins, mycelia collected from liquid cultures of six days old *P. infestans*, three days old *P. sojae*, and three days old *M. oryzae* were dried by filter paper, and ground with mortars and pestles in liquid nitrogen. 800 µl lysis buffer (1% SDS in TE buffer) was added into every 100 mg pulverized oomycete and fungal tissue. Lysates were mixed by vortex for 30 min at 4°C. 200 µl supernatants were collected after spun prepared for western blot. Protein samples were mixed with protein loading buffer (Beyotime, P0015) and denatured at 95°C for 10 min. To express PsH3 in *E. coli*, the target gene (Ps_322070) was PCR amplified from *P. sojae*, and was cloned into the plasmid pET32a using Vazyme ClonExpress II One Step Cloning Kit C112. The resulting plasmid pET32a-PsH3 was introduced in *E. coli* BL21. The protein was expressed at 30°C for 8 hours in LB medium with 0.5 mM isopropyl β-D-Thiogalactoside (IPTG) according to the manufacture instruction [68]. After sonication and centrifugation, about 100 µl supernatant containing *E. coli* crude protein extract was used for western blot. Antibodies H3K36me3 (abcam, ab9050), H3K27me3 (Millipore, 07-449), H3 (abcam, ab1791), H3K4me3 (abcam, ab8580) and H3K9me3 (abcam, ab8898) were used as primary antibodies against the relevant histone H3 modifications, and goat-anti-rabbit IRDye 800CW antibody (Odyssey, no. 926-32211, Li-Cor) was as a secondary antibody. The signals were exposure by laser imaging system (Odyssey, LI-COR company).

Phylogenetic analysis

H3 protein sequences from human (HsH3.1, accession number: NP_003520.1, HsH3.2, NP_066403.2, HsH3.3/NP_002098.1, HsCenP-A/NP_001035891.1), *Arabidopsis thaliana* (AtH3.1/NP_563838.1, AtH3.3/ NP_001329167.1, AtCenH3/NP_009564.1), *Neurospora crassa* (NcH3/CAA25761.1) and *Saccharomyces cerevisiae* (ScH3/NP_009564.1) were downloaded from NCBI. *Phytophthora* H2A sequence were from published paper [69]. Homologous gene blast was obtained by Seqhunter2 with an E value of 10^{-5} . Sequence alignment was performed by Bioedit software [70], and phylogenetic analysis was conducted by MEGA5 using neighbor-joining model and 10000 bootstrap replicates [71].

Native ChIP-seq, ATAC-seq and RNA-seq

Native ChIP experiments were performed as previously described [29, 36, 72]. Input and immunoprecipitated DNA samples were sequenced by BGI company as 50SE. To prepare ATAC-seq, *P. infestans* protoplasts were isolated according to refrenece [73]. Protoplast was stored at -80°C using Nalgene 5100-0001C, and then sent to BGI for treatment [74]. In brief, lysing the cells and keep the cells on the ice all the time. Add Tn5 transposase to the cell suspension after cell lysis and then purify it. The DNA fragments were sequenced in BGI company as PE150. Infection samples (2 dpi and 3 dpi) were prepared as described [75]. RNA were extracted using Omega Total RNA Kit I according

to the manufacturer's manual, RNA-seq libraries were prepared by BGI company and sequenced by BGISEQ-500.

High-throughput sequencing data analysis

Both ChIP-seq and ATAC-seq reads were polished by BGI prior to be released, and thus were mapped to the *P. infestans* reference genome directly using Bowtie2 (v2.3.5.1) [76] (see S6 Fig. for mappability). The aligned bam files were sorted and indexed by samtools (version: 1.7) [77]. The ChIP-ed and input samples were analyzed with DeepTools (v3.4.3) [78] “bamCompare” to calculate normalized ChIP signals ($\log_2[\text{ChIP}_{\text{RPKM}}/\text{Input}_{\text{RPKM}}]$),. ChIP-seq data was visualized using TBtools [79] and Integrative Genome Viewer (IGV, v2.8.0) (<https://software.broadinstitute.org/software/igv/>) [80]. ChIP-seq peaks were defined by MACS2 employing a default model for H3K4me3 , and a “--broad” model for H3K9me3, H3K27me3, and H3K36me3 [81, 82]. Peak overlaps were conducted by “intersectBed” in bedtools [83]. To visualize the ATAC-seq reads, BAM files were converted to bigwig format using “bamCompare” in DeepTools with RPKM normalization. ATAC-seq data was visualized using TBtools [79] and IGV v2.8.0 [80]. The ATAC-seq reads were shifted +4 on the positive strands and -5 on the negative strands using deepTools software, and then ATAC-seq peaks were defined by MACS2 in default model. Peak overlap was conducted by “intersectBed” in bedtools. RNA-seq reads were polished by BGI prior to be released. To generate mRNA profiles,

the RNA-seq reads were aligned to the genomes using HISAT2 (version 2.1.0) [84] , and the resulting files (.bam) were sorted and indexed by samtools (version 1.9) [77]. The bam file was converted to .tdf for visualization using IGV. Gene expression data were calculated by StringTie v2.1.2 [85] and was presented as FPKM values.

Other analysis

PCA plot and heatmap clustering of ChIP-seq and ATAC-seq was using “plotPCA” and “plotCorrelation” in deepTools. Overlap of ATAC peak and ChIP-seq were conducted by “intersectBed” in bedtools. The Pearson correlation computation was used in heatmap clustering. The chromatin state was defined by chromHMM [86] in “BinarizeBed” model using MACS2 peak files. GO enrichment was analyzed by R package clusterprofile v3.14.3 by universal enrichment analyzer “enricher” and “compareCluster” [87]. Gene conservation analysis among species was performed by OrthoMCL v2.0.9 by default parameters [88]. Protein conservation was based on 17 eukaryotic species including oomycetes, fungi, plants and animals, and divided the *P. infestans* coding-genes into four groups (Eukaryote, Oomycete, *Phytophthora* and *P. infestans*), each of which has 1157, 3064, 5680 and 2483 genes. Eukaryote group contains protein-coding genes that conserved among all 17 species. Oomycete group contains protein-coding genes that conserved among 10 oomycetes besides genes in eukaryote group.

504 *Phytophthora* group and *P. infestans* specific groups can be inferred like this. The
 505 Sankey plot was performed by R package networkD3 [89].

506

507 **Funding**

508 This work was supported by the National Natural Science Foundation of China
 509 (NSFC) 31721004 and 32100158, China Postdoctoral Science Foundation 2019TQ0156
 510 and Natural Science Foundation of Jiangsu Province BK20200538.

511

512 **Competing interests**

513 The authors have declared that no competing interests exist.

514

515 **Acknowledgments**

516 We are grateful to the Bioinformatics Center at Nanjing Agricultural University for
 517 support of the bioinformatics analysis. We thank the NJAU oomycete research group for
 518 valuable discussion.

519

520

References

1. Kinealy C. This great calamity : the Irish famine 1845-52. 2006.
2. Turner RS. After the famine: Plant pathology, *Phytophthora infestans*, and the late blight of potatoes, 1845—1960. *Historical Studies in the Physical and Biological Sciences*. 2005;35(2):341-70. doi: 10.1525/hsps.2005.35.2.341.
3. McDonough T. Mapping the Great Irish Famine. JSTOR; 2001.
4. Fones HN, Bebber DP, Chaloner TM, Kay WT, Steinberg G, Gurr SJ. Threats to global food security from emerging fungal and oomycete crop pathogens. *Nature Food*. 2020;1(6):332-42. doi: 10.1038/s43016-020-0075-0.
5. Savary S, Willocquet L, Pethybridge SJ, Esker P, McRoberts N, Nelson A. The global burden of pathogens and pests on major food crops. *Nat Ecol Evol*. 2019;3(3):430-9. Epub 2019/02/06. doi: 10.1038/s41559-018-0793-y. PubMed PMID: 30718852.
6. Campos H, Ortiz O. The potato crop: its agricultural, nutritional and social contribution to humankind: Springer Nature; 2020.
7. Burki F, Roger AJ, Brown MW, Simpson AGB. The New Tree of Eukaryotes. *Trends in Ecology & Evolution*. 2020;35(1):43-55. doi: 10.1016/j.tree.2019.08.008.
8. Thines M, Kamoun S. Oomycete-plant coevolution: recent advances and future prospects. *Current opinion in plant biology*. 2010;13(4):427-33. doi: 10.1016/j.pbi.2010.04.001. PubMed PMID: 20447858.
9. Jiang RH, Tyler BM. Mechanisms and evolution of virulence in oomycetes. *Annual review of*

541 phytopathology. 2012;50:295-318. doi: 10.1146/annurev-phyto-081211-172912. PubMed PMID:
542 22920560.

543 10. Wawra S, Belmonte R, Lobach L, Saraiva M, Willems A, van West P. Secretion, delivery and
544 function of oomycete effector proteins. Current opinion in microbiology. 2012;15(6):685-91. Epub
545 2012/11/28. doi: 10.1016/j.mib.2012.10.008. PubMed PMID: 23177095.

546 11. Haas BJ, Kamoun S, Zody MC, Jiang RH, Handsaker RE, Cano LM, et al. Genome
547 sequence and analysis of the Irish potato famine pathogen *Phytophthora infestans*. Nature.
548 2009;461(7262):393-8. Epub 2009/09/11. doi: 10.1038/nature08358. PubMed PMID: 19741609.

549 12. Raffaele S, Kamoun S. Genome evolution in filamentous plant pathogens: why bigger can be
550 better. Nature reviews Microbiology. 2012;10(6):417-30. doi: 10.1038/nrmicro2790. PubMed
551 PMID: 22565130.

552 13. Dong S, Raffaele S, Kamoun S. The two-speed genomes of filamentous pathogens: waltz
553 with plants. Current opinion in genetics & development. 2015;35:57-65. doi:
554 10.1016/j.gde.2015.09.001. PubMed PMID: 26451981.

555 14. Birch PR, Boevink PC, Gilroy EM, Hein I, Pritchard L, Whisson SC. Oomycete RXLR
556 effectors: delivery, functional redundancy and durable disease resistance. Current opinion in plant
557 biology. 2008;11(4):373-9.

558 15. Goldberg AD, Allis CD, Bernstein E. Epigenetics: a landscape takes shape. Cell.
559 2007;128(4):635-8. Epub 2007/02/27. doi: 10.1016/j.cell.2007.02.006. PubMed PMID: 17320500.

560 16. Allis CD, Jenuwein T. The molecular hallmarks of epigenetic control. Nature reviews

561 Genetics. 2016;17(8):487-500. Epub 2016/06/28. doi: 10.1038/nrg.2016.59. PubMed PMID:
562 27346641.

563 17. Jiang L, Mu J, Zhang Q, Ni T, Srinivasan P, Rayavara K, et al. PfSETvs methylation of
564 histone H3K36 represses virulence genes in Plasmodium falciparum. Nature.
565 2013;499(7457):223-7. Epub 2013/07/05. doi: 10.1038/nature12361. PubMed PMID: 23823717;
566 PubMed Central PMCID: PMC3770130.

567 18. Scaduto CM, Kabrawala S, Thomson GJ, Scheving W, Ly A, Anderson MZ, et al. Epigenetic
568 control of pheromone MAPK signaling determines sexual fecundity in Candida albicans.
569 Proceedings of the National Academy of Sciences of the United States of America.
570 2017;114(52):13780-5. doi: 10.1073/PNAS.1711141115.

571 19. Mittelsten Scheid O, Zhang W, Huang J, Cook DE. Histone modification dynamics at H3K27
572 are associated with altered transcription of in planta induced genes in Magnaporthe oryzae. PLoS
573 genetics. 2021;17(2):e1009376. doi: 10.1371/journal.pgen.1009376.

574 20. Rai LS, Singha R, Brahma P, Sanyal K. Epigenetic determinants of phenotypic plasticity in
575 Candida albicans. Fungal Biology Reviews. 2018;32(1):10-9. doi: 10.1016/j.fbr.2017.07.002.

576 21. Calo S, Shertz-Wall C, Lee SC, Bastidas RJ, Nicolas FE, Granek JA, et al. Antifungal drug
577 resistance evoked via RNAi-dependent epimutations. Nature. 2014;513(7519):555-8. doi:
578 10.1038/nature13575. PubMed PMID: 25079329; PubMed Central PMCID: PMC4177005.

579 22. Connolly LR, Smith KM, Freitag M. The Fusarium graminearum histone H3 K27
580 methyltransferase KMT6 regulates development and expression of secondary metabolite gene

581 clusters. PLoS genetics. 2013;9(10):e1003916. Epub 2013/11/10. doi:
582 10.1371/journal.pgen.1003916. PubMed PMID: 24204317; PubMed Central PMCID:
583 PMC3814326.

584 23. Galazka JM, Freitag M. Variability of chromosome structure in pathogenic fungi--of 'ends and
585 odds'. Current opinion in microbiology. 2014;20:19-26. doi: 10.1016/j.mib.2014.04.002. PubMed
586 PMID: 24835423; PubMed Central PMCID: PMC4133287.

587 24. Liu Y, Liu N, Yin Y, Chen Y, Jiang J, Ma Z. Histone H3K4 methylation regulates hyphal
588 growth, secondary metabolism and multiple stress responses in *Fusarium graminearum*.
589 Environmental microbiology. 2015;17(11):4615-30. doi: 10.1111/1462-2920.12993. PubMed
590 PMID: 26234386.

591 25. Collemare J, Seidl MF. Chromatin-dependent regulation of secondary metabolite
592 biosynthesis in fungi : is the picture complete? *fems microbiology reviews*. 2019;43(6):591-607.
593 doi: 10.1093/FEMSRE/FUZ018.

594 26. Torres-Garcia S, Yaseen I, Shukla M, Audergon PNCB, White SA, Pidoux AL, et al.
595 Epigenetic gene silencing by heterochromatin primes fungal resistance. *Nature*. 2020. doi:
596 10.1038/s41586-020-2706-x.

597 27. Scherf A, Lopez-Rubio JJ, Riviere L. Antigenic Variation in *Plasmodium falciparum*. Annual
598 review of microbiology. 2008;62(1):445-70. doi: 10.1146/annurev.micro.61.080706.093134.
599 PubMed PMID: WOS:000259968000024.

600 28. Chen H, Shu H, Wang L, Zhang F, Li X, Ochola SO, et al. *Phytophthora methylomes* are

601 modulated by 6mA methyltransferases and associated with adaptive genome regions. *Genome*
602 *Biol.* 2018;19(1):181. doi: 10.1186/s13059-018-1564-4. PubMed PMID: 30382931; PubMed
603 Central PMCID: PMC6211444.

604 29. Fang Y, Coelho MA, Shu H, Schotanus K, Thimmappa BC, Yadav V, et al. Long transposon-
605 rich centromeres in an oomycete reveal divergence of centromere features in Stramenopila-
606 Alveolata-Rhizaria lineages. *PLoS genetics.* 2020;16(3):e1008646. doi:
607 10.1371/journal.pgen.1008646. PubMed PMID: 32150559.

608 30. Wang X, Guo L, Han M, Shan K. Diversity, evolution and expression profiles of histone
609 acetyltransferases and deacetylases in oomycetes. *BMC genomics.* 2016;17(1):927.

610 31. Wang X-W, Lv J-L, Shi Y-R, Guo L-Y. Comparative Transcriptome Analysis Revealed Genes
611 Regulated by Histone Acetylation and Genes Related to Sex Hormone Biosynthesis in
612 *Phytophthora infestans*. *Frontiers in genetics.* 2020;11. doi: 10.3389/fgene.2020.00508.

613 32. Tzelepis G, Hoden KP, Fogelqvist J, Asman AKM, Vetukuri RR, Dixelius C. Dominance of
614 Mating Type A1 and Indication of Epigenetic Effects During Early Stages of Mating in
615 *Phytophthora infestans*. *Front Microbiol.* 2020;11:252. Epub 2020/03/11. doi:
616 10.3389/fmicb.2020.00252. PubMed PMID: 32153537; PubMed Central PMCID:
617 PMCPMC7046690.

618 33. Qutob D, Chapman BP, Gijzen M. Transgenerational gene silencing causes gain of virulence
619 in a plant pathogen. *Nature communications.* 2013;4:1349. Epub 2013/01/17. doi:
620 10.1038/ncomms2354. PubMed PMID: 23322037; PubMed Central PMCID: PMC3562452.

621 34. Kasuga T, Gijzen M. Epigenetics and the evolution of virulence. Trends in microbiology.
622 2013;21(11):575-82. Epub 2013/10/08. doi: 10.1016/j.tim.2013.09.003. PubMed PMID: 24095304.

623 35. Shrestha SD, Chapman P, Zhang Y, Gijzen M. Strain Specific Factors Control Effector Gene
624 Silencing in *Phytophthora sojae*. PloS one. 2016;11(3):e0150530. doi:
625 10.1371/journal.pone.0150530. PubMed PMID: 26930612; PubMed Central PMCID:
626 PMC4773254.

627 36. Wang L, Chen H, Li J, Shu H, Zhang X, Wang Y, et al. Effector gene silencing mediated by
628 histone methylation underpins host adaptation in an oomycete plant pathogen. Nucleic Acids Res.
629 2020;48(4):1790-9. doi: 10.1093/nar/gkz1160. PubMed PMID: 31819959; PubMed Central
630 PMCID: PMC7039004.

631 37. van West P, Kamoun S, van 't Klooster JW, Govers F. Internuclear gene silencing in
632 *Phytophthora infestans*. Molecular cell. 1999;3(3):339-48. Epub 1999/04/13. doi: 10.1016/s1097-
633 2765(00)80461-x. PubMed PMID: 10198636.

634 38. van West P, Shepherd SJ, Walker CA, Li S, Appiah AA, Grenville-Briggs LJ, et al.
635 Internuclear gene silencing in *Phytophthora infestans* is established through chromatin
636 remodelling. Microbiology. 2008;154(Pt 5):1482-90. doi: 10.1099/mic.0.2007/015545-0. PubMed
637 PMID: 18451057.

638 39. Fang Y, Jang HS, Watson GW, Wellappili DP, Tyler BM. Distinctive nuclear localization
639 signals in the oomycete *Phytophthora sojae*. Front Microbiol. 2017;8:10. Epub 2017/02/18. doi:
640 10.3389/fmicb.2017.00010. PubMed PMID: 28210240; PubMed Central PMCID:

641 PMCPMC5288373.

642 40. Clark DP, Pazdernik NJ. Chapter 8 - Genomics and Gene Expression. In: Clark DP,
643 Pazdernik NJ, editors. Biotechnology (Second Edition). Boston: Academic Cell; 2016. p. 249-94.

644 41. Fang YF, Coelho MA, Shu H, Schotanus K, Thimmappa BC, Yadav V, et al. Long
645 transposon-rich centromeres in an oomycete reveal divergence of centromere features in
646 Stramenopila-Alveolata-Rhizaria lineages. bioRxiv. 2019:765032. doi: 10.1101/765032.

647 42. Raffaele S, Farrer RA, Cano LM, Studholme DJ, MacLean D, Thines M, et al. Genome
648 evolution following host jumps in the Irish potato famine pathogen lineage. Science.
649 2010;330(6010):1540-3. Epub 2010/12/15. doi: 10.1126/science.1193070. PubMed PMID:
650 21148391.

651 43. Wang Y, Tyler BM, Wang Y. Defense and Counterdefense During Plant-Pathogenic
652 Oomycete Infection. Annual review of microbiology. 2019;73:667-96. Epub 2019/06/22. doi:
653 10.1146/annurev-micro-020518-120022. PubMed PMID: 31226025.

654 44. Vleeshouwers VG, Raffaele S, Vossen JH, Champouret N, Oliva R, Segretin ME, et al.
655 Understanding and exploiting late blight resistance in the age of effectors. Annual review of
656 phytopathology. 2011;49:507-31. Epub 2011/06/15. doi: 10.1146/annurev-phyto-072910-095326.
657 PubMed PMID: 21663437.

658 45. Sabbadin F, Urresti S, Henrissat B, Avrova AO, Welsh LRJ, Lindley PJ, et al. Secreted
659 pectin monooxygenases drive plant infection by pathogenic oomycetes. science.
660 2021;373(6556):774-9. doi: 10.1126/SCIENCE.ABJ1342.

661 46. Gao C, Xu H, Huang J, Sun B, Zhang F, Savage Z, et al. Pathogen manipulation of
662 chloroplast function triggers a light-dependent immune recognition. *Proceedings of the National*
663 *Academy of Sciences of the United States of America*. 2020;117(17):9613-20. doi:
664 10.1073/PNAS.2002759117.

665 47. Yang L, McLellan H, Naqvi S, He Q, Boevink PC, Armstrong M, et al. Potato NPH3/RPT2-
666 like protein StNRL1, targeted by a *Phytophthora infestans* RXLR effector, is a susceptibility factor.
667 *Plant physiology*. 2016;171(1):645-57. doi: 10.1104/PP.16.00178.

668 48. Stefańczyk E, Sobkowiak S, Brylińska M, Śliwka J. Expression of the Potato Late Blight
669 Resistance Gene Rpi-phu1 and *Phytophthora infestans* Effectors in the Compatible and
670 Incompatible Interactions in Potato. *phytopathology*. 2017;107(6):740-8. doi: 10.1094/PHYTO-09-
671 16-0328-R.

672 49. Zheng L, Li C, Ma X, Zhou H, Liu Y, Wang P, et al. Functional interplay of histone lysine 2-
673 hydroxyisobutyrylation and acetylation in *Arabidopsis* under dark-induced starvation. *Nucleic*
674 *Acids Research*. 2021;49(13):7347-60. doi: 10.1093/nar/gkab536.

675 50. Tropberger P, Pott S, Keller C, Kamieniarz-Gdula K, Caron M, Richter F, et al. Regulation of
676 transcription through acetylation of H3K122 on the lateral surface of the histone octamer. *Cell*.
677 2013;152(4):859-72.

678 51. Grau-Bové X, Navarrete C, Chiva C, Pribasniig T, Antó M, Torruella G, et al. Comparative
679 proteogenomics deciphers the origin and evolution of eukaryotic chromatin. 2021.

680 52. Bourdareau S, Tirichine L, Lombard B, Loew D, Scornet D, Wu Y, et al. Histone

681 modifications during the life cycle of the brown alga *Ectocarpus*. *Genome Biology*. 2021;22(1). doi:
682 10.1186/s13059-020-02216-8.

683 53. Liu B, Liu Y, Wang B, Luo Q, Shi J, Gan J, et al. The transcription factor OsSUF4 interacts
684 with SDG725 in promoting H3K36me3 establishment. *Nature communications*. 2019;10(1). doi:
685 10.1038/s41467-019-10850-5.

686 54. Schotanus K, Soyer JL, Connolly LR, Grandaubert J, Happel P, Smith KM, et al. Histone
687 modifications rather than the novel regional centromeres of *Zymoseptoria tritici* distinguish core
688 and accessory chromosomes. *Epigenetics & Chromatin*. 2015;8(1). doi: 10.1186/s13072-015-
689 0033-5.

690 55. Cook DE, Kramer HM, Torres DE, Seidl MF, Thomma BP. A unique chromatin profile defines
691 adaptive genomic regions in a fungal plant pathogen. *Elife*. 2020;9. doi: 10.7554/eLife.62208.
692 PubMed PMID: 33337321.

693 56. Möller M, Schotanus K, Soyer JL, Haueisen J, Happ K, Stralucke M, et al. Destabilization of
694 chromosome structure by histone H3 lysine 27 methylation. *PLoS genetics*. 2019;15(4). doi:
695 10.1371/JOURNAL.PGEN.1008093.

696 57. Rodgers-Melnick E, Vera DL, Bass HW, Buckler ES. Open chromatin reveals the functional
697 maize genome. *Proceedings of the National Academy of Sciences*. 2016;113(22):E3177-E84. doi:
698 10.1073/pnas.1525244113.

699 58. Spruce C, Dlamini S, Ananda G, Bronkema N, Tian H, Paigen K, et al. HELLS and PRDM9
700 form a pioneer complex to open chromatin at meiotic recombination hot spots. *Genes &*

701 Development. 2020;34(5-6):398-412. doi: 10.1101/gad.333542.119.

702 59. Monroe JG, Srikant T, Carbonell-Bejerano P, Exposito-Alonso M, Weng M-L, Rutter MT, et
703 al. Mutation bias shapes gene evolution in *Arabidopsis thaliana*. *BioRxiv*. 2020. doi:
704 10.1101/2020.06.17.156752.

705 60. Chen H, Raffaele S, Dong S. Silent control: microbial plant pathogens evade host immunity
706 without coding sequence changes. *FEMS Microbiol Rev*. 2021. Epub 2021/01/14. doi:
707 10.1093/femsre/fuab002. PubMed PMID: 33440001.

708 61. Soyer JL, El Ghalid M, Glaser N, Ollivier B, Linglin J, Grandaubert J, et al. Epigenetic control
709 of effector gene expression in the plant pathogenic fungus *Leptosphaeria maculans*. *PLoS*
710 *genetics*. 2014;10(3):e1004227. Epub 2014/03/08. doi: 10.1371/journal.pgen.1004227. PubMed
711 PMID: 24603691; PubMed Central PMCID: PMC3945186.

712 62. Fouche S, Plissonneau C, Croll D. The birth and death of effectors in rapidly evolving
713 filamentous pathogen genomes. *Current opinion in microbiology*. 2018;46:34-42. doi:
714 10.1016/j.mib.2018.01.020. PubMed PMID: 29455143.

715 63. Roy S, Ernst J, Kharchenko PV, Kheradpour P, Negre N, Eaton ML, et al. Identification of
716 Functional Elements and Regulatory Circuits by *Drosophila* modENCODE. *Science*.
717 2010;330(6012):1787-97. doi: 10.1126/science.1198374.

718 64. Gerstein MB, Lu ZJ, Van Nostrand EL, Cheng C, Arshinoff BI, Liu T, et al. Integrative
719 Analysis of the *Caenorhabditis elegans* Genome by the modENCODE Project. *Science*.
720 2010;330(6012):1775-87. doi: 10.1126/science.1196914.

721 65. Shen Y, Yue F, McCleary DF, Ye Z, Edsall L, Kuan S, et al. A map of the cis-regulatory
722 sequences in the mouse genome. *Nature*. 2012;488(7409):116-20. doi: 10.1038/nature11243.

723 66. Qian B, Su X, Ye Z, Liu X, Liu M, Shen D, et al. MoErv29 promotes apoplastic effector
724 secretion contributing to virulence of the rice blast fungus *Magnaporthe oryzae*. *new phytologist*.
725 2021. doi: 10.1111/NPH.17851.

726 67. Chi H, Liu C, Yang H, Zeng W-F, Wu L, Zhou W-J, et al. Comprehensive identification of
727 peptides in tandem mass spectra using an efficient open search engine. *Nature biotechnology*.
728 2018;36(11):1059-61. doi: 10.1038/NBT.4236.

729 68. Sambrook J, Fritsch EF, Maniatis T. *Molecular Cloning: A Laboratory Manual* 2001 1/15/2001.

730 69. Wang X, Guo L. Sequence analysis and expression patterns of histone H2A variants in
731 *Phytophthora infestans*. *chinese journal of biotechnology*. 2016;32(11):1564-75. doi:
732 10.13345/J.CJB.160109.

733 70. Alzohairy AM. BioEdit: An important software for molecular biology. 2011.

734 71. Tamura K, Peterson DS, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: Molecular
735 Evolutionary Genetics Analysis Using Maximum Likelihood, Evolutionary Distance, and Maximum
736 Parsimony Methods. *Molecular biology and evolution*. 2011;28(10):2731-9.

737 72. Li HY, Wang HN, Jing MF, Zhu JY, Guo BD, Wang Y, et al. A *Phytophthora* effector recruits
738 a host cytoplasmic transacetylase into nuclear speckles to enhance plant susceptibility. *Elife*.
739 2018;7. doi: ARTN e40039
740 10.7554/eLife.40039. PubMed PMID: WOS:000450804600001.

741 73. Zhao Y, Li Y, Qiu M, Ma W, Wang Y. Generating Gene Silenced Mutants in *Phytophthora*
742 *sojae*. In: Ma W, Wolpert T, editors. *Plant Pathogenic Fungi and Oomycetes: Methods and*
743 *Protocols*. New York, NY: Springer New York; 2018. p. 275-86.

744 74. Huang L, Li X, Dong L, Wang B, Pan L. Profiling of chromatin accessibility identifies
745 transcription factor binding sites across the genome of *Aspergillus* species. *bmc biology*.
746 2021;19(1). doi: 10.1186/S12915-021-01114-0.

747 75. Zhang F, Chen H, Zhang XJ, Gao CY, Huang J, Lu L, et al. Genome Analysis of Two Newly
748 Emerged Potato Late Blight Isolates Sheds Light on Pathogen Adaptation and Provides Tools for
749 Disease Management. *Phytopathology*. 2021;111(1):96-107. Epub 2020/10/08. doi:
750 10.1094/Phyto-05-20-0208-Fi. PubMed PMID: WOS:000614285700009.

751 76. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nature methods*.
752 2012;9(4):357-9. doi: 10.1038/NMETH.1923.

753 77. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
754 Alignment/Map format and SAMtools. *bioinformatics*. 2009;25(16):2078-9. doi:
755 10.1093/BIOINFORMATICS/BTP352.

756 78. Ramírez F, Dündar F, Diehl S, Grüning BA, Manke T. deepTools: a flexible platform for
757 exploring deep-sequencing data. *Nucleic acids research*. 2014;42:187-91. doi:
758 10.1093/NAR/GKU365.

759 79. Chen C, Chen H, Zhang Y, Thomas HR, Frank MH, He Y, et al. TBtools: An Integrative
760 Toolkit Developed for Interactive Analyses of Big Biological Data. *Molecular plant*.

2020;13(8):1194-202. doi: 10.1016/J.MOLP.2020.06.009.

80. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al. Integrative genomics viewer. *Nature biotechnology*. 2011;29(1):24-6. doi: 10.1038/NBT.1754.

81. Gaspar JM. Improved peak-calling with MACS2. 2018.

82. Zhang Y, Liu T, Meyer CA, Eeckhoutte J, Johnson DS, Bernstein BE, et al. Model-based Analysis of ChIP-Seq (MACS). *genome biology*. 2008;9(9):1-9. doi: 10.1186/GB-2008-9-9-R137.

83. Quinlan AR, Hall IMJB. BEDTools: a flexible suite of utilities for comparing genomic features. 2010;26(6):841-2.

84. Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory requirements. *Nature methods*. 2015;12(4):357-60. doi: 10.1038/NMETH.3317.

85. Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nature biotechnology*. 2015;33(3):290-5. doi: 10.1038/NBT.3122.

86. Ernst J, Kellis M. ChromHMM: automating chromatin-state discovery and characterization. *Nature methods*. 2012;9(3):215-6. doi: 10.1038/nmeth.1906.

87. Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R Package for Comparing Biological Themes Among Gene Clusters. *omics a journal of integrative biology*. 2012;16(5):284-7. doi: 10.1089/OMI.2011.0118.

88. Li L, Stoeckert CJ, Roos DS. OrthoMCL: identification of ortholog groups for eukaryotic genomes. *genome research*. 2003;13(9):2178-89. doi: 10.1101/GR.1224503.

781 89. Allaire JJ, Gandrud C, Russell K, Yetman CJ. D3 JavaScript Network Graphs from R [R
782 package networkD3 version 0.4]. 2017.

783

Figure legends

Fig. 1 Histone H3 modifications identified in *P. infestans*. Schematic showing various *P. infestans* H3 (PiH3) PTMs that was detected by HPLC-MS/MS. Numbers at top of the PiH3 amino acid sequence indicate the positions of residues. Dots with different colors denote different types of PTMs, and the red ones at the top of K represent lysine trimethylation.

Fig. 2 Distribution of four histone H3 methylations and chromatin accessibility.

(A). Heatmap showed the distribution of ChIP-seq and ATAC-seq peak. upstream was defined as -800 bp before start code, downstream was defined as +400 bp after stop code. Submits of peaks in S6C Fig. were used. (B). line plot showed the distribution of four histone H3 methylations and chromatin accessibility across gene, -1 kb to +1 kb region were included. (C). Representative snapshot of H3 methylations and ATAC-seq statue. The snapshot was collected from IGV software. (D), Accessible chromatin, H3K4me3 and H3K36me3 are associated with highly expressed genes, meanwhile, H3K9me3 and H3K27me3 are associated silenced genes.

Fig. 3 Distribution of H3 methylations and chromatin accessibility reflects the bipartite genome structure.

Heatmap analyses showed higher H3K9me3 and H3K27me3 in GSR, meanwhile, higher ATAC-seq, H3K4me3 and H3K36me3 density in GDR. The dotted line highlights GSR. Gradient color represents gene number in gene density plot, and gradient color in other plot represent the average normalized ATAC-seq or H3 methylation signal.

Fig. 4 Histone H3 methylations and chromatin accessibility are associated with evolution of the protein-coding genes.

H3K9me3 and H3K27me3 were abundant in *P. infestans* specific genes (A) and fast evolved genes (B) with generally condensed chromatin, whereas higher H3K4me3 and H3K36me3 preferably associated with conserved genes which were located in more accessible chromatin regions. In (A), Pink, purple, blue and green represent eukaryote, oomycete, Phytophthora, and *P. infestans* specific gene groups, respectively. In (B), all protein-coding genes were divided into two categories based on their dN/dS ratios, their methylation and ATAC-seq density were compared. The number of genes in two groups were 14291 ($0 < dN/dS < 1$) and 1213 ($dN/dS > 1$), respectively.

Fig. 5 Virulence genes were enriched in OC state.

A. Chromatin state definitions, open chromatin (OC), strong transcription region (ST), H3K9me3 dominant repression region (H3K9DR), H3K27me3 dominant repression

region (H3K27DR), quiescent state (Quies). B-C, genome coverage and composition (emission probability) of histone marks and chromatin accessibility were presented. D, Enrichments probability of different gene categories in each chromatin states measured by chromHMM. E, Secretome genes were divided into CAZyme, CRN, SCR, NLP, RxLR and other genes, and Sankey plot showed chromatin state distribution of those gene groups. F, Percentage of gene in OC states compared with background group. Grey bars were calculated as (gene number of corresponding gene group)/ (total gene number of *P. infestans*). Purple color bars were calculated as (number of gene in corresponding gene groups associated with OC state)/ (total number of genes associated in OC state). P-value was calculated by Chi-squared test. G, Gene expression heatmap of OC state and ST state associated secreted virulence genes in mycelium (MY), 2-day post incubation (2 dpi) and 3-day post incubation (3 dpi) stages. Expression heatmap were normalized by $\log_2(\text{FPKM}+1)$ and clustered in row scale by Tbttools. H, Summary of percentage by gene upregulation fold change in OC state and ST state associated secreted virulence genes, the gene upregulation fold change was calculated using row FPKM value.

S1 Fig. Sequences analysis of *P. infestans* H3 homologs.

(A), Nucleotide sequences alignment of four *P. infestans* H3 homologs. (B), Amino acid sequence alignment of four *P. infestans* H3 homologs.

S2 Fig. Phylogenetic and sequence analysis showed conserved Phytophthora H3 homologs.

(A), Phylogenetic analysis of Phytophthora H3 homologs. H3 homologs from human (HsH3.1/accession number: NP_003520.1, HsH3.2/ NP_066403.2, HsH3.3/NP_002098.1, HsCenP-A/NP_001035891.1), *Arabidopsis thaliana* (AtH3.1/NP_563838.1, AtH3.3/ NP_001329167.1, AtCenH3/NP_009564.1) and budding yeast (ScH3/NP_009564.1) were included in the study., H2A was set as an outgroup. (B), Alignment of H3 variants from *P. infestans* (PITG_05675 and PITG_03551) and *P. sojae* H3 (Ps_284752, Ps_322070 and Ps_476994) with HsH3.1, HsH3.2, AtH3.1, ScH3 and NcH3.

S3 Fig. PTM variants in Phytophthora H3.

(A), amino acid substitutions of two H3 forms were compared and different PTMs were marked. The numbers indicated amino acid position, amino acids at each site were listed and different colors indicated PTMs. (B), Individual mass spectrums of five PTM variants were shown.

S4 Fig. Summary of H3 lysine acetylation and methylation among seven organisms.

Highly conserved lysine were listed, red color indicated this PTM was detected or reported, blue color indicated this PTM was not detected or reported. We used grey color to mark the residue at 53 in human, for this site mutated to R.

S5 Fig. H3K4me3, H3K9me3, H3K27me3 and H3K36me3 were detected in *P. infestans*.

(A), MS spectra of H3K4me3, H3K9me3, H3K27me3 and H3K36me3 in *P. infestans*. (B), H3, H3K4me3, H3K9me3, H3K27me3 and H3K36me3 were detected by WB. *P. infestans*, *P. sojae* and *M. oryzae* total protein were detected by five different antibodies, and prokaryotic expressed H3 protein of *P. sojae* is the negative control that cannot be modified.

S6 Fig. The good quality and correlation between biological replicates of different histone modifications and ATAC-seq in this study.

(A). The overall analysis of ChIP-seq data and ATAC-seq data. (B). Highly repeatability of two replicates. The genome was divided into 2kb bins and RPKM values of each bin were used to calculate Pearson correlation coefficients. (C). Highly overlapped peak of two replicates.

S7 Fig. TE region have higher H3K9me3 and H3K27me3.

(A). Methylation level of gene and TE were compared. Methylation level was calculated as $\log_2(\text{IPRPKM}/\text{inputRPKM})$. P values are calculated with the two-sample Kolmogorov-

Smirnov test. (B). H3K9me3 and H3K27me3 are preferentially associated with LTR and non-LTR type transposon in *P. infestans*. Intersected percentage of peaks length and TE length were calculated.

S8 Fig. PCA and heatmap clustering revealed overall distinct features between ATAC-seq and ChIP-seq reads.

PCA analysis (A) and heatmap clustering (B) of bw file generated by deepTools. each dot represents one replicates. The Pearson correlation computation was used in (B).

S9 Fig. Unique role of chromatin accessibility in chromatin state defining compared with those four PTMs by peak analysis.

ATAC peaks were divided into ATAC peak-alone, ATAC and H3K4me3 peak cobound, and etc. The number of peaks were 10666, 3334, 1311, 2496, 1496, 331, 486 and 663, respectively. Overlapped peaks defined in S6C Fig. were used here.

S10 Fig. Genes under positive selection have higher H3K27me3 and H3K9me3 in *P. infestans*.

(A), Conserved genes have lower dN/dS value, and *P. infestans* specific gene is under highest positive selection. P values are calculated with the two-sample Kolmogorov-Smirnov test. (B). GSR has higher dN/dS ratio. Gradient color represents dN/dS ratio. (C), Genes under positive selection have higher H3K9me3 and H3K27me3. RPKM value from -1 kb to 1 kb region of each gene were collected and the methylation level was

896 calculated as $\log_2(\text{IP}_{\text{RPKM}}/\text{input}_{\text{RPKM}})$. P values are calculated with the two-sample
897 Kolmogorov-Smirnov test.

898 **S11 Fig. Comparison of the GO enrichment result of methylated genes.**

899 **S12 Fig. Genes encoding secretome protein and RxLR effectors harbor higher**
900 **H3K9me3 and H3K27me3, otherwise, ATAC-seq, H3K4me3 and H3K36me3 is lowly**
901 **on these type of genes.**

902 (A). comparison of ATAC-seq and four histone methylation signal in core, secretome and
903 RxLR genes. P values are calculated with the two-sample Kolmogorov-Smirnov test. (B),
904 Representative snapshot of silenced pectinase and RxLR gene cluster. The snapshot
905 was collected from IGV software, and specific loci were marked by blue frame.

906 **S13 Fig. GO enrichment result of genes in OC, ST, H3K9DR and H3K27DR**
907 **chromatin states.**

908

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5