1 Transcriptome analysis of diverse *Plasmodium falciparum* clinical isolates
2 identifies genes correlating with highly variable expression of merozoite
3 surface protein MSPDBL2

4

5 Suzanne E. Hocking ¹, Lindsay B. Stewart ¹, Sarah J. Tarr ¹, Aline Freville ¹, Kevin K. A. Tetteh
6 ¹, Ambroise D. Ahouidi ², Alfred Amambua-Ngwa ³, Mahamadou Diakite ⁴, Gordon A.
7 Awandare ⁵, David J. Conway ^{1*}

8

9 ¹ Department of Infection Biology, London School of Hygiene and Tropical Medicine, Keppel
10 St, London WC1E 7HT

11 ² Le Dantec Hospital, Université Cheikh Anta Diop, Dakar, Senegal

12 ³ MRC Unit The Gambia at London School of Hygiene and Tropical Medicine (LSHTM), Fajara,
13 Banjul, The Gambia

14 ⁴ Malaria Research and Training Center, University of Bamako, Mali

15 ⁵ West African Centre for Cell Biology of Infectious Pathogens, Department of Biochemistry,
16 Cell and Molecular Biology, University of Ghana, Legon, Ghana

17

18 * Corresponding : Email david.conway@lshtm.ac.uk ; Phone + 44 20 7927 2331.

19

20 Running title: Variable merozoite protein and transcriptome

Abstract

The merozoite surface protein MSPDBL2 of *Plasmodium falciparum* is under strong balancing selection and is a target of naturally acquired antibodies. Remarkably, MSPDBL2 is expressed in only a minority of mature schizonts of any cultured parasite line, and *mSPdbl2* gene transcription increases in response to overexpression of the gametocyte development inducer GDV1, so it is important to understand its natural expression. Here, MSPDBL2 in mature schizonts was analysed in the first *ex vivo* culture cycle of 96 clinical isolates from four populations with varying levels of infection endemicity in different West African countries, by immunofluorescence microscopy with antibodies against a conserved region of the protein. In most isolates, less than 1% of mature schizonts were positive for MSPDBL2 (median of 0.6% overall), but the frequency distribution was highly skewed as nine isolates had more than 3% schizonts positive and one had 73% positive. To investigate whether expression of other gene loci correlated with MSPDBL2 expression, whole transcriptome sequencing was performed on schizont-enriched material from 17 of the clinical isolates with a wide range of proportions of schizonts positive. Transcripts of particular parasite genes were highly significantly positively correlated with MSPDBL2 positivity in schizonts as well as with *mSPdbl2* gene transcript levels, with overrepresentation of genes previously implicated as likely to be involved in gametocytogenesis, but not including the gametocytogenesis master regulator *ap2g*. Although MSPDBL2 is apparently not directly involved in sexual commitment, it marks a co-occurring developmental subpopulation that may be functionally distinct within blood stage infections.

42 Introduction

43

44 The *Plasmodium falciparum* merozoite surface protein MSPDBL2 is one of two MSP3-like
 45 proteins with a duffy-binding like (DBL) domain, expressed in schizonts and co-localised with
 46 MSP1 on the merozoite cell surface (1, 2). MSPDBL2 is not directly membrane-bound, but
 47 appears complexed with MSP1, and can bind to the surface of uninfected erythrocytes (1),
 48 suggesting a potential role in invasion. The *mspdbl2* gene occurs at a single locus on
 49 chromosome 10 with an intact coding sequence in all *P. falciparum* isolates, although it has
 50 multiple stop codons in the related chimpanzee parasite *P. reichenowi* indicating that it is
 51 not functional in that species (3, 4), and an orthologue only exists within members of the
 52 *Laverania* sub-genus but not in other malaria parasites (5). The gene is highly polymorphic
 53 within *P. falciparum*, and analysis of allele frequency distributions in endemic populations
 54 indicates that diverse alleles are maintained by strong balancing selection (3, 6, 7),
 55 suggesting that it may be a target of immune selection.

56

57 MSPDBL2 is a target of naturally acquired antibody responses, and an endemic population
 58 cohort study indicates some association with reduced prospective risk of clinical malaria (8),
 59 while another study showing purified IgG inhibits parasites in culture (9). However, the
 60 vaccine candidacy of MSPDBL2 is uncertain, as it is not only polymorphic but also highly
 61 variable in expression. Analysis of schizont-rich cultures of clinical isolates and long-term
 62 adapted *P. falciparum* laboratory lines has revealed very low *mspdbl2* transcript levels in
 63 most isolates assayed by RT-qPCR (6), or by whole transcriptome analysis (10). Strikingly,
 64 MSPDBL2 protein expression is restricted to a small proportion of mature schizonts in each
 65 laboratory-adapted parasite line tested by immunofluorescence with specific antibodies (6),
 66 although frequencies of MSPDBL2 expression in schizonts of clinical isolates have not been
 67 reported. Consistent with being expressed in only a small minority of parasites,
 68 experimental disruption of the *mspdbl2* gene does not affect overall asexual parasite growth
 69 rates in culture (11).

70

71 If MSPDBL2 is restricted to a functionally-important parasite subpopulation, it might still be
 72 a target to be considered for vaccination. Interestingly, MSPDBL2 has been shown to bind to
 73 the Fc region of IgM, although the function of this is unknown (12). Separately,

overexpression of *mspdbl2* has been reported to enhance parasite survival in the presence of some antimalarial drugs in culture (13). Significantly, it has been suggested that *mspdbl2* expression may be correlated with parasite sexual differentiation, as over-expression of the *gdv1* gene in an engineered parasite clone results in marked increase in transcription of *mspdbl2* as well as genes known to be involved in switching to gametocyte development (14). This indicates the importance of studying variation in *mspdbl2* gene expression as well as MSPDBL2 protein expression in schizonts of clinical isolates, and explore whether this is associated with markers of parasite commitment to gametocyte development *in vivo*. It is clear that there is great variation in proportions of parasites committed to gametocyte development among different infections from a single endemic area (15), and transcript markers of ring stage parasites that will develop as gametocytes (16), but apart from the master regulator transcription factor AP2-G (17) there are no confirmed markers of sexually committed schizonts in the previous cycle (18, 19). This study investigates the frequency distribution of expression of MSPDBL2 protein in schizonts of clinical isolates from endemic populations, and tests for correlations with other genes expressed in clinical isolate transcriptomes, including those previously implicated as associated with the gametocytogenesis pathway.

91

92 **Materials and Methods**

93 ***P. falciparum* clinical isolates**

For analysis of parasite phenotypic and expression variation in natural infections, clinical malaria cases attending local health facilities in Ghana (Kintampo), Mali (Nioro du Sahel), Senegal (Pikine), and The Gambia (Basse) were investigated. The level of malaria infection endemicity varied among these different populations in West Africa, being higher at the sites in Ghana and Mali than the sites in Senegal and The Gambia, which is also reflected in more complex mixed genome *P. falciparum* infections at the more highly endemic sites, as previously shown for these populations (20). Patients aged between 2 and 14 years were eligible if they had uncomplicated clinical malaria, had not taken antimalarial drugs in the 72 hours preceding sample collection and tested positive for *P. falciparum* malaria by lateral flow rapid diagnostic test and slide microscopy. Written informed consent was obtained from parents or legal guardians of participating children and additional assent was received from participating children. Up to 5ml of venous blood was collected in heparinised anti-

coagulation BD Vacutainer® tubes (BD Biosciences), and a proportion of the erythrocytes were cryopreserved in glycerolyte and stored at -80°C or in liquid nitrogen before shipment on dry ice to the London School of Hygiene and Tropical Medicine for subsequent culture and laboratory analysis. Ethical approval for the collection and analysis of clinical samples was granted by the Ethics Committees of the Ministry of Health in Senegal, the Ministry of Health in Mali, the Ghana Health Service, the Noguchi Memorial Institute for Medical Research, University of Ghana, Kintampo Health Research Centre, MRC Gambia, and the London School of Hygiene and Tropical Medicine.

***P. falciparum* schizont preparations from the first cycle of *ex vivo* culture**

The clinical blood samples were thawed in batches of eight at a time and introduced into culture *ex vivo*, all isolates being processed in culture in a single laboratory, as described for a previous study of the first *ex vivo* malaria parasite generation from similar clinical samples (21). Giemsa-stained thin blood films were prepared for each isolate initially upon thawing, and later during the second day of culture to assess the developmental progression of parasites into schizogony. Isolates containing schizonts in culture on the second day after thawing were enriched for schizonts by magnetic MACS® separation, and parasites were then allowed to mature in the presence of E64 for 4 hours to prevent schizont rupture, following which cells were harvested by centrifugation, using methods similar to those previously applied to studies of schizonts in continuously cultured parasite lines (10). Erythrocytes containing matured schizonts were prepared for immunofluorescence assays by washing and resuspending in 1% BSA and spotting into individual wells of 12-well slides (Hendley-Essex), air dried and stored with desiccant at -40°C until assay, as performed in previous analysis of schizonts in cultured parasite lines (6).

Analysis of MSPDBL2 expression in schizonts by immunofluorescence

Staining of schizonts positive for MSPDBL2 by immunofluorescence was performed with the identical method previously used on laboratory-adapted *P. falciparum* isolates, involving incubation with polyclonal mouse serum specific for the conserved N-terminal portion of MSPDBL2, and goat anti-mouse IgG Alexa Fluor® 555 secondary antibody, with Vectashield® mounting fluid containing DAPI to visualise nuclei (6). For each isolate, approximately 1000 mature schizonts (each containing at least 8 nuclei) were counted using DAPI and scored for

MSPDBL2 expression using a manual Leica fluorescence microscope with a 100x objective. MSPDBL2 expression was always clearly brightly positive or entirely negative in each individual mature schizont, so that counts of numbers and proportions positive in each preparation were recorded, using same process as described previously (6).

RNA-seq of schizont-enriched samples of clinical isolates

Parasite schizont-enriched material from individual isolates was stored in either TRIzol® or RNA^{later}™ (Thermo Fisher Scientific, MA, USA), and RNA was extracted by phenol-chloroform and cleaned up using the NucleoSpin® RNA XS extraction kit (Macherey-Nagel, Germany). Samples showing successful RNA extraction after checking by Bioanalyzer electrophoresis were reverse transcribed and cDNA amplified using the SmartSeq® v4 Ultra® Low Input RNA Kit for Sequencing (Takara Bio. Inc., Shiga Prefecture, Japan). Successfully amplified samples were prepared for paired end short-read sequencing using the Nextera XT Library Prep kit (Illumina, California, USA), individual libraries being pooled in equimolar amounts at 4nM with up to 12 per pool, and sequencing was performed on an Illumina MiSeq using the 150-cycle MiSeq reagent kit v3. RNA from isolate INV236 which had the highest proportion of MSPDBL2-positive schizonts (73%) was prepared and sequenced as a priority, and following this RNA was extracted from 38 other isolates with varying proportions of MSPDBL2-positive schizonts, of which 25 showed expected cDNA size range profile after reverse transcription and amplification, and 16 of these were selected for sequencing as they had RNA quality RIN score > 6. This yielded a set of 17 isolates with RNA-seq data and matched IFA data on proportions of MSPDBL2-positive schizonts.

Following procedures previously used for RNA-seq analysis of schizont-enriched *P. falciparum* cultures of other isolates (10), whole transcriptome short read sequence data were assembled by alignment mapping to the *P. falciparum* 3D7 version 3.0 reference genome (22) using HISAT2 (23). Gene transcript levels were assessed using the FPKM metric (Fragments Per Kilobase of transcript per Million mapped reads, the number of reads mapping to each gene normalised for the size of the sequencing library and for gene length). Data were analysed using the R package DESeq2 (24), using a masked GFF annotation file that removed the *var*, *rifin*, and *stevor* gene families from analysis, as described for analysis of previous schizont transcriptome data (10). In addition to exclusion of these three sub-

telomeric gene families from analysis, portions of other protein-coding genes that show high allelic diversity (including the highly polymorphic central region of the *mspdbl2* gene) were masked to ensure mapping occurred only in conserved regions of those genes to minimise allelic bias among samples, as previously described for other data (10). Prior to analysis, three independent studies including proteomic or transcriptomic analyses were consulted (14, 25, 26) to identify *P. falciparum* genes considered to be potentially associated with gametocytogenesis, enabling compilation of a list of 119 genes (Supplementary Table S1) used as a set for conducting exploratory correlative analyses of the RNA-seq data within this study. To avoid discovery bias in the present study this needed to be a static list fixed prior to analysis, but is not a reference list for other studies as ongoing research means that any such compilations should be updated and can be subject to different criteria.

Statistical analyses

Tests for significance of correlations between different variables (including proportions of schizonts positive for MSPDBL2, and individual gene FPKM values), or estimations of odds ratios and significance of associations between categorical variables, were conducted using a combination of R, Epi-Info and Prism software. Differential gene expression analysis was carried out in DESeq2 (24), focusing on the distributions of derived FPKM values for each gene as defined above.

Results

Wide variation in proportions of schizonts expressing MSPDBL2 in clinical isolates

MSPDBL2 protein expression was examined in mature schizonts (each with at least eight nuclei) developed in the first cultured *ex vivo* cycle of each of 96 clinical isolates sampled from malaria patients in four different endemic countries in West Africa. In most isolates, less than 1% of all mature schizonts were positive for MSPDBL2 (overall median of 0.6%), but the frequency distribution was highly skewed as some had much higher proportions (Figure 1 and Supplementary Table S2). Nine isolates had more than 3% schizonts positive, including one that had 73% positive. There were no significant differences in the distributions among different countries (Kruskal-Wallis test and pairwise Mann-Whitney tests non-significant), although the two isolates with highest proportions were from Senegal

(Figure 1). The overall distribution was compared to that previously reported for a panel of 12 long-term laboratory-adapted *P. falciparum* lines originally isolated from more diverse sources (6), and this was not significantly different (Mann-Whitney test, $P = 0.52$). This shows that a wide range of MSPDBL2-positive schizonts, with most isolates having very low proportions positive, is a natural feature of expression in endemic *P. falciparum* populations rather than one which has been selected by laboratory culture.

Transcriptomes of schizont-enriched *ex vivo* cultures with a wide range in proportions of MSPDBL2-positive schizonts

To explore whether particular parasite gene transcripts are associated with the proportions of MSPDBL2-positive frequencies, whole transcriptome RNA-seq analysis was performed for 17 of the clinical isolates that had sufficient schizont-enriched material, representing a wide range of MSPDBL2 expression (zero to 73% of schizonts positive, Supplementary Table S1). Sequencing of the cDNA libraries yielded a mean of 2.1×10^6 Illumina short reads for each isolate, and most of these reads aligned to the 3D7 reference genome sequence (Supplementary Table S3), enabling read depth analysis of relative expression of individual genes after normalisation for the total number of reads for each isolate (Supplementary Figure S1). The FPKM values across all genes were first compared to published RNA-seq data from tightly synchronised *P. falciparum* 3D7 parasites sampled at seven timepoints post invasion (0, 8, 16, 24, 32, 40, and 48 hours, the last of which may include some next cycle reinvasion) (27), confirming that Spearman's rank correlations expression profiles in all samples had strongest correlations with schizont stage parasites (13 isolates correlated most strongly with the 40-hour timepoint and four isolates with the 32-hour timepoint, Supplementary Figure S1).

Gene transcripts correlating with variable proportions of MSPDBL2-positive schizonts among clinical isolates

Among the 17 isolates with RNA-seq data, individual gene FPKM relative expression values were tested for correlation with MSPDBL2 IFA expression. To scan for significantly correlated genes, a P value cut off of <0.001 was used which identified 52 genes with increased expression (including the *mispdb2* gene which was by far the most positively

correlated as expected, Supplementary Table S4), and 130 genes with negatively correlated expression (Supplementary Table S5). Aside from *mispdbl2* itself, 12 (24%) of the other 51 genes with higher expression were previously listed as having known or suspected roles in gametocytogenesis, whereas of the 130 genes with lower expression, only 10 (7%) were listed with known or suspected gametocytogenesis involvement. An odds ratio of 3.7 (95% CI, 1.5 - 9.2, $P = 0.0054$) on these proportions indicates a significant skew in gametocytogenesis-related genes being more likely to be positively rather than negatively correlated with proportions of MSPDBL2-positive schizonts.

Using a more stringent correlation value cut off of $P < 10^{-4}$ to focus on genes having most highly significant correlations with proportions of schizonts expressing MSPDBL2, 19 genes are identified as positively correlated, of which 9 (47%) were previously indicated as having known or suspected roles in gametocytogenesis (Table 1). At this level of correlation significance, 51 genes are negatively correlated (Supplementary Table S3), of which only one (2%) was previously indicated as gametocytogenesis-related. This indicates a very highly significant skew in gametocytogenesis-related genes being positively rather than negatively correlated with proportions of MSPDBL2-positive schizonts, yielding an odds ratio of 45.0 (95% CI, 5.1 – 396.0, $P = 1.2 \times 10^{-5}$).

Independently, a previous study has investigated the transcriptomic profiles of cultured schizonts of the transgenic *P. falciparum* parasite line 164/TdTom, comparing preparations enriched for sexually-committed versus asexually-committed schizonts (28). The data from this study on the PlasmoDB genomics resource site (29) shows relative transcript levels accessible for all except one of the 19 genes that were most highly positively correlated with MSPDBL2 expression in the present study. Fifteen (83%) of these 18 genes had higher expression in the sexually-committed schizont preparation compared to the asexual schizont preparation, a significantly positive skew compared to random expectations ($P < 0.05$).

Genes expressed in correlation with *mispdbl2* gene expression in schizont-enriched cultures of clinical isolates

To complement the above scan based on proportions of MSPDBL2-positive schizonts, the varying transcript levels of *mispdbl2* (FPKM values) among the clinical isolates were analysed to scan for other genes with correlated expression. Using a correlation significance value of $P < 0.001$ as cut-off identified 41 genes with positively correlated expression (Supplementary Table S6), many of which were also correlated with proportions of MSPDBL2-positive schizonts (16 correlated at $P < 0.001$, 30 correlated at $P < 0.01$), and 31 genes had negatively correlated expression (Supplementary Table S7).

Of the 41 genes positively correlated with *mispdbl2* transcript expression, 11 (27%) were previously identified as potentially gametocytogenesis-related, in comparison to only 3 (10%) of the 31 negatively correlated genes, giving an odds ratio of 3.4 (95% CI, 0.9 – 13.6, $P = 0.06$). Using a higher level of cut-off for correlation significance ($P < 10^{-4}$), 19 genes were positively correlated with transcript levels of *mispdbl2* (Table 2), of which 8 (42%) were previously identified as potentially gametocytogenesis-related, in comparison to 2 (15%) of 13 genes negatively associated with *mispdbl2*, giving an odds ratio of 4.0 (95% CI, 0.7 – 23.3, $P = 0.11$). In summary, the analysis based on *mispdbl2* transcript levels gives broadly similar results to the analysis based on MSPDBL2 positive schizont proportions, but the excess proportions of positively correlated gametocyte-related genes are less significant.

Discussion

This study shows that *P. falciparum* in diverse clinical isolates have a wide range of MSPDBL2 expression positivity in mature schizonts in the first *ex vivo* cycle of development, with most isolates having very low proportions positive. The frequency distribution is similar to that previously described for a smaller number of *P. falciparum* laboratory-adapted lines that had been cultured for many years (6), indicating that this is not a result of selection by culture adaptation. Furthermore, the frequency distribution was similar in isolates from each of the four populations sampled here, which have different levels of malaria infection endemicity within West Africa (20, 30), indicating that parasite populations maintain the wide range of MSPDBL2 expression variation within different endemic environments.

Results of the RNA-seq analyses here indicate that MSPDBL2 protein and gene expression in schizonts in the first cycle of development from clinical isolates is significantly correlated with the expression of other particular genes within the *ex vivo* cultures. Particularly, many of the most strongly correlated genes were previously implicated as having known or suspected involvement in the process of gametocytogenesis. This is consistent with expectations from a functional study on effects of the gametocyte development gene *gdv1* in assays of an engineered parasite line with highly induced expression of GDV1 (14), which showed significantly increased transcription of *ap2g* as expected, and also *mispdbl2*, as well as a PHISTa gene (PF3D7_1477700). A separate study indicated that levels of the protein encoded by *pfg14_748* increase as parasites develop along the gametocytogenesis pathway, being detectable alongside the early gametocytogenesis marker Pfs16 in parasite cultures before gametocytes were observed to develop (31). In the present study, *pfg14_748* had expression strongly correlated with *mispdbl2*, but although it was previously shown to be induced by GDV1 (14) it is apparently not dependent on expression of AP2-G (25). Several other genes which correlated with *mispdbl2* expression in the present study (including the nucleoprotein gene *nup116*, and the early gametocyte development marker *gexp02*) have been identified as being upregulated by *ap2-g* (25), but *mispdbl2* was not itself identified to be upregulated by *ap2-g*.

Genes correlating with MSPDBL2 expression in clinical isolates include members of the GEXP family encoding proteins involved in protein export occurring during gametocytogenesis (26), particularly *gexp02* (PF3D7_1102500) and *gexp04* (PF3D7_1372100). As well as *gexp02* being known to result from induced *gdv1* expression (14), it has also been shown that *gexp02* is de-repressed in parasites which have conditional knock out of heterochromatin protein 1 (HP1), presumably due to the resulting activation of *gdv1* (32). However, the correlating expression with *gexp02* and other gametocytogenesis genes here does not indicate that *mispdbl2* is active in the process of sexual commitment or that it is a specific marker. It should be noted that relevant analysis of *gdv1* transcript levels are not possible from the double-stranded cDNA transcriptome data in the present study, as *gdv1* is usually repressed by abundant antisense transcripts initiated from the 3'-intergenic region (14, 33, 34), and ratios of sense to antisense transcripts could only be determined by strand-specific sequencing.

Clearly, if *mSPDBL2* is directly involved in gametocytogenesis, it would be expected that *ap2-g* would be identified as differentially expressed in isolates with higher *mSPDBL2* expression. As this was not the case in this study, it is alternatively possible that the expression of *mSPDBL2* occurs in some parasites in parallel to the sexual commitment process. It is important to note that the correlation with some involved in gametocytogenesis in the present study is at the bulk transcriptome level, so that *mSPDBL2* expression might not be within the same individual parasites, but ones that tend to occur in the same bulk population. The only published single cell transcriptome data on *P. falciparum* schizonts are from a laboratory clone in which hardly any schizonts express MSPDBL2, so are not informative on co-expression between the *mSPDBL2* gene and others (6, 35). Future insights from single cell transcriptome data will require analysis of *P. falciparum* isolates that have a substantial proportion of schizonts expressing MSPDBL2.

Functional studies would be required to determine whether MSPDBL2 is in any way associated with the sexual commitment process, or whether it is only correlated at the population level within infections or within cultures due to being co-incidentally upregulated by *gdfv1* (14). As MSPDBL2 is a merozoite surface protein expressed in late schizonts, the current study analysed transcriptomes of schizont-enriched preparations from *ex vivo* culture, as focused on in few other studies of *P. falciparum* clinical isolates (10, 36). Other studies have analysed parasite transcriptomes from the earlier stages of intra-erythrocytic development that are present in peripheral circulation (37-40), and a temporal analysis of parasite development through to mature schizonts may be needed to identify gene products with which MSPDBL2 is functionally linked, and resolve whether it is a marker of an important parasite subpopulation that could be targeted for vaccination.

Acknowledgements

We are grateful to the malaria patients and clinical staff for participation in the study. The sample collection was facilitated by staff at Kintampo Health Research Centre in Ghana, at the National Institute for Public Health in Guinea, at Nioro du Sahel Health Centre in Mali and at Pikine Health Centre in Senegal. We appreciate the support of colleagues at the Medical Research Council Unit in The Gambia, the Laboratory of Bacteriology and Virology,

359 Le Dantec Hospital in Senegal, The University of Bamako in Mali, Noguchi Memorial Institute
 360 for Medical Research and the University of Ghana, and the London School of Hygiene and
 361 Tropical Medicine in enabling this work. This study was supported by an ERC Advanced
 362 Award (grant AdG-2011-294428), a Leverhulme-Royal Society Africa Award (grant
 363 AA110050), a BBSRC PhD studentship within the London Interdisciplinary Doctoral Training
 364 Programme (www.lido-dtp.ac.uk), and an MRC project grant (MR/S009760/1).
 365
 366

Figure 1. Variable proportions of *P. falciparum* schizonts expressing MSPDBL2 in *ex vivo* cultures of 96 clinical isolates from four West African countries. Samples were from malaria patients in Ghana n=42, The Gambia n=8, Senegal n=25, and Mali n=21. In each isolate, approximately 1000 mature schizonts containing eight or more nuclei were scored in immunofluorescence assays using polyclonal mouse serum against a conserved N-terminal region of MSPDBL2. Eighteen of the isolates had no schizonts positive, and the median across all isolates was only 0.6%, but proportions were highly skewed and nine isolates had more than 3% of schizonts positive. Although the two isolates with highest proportions were from Senegal, there were no significant differences in the overall distributions among different countries by non-parametric rank sum tests.

378

379 **Figure 2. Transcriptome analysis identifies *P. falciparum* genes showing correlation with**
 380 **MSPDBL2 expression in *ex vivo* cultured clinical isolates.** RNA-seq was performed on
 381 schizont-enriched cultures of 17 West African clinical isolate in the first *ex vivo* cycle, and
 382 across the isolates FPKM transcript levels of each gene were tested for correlation with the
 383 proportion of schizonts expressing MSPDBL2. Blue shading indicates those that have the
 384 most significant differential gene expression (DGE, $P < 10^{-4}$), and genes positively correlated
 385 at this level of significance are listed in Table 1. A broader set of genes correlated positively
 386 at a slightly lower level of individual significance ($P < 0.001$, Supplementary Table S4), and
 387 those that are negatively correlated at that level of significance are also listed separately
 388 (Supplementary Table S5).

Table 1. *P. falciparum* genes with most highly significantly increased expression ($P < 10^{-4}$) correlating with proportions of MSPDBL2-positive schizonts in clinical isolates.

Gene ID	P-value	Gene product description
<i>PF3D7_1036300</i>	5.4×10^{-25}	<i>duffy binding-like merozoite surface protein 2 (mspdbl2)</i>
PF3D7_1476600	6.7×10^{-08}	<i>Plasmodium</i> exported protein
PF3D7_1474000	7.6×10^{-08}	conserved <i>Plasmodium</i> protein
PF3D7_1102500	1.3×10^{-07}	<i>gexp02</i> , <i>Plasmodium</i> exported protein (PHISTb)
PF3D7_1461800	3.0×10^{-06}	conserved <i>Plasmodium</i> protein
PF3D7_1445700	1.4×10^{-05}	conserved <i>Plasmodium</i> protein
PF3D7_0814200	1.7×10^{-05}	DNA/RNA-binding protein Alba 1
PF3D7_1466200	2.0×10^{-05}	early gametocyte enriched phosphoprotein EGXP
PF3D7_0114000	2.1×10^{-05}	GEXP06, exported protein family 1
PF3D7_1372100	2.1×10^{-05}	<i>Plasmodium</i> exported protein (PHISTb)
PF3D7_0215000	2.2×10^{-05}	acyl-CoA synthetase
PF3D7_1362700	2.6×10^{-05}	conserved <i>Plasmodium</i> protein
PF3D7_1473700	2.8×10^{-05}	nucleoporin NUP116/NSP116, putative
PF3D7_0829400	2.8×10^{-05}	prolyl 4-hydroxylase subunit alpha, putative
PF3D7_1027300	5.2×10^{-05}	Peroxiredoxin, nuclear protein
PF3D7_0515000	6.2×10^{-05}	pre-mRNA-splicing factor CWC2, putative
PF3D7_1346800	6.2×10^{-05}	Pfs47, 6-cysteine protein
PF3D7_1132600	6.4×10^{-05}	pre-mRNA-splicing factor 38A, putative
PF3D7_1477700	9.1×10^{-05}	<i>Pfg14.748</i> , <i>Plasmodium</i> exported protein (PHISTa)
PF3D7_1431400	9.9×10^{-05}	surface-related antigen SRA

The most highly correlated transcript is *mspdbl2* itself, as expected. Of the other 19 most significantly correlated genes, 9 (47%) highlighted in bold were previously implicated as potentially gametocytogenesis-related (Supplementary Table S1). Additional genes that positively correlated at the $P < 0.001$ significance level are listed in Supplementary Table S4.

Table 2. *P. falciparum* genes with most significantly increased expression ($P < 10^{-4}$) correlating with *mSPDBL2* transcript levels measured by FPKM in transcriptomes of clinical isolates.

Gene ID	P-value	Gene product description
PF3D7_0114000*	2.6×10^{-9}	GEXP06, exported protein family 1
PF3D7_1362700*	2.6×10^{-8}	conserved <i>Plasmodium</i> protein
PF3D7_1466200*	2.2×10^{-7}	early gametocyte enriched phosphoprotein EGXP
PF3D7_1472200	3.1×10^{-7}	histone deacetylase, putative
PF3D7_1467600*	6.0×10^{-7}	conserved <i>Plasmodium</i> protein
PF3D7_0214300*	1.1×10^{-6}	conserved <i>Plasmodium</i> protein
PF3D7_1027300*	2.3×10^{-6}	peroxiredoxin
PF3D7_1461800*	2.5×10^{-6}	conserved <i>Plasmodium</i> protein
PF3D7_1473700*	2.9×10^{-6}	nucleoporin NUP116/NSP116, put
PF3D7_1361200*	8.4×10^{-6}	conserved <i>Plasmodium</i> protein
PF3D7_1474000*	1.0×10^{-5}	conserved <i>Plasmodium</i> protein
PF3D7_0501400	1.5×10^{-5}	interspersed repeat antigen
PF3D7_0801900	2.0×10^{-5}	lysine-specific histone demethylase, put
PF3D7_1408200	4.9×10^{-5}	AP2 domain transcription factor AP2-G2
PF3D7_0207800	5.3×10^{-5}	serine repeat antigen 3
PF3D7_1235300	7.1×10^{-5}	CCR4-NOT transcription complex s4, putative
PF3D7_0519500	7.4×10^{-5}	CCR4 domain-containing protein 1, putative
PF3D7_1228300	7.5×10^{-5}	NIMA related kinase 1
PF3D7_1134600	8.7×10^{-5}	zinc finger protein, putative

Genes highlighted bold have known or suspected roles in gametocytogenesis (indicated by prior listing from previous studies, Supplementary Table S1). * indicates genes that were also identified as having higher expression correlating to MSPDBL2 protein expression in schizonts at significance of $P < 0.001$ (Supplementary Table S4).

References

1. Hodder AN, Czabotar PE, Uboldi AD, Clarke OB, Lin CS, Healer J, Smith BJ, Cowman AF. 2012. Insights into Duffy binding-like domains through the crystal structure and function of the merozoite surface protein MSPDBL2 from *Plasmodium falciparum*. J Biol Chem 287:32922-39.
2. Singh S, Soe S, Weisman S, Barnwell JW, Perignon JL, Druilhe P. 2009. A conserved multi-gene family induces cross-reactive antibodies effective in defense against *Plasmodium falciparum*. PLoS One 4:e5410.
3. Ochola LI, Tetteh KK, Stewart LB, Riitho V, Marsh K, Conway DJ. 2010. Allele frequency-based and polymorphism-versus-divergence indices of balancing selection in a new filtered set of polymorphic genes in *Plasmodium falciparum*. Mol Biol Evol 27:2344-51.
4. Otto TD, Rayner JC, Bohme U, Pain A, Spottiswoode N, Sanders M, Quail M, Ollomo B, Renaud F, Thomas AW, Prugnolle F, Conway DJ, Newbold C, Berriman M. 2014. The genomes of chimpanzee malaria parasites reveal possible pathways of adaptation to human hosts. Nature Communications 5:4754.
5. Otto TD, Bohme U, Sanders M, Reid A, Bruske EI, Duffy CW, Bull PC, Pearson RD, Abdi A, Dimonte S, Stewart LB, Campino S, Kekre M, Hamilton WL, Claessens A, Volkman SK, Ndiaye D, Amambua-Ngwa A, Diakite M, Fairhurst RM, Conway DJ, Franck M, Newbold CI, Berriman M. 2018. Long read assemblies of geographically dispersed *Plasmodium falciparum* isolates reveal highly structured subtelomeres. Wellcome Open Res 3:52.
6. Amambua-Ngwa A, Tetteh KK, Manske M, Gomez-Escobar N, Stewart LB, Deerhake ME, Cheeseman IH, Newbold CI, Holder AA, Knuepfer E, Janha O, Jallow M, Campino S, Macinnis B, Kwiatkowski DP, Conway DJ. 2012. Population genomic scan for candidate signatures of balancing selection to guide antigen characterization in malaria parasites. PLoS Genet 8:e1002992.
7. Mobegi VA, Duffy CW, Amambua-Ngwa A, Loua KM, Laman E, Nwakanma DC, MacInnis B, Aspelng-Jones H, Murray L, Clark TG, Kwiatkowski DP, Conway DJ. 2014. Genome-wide analysis of selection on the malaria parasite *Plasmodium falciparum* in West African populations of differing infection endemicity. Mol Biol Evol 31:1490-9.
8. Tetteh KK, Osier FH, Salanti A, Kamuyu G, Drought L, Faily M, Martin C, Marsh K, Conway DJ. 2013. Analysis of antibodies to newly described *Plasmodium falciparum* merozoite antigens supports MSPDBL2 as a predicted target of naturally acquired immunity. Infect Immun 81:3835-42.
9. Chiu CY, Hodder AN, Lin CS, Hill DL, Li Wai Suen CS, Schofield L, Siba PM, Mueller I, Cowman AF, Hansen DS. 2015. Antibodies to the *Plasmodium falciparum* proteins MSPDBL1 and MSPDBL2 opsonize merozoites, inhibit parasite growth, and predict protection from clinical malaria. J Infect Dis 212:406-415.
10. Tarr SJ, Diaz-Ingelmo O, Stewart LB, Hocking SE, Murray L, Duffy CW, Otto TD, Chappell L, Rayner JC, Awandare GA, Conway DJ. 2018. Schizont transcriptome variation among clinical isolates and laboratory-adapted clones of the malaria parasite *Plasmodium falciparum*. BMC Genomics 19:894.
11. Zhang M, Wang C, Otto TD, Oberstaller J, Liao X, Adapa SR, Udenze K, Bronner IF, Casandra D, Mayho M, Brown J, Li S, Swanson J, Rayner JC, Jiang RHY, Adams JH. 2018. Uncovering the essential genes of the human malaria parasite *Plasmodium falciparum* by saturation mutagenesis. Science 360.

- 455 12. Crosnier C, Iqbal Z, Knuepfer E, Maciucă S, Perrin AJ, Kamuyu G, Goulding D,
456 Bustamante LY, Miles A, Moore SC, Dougan G, Holder AA, Kwiatkowski DP,
457 Rayner JC, Pleass RJ, Wright GJ. 2016. Binding of *Plasmodium falciparum*
458 Merozoite Surface Proteins DBLMSP and DBLMSP2 to Human Immunoglobulin M
459 Is Conserved among Broadly Diverged Sequence Variants. *J Biol Chem* 291:14285-
460 14299.
- 461 13. Van Tyne D, Uboldi AD, Healer J, Cowman AF, Wirth DF. 2013. Modulation of
462 PF10_0355 (MSPDBL2) alters *Plasmodium falciparum* response to antimalarial
463 drugs. *Antimicrob Agents Chemother* 57:2937-41.
- 464 14. Filarsky M, Fraschka SA, Niederwieser I, Brancucci NMB, Carrington E, Carrio E,
465 Moes S, Jenoe P, Bartfai R, Voss TS. 2018. GDV1 induces sexual commitment of
466 malaria parasites by antagonizing HP1-dependent gene silencing. *Science* 359:1259-
467 1263.
- 468 15. Usui M, Prajapati SK, Ayanful-Torgby R, Acquah FK, Cudjoe E, Kakaney C,
469 Amponsah JA, Obboh EK, Reddy DK, Barbeau MC, Simons LM, Czesny B,
470 Raiciulescu S, Olsen C, Abuaku BK, Amoah LE, Williamson KC. 2019. *Plasmodium*
471 *falciparum* sexual differentiation in malaria patients is associated with host factors
472 and GDV1-dependent genes. *Nat Commun* 10:2140.
- 473 16. Prajapati SK, Ayanful-Torgby R, Pava Z, Barbeau MC, Acquah FK, Cudjoe E,
474 Kakaney C, Amponsah JA, Obboh E, Ahmed AE, Abuaku BK, McCarthy JS, Amoah
475 LE, Williamson KC. 2020. The transcriptome of circulating sexually committed
476 *Plasmodium falciparum* ring stage parasites forecasts malaria transmission potential.
477 *Nat Commun* 11:6159.
- 478 17. Kafsack BF, Rovira-Graells N, Clark TG, Bancells C, Crowley VM, Campino SG,
479 Williams AE, Drought LG, Kwiatkowski DP, Baker DA, Cortes A, Llinas M. 2014. A
480 transcriptional switch underlies commitment to sexual development in malaria
481 parasites. *Nature* 507:248-52.
- 482 18. Bancells C, Llorca-Batlle O, Poran A, Notzel C, Rovira-Graells N, Elemento O,
483 Kafsack BFC, Cortes A. 2019. Revisiting the initial steps of sexual development in
484 the malaria parasite *Plasmodium falciparum*. *Nat Microbiol* 4:144-154.
- 485 19. Llorca-Batlle O, Michel-Todo L, Witmer K, Toda H, Fernandez-Becerra C, Baum J,
486 Cortes A. 2020. Conditional expression of PfAP2-G for controlled massive sexual
487 conversion in *Plasmodium falciparum*. *Sci Adv* 6:eaaz5057.
- 488 20. Duffy CW, Amambua-Ngwa A, Ahouidi AD, Diakite M, Awandare GA, Ba H, Tarr
489 SJ, Murray L, Stewart LB, D'Alessandro U, Otto TD, Kwiatkowski DP, Conway DJ.
490 2018. Multi-population genomic analysis of malaria parasites indicates local selection
491 and differentiation at the *gdlv1* locus regulating sexual development. *Sci Rep* 8:15763.
- 492 21. Bowyer PW, Stewart LB, Aspelting-Jones H, Mensah-Brown HE, Ahouidi AD,
493 Amambua-Ngwa A, Awandare GA, Conway DJ. 2015. Variation in *Plasmodium*
494 *falciparum* erythrocyte invasion phenotypes and merozoite ligand gene expression
495 across different populations in areas of malaria endemicity. *Infect Immun* 83:2575-82.
- 496 22. Bohme U, Otto TD, Sanders M, Newbold CI, Berriman M. 2019. Progression of the
497 canonical reference malaria parasite genome from 2002-2019. *Wellcome Open Res*
498 4:58.
- 499 23. Kim D, Langmead B, Salzberg SL. 2015. HISAT: a fast spliced aligner with low
500 memory requirements. *Nat Methods* 12:357-60.
- 501 24. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and
502 dispersion for RNA-seq data with DESeq2. *Genome Biol* 15:550.

25. Josling GA, Russell TJ, Venezia J, Orchard L, van Biljon R, Painter HJ, Llinas M. 2020. Dissecting the role of PfAP2-G in malaria gametocytogenesis. *Nat Commun* 11:1503.
26. Silvestrini F, Lasonder E, Olivieri A, Camarda G, van Schaijk B, Sanchez M, Younis S, Sauerwein R, Alano P. 2010. Protein export marks the early phase of gametocytogenesis of the human malaria parasite *Plasmodium falciparum*. *Mol Cell Proteomics* 9:1437-48.
27. Otto TD, Wilinski D, Assefa S, Keane TM, Sarry LR, Bohme U, Lemieux J, Barrell B, Pain A, Berriman M, Newbold C, Llinas M. 2010. New insights into the blood-stage transcriptome of *Plasmodium falciparum* using RNA-Seq. *Mol Microbiol* 76:12-24.
28. Pelle KG, Oh K, Buchholz K, Narasimhan V, Joice R, Milner DA, Brancucci NM, Ma S, Voss TS, Ketman K, Seydel KB, Taylor TE, Barteneva NS, Huttenhower C, Marti M. 2015. Transcriptional profiling defines dynamics of parasite tissue sequestration during malaria infection. *Genome Med* 7:19.
29. Aurecochea C, Brestelli J, Brunk BP, Dommer J, Fischer S, Gajria B, Gao X, Gingle A, Grant G, Harb OS, Heiges M, Innamorato F, Iodice J, Kissinger JC, Kraemer E, Li W, Miller JA, Nayak V, Pennington C, Pinney DF, Roos DS, Ross C, Stoeckert CJ, Jr., Treatman C, Wang H. 2009. PlasmoDB: a functional genomic database for malaria parasites. *Nucleic Acids Res* 37:D539-43.
30. WHO. 2021. World Malaria Report 2021. Geneva.
31. Eksi S, Haile Y, Furuya T, Ma L, Su X, Williamson KC. 2005. Identification of a subtelomeric gene family expressed during the asexual-sexual stage transition in *Plasmodium falciparum*. *Mol Biochem Parasitol* 143:90-9.
32. Brancucci NMB, Bertschi NL, Zhu L, Niederwieser I, Chin WH, Wampfler R, Freymond C, Rottmann M, Felger I, Bozdech Z, Voss TS. 2014. Heterochromatin protein 1 secures survival and transmission of malaria parasites. *Cell Host Microbe* 16:165-176.
33. Broadbent KM, Broadbent JC, Ribacke U, Wirth D, Rinn JL, Sabeti PC. 2015. Strand-specific RNA sequencing in *Plasmodium falciparum* malaria identifies developmentally regulated long non-coding RNA and circular RNA. *BMC Genomics* 16:454.
34. Lopez-Barragan MJ, Lemieux J, Quinones M, Williamson KC, Molina-Cruz A, Cui K, Barillas-Mury C, Zhao K, Su XZ. 2011. Directional gene expression and antisense transcripts in sexual and asexual stages of *Plasmodium falciparum*. *BMC Genomics* 12:587.
35. Howick VM, Russell AJC, Andrews T, Heaton H, Reid AJ, Natarajan K, Butungi H, Metcalf T, Verzier LH, Rayner JC, Berriman M, Herren JK, Billker O, Hemberg M, Talman AM, Lawniczak MKN. 2019. The Malaria Cell Atlas: Single parasite transcriptomes across the complete *Plasmodium* life cycle. *Science* 365.
36. Lemieux JE, Gomez-Escobar N, Feller A, Carret C, Amambua-Ngwa A, Pinches R, Day F, Kyes SA, Conway DJ, Holmes CC, Newbold CI. 2009. Statistical estimation of cell-cycle progression and lineage commitment in *Plasmodium falciparum* reveals a homogeneous pattern of transcription in *ex vivo* culture. *Proc Natl Acad Sci U S A* 106:7559-64.
37. Daily JP, Scanfeld D, Pochet N, Le Roch K, Plouffe D, Kamal M, Sarr O, Mboup S, Ndir O, Wypij D, Levasseur K, Thomas E, Tamayo P, Dong C, Zhou Y, Lander ES, Ndiaye D, Wirth D, Winzeler EA, Mesirov JP, Regev A. 2007. Distinct physiological states of *Plasmodium falciparum* in malaria-infected patients. *Nature* 450:1091-5.

- 552 38. Lee HJ, Georgiadou A, Walther M, Nwakanma D, Stewart LB, Levin M, Otto TD,
553 Conway DJ, Coin LJ, Cunningham AJ. 2018. Integrated pathogen load and dual
554 transcriptome analysis of systemic host-pathogen interactions in severe malaria. *Sci*
555 *Transl Med* 10.
- 556 39. Yamagishi J, Natori A, Tolba ME, Mongan AE, Sugimoto C, Katayama T,
557 Kawashima S, Makalowski W, Maeda R, Eshita Y, Tuda J, Suzuki Y. 2014.
558 Interactive transcriptome analysis of malaria patients and infecting *Plasmodium*
559 *falciparum*. *Genome Res* 24:1433-44.
- 560 40. Mok S, Ashley EA, Ferreira PE, Zhu L, Lin Z, Yeo T, Chotivanich K, Imwong M,
561 Pukrittayakamee S, Dhorda M, Nguon C, Lim P, Amaratunga C, Suon S, Hien TT,
562 Htut Y, Faiz MA, Onyamboko MA, Mayxay M, Newton PN, Tripura R, Woodrow
563 CJ, Miotto O, Kwiatkowski DP, Nosten F, Day NP, Preiser PR, White NJ, Dondorp
564 AM, Fairhurst RM, Bozdech Z. 2015. Population transcriptomics of human malaria
565 parasites reveals the mechanism of artemisinin resistance. *Science* 347:431-5.
566