Competition contributes to both warm and cool range edges

Shengman Lyu^{1*}, Jake M. Alexander¹

¹ Institute of Integrative Biology, ETH Zürich, 8092 Zürich, Switzerland

* Correspondence: shengman.lyu@usys.ethz.ch

ORCID

Shengman Lyu <https://orcid.org/0000-0003-0221-5315>

Jake M. Alexander <https://orcid.org/0000-0003-2226-7913>

Abstract

Competition plays an important role in shaping species' spatial distributions. However, it remains unclear where and how competition regulates species' range limits. In an experiment with plants originating from low and high elevations and conducted across an elevation gradient in the Swiss Alps, we found that the ability of both lowland and highland species to persist under competition was reduced beyond versus within their elevation ranges, suggesting that competition helps set both lower and upper elevation range limits of these species. Furthermore, the reduced ability of pairs of lowland or highland species to coexist beyond their range edges was mainly driven by diminishing niche differences; changes in both niche differences and relative fitness differences drove weakening competitive dominance of lowland over highland species with increasing elevation. These results highlight the need to account for competitive interactions and investigate underlying coexistence mechanisms to understand current and future species distributions.

Keywords

coexistence, competition, elevation gradient, relative fitness difference, niche difference, integral projection model, perennial plant, population growth

Introduction

Climate and biotic interactions, in particular competition, are key factors constraining species' geographic ranges^{1,2,3}. According to niche theory, a focal species is able to overcome abiotic and biotic constraints on population growth to maintain persistent populations within, but not beyond, its range edge⁴. This hypothesis has been widely supported by empirical studies. For instance, transplant experiments have often found declines in fitness when species are transplanted beyond their range edges, with greater declines in fitness when individuals are transplanted into sites where other species are present (see ref. 5 for a review). Another longstanding hypothesis regarding species' range limits is that competition is especially important for setting warm range edges, such as those at low elevation or latitude^{1,2,6,7}. However, while some transplant studies have found patterns consistent with this hypothesis⁵, other studies indicate that competition is also important for setting cool range edges such as those at high elevation or latitude^{8,9,10}. Thus, although the general importance of competition in shaping distributions is well established, how this varies across environmental gradients, as well as the underlying processes involved, remain unclear.

We can use tools and insights from community ecology theory to better understand when competition contributes to setting range limits^{11,12}. From the perspective of coexistence theory, within its range a population of a focal species is predicted to be able to grow from low density when other species are at their equilibrium abundance (i.e., invasion population growth rate, $\ln(\lambda_{\text{invasion}}) > 0$) and thus persist with other community members. When beyond its range limit, the same species is predicted to be unable to maintain persistent populations (i.e., $\ln(\lambda_{\text{invasion}}) < 0$), resulting in competitive exclusion¹¹, even if the species would be able to persist in the absence of other competitors (i.e., intrinsic population growth rate, $\ln(\lambda_{\text{intrinsic}}) > 0$). In other words, the ability of a species to coexist with its neighbours is expected to be reduced towards its range edge.

A key advantage of conceiving range limits as a coexistence problem is that we can begin to unpack the processes through which competitors influence species' range limits¹². Modern coexistence theory predicts that coexistence is determined by two sets of differences between competing species¹³. Differences in competitive ability (i.e., relative fitness differences) reflect both intrinsic differences in species' demographic performance and/or their sensitivity to competition, and drive competitive exclusion¹⁴; species' differences that reduce the intensity of interspecific relative to intraspecific competition (i.e., niche differences) promote stable coexistence¹⁵. These can result for example from resource partitioning or specialist natural enemies¹⁵. Accordingly, competition could shape range

limits through two pathways. Firstly, a focal species may be competitively eliminated outside of its range because it encounters other species with which it has greater niche overlap compared to species within its range. For example, the replacement of closely related, ecologically similar species across elevation gradients has been explained by interspecific competition for shared habitats¹⁶. Similarly, changing environmental conditions towards range edges can also reduce niche differences between currently co-existing species (i.e., decreased niche differences toward the range edge, see ref. 17 for a similar example), increasing the likelihood of competitive exclusion.

The second pathway through which competition could contribute to range limits is via changes to relative fitness differences. A focal species may coexist stably with other species within its range but be competitively excluded beyond its range edge because of reduced competitive ability. On the one hand, reduced competitive ability could arise because the focal species has a reduced ability to tolerate environmental conditions outside of its range in the absence of other competitors (i.e., a diminished $\lambda_{\text{intrinsic}}$). On the other hand, the focal species may become more sensitive to competition, or its competitors exert stronger effects beyond versus within its range^{15, 18}. Taking an elevation gradient as an example, a high elevation plant could be restricted to higher elevations by competition from lower elevation species that are better competitors for shared resources, like light¹⁹, under a warmer climate^{20, 21}; but even competition with co-occurring high elevation species might generate a range limit, if at least some of those species become relatively stronger competitors under warmer conditions at lower elevation (see refs. 22 and 23 for similar examples).

Here, we evaluate the hypothesis that species experience a reduced ability to coexist with neighbours beyond versus within their range edges, and examine the contributions of niche and relative fitness differences to changing coexistence across an elevation gradient. To address this, we conducted a field experiment interacting seven low and seven high elevation species with limited range overlap (hereafter lowland and highland species) in three sites across an elevation gradient. The low (890 m a.s.l.) site was located within the current elevation range of lowland species and beyond the lower range edge of highland species, while the high site (1900 m) was located within the elevation range of highland species and beyond the upper range edge of lowland species, and the middle site (1400 m) was located near the shared range edges of both groups. In each site, we parameterized integral projection models (IPMs) to estimate population growth rates (λ) both in the absence (i.e., intrinsic population growth rates) and presence (i.e., invasion population growth rates) of neighbour species. Our approach enabled us to quantify coexistence determinants (i.e., niche differences and relative fitness differences) and predict the outcomes of competition

between species pairs. We ask two main questions: (1) How does the ability of lowland and highland species to persist with competitors change across an elevation gradient? (2) To what extent are any changes in the strength of coexistence driven by changes in niche differences and changes in relative fitness differences? Our study shows that both lowland and highland species display a reduced ability to persist with competitors beyond their range edges, suggesting that competition helps set range limits both at low (i.e., the lower range edges of highland species) and high (i.e., the upper range edges of lowland species) elevations. Furthermore, we show that competition can influence species' elevation ranges through changes to both niche differences and relative fitness differences, but that their relative importance varies depending on the origin of the competitors. These results therefore highlight the importance of considering competition and its underlying mechanisms to understand species' distributions across environmental gradients, and ultimately to forecast range dynamics under changing environments.

Results

Declines in intrinsic and invasion population growth rates towards range edges

Lowland and highland species displayed distinct responses to elevation in the absence of competition (Fig. 1a). The intrinsic population growth rates of both lowland and highland species were projected to decline beyond elevational range edges (test based on the mean of 500 bootstraps; elevation $\times$ species origin: $F_{1, 41} = 7.062$, $P = 0.008$; Supplementary Table 7; median and 95% CI of elevation $\times$ species origin interaction across bootstrap replicates: 0.0011, 0.0008–0.0013). Interestingly, most species were projected to be able to persist in the absence of neighbours across the whole elevation gradient, regardless of their elevation origin (i.e., $\ln(\lambda_{\text{intrinsic}}) > 0$; Supplementary Fig. 6).

In the presence of competition, the projected population growth rates displayed greater declines beyond the range edges of both lowland and highland species (Fig. 1b; test based on the mean of 500 bootstraps; elevation $\times$ species origin: $F_{1, 300} = 21.215$, $P < 0.0001$; Supplementary Table 7; median and 95% CI of elevation $\times$ species origin interaction across bootstrap replicates: 0.0008, 0.0005 – 0.0011). When competing against other species, highland species were on average predicted to be able to persist at the high site, that is within their range, but not beyond their lower range edges at the low elevation site. In contrast, the projected ability of lowland species to persist under competition decreased with increasing elevation, though most of them were predicted to be able to persist in all three sites (Fig. 1b; Supplementary Fig. 6).

Changing outcomes of competition across the elevation gradient

The predicted ability of species to coexist changed across the elevation gradient, with effects that depended on competitor identity (Fig. 2). For sympatric pairs (i.e., lowland-lowland and highland-highland pairs), coexistence was more prevalent within than outside of the current ranges. On average across all bootstrap replicates, 63% of all eight and 67% of all five highland-highland pairs were predicted to coexist at the high and middle site versus only 40% at the low site (Fig. 2c). Predictions of coexistence of lowland-lowland pairs were more common and displayed only few changes across the sites, with 78% of all 10 lowland-lowland pairs predicted to coexist at the low and high sites and 50% at the middle site (Fig. 2a). Allopatric pairs (i.e., lowland-highland pairs) were more likely to coexist at the middle elevation near shared range edges, with 58% of all 13 pairs predicted to coexist at the middle site and 46% and 44% at the low and high sites, respectively (Fig. 2b).

These findings based on tallies of coexistence outcomes were supported by an analysis of a metric measuring the strength of coexistence (i.e., the coexistence metric, see Methods). These showed that both lowland-lowland and highland-highland pairs displayed weakened coexistence (i.e., decreasing coexistence metric) beyond their elevation ranges, while lowland-highland pairs displayed the greatest ability to coexistence at the middle elevation (Fig. 2d-f; test based on the mean of 500 bootstraps: elevation $\times$ competitor identity: $F_{2, 90} = 11.011$, $P = 0.004$; Supplementary Table 8). The interaction was mainly driven by the opposing responses of lowland and highland pairs, although this contrast was not significantly different from zero after error propagation (individual tests for each bootstrap replicate; median and 95% CI: -0.0006, -0.0017 – 0.00004).

Niche differences and relative fitness differences jointly mediate changing coexistence

Changes in the magnitude of both niche and relative fitness differences mediated variation in coexistence across the elevation gradient (Fig. 3). For sympatric pairs, our results showed that both lowland and highland species displayed greater niche differences (i.e., smaller niche overlap) in sites that lay within their range (Fig. 3a&c; test based on the mean of 500 bootstraps; elevation $\times$ competitor identity: $F_{2, 90} = 11.603$, $P = 0.003$; Supplementary Table 8), although this result was not significant after error propagation (individual tests for each bootstrap replicate; median and 95% CI of the contrast between lowland and highland pairs: 0.0005, -0.00005 – 0.0012). In addition, reduced relative fitness differences towards high elevation also contributed to strengthened coexistence of highland species within their range (Fig. 3f; test based on the mean of 500 bootstraps; $F_{1, 17} = 5.179$, $P = 0.023$), although this

result was not significant after error propagation (individual tests for each bootstrap replicate; median and 95% CI: -0.0002, -0.0006 – 0.0001). No significant trend on relative fitness differences was seen for lowland-lowland pairs ($F_{1,29} = 0.192$, $P = 0.661$) or when pooling all pairs together and taking absolute values (i.e., absolute differences in competitive ability irrespective of competitor origin; test based on the mean of 500 bootstraps, $F_{1,90} = 1.406$, $P = 0.236$; Supplementary Table 8).

For parapatric pairs, lowland species were predicted to be competitively dominant over highland species in general (i.e., $\ln(\text{fitness difference}) < 0$), but the competitive dominance was predicted to decrease towards high elevations, where some highland species were predicted to be dominant (Fig. 3e; test based on the mean of 500 bootstraps, $F_{1,44} = 3.379$, $P = 0.046$). However, this result was not significant after error propagation (individual tests for each bootstrap replicate; median and 95% CI of the slope: 0.0001, -0.0002 – 0.0005). Niche differences between lowland and highland species tended to be greatest at middle elevation (Fig. 3b), contributing to the greatest ability to coexist of at the middle site in conjunction with reduced relative fitness differences (Fig. 2e).

Discussion

Although the role of competition in shaping species' spatial distribution is well acknowledged^{2, 3}, how this role varies across environmental gradients, and by which processes (e.g., through changes in niche and relative fitness differences) competition regulates species' range limits, remain unclear^{6, 12}. Here we found that the ability of species to persist in the presence of competition was weakened beyond versus within their elevational ranges, supporting the role of competition in shaping species' elevation distribution limits. However, in contrast to the conventional view that competition is more important in abiotically benign environments, such as at low elevation or latitude (e.g., as predicted by the stress-gradient hypothesis^{6, 7, 24}), our results indicate that competition can also be important towards high elevation range limits. There are at least two aspects that distinguish our study from previous work that can partly help explain this discrepancy. Firstly, we focused on the population-level outcomes of competition rather than the individual-level strength of competition, as most previous empirical studies have done^{5, 25}. Both theoretical and empirical work demonstrate that competition can be equally important for population dynamics in harsh and benign environments¹⁰ and its effects cannot be completely understood by focusing solely on the strength of competition^{9, 26}. For example, harsh environmental conditions could reduce population growth rates directly, making species less tolerant of interspecific competition and competitive exclusion correspondingly more likely to occur, despite any reduced intensity or prevalence of interspecific competition itself^{9, 27, 28}.

Secondly, we grew the same set of species in each experimental site, which allowed us to isolate changes in the intensity of competition from turnover in community composition across the environmental gradient⁶. Though this to some extent divorces our study from the effects of the complex interactions that play-out in natural multispecies communities (see ref. 29 and below), it shows that both changing intensity of interactions and changing identity of species contribute to the changing competition experienced by a focal species towards its range edge, and we obtained further insight into these processes through analyses of changes in niche and fitness differences across elevation.

Although effects of niche and relative fitness differences cannot be understood in isolation¹³, for our species, changes in niche differences appeared to be especially important for mediating changing coexistence between species across the elevation gradient. Sympatric species (i.e., lowland-lowland and highland-highland pairs) possessed the greater niche differences within versus outside of their elevational range, while parapatric species (i.e., lowland-highland pairs) displayed the greatest niche differences at the middle elevation site near to their shared range margins. So far, there have been few a priori predictions made about how niche differences should change across environmental gradients. Nonetheless, our finding of greater niche differences between sympatric species within than beyond their range is consistent with a recent study showing reduced niche differences under drought conditions due to greater positive effects of drought on the magnitude of interspecific relative to intraspecific competition³⁰. We can only speculate on the biological processes driving the distinct responses of niche differences without further information (e.g., from functional traits³¹). However, one possible explanation could be temporal overlap in phenology. Specifically, the flowering phenology of lowland and highland species overlapped most at the high and low elevation sites, respectively, while the overlap between lowland and highland species was smallest at the middle elevation site (Supplementary Fig. 7). Thus, the phenological overlap between species was associated with the magnitude of niche differences, with greater phenology overlap leading to smaller niche differences and vice versa^{32, 33}.

Lowland species were in general competitively superior over highland species, but this superiority decreased significantly with increasing elevation, and the competitive dominance of some species pairs was switched in favour of highland plants at the high elevation sites (i.e., $\ln(\text{fitness difference}) > 0$ in Fig. 3e). Nonetheless, in most cases the competitive advantage of highland species at the high site was not great enough to exclude the lowland species; possible reasons for this are discussed below. The competitive dominance of lowland species is consistent with general expectations⁶ (but see ref. 20), as well as empirical evidence showing that lowland species can exert stronger competitive effects on

alpine focal plants than other alpine neighbours²¹. Greater competitive superiority of species within their home range could result either from their higher intrinsic population growth, in particular for highland species, or from the fact that species are more resistant to competition within versus beyond their range^{18, 32}, though we cannot isolate the relative contribution of these two processes with the data at hand. Similarly, the increased relative fitness differences of highland species towards low elevation could result from amplified differences in intrinsic population growth or competitive responses³⁰. Taken together, our results demonstrate that the range limits of our focal species were influenced by competitors from both within and outside of their range (i.e., sympatric and parapatric species) and mediated through both niche differences and relative fitness differences. These results speak to the need to broadly investigate how environmental conditions impact niche and relative fitness differences and how they jointly structure communities and species' distributions across environmental gradients¹².

Although we found significant effects of elevation on the strength of coexistence (i.e., the coexistence metric), we observed relatively few cases in which coexistence outcomes were qualitatively altered across the elevation gradient. For example, most lowland species could still maintain relatively large and positive invasion growth rates, and thus were predicted to persist, at the high site. The high site is above the current range limit of all lowland species in this region, and so it is surprising that the climatic and biotic environment appears to be suitable for them. This might be because while average climate conditions at high elevation support the growth of our lowland plants, climate extremes do not³⁴; if extreme conditions did not occur during the experiment, then climate limitation would not be observed. Another possible explanation for lowland species' persistence at high elevation is disequilibrium between their distributions and current climate. It has been widely reported that the rate of species' upwards migration across elevation gradients lags behind the pace of climatic warming^{35, 36, 37}. For example, the observed average upward shifts of 183 plant species over the past 50 years in the European Alps was 30 m³⁸, while isoclines in the same region have shifted upwards by 310-355 m³⁵. Disequilibrium in the distributions of our lowland species might be due to their limited dispersal capacities and the often low rate of germination and establishment of perennial species³⁶. Added to this, the growing season temperatures over the course of our experiment generally exceeded the long-term average (<https://www.meteoswiss.admin.ch/>). Thus, the high site could already be within the potential upper limits of some of the lowland species included in our study. But even if this were the case, the relationship shown in Fig. 1 implies that lowland species ranges will be limited by competitors before they are limited by climate.

Manipulative experiments such as ours will tend to underestimate the prevalence of stable coexistence when they focus on pairwise competition and use homogeneous environmental conditions over a relatively short temporal scale (e.g., ref. 14). Firstly, such experiments might miss potentially important stabilizing effects that only emerge from interaction networks in which any given pair of species is not necessarily able to coexist (e.g., higher-order interactions^{29, 39}). Secondly, standardized, short term experiments might not be able to capture temporal and spatial variation that could be important for species coexistence (e.g., temporal and spatial storage effects⁴⁰). In our experiment, we used the same soil in the three sites to isolate effects of changing climate from changing soil conditions across the elevation gradient, which might have reduced opportunities for coexistence driven by interactions belowground (e.g., plant-soil feedbacks⁴¹). This failure to capture all possible coexistence mechanisms might partly explain the lack of stable coexistence between currently co-occurring species observed in our experiment. For example, we found that only 63% of highland species pairs were predicted to be able to coexist at the high site; small-scale environmental heterogeneity has been suggested to be especially important for maintaining the high diversity of alpine plants⁴². Thirdly, due to the short duration the experiment (four years), our results might be contingent on specific weather conditions over the course of the experiment, which may also explain the lack of coexistence of highland species within their range. Lastly, violations of the assumption that monocultures are at equilibrium abundance could affect our predictions of coexistence. In our experiment, 32% (11 of 34) of monocultures were predicted to meet this assumption, and those that did not (10 and 13 monocultures were predicted to be above and below equilibrium, respectively) were evenly distributed across the sites and so are unlikely to affect the general patterns we observed (Supplementary Fig. 5).

Our results challenge the conventional view predicting a diminished role for competition in structuring species' distributions and ecological communities under "harsh" environments at high elevation or latitude, whilst highlighting the importance of considering species' interactions to understand and project future species distributions^{27, 28}. More generally, our finding that both lowland and highland species experienced weakened coexistence towards their range edges (i.e., the upper ranges of lowland species and the lower ranges of highland species, respectively) suggests that competition might be important for setting both species' lower and upper elevation limits¹⁰. An important role for competition at high elevation could help explain the limited ability of some species to shift their distribution limits to higher elevation to track climate warming^{35, 36}. However, our results suggest that continuous warming may facilitate the upwards shifts of lowland species and the local extinction of alpine species⁴³ by removing the competitive advantage that alpine species

possess under cooler climate. In addition, our results suggest that climatic warming may destabilize alpine communities by reducing niche differences and amplifying fitness differences between highland species.

Although modern coexistence theory provides a useful framework for understanding and predicting species coexistence across environmental gradients, niche differences and relative fitness differences are only phenomenological syntheses of biological processes underling species interactions. Future work should seek to uncover the mechanisms through which competition influences species' distributions along environmental gradients by manipulating specific abiotic and biotic factors and gathering data on species' physiology, morphology or life-history strategies (i.e., functional traits³¹). Future studies could also look at whether competition sets range limits across entire species ranges, including upper and lower edges of the same species, and how complex interaction networks in multispecies communities contribute to setting range limits. Such studies would help to provide more mechanistic insights into the role of species interactions in regulating species' distributions and bolster our ability to more accurately forecast the impacts of environmental change on species' distributions and community dynamics.

Materials and Methods

Study area and species

We selected three sites across an elevation gradient in the western Swiss Alps (Bex, Canton de Vaud), situated at 890, 1400 and 1900 m above sea level (hereafter, the low, middle, and high sites; Supplementary Fig. 1). The three sites span a temperature gradient ranging from 2.5 to 9.6 °C (mean annual temperature from 1981 to 2015 ⁴⁴; Supplementary Table 1). With increasing elevation, soil moisture increased, and growing season length was shortened by a longer snow-covered period, as measured from July 2019 to June 2020 (Supplementary Fig. 2). All sites were established on south facing and shallow slopes in pasture and fenced to exclude livestock.

We included 14 herbaceous focal species that frequently occur in this region, half of which originated from low elevation (hereafter lowland species) and half from high elevation (highland species, Supplementary Table 2). Lowland species had upper range limits (defined as the 90th percentile of their elevation distribution) below 1500 m (with the exception of *Plantago lanceolata*, with a 90th percentile of 1657 m), while highland species had lower range limits (defined as the 10th percentile of their elevation distribution) above 1500 m, based on a dataset of 550 vegetation plots from the study area ⁴⁵. These species consisted of 12 perennial and two biennial species, which are the dominant life histories in this region. Species were selected to include a range of functional types (7 forbs, 4 grasses, 3 legumes) and functional traits (based on plant height, specific leaf area and seed mass). Seeds were obtained from regional suppliers given the large quantities that were needed to establish the experiment (Supplementary Table 2).

Field competition experiment

We designed a field experiment to study the effects of elevation on population growth rates and competitive outcomes by growing focal plants either without competition or competing with a background monoculture of the same or another species (Supplementary Fig. 1). In spring 2017, we established 18 plots (1.6 m x 1 m, 0.2 m deep) at each of the three field sites, lined with wire mesh to exclude rodents (except at the high site) and with weed-suppressing fabric on the sides to prevent roots growing in from outside. To control for soil effects, the beds were then filled with a silt loam soil that originated from a nutrient poor meadow at 1000 m a.s.l. within the study area. Four plots were maintained as bare soil plots (non-competition plots). The other 14 plots received 9 g m⁻² of viable seeds of each species, which allowed the establishment of a monoculture of relatively high density (competition

plots). We then periodically weeded the plots to maintain monocultures over the course of the experiment. All species except for two (*Arnica montana* and *Daucus carota*) successfully established monocultures, of which 11 species (including six lowland species and five highland species) were fully established by autumn 2017. We then resowed the other plots, which subsequently established either in spring 2018 (*Poa trivialis* and *Poa alpina* in the low site and *Bromus erectus* in the middle site) or autumn 2018 (*Aster alpinus*, *P. trivialis* and *P. alpina* in the middle site and *Sesleria caerulea* in the low and high sites). Species that failed to establish were included only as focal species for the calculation of invasion population growth rates (i.e., the density was low for *A. montana* and *D. carota* in all sites, *Trifolium badium* in the low site and *S. caerulea* in the middle site, probably due to high mortality rates caused by drought).

We first raised seedlings of each species in a greenhouse for six weeks on standard compost and then transplanted them into the field (Supplementary Fig. 1). To test for responses to elevation in the absence of competition, focal plants were transplanted into non-competition plots at 25 cm apart in autumn 2017 ($n = 9$ per species and site). To test for effects of competition, we transplanted focal individuals into established plots with 14 cm spacing ($n = 9$ per focal species, competitor and site). Focal plants that died within two weeks of transplanting (ca. 5%) were replaced, assuming mortality was caused by transplant shock. Note that we transplanted focal plants into plots only when the background monocultures were fully established. In 2018 and 2019, we replaced dead focal individuals in spring and autumn (ca. 10% each time). The full design included 56 unique interspecific pairs in each site accounting for 61% of all $14 \times 13 = 91$ possible pairwise combinations. These pairs were selected to evenly sample differences in functional trait space based on a pilot analysis using plant height, specific leaf area and seed mass obtained from the LEDA dataset⁴⁶. Each focal species competed against four lowland and four highland species, yielding 14 lowland-lowland and highland-highland pairs and 28 lowland-highland pairs. Across all three sites, this design resulted in $N = 3780$ individuals in total ($[56 \text{ interspecific pairs} \times 2 + 14 \text{ intraspecific pairs} + 14 \text{ non-competition}] \times 9 \text{ individuals} \times 3 \text{ sites}$).

Demographic data

We followed each focal individual between 2017 and 2020 to monitor individual-based demographic performance (i.e., vital rates; Supplementary Fig. 4). Survival was monitored twice a year at the beginning and the end of the growing season. Towards the end of the growing season in each year (August-September), we measured all individuals to record plant size, whether they flowered, and to estimate seed production on flowering individuals. To estimate focal plant size, we measured size-related morphological traits on all focal

individuals at each census (i.e., the number and/or length of flowering stalks, leaves or ramets, depending on species) and estimated dry aboveground biomass using regression models fitted using collected plant samples (mean $R^2 = 0.871$; Supplementary Table 3; Supplementary Methods). To estimate seed production, we counted the number and measured the size of fruits on reproductive individuals; we then estimated the number of seeds produced by each individual using regression models fitted using intact fruits of each species collected at the early fruiting stage on background plants (mean $R^2 = 0.806$; Supplementary Table 4; Supplementary Methods). We conducted a separate experiment to estimate germination and recruitment of each species in each site (Supplementary Methods).

Population modelling

To estimate population growth rates (λ), we built integral projection models to incorporate multiple vital rates across the life cycle⁴⁷ (see Supplementary Table 5 for model structure and parameters). Separate IPMs were built to estimate intrinsic growth rates using plants growing in the absence of competition (in non-competition plots) and invasion growth rates using plants growing within the background monocultures (in competition plots), under the assumption that monocultures were at equilibrium (see Supplementary Fig. 5 for a test of this assumption) and that focal individuals did not interact with each other but only with the background species. We used plant size (i.e., estimated dry aboveground biomass, log scale) as a continuous state variable and fitted linear models to estimate vital rate parameters by combining multiple-year demographic data over three censuses (i.e., 2017-2018, 2018-2019 and 2019-2020; see Supplementary Methods for consideration of more complex models). We modeled the probability of survival, flowering, and seedling establishment using generalized linear models with a binomial distribution, modeled growth and seed production using general linear models and described the offspring size distribution using Gaussian probability density functions. We modeled seed germination, seedling establishment and the seedling size distribution as size-independent functions, assuming they are unaffected by maternal size (Supplementary Fig. 4; Supplementary Table 5). For each vital rate of each species, we selected the best-fitted vital rate model by comparing all nested models of the full models using the Akaike information criterion corrected for small samples (AICc), which allowed us to avoid overfitting models and to borrow strength across competitor species and sites in cases where full models were outperformed by reduced models (Supplementary Methods; Supplementary Table 6; Supplementary Data 1).

We calculated population growth rates (λ) as the dominant eigenvalue of the IPMs, which represents the discrete per-capita growth rate (i.e., $N_{t+1} = \lambda N_t$)⁴⁷. To evaluate the uncertainty around λ , we performed parametric bootstraps for size-dependent vital rates (i.e., survival, growth, flowering, and fecundity). Specifically, we resampled the parameters of each vital rate model using multivariate normal distributions based on their means and covariance matrices⁴⁸. We then fitted all IPMs and estimated λ for each of the 500 bootstrap replicates (Supplementary Data 2).

Estimation of niche differences, relative fitness differences and coexistence outcomes

We quantified niche and relative fitness differences and predicted coexistence outcomes following the method of Carroll et al.⁴⁹. This method is based on species' sensitivity to competition defined as the proportional reduction of the growth rate of a focal species i when invading a population of a competitor species j that is at its single-species equilibrium, and is mathematically equivalent to one minus the response ratio:

$$S_{ij} = 1 - \frac{\ln(\lambda_{ij})}{\ln(\lambda_i)}$$

where λ_{ij} denotes the invasion growth rate of focal species i and λ_i is its intrinsic growth rate. The natural logarithm of discrete population growth rates λ estimated from IPMs are equivalent to per capita growth rate in continuous population growth models⁵⁰, and this transformation makes sensitivities compatible with the coexistence analysis described below. Sensitivity is > 0 for antagonistic interactions, with higher values equating to stronger competition, while facilitative interactions lead to negative sensitivities.

For a pair of species, modern coexistence theory predicts that niche differences (ND) promote coexistence by reducing the intensity of interspecific competition experienced by both species. Therefore, a pair of species with a large niche difference should display small mean sensitivities to competition from each other. Consequently, niche differences can be calculated as one minus the geometric mean of the two sensitivities (i.e., niche overlap). In contrast, relative fitness differences (RFD) quantify the degree of asymmetry in species' competitive abilities. Therefore, a pair of species with a large fitness difference should display large differences in their sensitivities to competition from each other, as quantified as the geometric standard deviation of sensitivities⁴⁹:

$$ND = 1 - \sqrt{S_{ij}S_{ji}}$$

$$RFD = \sqrt{S_{ji}/S_{ij}}$$

There are three possible outcomes of competition between a given pair of species: stable coexistence, a priority effect, and competitive exclusion. These can be quantified based on either invasion criteria or the relative magnitude of niche differences versus relative fitness differences^{15, 51}. Stable coexistence is only possible when both species are able to invade each other's equilibrium populations; this condition is met when $ND > 0$ and $RFD < \frac{1}{1-ND}$ ⁴⁹, which is equivalent to $\frac{1}{RFD(1-ND)} > 1$, with greater values indicating more stable coexistence and providing a metric for the strength of coexistence (i.e., coexistence metric²⁶). When neither species can invade when rare, then priority effects occur, meaning that whichever species is initially established within a community has an advantage and excludes the other. This could happen when a species pair has a small niche difference and a small relative fitness difference, that is $ND < 0$ and $RFD < \frac{1}{1-ND}$ ⁵¹. Otherwise, if only one species can invade, then it is predicted to competitively exclude the inferior competitor; this occurs when relative fitness differences are big enough to override the stabilizing effect of niche differences, that is when $RFD > \frac{1}{1-ND}$. We quantified competitive outcomes and coexistence metrics for each of the 500 bootstrap replicates of the dataset (Supplementary Data 3).

Note that we excluded facilitative interactions that were present in 13% of all pairs because the equations for niche differences and relative fitness differences are not compatible with negative values of sensitivity. We quantified the coexistence determinants of species pairs in cases where either one or both of the species were predicted to be unable to persist in the absence of neighbours (i.e., $\ln(\lambda_{intrinsic}) < 0$; 8% of all pairs). This was because the quantification of niche and fitness differences were based on the competitive impacts on population growth (i.e., sensitivity to competition) and independent of the actual estimates of population growth rates. A further analyses after excluding these pairs produced very similar results that did not change our conclusions (data not shown).

Statistical analysis

We tested the effect of elevation on species' intrinsic and invasion growth rates, and whether these effects differed between lowland and highland species, by fitting mixed effects models with elevation (continuous), species origin (i.e., lowland or highland) and their interaction as fixed effects and focal species identity as a random effect. To test whether coexistence was strengthened at high versus low elevation, we fitted a mixed-effects model of the

coexistence metric as explained by elevation and competitor identity (i.e., the three types of pairs: lowland-lowland, lowland-highland and highland-highland pairs), with species-pair as a random effect. We fitted similar models for niche and relative fitness differences (absolute values) to gain insights into whether changes in coexistence were driven predominantly by variation in niche or relative fitness differences. Additionally, we tested the effect of elevation on relative fitness differences for each type of pairs separately and used the actual values of lowland-highland pairs since we were interested not only in the magnitude of relative fitness differences but also the direction of effects (i.e., whether lowland or highland species were dominant competitors). The response variables in these models were log-transformed to meet model assumptions. Note that we therefore had to use niche overlap (1-ND) for the log-transformation instead of niche differences, because niche differences included negative values. Firstly, we performed these analyses based on the mean of the 500 bootstrap replicates for a given focal species/background competitor combination and used likelihood ratio tests to determine the significance of fixed effects. To account for the uncertainty around estimated λ , we also performed individual tests for each bootstrap replicate of the dataset and determined the significance of fixed terms based on whether the 95% confidence interval of estimated coefficients, or contrasts of interest, included zero. All population modelling and statistical analysis were conducted in R version 4.0.3⁵².

Data availability

A complete comparison of vital rate models can be found in Supplementary Data 1. Estimated population growth rates of all integral projection models can be found in Supplementary Data 2. Estimates of competitive outcomes and coexistence coefficients can be found in Supplementary Data 3. Supplementary Data 2&3 were used to produce Figs. 1-3. These data are available on Figshare: <https://figshare.com/s/cb9e034633f47739466b>

Code availability

All R scripts used for modelling and statistical analyses are available on GitHub: <https://github.com/ShengmanLyu/competition-contributes-to-range-limits>

Acknowledgements

We thank Loïc Liberati, Tim Murray, Jessica Joaquim, Cuicui Zhou, Yan Hess, Kai-Hsiu Chen, Megan Stamp and other members of the Plant Ecology group at ETH Zürich for help with field work. We thank the Commune de Bex and Jean-Louis Putallaz for access to the field sites. We also thank Antoine Guisan and the ECOSPAT group at the University of Lausanne for providing plant community and climatic data. We thank Jonathan Levine and

521 the Plant Ecology group for their feedback on an earlier version of the manuscript. S.L.
 522 acknowledges the Chinese Scholarship Council for financial support (No.201706100184).
 523 J.M.A. received funding from the European Union's Horizon 2020 research and innovation
 524 programme under grant agreement No. 678841.

525 **Author contributions**

526 J.M.A. designed the experiment. S.L. and J.M.A. collected data. S.L. conducted the
 527 modelling and data analysis. S.L. wrote the manuscript with input from J.M.A.

528 **Competing interests**

529 The authors declare no competing interests.

530

References

1. Darwin C. *On the Origin of species*. London: British Museum Natural History
Cambridge: Cambridge University Press (1859).
2. MacArthur RH. *Geographical ecology : patterns in the distribution of species*. Harper & Row
(1972).
3. Wisz MS, *et al*. The role of biotic interactions in shaping distributions and realised
assemblages of species: implications for species distribution modelling. *Biological Reviews*
88, 15-30 (2013).
4. Holt RD. Bringing the Hutchinsonian niche into the 21st century: Ecological and evolutionary
perspectives. *Proceedings of the National Academy of Sciences* **106**, 19659-19665 (2009).
5. Hargreaves AL, Samis KE, Eckert CG, Associate Editor: Oswald JS, Editor: Judith LB. Are
Species Range Limits Simply Niche Limits Writ Large? A Review of Transplant Experiments
beyond the Range. *The American Naturalist* **183**, 157-173 (2014).
6. Louthan AM, Doak DF, Angert AL. Where and When do Species Interactions Set Range
Limits? *Trends in Ecology & Evolution* **30**, 780-792 (2015).
7. Paquette A, Hargreaves AL. Biotic interactions are more often important at species' warm
versus cool range edges. *Ecology Letters* **n/a**, (2021).
8. Liang E, *et al*. Species interactions slow warming-induced upward shifts of treelines on the
Tibetan Plateau. *Proceedings of the National Academy of Sciences* **113**, 4380-4385 (2016).
9. Chesson P, Huntly N. The Roles of Harsh and Fluctuating Conditions in the Dynamics of
Ecological Communities. *The American Naturalist* **150**, 519-553 (1997).
10. Chan S-F, Shih W-K, Chang A-Y, Shen S-F, Chen I-C. Contrasting forms of competition set
elevational range limits of species. *Ecology Letters* **22**, 1668-1679 (2019).
11. Godsoe W, Jankowski J, Holt RD, Gravel D. Integrating Biogeography with Contemporary
Niche Theory. *Trends in Ecology & Evolution* **32**, 488-499 (2017).
12. Alexander JM, Diez JM, Usinowicz J, Hart SP. Species' Distributions as a Coexistence
Problem: A Response to Godsoe et al. *Trends in Ecology & Evolution* **33**, 144-145 (2018).
13. Barabás G, D'Andrea R, Stump SM. Chesson's coexistence theory. *Ecological Monographs*
88, 277-303 (2018).
14. Levine JM, HilleRisLambers J. The importance of niches for the maintenance of species
diversity. *Nature* **461**, 254-257 (2009).

- 576 15. Chesson P. Mechanisms of Maintenance of Species Diversity. *Annual Review of Ecology and*
577 *Systematics* **31**, 343-366 (2000).
578
- 579 16. Diamond JM. Distributional Ecology of New Guinea Birds: Recent ecological and
580 biogeographical theories can be tested on the bird communities of New Guinea. *Science* **179**,
581 759-769 (1973).
582
- 583 17. Harpole WS, Tilman D. Grassland species loss resulting from reduced niche dimension.
584 *Nature* **446**, 791-793 (2007).
585
- 586 18. Hart SP, Freckleton RP, Levine JM. How to quantify competitive ability. *Journal of Ecology*
587 **106**, 1902-1909 (2018).
588
- 589 19. Vittoz P, Randin C, Dutoit A, Bonnet F, Hegg O. Low impact of climate change on subalpine
590 grasslands in the Swiss Northern Alps. *Global Change Biology* **15**, 209-220 (2009).
591
- 592 20. Freeman BG. Lower elevation animal species do not tend to be better competitors than their
593 higher elevation relatives. *Global Ecology and Biogeography* **29**, 171-181 (2020).
594
- 595 21. Alexander JM, Diez JM, Levine JM. Novel competitors shape species' responses to climate
596 change. *Nature* **525**, 515-518 (2015).
597
- 598 22. Lax S, Abreu CI, Gore J. Higher temperatures generically favour slower-growing bacterial
599 species in multispecies communities. *Nature Ecology & Evolution* **4**, 560-567 (2020).
600
- 601 23. Davis AJ, Jenkinson LS, Lawton JH, Shorrocks B, Wood S. Making mistakes when predicting
602 shifts in species range in response to global warming. *Nature* **391**, 783-786 (1998).
603
- 604 24. Bertness MD, Callaway R. Positive interactions in communities. *Trends in Ecology &*
605 *Evolution* **9**, 191-193 (1994).
606
- 607 25. He Q, Bertness MD, Altieri AH. Global shifts towards positive species interactions with
608 increasing environmental stress. *Ecol Lett* **16**, 695-706 (2013).
609
- 610 26. Germain RM, Mayfield MM, Gilbert B. The "filtering" metaphor revisited: competition and
611 environment jointly structure invasibility and coexistence. *Biology Letters* **14**, 20180460
612 (2018).
613
- 614 27. Violle C, Pu Z, Jiang L. Experimental demonstration of the importance of competition under
615 disturbance. *Proceedings of the National Academy of Sciences* **107**, 12925-12929 (2010).
616
- 617 28. Hart SP, Marshall DJ. Environmental stress, facilitation, competition, and coexistence.
618 *Ecology* **94**, 2719-2731 (2013).
619
- 620 29. Levine JM, Bascompte J, Adler PB, Allesina S. Beyond pairwise mechanisms of species
621 coexistence in complex communities. *Nature* **546**, 56 (2017).

622

- 623 30. Matías L, Godoy O, Gómez-Aparicio L, Pérez-Ramos IM. An experimental extreme drought
624 reduces the likelihood of species to coexist despite increasing intransitivity in competitive
625 networks. *Journal of Ecology* **106**, 826-837 (2018).
626
- 627 31. Perez-Ramos IM, Matias L, Gomez Aparicio L, Godoy O. Functional traits and phenotypic
628 plasticity modulate species coexistence across contrasting environments. *bioRxiv*, 539619
629 (2019).
630
- 631 32. Godoy O, Levine JM. Phenology effects on invasion success: insights from coupling field
632 experiments to coexistence theory. *Ecology* **95**, 726-736 (2014).
633
- 634 33. Alexander JM, Levine JM. Earlier phenology of a nonnative plant increases impacts on native
635 competitors. *Proceedings of the National Academy of Sciences* **116**, 6199-6204 (2019).
636
- 637 34. Zimmermann NE, *et al.* Climatic extremes improve predictions of spatial patterns of tree
638 species. *Proceedings of the National Academy of Sciences* **106**, 19723-19728 (2009).
639
- 640 35. Vitasse Y, *et al.* Phenological and elevational shifts of plants, animals and fungi under climate
641 change in the European Alps. *Biological Reviews* **n/a**, (2021).
642
- 643 36. Alexander JM, *et al.* Lags in the response of mountain plant communities to climate change.
644 *Global Change Biology* **24**, 563-579 (2018).
645
- 646 37. Rumpf SB, *et al.* Extinction debts and colonization credits of non-forest plants in the
647 European Alps. *Nature Communications* **10**, 4293 (2019).
648
- 649 38. Rumpf SB, *et al.* Range dynamics of mountain plants decrease with elevation. *Proceedings of*
650 *the National Academy of Sciences* **115**, 1848-1853 (2018).
651
- 652 39. Angulo MT, Kelley A, Montejano L, Song C, Saavedra S. Coexistence holes characterize the
653 assembly and disassembly of multispecies systems. *Nature Ecology & Evolution*, (2021).
654
- 655 40. Cáceres CE. Temporal variation, dormancy, and coexistence: A field test of the storage effect.
656 *Proceedings of the National Academy of Sciences* **94**, 9171-9175 (1997).
657
- 658 41. Ke P-J, Wan J. Effects of soil microbes on plant competition: a perspective from modern
659 coexistence theory. *Ecological Monographs* **n/a**, e01391 (2019).
660
- 661 42. Körner C. *Alpine plant life : functional plant ecology of high mountain ecosystems*, Softcover
662 reprint of the hardcover 2nd ed. 2003 edn. Springer (2003).
663
- 664 43. Nomoto HA, Alexander JM. Drivers of local extinction risk in alpine plants under warming
665 climate. *Ecology Letters* **24**, 1157-1166 (2021).
666

667 44. Scherrer D, Vitasse Y, Guisan A, Wohlgemuth T, Lischke H. Competition and demography
668 rather than dispersal limitation slow down upward shifts of trees' upper elevation limits in the
669 Alps. *Journal of Ecology* **108**, 2416-2430 (2020).
670

671 45. Randin CF, *et al.* Climate change and plant distribution: local models predict high-elevation
672 persistence. *Global Change Biology* **15**, 1557-1569 (2009).
673

674 46. Kleyer M, *et al.* The LEDA Traitbase: a database of life-history traits of the Northwest
675 European flora. *Journal of Ecology* **96**, 1266-1274 (2008).
676

677 47. Ellner SP, Childs DZ, Rees M. *Data-Driven Modelling of Structured Populations : A Practical*
678 *Guide to the Integral Projection Model*. Cham : Springer (2016).
679

680 48. Reed PB, *et al.* Climate manipulations differentially affect plant population dynamics within
681 versus beyond northern range limits. *Journal of Ecology* **109**, 664-675 (2021).
682

683 49. Carroll IT, Cardinale BJ, Nisbet RM. Niche and fitness differences relate the maintenance of
684 diversity to ecosystem function. **92**, 1157-1165 (2011).
685

686 50. Chesson P. Multispecies Competition in Variable Environments. *Theoretical Population*
687 *Biology* **45**, 227-276 (1994).
688

689 51. Grainger TN, Letten AD, Gilbert B, Fukami T. Applying modern coexistence theory to priority
690 effects. *Proceedings of the National Academy of Sciences* **116**, 6205-6210 (2019).
691

692 52. R Core Team. R: A language and environment for statistical computing.). R Foundation for
693 Statistical Computing (2020).
694
695
696

Figure 1. Population growth rates across the elevation gradient. Effects of elevation on intrinsic (**a**, in the absence of neighbours) and invasion (**b**, in the presence of neighbours) population growth rate of lowland (orange) and highland (blue) focal species. Populations are predicted to persist when $\ln(\lambda) > 0$ (dashed horizontal line). The points show average ($\pm$ SE) population growth rates across species. Solid lines show significant relationships (test based on the mean of 500 bootstrap replicates; $P < 0.05$). Orange and blue shading indicates the 95% quantiles of fitted relationships across 500 bootstrap replicates of the dataset.

Figure 2. Competition outcomes across the elevation gradient. The estimated competitive outcomes (a-c) and strength of coexistence (i.e., coexistence metric, log-transformed; d-f) of lowland-lowland (orange), lowland-highland (green), and highland-highland (blue) species pairs in the three sites. In a-c, grey bars represent stable coexistence while white bars represent competitive exclusion and priority effects, and error bars indicate mean and 95% confidence intervals based on 500 bootstrap replicates. In d-f, solid lines indicate a significant interaction of elevation x competitor identity (test based on the mean of 500 bootstraps; $P < 0.05$), and the points are averages ($\pm$ SE) across species pairs. The shading indicates the 95% quantiles of fitted relationships across 500 bootstrap replicates of the dataset.

Figure 3. Niche differences and relative fitness differences across the elevation

gradient. Estimates of the niche differences (expressed as niche overlap, i.e. $1 - \text{niche differences}$; see Methods, **a-c**) and relative fitness differences (**d-f**) of lowland-lowland (orange), lowland-highland (green) and highland-highland (blue) species pairs. Solid lines indicate the significant interactions of elevation x competitor identity for niche differences and significant effects of elevation on relative fitness differences for each type of pair separately (test based on the mean of 500 bootstrap replicates; $P < 0.05$) while dotted lines indicate non-significant relationships ($P > 0.05$). The points are averages ($\pm$ SE) across species pairs. The shading indicates the 95% quantiles of fitted relationships across 500 bootstrap replicates of the dataset. In **d-f**, absolute values of fitness differences (log-transformed) are shown for lowland-lowland and highland-highland pairs, while actual values are shown for lowland-highland pairs, with negative values indicating lowland species are dominant competitors.