Title: Trends in *Arabidopsis* Research post genome sequencing- A Scientometric study

**Sandeep Kumar^{1,2\$}, Amar Kant Kushwaha^{1,3}, Thribhuvan R^{1,4}, Balakrishnan M¹,
Krishnan P^{1\$}**

¹ICAR-National Academy of Agricultural Research and Management, Hyderabad, India

²Quality and Productivity Improvement Division, ICAR-Indian Institute of Natural Resins and
Gums, Ranchi, India

³Division of Crop Improvement and Biotechnology, ICAR-Central Institute for Subtropical
Horticulture, Rehmankhera, P.O. Kakori, (Lucknow), India, 226101

⁴Division of Crop Improvement, ICAR-Central Research Institute for Jute and Allied fibres,
Barrackpore, India

\$ Corresponding authors

Sandeep Kumar & Krishnan P

Mail: Sandeep.kumar8@icar.gov.in; krishnanars@naarm.org.in

ICAR-National Academy of Agricultural Research and Management,
Hyderabad, Telangana, India-500030

1. ABSTRACT:

Arabidopsis thaliana, a model plant, is intensively researched because of the intrinsic advantages associated with its life cycle, genetics, and other characteristics. In the present study, we analysed the publication data of research work done on *A. thaliana* during 2001-2020, a period when the whole genome sequence of the plant was available to the researchers. The current meta-analysis showed that globally, 31965 research papers were published on *Arabidopsis* during the study period. The USA topped the countries with the maximum share of publications (28.97%), followed by China, Japan, and West European countries. The analysis showed a temporal shift in the number of publications from different countries and their research focus. After 2013, the research output from China was higher than that from the USA, and there was a shift in focus from developmental biology to stress-related topics. Plant Physiology journal carried the most research work (2490 publications) on *Arabidopsis*, which was followed by Plant Cell (2376), the Plant Journal (2260) and the Journal of Experimental Botany (1121). However, there was a progressive decline in the publication in these journals, and a shift was evident in favour of open access journals like Frontiers in Plant Science. The publication and citation numbers also show ongoing *Arabidopsis* research's relevance to plant sciences in general, especially in curiosity-driven and discovery-based science. This article delves into the patterns in the prominence of research areas, ideas and foci for the future *Arabidopsis* research roadmap.

2. Introduction

Arabidopsis thaliana is a plant native to the Eurasian and African region, where it is generally considered a weed that often grows along the roadsides. Researchers across the globe have used this plant to study basic plant biology. It was first described by a German physician named Johannes Thal, on whose name Carl Linnaeus named it *Arabis thaliana*. The name *Arabidopsis thaliana* was given by Gustav Heynhold, who created a new genus by the name of *Arabidopsis* (Meyerowitz, 2001). It is self-pollinated, annual and grows up to 20-25cm. The key qualities, including small plant size, a relatively shorter life cycle period, seed setting in large quantities and smaller genome size with fewer chromosome number qualifies it as the best candidate as a model plant and amenable for genetic and physiological studies (Rédei, 1970; Somerville and Koornneef, 2002; Koornneef and Meinke, 2010). Primary emphasis was given to the screening and identification of auxotrophic mutants (Rédei, 1975), as a result of which the first *Arabidopsis* gene sequences came out in 1986 (Chang and Meyerowitz, 1986). After that, T-DNA-mediated transformation and vacuum infiltration made it more amenable to studies (Koncz *et al.*, 1989). Eventually *Arabidopsis* turned to be a plant amenable for generating gene mutations and implementing DNA transformation protocols. The culmination of these projects was the National Science Foundation (NSF) project on *Arabidopsis* gene identification and whole-genome sequencing (*Arabidopsis* Genome Initiative, 2000). This project was one of many initiatives of genome sequencing. Various technological advancements happened after the completion of *Arabidopsis* genome sequencing which shaped the biological research in many organisms. The genome sequencing brought a tremendous pace and shift in *Arabidopsis* research (Bevan and Walsh, 2005). Multinational *Arabidopsis* Steering Committee (MASC) has been actively involved in *Arabidopsis* research and constantly prepares the roadmaps for *Arabidopsis* research (<http://arabidopsisresearch.org/index.php/en/about>). The research interest in *Arabidopsis* as a model plant in the emerging research scenarios has been dwindling following the emergence of high throughput sequencing and other genomics tools (Fig. 1). In addition, there has been an increased application of these newer tools to non-model plants. The current research was designed to analyse the trends in research on the *Arabidopsis thaliana*, in the post-genome sequencing period, with the specific objectives of

- (i) Mapping the variation in the theme areas of *Arabidopsis* research and
- (ii) Profile the trends in research networks at various levels *viz.*, authors, organisations and countries.

Fig. 1 Timeline of key technological advancements in Genome studies after *A. thaliana* genome Sequencing. (Adapted from *Nature Milestones: Genomic Sequencing* <https://www.nature.com/collections/genomic-sequencing-milestones> (2021))

4. Materials and methods

4.1 Data collection and analysis

This study uses the bibliometric methodology to quantify scientific collaborations in *Arabidopsis* research throughout the world. The Web of Science (WoS) database was used for literature search on *Arabidopsis*. Web of Science is a website published by Clarivate Analytics that offers subscription-based access to numerous databases that provide detailed citation data for a wide range of academic subjects (Li *et al.*, 2018). The search string and filters used to select the publications were as follows (T.I. = (*Arabidopsis*)) AND DOCUMENT TYPES: (Article) Timespan: 2001-2020. Indexes: SCI-EXPANDED, CPCI-S, ESCI. The year 2001 was chosen as the starting point for the survey since *Arabidopsis* genome sequencing had been finished by 2000 (*Arabidopsis* Genome Initiative, 2000). Conference proceedings and books/book chapters were excluded from the study. The search phrase was chosen based on our investigation of existing literature and discussions among authors and subject matter experts. The *thaliana* was not added to the string as our preliminary scanning showed that many publications did not use the full binomial name in the titles and used only '*Arabidopsis*'.

The data retrieved from the WoS database was exported in text format, with each record containing the bibliometric data: title name, journal names, author names with Web of Science affiliation, number of citations, document kinds, year of publication for each article, authors, Web of Science keywords, and abstracts. The WoS database was also utilised for publication

trends over the years, for which five year moving average of publications was used. The Scientometric/bibliometric parameters like the total number of publications and citations, average citations per article (ACPP), h-index, research areas, collaborating countries and funding agencies were extracted from the WoS platform for 20 years (2000-2020).

4.3 Network analysis:

The co-authorship and keyword occurrence maps were generated using the VOS viewer version 1.6.16 using metadata generated from the web of science ([van Eck and Waltman, 2011](http://www.vosviewer.com/)) (<http://www.vosviewer.com/>). For the network analysis, top 100 results were selected and used in case of both institutional collaboration and year wise keyword occurrence. Bibliometrix, an R-based tool, was used for data filtering, research theme evolution and identifying journals based on Bradford's law wherein the metadata was fed in BibTeX files (<http://www.bibliometrix.org>) (Aria and Cuccurullo, 2017).

5. Results & Discussion

5.1 Global output of research on *Arabidopsis*

The publication of the *Arabidopsis* genome led to a significant rise in the *Arabidopsis* research as it expanded the area of research from only isozyme and protein study to functional genomics (Mysore *et al.*, 2001; Nath Radhamony *et al.*, 2005; Bevan and Walsh, 2005). The study showed that 31965 research articles were published on the subject during the 20 years from 2001 to 2020 (Fig. 2). One intriguing element of the total number of articles was a five-year consecutive decline in the total number of publications following 2013, except for 2019/2020. This decreasing trend in recent years can be attributed to increased emphasis on sequencing of commercially significant species such as rice, maize, pigeon pea, rapeseed, tomato and many others (International Rice Genome Sequencing Project, 2005; Schnable *et al.*, 2009; Varshney *et al.*, 2011; Wang *et al.*, 2011). This paradigm shift was made possible due to the remarkable progress of sequencing techniques appropriate for diverse purposes, from high cost and poor throughput to low cost and high throughput (Türktaş *et al.*, 2015).

In the keyword co-occurrence, we found a clear shift from developmental pathways related to the peripheral pathways like stress responses and the emergence of some new keywords like transcriptome and proteomics in the last decade (Fig. 3). It also demonstrates the widespread usage of various next-generation genomic tools and techniques in the previous decade (2011-2020). The *Arabidopsis* gene functions have also been elucidated in the major signalling

pathways (Alonso and Stepanova, 2004; Atkinson *et al.*, 2013). Bolker (2012) emphasised on the need to go beyond model organisms to get complete picture of various biochemical and signalling pathways in taxonomic groups. Moreover, the elucidation of biological function through *Arabidopsis* has trickled into the agriculturally important crop sp. for their validation and utilisation (Priyanka *et al.*, 2010; Manmathan *et al.*, 2013; Mishra *et al.*, 2014).

Figure 2. Research output in *Arabidopsis* over the years

5.2 Highly cited studies on Arabidopsis

After whole-genome sequencing of *Arabidopsis*, the research papers on insertional mutagenesis in *Arabidopsis thaliana* received more citations, thus indicating the research trends (Table 1). Before sequencing, it was tough to create a loss of function mutation. Till the sequencing of the first plant, only unicellular organisms were the subject for the creation of loss of mutation for understanding the function of the gene(s) (Giaever *et al.*, 2002) and gene silencing and RNAi were utilised for understanding the developmental role of genes in *C. elegans* with their limitations of unstable heritability, residual expression of genes and low throughput (Ashrafi *et al.*, 2003). After publication of the paper titled "Genome-wide Insertional mutagenesis of *Arabidopsis thaliana*" by Alonso *et al.* (2003) depicting the technique overcoming the limitation of earlier methods, work started in *Arabidopsis* mutant generation and their use in research. The article was found to influence a large stream of research, as indicated by the maximum citation (3790) that any *Arabidopsis* paper received in the last two decades and laid the foundations of functional biology in *Arabidopsis*.

Similarly, with the advancement in sequencing techniques, epigenomes were targeted to decipher the developmental pathway. To detect such epigenetic modification, Cokus *et al* (2008) came up with a new approach, BS-seq (Bisulphite sequencing), combining sequence information and bisulphate treatment for mapping the methylation sites in the whole genome. Similarly, in the same area, Lister *et al* (2008) mapped the methylation sites by sequencing smRNA, methylcytosine to decipher the role in the developmental pathway. The significance of these two articles is indicated by the citations received by them, viz., 1516 and 1363, respectively. Some other papers featured on the top 10 list by the number of citations were related to transcript normalisation, transient gene expression and microarray database. In short, the protocol and database development paper featured prominently on the most cited list. It is in line with the evolution of *Arabidopsis* as a model plant, evidenced by the fact that novel tools and techniques to decipher plant biological functions has been developed in *Arabidopsis* and later extended to the other crop plants (Koornneef and Meinke, 2010).

Figure 3: Thematic evolution based on the major keywords co-occurrence in two decades (2001-10) & (2011-20)

164 **Table 1:** Highly cited research publications on *Arabidopsis* since its full genome was sequenced

No	Title of the paper	Author and year	Affiliation of corresponding author	Journal	Citations (WoS)	
1.	Genome wide insertional mutagenesis of <i>Arabidopsis thaliana</i>	Alonso J.M. et al., 2003	Salk Inst Biol Studies, Genome Anal Lab, 10010 N Torrey Pines Rd, La Jolla, CA 92037 USA.	Science	3790	165 166 167 168 169 170 171
2.	<i>Arabidopsis</i> mesophyll protoplasts: a versatile cell system for transient gene expression analysis	Yoo S.D., Cho Y.H. and Sheen J. (2007)	Massachusetts Gen Hosp, Dept Mol Biol, Boston, MA 02114 USA.	Nature Protocols	2344	172 173 174
3.	Genome-wide identification and testing of superior reference genes for transcript normalisation in <i>Arabidopsis</i>	Czechowski, T. et al., 2005	Max Planck Inst Mol Plant Physiol, D-14476 Potsdam, Germany.	Plant Physiology	2018	175 176 177 178 179
4.	GENEVESTIGATOR. <i>Arabidopsis</i> microarray database and analysis toolbox	Zimmermann P. et al., 2004	ETH, Inst Plant Sci, CH-8092 Zurich, Switzerland.	Plant Physiology	1888	180 181 182
5.	A gene expression map of <i>Arabidopsis thaliana</i> development	Schmid, M. et al., 2005	Max Planck Inst Dev Biol, Spemannstr 37-39, D-72076 Tübingen, Germany	Nature Genetics	1787	183 184 185 186
6.	MAP kinase signalling cascade in <i>Arabidopsis</i> innate immunity	Asai, T. et al., 2002	Harvard Univ, Sch Med, Dept Genet, Boston, MA 02114 USA.	Nature	1684	187 188 189 190

5.3 Top countries working on the Arabidopsis research

Based on the number of publications, America contributed 28.97% of the total publications, followed by China, Germany, and Japan with 21.23%, 14.11% and 12.06% of the publications, respectively. Table 2 shows the top 10 countries and their per cent contribution, along with other publication metrics *viz.*, h-index, T.C. (Total Citations) and ACPP (Average Citation Per Publication). The h-index, which indirectly represents the research influence, was higher for publications from the USA (133), which was followed by Germany (223), England (204) and Japan (200). Though China has the second-highest per cent share in total research papers, its h-index is relatively lower, i.e., 166 and the lowest ACPP score of 29.96 compared to other top 10 countries. The country collaboration map shown in Fig. 5 provides a clear picture regarding the collaborations. The United States of America and the countries from Western Europe like the UK, France and Germany showed a very high collaboration and scientific output. Some African and Baltic countries, however, were missing from the Arabidopsis research map entirely. The countries leading the research in terms of the number of publications were part of the genome sequencing consortium or those collaborating with them (https://www.nsf.gov/bio/pubs/reports/arab_prog78.htm). Though China was not the part as they were not much into *Arabidopsis* research before 1995 (Chen *et al.*, 2006), its research output has been high in recent years (Fig 4). A plausible explanation of this is the acquisition of natural and mutant stocks of Arabidopsis seeds post-genome sequencing (Provart *et al.*, 2016).

Regarding subject areas, most publications are featured in the plant sciences and developmental biology across the countries (Table 3). Cell biology and developmental biology were laid less stress in China, possibly due to more research being done later. However, the research areas of publication journals are indicative, and there might exist many cross-cutting and multidisciplinary studies which do not follow this classification.

Figure 4: Country-wise publication over the different period

Table 2: Top 10 countries which published most research papers on *Arabidopsis* in the world during 2001-2020

No .	Country	TP	% Share to N	h-index	ACPP	TC	WSC	% Share of self-citation
1	USA	9262	28.97%	313	75.92	703214	640485	8.9
2	PR China	6787	21.23%	166	29.96	203311	179953	11.5
3	Germany	4510	14.11%	223	66.71	300856	279745	7.0
4	Japan	3857	12.06%	200	56.75	218899	204002	6.8
5	France	2613	8.17%	188	64.89	169568	160796	5.2
6	England	2447	7.65%	204	75.74	185338	177949	4.0
7	S. Korea	1685	5.27%	122	39.78	67025	63347	5.5
8	Canada	1422	4.44%	127	51.15	72733	69609	4.3
9	Spain	1333	4.17%	129	56.49	75304	72118	4.2
10	Australia	979	3.06%	137	64.65	63293	61007	3.6

TP=Total publications, TC= Total citations, WSC=Without-self citation, ACPP= average citation per paper

218

219 **Figure 5:** Country-wise Scientific production based on the number of publications and Interaction among countries publishing articles on *A.*
 220 *thaliana*. The blue colour's intensity corresponds to the total number of publications from the corresponding country, and the red lines
 221 highlight the interactions among countries (co-authorship).

Table 3: Top 10 Country vs Subject field wise (No. of publications)

Country /Region	Subject							Total *
	Plant Sciences	Bio- chemistr y and Molecul ar biology	Cell Biology	Muti- disciplinar y Sciences	Geneti c heredit y	Bio- technolo gy applied micro- biology	Develop mental biology	
USA	5549	3014	1760	1053	795	419	243	9262
China	4351	1976	753	616	410	361	55	6787
Germany	2885	1479	857	407	313	131	106	4510
Japan	2559	1206	1031	285	194	366	75	3857
France	1583	885	507	223	193	86	59	2613
England	1495	770	477	257	158	88	66	2447
S. Korea	1080	595	320	96	65	110	17	1685
Canada	1011	367	191	89	114	94	30	1422
Spain	880	392	194	133	82	57	40	1333
Australia	689	251	149	90	45	28	18	979

*The total number of articles in sub-subjects do not add up as only the five major subjects are mentioned here, and there exist many journals which fall under more than one category

5.4 Leading institutions on the Arabidopsis research:

Collaborations between different institutes revealed the close associations between American and European institutes (Fig. 5). The top three institutions with the most publications were Chinese Academy of Science, University of California, and University of Tokyo. The institutes in Asia had relatively weak partnerships with other countries. The North American and European institutes were active in the early research periods as visible from the corresponding colour. The Chinese Academy of Science and other Chinese Universities have published more papers than those from Japan and Western countries.

The highest papers feature the Chinese Academy of Sciences (Fig. 6). However, the articles from western Europe had relatively high-quality parameters, e.g. Chinese Academy of science

has 1856 total publications, but the average citation per paper is lower (41.69) than the U.K. Research Innovation (UKRI) which has only 779 total papers but 90.03 average citations per paper.

5.5 Top Journals for Arabidopsis Research:

Plant Physiology (2376), Plant cell (2490) and the Plant Journal (2260) were the top three journals in terms of the number of relevant publications, h-index, and total citations (Table 4). Bradford law is an essential consideration that decides the importance of different journals based on diminishing citation of relevant articles over the year, following Pareto distribution (Nash-Stewart *et al.*, 2012). Here it can be easily seen that Plant Physiology, Plant Cell, Plant Journal, Journal of Experimental Botany, Plant and Cell Physiology, Frontiers in Plant Science and PNAS were the core sources that carried most publications on Arabidopsis and thus represent the choice journals for the researchers (Fig. 6). The annual pattern of publication on Arabidopsis in these journals showed a shift in the preferred journals over the years (Fig. 7), i.e. a decline in the publication in the Plant cell, plant physiology and the plant journal; however, open publications models of frontiers in plant sciences have gained in popularity. As a result, none of the top three journals based on the total number of publications was present among the top three for the recent year, i.e. 2020. This trend is probably due to the citation advantage of an article in open access (O.A.) due to an increase in the number of readers citing open access articles (Cullen and Chawner, 2011). The radical initiative to O.A. rather than a subscription with 'Plans S' (science, speed, solution, shock) by 11 funding agencies in Europe indicates such implication (Else, 2018).

256

257 **Figure 6:** Collaboration among different institutions working on *Arabidopsis* in the first decade (2001-2010) vs second decade (2011-2020)

Figure 7: Quantitative and qualitative parameters in the research backed by different funding organisations (2001-2020)

Table 4: Top journals that published the majority of Arabidopsis research articles from 2001 to 2020

No.	Title of the journal	TP	% share	TC	WCS	H-Index	ACP P	IF (2020)
1	<i>Plant Physiology</i>	2490	7.79	226960	220788	204	91.15	6.902
2	<i>Plant Cell</i>	2376	7.43	287284	277416	248	120.21	9.618
3	<i>Plant Journal</i>	2260	7.07	179334	174911	190	79.35	6.141
4	<i>Journal of Experimental Botany</i>	1121	3.51	44776	44089	99	39.94	5.908
5	<i>Plant and Cell Physiology</i>	999	3.12	46314	45320	105	46.36	4.062
6	<i>Frontiers in Plant Science</i>	950	2.97	13016	12696	44	13.7	4.407
7	<i>PNAS</i>	878	2.75	102939	101908	180	117.24	9.412
8	<i>PLoS ONE</i>	829	2.59	23548	23269	67	28.41	2.740
9.	<i>Plant Signaling Behaviour</i>	804	2.52	8810	8728	36	10.96	1.671
10.	<i>Plant Molecular Biology</i>	677	.12	33566	33234	90	49.58	3.302

Figure 8: The Bradford law displaying key Arabidopsis journals

Figure 9: Performance of core journals (as defined by Bradford's law) over 2001-2020 period based on total number of annual articles on 'Arabidopsis'

Conclusion

We set out to determine the trajectory of Arabidopsis research in various areas. With the sequencing of the Arabidopsis genome, there was a rise in interest in the research, but it decreased after the advent and popularisation of new sequencing tools. Publication in Arabidopsis increased steadily in the first two quarters after the genome sequencing, reducing in the third quarter. Primary research interests spanned from mutant identification to protein investigation, as seen by co-word analysis, whereas gene expression experiments were constant throughout 20 years. Some of the highly cited articles were related to developing and validating novel techniques in plant science research.

Furthermore, Asian countries that have shown a strong interest in Arabidopsis research must collaborate closely with their western counterparts to improve the quality and reach. Quality is a significant consideration, as evidenced by the low number of citations, particularly from China, which has shown a rising trend in the number of publications. The funds all over the world are depicted more toward the number of publications than ACPP. The target journal should be criteria based on impact factors for grants by the funding agency, depicting quality (IF) rather than quantity (T.P.). This can result in good ACPP score, although ACCP is affected by several other factors also viz., journal rank, primary/reporting author's h-index, SCImago journal rank (SJR) and source normalised impact per paper (SNIP) (Yaminfirooz and Ardali, 2018). This can be represented from the author collaboration point of view as it also has a significant correlation with citation, and it has been witnessed that lower collaboration among Asian countries compared to among European countries and USA. A substantial part of Arabidopsis papers has gone to three leading journals: Plant Physiology, Plant Cell, and Plant Journal. The growing trend toward open access publication is unmistakably the way forward for journals seeking to remain relevant.

Conflict of interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Bibliography

- Alonso, J.M., Stepanova, A.N., Leisse, T.J., et al.** (2003) Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science*, **301**, 653–657.
- Alonso, J.M. and Stepanova, A.N.** (2004) The ethylene signaling pathway. *Science*, **306**, 1513–1515.
- Arabidopsis Genome Initiative** (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature*, **408**, 796–815.
- Aria, M. and Cuccurullo, C.** (2017) bibliometrix : An R-tool for comprehensive science mapping analysis. *Journal of Informetrics*, **11**, 959–975.
- Ashrafi, K., Chang, F.Y., Watts, J.L., Fraser, A.G., Kamath, R.S., Ahringer, J. and Ruvkun, G.** (2003) Genome-wide RNAi analysis of *Caenorhabditis elegans* fat regulatory genes. *Nature*, **421**, 268–272.
- Atkinson, N.J., Lilley, C.J. and Urwin, P.E.** (2013) Identification of genes involved in the response of *Arabidopsis* to simultaneous biotic and abiotic stresses. *Plant Physiol.*, **162**, 2028–2041.
- Bevan, M. and Walsh, S.** (2005) The *Arabidopsis* genome: a foundation for plant research. *Genome Res.*, **15**, 1632–1642.
- Bolker, J.** (2012) Model organisms: There’s more to life than rats and flies. *Nature*, **491**, 31–33.
- Chang, C. and Meyerowitz, E.M.** (1986) Molecular cloning and DNA sequence of the *Arabidopsis thaliana* alcohol dehydrogenase gene. *Proc Natl Acad Sci USA*, **83**, 1408–1412.
- Chen, H., Karplus, V.J., Ma, H. and Deng, X.W.** (2006) Plant biology research comes of age in China. *Plant Cell*, **18**, 2855–2864.
- Cokus, S.J., Feng, S., Zhang, X., et al.** (2008) Shotgun bisulphite sequencing of the *Arabidopsis* genome reveals DNA methylation patterning. *Nature*, **452**, 215–219.
- Cullen, R. and Chawner, B.** (2011) Institutional Repositories, Open Access, and Scholarly Communication: A Study of Conflicting Paradigms. *The Journal of Academic Librarianship*, **37**, 460–470.
- Else, H.** (2018) Radical open-access plan could spell end to journal subscriptions. *Nature*, **561**, 17–18.
- Giaever, G., Chu, A.M., Ni, L., et al.** (2002) Functional profiling of the *Saccharomyces cerevisiae* genome. *Nature*, **418**, 387–391.

345 **International Rice Genome Sequencing Project** (2005) The map-based sequence of the
346 rice genome. *Nature*, **436**, 793–800.

347 **Koncz, C., Martini, N., Mayerhofer, R., Koncz-Kalman, Z., Körber, H., Redei, G.P. and**
348 **Schell, J.** (1989) High-frequency T-DNA-mediated gene tagging in plants. *Proc Natl Acad*
349 *Sci USA*, **86**, 8467–8471.

350 **Koornneef, M. and Meinke, D.** (2010) The development of Arabidopsis as a model plant.
351 *Plant J.*, **61**, 909–921.

352 **Li, K., Rollins, J. and Yan, E.** (2018) Web of Science use in published research and review
353 papers 1997–2017: a selective, dynamic, cross-domain, content-based analysis.
354 *Scientometrics*, **115**, 1–20.

355 **Lister, R., O'Malley, R.C., Tonti-Filippini, J., Gregory, B.D., Berry, C.C., Millar, A.H.**
356 **and Ecker, J.R.** (2008) Highly integrated single-base resolution maps of the epigenome in
357 Arabidopsis. *Cell*, **133**, 523–536.

358 **Manmathan, H., Shaner, D., Snelling, J., Tisserat, N. and Lapitan, N.** (2013) Virus-
359 induced gene silencing of Arabidopsis thaliana gene homologues in wheat identifies genes
360 conferring improved drought tolerance. *J. Exp. Bot.*, **64**, 1381–1392.

361 **Meyerowitz, E.M.** (2001) Prehistory and history of Arabidopsis research. *Plant Physiol.*,
362 **125**, 15–19.

363 **Mishra, S., Shukla, A., Upadhyay, S., et al.** (2014) Identification, occurrence, and
364 validation of DRE and ABRE Cis-regulatory motifs in the promoter regions of genes of
365 Arabidopsis thaliana. *J. Integr. Plant Biol.*, **56**, 388–399.

366 **Mysore, K.S., Tuori, R.P. and Martin, G.B.** (2001) Arabidopsis genome sequence as a tool
367 for functional genomics in tomato. *Genome Biol.*, **2**, REVIEWS1003.

368 **Nash-Stewart, C.E., Kruesi, L.M. and Del Mar, C.B.** (2012) Does Bradford's Law of
369 Scattering predict the size of the literature in Cochrane Reviews? *Journal of the Medical*
370 *Library Association: JMLA*, **100**, 135.

371 **Nath Radhamony, R., Mohan Prasad, A. and Srinivasan, R.** (2005) T-DNA insertional
372 mutagenesis in Arabidopsis: a tool for functional genomics. *Electron. J. Biotechnol.*, **8**.

373 **Priyanka, B., Sekhar, K., Sunita, T., Reddy, V.D. and Rao, K.V.** (2010) Characterization
374 of expressed sequence tags (ESTs) of pigeonpea (*Cajanus cajan* L.) and functional
375 validation of selected genes for abiotic stress tolerance in Arabidopsis thaliana. *Molecular*
376 *Genetics and Genomics*, **283**, 273–287.

377 **Provart, N.J., Alonso, J., Assmann, S.M., et al.** (2016) 50 years of Arabidopsis research:
378 highlights and future directions. *New Phytol.*, **209**, 921–944.

379 **Rédei, G.P.** (1970) Arabidopsis thaliana (L.) Heynh. A review of the genetics and biology.
380 *Bibliographia Genetica.*, **20**.

381 **Rédei, G.P.** (1975) Induction of auxotrophic mutations in plants. In *Genetic manipulations*
382 *with plant material*. Springer, pp. 329–350.

383 **Schnable, P.S., Ware, D., Fulton, R.S., et al.** (2009) The B73 maize genome: complexity,
384 diversity, and dynamics. *Science*, **326**, 1112–1115.

385 **Somerville, C. and Koornneef, M.** (2002) A fortunate choice: the history of Arabidopsis as
386 a model plant. *Nat. Rev. Genet.*, **3**, 883–889.

387 **Türktaş, M., Yücebılgılı Kurtoğlu, K., Dorado, G., Zhang, B., Hernandez, P. and Ünver,**
388 **T.** (2015) Sequencing of plant genomes – a review. *Turk. J. Agric. For.*, **39**, 361–376.

389 **Varshney, R.K., Chen, W., Li, Y., et al.** (2011) Draft genome sequence of pigeonpea
390 (*Cajanus cajan*), an orphan legume crop of resource-poor farmers. *Nat. Biotechnol.*, **30**,
391 83–89.

392 **Wang, Xiaowu, Wang, H., Wang, Jun, et al.** (2011) The genome of the mesopolyploid crop
393 species *Brassica rapa*. *Nat. Genet.*, **43**, 1035–1039.

394 **Yaminfirooz, M. and Ardali, F.R.** (2018) Identifying the Factors Affecting Papers’
395 Citability in the Field of Medicine: an Evidence-based Approach Using 200 Highly and
396 Lowly-cited Papers. *Acta Inform. Med.*, **26**, 10–14.