The *Eucalyptus grandis* chloroplast proteome: leaf development and seasonal variations

Amanda Cristina Baldassi^{1¶} and Tiago Santana Balbuena^{1*&}

¹ Department of Agricultural, Livestock and Environmental Biotechnology, São Paulo State University (UNESP), School of Agriculture and Veterinary Sciences, Jaboticabal, São Paulo, Brazil.

*Corresponding author

E-mail: tiago.balbuena@unesp.br (TSB)

¶ This author carried out the experiment, processed and analyzed the data and wrote the manuscript with support from T. S. B. All authors discussed the results and contributed to the final manuscript.

& This author designed the study, supervised the project and contributed to the interpretation of the results. All authors discussed the results and contributed to the final manuscript.

Abstract

Chloroplast metabolism is very sensitive to environmental fluctuations and is intimately related to plant leaf development. Characterization of the chloroplast proteome dynamics may contribute to enlarge the understanding on plant adaptation to different climate scenarios and leaf development processes. Herein, we carried out a discovery-driven proteome analysis of the *Eucalyptus grandis* chloroplast proteome during leaf maturation and throughout different seasons of the year. The chloroplast proteome from young leaves differed the most from all assessed samples. Most up-regulated proteins identified in mature and young leaves were those related to catabolic-redox signaling and biogenesis processes, respectively. Seasonal dynamics revealed unique proteome features in the autumn and spring periods. The most abundant chloroplast protein in

humid (wet) seasons (spring and summer) was a small subunit of RuBisCO, while in the dry periods (fall and winter) the proteins that showed the most pronounced accumulation were associated with photo-oxidative damage, Calvin cycle, shikimate pathway, and detoxification. Our investigation of the chloroplast proteome dynamics during leaf development revealed significant alterations in relation to the maturation event. Our findings also suggest that transient seasons induced the most pronounced chloroplast proteome changes over the year. This study contributes to a more comprehensive understanding on the subcellular mechanisms that lead to plant leaf adaptation and ultimately to *Eucalyptus grandis* productivity. Mass spectrometric data are available via ProteomeXchange under identifier PXD029004.

Introduction

Chloroplast is a subcellular compartment in which solar radiation is used to generate sugar compounds in a complex process named photosynthesis. It is also the cellular compartment that hosts several other metabolic processes that are essential for plant survival, such as synthesis of lipids, fatty acids, vitamins, hormones, secondary metabolites, in addition to most of the proteogenic amino acids found in plant cells, like methionine, cysteine, and threonine [1-5].

Plastids are composed of inner and outer membrane, stroma, thylakoidal membrane, and lumen [5]. Chloroplast genome encodes approximately 120-130 proteins, whilst most of the chloroplast proteome, comprising approximately 2000 proteins, are synthesized in the cytosol, and imported into the organelle through the envelope membranes [6-7]. Chloroplast genome varies from 107 kb to 218 kb in size depending on the plant species [8]. In *Eucalyptus grandis*, the chloroplast genome represents approximately 160 kb [9-10].

As chloroplasts are very sensitive to environmental fluctuations, dynamic proteome changes may represent a key factor for understanding their responses to abiotic stresses [11]. The most recent chloroplast proteomics studies have aimed to identify (1) changes induced by biotic and abiotic stresses [12-16], (2) mechanisms of remodeling and reprogramming of plant metabolism [17,18], (3) biogenesis and senescence [19-21], (4) chloroplast development [22,23], in addition to (5) novel techniques for increasing proteome coverage [24-26].

Chloroplast biogenesis starts from proplastids, which, upon exposure to light, turn into mature chloroplasts with functional electron transport carriers, membrane channel proteins, and stromal enzymes that are able to capture, assimilate, and store carbon [27]. During senescence, they differentiate from gerontoplasts, which lead to leaf yellowing, necrosis, and death [28]. Thereby, plastid development is intimately related to the

development of the whole plant at different stages [32]. The manipulation of several agronomic traits, *e.g.*, photosynthetic rate, stomatal conductance, chlorophyll content, and PSII photochemical, has been reported to delay the senescence process and increase plant biomass, seed yield, seed protein composition, and tolerance to abiotic stresses [33, 34, 35].

Plants are sessile organisms, which means that they are subjected to seasonal climate alterations and adverse environmental conditions. Seasonal climate variations can drive plant growth and development, favoring a metabolic adaptation of the life cycle with changes in the environment [36]. Strong seasonal effects have been observed on the growth of perennial plants due to varying availability of water, light, nutrients, and temperature [37, 38]. In *Populus* species, for instance, seasonal phenology is largely related to changes in growth and development, such as bud flush in spring followed by vegetative growth in the summer; while during the fall, the trees present growth inhibition and bud formation, followed by leaf senescence, abscission, and cold acclimation before the winter season [39]. Regarding photosynthetic physiological changes, studies have revealed that the quantum efficiency seasonal pattern shows minimal values in the spring, increasing during the summer, and reaching maximal values in the fall followed by a decrease in the winter [40, 41].

Understanding functional and regulatory networks based on leaf development process and seasonal metabolic changes can provide valuable information for increasing plant productivity and learning the mechanisms that lead to plant adaptation. This is the first exploratory investigation on proteome changes during leaf development and seasonal variations in *Eucalyptus grandis* young plants on an organellar level. This research aimed to carry out an exploratory analysis of chloroplast alterations during leaf development and seasonal variations in the *Eucalyptus grandis* chloroplast proteome.

Material and Methods

Plant cultivation and sampling

Eucalyptus grandis plants were cultivated at the “Horto Florestal” of the São Paulo State University (Jaboticabal – SP, Brazil) using a randomized block design experiment with 2m x 3m spacing. Fertilization and irrigation were carried out twice a month and three times a week, respectively. The region climate is defined as tropical and characterized by rainfalls concentrated in the summer and dry winter [42]. Agrometeorological data were collected daily at the UNESP environmental station (S1 Table).

For the leaf developmental assay, three biological replicates, comprising five randomly selected branches, were divided into three different regions (young, middle, and mature), according to the fluorescence data (FV/FM) and chlorophyll relative quantification (CCI), and used to assess proteome changes throughout development (S1A Fig). For the chloroplast proteome seasonal variation assay, four biological replicates, comprising forty leaves from the first until the fifth node, were assessed in all four seasons from August/2019 to August/2020 (S1B Fig). For both assays using leaf sampling, plant material was immediately transferred to ice before isolating the chloroplasts, as described in the next sections.

Fluorescence Measurement and Chlorophyll Relative Quantification

Fluorescence analysis was performed on the fluorometer FluorPen FP 100 (Photon Systems Instruments) assessing the adaxial surface of leaves from different regions of the 15 branches. The fluorescence values obtained directly from the device were used to assess the potential quantum efficiency of photosystem II (FV/FM). Relative chlorophyll was quantified on CCM-200 equipment (Opti-Sciences), and the adaxial surface of leaves from different regions of the 15 branches was examined. The CCI

(chlorophyll content index) values obtained directly from the device were used to assess relative chlorophyll quantification. Statistical analysis was performed according to the ANOVA test ($p < 0.05$), and Tukey's test was used for comparison among different leaves.

Chloroplast proteome isolation

Aliquots of six grams of leaves were collected, fine sliced, and transferred to tubes containing 20 mL of chloroplast isolation buffer (CIB, CP-ISO-Sigma Aldrich). The leaf tissue was homogenized (Omni GLH Turrax) and filtered through a nylon membrane. The processed material was centrifuged, and the chloroplasts were isolated in an 80%/40% (1:2 v/v) Percoll density gradient. After washing, the chloroplast pellet was resuspended in a protein extraction buffer (Thiourea 2M; CHAPS; Tris-HCl 1M, pH 8; Glycerol 50%; Triton X-100; DTT 1M and Urea 8M) and the lysis of intact chloroplasts was carried out through 20 cycles of sonication. After centrifugation, the supernatant was precipitated in cold acetone for 16 hours at -30°C , and the protein pellet obtained was solubilized in Tris buffer (125 mM Tris pH 6.8, 20% glycerol, 1% SDS, 0.1% DTT).

Total leaf proteome isolation

To assess the enrichment of the chloroplast subproteome in relation to the total leaf proteome, intact leaves were frozen in -80°C and grinded in Tris extraction buffer (125 mM Tris, pH 6.8, 20% glycerol, 1% SDS, 0.1% DTT). Upon centrifugation at 4°C for 15 min, the supernatant was transferred to a clean tube and the proteins were precipitated in cold acetone for 16 hours at -30 °C. Finally, the proteins were solubilized in Tris buffer before protein quantification assay.

Protein quantification assay and SDS-PAGE

Sample quantifications followed the Bradford method using a BSA curve as reference. Aliquots of 50 µg of proteins were separated for 1 hour by denaturing electrophoresis on a polyacrylamide gel containing sodium dodecyl sulfate (SDS-PAGE) to obtain a single band. The electrophoretic system was composed of a stacking gel (5%) and a separation gel (12.5%). The single protein band was excised from the gel and subjected to in-gel digestion.

Mass spectrometry analysis

Proteins were in-gel digested using trypsin at 1:10 proportion (trypsin:protein) [43]. To clean up the peptide samples, extracts were filtered in an acetate cellulose membrane (Spin-X, Corning) by

centrifugation for 7 minutes. The flow-through was collected using a clean tube, concentrated, and then desalted with ZipTip Pipette Tips (Millipore) according to the manufacturer's protocol. The amount of 1 µg of extracted peptides was analyzed through liquid chromatography coupled to a tandem mass spectrometer. Tryptic peptides were resuspended in H₂O containing 0.1% formic acid and separated in an EASY-nLC1000 chromatographic system (Thermo Fisher Scientific) for 95 minutes using a C18 nano-column under a constant flow of 400 nL/min and increasing concentrations of acetonitrile (7 to 30% over 70 min and then 70% in 25 min). The peptides were ionized through electrospray (3.5 kV) and the mass spectrometer (Q-Exactive, Thermo Fisher Scientific) was operated in data-dependent acquisition mode according to a top 20 acquisition method with dynamic exclusion time adjusted to five seconds. Full scans and MS/MS scans were acquired with resolutions equal to 70,000 and 35,000 FWHM (full width at half maximum), respectively. The MS/MS spectra were generated from HCD fragmentation (high-energy collision dissociation) of the precursor peptide ions isolated under 35 eV collision energy.

Protein identification

The peptides and protein inferences were identified through stringent searches of experimental data against the protein database *Eucalyptus grandis* v.2.0 (<http://www.phytozome.net/>) on MaxQuant (v. 1.6.3.3)

[44]based on dedicated search parameters: 20 ppm tolerance for precursor ions, oxidation of methionines and carbamidomethylation of cysteines – selected as dynamic and static modifications, respectively. Maximum missed cleavages of 2 and false-discovery rate adjusted to 1%. Relative protein of the chloroplast proteins were quantified based on the label free-based approach (LFQ intensity). Relative abundance of total leaf proteins was obtained following the NSAF ("normalized spectral abundance factor") approach [45]. Mass spectrometry proteomics data are currently stored on the ProteomeXchange Consortium via PRIDE partner repository under dataset identifier PXD029004.

It is worth noting that the ProteomeXchange dataset PXD029004 has not yet been publicly released, therefore, reviewers may require the following information to access mass spectrometric data:

Username: reviewer_pxd029004@ebi.ac.uk

Password: WwsYFMTy

Statistical analysis and functional annotation

Data processing and statistical analysis were carried out on the Perseus software (v. 1.6.15.0) [46]. Firstly, all contaminants and reverse sequences were removed from data analysis. Subsequently, label-free

quantification data (LFQ) were transformed into log2 scale, and a protein was considered a true hit if identified in at least two out of three replicates. Missing values imputation was carried out from the normal distribution curve from each biological replicate. The means of ANOVA test and Tukey comparison test ($p < 0.05$) were used in the comparative analyses. Multivariate analyses were carried out with differentially abundant proteins for both assays on the Perseus software. For PCA and hierarchical clustering, data were z-score normalized. Hierarchical clustering was performed using Euclidean distance and complete linkage.

Functional annotation of the differentially abundant proteins was carried out using the Blast2GO platform ($p < 0.05$) [47]. the ten most representative level-four biological processes were considered in the functional characterization aiming to balance between a comprehensive overview and detailed information on the targeted biological process. Level four was chosen for including photosynthesis, an important process in the context of leaves and chloroplasts.

Subcellular prediction

In silico analysis was performed using predictors of subcellular location and chloroplast proteome repositories to assess and confirm the cellular location of the identified proteins. For this purpose, a protein was considered plastidial if identified by at least one of the subcellular

prediction algorithms – ChloroP [48] and Predotar [49] – or by presenting at least one homologous protein in any of the databases used – PPDB [50] or AT-Chloro [51].

Results and Discussion

Before performing any description or comparative proteome analysis, we decided to verify the efficiency of the chloroplast enrichment strategy used herein. For this purpose, we compared the abundance levels, in terms of NSAF and LFQ intensities, of a well-known chloroplast proteoform Eucgr.C03525.1 (Ribulose-bisphosphate carboxylase – RBCL) against five potential mitochondrial contaminants commonly found in chloroplast isolations (Table 1). Quantitative analysis indicated that the chloroplast proteoform was enriched upon the isolation process, whereas the mitochondrial proteoforms presented a decrease in the abundance levels. These results indicated that our chloroplast isolation and enrichment strategy was effective in reducing the number of cross-contaminants, as well as in the enrichment process of chloroplast-related proteins in the target samples.

Table 1. Enrichment analysis of chloroplast proteins in *Eucalyptus grandis* using a comparative approach between total leaf protein extract and chloroplast protein

266 **extract.** Local: Localization, M: Mitochondria, C: Chloroplast, Leaf: Total leaf protein
 267 extract, Chloro: Chloroplast protein extract, FD: Fold-Change.

Description ^a	Loc ^b	NSAF ^c				LFQ intensity ^d			
		Leaf	Chloro	FD		Leaf	Chloro	FD	
Contaminants									
Malate dehydrogenase (Eucgr.F01209.1)	M	0.0036	0.0010	3.52	↓	3.E+08	2.E+07	15.39	↓
Isocitrate dehydrogenase (Eucgr.A01135.1)	M	0.0010	0.0005	2.27	↓	2.E+08	1.E+07	10.98	↓
Citrate synthase (Eucgr.G03412.1)	M	0.0004	0.0000	10.98	↓	3.E+07	2.E+06	16.04	↓
Fumarase (Eucgr.H01081.1)	M	0.0001	0.0000	3.65	↓	7.E+06	4.E+05	20.04	↓
Succinate dehydrogenase (Eucgr.F00243.1)	M	0.0002	0.0002	1.02	↓	2.E+06	2.E+06	1.13	↓
Positive control									
Ribulose-bisphosphate carboxylase-large (Eucgr.C03525.1)	C	0.0155	0.0276	1.78	↑	3.E+09	3.E+09	1.19	↑
268	^a Functional Annotation and protein identifier according to the E. grandis v 2.0 annotation								
269	^b Subcellular protein location								
270	^c Relative protein expression by the NSAF approach								
271	^d Relative protein expression by the LFQ intensity approach								
272									

273 **Photosynthetic proteins comprise most of the *Eucalyptus***
 274 ***grandis* chloroplast proteome.**

275 Aiming to illustrate a general overview of the chloroplast proteome
 276 isolated from *Eucalyptus grandis* leaves, we compiled the identification
 277 datasets from both leaf development and seasonal experiments. The
 278 combined analysis resulted in a stringent, non-redundant dataset of 431
 279 chloroplast proteins (2S Table). Gene ontology analysis indicated that most
 280 of the isolated proteins were involved in the “organonitrogen compound
 281 metabolic process”, “organic substance biosynthetic process”, and
 282 “macromolecule metabolic process” (Fig 1A). Although these biological
 283 processes were the most prominent in terms of protein identifications, an
 284 MS1-based quantitative analysis indicated that the “photosynthesis

process” represented 11% of the isolated chloroplast proteomes (Fig 1B). The fraction of “photosynthetic process” related to proteins becomes even more pronounced within the isolated proteome upon the analysis of the most abundant proteins identified in the entire experiment (Fig 1C). The RuBisCO small chain (Eucgr.J01502.2.p) was the most abundant protein identified herein, showing approximately 4000 MS/MS counts. The proteins involved in the photosystem II structure were also highly abundant: two extrinsic proteins (PsbO-2 – Eucgr.I01025.1.p and PsbQ-2 – Eucgr.D00854.1.p) that form the oxygen-evolving complex (OEC) and a light-harvesting complex protein from the PSII (Eucgr.F04099.1.p) corresponding to 1500 to 3000 MS/MS counts (Fig 1C).

Regarding the suborganellar localization of the 431 chloroplast proteins, 137 proteins were present in the stroma compartment, whilst 135 were thylakoidal proteins. A total of 87 proteins were predicted *in silico* to be located in the chloroplast outer membrane, and 13 identifications were found to be part of the chloroplast inner membrane. Finally, 52 proteins were located in the chloroplast lumen and 7 proteins herein identified had been previously reported as part of the plastoglobule structure (Fig 1D).

Fig 1. The *Eucalyptus grandis* chloroplast proteome. Classification analysis, in terms of the number of identified proteins (A) and terms of the percentage of identified proteins (B), upon categorization using the “biological process” ontology; protein expression analysis of the Gene

Ontology terms (C); number of protein identifications per chloroplast sub-cellular compartment (D).

Chloroplast proteome reflects the physiological status in developing leaves

Based on the varying Fv/Fm ratios and the differences observed from the relative chlorophyll quantitation analysis (S2 Fig), we isolated the chloroplast proteome from young, middle, and mature leaves. Quantitative data from the isolated proteins were used to demonstrate the changes within this subcellular structure throughout the leaf development process. Pairwise analysis indicated that 61 proteins were considered to be differentially abundant over the development process, with the young chloroplast proteome as the most dissimilar of the three different regions (2A Fig). Such result is in accordance with physiological evidence highlighted in S1 Fig, as the young leaf tissue presented the most significant variations concerning middle and mature regions. A study on sugarcane leaves have also demonstrated that physiological parameters, such as chlorophyll content, nitrogen content, and RubisCO activity, were greater and more consistent between mature and transition tissues than in young regions [52].

Clustering analysis allowed to classify the differentially abundant proteins (DAP) in three distinctive groups: up-regulated – 17 proteins with higher abundance over leaf development, down-regulated – a set of 29

proteins with lower abundance along the studied process, and an undefined group, comprising 15 proteins with an unclear abundance pattern (2B Fig).

Fig 2. Multivariate analyses of the 61 differentially abundant chloroplast proteins isolated from young, middle, and mature *Eucalyptus grandis* leaves. Principal component analysis (A), clustering expression analysis (B), and functional categorization of the differentially-regulated chloroplast proteins between young (down-regulated) and mature (up-regulated) (C). The z-score indicates the distance of the data compared to the mean value.

Proteins with higher abundance throughout leaf development comprised those involved in the antioxidant metabolism (APX4, CAT2, tAPX, and SODFe2), stress responses/protection against lipid peroxidation (IDH-V, MPC1, and CHL), and photosynthesis (PSBQ-2, PSBO-2, GAPB, and FTRA2). Additionally, proteins involved in post-transcriptional modifications (CBR), regulation of PSII (TLP), amino acid metabolism (PYD2), photorespiration (GCT), carboxylation, and CO₂ diffusion (CA1) and plastoglobuli development (FIB4) were also found to be up-regulated. Conversely, antioxidants and detoxification (TPX, AOR 1, and AOR2), and photosynthesis (RCA, RBCS1, RBCS2, and PSBS) were observed during leaf development in addition to lower abundance, in ribosomal proteins (RPL13, PRPL1, RBD1, RBD2, and RPS1), which are related to different biosynthetic (CRD1, PORA, THI1, PLP and IPP2, GSA1, LPD1, and FKB), folding, modification, and transport (TPR, HSC70-2 and CLPC1,

CURT1C, ENR1, LTP1, and LTP2) processes,. Proteins with unclear abundance pattern or those presenting a transient increase or decrease in the abundance levels were found to be involved in the photosynthetic process and electron transport chain (LHB1, PSAF, and THOL), in addition to folding, modification, and transport (HSP70, HSP21, ENR2, OEP24A, and RABE1B) and biosynthetic processes (NDPK1, MEE32, and PSAT). Proteins responsible for peptide degradation/repair of photosynthetic machinery (PREP2 and FtsH), oxidoreductase, and protease (CHLP and ASP) were also clustered within such heterogeneous group (S3 Table).

Aiming to highlight changes in the chloroplast metabolic processes along leaf development, up and down-regulated proteins were cross-referenced to corresponding gene ontology terms. Functional categories revealed differences between the chloroplast protein groups (2C Fig). Leaf development induced “cellular catabolic processes” in *Eucalyptus grandis* chloroplasts for being the most representative category within the up-regulated group. Responses to stress, mainly to oxidative stress and cellular detoxification, were the most abundant functional category in up-regulated throughout leaf development, based on the maturity of the photosynthetic apparatus and the higher photosynthesis rates in young leaves. In non-stressing conditions, mature leaves are expected to maintain cellular homeostasis with a dynamic balance between the light energy absorbed and energy consumption [53]. In mature leaves, generation of reactive oxygen

species and changes in the redox state in the photosynthetic electron transport chain are important indicators of operational retrograde signals. The function of operational retrograde signaling differs from biogenic signaling as it focuses on the adjustments and cellular homeostasis of the chloroplast in response to different perturbations and developmental cues [54].

Conversely, the most representative functional class identified in the chloroplast down-regulated protein group (*i.e.*, abundant in young leaves) were involved with cellular and organic biosynthetic processes, being the most representative in the “organic substance biosynthetic process” and the “nitrogen compound metabolic process”. These results are under the developmental stage of the young leaves used here. As plastids are intimately related to developmental and environmental signals, chloroplasts from young leaves were expected to reflect the biogenesis processes, which involves large-scale gene transcription followed by synthesis, importing, and assemblies of biogenesis proteins, such as those involved in plastid translation apparatus and enzymes in the synthesis of isoprenoids and tetrapyrroles [55]. In turn, in mature leaves, redox-related proteins comprised most of the differentially regulated proteins; the only category suggested by gene ontology analysis in young leaves for “cell redox homeostasis” comprised only 3% of all down-regulated proteins. A study on the developmental dynamics of maize leaves also found up-regulated

genes that encode enzymes involved in diverse biosynthetic processes in young leaves. Similarly, in mature leaves, the up-regulated genes predominant encoded Calvin cycle, redox regulation, and the light reactions of photosynthesis [55].

For better insights into the changing patterns of the up-and down-regulated proteins, we classified the chloroplast proteins into three expression trends, as shown in Fig 3. The early responsive proteins comprised the subset with a drastic change in the abundance level in middle leaf tissues, remaining at similar levels in both mature and young leaves. Conversely, late responsive proteins showed a drastic abundance change only in the contrasting leaf tissue (young or mature leaves). We considered as almost linear responsive the proteins without drastic changes throughout the leaf development process, but with a significant change between mature and young tissues.

Early up-regulated proteins (*i.e.*, early increased abundance along leaf development) were identified as follows: Mitochondrial pyruvate carrier, Isocitrate dehydrogenase V, Ferredoxin/thioredoxin reductase subunit A2, Pyrimidine 2, Carbonic anhydrase 1, and Glycine cleavage T-protein family (MPC1, IDH-V, FTRA2, PYD2, CA1, and GCT). The most pronounced increase occurred in the Pyrimidine 2 (PYD2) protein from young to middle tissues. This protein acts in beta-alanine biosynthesis and nitrogen recycling from pyrimidines to general nitrogen metabolism [56].

Late up-regulated proteins (*i.e.*, late increased abundance along leaf development) were identified as follows: Photosystem II subunit Q-2, Glyceraldehyde-3-phosphate dehydrogenase B subunit, and Chloroplast lipocalin (PSBQ-2, GAPB, CHL). The most pronounced increase in abundance chloroplast lipocalin (CHL) presented. This protein acts in the protection of thylakoidal membrane lipids against reactive oxygen species, especially singlet oxygen, induced by high light and drought stresses occurring mainly at early stress stages in *Arabidopsis* [57]. It has also been reported that stress conditions cause the CHL to protect the thylakoid membrane by facilitating a sustained non-photochemical quenching (NPQ) in the LHCII in *Arabidopsis* [58]. A nearly increasing linear responsive pattern throughout leaf development was found in the following chloroplast proteins: Ascorbate peroxidase 4, Catalase 2, Thylakoidal ascorbate peroxidase, Fe superoxide dismutase 2, Thylakoid lumen 18.3 kDa protein, Photosystem II subunit O-2, Chloroplast RNA binding, and Plastid-lipid associated protein PAP/fibrillin (APX4, CAT2, tAPX, SODFe2, TLP, PSBO-2, CBR, and FIB4).

Early down-regulated proteins (*i.e.* early decreased abundance along leaf development) was the largest trend group according to our categorization (16 proteins) comprising proteins involved in the biosynthesis of fatty acids, isoprenoids, and amino acids (LPD1, IPP2, and PLP), chloroplast RNA binding proteins (RBD1, RBD2, and DAG),

ribosomal proteins acting in the biosynthesis of thylakoid membrane and thylakoid curvature (RPS1 and CURT1C), transport, modification and import into the chloroplast (LTP2, TPR, and CLPC1), proteins associated with detoxification (TPX1, AOR1, and AOR2), binding cofactors (ENR1), and sucrose metabolism (ALD). a nearly linear down-regulated responsive trend appeared in 10 protein sequences showed that act in carbon fixation RubisCO (RCA, RBCS1, and RBCS2), chlorophyll biosynthesis (CRD1, PORA, GSA1), transport and folding (LTP1 and HSC70-2), and structural ribosomal proteins (RPL13 and PRPL1). In turn, only two proteins showed a late down-regulated pattern, pfkB-like carbohydrate kinase family protein and Thiazole biosynthetic enzyme, chloroplast (FKB and THI1).

Fig 3. Expression trends of the differentially regulated chloroplast proteins. Up-regulated proteins increased abundance throughout leaf development, while down-regulated proteins decreased abundance during leaf development.

In conclusion, while investigating the chloroplast proteome changes over leaf development, we found significant alterations in the target proteome regarding the maturation process. Out of the 431 proteins isolated from the chloroplast material, 61 showed significant changes in the abundance throughout leaf development. There was no clear bias towards a unique change in the proteome composition; instead, we found distinct patterns of protein accumulation. In addition to pinpointing proteins with

pronounced changes ranging the studied phenomenon, through a quantitative proteomics approach, we also specified the most particular biological and metabolic processes involving leaf development. However, as the proteome of any living entity is constantly changing, we also decided to investigate the undisturbed (*i.e.*, without inducing anthropogenic alterations due to external stimulus) sub-cellular proteome dynamics by analyzing the seasonal changes occurring in the chloroplast proteome of *Eucalyptus grandis* plants cultivated at the State of Sao Paulo Forest Nursery between September 2019 and August 2020.

Transient seasons induce the most pronounced chloroplast proteome changes

Meteorological data revealed only small changes in the mean and maximum temperatures recorded throughout the study period, with the minimum diurnal temperature ranges observed during the summer season. Precipitation analysis revealed extremely dry fall and winter seasons and mean monthly rainfall of almost 10 mm over the summer (Fig 4A). Multivariate analysis of the *E. grandis* chloroplast proteomes suggested that both the fall- and the spring-related samples presented the most dissimilar profiles of the studied seasons (Fig 4B). This is likely to be related to the phenological transitions commonly occurring in the species

during such periods. In *Populus* species, phenology is largely related to seasonal changes in growth and development, such as bud flush in spring followed by vegetative growth in summer. In turn, during the fall, the trees presented growth cessation and bud formation, followed by leaf senescence and abscission, in addition to cold acclimation for the winter [39]. Our comparative proteome analysis revealed that 41 proteins were differentially regulated across the seasons of the year in *E. grandis* (4S Table).

Fig 4. Meteorological data recorded at the Sao Paulo State University Forest Nursery during *Eucalyptus grandis* cultivation. Changes in temperature and rainfall rates between August 2019 and September 2020 (A). Multivariate analysis of the chloroplast proteome samples collected in the summer, fall, winter, and spring (B).

Hierarchical clustering analysis of DAP proteins suggests 5 different clusters according to their abundance patterns (Fig 5A). As the transition seasons (fall and spring) presented the most dissimilar profiles, it is worth highlighting the clusters that showed the most expressive differences in these specific periods.

Cluster 1 comprises 10 identifications with the lowest abundance values in the spring season. Proteins are related to light-harvesting complex, carbonyl detoxification, folding and modifications, carbohydrate biosynthesis, hydrolases, amino acid precursors, and ATP carriers. Conversely, Cluster 5 is composed of 6 proteins that showed higher

abundance in the spring. Most of these identifications are associated with RuBisCO regulation to enhance photosynthetic capacity. Other identifications are responsible for lipid transport, protein modification, and translation (Fig 5B).

Other studies have revealed that quantum efficiency seasonal pattern has minimal values in the spring, increasing over the summer, reaching maximal values in the fall, and finally decreasing in the winter [40, 41]. Such scenario is demonstrated in Cluster 1, with proteins LHB1 and LHCB4, which participate in the first steps of photosynthesis and are very efficient in capturing light energy and transforming light into chemical energy [59]. Photosynthetic capacity in plants has also been reported to fluctuate with increased levels throughout leaf development, reaching its peak in the spring or early summer, and remaining stable or gradually declining over the summer [60-62]. Such pattern appears in Cluster 5, with RuBisCO-related proteins (RCA, RBCS1, and 2). Leaf fluorescence and photosynthesis are expected to increase over the spring, especially in chlorophyll content. In the fall, these parameters are reduced when associated with leaf physiological properties such as leaf aging and senescence [63].

Cluster 4 presented the highest abundance for the fall season (Fig 5B), with 7 identifications associated with the following processes: PSII regulation, chlorophyll biosynthesis, translation, phospholipid transport; in

addition to two histones that play an important role in regulating transcription and DNA repair and replication.

As previously described, trees undergo growth cessation and establish dormancy before the onset of winter over a transition season. Low temperatures were found to facilitate photoinhibition but not necessarily to cause it [64]; in addition, downregulation of Calvin-Benson cycle activity and gas exchange precedes the reorganization of photosynthetic apparatus in thylakoid membrane during the fall [65]. All these events are partially related to PSII regulation and corroborate our analysis, where we found an up-regulated protein related to the regulation of PSII in Cluster 4. Furthermore, a plant epigenome is very responsive to changes in the environment, and variations in epigenetic silencing mechanisms are among the key factors that assist plants to adapt to different climate conditions [66]. DNA methylation and histone modifications associated with chromatin remodeling were reported to be the major changes occurring throughout dormancy season transitions [67]. The up-regulation of two histones in the fall occurring in Cluster 4 could be associated with these epigenetics events.

Fig 5. Differentially abundant proteins identified in *Eucalyptus grandis* chloroplasts in different seasons of the year. Hierarchical clustering (A) and protein abundance trends (B). The z-score indicates the distance of the data compared to the mean value.

Aiming to investigate which chloroplast proteins had the most pronounced synthesis in each season, we carried out a fold-change analysis, as shown in Fig 6. The red boxes indicate proteins with significant change equal or higher than 2-folds, blue boxes show significant change equal or higher than 5-folds, and white boxes point to proteins with change lower than 2-folds.

The most significant abundance increase had in both wet seasons (spring and summer) occurred in the RBCS1 protein. This is one of the small subunits of the RuBisCO enzyme, whose down-regulation has been observed in *Cistus albidus* [68] in drought periods, such as fall and winter,. In *Lolium perenne*, the RbcS transcripts reached the highest levels over the spring, gradually decreasing between the summer and winter [69]. Similarly, all photosynthetic parameters measured in *Quercus coccinea* Muench (scarlet oak) showed a peak in the late summer, followed by a sharp decline in the fall [70].

Fig 6. Most differentially abundant chloroplast proteins isolated from *Eucalyptus grandis* leaves in studies seasons. Red boxes indicate proteins with significant change higher than 2-folds. Blue boxes indicate a significant change higher than 5-folds. White boxes indicate proteins with change lower than 2-folds.

In contrast, the the most abundant proteins in both dry seasons (fakk and winter) were MPH1, PORA, MEE32, TPI, and AOR. MPH1 (proline-rich family protein) is a PSII-associated proline-rich protein that

participates in the maintenance of normal PSII activity and protection of PSII from photooxidative damage under excessive light conditions [71]. The PORA protein (Protochlorophyllide oxidoreductase A) catalyzes the reduction of protochlorophyllide to chlorophyllide, subsequently producing chlorophyll a and b [72,73]. In addition to its role in chlorophyll biosynthesis, PORA can also exert a photoprotective action in the transition from dark to light [74], whereas the *Triosephosphate isomerase* enzyme (TPI) interconverts glyceraldehyde 3-phosphate (G3P) in the Calvin cycle placed in chloroplasts. Up-regulation of TPI in the winter has been reported in experiments performed on tubers and *Eucalyptus grandis* bark [75,76].

The bifunctional enzyme MEE32 (dehydroquinase dehydratase, putative/shikimate dehydrogenase) is responsible for steps three and four of the shikimate pathways, catalyzing the dehydration of dehydroquinase to dehydroshikimate and the reversible reduction of dehydroshikimate to shikimate, respectively. Transgenic lines of tobacco with suppressed MEE32 gene displayed severe growth retardation and reduced content of aromatic amino acids and downstream products such as lignin and chlorogenic acid [77]. In *Eucalyptus camaldulensis*, it confers aluminum tolerance [78]. AOR protein (Oxidoreductase, zinc-binding dehydrogenase family protein) detoxify α , β -unsaturated carbonyls by reducing a highly electrophilic α , β -unsaturated bond using NAD(P)H [79, 80]. It favors the detoxifying of stromal lipid peroxide-derived reactive carbonyls (RCs)

produced under oxidative stress in chloroplasts [80], in addition to protecting dark respiration and supporting plant growth during the night [81].

In conclusion, our findings suggest that transient seasons (spring and fall) induce the most pronounced chloroplast proteome changes over the year. Out of the 431 proteins isolated in the chloroplast proteome, 41 showed significant abundance changes in different seasons. We also found distinct patterns of protein abundance among the seasons and were able to pinpoint the most abundant proteins in both the dry and wet periods of the year.

The importance of studying chloroplasts metabolic changes

Chloroplast biogenesis starts with proplastids that are exposed to light and then transformed into mature chloroplasts, which differentiate from gerontoplasts during senescence. Leaf senescence can be triggered by both external and internal factors. Internal factors include age, reproductive growth, redox metabolism, and hormone levels [82-84]. Leaf premature senescence can cause productivity losses due to lower assimilatory capacity [85]. Investigating the complex molecular network behind the developmental processes is essential for understanding the continual transition of functional and regulatory metabolic pathways occurring multiple times throughout the perennial plants lifespan. It is also quite

relevant for agriculture because [86-87] it may open new possibilities to increase crop yield and establish strategies to maximize biomass production.

An inter-organellar transcriptomic analysis of leaf senescence in *Arabidopsis* indicated that the chloroplast transcriptome exhibited the most pronounced changes throughout the plant lifespan [88]. These findings indicate that plastid compartment could be an important target to plant breeding programs that intend to delay leaf senescence and maximize biomass production. Regarding chloroplast metabolism, a large number of differentially expressed transcripts were identified in mature to senescence stages, and the biological processes associated with biogenesis at the growth to maturation stage showed higher expression, suggesting robust transcriptional coordination during early leaf development and more dynamic biochemical transitions during the senescence process [88]. At the protein level, we also observed higher abundance in biogenesis processes in young leaf tissue; in contrast, we detected a larger number of differential abundance proteins for the young stage rather than in mature tissues. This indicates that protein-level dynamics diverges from transcriptome dynamics. The plants also presented a strong translational coordination during early leaf development, and such biochemical networks seem more active during early stages.

Plants are sessile organisms and need to develop survival strategies due to changing environmental conditions. Perennial plants have been described to be strongly affected by seasonally varying availability of nutrients, water, light, and temperatures [37, 38]. Seasonal climate variations can drive plant growth and development, favoring a metabolic adaptation of the life cycle with changes in the environment [36].

Our findings suggest that transient seasons (spring and fall) induce the most pronounced chloroplast proteome changes over the year in *Eucalyptus grandis* cultivated in tropical climate. For the fall, we found higher abundance of proteins related to epigenetic events that are very sensitive to external environment changes. Such epigenetic mechanisms are among the factors that help plants adapt to different environmental climates [66]. The combination of seasonal proteomics and epigenomics studies in natural environments can be important to reveal seasonal dynamics regulation at each level on plant responses plasticity.

Moreover, agrometeorological data revealed extremely dry autumn and winter seasons and mean monthly rainfall of almost 10 mm over the summer. Photosynthetic organisms have established several strategies to cope with dry environments, including short life cycle, enhanced water uptake and reduced water loss, osmotic adjustment, and antioxidant capacity [89]. *Eucalyptus* species have been reported to have developed a strategy involving lower photosynthetic activity during dry periods

associated with resource reallocation through major changes in the gene expression of primary metabolism [90]. Identifying these metabolic changes allows to further investigate and understand the mechanisms behind plants adaptation and assist tree breeding programs.

Final Remarks

Investigation of *Eucalyptus grandis* chloroplast proteome provided a comprehensive resource for understanding different processes underlying age-dependent and seasonal variation in an organellar level. Our chloroplast isolation and enrichment strategies were effective at reducing the number of cross-contaminants and allowed a deep investigation of the changes occurring in the chloroplast proteome throughout leaf development and the seasons of the year. Distinct patterns of protein accumulation were detected in both assays. The *Eucalyptus grandis* chloroplast proteome seems prone to catabolic and redox processes as leaves age, while the accumulation of proteins involved in biosynthetic processes are down-regulated. Most pronounced proteome changes occurred in the transient seasons of the year with a clear proteome bias towards photosynthesis over the wet seasons.

A deeper understanding on age-associated biological networks is important for developing strategies to maximize the biomass production.

Furthermore, analyzing seasonal dynamics unravels the molecular mechanisms underlying metabolism plasticity and plant adaptation in spatiotemporal networks.

References

1. Mifflin B, Lea, P. Amino acid metabolism. Annu. Rev. Plant Biol. 1977; (28):299-29.
2. Ravanel S, Gakière B, Job D, Douce R. The specific features of methionine biosynthesis and metabolism in plants. Proc. Natl. Acad. Sci. 1998; 95(13):7805-12.
3. Hesse H, Hoefgen R. Molecular aspects of methionine biosynthesis. Trends Plant Sci. 2003;(8):259-62.
4. Kamal AHM, Cho K, Choi JS, Bae KH, Komatsu S, Uozumi N, Woo SH. The wheat chloroplastic proteome. J Proteomics. 2013; (93):326-42.
5. Van Wijk KJ. Plastid proteomics. Plant Physiol Biochem. 2004; 42 (12):963-77.
6. Keegstra K, Cline K. Protein import and routing systems of chloroplasts. Plant cell. 1999; 11(4):557-70.
7. Rossig C, Reinbothe C, Gray J, Valdes O, Von Wettstein D, Reinbothe S. Three proteins mediate import of transit sequence-less precursors into the inner envelope of chloroplasts in *Arabidopsis thaliana*. Proc. Natl. Acad. Sci. USA. 2013; 110:19962-67.
8. Daniell H, Lin CS, Yu M, Chang WJ. Chloroplast genomes: diversity, evolution, and applications in genetic engineering. Genome biol. 2016; 17(1):134.
9. Myburg AA, Grattapaglia D, Tuskan GA, Hellsten U, Hayes RD, Grimwood J, et al. The genome of *Eucalyptus grandis*. Nature. 2014; 510(7505):356-62.
10. Pinard D, Myburg AA, Mizrahi E. The plastid and mitochondrial genomes of *Eucalyptus grandis*. BMC Genomics. 2019;20:132.
11. Taylor NL, Tan YF, Jacoby RP, Millar AH. Abiotic environmental stress induced changes in the *Arabidopsis thaliana* chloroplast,

- mitochondria and peroxisome proteomes. J Proteomics. 2009;72:367–78.
12. Tamburino R, Vitale M, Ruggiero A, Sassi M, Sannino L, Arena S, et al. Chloroplast proteome response to drought stress and recovery in tomato (*Solanum lycopersicum* L.). BMC Plant Biol. 2017;17(1):40.
13. Chang L, Wang L, Peng C, Tong Z, Wang D, Ding G, et al. The chloroplast proteome response to drought stress in cassava leaves. Plant Physiol Biochem. 2019;142:351-62.
14. Megias E, Do Carmo LST, Nicolini C, Silva LP, Blawid R, Nagata T, Mehta A. Chloroplast Proteome of *Nicotiana benthamiana* Infected by Tomato Blistering Mosaic Virus. Protein J. 2018;37(3):290-99.
15. Zhu D, Luo F, Zou R, Liu J, Yan Y. Integrated physiological and chloroplast proteome analysis of wheat seedling leaves under salt and osmotic stresses. J Proteomics. 2021;234:104097.
16. Herrmann HA, Dyson BC, Miller MAE, Schwartz JM, Johnson GN. Metabolic flux from the chloroplast provides signals controlling photosynthetic acclimation to cold in *Arabidopsis thaliana*. Plant Cell Environ. 2021; 44(1):171-85.
17. Lande NV, Barua P, Gayen D, Kumar S, Varshney S, Sengupta S, Chakraborty S, Chakraborty N. Dehydration-induced alterations in chloroplast proteome and reprogramming of cellular metabolism in developing chickpea delineate interrelated adaptive responses. Plant Physiol Biochem. 2020a;146:337-48.
18. Watson SJ, Sowden RG, Jarvis P. Abiotic stress-induced chloroplast proteome remodelling: a mechanistic overview. J Exp Bot. 2018;69(11):2773-81.
19. Xiong E, Li Z, Zhang C, Zhang J, Liu Y, Peng T, Chen Z, Zhao Q. A study of leaf-senescence genes in rice based on a combination of genomics, proteomics and bioinformatics. Brief Bioinform. 2021;22(4):1–35.
20. Pipitone R, Eicke S, Pfister B, Glauser G, Falconet D, Uwizeye C, et al. A multifaceted analysis reveals two distinct phases of chloroplast biogenesis during de-etiolation in *Arabidopsis*. eLife. eLife Sciences Publication. 2021; 10.

21. Huokko T, Ni T, Dykes GF, Simpson DM, Brownridge P, Conradi FD, et al. Probing the biogenesis pathway and dynamics of thylakoid membranes. *Nat Commun.* 2021; 12 (1):3475.
22. Mielke K, Wagner R, Mishra LS, Demir F, Perrar A, Huesgen PF, Funk C. Abundance of metalloprotease FtsH12 modulates chloroplast development in *Arabidopsis thaliana*. *J Exp Bot.* 2021;72(9):3455-73.
23. Flannery SE, Hepworth C, Wood WHJ, Pastorelli F, Hunter CN, Dickman MJ, et al. Developmental acclimation of the thylakoid proteome to light intensity in *Arabidopsis*. *Plant J.* 2021;105 (1):223-44.
24. Bouchnak I, Brugière S, Moyet L, Le Gall S, Salvi D, Kuntz M, et al. Unraveling Hidden Components of the Chloroplast Envelope Proteome: Opportunities and Limits of Better MS Sensitivity. *Mol Cell Proteomics.* 2019;18 (7):1285-06.
25. Lande NV, Barua P, Gayen D, Kumar S, Chakraborty S, Chakraborty N. Proteomic dissection of the chloroplast: Moving beyond photosynthesis. *J Proteomics.* 2020b;212:103542.
26. Bayer RG, Stael S, Teige M. Chloroplast Isolation and Enrichment of Low-Abundance Proteins by Affinity Chromatography for Identification in Complex Proteomes. *Methods Mol Biol.* 2021;2261:535-47.
27. Biswal B, Pandey J. Development of Chloroplast Biogenesis, Senescence, and Regulations. *Handbook of photosynthesis*. Third edition. CRC press; 2016.
28. Thomas H. Senescence, aging and death of the whole plant. *New Phytol.* 2013;197: 696–11.
29. Rottet S, Besagni C, Kessler F. The role of plastoglobules in thylakoid lipid remodeling during plant development. *Biochim Biophys Acta Bioenerg.* 2015;1847(9):889-99.
30. Andriankaja M, Dhondt S, De Bodt S, Vanhaeren H, Coppens F, De Milde L, et al. Exit from proliferation during leaf development in *Arabidopsis thaliana*: a not-so-gradual process. *Dev Cell.* 2012;22: 64–78.
31. Thatcher SR, Burd S, Wright C, Lers A, Green PJ. Differential expression of miRNAs and their target genes in senescing leaves and

- siliques: insights from deep sequencing of small RNAs and cleaved target RNAs. *Plant Cell Environ.* 2015; 38:188–200.
32. Biswal B, Biswal UC, Raval MK. Chloroplast biogenesis: From proplastid to gerontoplast. Dordrecht: Kluwer Academic Publishers; 2003.
33. Xue ZC, Gao HY, Zhao SJ. Effects of cadmium on the photosynthetic activity in mature and young leaves of soybean plants. *Environ. Sci. Pollut. Res.* 2014;(21):4656-64.
34. Mlinarić S, Dunić JA, Štolfa I, Cesar V, Lepeduš H. High irradiation and increased temperature induce different strategies for competent photosynthesis in young and mature fig leaves. *S. Afr. J. Bot.* 2016;(103):25-31.
35. Guo Y, Gan SS. Translational researches on leaf senescence for enhancing plant productivity and quality. *J Exp Bot.* 2014;65(14):3901-13.
36. Donohue K. Why ontogeny matters during adaptation: developmental niche construction and pleiotropy across the life cycle in *Arabidopsis thaliana*. *Evolution.* 2014;68:32–47.
37. Kimberley MO, Richardson B. Importance of seasonal growth patterns in modelling interactions between radiata pine and some common weed species, *Can J For Res.* 2004;(34):184-94.
38. Nazarova GG. Effects of seasonal, ontogenetic, and genetic factors on lifespan of male and female progeny of *Arvicola amphibious*. *Front Genet.* 2013;(4):100.
39. Dillen SY, Rood RSB. Ceulemans Growth and physiology. *Genetics and Genomics of Populus.* 2010;8:39-66.
40. Van Goethem D, Potters G, De Smedt S, Gu L, Samson R. Seasonal, diurnal and vertical variation in photosynthetic parameters in *Phyllostachys humilis* bamboo plants. *Photosynth Res.* 2014;120:331–46.
41. Yang HB, An SQ, Sun OJ, Shi ZM, She XS, Sun QY, Liu SR. Seasonal variation and correlation with environmental factors of photosynthesis and water use efficiency of *Juglans regia* and *Ziziphus jujuba*. *J Integr Plant Biol.* 2008;50:210–22.
42. Alvares CA, Stape JL, Sentelhas PC, Gonçalves, JLM, Sparovek, G. Köppen's climate classification map for Brazil. *Meteorologische Zeitschrift.* 2013;22(6):711–28.

43. A. Shevchenko A, Tomas H, Havlis J, Olsen JV, Mann M. In-gel digestion for mass spectrometric characterization of proteins and proteomes. *Nature Protocols*. 2007; 1(6):2856–60.
44. Tyanova S, Temu T, Cox J. The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. *Nat Protocols*. 2016a;(11):2301–19.
45. Paoletti AC, Parmely TJ, Tomomori-Sato C, Sato S, Zhu D, Conaway, RC, et al. Quantitative proteomic analysis of distinct mammalian Mediator complexes using normalized spectral abundance factors. *Proc Natl Acad Sci U S A*. 2006;103(50):18928–33.
46. Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, et al. The Perseus computational platform for comprehensive analysis of (prote)omics data. *Nat Methods*. 2016b;13:731–40.
47. Götz S, Garcia-Gomez JM, Terol J, Williams TD, Nagaraj SH, Nueda MJ, et al. High throughput functional annotation and data mining with the blast2go suite. *Nucl. Acids Res*. 2008;36:3420–35.
48. Emanuelsson O, Nielsen H, von Heijne G. ChloroP, a neural network-based method for predicting chloroplast transit peptides and their cleavage sites. *Protein Sci*. 1999;8(5):978-84.
49. Small I, Peeters N, Legeai F, Lurin C. Predotar: A tool for rapidly screening proteomes for N-terminal targeting sequences. *Proteomics*. 2004 Jun;4(6):1581-90.
50. Sun Q, Zybaïlov B, Majeran W, Friso G, Olinares PD, van Wijk KJ. PPDB, the Plant Proteomics Database at Cornell. *Nucleic Acids Res*. 2009 Jan;37(Database issue):D969-74.
51. Salvi D, Bournais S, Moyet L, Bouchnak I, Kuntz M, Bruley C, Rolland N. AT_CHLORO: The First Step When Looking for Information About Subplastidial Localization of Proteins. *Methods Mol Biol*. 2018;1829:395-406.
52. Mattiello L, Riaño-Pachón DM, Martins MC, Da Cruz LP, Bassi D, Marchiori PE, et al. Physiological and transcriptional analyses of developmental stages along sugarcane leaf. *BMC plant biology*, 2015;15(300).
53. Biswal B, Joshi PN, Raval MK, Biswal UC. Photosynthesis, a global sensor of environmental stress in green plants: Stress signaling and adaptation. *Curr. Sci*. 2011;101:47–56.

- 54.Chan KX, Phua SY, Crisp P, McQuinn R, Pogson BJ. Learning the languages of the chloroplast: retrograde signaling and beyond. *Annu Rev Plant Biol.* 2016;(67):25–53.
- 55.Li P, Ponnala L, Gandotra N, Wang L, Si Y, Tausta SL, et al. The developmental dynamics of the maize leaf transcriptome. *Nat Genet.* 2010;42:1060–67.
- 56.Zrenner R, Riegler H, Marquard CR, Lange PR, Geserick C, Bartosz CE, et al. A functional analysis of the pyrimidine catabolic pathway in *Arabidopsis*. *New phytologist.* 2009;183(1):117–32.
- 57.Levesque-Tremblay G, Havaux M, Ouellet F. The chloroplastic lipocalin AtCHL prevents lipid peroxidation and protects *Arabidopsis* against oxidative stress. *Plant J.* 2009;60(4):691-702.
- 58.Malnoë A, Schultink A, Shahrasbi S, Rumeau D, Havaux M, Niyogi KK. The Plastid Lipocalin LCNP Is Required for Sustained Photoprotective Energy Dissipation in *Arabidopsis*. *Plant cell.* 2018;30(1):196–208.
- 59.Cheng YC, Fleming GR. Dynamics of light harvesting in photosynthesis. *Annu Rev Phys Chem.* 2009;60:241-62.
- 60.Bauerle WL, R. Oren, D. A. Way, S. S. Qian, P. C. Stoy, P. E. Thornton. Pattern of photosynthetic capacity and the implications for carbon cycling. *Proc Natl Acad Sci USA.* 2012;109: 8612–17.
- 61.Medvigy D, Jeong SJ, Clark KL, Skowronski NS, Schäfer KV. Effects of seasonal variation of photosynthetic capacity on the carbon fluxes of a temperate deciduous forest. *Journal of Geophysical Research – Biogeosciences.* 2013;118:1703–14.
- 62.Xu L, Baldocchi DD. Seasonal trends in photosynthetic parameters and stomatal conductance of blue oak (*Quercus douglasii*) under prolonged summer drought and high temperature. *Tree Physiology.* 2003;23:865–77.
- 63.Yang H, Yang X, Zhang Y, Heskell MA, Lu X, Munger JW, Sun S, Tang J. Chlorophyll fluorescence tracks seasonal variations of photosynthesis from leaf to canopy in a temperate forest. *Glob Change Biol.* 2017;23:2874-86.
- 64.Sello S, Meneghesso A, Alboresi A, Baldan B, Morosinotto T. Plant biodiversity and regulation of photosynthesis in the natural environment. *Planta.* 2019;249:1217–28.

65. Chang CY, Unda F, Zubilewich A, Mansfield SD, Ensminger I. Sensitivity of cold acclimation to elevated autumn temperature in field-grown *Pinus strobus* seedlings. *Front. Plant Sci.* 2015;6:165.
66. Baulcombe DC, Dean C. Epigenetic regulation in plant responses to the environment. *Cold Spring Harb Perspect Biol.* 2014;6(9):a019471.
67. Howe GT, Horvath DP, Dharmawardhana P, Priest HD, Mockler TC, Strauss SH. Extensive transcriptome changes during natural onset and release of vegetative bud dormancy in *Populus*. *Front. Plant Sci.* 2015;6:989.
68. Brossa R, Pintó-Marijuan M, Francisco R, López-Carbonell M, Chaves MM, Alegre L. Redox proteomics and physiological responses in *Cistus albidus* shrubs subjected to long-term summer drought followed by recovery. *Planta.* 2015;241(4):803-22.
69. Sathish P, Withana N, Biswas M, Bryant C, Templeton K, Al-Wahb M, et al. Transcriptome analysis reveals season-specific rbcS gene expression profiles in diploid perennial ryegrass (*Lolium perenne* L.). *Plant Biotechnol J.* 2007;5(1):146-61.
70. Burnett AC, Serbin SP, Lamour J, Anderson J, Davidson KJ, Yang D, Rogers A. Seasonal trends in photosynthesis and leaf traits in scarlet oak, *Tree Physiology.* 2021; 41(8):1413–24.
71. Liu J, Last RL. MPH1 is a thylakoid membrane protein involved in protecting photosystem II from photodamage in land plants. *Plant Signal Behav.* 2015;10(10):e1076602.
72. Buhr F, Bakkouri M, Valdez O, Pollmann S, Lebedev N, Reinbothe S, Reinbothe C. Photoprotective role of NADPH: protochlorophyllide oxidoreductase A. *Proc Natl Acad Sci USA.* 2008;105:12629–34.
73. Heyes DJ, Hunter CN. Making light work of enzyme catalysis: protochlorophyllide oxidoreductase. *Trends Biochem Sci.* 2005;(30):642–649.
74. Paddock T, Lima D, Mason ME, Apel K, Armstrong GA. Arabidopsis light-dependent protochlorophyllide oxidoreductase A (PORA) is essential for normal plant growth and development. *Plant Mol Biol.* 2012;78:447–60.
75. Budzinski IGF, Moon DH, Morosini JS, Lindén P, Bragatto J, Moritz T, et al. Integrated analysis of gene expression from carbon

- metabolism, proteome and metabolome, reveals altered primary metabolism in *Eucalyptus grandis* bark, in response to seasonal variation. BMC Plant Biol. 2016;16:149.
- 76.Jungsukcharoen J, Chokchaichamnankit D, Srisomsap C, Cherdshewasart W, Sangvanich P. Proteome analysis of *Pueraria mirifica* tubers collected in different seasons. Biosci Biotechnol Biochem. 2016; 80(6):1070–108.
- 77.Ding L, Hofius D, Hajirezaei MR, Fernie AR, Börnke F, Sonnewald U. Functional analysis of the essential bifunctional tobacco enzyme 3-dehydroquinase dehydratase/shikimate dehydrogenase in transgenic tobacco plants. J Exp Bot. 2007; 58(8):2053-67.
- 78.Tahara K, Nishiguchi M, Funke ES, Miyazawa S, Miyama T, Milkowski C. Dehydroquinase dehydratase/shikimate dehydrogenases involved in gallate biosynthesis of the aluminum-tolerant tree species *Eucalyptus camaldulensis*. Planta. 2021; 253(3).
- 79.Mano J, Torii Y, Hayashi S, Takimoto K, Matsui K, Nakamura K, et al. The NADPH: quinone oxidoreductase P1- ζ -crystallin in *Arabidopsis* catalyzes the α,β -hydrogenation of 2-alkenals: detoxication of the lipid peroxide-derived reactive aldehydes. Plant Cell Physiol.2002; (43):1445–55.
- 80.Yamauchi Y, Hasegawa A, Mizutani M, Sugimoto Y. Chloroplastic NADPH-dependent alkenal/one oxidoreductase contributes to the detoxification of reactive carbonyls produced under oxidative stress. FEBS Letters. 2012;586(8):1208-13.
- 81.Takagi D, Ifuku K, Ikeda K, Inoue KI, Park P, Tamoi M, et al. Suppression of Chloroplastic Alkenal/One Oxidoreductase Represses the Carbon Catabolic Pathway in *Arabidopsis* Leaves during Night. Plant Physiol. 2016;170(4):2024-39.
- 82.Mayta ML, Hajirezaei MR, Carrillo N, Lodeyro AF. Leaf Senescence: The Chloroplast Connection Comes of Age. Plants (Basel). 2019 Nov 12;8(11):495.
- 83.Gan S. Mitotic and post-mitotic senescence in plants. Science's SAGE KE 2003(38), RE7.
- 84.Jibrán R, Hunter D, Dijkwel P. Hormonal regulation of leaf senescence through integration of developmental and stress signals. Plant Molecular Biology. 2013; 82: 547–61.

85. Egli DB. Seed-fill duration and yield of grain crops. In: Advances in Agronomy. San Diego: Elsevier Academic Press Inc. 2004; 83:243–279.
86. Gregersen PL, Culetic A, Boschian L, Krupinska K. Plant senescence and crop productivity. Plant Mol. Biol. 2013;82:603–22.
87. Moschen S, Higgins J, Di Rienzo JA, Heinz RA, Paniego N, Fernández P. Network and biosignature analysis for the integration of transcriptomic and metabolomic data to characterize leaf senescence process in sunflower. BMC Bioinform. 2016;17:174.
88. Woo HR, Koo HJ, Kim J, Jeong H, Yang JO, Lee IH, et al. Programming of Plant Leaf Senescence with Temporal and Inter-Organellar Coordination of Transcriptome in *Arabidopsis*, Plant Physiology. 2016;171(1):452–67.
89. Zhang Q. Strategies for developing Green Super Rice. Proc. Natl. Acad. Sci. USA. 2007;104:16402–09.
90. Villar E, Klopp C, Noirot C. et al. RNA-Seq reveals genotype-specific molecular responses to water deficit in *Eucalyptus*. BMC Genomics. 2011;12: 538.

Supporting information

S1 Fig. *Eucalyptus grandis* sampling. Leaf developmental assay, branches were divided into 3 different regions (young, middle, and mature), according to the fluorescence data (FV/FM) and chlorophyll relative quantification (CCI) (A). Proteome seasonal variation assay, forty leaves were collected from the first until the fifth node in all four seasons (Spring, Summer, Fall and Winter) (B).

S2 Fig. *Eucalyptus grandis* leaf responses. Chlorophyll content index (CCI) (A) and Quantum efficiency of photosystem II (FV/FM) (B). Leaves were isolated from young, middle, and mature regions as described in the Material and Methods section. Different letters indicate significant differences according to Tukey's test ($p < 0.05$).

S1 Table. Meteorological data recorded at the Sao Paulo State University Forest Nursery during *Eucalyptus grandis* cultivation.

S2 Table. *Eucalyptus grandis* non-redundant protein dataset

S3 Table. Expression profile of the differentially regulated proteins identified in the *Eucalyptus grandis* chloroplasts during leaf development. Proteins were classified according to their abundance profile: “up-regulated” - proteins with the highest abundance in mature leaves; “down-regulated” – proteins with the lowest abundance in young leaves; "undefined" – proteins with undefined profile pattern.

S4 Table. Differentially abundant proteins according to the Tukey’s test ($p < 0.05$) in *Eucalyptus grandis* chloroplasts isolated in different seasons of the year classified per protein expression pattern.

Figure 1

Figure 2

Figure 3

A **B**

Figure 4

Figure 5

ID	Accession	Description	Humid seasons		Dry seasons	
			Spring	Summer	Autumn	Winter
1	Eucgr.A00746.1	LTP1				
2	Eucgr.K03160.1	HTA12				
3	Eucgr.A02448.1	ENR1				
4	Eucgr.H04433.2	G-H2AX				
5	Eucgr.B02310.1	RCA				
6	Eucgr.B03013.1	ENR2				
7	Eucgr.C01875.1	ASP5				
8	Eucgr.C02720.2	LHB1				
9	Eucgr.D00322.1	RPS5				
10	Eucgr.D00609.2	SCO1				
11	Eucgr.D02023.1	TKL				
12	Eucgr.D02467.1	PETA				
13	Eucgr.E02606.1	PORA				
14	Eucgr.F01098.1	LHCB4.3				
15	Eucgr.G03060.1	MPH1				
16	Eucgr.G03205.2	AAC2				
17	Eucgr.G03388.1	CLPD				
18	Eucgr.H00243.3	RRF				
19	Eucgr.H01473.3	APTF				
20	Eucgr.H02991.1	MEE32				
21	Eucgr.H04427.1	GAPC2				
22	Eucgr.H04673.1	PRPL29				
23	Eucgr.I00441.1	ZEP				
24	Eucgr.I00922.1	TPI				
25	Eucgr.I01374.1	RPL27				
26	Eucgr.I01643.1	AOR				
27	Eucgr.I01808.1	HSC70-2				
28	Eucgr.I02771.1	HSC70				
29	Eucgr.J00025.1	HCF136				
30	Eucgr.J00865.1	RPL14				
31	Eucgr.J02666.1	ATPB				
32	Eucgr.J02738.1	TPR				
33	Eucgr.K00402.3	PDIL2-2				
34	Eucgr.K00732.1	AHRI				
35	Eucgr.K01105.1	LTP2				
36	Eucgr.K01283.1	ENR3				
37	Eucgr.K01490.1	SFGH				
38	Eucgr.K01713.1	RBCS2				
39	Eucgr.K02223.1	HAD				
40	Eucgr.K02765.1	GCT				
41	Eucgr.L02773.1					

Figure 6