Are tropical hummingbird pollination networks resistant to experimental extirpation of a common flowering plant?

Kara G. Leimberger^{1, *}, Adam S. Hadley^{1,2}, Sarah J.K. Frey^{1,3}, and Matthew G. Betts¹

¹Forest Biodiversity Research Network, Department of Forest Ecosystems and Society, Oregon State University, Corvallis, Oregon, USA

²Biodiversity Section, Department of Natural Resources and Energy Development, Fredericton, New Brunswick, Canada

³Department of Animal and Rangeland Sciences, Oregon State University, Corvallis, Oregon, USA

* Author for correspondence (E-mail: kara.leimberger@oregonstate.edu).

ABSTRACT

Plants and pollinators are experiencing parallel declines worldwide, despite theory predicting that pollination networks should withstand disturbance due to redundancy of pollinators, rarity of interactions between specialists, and flexible pollinator foraging behavior (rewiring). Experiments evaluating the stability of plant-pollinator networks may help resolve this incongruity but remain uncommon. In this study, we simulated the extirpation of a hummingbird-pollinated understory plant, *Heliconia tortuosa*, from tropical forest fragments using a replicated Before-After-Control-Impact experimental design while quantifying hummingbird space use (383 hummingbird captures and 72 radio-tagged individuals), floral visitation rates (6,759 visitations from 20,725 observation hours), and plant pollination success (529 flowers). To complement this experimental approach, we also examined these responses across a natural gradient in *H. tortuosa* density. We expected that declines of *H. tortuosa* would either result in (a) network collapse, in which hummingbirds vacate fragments and compromise the reproductive success of other flowering plants, or (b) increased hummingbird reliance on alternative resources, leading to sustained fragment use. We also hypothesized that landscape and local context (i.e., connectivity to additional forest area and alternative resource abundance) could mediate hummingbird responses to *H. tortuosa* declines; for example, connectivity could facilitate hummingbird visitation to areas of reduced food availability. In our removal experiment, hummingbird persistence and plant pollination success were remarkably resistant to loss of *H. tortuosa*, a locally common plant species representing >40% of the available nectar resources on average. However, naturally low *H. tortuosa* densities were associated with reduced floral visitation rates and decreased pollination. Although landscape context (connectivity) led to higher hummingbird abundance, we found little evidence that connectivity or resource availability mediated hummingbird responses to declines in *H. tortuosa* availability. The exact

mechanisms enabling short-term hummingbird persistence after resource removal remain unclear, as we did not discover evidence of rewiring. Physiological adaptations (e.g., torpor and insectivory), may have contributed to hummingbird resilience, perhaps alongside high movement abilities that facilitated visitation to spatially dispersed floral resources. With the important caution that short-term experiments may not realistically emulate natural extinction processes, our study provides partial support for theoretical predictions that pollination networks may be relatively robust to plant species loss.

Key words: mutualistic networks, ecosystem function, foraging ecology, functional connectivity, fragmentation, Trochilidae

INTRODUCTION

Species loss can have cascading consequences for the broader community, including coextinctions of dependent species and declines in ecosystem function (Colwell, Dunn, and Harris 2012; Koh et al. 2004). Linked extinctions may be especially likely for mutualistic interactions, such as those between plants and pollinators (Dunn et al. 2009; Aslan et al. 2013). For instance, plants that are not adequately pollinated produce fewer seeds, accruing extinction debt that becomes realized when the existing generation cannot be replaced by new recruits (Ashman et al. 2004; Vellend et al. 2006; Kuussaari et al. 2009). This repeated reproductive failure will eventually cause demographic collapse of plant populations (Anderson et al. 2011; Phillips et al. 2015), which could then hasten the decline of animals that consume pollen or nectar (Biesmeijer et al. 2006; Pauw and Hawkins 2011; Weiner et al. 2014; Scheper et al. 2014). Indeed, plants and pollinators are experiencing declines worldwide (Biesmeijer et al. 2006; Potts et al. 2010; Regan et al. 2015), which not only imperils the essential ecosystem services on

which humans depend but also threatens the maintenance of global biodiversity (Allen-Wardell et al. 1998; Klein et al. 2007; Ollerton, Winfree, and Tarrant 2011).

Although the underlying reasons for plant and pollinator declines are not fully understood, one prominent hypothesis invokes the landscape-level processes of habitat loss and fragmentation (Kearns, Inouye, and Waser 1998; Aguilar et al. 2006; Potts et al. 2010; Hadley and Betts 2012). Habitat destruction often creates small habitat patches unable to sustain large populations, which can increase rates of local extinction (Hanski 1998), reduce functional connectivity (sensu P. D. Taylor et al. 1993), and impede pollinator movement (Hadley and Betts 2012). Likely due to these combined mechanisms, land-use change and habitat isolation have been repeatedly linked to declines in pollinator diversity (Winfree, Bartomeus, and Cariveau 2011; Montero-Castaño and Vilà 2012; Senapathi et al. 2015) and plant reproduction (Steffan-Dewenter and Tscharrntke 1999; Taki, Kevan, and Ascher 2007; Farwig et al. 2009; Garibaldi et al. 2011).

However, parallel declines of pollinators and plants are at odds with theoretical predictions suggesting that species loss should generally not result in coextinction of interaction partners (Memmott, Waser, and Price 2004; Bascompte, Jordano, and Olesen 2006). Tolerance to subsequent extinctions (i.e., robustness) primarily arises from the expectation that interaction networks exhibit strongly nested topologies, such that specialists interact with subsets of partners linked to generalists (Bascompte et al. 2003). Under this interaction pattern, connections between specialists are uncommon, so the functional role of a specialist can be absorbed by its more abundant generalist partners (Bascompte et al. 2003; Bascompte and Jordano 2007). In addition to the redundancy afforded by a nested structure, models suggest that flexibility in pollinator foraging behavior – i.e., rewiring to use alternative resources – may buffer networks against

disturbance (Kaiser-Bunbury et al. 2010; Ramos-Jiliberto et al. 2012; Valdovinos et al. 2013; Timóteo et al. 2016).

Despite these overall predictions and mechanisms of robustness, pollination networks may be vulnerable to habitat loss and fragmentation under certain conditions. Although systematic loss of generalist pollinators may be less realistic than extinctions of rare, specialist pollinators (Aizen, Sabatino, and Tylianakis 2012), simulation studies have revealed that removal of highly connected, generalist species thwarts a network’s capacity to buffer perturbations, leading to rapid partner extinction (Memmott, Waser, and Price 2004; Kaiser-Bunbury et al. 2010; Traveset, Tur, and Eguíluz 2017). Thus, if external stressors such as habitat fragmentation threaten highly connected species (‘topological keystones’ sensu Jordán 2009), an extinction cascade could occur – particularly without rewiring by pollinators. However, uncertainty about the robustness of real-world plant-pollinator networks still exists because relatively few studies have quantified pollinator responses to deletions of other species (Appendix S1: Table S1).

In this study, we tested whether declines of a hummingbird-pollinated plant (*Heliconia tortuosa*) could have cascading consequences for communities of hummingbirds and flowering plants within a fragmented tropical forest landscape – or whether rewiring might forestall network collapse. In Neotropical forests, the genus *Heliconia* (Heliconiaceae) comprises understory herbs that have been suggested to function as “keystone mutualists” (Gilbert 1980). Large, brightly colored *Heliconia* inflorescences not only provide nectar to large-bodied hermit hummingbirds (Trochilidae: Phaethornithinae) throughout the year, but also support a variety of other hummingbird species during periods of resource scarcity (Stiles 1975; 1985). Because most hummingbird species visit multiple plant species (Rodríguez-Flores et al. 2019), loss of *H.*

tortuosa and its associated hummingbird partners could jeopardize the reproduction of co-occurring plants, thus generating an extinction cascade (Gilbert 1980). Although *H. tortuosa* is locally common in the mid-elevation, premontane tropical forests of Costa Rica, Hadley *et al.* (2014) found that *H. tortuosa* in smaller forest fragments produced fewer seeds, possibly due to short-distance pollen transfer associated with pollinator movement limitation. Although numerous hummingbird species visit *H. tortuosa* opportunistically, its primary pollinators are the green hermit (*Phaethornis guy*) and violet sabrewing (*Campylopterus hemileucurus*) (Stiles 1979; Wolf and Stiles 1989; J. Taylor and White 2007; Betts, Hadley, and Kress 2015). These relatively large, forest-associated hummingbirds are high-reward trapliners (*sensu* Feinsinger and Colwell 1978) that visit spatially dispersed, nectar-rich floral resources in a routine sequence (Borgella, Snow, and Gavin 2001). However, their movements become restricted in fragmented habitats (Hadley and Betts 2009; Volpe *et al.* 2016; Kormann *et al.* 2016).

To assess the potential consequences of *H. tortuosa* declines, we measured hummingbird space use, floral visitation rates, and plant pollination success before and after temporary experimental removal of *H. tortuosa* within forest fragments. Additionally, we assessed how these responses varied across a natural gradient in *Heliconia* density. We developed two alternative hypotheses about how hummingbirds would respond to declines of *H. tortuosa* (hereafter *Heliconia*) and potentially affect the pollination of co-occurring plant species (Table 1). Under the *robust network hypothesis* (**H1a**), hummingbirds do not emigrate from forest fragments following *Heliconia* decline, leading to uninterrupted visitation to remaining plants. Under this scenario, we hypothesized that hummingbirds compensate for *Heliconia* extinction by increasing their use of non-*Heliconia* floral resources (rewiring). Alternatively, hummingbirds may be highly dependent on *H. tortuosa* and inflexible in their foraging choices; therefore, local

Heliconia declines should reduce nectar availability such that hummingbirds cannot afford the energetic costs to remain in (or travel to) focal forest fragments, leading to decreases in fragment use and pollination success (**H1b**: *parallel extinction hypothesis*). Under both hypotheses, we expected hummingbirds to change their behavior due to their exceptionally rapid metabolisms, high dependency on nectar, and limited fat stores (Suarez 1992; Powers, Brown, and Van Hook 2003). Additionally, we hypothesized that landscape and local context could mediate the influence of *Heliconia* declines on hummingbird behavior (Table 1). Connectivity to additional forest area (*connectivity hypothesis*, **H2**) may sustain hummingbird fragment use by facilitating travel to fragments (*rescue hypothesis*, **H2a**) or promote steeper declines in fragment use by facilitating hummingbird departure (*abandonment hypothesis*, **H2b**). Further, greater local availability of non-*Heliconia* floral resources may promote hummingbird persistence through increased opportunities for rewiring (*resource hypothesis*, **H3**). Lastly, we expected that changes in *Heliconia* density would affect green hermits and violet sabrewings more strongly than other hummingbird species (**H4**). Green hermits and violet sabrewings (hereafter ‘*Heliconia* specialists’) inhabit the forest understory, have bills that are morphologically specialized to long, curved *H. tortuosa* flowers, and are thought to visit *H. tortuosa* more frequently than other hummingbirds (Borgella, Snow, and Gavin 2001; Betts, Hadley, and Kress 2015).

To our knowledge, this study not only constitutes one of most highly replicated, large-scale removal experiments conducted within the context of plant-pollinator interaction networks (Appendix S1: Table S1), but also represents the first test of how landscape and local context influence hummingbird responses to declines in a common floral resource. Importantly, understanding mechanisms of pollinator persistence in fragmented landscapes has implications for conservation planning, since fragmentation effects may be ameliorated by habitat restoration,

preservation of riparian corridors, and/or implementation of living fences (Renjifo 2001; Kormann et al. 2016).

METHODS

Study area

This study was conducted in southern Costa Rica within a ~122 km² area surrounding the Las Cruces Biological Station (Coto Brus Canton, Puntarenas Province, 8°47'7'' N, 82°57'32'' W, Fig. 1: A-B). This study area comprises cattle pastures, coffee plantations, and fragments of tropical premontane wet forest created during a period of extensive land clearing concentrated between 1960 and 1980 (Zahawi, Duran, and Kormann 2015). Remnant forest fragments (~30% of the study area) span a wide range of sizes (<1 to >1000 ha), with otherwise isolated fragments often connected by narrow riparian corridors and living fences (Hadley and Betts 2009; Zahawi, Duran, and Kormann 2015). Mean annual temperature at the Las Cruces Biological Station is ~21 °C, mean annual rainfall is 3500-4000 mm, and a dry season occurs from December to March (Borgella, Snow, and Gavin 2001; Zahawi, Duran, and Kormann 2015).

Study sites

We selected 14 focal fragments embedded within focal landscapes of 1000 m radius ('sites', Fig. 1: A, C). We chose the spatial extent of focal landscapes *a priori* based on the maximum estimated daily movement distance for large, forest-associated hummingbird species in this study system (Volpe et al. 2014). We identified potential sites using a stratified design with two categories: 'connected to forest' or 'unconnected to forest', based on visual examination of satellite imagery and previous work in this study area (Hadley et al. 2014; 2018; Kormann et al. 2016). This design was only for site establishment, as we later quantified connectivity as a continuous variable (see '*Quantifying connectivity*' below). Sites were then

included based on presence of flowering *Heliconia*, landowner permission, vehicle accessibility, and navigable terrain. To ensure site accessibility, we conducted our study during the dry season and beginning of the rainy season (February – May), because localized flooding can occur during periods of heavy rainfall. Elevation of sites ranged from ~900-1500 m (Appendix S2: Table S1).

Study design

Using the 14 focal-patch landscapes, we repeated the *Heliconia* removal experiment 16 times across three years (2016-2018). Each iteration of the experiment followed a BACI (Before-After-Control-Impact) design and used two sites paired based on elevation and connectivity (Appendix S2: Table S2). One of the paired sites ('treatment site') received the treatment of *Heliconia* removal, and we designated the other site as a control ($N = 16$ control replicates, $N = 16$ treatment replicates). Most sites were used multiple times across the three-year study period, but only once per year (Appendix S2: Tables S2-S3). Initial treatment assignments were random, and sites used multiple times received the opposite treatment of the previous year (except in 3 of 16 instances); this design reduces the likelihood that treatment-control differences were due to factors associated with individual sites. To prevent hummingbirds from the treatment site relocating to the control site, paired sites were located at least 1000 m apart (mean $\pm$ SD: 2266 $\pm$ 1511 m, median: 3086 m).

Each experimental iteration lasted nine days and began with a period of pre-treatment data collection in each site (Fig. 2A). Starting around noon on the fourth day, we removed *Heliconia* from the treatment site and designated the following 36 hours as a behavioral adjustment period during which no data were collected. Although this adjustment period was necessarily relatively short due to the limited battery lifespan of hummingbird radio-transmitters (see 'Radio telemetry'), we expected hummingbirds to respond rapidly due to their exceptionally

high metabolisms and limited fat stores (Suarez 1992; Powers, Brown, and Van Hook 2003). To ensure that *Heliconia* inflorescences were not overlooked during removal, field technicians positioned themselves ~2-3 meters apart and slowly walked through the forest while covering all *Heliconia* inflorescences (mature and immature) with dark-colored fabric bags (Fig. 2B). These bags prevented hummingbird access, obscured flowers from view, and allowed us to avoid long-term changes to floral resources associated with destructive sampling methods (e.g., cutting). The location of each *Heliconia* plant was recorded using a handheld GPS unit, and we retrieved the covering bags at the end of the experimental period. In each site, we also recorded the location and identity of all non-*Heliconia* floral resources available to hummingbirds (see ‘*Quantifying resource availability*’). The areas where we removed *Heliconia* and/or surveyed for floral resources are referred to as ‘focal areas’ (Fig. 1D), which we delineated using 100% minimum convex polygons. Focal areas also encompass locations where we captured hummingbirds, installed trail cameras, and set up plant arrays (Fig. 1D, see ‘*Response variables*’).

Response variables

Hummingbird captures

Within each focal area, we captured hummingbirds at the beginning and end of the experimental period (Fig. 2A) by deploying 5 - 20 mist nets (12-m and 6-m in length) near flowering plants visited by hummingbirds (e.g., the genera *Heliconia*, *Centropogon*, *Palicourea*, *Musa*). Nets were opened 30 minutes after sunrise and remained open for 2 - 5 hours per session. Capture effort (net-hours) and net locations were kept constant for both capture sessions, except in one instance (Appendix S2: Figure S1). Following the North American Banding Council’s hummingbird protocol (Russell and Russell 2001), we tagged hummingbirds with an aluminum

tarsus band, measured body mass to the nearest 0.01 g using an electronic balance, and measured wing length to the nearest 0.5 mm using a wing rule. Tarsus bands allowed detection of recaptured hummingbirds within and between capture sessions; individuals recaptured during the same capture session were only counted once. In 2017-2018, we also marked hummingbirds' heads with unique color combinations of nail polish following Hadley *et al.* (2014); this additional mark allowed us to identify individual hummingbirds seen on trail cameras (see '*Camera observations*'). All birds were handled in accordance with Oregon State University Animal Care and Use Protocols (ACUP #4655, #5015).

Radio telemetry

We attached miniaturized radio transmitters (0.15-0.25g, Blackburn Transmitters, Nacogdoches, TX) to 72 hummingbirds according to the methods of Hadley & Betts (2009). We then followed hummingbirds on foot using portable receivers (Wildlife Materials Inc TRX 1000S) and handheld Yagi antennae, recorded locations continuously when the bird was found within receiver range (~250 m in the absence of topographical barriers). In total, we completed 447 hours of telemetry observation, averaging 3.1 ± 0.8 hours per session (typically ~8:00-11:00) and 5.3 ± 0.9 sessions per site (mean $\pm$ SD). From the recorded locations, we estimated the proportion of observation time that radio-tagged birds spent in the focal area (Appendix S2). Thirty-six hummingbirds spent time in the focal area during the 'pre' period and were included in analyses of the *Heliconia* removal experiment; for a comprehensive summary of radio-tracking outcomes, see Appendix S2 (Table S4).

Floral visitation rates

To understand hummingbird resource use within each site and to monitor for the presence of marked hummingbirds, we positioned trail cameras (PlotWatcher Pro, Day 6 Outdoors) near

inflorescences of plant species known or suspected to be visited by hummingbirds. Effort within each site was concentrated at two locations ('stations', Fig. 1D) located on roughly opposite ends of the focal area (mean $\pm$ SD distance between stations: 60 ± 44 m). Stations were centered around a focal *Heliconia* plant that we expected to flower throughout the experimental period, was not in an extremely steep area, and was >10 m from the fragment edge. The two focal *Heliconia* plants remained uncovered throughout the entire experimental period to assess how visitation rates changed after declines in surrounding *Heliconia* density. To partially standardize plant species composition between control and treatment replicates, we added a floral array comprising ~3-5 additional plant species at each station (see Appendix S2: Table S5 and 'Pollination success').

Trail cameras took one photograph per second during daylight hours (5:30-17:30) and automatically combined these photographs into a time-lapse video. We reviewed videos for hummingbird sightings using the motion detection program MotionMeerkat (Weinstein 2015) or by manually watching the sped-up video. When hummingbirds were detected on camera, we identified the species, recorded any color marks, and noted whether they visited any flowers. From video, we also recorded the number of open flowers available per day. Dates with no available flowers were excluded from analysis, yielding 20,735 total hours of video (Appendix S2: Table S6).

Pollination success

To investigate whether *Heliconia* declines influenced pollination success, we examined styles for the presence of pollen tubes, which indicate pollen grain germination and subsequent growth toward the flower ovule(s) (Kearns and Inouye 1993). We focused our pollen tube investigation on the two focal *Heliconia* plants per replicate, as well as several species from the

focal arrays (Appendix S2: Table S5). Throughout the experimental period, we collected flowers from these focal plants every 1-2 days, which corresponds to the maximum floral longevity for most of the selected species (*Heliconia tortuosa*, *Hamelia patens*: Stratton 1989; *Stachytarpheta frantzii*: Thomas-Granger 2003). Styles were fixed in formalin-acetic acid-alcohol for at least 24 hours, stained with aniline blue, and examined for pollen tubes using epifluorescence microscopy (Kearns and Inouye 1993; Betts, Hadley, and Kress 2015). Due to low pollen tube presence and/or challenges with pollen tube visualization, we were only able to analyze two plant species: *H. tortuosa* (N = 327 styles) and *H. patens* (N = 202 styles). Additional details about style collection and lab procedures are available in Appendix S2.

Additional response variables (Heliconia removal experiment only)

In addition to the primary response variables described above, we examined several other metrics for the *Heliconia* removal experiment: (1) recapture probability (i.e., proportion of birds captured during the ‘pre’ period and recaptured during the ‘post’ period), (2) floral visitation rates from individual birds marked with nail polish, and (3) hummingbird body mass. The first two response variables assessed the possibility that hummingbirds caught during the initial capture sessions were rapidly replaced by new individuals, potentially obscuring treatment effects. To explore whether birds that remained in (or returned to) treatment focal areas were paying an energetic cost, we also analyzed changes in the body mass of hummingbirds captured during both experimental periods. To compare body mass change across hummingbirds of different structural sizes, we calculated body mass relative to the mass predicted by species-specific allometric equations (i.e., $\ln(\text{body mass}) \sim \ln(\text{wing length})$, Appendix S2: Fig. S4).

Quantifying connectivity

Within each focal landscape, we used Google Earth Pro to digitize closed-canopy forest and regenerating forest (hereafter ‘forest’) from high resolution (1 m) Airbus/CNES satellite imagery. These two forest types were combined because functional connectivity experiments suggest that regenerating forest is nearly equivalent to closed-canopy forest in facilitating hummingbird movement (Jackson 2017). We then delineated connected areas of forest (‘patches’) assuming a hummingbird gap-crossing ability of 50 m (Volpe et al. 2016; Kormann et al. 2016). Lastly, we quantified functional connectivity as the amount of reachable forest within each focal landscape, following a modified version of Hanski’s (1994) incidence function model for metapopulation connectivity. Substituting forested pixels for patches used in the original model, we assigned exponentially lower importance to pixels located farther away from the focal area (Fig. 1E, Appendix S2: Fig. S5). These weights were calculated as:

$$W_j = \exp(-\alpha d_{ij}), \quad (\text{Eq. 1})$$

where α is the decay rate parameter that scales the effect of distance and d is the shortest cumulative distance between the focal area centroid i and each forested pixel j , staying within the patch boundary. The decay rate parameter is inversely related to the average movement distance of the focal species (Moilanen and Nieminen 2002) and was calculated using the mean home range length of green hermit hummingbirds (282 m: Volpe et al. 2016). Then, to calculate functional connectivity, we then summed the weighted pixel areas as follows:

$$S_i = \sum W_j A_j, \quad (\text{Eq. 2})$$

where W_j is the weight of each forested pixel and A is the area of each forested pixel. Geospatial processing was conducted using Google Earth Engine through the JavaScript API (Gorelick et al. 2017).

Quantifying resource availability

From the floral resource surveys (see ‘*Study design*’), we estimated the energetic value of the *Heliconia* flowers and the remaining non-*Heliconia* resources. Briefly, we converted resource counts (i.e., number flowers and inflorescences) into calories, the most biologically relevant unit for hummingbird energetics (e.g., Wolf, Hainsworth, and Stiles 1972; Stiles 1975). To calculate calories, we measured nectar volume and concentration from >1400 flowers ($N = 38$ species) growing in the focal landscape and/or the Wilson Botanical Garden at the Las Cruces Biological Station, supplementing these measurements with literature data when necessary. We then converted average nectar volume and concentration per flower to calories per plant (see Appendix S2 for full details). Because tropical hummingbirds generally interact with subsets of the available floral resources (e.g., Stiles 1985; Maglianesi et al. 2014; Vizentin-Bugoni, Maruyama, and Sazima 2014), we then generated estimates of caloric availability specific to each hummingbird group examined. These estimates were developed prior to any analysis based on relative visitation rates of each hummingbird species to 37 flowering plant species (Fig. 3). We divided these calorie estimates by focal area size, then natural log-transformed the resulting densities. Floral corolla length was measured from flowers collected in the field, supplemented with literature values when necessary (Appendix S2: Table S16).

Statistical analysis

We used linear mixed models (LMMs) and generalized linear mixed models (GLMMs) to analyze how experimental *Heliconia* extinction and natural variation in *Heliconia* density influenced our primary response variables: (i) captured hummingbird abundance, (ii) proportion of time spent in focal area, (iii) visitation rates to flowering plants (*Heliconia* and non-*Heliconia*), and (iv) plant pollination success. Analyses of the *Heliconia* removal experiment

also included several additional, individual-level response variables (see ‘*Additional response variables*’). To determine whether declines in *Heliconia* affected *Heliconia* specialists more strongly than other hummingbird species (**H4**), we analyzed all hummingbird responses with (i) the full dataset containing all species, and (ii) a dataset including green hermits and violet sabrewings.

We modeled each response variable using appropriate statistical distributions and controlled for variable effort when applicable (Appendix S2: Tables S17-S18). All statistical analyses were conducted in R version 4.1.1 (R Core Team 2021) using ‘glmmTMB’ (Brooks et al. 2017). For LMMs, we checked for normality, homoscedasticity, and linearity using the ‘performance’ package (Lüdtke et al. 2021). For GLMMs, we checked for overdispersion, zero-inflation, and linearity by visually inspecting residual plots from ‘DHARMa’ (Hartig 2020). Estimated marginal means were calculated using ‘emmeans’ (Lenth 2020) and visualized using the ‘ggemmeans’ function within ‘ggeffects’ (Lüdtke 2018). Results were considered statistically significant if $P < 0.05$.

Heliconia removal experiment

In a BACI experimental design, an overall treatment effect is indicated by a statistically significant interaction between the variables representing treatment and experimental period (Underwood 1994; McDonald, Erickson, and McDonald 2000). We adopted this standard analytical approach rather than calculating change scores (e.g., pre-post differences) because change scores are nearly always negatively correlated with baseline measurements; when baseline measurements also covary with predictor variables of interest, misleading (i.e., spuriously negative) relationships can emerge (Allison 1990). Therefore, we tested the *parallel*

extinction hypothesis (H1a) and *robust network hypothesis (H1b)* using a two-way interaction between *Heliconia* removal and period (pre *versus* post) for all response variables measured during both experimental periods (recaptures were measured in the post period only). All models included a random intercept for ‘Site’. Except when modeling recapture probability, we also included a nested random effect for ‘Replicate ID’ (Site-Year), which paired pre-post observations within each replicate. Additional nested random effects, namely for plant species and individual birds, were included where applicable (Appendix S2: Table S17). Because we were interested in pre-post changes in our response variables, we only included covariates that varied between pre and post periods. For example, when analyzing hummingbird visitation rates, we accounted for the attractiveness of a given focal plant by including a covariate for the number of open flowers per day, averaged across the experimental period. Since the magnitude of our experimental *Heliconia* extinction varied widely (see ‘Results’), we ran parallel models using a categorical treatment variable (Control/Treatment) and a quantitative treatment variable (Calories removed/hectare). Because results were qualitatively similar, only results from the categorical approach are presented in the main text.

Natural variation in Heliconia density

To complement the *Heliconia* removal experiment, we also examined hummingbird responses over a natural gradient in *Heliconia* availability using data from the pre period of the experimental sites. This ‘space-for-time’ approach elucidates how hummingbird space use and foraging behavior respond to natural variation present in the study area – presumably emerging over longer time scales than those of our removal experiment. We also used this approach to test the connectivity hypothesis and resource hypothesis by modeling the interaction term between

Heliconia density and either connectivity (**H2**) or non-*Heliconia* resource availability (**H3**). These hypotheses were not tested using the experimental framework due to limited sample size (and thus statistical power) to enable estimation of a three-way interaction between treatment, experimental period, and resources/connectivity. Because resource availability data were not collected in 2016 control replicates, these replicates ($N = 6$) were excluded from this analysis. In these analyses, we included several additional covariates not included in the analyses of the *Heliconia* removal experiment. First, we used estimates of non-*Heliconia* resource availability tailored to each hummingbird group being examined. Because resource availability was moderately correlated with connectivity (Pearson's $r = 0.57$, $P = 0.002$, Appendix S2: Figure S6), we included both covariates in all models to statistically control for their independent effects. Analyses of radio telemetry data (i.e., proportion of time spent in focal area) included a covariate for the focal area size. When analyzing visitation rates of *Heliconia* specialists to non-*Heliconia* resources, we also included plant corolla length as a covariate, since these relatively long-billed hummingbirds (mean bill lengths of 40 and 33 mm: Borgella, Snow, and Gavin 2001) are expected to select plant species with long corolla tubes (e.g., Maglianesi et al. 2014; Vizentin-Bugoni, Maruyama, and Sazima 2014; Dalsgaard et al. 2021). We included random effects for Site in all models; additional random effects were included where applicable (Appendix S2: Table S18).

RESULTS

Across all study sites and years, *Heliconia* density ranged from 3 to 1131 inflorescences per ha (mean $\pm$ SD: 186 ± 228 , median: 145, Appendix S3: Table S1). The magnitude of our experiment therefore varied widely; we removed between 8 and 526 *Heliconia* inflorescences per treatment replicate (mean $\pm$ SD: 172 ± 138 , median: 138, Appendix S3: Table S2). These

inflorescences represented, on average, 96% ($\pm 5.5\%$) of the *Heliconia* inflorescences encountered during removal (two focal plants remained uncovered and sometimes plant inaccessibility limited our capacity to cover). Sizes of focal treatment areas ranged from 0.27 to 7.49 ha (mean $\pm$ SD: 1.7 ± 1.9 , median: 1.25, Appendix S3: Table S2), depending on the size of the forest fragment and the navigability of the terrain. We estimate that, on average, the covered *Heliconia* plants supplied 42.5% ($\pm 30.9\%$, range: 0.9 – 90.4%) of the calories available for the entire hummingbird community ($N = 19$ species, Appendix S3: Table S3) and at least 65.5% ($\pm 31.3\%$, range: 2.4-98%) of calories for *Heliconia* specialists (Fig. 2C, Appendix S3: Table S4). Focal *Heliconia* plants were visited by hummingbirds 6.4 ± 5.7 times per day (range: 0-25, median: 5), with green hermits visiting most frequently (75% of 1,065 sightings).

Overall responses to changes in Heliconia density (H1)

Hummingbird behavior and pollination success were remarkably resistant to experimentally induced *Heliconia* extinction. Despite the magnitude of *Heliconia* removal and confirmed use of *Heliconia* by hummingbirds, we did not detect an overall effect of local *Heliconia* removal hummingbird space use, floral visitation rates, or plant pollination success. However, when examining these response variables across a natural gradient in *Heliconia* availability (i.e., the space-for-time approach), we found that floral visitation rates and pollination success declined as *Heliconia* became scarce. We provide detailed evidence for these findings below.

Hummingbird captures & radio telemetry

Overall, we did not find that experimental *Heliconia* removal decreased hummingbird abundance or the proportion of time that radio-tagged hummingbirds spent in the focal area. Although we caught 23% fewer hummingbirds in the ‘post’ capture session than the ‘pre’

capture session in treatment replicates (95% CI: -41% to 1%), this pre-post decline fell within the variability of the control replicates (all species: $z = -1.11$, $P = 0.27$; *Heliconia* specialists: $z = -0.5$, $P = 0.62$; Fig. 4: A-B; Appendix S3: Table S5). Similarly, although radio-tagged hummingbirds spent 25-30% less time in focal areas during the ‘post’ period, we found no evidence that pre-post changes in space use were linked to our experimental manipulation (95% CI, treatment: -56% to 11%; 95% CI, control: -56% to 28%). Results were similar when examining all species together (Fig. 4C) and *Heliconia* specialists separately (Fig. 4D). We also found no evidence that natural variation in *Heliconia* density influenced the hummingbird space use, quantified either with mist net captures or radio telemetry (Appendix S3: Tables S5-S6).

When examining individual-level hummingbird responses, we also did not detect changes associated with *Heliconia* removal. On average, the estimated probability of recapturing a hummingbird from the ‘pre’ capture session was 18% in control replicates (95% CI: 11% to 29%), compared to 14% in treatment replicates (95% CI: 9% to 23%). This difference was not statistically significant, and recaptures of *Heliconia* specialists showed qualitatively similar results (Appendix S3: Fig. S1: A-B, Table S7). Among the individual hummingbirds caught during both sessions, we also found no evidence that pre-post changes in the relative body mass differed between control and treatment replicates (Appendix S3: Table S8).

Floral visitation rates & pollination success

Using trail cameras, we continuously monitored hummingbird visitation at the two focal *Heliconia* inflorescences that remained uncovered throughout the experimental period, as well as numerous other flowering plants (mean $\pm$ SD: 9.3 ± 2 species/replicate; range: 6-14). Under the *robust network hypothesis*, we expected hummingbirds to rewire and increasingly use alternative (non-*Heliconia*) resources. However, we did not find that pre-post changes in hummingbird

visitation rate differed between control and treatment replicates, either when examining the focal *Heliconia* plants ($z = 0.83$, $P = 0.41$, Fig. 5: A-B, Appendix S3: Table S9) or non-*Heliconia* plant species ($z = -0.69$, $P = 0.49$, Fig. 6: A-B, Appendix S3: Table S10). Responses by *Heliconia* specialists mirrored the responses observed for all species, although these species visited non-*Heliconia* species relatively infrequently (<1 visit/day, on average: Fig. 6B). Results were similar when examining pre-post changes in visitation rates for individual color-marked hummingbirds observed on camera (Appendix S3: Fig. S1: C-F, Tables S11-S12).

In contrast to the experimental results, we found that natural variation in *Heliconia* availability strongly influenced visitation to focal *Heliconia* plants (Fig. 5: D-E). For every standard deviation decrease in the caloric density of *Heliconia*, we found that overall *Heliconia* visitation rates decreased by 49% (95% CI: 15% to 93%, $z = 3.03$, $P = 0.0024$, Appendix S3: Table S9). Visitation to non-*Heliconia* plants also declined when natural *Heliconia* availability was low, but only for *Heliconia* specialists ($z = 2.02$, $P = 0.044$, Fig. 6E, Appendix S3: Table S10).

We found no evidence that experimental *Heliconia* removal influenced pollination success for the focal *Heliconia* plants ($z = 0.23$, $P = 0.82$, Fig. 5C, Appendix S3: Table S13) or another plant species from the floral arrays (*Hamelia patens*: $z = 0.19$, $P = 0.85$, Appendix S3: Table S13). However, when *Heliconia* availability was naturally low, we observed declines in the pollination success of the focal *Heliconia* plants; for every standard deviation decrease in the caloric density of *Heliconia*, the odds of pollination declined by 2.4 times (95% CI: 1.1 times to 5.4 times, $z = 2.2$, $P = 0.028$, Fig. 5F).

Effects of connectivity (H2) and resource availability (H3)

Hummingbird captures & radio telemetry

Connectivity influenced hummingbird abundance; hummingbird captures increased by 6% for every one-unit increase in connectivity (95% CI: +1% to +12%, $z = 2.31$, $P = 0.021$, Appendix S3: Table S5). Further, radio-tagged birds spent more time in focal areas when alternative (non-*Heliconia*) resources were abundant; the odds of being in the focal area increased by 2.3 times (95% CI: 1.3 to 4.1 times) for every standard deviation increase in caloric density (all species only: $z = 2.78$, $P = 0.0055$, Appendix S3: Table S6). However, neither connectivity or resource availability appeared to influence hummingbird responses to *Heliconia* declines, either with experimental removal or across the natural density gradient ($P > 0.05$ for all interactions of interest, Appendix S3: Tables S5-S6).

Floral visitation rates & pollination success

We did not observe any statistically significant main effects of connectivity on hummingbird visitation rates, either to focal *Heliconia* plants or non-*Heliconia* species. However, after *Heliconia* removal, visitation rates to focal *Heliconia* plants by *Heliconia* specialists declined more sharply when connectivity was low ($z = -1.76$, $P = 0.08$, Appendix S3: Fig. S2, Table S9). This potential connectivity effect did not emerge when examining *Heliconia* pollination success ($z = -0.55$, $P = 0.58$). Although we expected that interaction rewiring might be more likely to occur when alternative resources were abundant, we did not find that *Heliconia* declines in otherwise resource-rich areas caused hummingbirds to rely increasingly on non-*Heliconia* species (Appendix S3: Table S10).

DISCUSSION

In this study, we experimentally induced the local extinction of a locally abundant tropical plant, *Heliconia tortuosa*, to assess the robustness of plant-hummingbird interaction

networks in a fragmented tropical landscape. Additionally, across a natural gradient in *Heliconia* density, we assessed whether low *Heliconia* abundance would lead to parallel declines in hummingbirds and their pollination services – and whether landscape context (connectivity) and local context (alternative resource availability) could buffer against potential declines. Based on suite of complementary response variables we found that, even when very high proportions of total available calories were experimentally removed (~43-66% on average), hummingbird responses fell within the natural variability of control replicates. As might be expected with such small changes in hummingbird behavior, we did not observe any cascading effects of experimental manipulation on pollination success of flowering plants, which supports the *robust network hypothesis* (**H1b**). On the other hand, when examining hummingbird responses and pollination success across a natural gradient in *Heliconia* density, we found evidence for the *parallel extinction hypothesis* (**H1a**). In sites where *Heliconia* density was low, hummingbirds visited individual *Heliconia* plants less frequently, and these plants experienced reduced pollination success. Other flowering plants also exhibited lower visitation rates from *Heliconia*-associated hummingbirds, suggesting that high *Heliconia* densities may attract these hummingbirds (and/or incentivize repeat visitation) and therefore facilitate pollination for other species.

Our finding that that experimental removal of *Heliconia* did not elicit parallel extinctions in pollinators aligns with theoretical predictions (Memmott, Waser, and Price 2004; Kaiser-Bunbury et al. 2010) and several other experiments that removed plant species from mutualistic networks (Ferrero et al. 2013; Goldstein and Zych 2016; Costa et al. 2018). However, if hummingbirds remained in forest fragments despite *Heliconia* declines, we assumed that they would rewire to use alternative floral resources; hummingbirds as a taxonomic group are highly

opportunistic foragers (Arizmendi and Ornelas 1990; Maruyama et al. 2013) and the most frequent *Heliconia* visitor, the green hermit, has a relatively wide niche breadth (Fig. 3, see also Borgella, Snow, and Gavin 2001; Betts, Hadley, and Kress 2015). Unexpectedly, we found no evidence that hummingbirds increasingly visited non-*Heliconia* resources when *Heliconia* became less available, suggesting that the mechanisms underlying network robustness in this study may not necessarily align with the rewiring observed in previous empirical tests (e.g., Timóteo et al. 2016; Kaiser-Bunbury et al. 2017; Costa et al. 2018). This apparent lack of foraging flexibility may emerge from various constraints, such as morphological trait-matching, search costs, and/or competition from other hummingbirds (see also Weinstein and Graham 2017). For example, high levels of exploitative competition for non-defendable resources may have discouraged hummingbirds from modifying their trapline and risking overlap with another bird's foraging route (Young 1971; Gill 1988; Ohashi and Thomson 2009). Lastly, is possible that fine-scale rewiring did occur but was not detected using our analytical approach, which aggregated visitation rates to the level of hummingbird species; further analysis examining the diversity of pollen grains carried by individual hummingbirds may shed light on the possibility of rewiring at finer scales.

In the absence of rewiring, the resilience in hummingbird behavior, abundance, and pollination services following experimental *Heliconia* removal is surprising. Brief pauses in nectar intake can disrupt a hummingbird's daily energy balance, given their high metabolic rate and limited energy reserves (Calder and Booser 1973; Tooze and Gass 1985; Brice 1992). Moreover, previous nectar manipulations with hummingbirds – albeit within defended territories – led to rapid changes in territory size, foraging patterns, and/or defense behavior, generally within a day (e.g., Hixon, Carpenter, and Paton 1983; Gass and Sutherland 1985; Eberhard and

Ewald 1994; Temeles et al. 2004). One explanation for our result is that hummingbirds permanently emigrated from fragments after *Heliconia* removal but, because of high intruder pressure, were quickly replaced by new individuals (Stiles and Wolf 1970). However, we monitored individual birds using several different approaches (radio telemetry, recaptures, visitation rates) and found no evidence that *Heliconia* removal caused emigration and subsequent replacement. An alternative possibility is that hummingbirds relied on physiological adaptations, such as a torpor and/or insectivory, to maintain body condition despite large declines in nectar availability – resulting in the stable body mass measurements we observed in recaptured hummingbirds. Torpor is an energy-saving adaptation characterized by declines in body temperature and metabolic rate (Hainsworth, Collins, and Wolf 1977; Spence and Tingley 2021), and even tropical hummingbirds inhabiting low-to-mid elevations appear to possess this capability, including the green hermit (Krüger, Prinzinger, and Schuchmann 1982; Bech et al. 1997; Shankar et al. 2020; Schuchmann and Prinzinger 1988). Additionally, hummingbirds are highly insectivorous and, although they cannot completely replace nectar with insects (Brice 1992; Stiles 1995), they may supplement their diet with arthropods when nectar becomes scarce (Young 1971; Kuban and Neill 1980; Montgomerie and Redsell 1980; Hazlehurst and Karubian 2018). Finding torpid hummingbirds at night is extremely challenging (Carpenter and Hixon 1988), but future research could explore changes in arthropod consumption using techniques such as DNA metabarcoding (Moran, Prosser, and Moran 2019).

Another explanation for hummingbird persistence following experimental *Heliconia* removal involves their high movement ability. *H. tortuosa* is a patchily distributed species that necessitates long-distance foraging movements, as a single clump does not make territoriality energetically profitable (Stiles 1975). Densities can also vary substantially across short distances;

for example, Stiles (1979) notes that some Costa Rican *Heliconia* species can be common in one watershed, yet nearly absent in an adjacent watershed < 1 km away (see also Bruna and Kress 2002). The most frequent *Heliconia* visitor, the green hermit, is therefore capable of traveling long distances (>500m) while foraging within home ranges of >150 ha, although home range sizes of ~2 ha are more typical (Volpe et al. 2014; 2016). Thus, returning to our focal areas (typically ~1.5 ha) to visit the few remaining *Heliconia* resources could be perceived as a relatively low-cost option for long-distance trappiners, at least in the short term. Notably, although highly mobile pollinators might buffer against network collapse (Volpe et al. 2016), they may be vulnerable to habitat loss and fragmentation once thresholds are crossed, leading to parallel pollinator declines observed in nature (Andrn 1994; Hadley and Betts 2012). This strategy is also unlikely to be profitable over longer time scales than measured by our experiment, which only removed *Heliconia* for five days.

The relatively short length of our removal experiment might also explain why – when examining hummingbird responses across a natural gradient in *Heliconia* availability – lower *Heliconia* densities were indeed associated with reduced floral visitation rates. Although a short adjustment period (<1 day) is standard in similar species removal experiments (e.g., Hixon, Carpenter, and Paton 1983; Brosi and Briggs 2013; Hazlehurst and Karubian 2018), extending the experimental timeline may have more realistically simulated a future extinction and provided more time for behavioral or demographic adjustment. If hummingbirds were not permanent residents of our focal areas – as suggested by our relatively low recapture rates (~15%) and proportion of time spent in the focal area (~20%) – then transient individuals might have not immediately noticed changes in *Heliconia* density and changed their foraging patterns accordingly. Likewise, hummingbirds might not have persisted had we removed *Heliconia* over

larger areas, because traveling to isolated plants would become even more energetically costly (Heinrich and Raven 1972). Unfortunately, covering (and then uncovering) *Heliconia* plants in steep terrain with dense understories limited the area we could manipulate.

We also acknowledge that our ability to detect experimental effects may have been limited by low statistical power, owing to high individual-level variation and limited sample size. Low statistical power may have also weakened our tests of the connectivity hypothesis and resource hypothesis, as those datasets only included a subset of the replicates included in the experimental analyses (resource availability data were not available for all sites). Still, we emphasize that the spatial scale of our experiment and amount of replication surpasses most other empirical studies simulating species extinctions within plant-pollinator networks (Appendix S1: Table S1). Nevertheless, future removal experiments with highly mobile pollinators should consider increasing the amount of replication, and, if possible, manipulate larger areas.

In the space-for-time component of our study, we found that visitation rates to individual plants decreased with natural declines in *Heliconia* density, suggesting that *Heliconia* and its pollinators might facilitate pollination of co-flowering plants in this study system. Indeed, facilitative effects arising from the overall attractiveness of the floral neighborhood are expected to operate most strongly in systems where plants are spatially dispersed and pollinator density is low (Schemske 1981; Ghazoul 2005; Moeller 2004), as is the case in the tropical forest understory. If *Heliconia* population densities decrease due to habitat fragmentation (Hadley et al. 2014), plants visited exclusively by *Heliconia*-associated hummingbirds might therefore experience a higher risk of reproductive failure (e.g., *Pitcairnia imbricata*, *Calathea guzmanoides*, *Aphelandra golfodulcensis*, Fig. 3). Whether visitation rates map onto pollination

success for these species remains an open question, since pollinator sharing may lead to heterospecific pollen transfer and pollen loss (Ashman and Arceo-Gómez 2013; Morales and Traveset 2008) – tipping the balance toward interspecific competition rather than facilitation (Ghazoul 2005; Mitchell et al. 2009; Rathcke 1983). However, we did discover a positive relationship between visitation rate and pollination success for *Heliconia*. This relationship has been predicted for other *Heliconia* species but yielded mixed results (Stein et al. 2013; Bruna et al. 2004), in accordance with accumulating evidence that facilitation effects can be highly context-specific (Bergamo et al. 2020). Nevertheless, this finding suggests that any reproductive declines of *Heliconia* in this study system might accelerate once extinction debt is realized and conspecific density begins to decline.

CONCLUSIONS

Although pollinators seemed able to persist during our experiments that removed floral resources for a short period of time (several days), this persistence may not be sustainable or reflect the outcome of a natural extinction process. Indeed, we observed declines in hummingbird visitation when *Heliconia* availability was naturally low; this reduced hummingbird visitation had cascading consequences for pollination success. Thus, longer-term experiments could be key for predicting consequences of species loss and understanding mechanisms of network robustness – at least for pollinator species that forage at large spatial scales. Our study also suggests that, over the short term, behavioral flexibility to use alternative floral resources might be limited, even in taxonomic groups thought to be highly opportunistic foragers. Thus, even rewiring rules with strong biological underpinnings (e.g., Vizentin-Bugoni et al. 2020) may not necessarily predict real-world responses to species loss, perhaps due to additional constraints such as competition. To further understand the mechanisms underlying

network robustness or collapse, we encourage future experiments that consider how temporal scale, species traits (e.g., movement ability), competition, and landscape context interact to determine the outcome of species extinctions. These experiments could be especially valuable at geographic range limits, where climate change will likely lead to species replacements (HilleRisLambers et al. 2013), and in the tropics where landscape change and species loss are most rapid (Hansen et al. 2013).

ACKNOWLEDGMENTS

This material is based upon work supported by the National Science Foundation (NSF) Graduate Research Fellowship Program under Grant No. 1840998 to KGL. Funding for the experiment was provided primarily by NSF-DEB-1457837 to MGB and ASH, with additional grants to KGL from the American Ornithological Society and Sigma Xi. We also thank the Las Cruces Biological Station and the private landowners who allowed us to work on their property. Field assistance was provided by Jorge Araya Paniagua, Michael Atencio Picado, Marion Donald, Dustin Gannon, Jessica Greer, Matt Hadley, Urs Kormann, Daniel Moreno, Mauricio Paniagua Castro, Ignacio Ruiz Ilama, Esteban Sandi Paniagua, Andrés Solorzano Vargas, Laura Sutcliffe, and Felipe Torres Vanegas. Videos from trail cameras were reviewed with the help of Marion Donald, Colleen Franklin, Jessica Greer, Zoe Griffith, Elena Hart, Billy Hilgert, Zachary Kendall, Jeremy Lee, Ana Medina Roman, Briley Mullin, Amber Newell, Thanh Nguyen, Zoe Sallada, Allison Simoni, Claire Woods, and Mel Xiao. Nozomi Birkett and Marion Donald helped collect nectar measurements, and Chandler Spruill assisted with pollen tube processing. Ariel Muldoon provided valuable statistical advice, and Mark Novak provided feedback on an earlier version of the manuscript. Response variable icons were created from the following Noun Project images: Hummingbird by Agne Alesiute, net by Guillaume Kurkdjian, Stopwatch by

Thomas Marijinissen, leaves by Tatiana, Camera by Stanislav Levin, and Microscope by Gregor Cresnar.

LITERATURE CITED

Aguilar, R., L. Ashworth, L. Galetto, and M.A. Aizen. 2006. “Plant Reproductive Susceptibility to Habitat Fragmentation: Review and Synthesis through a Meta-Analysis.” *Ecology Letters* 9 (8): 968–80.

Aizen, M.A., M. Sabatino, and J.M. Tylianakis. 2012. “Specialization and Rarity Predict Nonrandom Loss of Interactions from Mutualist Networks.” *Science* 335 (6075): 1486–89.

Allen-Wardell, G., P. Bernhardt, R. Bitner, A. Burquez, S. Buchmann, J. Cane, P.A. Cox, et al. 1998. “The Potential Consequences of Pollinator Declines on the Conservation of Biodiversity and Stability of Food Crop Yields.” *Conservation Biology* 12 (1): 8–17.

Allison, P.D. 1990. “Change Scores as Dependent Variables in Regression Analysis.” *Sociological Methodology* 20: 93–114.

Anderson, S.H., D. Kelly, J.J. Ladley, S. Molloy, and J. Terry. 2011. “Cascading Effects of Bird Functional Extinction Reduce Pollination and Plant Density.” *Science* 331 (6020): 1068–71.

Andrén, H. 1994. “Effects of Habitat Fragmentation on Birds and Mammals in Landscapes with Different Proportions of Suitable Habitat: A Review.” *Oikos* 71 (3): 355–66.

Arizmendi, M.C., and J.F. Ornelas. 1990. “Hummingbirds and Their Floral Resources in a Tropical Dry Forest in Mexico.” *Biotropica* 22 (2): 172–80.

675 Ashman, T.L., and G. Arceo-Gómez. 2013. “Toward a Predictive Understanding of the Fitness
676 Costs of Heterospecific Pollen Receipt and Its Importance in Co-Flowering
677 Communities.” *American Journal of Botany* 100 (6): 1061–70.

678 Ashman, T.L., T.M. Knight, J.A. Steets, P. Amarasekare, M. Burd, D.R. Campbell, M.R.
679 Dudash, M.O. Johnston, S.J. Mazer, and R.J. Mitchell. 2004. “Pollen Limitation of Plant
680 Reproduction: Ecological and Evolutionary Causes and Consequences.” *Ecology* 85 (9):
681 2408–21.

682 Aslan, C.E., E.S. Zavaleta, B. Tershy, and D. Croll. 2013. “Mutualism Disruption Threatens
683 Global Plant Biodiversity: A Systematic Review.” *PLOS ONE* 8 (6): e66993.

684 Bascompte, J., and P. Jordano. 2007. “Plant-Animal Mutualistic Networks: The Architecture of
685 Biodiversity.” *Annual Review of Ecology, Evolution, and Systematics* 38 (1): 567–93.

686 Bascompte, J., P. Jordano, C.J. Melián, and J.M. Olesen. 2003. “The Nested Assembly of Plant–
687 Animal Mutualistic Networks.” *Proceedings of the National Academy of Sciences* 100
688 (16): 9383–87.

689 Bascompte, J., P. Jordano, and J.M. Olesen. 2006. “Asymmetric Coevolutionary Networks
690 Facilitate Biodiversity Maintenance.” *Science* 312 (5772): 431–33.

691 Bech, C., A.S. Abe, J.F. Steffensen, M. Berger, and J.E.P.W. Bicudo. 1997. “Torpor in Three
692 Species of Brazilian Hummingbirds under Semi-Natural Conditions.” *The Condor* 99 (3):
693 780–88.

694 Bergamo, P.J., N.S. Streher, A. Traveset, M. Wolowski, and M. Sazima. 2020. “Pollination
695 Outcomes Reveal Negative Density-Dependence Coupled with Interspecific Facilitation
696 among Plants.” *Ecology Letters* 23 (1): 129–39.

697 Betts, M.G., A.S. Hadley, and W.J. Kress. 2015. “Pollinator Recognition by a Keystone Tropical
698 Plant.” *Proceedings of the National Academy of Sciences* 112 (11): 3433–38.

699 Biesmeijer, J.C., S.P.M. Roberts, M. Reemer, R. Ohlemüller, M. Edwards, T. Peeters, A.P.
700 Schaffers, et al. 2006. “Parallel Declines in Pollinators and Insect-Pollinated Plants in
701 Britain and the Netherlands.” *Science* 313 (5785): 351–54.

702 Borgella, R., A.A. Snow, and T.A. Gavin. 2001. “Species Richness and Pollen Loads of
703 Hummingbirds Using Forest Fragments in Southern Costa Rica.” *Biotropica* 33 (1): 90–
704 109.

705 Brice, A.T. 1992. “The Essentiality of Nectar and Arthropods in the Diet of the Anna’s
706 Hummingbird (Calypte Anna).” *Comparative Biochemistry and Physiology Part A:
707 Physiology* 101 (1): 151–55.

708 Brooks, M.E., K. Kristensen, K.J. van Benthem, A. Magnusson, C.W. Berg, A. Nielsen, H.J.
709 Skaug, M. Maechler, and B.M. Bolker. 2017. “GlmTMB Balances Speed and
710 Flexibility Among Packages for Zero-Inflated Generalized Linear Mixed Modeling.” *The
711 R Journal* 9 (2): 378–400.

712 Brosi, B.J., and H.M. Briggs. 2013. “Single Pollinator Species Losses Reduce Floral Fidelity and
713 Plant Reproductive Function.” *Proceedings of the National Academy of Sciences* 110
714 (32): 13044–48.

715 Bruna, E.M., and W.J. Kress. 2002. “Habitat Fragmentation and the Demographic Structure of
716 an Amazonian Understory Herb (*Heliconia Acuminata*).” *Conservation Biology* 16 (5):
717 1256–66.

718 Bruna, E.M., W.J. Kress, F. Marques, and O.F. da Silva. 2004. “Heliconia Acuminata
719 Reproductive Success Is Independent of Local Floral Density.” *Acta Amazonica* 34 (3):
720 467–71.

721 Calder, W.A., and J. Booser. 1973. “Hypothermia of Broad-Tailed Hummingbirds during
722 Incubation in Nature with Ecological Correlations.” *Science* 180 (4087): 751–53.

723 Carpenter, F.L., and M.A. Hixon. 1988. “A New Function for Torpor: Fat Conservation in a
724 Wild Migrant Hummingbird.” *The Condor* 90 (2): 373–78.

725 Colwell, R.K., R.R. Dunn, and N.C. Harris. 2012. “Coextinction and Persistence of Dependent
726 Species in a Changing World.” *Annual Review of Ecology, Evolution, and Systematics* 43
727 (1): 183–203.

728 Costa, J.M., J.A. Ramos, L.P. da Silva, S. Timóteo, P. Andrade, P.M. Araújo, C. Carneiro, et al.
729 2018. “Rewiring of Experimentally Disturbed Seed Dispersal Networks Might Lead to
730 Unexpected Network Configurations.” *Basic and Applied Ecology* 30: 11–22.

731 Dalsgaard, B., P.K. Maruyama, J. Sonne, K. Hansen, T.B. Zanata, S. Abrahamczyk, R. Alarcón,
732 et al. 2021. “The Influence of Biogeographical and Evolutionary Histories on
733 Morphological Trait-Matching and Resource Specialization in Mutualistic
734 Hummingbird–Plant Networks.” *Functional Ecology* 35 (5): 1120–33.

735 Dunn, R.R., N.C. Harris, R.K. Colwell, L.P. Koh, and N.S. Sodhi. 2009. “The Sixth Mass
736 Coextinction: Are Most Endangered Species Parasites and Mutualists?” *Proceedings of*
737 *the Royal Society B: Biological Sciences* 276 (1670): 3037–45.

738 Eberhard, J.R., and P.W. Ewald. 1994. “Food Availability, Intrusion Pressure and Territory Size:
739 An Experimental Study of Anna’s Hummingbirds (*Calypte Anna*).” *Behavioral Ecology*
740 *and Sociobiology* 34 (1): 11–18.

Farwig, N., D. Bailey, E. Bochud, J.D. Herrmann, E. Kindler, N. Reusser, C. Schüepp, and M.H. Schmidt-Entling. 2009. "Isolation from Forest Reduces Pollination, Seed Predation and Insect Scavenging in Swiss Farmland." *Landscape Ecology* 24 (7): 919–27.

Feinsinger, P., and R.K. Colwell. 1978. "Community Organization among Neotropical Nectar-Feeding Birds." *American Zoologist* 18 (4): 779–95.

Ferrero, V., S. Castro, J. Costa, P. Acuña, L. Navarro, and J. Loureiro. 2013. "Effect of Invader Removal: Pollinators Stay but Some Native Plants Miss Their New Friend." *Biological Invasions* 15 (10): 2347–58.

Garibaldi, L.A., I. Steffan-Dewenter, C. Kremen, J.M. Morales, R. Bommarco, S.A. Cunningham, L.G. Carvalheiro, N.P. Chacoff, J.H. Dudenhöffer, and S.S. Greenleaf. 2011. "Stability of Pollination Services Decreases with Isolation from Natural Areas despite Honey Bee Visits." *Ecology Letters* 14 (10): 1062–72.

Gass, C.L., and G.D. Sutherland. 1985. "Specialization by Territorial Hummingbirds on Experimentally Enriched Patches of Flowers: Energetic Profitability and Learning." *Canadian Journal of Zoology* 63 (9): 2125–33.

Ghazoul, J. 2005. "Pollen and Seed Dispersal among Dispersed Plants." *Biological Reviews* 80 (03): 413–43.

Gilbert, L.E. 1980. "Food Web Organization and Conservation of Neotropical Diversity." In *Conservation Biology: An Evolutionary-Ecological Perspective*, edited by M.E. Soulé and B.A. Wilcox, 11–33. Sunderland: Sinauer.

Gill, F.B. 1988. "Trapline Foraging by Hermit Hummingbirds: Competition for an Undefended, Renewable Resource." *Ecology* 69 (6): 1933–42.

763 Goldstein, J., and M. Zych. 2016. “What If We Lose a Hub? Experimental Testing of Pollination
764 Network Resilience to Removal of Keystone Floral Resources.” *Arthropod-Plant*
765 *Interactions* 10 (3): 263–71.

766 Gorelick, N., M. Hancher, M. Dixon, S. Ilyushchenko, D. Thau, and R. Moore. 2017. “Google
767 Earth Engine: Planetary-Scale Geospatial Analysis for Everyone.” *Remote Sensing of*
768 *Environment* 202: 18–27.

769 Hadley, A.S., and M.G. Betts. 2009. “Tropical Deforestation Alters Hummingbird Movement
770 Patterns.” *Biology Letters* 5 (2): 207–10.

771 ———. 2012. “The Effects of Landscape Fragmentation on Pollination Dynamics: Absence of
772 Evidence Not Evidence of Absence.” *Biological Reviews* 87 (3): 526–44.

773 Hadley, A.S., S.J.K. Frey, W.D. Robinson, and M.G. Betts. 2018. “Forest Fragmentation and
774 Loss Reduce Richness, Availability, and Specialization in Tropical Hummingbird
775 Communities.” *Biotropica* 50 (1): 74–83.

776 Hadley, A.S., S.J.K. Frey, W.D. Robinson, W.J. Kress, and M.G. Betts. 2014. “Tropical Forest
777 Fragmentation Limits Pollination of a Keystone Understory Herb.” *Ecology* 95 (8):
778 2202–12.

779 Hainsworth, F.R., B.G. Collins, and L.L. Wolf. 1977. “The Function of Torpor in
780 Hummingbirds.” *Physiological Zoology* 50 (3): 215–22.

781 Hanski, I. 1994. “Patch-Occupancy Dynamics in Fragmented Landscapes.” *Trends in Ecology &*
782 *Evolution* 9 (4): 131–35.

783 ———. 1998. “Metapopulation Dynamics.” *Nature* 396 (6706): 41–49.

784 Hartig, F. 2020. *DHARMA: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)*
785 *Regression Models*. <https://CRAN.R-project.org/package=DHARMA>.

Hazlehurst, J.A., and J.O. Karubian. 2018. “Impacts of Nectar Robbing on the Foraging Ecology of a Territorial Hummingbird.” *Behavioural Processes* 149: 27–34.

Heinrich, B., and P.H. Raven. 1972. “Energetics and Pollination Ecology.” *Science* 176 (4035): 597–602.

HilleRisLambers, J., M.A. Harsch, A.K. Ettinger, K.R. Ford, and E.J. Theobald. 2013. “How Will Biotic Interactions Influence Climate Change–Induced Range Shifts?” *Annals of the New York Academy of Sciences* 1297 (1): 112–25.

Hixon, M.A., F.L. Carpenter, and D.C. Paton. 1983. “Territory Area, Flower Density, and Time Budgeting in Hummingbirds: An Experimental and Theoretical Analysis.” *The American Naturalist* 122 (3): 366–91.

Jackson, E.R. 2017. “Does the Matrix Matter? Using Radio Frequency Identification (RFID) Technology to Examine Hummingbird Movements in Fragmented Tropical Landscapes.” Master’s thesis, Oregon State University.
https://ir.library.oregonstate.edu/concern/graduate_thesis_or_dissertations/cj82kd06n.

Jordán, F. 2009. “Keystone Species and Food Webs.” *Philosophical Transactions of the Royal Society B: Biological Sciences* 364 (1524): 1733–41.

Kaiser-Bunbury, C.N., J. Mougai, A.E. Whittington, T. Valentin, R. Gabriel, J.M. Olesen, and N. Blüthgen. 2017. “Ecosystem Restoration Strengthens Pollination Network Resilience and Function.” *Nature* 542 (7640): 223–27.

Kaiser-Bunbury, C.N., S. Muff, J. Memmott, C.B. Müller, and A. Caflisch. 2010. “The Robustness of Pollination Networks to the Loss of Species and Interactions: A Quantitative Approach Incorporating Pollinator Behaviour.” *Ecology Letters* 13 (4): 442–52.

809 Kearns, C.A., and D.W. Inouye. 1993. *Techniques for Pollination Biologists*. Niwot: University
810 Press of Colorado.

811 Kearns, C.A., D.W. Inouye, and N.M. Waser. 1998. “Endangered Mutualisms: The Conservation
812 of Plant-Pollinator Interactions.” *Annual Review of Ecology and Systematics* 29 (1): 83–
813 112.

814 Klein, A.-M., B.E. Vaissière, J.H. Cane, I. Steffan-Dewenter, S.A. Cunningham, C. Kremen, and
815 T. Tscharntke. 2007. “Importance of Pollinators in Changing Landscapes for World
816 Crops.” *Proceedings of the Royal Society B: Biological Sciences* 274 (1608): 303–13.

817 Koh, L.P., R.R. Dunn, N.S. Sodhi, R.K. Colwell, H.C. Proctor, and V.S. Smith. 2004. “Species
818 Coextinctions and the Biodiversity Crisis.” *Science* 305 (5690): 1632–34.

819 Kormann, U., C. Scherber, T. Tscharntke, N. Klein, M. Larbig, J.J. Valente, A.S. Hadley, and
820 M.G. Betts. 2016. “Corridors Restore Animal-Mediated Pollination in Fragmented
821 Tropical Forest Landscapes.” *Proceedings of the Royal Society B: Biological Sciences*
822 283 (1823): 20152347.

823 Krüger, K., R. Prinzinger, and K.-L. Schuchmann. 1982. “Torpor and Metabolism in
824 Hummingbirds.” *Comparative Biochemistry and Physiology Part A: Physiology* 73 (4):
825 679–89.

826 Kuban, J.F., and R.L. Neill. 1980. “Feeding Ecology of Hummingbirds in the Highlands of the
827 Chisos Mountains, Texas.” *The Condor* 82 (2): 180–85.

828 Kuussaari, M., R. Bommarco, R.K. Heikkinen, A. Helm, J. Krauss, R. Lindborg, E. Öckinger, et
829 al. 2009. “Extinction Debt: A Challenge for Biodiversity Conservation.” *Trends in*
830 *Ecology & Evolution* 24 (10): 564–71.

831 Lenth, R. 2020. *Emmeans: Estimated Marginal Means, Aka Least-Squares Means*.
832 <https://CRAN.R-project.org/package=emmeans>.
833 Lüdtke, D. 2018. “Ggeffects: Tidy Data Frames of Marginal Effects from Regression Models.”
834 *Journal of Open Source Software* 3 (26): 772.
835 Lüdtke, D., M. Ben-Shachar, I. Patil, P. Waggoner, and D. Makowski. 2021. “Performance: An
836 R Package for Assessment, Comparison and Testing of Statistical Models.” *Journal of*
837 *Open Source Software* 6 (60): 3139.
838 Maglianesi, M.A., N. Blüthgen, K. Böhning-Gaese, and M. Schleuning. 2014. “Morphological
839 Traits Determine Specialization and Resource Use in Plant–Hummingbird Networks in
840 the Neotropics.” *Ecology* 95 (12): 3325–34.
841 Maruyama, P.K., G.M. Oliveira, C. Ferreira, B. Dalsgaard, and P.E. Oliveira. 2013. “Pollination
842 Syndromes Ignored: Importance of Non-Ornithophilous Flowers to Neotropical Savanna
843 Hummingbirds.” *Naturwissenschaften* 100 (11): 1061–68.
844 McDonald, T.L., W.P. Erickson, and L.L. McDonald. 2000. “Analysis of Count Data from
845 Before-after Control-Impact Studies.” *Journal of Agricultural, Biological, and*
846 *Environmental Statistics* 5 (3): 262–79.
847 Memmott, J., N.M. Waser, and M.V. Price. 2004. “Tolerance of Pollination Networks to Species
848 Extinctions.” *Proceedings of the Royal Society of London B: Biological Sciences* 271
849 (1557): 2605–11.
850 Mendenhall, C.D., and A.M. Wrona. 2018. “Improving Tree Cover Estimates for Fine-Scale
851 Landscape Ecology.” *Landscape Ecology* 33 (10): 1691–96.
852 Mitchell, R.J., R.J. Flanagan, B.J. Brown, N.M. Waser, and J.D. Karron. 2009. “New Frontiers in
853 Competition for Pollination.” *Annals of Botany* 103 (9): 1403–13.

854 Moeller, D.A. 2004. “Facilitative Interactions among Plants via Shared Pollinators.” *Ecology* 85
855 (12): 3289–3301.

856 Moilanen, A., and M. Nieminen. 2002. “Simple Connectivity Measures in Spatial Ecology.”
857 *Ecology* 83 (4): 1131–45.

858 Montero-Castaño, A., and M. Vilà. 2012. “Impact of Landscape Alteration and Invasions on
859 Pollinators: A Meta-Analysis.” *Journal of Ecology* 100 (4): 884–93.

860 Montgomerie, R.D., and C.A. Redsell. 1980. “A Nesting Hummingbird Feeding Solely on
861 Arthropods.” *The Condor* 82 (4): 463.

862 Morales, C.L., and A. Traveset. 2008. “Interspecific Pollen Transfer: Magnitude, Prevalence and
863 Consequences for Plant Fitness.” *Critical Reviews in Plant Sciences* 27 (4): 221–38.

864 Moran, A.J., S.W.J. Prosser, and J.A. Moran. 2019. “DNA Metabarcoding Allows Non-Invasive
865 Identification of Arthropod Prey Provisioned to Nestling Rufous Hummingbirds
866 (*Selasphorus rufus*).” *PeerJ* 7: e6596.

867 Ohashi, K., and J.D. Thomson. 2009. “Trapline Foraging by Pollinators: Its Ontogeny,
868 Economics and Possible Consequences for Plants.” *Annals of Botany* 103 (9): 1365–78.

869 Ollerton, J., R. Winfree, and S. Tarrant. 2011. “How Many Flowering Plants Are Pollinated by
870 Animals?” *Oikos* 120 (3): 321–26.

871 Pauw, A., and J.A. Hawkins. 2011. “Reconstruction of Historical Pollination Rates Reveals
872 Linked Declines of Pollinators and Plants.” *Oikos* 120 (3): 344–49.

873 Phillips, R.D., R. Peakall, B.A. Retter, K. Montgomery, M.H.M. Menz, B.J. Davis, C. Hayes,
874 G.R. Brown, N.D. Swarts, and K.W. Dixon. 2015. “Pollinator Rarity as a Threat to a
875 Plant with a Specialized Pollination System.” *Botanical Journal of the Linnean Society*
876 179 (3): 511–25.

877 Potts, S.G., J.C. Biesmeijer, C. Kremen, P. Neumann, O. Schweiger, and W.E. Kunin. 2010.
878 “Global Pollinator Declines: Trends, Impacts and Drivers.” *Trends in Ecology &*
879 *Evolution* 25 (6): 345–53.

880 Powers, D.R., A.R. Brown, and J.A. Van Hook. 2003. “Influence of Normal Daytime Fat
881 Deposition on Laboratory Measurements of Torpor Use in Territorial versus
882 Nonterritorial Hummingbirds.” *Physiological and Biochemical Zoology* 76 (3): 389–97.

883 R Core Team. 2021. *R: A Language and Environment for Statistical Computing*. Vienna,
884 Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>.

885 Ramos-Jiliberto, R., F.S. Valdovinos, P. Moisset de Espanés, and J.D. Flores. 2012.
886 “Topological Plasticity Increases Robustness of Mutualistic Networks.” *Journal of*
887 *Animal Ecology* 81 (4): 896–904.

888 Rathcke, B. 1983. “Competition and Facilitation among Plants for Pollination.” In *Pollination*
889 *Biology*, edited by L.A. Real, 305–29. New York: Academic Press.

890 Regan, E.C., L. Santini, L. Ingwall-King, M. Hoffmann, C. Rondinini, A. Symes, J. Taylor, and
891 S.H.M. Butchart. 2015. “Global Trends in the Status of Bird and Mammal Pollinators.”
892 *Conservation Letters* 8 (6): 397–403.

893 Renjifo, L.M. 2001. “Effect of Natural and Anthropogenic Landscape Matrices on the
894 Abundance of Subandean Bird Species.” *Ecological Applications* 11 (1): 14–31.

895 Rodríguez-Flores, C.I., J.F. Ornelas, S. Wethington, and M.C. Arizmendi. 2019. “Are
896 Hummingbirds Generalists or Specialists? Using Network Analysis to Explore the
897 Mechanisms Influencing Their Interaction with Nectar Resources.” *PLOS ONE* 14 (2):
898 e0211855.

Russell, S.M., and R.O. Russell. 2001. *The North American Banders' Manual for Banding Hummingbirds*. North American Banding Council.

Schemske, D.W. 1981. "Floral Convergence and Pollinator Sharing in Two Bee-Pollinated Tropical Herbs." *Ecology* 62 (4): 946–54.

Scheper, J., M. Reemer, R. van Kats, W.A. Ozinga, G.T.J. van der Linden, J.H.J. Schaminée, H. Siepel, and D. Kleijn. 2014. "Museum Specimens Reveal Loss of Pollen Host Plants as Key Factor Driving Wild Bee Decline in The Netherlands." *Proceedings of the National Academy of Sciences* 111 (49): 17552–57.

Schuchmann, K.-L., and R. Prinzinger. 1988. "Energy Metabolism, Nocturnal Torpor, and Respiration Frequency in a Green Hermit (*Phaethornis Guy*)." *Journal Für Ornithologie* 129 (4): 469–72.

Senapathi, D., L.G. Carvalheiro, J.C. Biesmeijer, C.-A. Dodson, R.L. Evans, M. McKerchar, R.D. Morton, et al. 2015. "The Impact of over 80 Years of Land Cover Changes on Bee and Wasp Pollinator Communities in England." *Proceedings of the Royal Society B: Biological Sciences* 282 (1806): 20150294.

Shankar, A., R.J. Schroeder, S.M. Wethington, C.H. Graham, and D.R. Powers. 2020. "Hummingbird Torpor in Context: Duration, More than Temperature, Is the Key to Nighttime Energy Savings." *Journal of Avian Biology* 51 (5): e02305.

Spence, A.R., and M.W. Tingley. 2021. "Body Size and Environment Influence Both Intraspecific and Interspecific Variation in Daily Torpor Use across Hummingbirds." *Functional Ecology* 35 (4): 870–83.

Steffan-Dewenter, I., and T. Tschardtke. 1999. "Effects of Habitat Isolation on Pollinator Communities and Seed Set." *Oecologia* 121 (3): 432–40.

922 Stein, K., M. Templin, I. Hensen, M. Fischer, D. Matthies, and M. Schleuning. 2013. “Negative
923 Effects of Conspecific Floral Density on Fruit Set of Two Neotropical Understory
924 Plants.” *Biotropica* 45 (3): 325–32.

925 Stiles, F.G. 1975. “Ecology, Flowering Phenology, and Hummingbird Pollination of Some Costa
926 Rican Heliconia Species.” *Ecology* 56 (2): 285–301.

927 ———. 1979. “Notes on the Natural History of Heliconia (Musaceae) in Costa Rica. Notas
928 Sobre La Historia Natural de Heliconia (Musaceae) En Costa Rica.” *Brenesia* 15: 151–
929 80.

930 ———. 1985. “Seasonal Patterns and Coevolution in the Hummingbird-Flower Community of a
931 Costa Rican Subtropical Forest.” *Ornithological Monographs* 36: 757–87.

932 ———. 1995. “Intraspecific and Interspecific Variation in Molt Patterns of Some Tropical
933 Hummingbirds.” *The Auk* 112 (1): 118–32.

934 Stiles, F.G., and L.L. Wolf. 1970. “Hummingbird Territoriality at a Tropical Flowering Tree.”
935 *The Auk* 87 (3): 467–91.

936 Stratton, D.A. 1989. “Longevity of Individual Flowers in a Costa Rican Cloud Forest: Ecological
937 Correlates and Phylogenetic Constraints.” *Biotropica* 21 (4): 308–18.

938 Suarez, R.K. 1992. “Hummingbird Flight: Sustaining the Highest Mass-Specific Metabolic Rates
939 among Vertebrates.” *Experientia* 48 (6): 565–70.

940 Taki, H., P.G. Kevan, and J.S. Ascher. 2007. “Landscape Effects of Forest Loss in a Pollination
941 System.” *Landscape Ecology* 22 (10): 1575–87.

942 Taylor, J., and S.A. White. 2007. “Observations of Hummingbird Feeding Behavior at Flowers
943 of Heliconia Beckneri and H. Tortuosa in Southern Costa Rica.” *Ornitologia Neotropical*
944 18: 133–38.

945 Taylor, P.D., L. Fahrig, K. Henein, and G. Merriam. 1993. “Connectivity Is a Vital Element of
946 Landscape Structure.” *Oikos*, 571–73.

947 Temeles, E.J., A.B. Muir, E.B. Slutsky, and M.N. Vitousek. 2004. “Effect of Food Reductions on
948 Territorial Behavior of Purple-Throated Caribs.” *The Condor* 106 (3): 691–95.

949 Thomas-Granger, K. 2003. “Light Sum, Nitrogen Level and Growth Retardant Effects on Flower
950 and Nectar Production in *Stachytarpheta Frantzii*.” Master’s thesis.

951 Timóteo, S., J.A. Ramos, I.P. Vaughan, and J. Memmott. 2016. “High Resilience of Seed
952 Dispersal Webs Highlighted by the Experimental Removal of the Dominant Disperser.”
953 *Current Biology* 26 (7): 910–15.

954 Tooze, Z.J., and C.L. Gass. 1985. “Responses of Rufous Hummingbirds to Midday Fasts.”
955 *Canadian Journal of Zoology* 63 (10): 2249–53.

956 Traveset, A., C. Tur, and V.M. Eguíluz. 2017. “Plant Survival and Keystone Pollinator Species
957 in Stochastic Coextinction Models: Role of Intrinsic Dependence on Animal-Pollination.”
958 *Scientific Reports* 7: 6915.

959 Underwood, A.J. 1994. “On Beyond BACI: Sampling Designs That Might Reliably Detect
960 Environmental Disturbances.” *Ecological Applications* 4 (1): 3–15.

961 Valdovinos, F.S., P. Moisset de Espanés, J.D. Flores, and R. Ramos-Jiliberto. 2013. “Adaptive
962 Foraging Allows the Maintenance of Biodiversity of Pollination Networks.” *Oikos* 122
963 (6): 907–17.

964 Vellend, M., K. Verheyen, H. Jacquemyn, A. Kolb, H.V. Calster, G. Peterken, and M. Hermy.
965 2006. “Extinction Debt of Forest Plants Persists for More Than a Century Following
966 Habitat Fragmentation.” *Ecology* 87 (3): 542–48.

967 Vizentin-Bugoni, J., V.J. Debastiani, V.A.G. Bastazini, P.K. Maruyama, and J.H. Sperry. 2020.
 968 “Including Rewiring in the Estimation of the Robustness of Mutualistic Networks.”
 969 *Methods in Ecology and Evolution* 11 (1): 106–16.

970 Vizentin-Bugoni, J., P.K. Maruyama, and M. Sazima. 2014. “Processes Entangling Interactions
 971 in Communities: Forbidden Links Are More Important than Abundance in a
 972 Hummingbird–Plant Network.” *Proceedings of the Royal Society B: Biological Sciences*
 973 281 (1780): 20132397.

974 Volpe, N.L., A.S. Hadley, W.D. Robinson, and M.G. Betts. 2014. “Functional Connectivity
 975 Experiments Reflect Routine Movement Behavior of a Tropical Hummingbird Species.”
 976 *Ecological Applications* 24 (8): 2122–31.

977 Volpe, N.L., W.D. Robinson, S.J.K. Frey, A.S. Hadley, and M.G. Betts. 2016. “Tropical Forest
 978 Fragmentation Limits Movements, but Not Occurrence of a Generalist Pollinator
 979 Species.” *PLOS ONE* 11 (12): e0167513.

980 Weiner, C.N., M. Werner, K.E. Linsenmair, and N. Blüthgen. 2014. “Land-Use Impacts on
 981 Plant–Pollinator Networks: Interaction Strength and Specialization Predict Pollinator
 982 Declines.” *Ecology* 95 (2): 466–74.

983 Weinstein, B.G. 2015. “MotionMeerkat: Integrating Motion Video Detection and Ecological
 984 Monitoring.” *Methods in Ecology and Evolution* 6 (3): 357–62.

985 Weinstein, B.G., and C.H. Graham. 2017. “Persistent Bill and Corolla Matching despite Shifting
 986 Temporal Resources in Tropical Hummingbird–Plant Interactions.” *Ecology Letters* 20
 987 (3): 326–35.

988 Winfree, R., I. Bartomeus, and D.P. Cariveau. 2011. “Native Pollinators in Anthropogenic
 989 Habitats.” *Annual Review of Ecology, Evolution, and Systematics* 42 (1): 1–22.

- 990 Wolf, L.L., F.R. Hainsworth, and F.G. Stiles. 1972. “Energetics of Foraging: Rate and Efficiency
991 of Nectar Extraction by Hummingbirds.” *Science* 176 (4041): 1351–52.
- 992 Wolf, L.L., and F.G. Stiles. 1989. “Adaptations for the ‘fail-Safe’ Pollination of Specialized
993 Ornithophilous Flowers.” *The American Midland Naturalist* 121 (1): 1–10.
- 994 Young, A.M. 1971. “Foraging for Insects by a Tropical Hummingbird.” *The Condor* 73 (1): 36–
995 45.
- 996 Zahawi, R.A., G. Duran, and U. Kormann. 2015. “Sixty-Seven Years of Land-Use Change in
997 Southern Costa Rica.” *PLOS ONE* 10 (11): e0143554.

998 **TABLES**

999 **Table 1.** Hypotheses testing whether declines of a locally common, tropical flowering plant (*Heliconia tortuosa*) affect hummingbird
1000 space use, hummingbird floral visitation rates, and plant pollination success. Predicted effects are indicated in the column below each
1001 hypothesis. To address Question 1, we (i) experimentally removed *H. tortuosa* using a BACI (Before-After-Control-Impact)
1002 experimental design and (ii) studied a natural gradient in *H. tortuosa* density. Under the experimental framework, predicted effects can
1003 be conceptualized as pre-post changes in treatment replicates, compared to to pre-post changes in control replicates. To address
1004 Question 2, we examined changes in hummingbird and plant responses across a natural gradient in *H. tortuosa* density only.

			Question 1: How do hummingbirds respond to declines in <i>Heliconia</i> , and are there cascading effects on the remaining plants?		Question 2: What factors mediate hummingbird responses to <i>Heliconia</i> declines?				
					Connectivity hypothesis: Rescue (H-2a)	Connectivity hypothesis: Abandonment (H-2b)	Resource hypothesis (H-3)		
			Response variable	Methodological approach	Robust network hypothesis (H-1a)	Parallel extinction hypothesis (H-1b)	<i>In highly connected forest fragments...</i>	<i>In highly connected forest fragments...</i>	<i>In forest fragments with high resource availability...</i>
Space use			Abundance	Mist net captures	No change	Decrease			
			Time spent in focal area	Radio telemetry					
Floral visitation rates			Visitation to focal <i>Heliconia</i> plants	Remote camera observations	No change or increase	Decrease	Smaller decrease	Larger decrease	Smaller decrease
			Visitation to non- <i>Heliconia</i> plants	Remote camera observations	Increase (rewiring)	Decrease			
Pollination services			Pollination success	Pollen tubes	No change or increase	Decrease			
1005									
1006									

1007 **FIGURES**

1008

Figure 1. This study was conducted within forest fragments in the landscape surrounding the Las Cruces Biological Station (A) in southern Costa Rica (B). Map in (A) illustrates forest (grey) and non-forest (white), with non-forest primarily comprising pasture. The dark grey forest layer was hand-digitized for this study; the light grey forest layer, available within the geographic boundary of Coto Brus Canton, was obtained from Mendenhall & Wrona (2018). We utilized fourteen focal landscapes of 1000 m radius ('sites', dashed circles), numbered here by increasing connectivity to other areas of forest. Most sites were used more than once across three years of study (2016-2018). Within a year, each site received either the *Heliconia* removal treatment or was assigned to be a control. Treatments were reversed in alternate years for a total sample size of 16 control replicates and 16 treatment replicates. An example site is depicted in (C). Red outlines in (C) and (D) represent a focal area in which we removed *Heliconia* (in treatment replicates) and surveyed floral resources (in all replicates). Within each focal area, we captured hummingbirds using mist nets (triangles in D) and surveyed plant-hummingbird interactions using trail cameras (icons in D). To quantify functional connectivity, we summed the total area of connected forest within the focal landscape, with connectedness based on 50-m gap crossing ability and pixels down-weighted based on distance to focal area (E; see Appendix S2 for details).

Figure 2. (A) During the *Heliconia* removal experiment, we measured four main response variables: captured hummingbird abundance, proportion of time radio-tagged hummingbirds spent in the focal area, floral visitation rates, and plant pollination success. Each response variable was measured during both experimental periods (pre and post); the sampling schedule for each is denoted with an 'X'. Each experimental replicate used two sites (control and treatment) sampled using this schedule, offset by one day. Starting around noon on the fourth day, we covered *Heliconia* inflorescences in the treatment site. (B) *Heliconia* inflorescences were covered with dark-colored bags that prevented hummingbirds from accessing any nectar present. (C) Our manipulation removed, on average, 42.5% ($\pm 30.9\%$ SD) of the calories available for the entire hummingbird community and 65.5% ($\pm 31.3\%$ SD) of the available calories for *Heliconia* specialists (green hermits and violet sabrewings). Mean values are denoted by red diamonds, and the darker black line represents the median. Lower and upper lines within boxplots represent the first and third quartiles. Whiskers extend from the first and third quartiles to the lowest and highest values within 1.5 times the interquartile range. Individual data points

- 1041 appear next to each boxplot. *Response variable icons in (A) were created using images from the*
- 1042 *Noun Project (see 'Acknowledgments'). Photos in (B) were provided by Marion Donald.*

Figure 3. The overall plant-hummingbird visitation network for forest fragments in the landscape surrounding the Las Cruces Biological Station (A), created from 15,196 hours of camera observations across 37 plant species, 14 study sites, and three field seasons (2016-2018). (B) The network in A, but subset to focus on visitation by green hermits (*P. guy*) and violet sabrewings (*C. hemileucurus*), the two most frequent visitors to *Heliconia*. The height of each rectangle represents the overall observation frequency for that species (larger rectangle = more observations), while width of the connecting links reflects relative visitation rate to different partners (wider link = higher visitation rate). Five plant species (*Calathea lutea*, *Ctenanthe dasycarpa*, *Erythrina poeppigiana*, *Maripa nicaraguensis*, *Mucuna globulifera*) received no hummingbird visitation during the observation period and are thus not represented in these networks.

Figure 4. Hummingbird space use before and after experimental local extinction of an understory plant, *Heliconia tortuosa*, within forest fragments in southern Costa Rica. Despite removing at least 43% (± 31 SD) of calories available to hummingbirds within focal areas of 1.7 ha (± 1.9 SD), we did not observe an effect of *Heliconia* removal on the number of hummingbirds captured (A-B) or the proportion of time that radio-tagged hummingbirds spent in the focal areas (C-D). Results are shown for all species analyzed (left) and green hermits and violet sabrewings (right). Estimated marginal means from GLMMs are presented alongside 95% confidence intervals. *P*-values refer to the interaction testing the effect of *Heliconia* removal, relative to control replicates; conceptually, a treatment effect is indicated by non-parallel lines. The estimated number of captured hummingbirds is calculated for 45.8 net-hours, the mean capture effort across all replicates.

Figure 5. Hummingbird visitation rates to focal *Heliconia* plants and *Heliconia* pollination success before and after experimental *Heliconia* removal (A-C) and across a natural gradient in *Heliconia* density (D-F). For hummingbird visitation rates, results are shown for the entire hummingbird community (A, D) and green hermits and violet sabrewings separately (B, E). Estimated marginal means from GLMMs are presented alongside 95% confidence intervals. (A-C) *P*-values refer to the interaction testing the effect of *Heliconia* removal, relative to control replicates; conceptually, a treatment effect is indicated by non-parallel lines. (D-F) Values of *Heliconia* availability (*x*-axis) are standard deviations (1 SD = 1.35), with zero representing the mean log-transformed caloric density (mean = 8). Note that the *x*-axis is reversed, such that declining visitation is associated with declining *Heliconia* availability. *P*-values refer to the coefficient for *Heliconia* availability.

1079

Figure 6. Hummingbird visitation rates to non-*Heliconia* plant species before and after experimental *Heliconia* removal (A-B) and across a natural gradient in *Heliconia* density (C-D). Results are shown for the entire hummingbird community (A, C) and green hermits and violet sabrewings separately (B, D). Estimated marginal means from GLMMs are presented alongside 95% confidence intervals. (A-B) *P*-values refer to the interaction testing the effect of *Heliconia* removal, relative to control replicates; conceptually, a treatment effect is indicated by non-parallel lines. (C-D) Values of *Heliconia* availability (*x*-axis) are standard deviations (1 SD = 1.35), with zero representing the mean log-transformed caloric density (mean = 8). Note that the *x*-axis is reversed, such that declining visitation is associated with declining *Heliconia* availability. *P*-values refer to the coefficient for *Heliconia* availability.