RNA targeting with CRISPR-Cas13a facilitates bacteriophage genome engineering

Jingwen Guan,¹ Agnès Oromí-Bosch,² Senén D. Mendoza,¹⁺ Shweta Karambelkar,¹ Joel Berry,² Joseph Bondy-Denomy^{1,3,4,*}

¹Department of Microbiology & Immunology, University of California, San Francisco, San Francisco, CA, USA

²Felix Biotechnology, South San Francisco, CA, USA

³Quantitative Biosciences Institute, University of California, San Francisco, San Francisco, CA, USA

⁴Innovative Genomics Institute, Berkeley, CA, USA

⁺Present address: Department of Biology, Massachusetts Institute of Technology, Cambridge, MA, USA

*Corresponding author: joseph.bondy-denomy@ucsf.edu

14 Graphical abstract

15

16

17 Abstract

18 The viruses that infect bacteria, bacteriophages (or phages), possess numerous genes of
 19 unknown function. Genetic tools are required to understand their biology and enhance their
 20 efficacy as antimicrobials. *Pseudomonas aeruginosa* jumbo phage Φ KZ and its relatives are a
 21 broad host range phage family that assemble a proteinaceous “phage nucleus” structure during
 22 infection. Due to the phage nucleus, DNA-targeting CRISPR-Cas is ineffective against this phage
 23 and thus there are currently no reverse genetic tools for this family. Here, we develop a DNA
 24 phage genome editing technology using the RNA-targeting CRISPR-Cas13a enzyme as a
 25 selection tool, an anti-CRISPR gene (*acrVIA1*) as a selectable marker, and homologous
 26 recombination. Precise insertion of foreign genes, gene deletions, and the addition of
 27 chromosomal fluorescent tags into the Φ KZ genome were achieved. Deletion of *phuZ*, which
 28 encodes a tubulin-like protein that centers the phage nucleus during infection, led to the
 29 mispositioning of the phage nucleus but surprisingly had no impact on phage replication, despite
 30 a proposed role in capsid trafficking. A chromosomal fluorescent tag placed on gp93, a proposed
 31 “inner body” protein in the phage head revealed a protein that is injected with the phage genome,
 32 localizes with the maturing phage nucleus, and is massively synthesized around the phage
 33 nucleus late in infection. Successful editing of two other phages that resist DNA-targeting
 34 CRISPR-Cas systems [OMKO1 (Φ KZ-like) and PaMx41] demonstrates the flexibility of this
 35 method. RNA-targeting Cas13a system holds great promise for becoming a universal genetic
 36 editing tool for intractable phages. This phage genetic engineering platform enables the
 37 systematic study of phage genes of unknown function and the precise modification of phages for
 38 use in a variety of applications.

Introduction

Bacteriophages are viruses that infect bacteria and can cause their lysis after replication. In recent decades, the rapid emergence of multi-antibiotic resistant bacterial pathogens and simultaneous decline in the discovery of new antibiotics has rekindled interest in the use of phages as alternative antimicrobial therapeutics (phage therapy) [1, 2]. Phages offer many advantages over antibiotics, including high specificity and efficient self-propagation in the presence of their bacterial host [3-5]. However, host range limitations and the rapid emergence of phage resistance in clinical strains present barriers for furthering phage therapy [1, 3, 6]. Phage genome engineering may help overcome these hurdles [7, 8]. Robust phage engineering tools can aid fundamental discoveries, broaden host range, enhance evasion of host antiviral defense systems, and reduce phage toxicity and immunogenicity [9-12]. Phage engineering techniques often utilize homologous recombination (HR) with a template plasmid [13, 14], coupled with a selective pressure such as CRISPR-Cas targeting. CRISPR-Cas systems (clustered regularly interspaced short palindromic repeats and CRISPR-associated proteins) are adaptive anti-phage immune systems in prokaryotes [15, 16]. CRISPR-Cas programmable targeting enables effective enrichment for phage recombinants by removing wild-type phages from the population and has been coupled with the integration of an anti-CRISPR gene as a selectable marker [17].

To date, all CRISPR-based screening tools used in phage engineering recognize and target phage genomic DNA. However, due to the everlasting evolutionary battle between bacteria and phages in nature, phages have amassed various strategies to circumvent DNA-targeting immunity, including anti-CRISPR proteins, DNA base modifications, and genome segregation [18, 19]. *P. aeruginosa* jumbo phage Φ KZ is resistant to a broad spectrum of DNA-targeting immune systems via assembly of a proteinaceous “phage nucleus” structure that shields phage DNA during replication [20, 21]. This phage family is thus a great candidate for use as a phage therapeutic and likely possesses other fascinating fundamental biology. However, no genetic tools are available for this phage family and most studies have relied on plasmid-based over-expression [22].

Although the phage nucleus prevents DNA-targeting, the mRNA-targeting CRISPR-Cas13a system (type VI-A) [23] effectively inhibits Φ KZ replication by degrading phage mRNA that is exported from the phage nucleus to the cytoplasm [20]. Here, we develop the CRISPR-Cas13a system as a novel genetic engineering approach for Φ KZ. Using Cas13a to target an essential transcript, we select for phage DNA that has undergone homologous recombination resulting in a desired genetic change along with the acquisition of an anti-Cas13a *trans* gene, *acrVIA1* (derived from Listeriophage Φ LS46 [24]), as a selectable marker. This approach allows us to precisely insert foreign gene fragments into the Φ KZ genome, knock out non-essential genes, and fuse fluorescent tags to individual genes. Importantly, the same guide can be used for any genomic manipulation of a single phage as engineered phages are identified based on the acquisition of the Cas13a inhibitor, not a change in the target sequence. Our work establishes a Cas13a-based phage engineering strategy that could be a universally powerful tool for engineering phages.

Results

Optimization of CRISPR-Cas13a for efficient phage targeting

Cas13a is an RNA-guided RNA nuclease that can block Φ KZ replication in *P. aeruginosa* PAO1. We previously targeted Φ KZ by expressing crRNA guides from a plasmid (Figure 1A, Version 1) [20]. For the effective elimination of WT phages in the population, we first sought to enhance activity of crRNA guides. We designed Version 2 (V2) with the repeat-spacer-repeat unit moved to the +1 transcription start site and the second direct repeat (DR) mutated to remove repeat homology (Figure 1A). To further stabilize the crRNA cassette, we omitted the second DR and generated V3 (Figure 1A), which could prevent recombination between the repeats. Using the same spacer, both V2 and V3 provided more robust defense against phage JBD30 and Φ KZ compared with V1 (Figure 1A). For simplicity, we selected the V3 cassette that has a single repeat to express crRNAs against Φ KZ. We designed multiple spacer sequences to target various Φ KZ gene transcripts. Strong targeting was observed for some crRNAs, to the point that escaper phages could be isolated, such as the two spacers matching *orf120* and *orf146* transcripts, but not all crRNAs were efficacious (Figure 1B and S1). Given the variability of targeting efficiency, for the remainder of this report, we use *orf120*-targeting gRNA#2 as our primary guide to screen for engineered phages. We refer the PAO1 strain simultaneously expressing Cas13a and *orf120*-gRNA#2 to as Cas13a counter-selection strain. We describe below how the same guide can be used to facilitate the engineering of distinct genomic loci.

Isolation of Φ KZ recombinants by CRISPR-Cas13a counter selection

To avoid disrupting any essential genes that are required for phage replication, we first attempted to insert *acrVIA1* immediately downstream of Φ KZ major capsid gene (*orf120*). A template DNA substrate for homologous recombination, composed of ~600-bp homology arms flanking *acrVIA1* was cloned into a plasmid, referred to as an editing plasmid (Figure 2A). After infecting a PAO1 strain with the editing plasmid to allow recombination, the phage lysate was then titrated on a lawn of the Cas13a counter-selection strain to eliminate WT phages. To screen for recombinants, individual plaques were examined for *acrVIA1* integration via PCR, showing that 8 out of the 16 tested phage plaques generated the expected 1.6 kbp band, which was not detectable in a WT plaque (Figure 2B). Amplification of the entire region also revealed the expected size increase, from ~1.7 kbp of WT to ~2.5 kbp of recombinants, and Sanger sequencing confirmed the correct integration junction (Figure 2C). Recombinant phages propagated well on hosts expressing crRNAs targeting other genomic loci due to the expression of *acrVIA1*, which abolishes Cas13a immunity regardless of the crRNA sequence (Figure 2D). Three randomly selected phages that escaped Cas13a targeting but screened negative for the *acrVIA1* integration contained genomic deletions ranging from 27 bp to 69 bp starting immediately downstream of the *orf120* stop codon, disrupting the protospacer (Figure S2). While the crRNA used is therefore not inescapable, the recombination efficiency to insert the selectable marker is clearly efficient enough to enable facile identification of the desired mutants.

To test the flexibility of this nascent genetic technology (Figure 2E) and generate new biological insights of phage Φ KZ, we next knocked out (or attempted to knock out) multiple genes; *phuZ* (*orf39*), *orf54*, *orf89-orf93*, *orf93*, *orf146*, *orf241*, and *orf241-orf242* (summarized in Table 1), in addition to attempting to add chromosomal fluorescent tags onto *orf54* and *orf93*. The successes, failures, and new insights gained are discussed below. Whole genome sequencing of two deletion mutants, Δ *phuZ* and Δ *orf93*, revealed no other mutations. The accuracy of this

system highlights an important advantage of adapting an RNA-targeting system to select for edits in DNA phage genomes, compared to direct DNA cleavage.

Characterization of PhuZ and gp93 using engineered Φ KZ mutants

PhuZ (gp39) is a tubulin homolog conserved across “group 1” jumbo phages and some megaphages [25, 26]. It assembles a bipolar spindle to center the phage nucleus during phage intracellular development [27, 28], and “treadmill” newly synthesized phage capsids from the cell inner membrane to the phage nucleus for DNA packaging [29]. These functions made us speculate that PhuZ might be essential for phage growth, however, this is not the case. Δ *phuZ* mutants exhibited a similar burst size (24 phage particles per infected bacterial cell vs. 19 of WT phage) under our experimental conditions. While cells infected with WT phages or Δ *phuZ* mutants complemented *in trans* had phage nuclei in the center of the cell ~80% of the time, the localization of the phage nucleus showed a wide distribution in cells infected by Δ *phuZ* mutants (Figure 3A-3B, Movie S1). This is consistent with the previous findings that *trans* over-expression of catalytic mutant PhuZ resulted in mispositioning of the phage nucleus [28, 30]. ~25% of mutant-infected cells still positioned the phage nucleus at the cell center (Figure 3B), a phenotype most commonly seen in shorter cells (*Pearson* correlation coefficient = 0.486, $p < 0.001$), in contrast with WT infection where no correlation was observed (*Pearson* correlation coefficient = 0.029, $p = 0.505$) (Figure 3C). Considering that PhuZ is proposed to traffic phage capsids from the cell inner membrane to the phage nucleus [29, 31], we speculate that this is only required under specific conditions or not at all. Taken together, these data suggested that PhuZ positions the phage nucleus at the cell center during infection, but the removal of PhuZ does not have a significant impact on Φ KZ growth in laboratory conditions.

orf93 encodes gp93, a high copy number “inner body (IB)” protein that is packaged in the phage head [32, 33]. Deletion of *orf93* also yielded viable phage with no obvious growth defect. We next analyzed an inserted fluorescent chromosomal label at the C-terminus of the protein, which is notable as the first chromosomal protein tag in this phage family. Labeled gp93 was observed in the mature virion (Figure 4A), as predicted by previous mass spectrometry studies [33]. Excitingly, time-lapse movies revealed the fluorescently labeled protein being injected with the phage DNA at the cell pole (Figure 4B, Movie S2) and subsequently translocating to the cell center where it remained bound to the phage nucleus. As new gp93-mNeonGreen was expressed from the phage genome, more and more green signals concentrated on the surface of the phage nucleus, while some foci appeared on the cell inner membrane. Finally, cells lysed and released fluorescent phage progeny. To confirm that the protein that appears to be injected was not rapidly synthesized *de novo*, we monitored the infection behaviors of WT phages loaded with gp93-mNeonGreen expressed from a plasmid during phage production, but where no new fluorescent protein could be made during infection (Figure 4C, Movie S3). Similar to the engineered phage, the phage particles were fluorescent, injected the labeled protein, and the green focus migrated from the cell pole to the cell center along with phage DNA on the surface of the phage nucleus until cell lysis. Therefore, the IB protein gp93 is not only packaged in the phage head, but may also play a role in phage development and maturation. The ability to chromosomally label phage proteins, as demonstrated here, will be beneficial for characterizing Φ KZ virion and cell biology in the future.

To assess whether the deletions of *phuZ* or *orf93* impact growth in a strain-dependent manner, we challenged a panel of 21 *P. aeruginosa* clinical strains with the mutant phages. Plaque assays showed that host ranges of both mutants were quite similar to WT (Figure 3D, S3),

suggesting that these knockouts, and Cas13a-mediated genetic engineering in general, has no impact on the Φ KZ host range.

The phage nucleus is primarily composed of gp54 [28]. We were unable to knockout or fluorescently label *orf54*, even when wild-type gp54 was provided by expressing from a plasmid *in trans*. The primers used to amplify the region of editing generated multiple bands for both deletion and tag-addition mutant variants (Figure S4A). N- or C-terminal fusion of gp54 with mCherry tags yielded similar results. Whole genome sequencing of an isolated *orf54* “pseudo knock-out” strain revealed that part of the editing plasmid was integrated upstream of *orf54*, while the *orf54* gene was left intact (Figure S4B). A similar attempt to delete the structural gene (*orf146*) and a cluster of IB genes (*orf89-orf93*) also failed, while deletion of accessory genes *orf241* and *orf241-242* succeeded but yielded no obvious phenotype. These results highlighted that the CRISPR-Cas13a counter selection system is a strong and efficient phage genome engineering tool, but the modification of phage essential genes remains challenging.

Precise genome engineering of clinical phage OMKO1

We next explored the versatility of our phage engineering platform by editing the genome of a clinical jumbo phage. We selected OMKO1, a *P. aeruginosa* phage with a ~280 kbp genome that has high sequence identity to Φ KZ (98.23%). OMKO1 can drive re-sensitization of surviving *P. aeruginosa* cells to small-molecule antibiotics. This phage has been used for phage therapy as emergency treatment for chronic infections caused by antibiotic resistant *P. aeruginosa* [34], and it is currently being tested in a phase I/II clinical trial (CYPHY, <https://clinicaltrials.gov/ct2/show/NCT04684641>). With this phage, we tested whether we could insert “DNA barcodes” without impacting host range, for downstream clinical applications. Insertion of a DNA tracking signature into clinical phages would enable differentiation from naturally occurring phages during the manufacturing process and following administration to patients.

Two engineered OMKO1 strains were generated, one with *acrVIA1* and a 120-nt barcode inserted downstream of the capsid gene, and another with *acrVIA1* integrated upstream of the shell gene (Figure 5A, Table 1). The presence of the desired inserts were confirmed with whole genome sequencing of both OMKO1 engineered strains, and no unintended genetic changes occurred. Moreover, both strains exhibited strong resistance to Cas13a targeting (Figure 5B), owing to the expression of *acrVIA1* from the phage genomes. The host range and virulence of the two engineered OMKO1 variants together with the parental phage was then assessed on 22 *P. aeruginosa* clinical strains (including PAO1). The experiment was performed in a microplate liquid assay, where phage variants were individually mixed with each host strain at a MOI~1 and MOI~0.01. All three phages displayed the same host range (Figure 5C, S5), capable of infecting and suppressing growth of 20/22 (91%) clinical strains tested. Infections at high MOI (MOI = 1) resulted in a broader host range and greater bacterial growth suppression, while low MOI (MOI = 0.01) infections were able to suppress cell growth of 12/22 (55%) hosts. All phages exhibited similar virulence across all hosts with small differences in 5/22 strains. In 2/22 hosts (FB-89 and FB-92), both engineered OMKO1 strains displayed significantly higher liquid assay scores than the parental phage ($p < 0.05$). While mutant OMKO1::Acr-BC had weaker virulence than the parental phage in 3/22 strains. Altogether, these results indicate that OMKO1’s host range was not affected, and virulence was impacted only modestly by inserting *acrVIA1* or *acrVIA1* and a barcode in the two selected genome locations under the tested conditions.

Application of CRISPR-Cas13a phage engineering to a small lytic Podophage

To evaluate the applicability of the CRISPR-Cas13a-mediated genome editing approach to other virulent phages, we selected *P. aeruginosa* phage PaMx41. PaMx41 is a Podophage isolated from environmental and sewage water samples in Central Mexico [35]. Its genome is approximately 43.5 kbp long and harbors 55 open reading frames (ORFs), ~70% of which have unknown function [36]. Remarkably, we discovered that PaMx41 appears to be resistant to many DNA-targeting CRISPR-Cas systems (Type I-C, II-A, and V-A) and showed partial sensitivity (~10-fold reduction in efficiency of plating) to Type I-F to a degree that is not sufficient for counter selection (Figure 6A). In contrast, when the transcripts of the major capsid gene (*orf11*) were targeted by CRISPR-Cas13a, PaMx41 exhibited strong sensitivities to specific crRNAs (Figure 6B). Following the same approach as we developed to engineer Φ KZ, we successfully replaced one hypothetical gene (*orf24*) and its downstream non-coding region with *acrVIA1* and isolated a pure mutant strain using an efficient gRNA (#5) (Figure 6B, Table 1). The mutant showed expected anti-CRISPR activities against different *orf11*-targeting crRNAs, indicating that the incorporated *acrVIA1* was expressed and properly functioning (Figure 6C). No other obvious phenotype was observed for the PaMx41 Δ *orf24* phage. Notably, initial PCR screening for recombinant plaques showed that 100% of surviving phages were desired recombinants, with no spontaneous escaper plaques. The data suggest that the RNA-targeting Cas13a system holds great promise for becoming a universal genetic editing tool to deal with previously intractable phages.

Discussion

We exploited the RNA-targeting CRISPR-Cas13a system in conjunction with homologous recombination to achieve genetic modification of jumbo phage Φ KZ, OMKO1, and phage PaMx41 that all resist DNA-targeting CRISPR-Cas systems. CRISPR-Cas13a-mediated counter-selection recovered rare (~10⁻⁵) phage recombinants from a large pool of wild-type phages. Many studies have uncovered that phages can hamper CRISPR-Cas activities, for example, by repressing transcription of endogenous CRISPR-Cas components [37, 38], possessing covalent DNA modifications [39-42], or encoding anti-CRISPR proteins (recently reviewed in [43]). Furthermore, the assembly of a proteinaceous nucleus-like structure that shields phage genomes from attack by distinct DNA-targeting nucleases [20, 21] represents the ultimate “anti-CRISPR/anti-RM” mechanism. Therefore, development of a phage genomic manipulation approaches that target a relatively constant and exposed molecule, mRNA, may provide a near-universal approach. Moreover, Cas13 is rarely encoded in bacteria and thus most phages are not expected to encode anti-Cas13a proteins.

Applying gene editing to this phage allowed us to query endogenous gene function and essentiality for the first time. We observed that Δ *phuZ* mutant phages mispositioned the phage nucleus during viral intracellular development. Previous studies revealed that newly assembled phage capsids trafficked along PhuZ filaments towards the phage nucleus for viral DNA packaging [29]. However, our work suggests that successful DNA loading into capsids is not dependent on PhuZ. Moreover, loss of PhuZ appeared not to affect burst size, in contrast with a previous report using over-expression of a catalytic mutant and microscopy to estimate burst size [30]. Overall, *phuZ* seems to be a *bona fide* nonessential gene for Φ KZ. The evolutionary advantage of encoding tubulin in this jumbo phage and many others requires further investigation. Furthermore, a chromosomal fluorescent label on gp93 demonstrated that it is packaged in the phage head, injected with the genome, and massively synthesized later in infection, with peri-

nuclear localization. The labeling not only allows us to visualize individual virions under the microscope, but also to observe the injection of this inner body protein into the host cell, which had been previously suggested with little evidence [33, 44].

One major challenge of using Cas13a in our experience has been the wide variability of crRNA efficacy. Future studies focusing on the optimization of crRNA design for phage targeting or perhaps the implementation of other RNA-targeting enzymes, such as Cas13 orthologues or Cas7-11 [45, 46] will be important. However, we circumvent this problem by implementing an anti-CRISPR selectable marker [17] to ensure that the same strong guide can be used for all genetic manipulations. The downside is that this limits the user to a single perturbation, however, double and triple mutants are possible in principle if one uses crRNAs specific to the site of editing or removes the *acr* gene from the genome.

Altogether, the RNA-targeting CRISPR-Cas13a counter-selection tool should be applicable to a broad range of phages and enable downstream high throughput phage engineering. The ability to precisely and efficiently generate synthetic phages with desired features will not only benefit phage therapeutic applications but also advance our understanding of fundamental phage biology and phage-bacteria interactions.

Acknowledgements

J.B.-D. was supported by the National Institutes of Health (R01GM127489), the Vallee Foundation, the Searle Scholarship, the Innovative Genomics Institute, and the UCSF Program for Breakthrough Biomedical Research funded in part by the Sandler Foundation. This work was also supported by research funds from Felix Biotechnology.

We are grateful to Luciano Marraffini (The Rockefeller University) for providing the plasmid pAM383. We thank Paul Turner (Yale University) for providing phage OMKO1 and *P. aeruginosa* clinical isolates. We thank Gabriel Guarneros Peña at Centro de Investigación y de Estudios Avanzados for providing phage PaMx41. We thank Tomer Rotstein for his generous assistance with NGS data processing and interpretation. We thank members of the Bondy-Denomy laboratory for productive conversations and generous suggestions for our work.

Declaration of interests

J.B.-D. is a scientific advisory board member of SNIPR Biome, Excision Biotherapeutics, and Leapfrog Bio, and a scientific advisory board member and co-founder of Acrigen Biosciences. The Bondy-Denomy lab receives research support from Felix Biotechnology.

Materials and methods

Strains, DNA oligonucleotides and plasmid constructions

All bacterial and phage strains, spacer sequences, and primers used in this study are listed in Tables S1, S2, and S3, respectively.

The crRNAs designed for CRISPR-Cas13 targeting were constructed in the pHERD30T backbone. The pHERD30T-crRNA Version 2 was constructed by thermal annealing of oligonucleotides oSDM465 and oSDM466 and phosphorylation by polynucleotide kinase (PNK). The annealed product was introduced by Gibson assembly into pHERD30T linearized by PCR using oligonucleotides oSDM457 and oSDM458. Proper construction of the expression vector was verified by Sanger sequencing. The pHERD30T-crRNA Version 3 was constructed just as for Version 2, but the crRNA-coding insert was instead composed of oligonucleotides oSDM455 and oSDM456. Both Version 2 and Version 3 of this plasmid were designed such that cleavage by BsaI would generate a linear plasmid that would accept annealed oligonucleotide spacers via ligation. Oligonucleotide pairs with repeat-specific overhangs encoding spacer sequences were annealed and phosphorylated using T4 polynucleotide kinase and then cloned into the BsaI-digested empty vectors. Cloning procedures were performed in commercial *E. coli* DH5α cells (New England Biolabs) according to the manufacturer's protocols. The resulting crRNA plasmids were electroporated into *P. aeruginosa* PAO1 strain harboring the *tn7::cas13a^{Lse}* (SDM084) on the chromosome as described previously [20]. Gene expression was induced by the addition of L-arabinose at a final concentration of 0.3% and isopropyl β-D-1-thiogalactopyranoside (IPTG) at a final concentration of 1 mM.

To construct template plasmids for homologous recombination, homology arms of >500 bp in length were amplified by PCR using the ΦKZ genomic DNA as the template. To prevent Cas13a cleavage, several synonymous mutations were introduced into the crRNA-targeting site of the left *orf120* homology arm by designing the reverse primer (JG064) to contain appropriate mismatches. The *acrVIA1* gene was amplified from plasmid pAM383 [24], a gift from Luciano Marraffini, The Rockefeller University. PCR products were purified and assembled as a recombineering substrate and then inserted into the NheI site of the pHERD30T backbone by Gibson Assembly (New England Biolabs) following the manufacturer's protocols. The resulting plasmids were transformed into PAO1 by electroporation.

Isolation of phage recombinants

Host strains bearing recombination plasmids were grown in LB supplemented with 10 mM MgSO₄ and 50 μg/ml gentamicin, at 37°C with aeration at 250 rpm. When OD₆₀₀ is around 2, Wild type ΦKZ was added into the culture at a MOI (multiplicity of infection) of 1 to allow infection to occur for ~18 hours. 2% volume of chloroform was added into the infection culture and left to shake gently on an orbital shaker at room temperature for 15 min, followed by centrifugation at 4,000 x g for 15 min to remove cell debris. The supernatant lysate was further treated with 2% of chloroform for 15 min and centrifuged again under the same conditions, followed by a 30-min treatment with DNase I (New England Biolabs) at 37°C. The resulting phage lysate containing both WT phages and recombinants are tittered on PAO1 strains bearing the CRISPR-Cas13a system with the most efficient crRNA (*orf120* guide#2) to screen for recombinants. Individual phage plaques were picked from top agar and purified for three rounds using the CRISPR counter-selection strain to ensure thorough removal of any remaining WT or escapers. Whether or not they are recombinant phages or Cas13a escaper phages were determined by PCR using

appropriate pairs of primers amplifying the modified regions of the phage genome. Identified phages were further confirmed and analyzed by sequencing the PCR products or the whole genomes and then stored at 4°C.

Phage plaque assay

Host strains were grown in LBM (LB supplemented with 10 mM MgSO₄), 50 µg/ml gentamicin, 1 mM IPTG and 0.3% arabinose inducers for gene expression, at 37°C with aeration at 250 rpm for overnight. Phage spotting assays were performed using 1.5% LB agar plates and 0.42% LB top agar, both of which contained 10 mM MgSO₄ and inducers. 100 µl of appropriate overnight culture was suspended in 3.5 ml of molten top agar and then poured onto an LB+10 mM MgSO₄ agar plate, leading to the growth of a bacterial lawn. After 10-15 min at room temperature, 2 µl of ten-fold serial dilutions of phages was spotted onto the solidified top agar. Plates were incubated overnight at 37°C. Plate images were obtained using Gel Doc EZ Gel Documentation System (BioRad) and Image Lab (BioRad) software.

Microplate Liquid Assay

Fresh overnight cultures were diluted to a cell concentration of 1×10⁸ cfu/ml in TSB media supplemented with 10mM MgSO₄. Phage lysates were added to reach a MOI of ~1 and ~0.01 in a Corning Costar 96-well clear flat-bottom microplate (Thermo Fisher Scientific) sealed with a Breathe-Easy® sealing membrane (Merck KGaA). After the infection cultures were incubated at room temperature for 20 min, plates were incubated at 37°C, 800 rpm for 8 hours in a BioTek LogPhase 600 plate reader (Agilent Technologies, Inc.). Cell growth was monitored by measuring OD₆₀₀ every 20 min. Each phage-host combination was performed in three biological replicates.

Data Analysis

Growth curves for each phage-host combination were obtained by plotting OD₆₀₀ after blank correction (baseline adjustment) against time. Each growth curve was transformed into a single numerical value by calculating the area under the curve (AUC) using the Trapezoid method. Then, AUCs were normalized as a percentage of the AUC of their corresponding uninfected control following the equation,

$$\text{Liquid assay score} = \frac{\text{AUC}(\text{positive control}) - \text{AUC}(\text{phage treatment})}{\text{AUC}(\text{positive control})} \times 100$$

The resulting value, defined as the “liquid assay score”, represents how well the phage strain can repress the growth of a bacterial population over the course of the 8-hour experiment. No inhibition of bacterial growth would result in a liquid assay score of 0, and complete suppression would translate into a score of 100. Liquid assay scores were averaged using data from three biological replicates.

Burst size measurement

Phage burst size was determined by one-step growth curve experiments. Briefly, the host PAO1 strain was grown in LB media to OD₆₀₀ ~0.4. 1 ml of the cell culture was then centrifuged at 4,000 × g for 2 min and the cell pellet was resuspended in 50 µl of fresh LBM. Appropriate amount of phages was mixed with the cell culture to achieve an MOI of 0.01 to limit to single infections. The mixture was incubated on ice for 20 min for phage adsorption and transferred to a 37°C heat block for 10 min to trigger phage DNA injection. The infection mixture was centrifuged

at 10,000 × g for 2 min. Transfer 10 µl of the supernatant into 990 µl of ice-cold SM buffer supplied with 2% chloroform. Titer to calculate the number of free phages. After discarding the supernatant to remove free phage particles, the pellet was resuspended in 1 ml of LBM, followed by incubation at 37°C with shaking at 250 rpm. Samples were collected at 10-min intervals until 90 min, and phage titer was determined immediately. Phage burst sizes were calculated by dividing the phage titers at ~50 min by the initial phage titers after subtracting free phages.

Single-cell infection assay

1 ml of host cells was grown in LBM (LB supplemented with 10 mM MgSO₄) and 50 µg/ml gentamicin (if necessary), at 37°C with aeration at 250 rpm for overnight. The overnight culture was diluted 1:100 into 5 ml of LBM and grown at 37°C with 250 rpm shaking until OD₆₀₀ ~0.4. Next, 1 ml of cell culture was collected by centrifugation at 3,000 × g for 2 min at room temperature and concentrated by 25-fold in fresh LBM. 10 µl of cells were then mixed with 10 µl of appropriate phage strains to reach an appropriate MOI, followed by incubation at 30°C for 10 min to allow for phage infection. The infection mixture was further diluted by 10-fold into 50 µl of fresh LBM at room temperature. 1 µl of the diluted culture was gently placed onto a piece of agarose pad (~1 mm thick) with 1:5 diluted LBM, arabinose (0.8%), and DAPI (5 µg/ml; Invitrogen#153, No. D1306). A coverslip (No.1.5, Fisher Scientific) was gently laid over the agarose pad and the sample was imaged under the fluorescence microscope at 30°C within a cage incubator to maintain temperature and humidity.

Fluorescence microscopy and imaging

Microscopy was performed on an inverted epifluorescence (Ti2-E, Nikon, Tokyo, Japan) equipped with the Perfect Focus System (PFS) and a Photometrics Prime 95B 25-mm camera. Image acquisition and processing were performed using Nikon Elements AR software. During a time-lapse movie, the specimen was typically imaged at a time interval of 5 min at the focal plane for 2.5~3 h, through channels of phase contrast (200 ms exposure, for cell recognition), blue (DAPI, 200 ms exposure, for phage DNA), and green (GFP, 300 ms exposure, for Gp93-mNeonGreen).

Next-generation sequencing (NGS)

To isolate phage genomic DNA, purified high titer lysates were treated with Benzonase Nuclease (Sigma) for 30 min at 37°C. Phage genomic DNA was extracted with a modified Wizard DNA Clean-Up kit (Promega) protocol. DNA samples were quantified with AccuGreen Broad Range dsDNA quantification kit (Biotium, USA) in a Qubit Fluorometer 2.0.

Purified phage genomic DNA was processed following Illumina DNA Preparation Protocol. Samples were sequenced on a MiSeq system (Illumina) with 300 cycles of paired-end sequencing, and loading concentration of 12 pM. Illumina short reads were downsampled to ~50-100× coverage and de novo assembled using SPAdes. The sequences of mutant phage strains were aligned to the reference genome in Geneious with the Mauve alignment algorithm to confirm the intended genomic edits.

The isolated *orf54* “pseudo knock-out” phage strain (“*orf54*”) was sequenced using long-read sequencing. DNA samples were processed using SQK-LSK109 kit (Oxford Nanopore Technologies, UK). Libraries were sequenced using an R10.3 flow cell until the desired number

415 of reads was achieved. Oxford Nanopore long reads were filtered for the longest high quality
416 reads using Nanofilt and de novo assembled using Flye.

Figure 1. Optimization of CRISPR-Cas13a crRNA expression vector for efficient phage interference.

(A) Schematic of three versions of CRISPR-Cas13a crRNA cassette (B) Efficiency of plaquing of three versions of crRNA cassette targeting two unrelated phage strains: JBD30 and ΦKZ. Wild-type phages were spotted in ten-fold serial dilutions (left to right) on a lawn of *P. aeruginosa* PAO1 expressing *LseCas13a* from the chromosome and harboring indicated crRNA expression vectors. The cassettes carried the same spacer sequences targeting transcripts of *orf38* of JBD30 and *orf54* of ΦKZ, respectively. (C) Cas13a interference efficiency of distinct crRNAs against ΦKZ using Version 3 crRNA expression vector. The most efficient crRNA that targets the *orf120* transcript was selected for genetic engineering of ΦKZ. All plaque assays were replicated three times yielding similar results.

Figure 2. Screening for ΦKZ recombinants by CRISPR-Cas13a counter selection.

(A) Schematic of WT ΦKZ and recombinant genomes at the editing site. The *acrVIA1* gene, shown as a red rectangle, was inserted downstream of *orf120*, with up- and downstream of homology arms (H) indicated by blue rectangles. Green stripes represent synonymous mutations that were introduced to the homology region to prevent crRNA targeting for recombinant phages. F and R indicate forward and reverse primers, respectively, being used to confirm the insertion of *acrVIA1*. (B) Recombinant phages were screened by PCR using primer F1 and R. (C) Recombinant phage plaques underwent 3 rounds of purification and were further confirmed by PCR using primer F2 and R. (D) Plaque assays showing robust anti-CRISPR activities acquired by recombinant ΦKZ against distinct crRNAs, owing to the successful expression and execution of the incorporated AcrVIA1. (E) Workflow of phage genome engineering using CRISPR-Cas13a. The strategy can be divided into three steps. In the first step, an editing plasmid is constructed to introduce desired genetic modifications, such as insertion, deletion, and tag-fusion, and *acrVIA1* flanked by up- and downstream homology arms matching the phage genome. This plasmid is transformed into a bacterial host strain, referred to as the editing strain, followed by infection by wild-type phages. In the second step, the infection culture is harvested, yielding a mixed phage lysate, containing WT phages, recombinants, and escaper mutants. In the last step, the lysate is plated on the selection strain harboring Cas13a on the bacterial chromosome and the most effective gRNA targeting WT phages. Recombinants and escaper mutants will be recovered to form visible plaques on the cell lawn, while WT phages are eliminated by CRISPR-Cas13a via cleavage of targeted phage mRNA. Recombinants are screened by PCR using appropriate primers and further confirmed by sequencing. Remarkably, AcrVIA1 produced by recombinant phages enables ineffectiveness of Cas13a regardless of any gRNA. Accordingly, the same guide can be used for any genomic manipulation of the same phage strain, greatly reducing the effort on seeking for efficient gRNAs to edit distinct genomic loci.

Figure 3. Absence of PhuZ causes mispositioning of the phage nucleus.

(A) Left: a representative cell infected by WT ΦKZ . Right: a representative cell infected by $\Delta phuZ$ mutant. Phage DNA is stained by DAPI and shown as blue signals. The phage nucleus is mispositioned near the cell polar region upon infection by $\Delta phuZ$, in contrast to the WT infection where the phage nucleus is centered. Scale bar denotes 2 μm . (B) Distribution of subcellular location of the phage nucleus in PAO1 cells infected by WT (blue, N = 521) and $\Delta phuZ$ (red, N = 503), and a PAO1 strain expressing wild-type PhuZ *in trans* (p-PhuZ) and infected by $\Delta phuZ$ (light green, N = 573). The diagram of an infected cell is shown on the top. The phage nucleus location is defined as the relative distance between the cell center and the nucleus center. (C) The phage nucleus location is plotted against the cell length for WT and $\Delta phuZ$. The phage nucleus position in $\Delta phuZ$ -infected cells positively correlates with the cell length with a *Pearson* correlation coefficient of 0.486, $p < 0.001$, whereas there is no such correlation for WT infection with a *Pearson* correlation coefficient of 0.029, $p = 0.505$. (D) Efficiency of plaquing of WT and $\Delta phuZ$ on representative *P. aeruginosa* clinical strains. (The full panel of plaque assays is presented in Figure S3)

Figure 4. Chromosomal fluorescent labeling of an inner body protein of ΦKZ.

(A) Visualization of individual phage particles under the fluorescence microscope. *orf39* encoding one of the major components of ΦKZ inner body is genetically fused to mNeonGreen. Each mutant phage particle is visible as a green focus (left), due to the packaging of gp93-mNeonGreen in the capsid. ΦKZ genomic DNA is labeled by DAPI (right). mNeonGreen and DAPI signals colocalize very well and individual virions are easily distinguishable. (B) Overlay images (phase-contrast and fluorescent channels) from a time-lapse movie depicting a representative PAO1 cell being infected by a gp93-mNeonGreen mutant phage. gp93-mNeonGreen and phage DNA are shown as green and blue signals, respectively. Red arrows point to newly assembled phage capsids, both originated on the cell inner membrane (20 min) and anchored on the phage nucleus surface for subsequent phage DNA packaging (25, 45, and 75 min). (C) Overlay images from a time-lapse movie showing a PAO1 cell being infected by a WT ΦKZ loaded with gp93-mNeonGreen fusions. Packaged gp93-mNeonGreen appears as a green focus and remains bound to phage DNA throughout the infection cycle. Scale bar denotes 2 μm.

Figure 5. Genetic engineering of therapeutic jumbo phage OMKO1.

(A) Schematic of genomes of two engineered OMKO1 variants where indicated gene fragments are integrated into the WT genome. OMKO1::Acr, *acrVIA1* is inserted upstream of the shell gene. OMKO1::Acr-BC, *acrVIA1* is inserted together with a barcode sequence (BC) downstream of the major capsid gene. (B) Plaque assays of WT OMKO1 and mutants. Both engineered variants exhibit robust resistance against distinct CRISPR-Cas13a crRNAs, suggesting that the incorporated AcrVIA1 is successfully functioning. (C) Determination of host range of WT OMKO1 and mutants on representative *P. aeruginosa* clinical strains by microplate liquid assay at MOI of 0.01 and 1. Data are presented as the mean liquid assay scores across three independent experiments. Asterisks (*) indicate significant difference between WT and mutants as determined by Students' T-test ($p < 0.05$). The color intensity of each phage-host combination reflects the liquid assay score, with the darker color the stronger intensity displaying a greater score. (The full table of plaque assays is shown in Figure S5)

Figure 6. Genetic engineering of Podophage PaMx41.

(A) Plaque assays showing PaMx41 resistance to a broad variety of DNA-targeting CRISPR-Cas systems. (B) PaMx41 exhibits significant sensitivity to CRISPR-Cas13a. gRNA#5 highlighted in the red frame has been used for PaMx41 genome engineering. (C) PaMx41 $\Delta orf24$ mutant strain is resistant to diverse crRNAs of CRISPR-Cas13a, due to the expression of AcrVIA1 from the phage genome.

Table 1. Summary of phage mutants engineered by CRISPR-Cas13a.

Phage	Gene No. /Genomic site	Identified protein	Modification	Plaque# ¹	screening%	Isolated or not?
ΦKZ	Upstream of <i>orf54</i>	-	Insertion	17	41.2%	Yes
	Downstream of <i>orf120</i>	-	Insertion	16	50.0%	
	<i>orf39</i>	PhuZ	Deletion	17	52.9%	
	<i>orf93</i>	Inner body protein	Deletion	20	20.0%	
			gp93-mNeonGreen fusion	22	18.2%	
	<i>orf241</i>	Hypothetical	Deletion	23	26.1%	
	<i>orf241, orf242</i>	Hypothetical	Double deletions	12	41.7%	No
	<i>orf54</i>	Shell	Deletion	15	20.0%	
			mCherry-gp54 fusion	24	8.3%	
			gp54-mCherry fusion	11	9.1%	
			gfp11-gp54 fusion	57	7.0%	
	<i>orf89, orf90, orf91, orf92, orf93</i>	Head structural proteins	Multiple deletions	17	11.8%	
	<i>orf146</i>	Structural protein	Deletion	86	0.0%	
OMKO1	Downstream of the capsid gene	-	Insertion with a barcode	24	70.8%	Yes
	Upstream of the shell gene	-	Insertion	12	50.0%	
PaMx41	<i>orf24</i>	Hypothetical	Deletion	8	100.0%	Yes

¹Number of plaques analyzed by PCR to screen for recombinants.

References

1. Kortright, K.E., Chan, B.K., Koff, J.L., and Turner, P.E. (2019). Phage Therapy: A Renewed Approach to Combat Antibiotic-Resistant Bacteria. *Cell Host Microbe* 25, 219-232.
2. Pires, D.P., Cleto, S., Sillankorva, S., Azeredo, J., and Lu, T.K. (2016). Genetically Engineered Phages: a Review of Advances over the Last Decade. *Microbiol Mol Biol Rev* 80, 523-543.
3. Doss, J., Culbertson, K., Hahn, D., Camacho, J., and Barekzi, N. (2017). A Review of Phage Therapy against Bacterial Pathogens of Aquatic and Terrestrial Organisms. *Viruses* 9.
4. Nobrega, F.L., Costa, A.R., Kluskens, L.D., and Azeredo, J. (2015). Revisiting phage therapy: new applications for old resources. *Trends Microbiol* 23, 185-191.
5. Lusiak-Szelachowska, M., Zaczek, M., Weber-Dabrowska, B., Miedzybrodzki, R., Klak, M., Fortuna, W., Letkiewicz, S., Rogoz, P., Szufnarowski, K., Jonczyk-Matysiak, E., et al. (2014). Phage Neutralization by Sera of Patients Receiving Phage Therapy. *Viral Immunol* 27, 295-304.
6. Weber-Dabrowska, B., Jonczyk-Matysiak, E., Zaczek, M., Lobočka, M., Lusiak-Szelachowska, M., and Gorski, A. (2016). Bacteriophage Procurement for Therapeutic Purposes. *Front Microbiol* 7, 1177.
7. Lu, T.K., and Koeris, M.S. (2011). The next generation of bacteriophage therapy. *Curr Opin Microbiol* 14, 524-531.
8. Lenneman, B.R., Fernbach, J., Loessner, M.J., Lu, T.K., and Kilcher, S. (2020). Enhancing phage therapy through synthetic biology and genome engineering. *Curr Opin Biotechnol* 68, 151-159.
9. Ando, H., Lemire, S., Pires, D.P., and Lu, T.K. (2015). Engineering Modular Viral Scaffolds for Targeted Bacterial Population Editing. *Cell Syst* 1, 187-196.
10. Mahichi, F., Synnott, A.J., Yamamichi, K., Osada, T., and Tanji, Y. (2009). Site-specific recombination of T2 phage using IP008 long tail fiber genes provides a targeted method for expanding host range while retaining lytic activity. *FEMS Microbiol Lett* 295, 211-217.
11. Matsuda, T., Freeman, T.A., Hilbert, D.W., Duff, M., Fuortes, M., Stapleton, P.P., and Daly, J.M. (2005). Lysis-deficient bacteriophage therapy decreases endotoxin and inflammatory mediator release and improves survival in a murine peritonitis model. *Surgery* 137, 639-646.
12. Monteiro, R., Pires, D.P., Costa, A.R., and Azeredo, J. (2019). Phage Therapy: Going Temperate? *Trends Microbiol* 27, 368-378.
13. Kilcher, S., and Loessner, M.J. (2019). Engineering Bacteriophages as Versatile Biologics. *Trends Microbiol* 27, 355-367.
14. Marinelli, L.J., Hatfull, G.F., and Piuri, M. (2012). Recombineering: A powerful tool for modification of bacteriophage genomes. *Bacteriophage* 2, 5-14.
15. Deveau, H., Garneau, J.E., and Moineau, S. (2010). CRISPR/Cas system and its role in phage-bacteria interactions. *Annu Rev Microbiol* 64, 475-493.
16. Hille, F., Richter, H., Wong, S.P., Bratovic, M., Ressel, S., and Charpentier, E. (2018). The Biology of CRISPR-Cas: Backward and Forward. *Cell* 172, 1239-1259.
17. Mayo-Munoz, D., He, F., Jorgensen, J.B., Madsen, P.K., Bhoobalan-Chitty, Y., and Peng, X. (2018). Anti-CRISPR-Based and CRISPR-Based Genome Editing of *Sulfolobus islandicus* Rod-Shaped Virus 2. *Viruses* 10.
18. Samson, J.E., Magadan, A.H., Sabri, M., and Moineau, S. (2013). Revenge of the phages: defeating bacterial defences. *Nat Rev Microbiol* 11, 675-687.
19. Malone, L.M., Birkholz, N., and Fineran, P.C. (2020). Conquering CRISPR: how phages overcome bacterial adaptive immunity. *Curr Opin Biotechnol* 68, 30-36.

20. Mendoza, S.D., Nieweglowska, E.S., Govindarajan, S., Leon, L.M., Berry, J.D., Tiwari, A., Chaikerasak, V., Pogliano, J., Agard, D.A., and Bondy-Denomy, J. (2020). A bacteriophage nucleus-like compartment shields DNA from CRISPR nucleases. *Nature* 577, 244-248.
21. Malone, L.M., Warring, S.L., Jackson, S.A., Warnecke, C., Gardner, P.P., Gummy, L.F., and Fineran, P.C. (2020). A jumbo phage that forms a nucleus-like structure evades CRISPR-Cas DNA targeting but is vulnerable to type III RNA-based immunity. *Nat Microbiol* 5, 48-55.
22. Guan, J., and Bondy-Denomy, J. (2020). Intracellular Organization by Jumbo Bacteriophages. *J Bacteriol* 203.
23. Abudayyeh, O.O., Gootenberg, J.S., Konermann, S., Joung, J., Slaymaker, I.M., Cox, D.B., Shmakov, S., Makarova, K.S., Semenova, E., Minakhin, L., et al. (2016). C2c2 is a single-component programmable RNA-guided RNA-targeting CRISPR effector. *Science* 353, aaf5573.
24. Meeske, A.J., Jia, N., Cassel, A.K., Kozlova, A., Liao, J., Wiedmann, M., Patel, D.J., and Marraffini, L.A. (2020). A phage-encoded anti-CRISPR enables complete evasion of type VI-A CRISPR-Cas immunity. *Science* 369, 54-59.
25. L, M.I., Anantharaman, V., Krishnan, A., Burroughs, A.M., and Aravind, L. (2021). Jumbo Phages: A Comparative Genomic Overview of Core Functions and Adaptions for Biological Conflicts. *Viruses* 13.
26. Al-Shayeb, B., Sachdeva, R., Chen, L.X., Ward, F., Munk, P., Devoto, A., Castelle, C.J., Olm, M.R., Bouma-Gregson, K., Amano, Y., et al. (2020). Clades of huge phages from across Earth's ecosystems. *Nature* 578, 425-431.
27. Aylett, C.H., Izore, T., Amos, L.A., and Lowe, J. (2013). Structure of the tubulin/FtsZ-like protein TubZ from *Pseudomonas* bacteriophage PhiKZ. *J Mol Biol* 425, 2164-2173.
28. Chaikerasak, V., Nguyen, K., Egan, M.E., Erb, M.L., Vavilina, A., and Pogliano, J. (2017). The Phage Nucleus and Tubulin Spindle Are Conserved among Large *Pseudomonas* Phages. *Cell Rep* 20, 1563-1571.
29. Chaikerasak, V., Khanna, K., Nguyen, K.T., Sugie, J., Egan, M.E., Erb, M.L., Vavilina, A., Nonejuie, P., Nieweglowska, E., Pogliano, K., et al. (2019). Viral Capsid Trafficking along Treadmilling Tubulin Filaments in Bacteria. *Cell* 177, 1771-1780 e1712.
30. Kraemer, J.A., Erb, M.L., Waddling, C.A., Montabana, E.A., Zehr, E.A., Wang, H., Nguyen, K., Pham, D.S., Agard, D.A., and Pogliano, J. (2012). A phage tubulin assembles dynamic filaments by an atypical mechanism to center viral DNA within the host cell. *Cell* 149, 1488-1499.
31. Chaikerasak, V., Nguyen, K., Khanna, K., Brilot, A.F., Erb, M.L., Coker, J.K., Vavilina, A., Newton, G.L., Buschauer, R., Pogliano, K., et al. (2017). Assembly of a nucleus-like structure during viral replication in bacteria. *Science* 355, 194-197.
32. Wu, W., Thomas, J.A., Cheng, N., Black, L.W., and Steven, A.C. (2012). Bubblegrams reveal the inner body of bacteriophage phiKZ. *Science* 335, 182.
33. Thomas, J.A., Weintraub, S.T., Wu, W., Winkler, D.C., Cheng, N., Steven, A.C., and Black, L.W. (2012). Extensive proteolysis of head and inner body proteins by a morphogenetic protease in the giant *Pseudomonas aeruginosa* phage phiKZ. *Mol Microbiol* 84, 324-339.
34. Chan, B.K., Turner, P.E., Kim, S., Mojibian, H.R., Eleftheriades, J.A., and Narayan, D. (2018). Phage treatment of an aortic graft infected with *Pseudomonas aeruginosa*. *Evol Med Public Health* 2018, 60-66.
35. Sepulveda-Robles, O., Kameyama, L., and Guarneros, G. (2012). High diversity and novel species of *Pseudomonas aeruginosa* bacteriophages. *Appl Environ Microbiol* 78, 4510-4515.

36. Cruz-Plancarte, I., Cazares, A., and Guarneros, G. (2016). Genomic and Transcriptional Mapping of PaMx41, Archetype of a New Lineage of Bacteriophages Infecting *Pseudomonas aeruginosa*. *Appl Environ Microbiol* 82, 6541-6547.
37. Skennerton, C.T., Angly, F.E., Breitbart, M., Bragg, L., He, S., McMahon, K.D., Hugenholtz, P., and Tyson, G.W. (2011). Phage encoded H-NS: a potential achilles heel in the bacterial defence system. *PLoS One* 6, e20095.
38. Pul, U., Wurm, R., Arslan, Z., Geissen, R., Hofmann, N., and Wagner, R. (2010). Identification and characterization of *E. coli* CRISPR-cas promoters and their silencing by H-NS. *Mol Microbiol* 75, 1495-1512.
39. Hampton, H.G., Watson, B.N.J., and Fineran, P.C. (2020). The arms race between bacteria and their phage foes. *Nature* 577, 327-336.
40. Vlot, M., Houkes, J., Lochs, S.J.A., Swarts, D.C., Zheng, P., Kunne, T., Mohanraju, P., Anders, C., Jinek, M., van der Oost, J., et al. (2018). Bacteriophage DNA glucosylation impairs target DNA binding by type I and II but not by type V CRISPR-Cas effector complexes. *Nucleic Acids Res* 46, 873-885.
41. Bryson, A.L., Hwang, Y., Sherrill-Mix, S., Wu, G.D., Lewis, J.D., Black, L., Clark, T.A., and Bushman, F.D. (2015). Covalent Modification of Bacteriophage T4 DNA Inhibits CRISPR-Cas9. *mBio* 6, e00648.
42. Liu, Y., Dai, L., Dong, J., Chen, C., Zhu, J., Rao, V.B., and Tao, P. (2020). Covalent Modifications of the Bacteriophage Genome Confer a Degree of Resistance to Bacterial CRISPR Systems. *J Virol* 94.
43. Davidson, A.R., Lu, W.T., Stanley, S.Y., Wang, J., Mejdani, M., Trost, C.N., Hicks, B.T., Lee, J., and Sontheimer, E.J. (2020). Anti-CRISPRs: Protein Inhibitors of CRISPR-Cas Systems. *Annu Rev Biochem* 89, 309-332.
44. Krylov, V.N., Smirnova, T.A., Minenkova, I.B., Plotnikova, T.G., Zhazikov, I.Z., and Khrenova, E.A. (1984). *Pseudomonas* bacteriophage phi KZ contains an inner body in its capsid. *Can J Microbiol* 30, 758-762.
45. van Beljouw, S.P.B., Haagsma, A.C., Rodriguez-Molina, A., van den Berg, D.F., Vink, J.N.A., and Brouns, S.J.J. (2021). The gRAMP CRISPR-Cas effector is an RNA endonuclease complexed with a caspase-like peptidase. *Science* 373, 1349-1353.
46. Ozcan, A., Krajcski, R., Ioannidi, E., Lee, B., Gardner, A., Makarova, K.S., Koonin, E.V., Abudayyeh, O.O., and Gootenberg, J.S. (2021). Programmable RNA targeting with the single-protein CRISPR effector Cas7-11. *Nature* 597, 720-725.

Supplementary Information

Figure S1. Plaque efficiency assays of distinct crRNAs of CRISPR-Cas13a targeting transcripts of diverse Φ KZ genes.

Different crRNAs show varied degrees of Cas13a interference efficiency against Φ KZ. NT, non-targeting.

Figure S2. Sequence alignment of wild type ΦKZ and three escaper mutants at the engineered genomic site.

Escaper mutants were isolated and verified by PCR and sequencing. The WT *orf120* sequence is highlighted in blue and the downstream region is highlighted in grey. The stop codon (TGA) of *orf120* is highlighted in green and Escaper #3 reconstitutes it to TAG. The sequence in the red frame matches the spacer sequence of the crRNA that was used to target and eliminate WT phages. Deletions were indicated by dashed lines and their corresponding numbers of absent base pairs.

Figure S3. Determination of host range of ΦKZ ΔphuZ and Δorf93 mutants by plaque assay on *P. aeruginosa* clinical strains.

Figure S4. Failure of genetic editing the shell gene (*orf54*) in Φ KZ.

(A) Schematic of genomes of WT Φ KZ and three mutated *orf54* variants, “ Δ *orf54*”, “*FP-orf54*”, and “*orf54-FP*”, at the editing site. *orf54*, *acrVIA1*, and fluorescent protein (FP) are shown as blue, red, and green rectangles, respectively. F and R indicate forward and reverse primers, respectively, for PCR confirmation of *orf54* engineering. (B) PCR confirmation of the indicated *orf54* mutants using their corresponding pair of primers. All three mutants generated multiple bands, including a band in the same size as the single band produced by WT. (C) Genome alignment of WT phage with the isolated *orf54* “pseudo knock-out” mutant (“ Δ *orf54*”). A gene cluster of ~ 7 kbp (*orf206* - *orf216*) was missing in the mutant, likely as a result of phage packaging capacity. The majority of the editing plasmid used to generate recombinants was at the editing site, leaving the *orf54* intact.

Figure S5. Host range assay of engineered OMKO1 variants.

Host ranges were determined by microplate liquid assay at MOI of 0.01 and 1 on 22 *P. aeruginosa* clinical strains. The values are presented as the mean liquid assay scores across three independent experiments. Asterisks (*) indicate significant difference between WT and engineered strains as determined by Students' T-test ($p < 0.05$). The color intensity of each phage-host combination reflects the liquid assay score, with the darker color the stronger intensity displaying a greater score.

693 **Movie S1. Time-lapse movie of a *P. aeruginosa* cell infected by a Φ KZ Δ *phuZ* mutant phage.**

694 Phage DNA is stained by DAPI and shown in blue.

695

696 **Movie S2. Time-lapse movie of a *P. aeruginosa* cell infected by a Φ KZ gp93-mNeonGreen**
697 **mutant phage.**

698 Phage DNA is stained by DAPI and shown in blue. gp93-mNeonGreen is shown in green.

699

700 **Movie S3. Time-lapse movie of a *P. aeruginosa* cell infected by a WT Φ KZ packaged with**
701 **gp93-mNeonGreen fusion proteins in the capsid.**

702 Phage DNA is stained by DAPI and shown in blue. gp93-mNeonGreen is shown in green.

Table S1. Bacterial strains and phages used in this work.

Name	Description	Source/Reference
Bacterial strains		
PAO1	<i>P. aeruginosa</i>	Lab collection
SDM084	PAO1 <i>tn7::Lse cas13a</i>	[20]
BF#	<i>P. aeruginosa</i> CF clinical strains	Paul Turner Lab
Phages		
ΦKZ		Alan Davidson Lab
JBD30		Alan Davidson Lab
OMKO1	Clinical <i>P. aeruginosa</i> phage	Paul Turner Lab
PaMx41		Gabriel Guarneros Peña at Centro de Investigación y de Estudios Avanzados (KU884563)

Table S2. crRNA sequences.

Phage	Target	Type	No.	Sequence (5' - 3')
JBD30	orf38	VI-A		acgccaatctctgctacgacggtc
ΦKZ	orf54	VI-A	#1	tcaaactcaagctactgctcctgc
			#2	atgctcggttcagcagatgccctac
			#3	ggtatggctggctctactcattgtt
	orf120		#1	cttaagccgtgagctggaaaggggt
			#2	gctgatcttaagccgtgagctgga
			#3	agcttgctgatcttaagccgtgag
	orf146		#1	ttcatactgggttgaaactaatga
			#2	ggcttctgcattgaagattatgt
PaMx41	orf11	VI-A	#1	ccaccctgaatcctcaagaactca
			#2	agactggtcgtatctcggctgcta
			#3	aaagagcgcgtatatctgaccgcc
			#4	gtcaagcgtcagactggctgatatc
			#5	tcaatcttaccgtcaagcgtcaga
			#6	ctcggctgctatcatcgctgcgaa
	orf04 orf28	I-C	#1	tgatacgaaggaaattgtcgcaattggcgataga
			#2	atttctgactttaatagggaggccgaattccata
	orf14 orf42	I-F	#1	tctcagcgtaatagtctgtttcttctggcag
			#2	ttcttctatttctccagaatctttccgcaat
	orf06 orf20	II-A	#1	tttattactgttgaccctgc
			#2	tttcagcatagttctatac
	orf23 orf46	V	#1	tggagaatgaagaaggaagga
			#2	aqaatatgccttgcataggataq

Table S3. Primers.

No.	Description	Name	Sequence (5' - 3')
JG003	mNeonGreen	mGreen-F	CCAATATTCTCGAGGGTGGCGGTGGCTCGATGGTGAGCAAGGGCG
JG004		mGreen-pHERD-R	GACTCTAGAGGATCCCCGGGTACCTTACTTGTACAGCTCGTCC
JG010	gp93	IB3-SacI-F	AGCATC GAATTCGAGCTCC ATGTCTCTACTTAAAATGCT
JG011		IB3-XhoI-R	AGCATC CTCGAG CTTCTCATCATTCTCAGGTG
JG075	H1 for insertion at gp120 3'UTR	pHERD-gp120H1-F	CCATGGGATCTGATAAGAATTC CCTAGG GTAATGCGTCCGACCTATCG
JG064		gp120-muH1stop-R	GATCATTTTTTATTCTACGTAGG CTCGAG CTAAGGTTTTAAGTCGGCAAGCTTACC
JG029	H2 for insertion at gp120 3'UTR	gp120-H2-KpnI-F	AGCATC GGTACC CATATG GCTGGAAAGGGTTACCGTTA
JG030		gp120-H2-Sall-R	AGCATC GTCGAC CAACAAAACGTGAGACAGGG
JG046	Check insertion at gp120 3' UTR	CheckFP-inphage-R	AGTGCTTCTTCTTCTGGACTAAGTC
JG047		CheckFP-inphage-F	GTGGTACTATCAGCCTATCCAAAGG
JG065	AcrVIA1	AcrVIA1-XhoI-F	AGGCCC CTCGAG CCTACGTAGGAATAAAAA ATGATCTACTATATAAAAG
JG066		AcrVIA1-NdeI-R	CGGCCC CATATG TTAATTTAGCTCCTCTTTTA
JG088	Check the existence of AcrVIA1	AcrVIA1-mid-F	TCCGTTCGCTCCCGATACGA
JG095	H1 for mCherry-gp54 fusion	pHERD-Op24NheI-F	GCCTTTTGCTGGCCTTTTGCTCACATAAG TTGAAGACCCGCCACTATATTTAGG
JG096		Op24H1-AcrVIA-R	CACCTTTTAAATCTTTTATATAGTAGATCAT TTAAATACCTTTACGATTATGG
JG097	AcrVIA1 for mCherry-gp54 fusion	Op24H1-AcrVIA-F	CAAGAAACCATAATCGTAAAGGTATTTAAA ATGATCTACTATATAAAAGA
JG098		AcrVIA-mCherry-R	CACCATTTTTTATTCTACGTAGGCCTAGG TTAATTTAGCTCCTCTTTTA
JG099	H2 for mCherry-gp54 fusion	AcrVIA-mCherry-F	AATTAACCTAGGCCTACGTAGGAATAAAAA ATGGTGAGCAAGGGCGAGGA
JG100		pHERD-gp54NheI-R	TGCTCTGCGAGGCTGGCCGATAAGCTAG ATGTGGGTCGCAATAGAGAACGGAG
JG104	H1 for gp54-gfp11 fusion	pHERD-gp54end-F	GCCTTTTGCTGGCCTTTTGCTCACATAAG GAGCAAGTCATCCTAGATGCA
JG105		gp54end-gfp11-R	CATGTGGTCACGCGAGCCGCCCGACCCGCC GTACCAGGTACCCGGTGCAT

JG106	H2 for gp39 deletion	KO39H2-F	ATTAACCTTATACTGGAGCCCTTCGG
JG107		pHERD-KO39H2-R	TGCTCTGCGAGGCTGGCCGATAAGCTAG CTTTTTAGCAGTGTTCTTTGCC
JG050	Check gp54 editing	gp54-mid-R	CAGTGGTCGGAGTCCAATGTAGATC
JG108		Check-recomb54-F	AGTTTGAGGACCAAGTGTAACACC
JG118		Check-KO54-R	GTGGGTTAATTAGGCTACGTACGTG
JG110	gp54	pHERD-gp54NheI-F	GCCTTTTGCTGGCCTTTTGCTCACATAAG CTGCATTTTATAAAAATACTG
JG111		pHERD-gp54NheI-R	TGCTCTGCGAGGCTGGCCGATAAGCTAG TTAGTACCAGGTACCCGGT
JG127		gp54-PstI-F	AGCATC CTGCAG ATGGCTGTTAACGAAAACGA
JG128		gp54-SacI-R	AGCGTC GAGCTC TTAGTACCAGGTACCCGGTGCAATTATAG
JG112	AcrVIA1 for gp54 deletion (pair with JG097)	AcrVIA-Op24H2-R	TAACATCCCTATCTACCGAGGTATTTCTTA TTAATTTAGCTCCTCTTTTA
JG113	H2 for gp54 deletion	AcrVIA-Op24H2-F	GCAAACAAATTTTAAAAGAGGAGCTAAATTAA TAGGAAATACCTCGGTAGATAGG
JG114		pHERD-Op24H2NheI-R	TGCTCTGCGAGGCTGGCCGATAAGCTAG CTTTCTCATCTGAGTTAC
JG119	H2 for gp93 deletion	KO93H2-F	CAAACATAATGAGGAACCTTCGG
JG120		pHERD-KO93H2-R	TGCTCTGCGAGGCTGGCCGATAAGCTAG CATAAAAGCAGCTGATTGTTA
JG121	Check gp93 deletion	Check-KOgp93-R	AGTTAGTACACGCTGTGCCGCT
JG122	gp39	pHERD-gp39-NheI-F	GCCTTTTGCTGGCCTTTTGCTCACATAAG ATTATATTAATCACAATGAGG
JG123		pHERD-gp39-NheI-R	TGCTCTGCGAGGCTGGCCGATAAGCTAG TTAATCGAATACAAGACCACTACT
JG124	H1 for gp241-gp242 deletion	pHERD-KO2412-H1Acr-F	GCCTTTTGCTGGCCTTTTGCTCACATAAG TTAAATAAAGATAACTTCTTTTC
JG125		KO2412-H1Acr-R	CCTTTTAGTTAATTTAATTTAGCTCCTCTTTTAAAATTTG
JG126	Check gp241-gp242 deletion	Check-KO2412-R	GGCTAAGTCTTTTTCTCGATACTGG
JG133	Operon#24	pHERD-Op24NheI-F	GCCTTTTGCTGGCCTTTTGCTCACATAAG TACGTGACGGAGCATTCTTAAATA
JG134		pHERD-Op24NheI-R	TGCTCTGCGAGGCTGGCCGATAAGCTAG CCCGAAGGAGCCGCTATATCAG

JG135	H1 for insertion at the 3'UTR of the capsid gene in OMKO1	Omk-capH1-GibsonF	GCCTTTTGCTGGCCTTTTGCTCACATAAG ATCGAAGGCGCACTATCCGCTG
JG136		Omk-capmuH1-GibsonR	GATCATTTTTTTATTCCCTACGTAGG GCTAGC CTAAGGTTTTTAAGTCGGCAAGCTTACC
JG137	H1 for insertion at gp54 5'UTR	pHERD-Op24lnH1-F	GCCTTTTGCTGGCCTTTTGCTCACATAAG GGAGCATTCTTAAATAAATTA
JG138		Op24lnH1-RBS-R	GATCATTGTATATTCTATTTGTAAT CTCGAG CGTAACGAACACTATGTT
JG139	AcrVIA1 for insertion at gp54 5'UTR	gp53RBS-AcrVIA1-F	CTCGAG ATTACAAATAGGAATATACA ATGATCTACTATATAAAAGA
JG140		AcrVIA1-Op24lnH2-R	CGCTATAATCTGCTATCAGCAGAT CCTAGG TTAATTTAGCTCCTCTTTTA
JG141	H2 for insertion at gp54 5'UTR	AcrVIA1-Op24lnH2-F	AGGAGCTAAATTAA CCTAGG ATCTGCTGATAGCAGATTATAGCG
JG142		pHERD-Op24lnH2-R	TGCTCTGCGAGGCTGGCCGATAAGCTAG GTTGTAAACGGTTCCTAG
JG146	H1 for gp146 deletion	pHERD-KOgp146H1-F	GCCTTTTGCTGGCCTTTTGCTCACATAAG AGCTATTGCAAATGCGGTACC
JG147		KOgp146H1-AcrR	TAGATCATTTTTATTTCTTGCTGTATAAGAA CATTATACTACTTCCTCAAGTTCCG
JG148	Check gp146 deletion	Check-gp146edit-R	AATCTATCCCAGTTCAAGCTAAGCC
JG152	AcrVIA1 for gp146 deletion	gp145RBS-AcrVIA1-F	TTCTTATACAGCAAGGAAATAAAA ATGATCTACTATATAAAAGAT
JG153		AcrVIA1-Op63lnH2-R	GGAGGGCAATATGTCTGTT TTAATTTAGCTCCTCTTTTAAAAT
JG154	H2 for gp146 deletion	AcrVIA1-Op63lnH2-F	AAAGAGGAGCTAAATTAA AACGACATATTGCCCTCCCTTCGG
JG155		pHERD-Op63lnH2-R	TGCTCTGCGAGGCTGGCCGATAAGCTAG ATTCCATTTATTATTCAATTAATAGCT
JG156	gp146	pMMB-gp146-GbF	GGAAACAGAATTAATTAAGCTTGCATGC CTGCAG ATGGCTTCTGCATTTGAAGA
JG157		pMMB-gp146-GbR	CCGCCAAAACAGCCAAGCTGAATTC GAGCTC TTAAATAACAGCACCTTTGG
JG160	H2 for gp241 deletion	AcrVIA1-KO241H2-F	CAAACAAATTTTAAAAGAGGAGCTAAATTAATAAGAGCATATGGCCCTCCCTTCGG
JG161		pHERD-KO241H2-R	TGCTCTGCGAGGCTGGCCGATAAGCTAG TAATAGTGGTATTAACCTTGTAG
JG162	Check gp241 deletion	Check-KOgp241-R	CGGGTGTAATCGTACGTAGATCAATCGC
JG166	H2 for gp241-gp242 deletion	KO2412-H2-F	GAGGAGCTAAATTAATTAATAACTAAAAGGTAAAAAATGAATAATC
JG167		pHERD-KO2412-H2-R	TGCTCTGCGAGGCTGGCCGATAAGCTAG CATCCTTATTATGGAAGCTATA

JG168	Check orf24 deletion in PaMx41	Check-PaM24-R	CTCGTGGTGATGTTAACCCTAGGG
JG169		Check-PaM24-F	CAACTCTGAATTCCGAAGTCTGAGGCC
JG170	Check gp93 deletion	Check-KOgp93-F	ACTCTTGTATATCCAATCTGTAGCGG
JG173	gp39	gp39-SacI-F	ACCCATGGGATCTGATAAGAATTCGAGCTC ATGATGTCTAAAGTAAAACTCG
JG174		gp39-KpnI-R	CAGGTCGACTCTAGAGGATCCCCGGGTACC TTAATCGAATACAAGACCACTACT
JG175	H1 for Operon#42 deletion	pHERD-KOop42H1-F	GCCTTTTGCTGGCCTTTTGCTCACATAAGTAACCTACCTTCTTTACTAATGCAGA ATAG
JG176		KOop42H1-R	CTTTTAAATCTTTTATATAGTAGATCAT TTTTCTCTTCCTATATGGGTTTGCAGAG
JG178	gfp1-10	pHERD-gfp10-GbF	ACCCATGGGATCTGATAAGAATTC GAGCTC ATGGTGAGCAAGGGCGAGGA
JG179		pHERD-gfp10-GbR	GGCCAGTGCCAAGCTTGCATGC CTGCAG CTTCTCGTTGGGGTCTTTGCTC
JG180	Check gfp11-gp54 and gp54-gfp11 fusion	gfp11-mid-F	TGGTCCTTCATGAGTATGTAAATGC
JG181	Check gp39 deletion	Check-KOgp39-R	CTTCAGCCCATTTGTCTACCCG
JG182	H1 for gp93-mNeonGreen fusion	gp93-mNGH1-Gb-F	GCCTTTTGCTGGCCTTTTGCTCACATAAG AATCCGAAGACTATTGCAGACTTCA
JG183		gp93-mNGH1-Gb-R	CTATATGGGTTTGCGGTTCTCATTATGTTTGTTA TTAATTGTACAGCTCGTCCAT
JG186	Check Operon#42 deletion	Check-KOop42-F	CACCAATAAAGTCACCAGTACGTCC
JG187	Check gp93-mNeonGreen fusion	gp93-mid-F	GCCCTAGCTGATTGGAATCCTGACG
oSDM 455	pHERD30T-crRNA Version 3	Lse_crRNA_V3_Top	ACTCTCTACTGTTTCTCCATGGTAAGAGACTACCTCTATATGAAAGAGGACTAAAA CCGA
oSDM 456		Lse_crRNA_V3_Bot	CGGGTACCGAGCTCGAATTCGAGACCGTACGTACGTGGTCTCGGTTTTAGTCCTC TTCA
oSDM 457		p30T_TSS-R	ATGGAGAAACAGTAGAGAGTTG

oSDM 458	pHERD30T- crRNA Version 2	p30T_TSS-F	GAATTCGAGCTCGGTACC
oSDM 465		LseCas13 crRNA V2 F	ACTCTCTACTGTTTCTCCATGGTAAGAGACTACCTCTATATGAAAGAGGACTAAAA CCGAGACCACGTACGTACGGTC
oSDM 466		LseCas13 crRNA V2 R	CGGGTACCGAGCTCGAATTCGTTTTGTGGAGATTCATATTCTCCAAGTCTCTTAC CGGAGACCGTACGTACGTGGTC