Transcriptional response of *Candida auris* to the Mrr1 inducers methylglyoxal and benomyl

Amy R. Biermann, Deborah A. Hogan*

Department of Microbiology and Immunology, Geisel School of Medicine at Dartmouth,
Hanover, NH 03755

*To whom correspondence should be addressed

Department of Microbiology and Immunology,

Geisel School of Medicine at Dartmouth

Rm 208 Vail Building, Hanover, NH 03755

E-mail: dhogan@dartmouth.edu

Tel: (603) 646-5372

Abstract

Candida auris is an urgent threat to human health due to its rapid spread in healthcare settings and its repeated development of multidrug resistance. Diseases that put individuals at a higher risk for *C. auris* infection, such as diabetes, kidney failure, or immunocompromising conditions, are associated with elevated levels of methylglyoxal (MG), a reactive dicarbonyl compound derived from several metabolic processes. In other *Candida* species, expression of MG reductase enzymes that catabolize and detoxify MG are controlled by Mrr1, a multidrug resistance-associated transcription factor, and MG induces Mrr1 activity. Here, we used transcriptomics and genetic assays to determine that *C. auris MRR1a* contributes to MG resistance, and that the main Mrr1a targets are an MG reductase and *MDR1*, which encodes a drug efflux protein. The *C. auris* Mrr1a regulon is smaller than Mrr1 regulons described in other species. In addition to MG, benomyl (BEN), a known Mrr1 stimulus, induces *C. auris* Mrr1 activity, and characterization of the *MRR1a*-dependent and independent transcriptional responses revealed substantial overlap in genes that were differentially expressed in response to each compound. Additionally, we found that an *MRR1* allele specific to one *C. auris* phylogenetic clade, clade III, encodes a hyperactive Mrr1 variant, and this activity correlated with higher MG resistance. *C. auris MRR1a* alleles were functional in *Candida lusitanae* and were inducible by BEN, but not by MG, suggesting that the two Mrr1 inducers act via different mechanisms. Together, the data presented in this work contribute to the understanding Mrr1 activity and MG resistance in *C. auris*.

43 Importance

44 *Candida auris* is a fungal pathogen that has spread since its identification in 2009 and is
 45 of concern due to its high incidence of resistance against multiple classes of antifungal drugs. In
 46 other *Candida* species, the transcription factor Mrr1 plays a major role in resistance against
 47 azole antifungals and other toxins. More recently, Mrr1 has been recognized to contribute to
 48 resistance to methylglyoxal (MG), a toxic metabolic byproduct. Here, we show that *C. auris*
 49 *MRR1a*, the closest ortholog to *MRR1* in other species, contributes to resistance to MG, and
 50 that Mrr1a strongly co-regulates expression of *MGD1*, encoding a methylglyoxal reductase
 51 enzyme and *MDR1*, encoding an efflux protein involved in resistance to azole drugs,
 52 antimicrobial peptides and bacterial products. We found that one major clade of *C. auris* has a
 53 constitutively active Mrr1 despite high azole resistance due to other mutations, and that this high
 54 Mrr1a activity correlates with higher MG resistance. Finally, we gain insights into the activities of
 55 MG and another Mrr1 inducer, benomyl, to better understand *C. auris* regulation of phenotypes
 56 relevant *in vivo*.

57

58 Keywords

59 *Candida auris*, *Candida*, methylglyoxal, benomyl, Mrr1, RNA-seq, transcriptomics

60

61 Introduction

62 Although *Candida albicans* has historically been the most prominent *Candida* species
 63 associated with both superficial and invasive fungal infections, worldwide incidence of non-
 64 *albicans Candida* (NAC) species is increasing (1-10). Of particular concern is *Candida auris*,
 65 which the CDC classifies as an urgent threat due to its relatively high frequency of resistance to

multiple different classes of drugs including amphotericin B, echinocandins and azoles (reviewed in (11)). Since its recognition as a novel *Candida* species in 2009, *C. auris*, has been reported in at least 40 countries (12-14). Whole-genome sequencing (WGS) analyses of *C. auris* isolates collected from across the globe indicate the concurrent emergence of four genetically distinct clades (15) with a potential fifth clade defined more recently (16). *C. auris* is thought to primarily colonize the skin (17-19) in addition to a diverse array of body sites, and most clinical isolates to date have been isolated from blood (20). Once *C. auris* has disseminated to the bloodstream, it can cause potentially fatal candidemia which has an estimated global mortality rate ranging from about 30 – 60% (15, 21, 22).

The resistance to azoles in *C. auris* is multifactorial; it has been shown that certain mutations in *ERG11* (15, 23-31) and overproduction of Cdr1p (32-36) contribute to resistance to fluconazole (FLZ). In multiple *Candida* species, the transcriptional regulator Mrr1 also plays a role in FLZ resistance (37-45), and Mayr and colleagues (46) found three *C. auris* homologs of the transcriptional regulator Mrr1, and showed that one of them, *MRR1a*, modestly affected fluconazole resistance. Previously, we demonstrated that in *Candida (Clavispora) lusitaniae*, which is more closely related to *C. auris* relative to other well-studied *Candida* species (12, 47), Mrr1 regulates the expression of *MDR1*, and overexpression of *MDR1* confers resistance to FLZ (40, 48-55), the host antimicrobial peptide histatin-5 (40, 56), bacterially produced phenazines (40), and other toxic compounds (57) in multiple *Candida* species. *C. lusitaniae* Mrr1 also regulates dozens of other genes with two of the most strongly regulated genes encoding methylglyoxal (MG) reductase enzymes, *MGD1* and *MGD2* (37, 40, 58). Mrr1 contributes to *C. lusitaniae* resistance to MG (58), which is a spontaneously formed dicarbonyl electrophile generated as a byproduct of several metabolic processes by all living cells (reviewed in (59)).

Via its carbonyl groups, MG reacts non-enzymatically with biomolecules, which can lead to cellular stress and toxicity (reviewed in (59)). Some of the risk factors (60-69) for candidiasis caused by *C. auris* or other *Candida* spp., such as diabetes (70-72), kidney disease (73-76), or septic shock (77), are associated with elevated MG in human serum. MG resistance across clinical isolates of the same *Candida* species, including *C. auris*, can vary (58).

Through specific regulators, MG and other reactive electrophiles induce stress responses in bacteria (78-80), plants (reviewed in (81)), mammals (reviewed in (82)), and the yeasts *Saccharomyces cerevisiae* (83-87) and *Schizosaccharomyces pombe* (88, 89) at subinhibitory concentrations. We found in *C. lusitanae*, MG induces expression of *MGD1* and *MGD2* as well as *MDR1*, through a mechanism that involved Mrr1 (58), and that MG increased fluconazole (FLZ) resistance. *C. auris* displays nosocomial transmission (61-63, 65-69), in part due to its resistance to high temperatures (90) and common surface antiseptics (91), and persistence on abiotic surfaces including latex and nitrile gloves (92), plastics (90), and axillary temperature probes (93). The factors that control *C. auris* stress resistance are not yet known.

In the present study, we show that *C. auris* *MRR1a* regulates resistance to MG and that MG is an inducer of Mrr1-regulated gene expression. Mrr1a regulates the gene orthologous to the methylglyoxal reductase genes *C. lusitanae* *MGD1* in addition to *MDR1*, which regulates FLZ efflux, but the Mrr1a regulon is smaller than that described for other species. Furthermore, we characterize Mrr1a in both Clade I and Clade III isolates and show that the Mrr1 variant in Clade III is constitutively active. Transcriptomics analysis shows that MG elicits a large transcriptional response that is similar in both Clade I and Clade III, and that there are commonalities in the responses elicited by MG and the Mrr1 inducer benomyl. These data support the model that Mrr1 is a regulator of MG resistance in coordination with efflux proteins

such as Mdr1 and provides the basis for future studies on the roles of Mrr1 and MG in survival of *C. auris* in hospital settings.

Results

Mrr1a regulates expression of orthologs to *MDR1* and *MGD1* in *C. auris* strain B11221 and is involved in MG resistance.

To determine whether the *C. auris* *MRR1* orthologs *MRR1a*, *MRR1b*, and *MRR1c* contributed to resistance to MG, we performed growth kinetic assays in YPD +/- 5 mM, 10 mM, or 15 mM MG. At MG concentrations of 10 mM (**Fig. 1A**) and 15 mM (**Fig. S1**), the *mrr1a*Δ mutant displayed a substantial growth defect relative to WT, while the *mrr1b*Δ and *mrr1c*Δ mutants exhibited growth comparable to WT. None of the mutants (*mrr1a*Δ, *mrr1b*Δ, and *mrr1c*Δ) differed from the parental isolate B11221 (WT) in YPD alone or in the presence of 5 mM MG (**Fig. S1**). Like *C. lusitanae*, the *C. auris* genome encodes multiple putative MG reductases; the closest orthologs to *MGD1* and *MGD2* were *CJI97_000658* and *CJI97_004624*, respectively, in the B11221 genome assembly (58) and we will henceforth refer to these genes as *MGD1* and *MGD2*. For reference, *MGD1* and *MGD2* correspond to B9J08_000656 and B9J08_004828 respectively. By quantitative real-time PCR (qRT-PCR), basal expression of *MGD1* was significantly decreased 24-fold in the *mrr1a*Δ mutant relative to B11221 WT (**Fig. 1B**), and expression of *MGD2* trended lower in the *mrr1a*Δ mutant (~1.2-fold) but this difference did not reach statistical significance (**Fig. 1C**). *MGD1* was also more highly expressed than *MGD2* in the WT B11221 as in *C. lusitanae* (58). Consistent with the transcriptional patterns, *C. auris* Mgd1 shares slightly more identity with *C. lusitanae* Mgd1 than does *C. auris* Mgd2 (63% identity versus 61% identity).

In the *C. auris* B11221 background, expression of *MDR1*, another target of Mrr1 in other species including *C. lusitaniae*, also depended on Mrr1a, as the *mrr1aΔ* mutant exhibited a significant 21-fold decrease in *MDR1* expression compared to the WT parent (**Fig. 1D**). These results indicate that in *C. auris* *MDR1* and *MGD1* are co-regulated, as has been reported in *C. albicans* (44, 45, 94-96), *C. parapsilosis* (97), and *C. lusitaniae* (37, 39, 40, 58, 98) and that higher expression of *MGD1* and/or *MDR1* contributes to growth in high concentrations of MG (**Fig. 1A**).

In *C. lusitaniae* and other *Candida* species, Mrr1 regulates dozens of genes in addition to *MDR1* and *MGD1* (37, 40). To further elucidate the Mrr1a regulon in *C. auris* isolate B11221, we performed an RNA-seq analysis of in B11221 WT and its *mrr1aΔ* derivative in cells from exponential phase cultures grown at 37°C in YPD. In the control condition (YPD + dH₂O), only four genes, including *MDR1* and *MGD1*, were differentially expressed between the two strains with the cutoff of a log₂ fold change (log₂FC) ≥ 1.00 or ≤ -1.00 and a p-value less than 0.05 (**Fig 1E** and **Supplementary File S1** for all data). *MGD1* and *MDR1* showed a 22- and 24-fold decrease, respectively, in *mrr1aΔ* compared to WT, consistent with our qRT-PCR data. *CJI97_005632*, which was 2.25-fold lower in *mrr1aΔ*, is orthologous to the *C. albicans* genes *RIM11* and *C2_04280W_A*, both of which are predicted to encode proteins with serine/threonine kinase activity, though it is worth noting that levels of the transcript were much lower than levels of *MDR1* and *MGD1*. *CJI97_000852*, which was 2.77-fold higher in *mrr1aΔ* than in WT, has 16 orthologs of diverse predicted or known functions in *C. albicans*, including *USO5*, *USO6*, and *RBF1* (**Fig. 1E** and **Supplementary File S1**). Notably, *MGD2* was not differentially expressed between B11221 WT and the *mrr1aΔ* mutant in our RNA-seq data (**Supplementary File S1**), consistent with our qRT-PCR results described above.

158

159 **Mrr1a regulates only *MDR1* and *MGD1* in response to MG and benomyl**

160 We have previously shown in *C. lusitaniae* that MG induces expression of the Mrr1-
 161 regulated genes *MGD1* and *MGD2* in an Mrr1-dependent manner, and *MDR1* in a partially Mrr1-
 162 dependent manner (58). To determine if MG would induce expression of *MGD1*, *MGD2*, and/or
 163 *MDR1* in *C. auris*, we purified RNA for qRT-PCR from exponential-phase cultures of B11221 WT
 164 and *mrr1aΔ* treated with 5 mM MG or an equal volume of dH₂O for 15 minutes. We found that
 165 MG treatment significantly enhanced expression of *MGD1* in WT by 2.4-fold but not in *mrr1aΔ*
 166 (**Fig. 2A**). *MGD1* was also induced by a 30-min treatment with 25 µg/mL benomyl (BEN), a
 167 known inducer of Mrr1-regulated genes in other *Candida* species (37, 41, 43, 95, 99-104), by
 168 7.5-fold in the WT (**Fig. 2A**). The different treatment times for MG and BEN were used to be
 169 consistent with previous studies using either compound in the related species *C. lusitaniae* (37,
 170 58). Expression of *MDR1* was also more highly induced by treatment with either MG or BEN in
 171 WT compared to the *mrr1aΔ* mutant by 6- and 14.5- fold respectively (**Fig. 2B**). Although *MDR1*
 172 expression was significantly induced by MG and BEN in the *mrr1aΔ*, transcript levels of *MDR1*
 173 were approximately 20-fold higher in the WT than in the *mrr1aΔ* under these conditions (**Fig.**
 174 **2B**), suggesting that Mrr1a is required for maximum expression of *MDR1* in response to stimuli.

175 To describe the complete Mrr1-dependent MG- and BEN- response regulon under our
 176 test conditions in *C. auris*, we also performed RNA-seq on exponential-phase cultures of B11221
 177 WT and *mrr1aΔ* treated with MG or BEN as described above. In B11221 WT, MG led to the
 178 upregulation of 319 genes and downregulation of 133 genes compared to the control condition
 179 (**Fig. 2C** and **Supplementary File S2**). In the *mrr1aΔ* mutant, MG led to the upregulation of 349
 180 genes and downregulation of 143 genes compared to the control condition (**Fig. S2A** and

181 **Supplementary File S2).** Consistent with our qRT-PCR data in **Fig. 2A**, MG induced expression
 182 of *MGD1* in the WT but not in the *mrr1a*Δ mutant (**Table S1** and **Supplementary File S2**).
 183 Although expression of *MDR1* was significantly induced by MG in both the WT and the *mrr1a*Δ
 184 mutant (**Table S1** and **Supplementary File S2**), levels of *MDR1* were substantially lower in the
 185 *mrr1a*Δ mutant even in the presence of MG (**Fig. 2D** and **Supplementary File S2**), also in
 186 agreement with our qRT-PCR data. *MGD1* and *MDR1* strongly stood out as the only two genes
 187 in the MG response that were strongly dependent on Mrr1a (**Fig. 2D**).

188 Treatment with BEN led to upregulation of 160 genes and downregulation of 163 genes
 189 in the WT (**Fig. 2E** and **Supplementary File S3**). In the *mrr1a*Δ mutant, 181 genes were
 190 upregulated, and 229 genes were downregulated in response to BEN (**Fig. S2B** and
 191 **Supplementary File S3**). Like MG, induction of *MGD1* by BEN was completely dependent on
 192 Mrr1a (**Table S1** and **Supplementary File S3**) and *MGD2* expression was not induced by BEN
 193 (**Supplementary File S3**). Expression of *MDR1* was also induced by BEN in both the WT and
 194 the *mrr1a*Δ mutant, but as with MG, *MDR1* levels in the *mrr1a*Δ mutant did not reach that of the
 195 WT even with BEN treatment (**Fig. 2F** and **Supplementary File S3**). Again, *MGD1* and *MDR1*,
 196 appear to be the only genes in *C. auris* whose induction of expression by either MG or BEN is
 197 dependent on Mrr1a. The Mrr1a-independent responses to MG and BEN are discussed further
 198 below.

199
 200 **B11221 has higher basal expression of *MDR1* and of putative MG reductase genes**
 201 **compared to the Clade I isolate AR0390.**

202 Many Clade III isolates, including B11221, contain an N647T single nucleotide
 203 polymorphism (SNP) in *MRR1a* (25, 105). In (105), this SNP was proposed to be a gain-of-

function mutation due to the resistance of Clade III isolates against azoffluxin, a novel antifungal compound that inhibits expression and activity of *C. auris* efflux pumps. As a first step to determine if there were activity differences between the Mrr1a variant that was found Clade III strains was different from that encoded by the alleles found in Clade I, II, and IV strains, we compared MG sensitivity of B11221 to that of Clade I isolate AR0390. Interestingly, AR0390 grew substantially better than B11221 in the YPD control but showed a greater reduction in growth in YPD with 5 mM MG than did B11221 (**Fig. S3**). At concentrations of 10 mM (**Fig. 3A**) and 15 mM MG (**Fig. S3**), AR0390 exhibited a profound growth defect compared to B11221. To determine if differences in MG sensitivity were due to differences in *MGD1* expression, we measured basal expression of *MGD1* and its co-regulated gene *MDR1* in B11221 and AR0390 using qRT-PCR. Both genes were significantly more highly expressed in B11221 by 42- and 4.2-fold, respectively (**Fig. 3B-C**).

To gain a deeper understanding of the broader transcriptional differences between B11221 and AR0390, we compared the basal global gene expression in YPD of the two strains using RNA-seq. First, we matched the 5227 syntenic orthologs between the genomes of B11221 and the Clade I reference strain B8441 to compare expression of each gene under the control condition. Of these, 755 genes were differentially expressed between B11221 and AR0390 in the control condition ($|\log_2FC| \geq 1.00$, FDR-corrected $p < 0.05$) (**Fig. 3D, Supplementary File S4**). The top twenty differentially expressed genes whose orthologs have known or predicted functions in *C. albicans* are reported in **Table S3**. Strikingly, the two genes which exhibited the largest difference in expression between B11221 and AR0390 were *MGD2* ($\log_2FC = 11.29$) and *MGD1* ($\log_2FC = 8.53$) (**Fig. 3D, Table S3, and Supplementary File S4**). A third gene with homology to MG reductases, *CJI97_001800/B9J08_002257*, was also more highly expressed

in B11221, although the log₂FC in expression of this gene in B11221 vs AR0390 was only 1.41 (Supplementary File S4). Low expression of *MGD1*, *MGD2*, and/or *B9J08_002257* may contribute to the severe growth defect of AR0390 in the presence of MG. Consistent with our qRT-PCR data, *MDR1* was also significantly more highly expressed in B11221 relative to AR0390 (log₂FC = 4.42) (Fig. 3D and Table S3). Although *MGD2* and *B9J08_002257* do not appear to be regulated by Mrr1a in our studies, it is nonetheless interesting to note the elevated expression of three putative MG reductases in the *MDR1*-overexpressing *C. auris* isolate B11221, as the co-expression of *MDR1* with at least one MG reductase has been reported in numerous studies in other *Candida* species (37, 40, 44, 45, 58, 94-97).

Clade III Mrr1a^{N647T} exhibits a gain-of-function phenotype compared to Clade I Mrr1a when expressed in *C. lusitaniae*.

To compare the activities of the proteins encoded by the *MRR1a* alleles of B11221 and AR0390 more directly, we heterologously expressed each allele, henceforth referred to as *CauMRR1a*^{N647T} and *CauMRR1a* respectively, independently in a *C. lusitaniae* *mrr1*Δ mutant previously generated and characterized by our lab (37, 40, 58). All three *C. lusitaniae* clones expressing *CauMRR1a*^{N647T} which we tested exhibited a four-fold increase in fluconazole (FLZ) MIC relative to the U04 *mrr1*Δ parent (16 μg/mL versus 4 μg/mL, Table S4), confirming that *C. auris* Clade III *MRR1a* can complement *MRR1*-dependent FLZ resistance in *C. lusitaniae* and adding support to the hypothesis that the N647T substitution in Clade III *MRR1a* confers increased activity. However, the FLZ MIC of the three tested *C. lusitaniae* clones expressing *CauMRR1a* did not differ from that of U04 *mrr1*Δ (4 μg/mL; Table S4), so FLZ MIC alone could not indicate whether this allele is functional in *C. lusitaniae*. One clone expressing each *C. auris*

250 *MRR1a* allele was chosen at random for the remaining experiments described in this paper:
 251 clone #1 for *CauMRR1a*^{N647T} and clone #5 for *CauMRR1a*. Using qRT-PCR, we then examined
 252 basal expression levels of *C. lusitaniae* *MGD1* (*CLUG_01281*) and *MDR1*
 253 (*CLUG_01938/CLUG_01939*) in the heterologous complements and the U04 *mrr1Δ* parent.
 254 Complementation with *CauMRR1a*^{N647T} conferred a significant increase in basal expression of
 255 both *MGD1* (**Fig. 3E**) and *MDR1* (**Fig. 3F**) compared to the *mrr1Δ* parent, while complementation
 256 with *CauMRR1a* led to a small, but significant, decrease in expression of both genes relative to
 257 *mrr1Δ* (**Fig. 3E-F**). These results are consistent with our previous observations that *C. lusitaniae*
 258 strains expressing certain Mrr1 variants with low basal activity demonstrate lower expression of
 259 some Mrr1-regulated genes, including *MDR1* and *MGD1*, compared to an isogenic *mrr1Δ* strain
 260 suggesting that Mrr1 has both repressing and activating roles (37, 58). Finally, we assessed the
 261 relative MG resistance of the isogenic *C. lusitaniae* strains expressing *CauMRR1a*^{N647T} or
 262 *CauMRR1a* and the U04 *mrr1Δ* parent. The *CauMRR1a*^{N647T} complement grew markedly better
 263 in 15 mM MG compared to U04 *mrr1Δ* whereas the *CauMRR1a* complement grew substantially
 264 worse than U04 *mrr1Δ* (**Fig. 3G**), consistent with the pattern of *MGD1* expression we observed
 265 in these strains via qRT-PCR. None of the *C. lusitaniae* strains demonstrated growth differences
 266 in the YPD control, or in the presence of MG at concentrations of 5 mM or 10 mM (**Fig. S4**).

267

268 **MG induces expression of *MGD1* and *MDR1* in *C. auris* B11221 and AR0390, but not**
 269 **in *C. lusitaniae* strains expressing *C. auris* *MRR1a* alleles.**

270 Next, we compared induction of *MGD1* and *MDR1* by MG in the *C. auris* strains B11221
 271 and AR0390 via qRT-PCR. MG significantly induced expression of *MGD1* by 2.4-fold in *C. auris*
 272 strain B11221 and by 4.0-fold in *C. auris* strain AR0390 (**Fig. 4A**) and expression of *MDR1* by

6.0-fold in B11221 and 9.3-fold in AR0390 (**Fig. 4B**). AR0390 displayed lower expression of both genes in MG, but a higher fold change compared to B11221, further supporting the hypothesis that the N647T allele is gain-of function.

Finally, we compared induction of *MGD1* and *MDR1* by MG in the isogenic *C. lusitaniae* strains expressing either *CauMRR1a*^{N647T} or *CauMRR1a* and the *mrr1Δ* parent. Additionally, we tested induction by BEN in these strains as a control. While the *mrr1Δ* parent exhibited a significant 1.8- fold induction of *MDR1*, neither *C. lusitaniae* strain expressing a *C. auris* Mrr1a allele demonstrated a significant change in *MGD1* or *MDR1* expression in response to MG (**Fig. 4C-D**), indicating that *C. auris* Mrr1a may repress *MRR1*-independent MG induction of *MDR1* in *C. lusitaniae* and that induction of *MGD1* by MG in *C. lusitaniae* requires a functional *MRR1* allele from its own species. Treatment with BEN led to significant increase in expression of *MGD1* (**Fig. 4E**) and *MDR1* (**Fig. 4F**) in all three *C. lusitaniae* strains. In response to BEN, *MGD1* was induced by 1.9-fold in *mrr1Δ*, 2.9-fold in the *CauMRR1a*^{N647T} complement, and 6.1-fold in the *CauMRR1a* complement (**Fig. 4E**). Likewise, expression of *MDR1* was induced by 2.3-fold in *mrr1Δ*, 3.5-fold in the *CauMRR1a*^{N647T} complement, and 5.0-fold in the *CauMRR1a* complement in response to BEN (**Fig. 4F**). The striking difference in the ability of the *C. lusitaniae* strains expressing *C. auris* *MRR1a* alleles to respond to BEN versus MG suggests that there are differences in the mechanisms by which BEN and MG induce Mrr1-dependent transcriptional activation and that MG induction of *C. auris* Mrr1a is not supported by *C. lusitaniae* factors. These potential differences are a topic of future study and may shed light on mechanisms of Mrr1 activation in *Candida* species.

MG and BEN induced Mrr1a-independent transcriptional responses in *C. auris*

296 We have previously observed heterogeneity in MG resistance as well as MG-induced FLZ
297 resistance among several *C. auris* isolates from different clades (58), and thus we were
298 interested in whether the overall transcriptional response to MG was more similar or different in
299 B11221 and AR0390. AR0390 had greater number of genes differentially expressed by MG
300 compared to B11221; 438 genes were significantly upregulated, and 242 genes were
301 significantly downregulated by MG (see **Fig. S5** for the volcano plot of all genes). More genes
302 had a larger fold change in response to MG in AR0390 compared to B11221, including *MGD1*
303 and *MDR1* (**Fig. 5A**), consistent with the qRT-PCR results in **Fig. 4A-B**. However, there was a
304 large overlap of 254 genes which were induced by MG in both strains (**Fig. 5B**), suggesting a
305 common response across these two genetically distinct clades. These commonly induced genes
306 include many with putative roles in amino acid biosynthesis; transmembrane transport; or
307 acquisition and usage of sulfur (**Fig. 5C** and **Table S5**). The complete comparison is available
308 in **Supplementary File S6**).

309 Only 68 genes with syntenic orthologs across both strains were commonly repressed by
310 MG (**Fig 5B**). These genes include some with putative roles in metal transport or carbohydrate
311 uptake and metabolism (**Fig. 5C** and **Table S6**). We did not observe obvious patterns in genes
312 that were only induced or repressed in one strain, and some genes that are listed as only induced
313 or repressed in one strain were close to the cutoff in the other strain.
314 The groups of genes that were differentially expressed in response to MG in both B11221 and
315 AR0390 were also evident in the response of B11221 to BEN as well as the response of the
316 *mrr1aΔ* mutant in response to MG and BEN. In B11221, a total of 46 genes exhibited significant
317 induction by both MG and BEN, including *MGD1* and *MDR1*. Many of the 44 other genes have
318 predicted roles in assimilation and biosynthesis of sulfur-containing compounds or xenobiotic

transport (**Fig. 5C** and **Table S1**). MG also induced expression of many genes with predicted roles in the biosynthesis of amino acids. The two genes most highly upregulated upon MG treatment, in terms of fold change, in this strain were orthologous to the arginine biosynthesis genes *ARG3* ($\log_2FC = 4.77$) and *ARG1* ($\log_2FC = 4.72$) (**Fig. 2C** and **Table S1**). Conversely, BEN had a limited effect on expression of amino acid biosynthesis genes (**Table S1**). There were also common themes among the genes that were significantly repressed by both MG and BEN in B11221. Genes that were repressed by both MG and BEN included four orthologs of the *HGT* glucose transporter family, five genes with a predicted role in uptake of iron and/or copper, and *ERG6*, which encodes an enzyme in the ergosterol biosynthesis pathway (**Fig. 5C** and **Table S2**). The genes that were repressed by only one stimulus, MG or BEN, also included those involved in ergosterol biosynthesis and the uptake of iron, copper, or glucose (**Fig. 5C**, **Table S2**). In general, the transcriptional response of the *mrr1a* Δ mutant to MG and BEN was similar to that of B11221 WT (**Fig. S2** and **Tables S1-S2**). The complete datasets for MG and BEN responses are available in **Supplementary Files S2** and **S3**, respectively.

Discussion

In this work, we have demonstrated that in *C. auris*, the zinc-cluster transcription factor Mrr1a, which is orthologous to Mrr1 in other *Candida* species, strongly regulates expression of a putative MG reductase *MGD1* in addition to *MDR1*, and that Mrr1a plays a role in MG resistance, highlighting a function of Mrr1 that is distinct from antifungal resistance. We also compared basal global gene expression in B11221 and AR0390 and found that *MDR1*, *MGD1*, and *MGD2* were among the genes significantly more highly expressed in B11221, consistent with the higher MG resistance of this isolate relative to AR0390. These differences were

explained by our finding that *MRR1a* from B11221 encoded a higher activity variant than that from AR0390 as evidenced by a higher FLZ MIC, higher expression of *MDR1* and *MGD1*, and higher MG resistance in the strain expressing *CauMRR1a*^{N647T} compared to the isogenic strain expressing *CauMRR1a*. The allele from B11221, which contains an N647T amino acid substitution (25, 105) which is in the central region of the regulator where other gain of function substitutions have been found. Both alleles result in induction of *MDR1* and *MGD1* in response to BEN but not to MG in *C. lusitaniae*, suggesting that these two compounds activate Mrr1-dependent transcription through different mechanisms.

Under the conditions tested, Mrr1a regulation in the *C. auris* B11221 background was mainly of *MGD1* and *MDR1*. Homologs of *MDR1* and at least one gene encoding a known or predicted MG reductase are co-regulated by Mrr1 in *C. albicans* (44, 45, 94-96), *C. parapsilosis* (97), and *C. lusitaniae* (37, 40, 58), suggesting that the co-regulation of these two genes has been conserved throughout multiple *Candida* species. Gaining a deeper understanding of the evolutionary and biochemical relationship between methylglyoxal reductases and efflux pumps, particularly Mdr1, may shed light on how *Candida* species sense and respond to environmental or physiological stresses, evade host defense mechanisms, and develop antifungal resistance. In all other *Candida* species with published Mrr1 regulons, however, Mrr1 appears to regulate expression of many more genes than the four we have described here in the *C. auris* strain B11221 (37, 40, 44, 45, 97). The surprisingly small number of *C. auris* genes whose expression was significantly altered by genetic deletion of *MRR1a* may be due to possible redundancy between *MRR1a* and the other two *MRR1* orthologs in *C. auris*, *MRR1b* and *MRR1c*, although further studies would be necessary to test this hypothesis. It is striking, however, that *MRR1a* alone seems to be necessary for expression and induction of *MGD1*, which is further supported

365 by our observation that only the *mrr1a*Δ mutant had a growth defect in MG compared to parental
366 B11221 (**Fig. 1A**).

367 Our demonstration of increased basal activity of the *CauMRR1a*^{N647T} allele compared to
368 the allele from AR0390 supports the hypothesis put forth by Iyer et al. that the N647T substitution
369 found in many Clade III isolates is a gain-of-function mutation (105). Furthermore, this may
370 explain why deletion of *MRR1a* leads to a mild decrease in azole resistance in B11221, but not
371 in the Clade IV isolate B11243 (46). In *C. albicans*, knocking out gain-of-function *MRR1* causes
372 a significant decrease in FLZ resistance, but knocking out *MRR1* with wild-type transcriptional
373 activity does not alter FLZ resistance (41, 44, 45, 106). Similarly, knocking out gain-of-function
374 *MRR1* in *C. lusitaniae* also decreases FLZ resistance, although knocking out *MRR1* alleles that
375 do not encode a constitutively active protein generally leads to increased FLZ resistance (37).

376 Although Mrr1a does not appear to play a major role in *C. auris* azole resistance (46), our
377 findings suggest that it contributes to resistance against MG, which may be encountered in the
378 host environment. We have previously shown that Mrr1 also contributes to MG resistance in *C.*
379 *lusitaniae* in a manner that is partially dependent on *MGD1* and *MGD2* (58). Indeed, gain-of-
380 function mutations in *MRR1* may arise in various *Candida* species due to selective pressures
381 other than azoles. In *C. lusitaniae*, we have reported the emergence of gain-of-function
382 mutations in *MRR1* among isolates from a patient with no prior history of clinical antifungal use
383 (40). In *C. auris*, most sequenced clade III isolates exhibit both the *MRR1a*^{N647T} allele and the
384 *ERG11*^{F126L} allele (25), the latter of which has been shown to be a major contributor to azole
385 resistance (31). Although it is not known whether the *MRR1a* or *ERG11* mutation occurred first
386 in the clade III lineage, it seems plausible that if the *ERG11* mutation did occur first, evolution of
387 the *MRR1a*^{N647T} allele in *C. auris* is likely to be the result of selection for *MGD1* expression and/or

an unknown role for Mdr1 that is unrelated to azole resistance. Therefore, we hypothesize that Mrr1 may act, either directly or indirectly, as a response regulator for carbonyl stress in *Candida* species, and future studies will investigate a possible role for Mrr1 in resistance against other physiologically relevant reactive carbonyl compounds.

Curiously, although both variants of *C. auris* Mrr1a were inducible by BEN when expressed in *C. lusitanae*, they were not inducible by MG under the conditions tested (**Fig. 4E-F**). One possible hypothesis for this observation is that Mrr1 must interact with at least one particular binding partner to induce transcription in response to MG, and that *C. auris* Mrr1a does not bind efficiently to this *C. lusitanae* Mrr1-binding protein or complex. Differential requirements for Mrr1-dependent transcriptional activation by chemical stressors have reported in *C. albicans*. For example, the transcription factor Mcm1 is required for Mrr1-dependent induction of *MDR1* in response to BEN but not to H₂O₂ (101), and the redox-sensing transcription factor Cap1 is required for *MDR1* induction by H₂O₂ and may play a role in *MDR1* induction by BEN (44). Furthermore, gain-of-function Mrr1 in *C. albicans* requires the Swi/Snf chromatin remodeling complex to maintain promoter occupancy, and the kinase Ssn3, which is a subunit of the Mediator complex, may act in opposition to Mrr1 or its coactivators (38). Thus, although *C. auris* Mrr1a can complement Mrr1-dependent basal and BEN-induced expression of *MDR1* and *MGD1* in *C. lusitanae*, it may be incompatible with certain elements of the *C. lusitanae* MG-responsive transcriptional machinery. Further studies on the differences between *C. lusitanae* and *C. auris* Mrr1, particularly in the presence of MG, may elucidate more detailed mechanisms of Mrr1 activation.

In general, we observed substantial upregulation of genes with predicted roles in transmembrane transport, sulfur metabolism, and amino acid biosynthesis in response to MG in

all three strains tested. Many genes downregulated in response to MG in all three strains have predicted roles in metal acquisition, particularly iron, and carbohydrate metabolism. In both B11221 WT and *mrr1a*Δ, BEN treatment led to differential expression of similar groups of genes as MG in addition to induction of genes with predicted roles in oxidative stress response. Our studies of the transcriptional response of *C. auris* to MG and BEN contribute to the understanding of how *Candida* species may adapt to oxidative and/or carbonyl stress, two types of stress that a pathogen is likely to encounter in the host environment. In humans, elevated serum MG has been reported in diabetes as well as in renal failure, which are both risk factors for *Candida* infection (107, 108). There is also evidence that neutrophils (109) and macrophages (110, 111) generate MG during the inflammatory response, consistent with elevated levels of MG in sepsis patients (77). In our transcriptomics analysis of three *C. auris* strains exposed to 5 mM MG for 15 min, upregulation of numerous genes involved in amino acid uptake, metabolism, and biosynthesis was one of the most striking responses to MG (**Tables S1** for comparison of MG and BEN and **S5** for the comparison of genes induced by MG in B11221 and/or AR0390). In particular, induction of *ARG* genes is interesting considering the report that *C. albicans* upregulates expression of arginine biosynthesis genes when phagocytosed by macrophages or in response to sublethal concentrations of hydrogen peroxide, tert-butyl hydroperoxide, or menadione *in vitro* (112). This induction of *ARG* genes in *C. albicans* by macrophages is dependent on the *gp91^{phox}* subunit of the macrophage oxidase, and thus is likely a direct response to oxidative stress rather than arginine depletion (112). In our dataset, *ARG3* and *ARG1* exhibited the highest log₂FC in response to MG in the B11221 background, independently of *MRR1a* (**Table S1**). We also observed, in all three *C. auris* strains, induction of several *MET* genes, which are involved in methionine synthesis and are an important branch of sulfur

assimilation in yeast. Other genes involved in sulfur acquisition and assimilation that were induced by MG include the sulfate importer *SUL2*, a gene orthologous to both *CYS3* and *STR3* of *S. cerevisiae*, and numerous genes associated with iron-sulfur cluster formation (**Table S1**). A gene orthologous to *MUP1* of *S. cerevisiae* and *C. albicans* was induced by MG in B11221 WT and AR0390 but fell short of the $\log_2\text{FC} \geq 1.00$ cutoff in *mrr1a* Δ (**Table S1** and **Supplementary File S2**). Induction of genes involved in sulfur metabolism, including the *MET* pathway, *SUL2*, *CYS3*, *STR3*, and *MUP1*, has previously been observed in *Saccharomyces cerevisiae* exposed to 1g/L acetaldehyde (113), another reactive aldehyde metabolite that is structurally similar to MG. Thus, sulfur acquisition and metabolism may be an important part of the carbonyl stress response in yeast.

In the B11221 background, we observed modest overlap in the genes and groups of genes that were up- or down- regulated in response to either MG or BEN. *MDR1* and *MGD1* were among the genes induced by both compounds, and induction of *MGD1* by either MG or BEN was completely dependent on *MRR1a*. Although BEN, which originated as an agricultural fungicide, is widely recognized as an inducer of expression of Mrr1-regulated genes in *Candida* species (37, 41, 43, 95, 99-104), the mechanism by which this induction occurs is not yet known. BEN is thought to cause oxidative stress in yeast (114, 115), which is consistent with our observation of an upregulation of genes with a predicted role in oxidative stress response in BEN-treated *C. auris* cultures (**Table S1**). Additionally, in mammalian cells, BEN exposure has been shown to inhibit aldehyde dehydrogenase enzymes (116-119), which may lead to an accumulation of reactive aldehydes, although this possible mechanism has not yet been investigated in fungi.

We also note similarities between the results of our study of MG- and BEN- treated *C. auris* and the recently published transcriptional analysis of the Clade I *C. auris* strain NCPF 8973 exposed to 75 μ M farnesol (120). In response to farnesol, the authors reported upregulation of many genes with predicted roles in transmembrane transport, such as *MDR1* and *CDR1*, and downregulation of numerous genes predicted to be involved in metal acquisition and homeostasis, including multiple ferric reductases and iron permeases (120). As farnesol may cause oxidative stress in *Candida* species (120-123) and in *S. cerevisiae* (124, 125), the overlap in transcriptional changes in response to MG, BEN, and farnesol likely provides valuable insight into how *C. auris* and other *Candida* species sense and adapt to physiologically relevant stressors. In fact, MG itself may serve as a stress signal in various organisms. In plants, for example, intracellular MG increases in response to drought (126, 127), salinity (126, 128-131), cold stress (126), heavy metals (128), or phosphorous deficiency (131), and overexpression of certain genes involved in MG detoxification has been shown to enhance salt tolerance in tobacco (126) and in *Brassica juncea* (132). Investigating whether MG detoxification is linked to abiotic stressors such as salt, temperature, or desiccation in *Candida* species would be an interesting avenue of future research, particularly in *C. auris* due to its persistence on hospital surfaces and high salt tolerance.

Methods

Strains, media, and growth conditions

The sources of all strains used in this study are listed in **Table S7**. All strains were stored long term in a final concentration of 25% glycerol at -80°C and freshly streaked onto yeast extract peptone dextrose (YPD) agar (10 g/L yeast extract, 20 g/L peptone, 2% glucose, 1.5% agar)

once every seven days and maintained at room temperature. Unless otherwise noted, all overnight cultures were grown in 5 mL YPD liquid medium (10 g/L yeast extract, 20 g/L peptone, 2% glucose) on a rotary wheel at 30°C. Media was supplemented with 25 µg/mL BEN (stock 10 mg/ml in DMSO) or 5 mM, 10 mM, or 15 mM MG (Sigma-Aldrich, 5.55 M) as noted. *E. coli* strains were grown in LB with 15 µg/mL gentamycin (gent).

Plasmids for complementation of *C. auris* *MRR1a*

Plasmids for complementing *C. auris* *MRR1a* into *C. lusitaniae* were created as follows: the open reading frame of *MRR1a* was amplified from the genomic DNA of *C. auris* isolates B11221 (for *CauMRR1a*^{N647T}) and AR0390 (for *CauMRR1a*) using a forward primer with homology to the 5' flank of *C. lusitaniae* *MRR1* and a reverse primer with homology to the 3' flank of *C. lusitaniae* *MRR1* for recombination into the *C. lusitaniae* *MRR1* complementation plasmid pMQ30^{MRR1-L1191H+Q1197*} (58). Plasmid pMQ30^{MRR1-L1191H+Q1197*} was digested with *Ascl* (New England BioLabs) and *AgeI*-HF (New England BioLabs). The PCR products and digested plasmid were cleaned using the Zymo DNA Clean & Concentrator kit (Zymo Research) and assembled using the *S. cerevisiae* recombination technique described in (133). Recombined plasmids were isolated from *S. cerevisiae* using a yeast plasmid miniprep kit (Zymo Research) before transformation into NEB®5-alpha competent *E. coli* (New England BioLabs). *E. coli* containing pMQ30-derived plasmids were selected for on LB containing 15 µg/mL gentamycin. Plasmids from *E. coli* were isolated using a Zyppy Plasmid Miniprep kit (Zymo Research) and subsequently verified by Sanger sequencing with the Dartmouth College Genomics and Molecular Biology Shared Resources Core. *MRR1a* complementation plasmids containing the correct sequences were linearized with *NotI*-HF (New England BioLabs), cleaned up with the

Zymo DNA Clean & Concentrator kit (Zymo Research) and eluted in molecular biology grade water (Corning) before transformation of 1.5 µg into *C. lusitaniae* strain U04 *mrr1*Δ as described below. All plasmids used and created in this study are listed in **Table S7**. All primers used in this study are listed in **Table S8**.

Transformation of *C. lusitaniae* with *C. auris* *MRR1a* complementation constructs

Mutants in *C. lusitaniae* were generated using an expression-free CRISPR-Cas9 method as previously described (37, 58, 134). In brief, cells suspended in 1M sorbitol were electroporated immediately following the addition of 1.5 µg of *C. auris* *MRR1a* complementation plasmid that had been previously linearized with NotI-HF (New England BioLabs) and Cas9 ribonucleoprotein containing crRNA targeting the *NAT1* gene. Transformants were selected on YPD agar containing 600 µg/ml hygromycin B (HygB). Successful transformants were identified via PCR of the *C. lusitaniae* *MRR1* locus as previously described (37, 58). CRISPR RNAs (crRNAs; IDT) and primers used to validate transformants are listed in **Table S8**.

Minimum Inhibitory Concentration (MIC) Assay

MIC assays for FLZ were performed in RPMI-1640 medium (Sigma, containing L-glutamine, 165 mM MOPS, 2% glucose at pH 7) as described in (40) and (58) using the broth microdilution method. The final concentration of FLZ in each well ranged from 64 µg/mL to 0.125 µg/mL. Plates were incubated at 35°C and scored for growth at 24 and 48 hours; the results are reported in **Table S4**. The MIC was defined as the drug concentration that abolished visible growth compared to a drug-free control.

Growth Kinetics

Growth kinetic assays were performed as previously described in (58). In brief, exponential-phase cultures of *C. auris* or *C. lusitaniae* were washed and diluted in dH₂O to an OD₆₀₀ of 1; 60 µL of each diluted cell suspension was added to 5 mL fresh YPD. To each well of a clear 96-well flat-bottom plate (Falcon) was added 100 µL of YPD or YPD with MG at twice the desired final concentration and 100 µL of cell inoculum in YPD. Plates were arranged in technical triplicate for each strain and condition and incubated in a Synergy Neo2 Microplate Reader (BioTek, USA) according to the following protocol: heat to 37°C, start kinetic, read OD₆₀₀ every 60 minutes for 36 hours, end kinetic. Results were calculated in Microsoft Excel and plotted in GraphPad Prism 9.0.0 (GraphPad Software).

Quantitative Real-Time PCR

Overnight cultures of *C. auris* or *C. lusitaniae* were diluted 1:50 into 5 mL fresh YPD, and grown to for four hours at 37°C. To each culture was added MG to a final concentration of 5 mM (4.5 µL stock), BEN to a final concentration of 25 µg/mL (12.5 µL stock), or 4.5 µL molecular biology grade dH₂O. Cultures were returned to the roller drum at 37°C for 15 min (MG or dH₂O) or 30 min (BEN), then centrifuged at 5000 rpm for 5 min. The differences in time of exposure in the experimental scheme was used to maintain consistency with published experiments in other species, and not because of known differences in kinetics of activity for the two inducers. RNA isolation, gDNA removal, cDNA synthesis, and quantitative real-time PCR were performed as previously described (40). Transcripts were normalized to *C. auris* or *C. lusitaniae* *ACT1* expression as appropriate. Results were calculated in Microsoft Excel and plotted in GraphPad Prism 9.0.0 (GraphPad Software). Primers are listed in **Table S8**.

548

549 **RNA sequencing**

550 Overnight cultures of *C. auris* were diluted to an OD₆₀₀ of 0.1 in 5 mL fresh, pre-warmed
551 YPD, and incubated on a roller drum at 37°C for 5-6 doublings (approx. 6 hours). Cultures were
552 diluted once more to an OD₆₀₀ of 1 in 5 mL fresh, pre-warmed YPD and returned to the roller
553 drum at 37°C for another 5-6 doublings. To each culture was added MG to a final concentration
554 of 5 mM (4.5 µL), BEN to a final concentration of 25 µg/mL (12.5 µL), or 4.5 µL molecular biology
555 grade dH₂O. Cultures were returned to the roller drum at 37°C for 15 min (MG or dH₂O) or 30
556 min (BEN), then centrifuged at 5000 rpm for 5 min. Supernatants were discarded and RNA
557 isolation was performed on cell pellets as described above for qRT-PCR. gDNA was removed
558 from RNA samples as described above. DNA-free RNA samples were sent to the Microbial
559 Genome Sequencing Center (<https://www.migscenter.com/>) for RNA sequencing.

560

561 **Analysis of RNAseq**

562 RNAseq data were analyzed by the Microbial Genome Sequencing Center
563 (<https://www.migscenter.com/>) as follows: Quality control and adapter trimming was performed
564 with bcl2fastq ([https://support.illumina.com/sequencing/sequencing_software/bcl2fastq-](https://support.illumina.com/sequencing/sequencing_software/bcl2fastq-conversion-software.html)
565 [conversion-software.html](https://support.illumina.com/sequencing/sequencing_software/bcl2fastq-conversion-software.html)). Read mapping was performed with HISAT2 (135). Read
566 quantification was performed using Subread's featureCounts (136) functionality. Read counts
567 were loaded into R (<https://www.R-project.org/>) and normalized using edgeR's (137) Trimmed
568 Mean of M values (TMM) algorithm. Subsequent values were then converted to counts per
569 million (cpm). Differential expression analysis was performed using edgeR's Quasi Linear F-
570 Test. In the supplementary files, sheets named "All Quantified Genes" contain the results of the

exact test for all genes in addition to the normalized counts per million. Differentially expressed genes were determined using the cutoff of $|\log FC| > 1$ and $p < .05$.

Identification of orthologs

Orthologs of *C. auris* genes in *C. albicans*, *C. lusitaniae*, and *S. cerevisiae*, as well as orthologs between B11221 and the Clade I reference strain B8441, were identified using FungiDB (<https://fungidb.org>) (138, 139).

Generation of Venn diagrams

Venn diagrams of differentially expressed genes across different strains and conditions were computed using the Venn diagram tool from UGent Bioinformatics & Evolutionary Genomics, which is accessible at <https://bioinformatics.psb.ugent.be/webtools/Venn/>.

Statistical Analysis and Figure preparation

All graphs were prepared with GraphPad Prism 9.0.0 (GraphPad Software). Ratio paired t-tests and one-way ANOVA tests were performed in Prism; details on each test are described in the corresponding figure legends. All p-values were two-tailed and $p < 0.05$ were considered significant for all analyses performed and are indicated with asterisks in the text: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Data availability

The data supporting the findings in this study are available within the paper and its supplemental material and are also available from the corresponding author upon request. The

raw sequence reads from the RNA-Seq analysis have been deposited into NCBI sequence read archive under BioProject PRJNA801628.

Acknowledgements

We thank Joachim Morschhäuser and the FDA-CDC Antimicrobial Resistance Isolate Bank for providing strains. We thank Judith Berman for the pGEM-*URA3* plasmid used for yeast cloning. We thank Elora Demers for primers.

Author contributions. ARB and DAH conceived and designed the experiments and wrote the paper. ARB performed the experiments. ARB and DAH analyzed the data.

Funding. This study was supported by grants R01 5R01 AI127548 to DAH. Core services were provided by STANTO19R0 to CFF RDP, P30-DK117469 to DartCF, and P20-GM113132 to BioMT. Sequencing services and specialized equipment were provided by the Genomics and Molecular Biology Shared Resource Core at Dartmouth, NCI Cancer Center Support Grant 5P30 CA023108-41. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Competing interests. The authors have declared that no competing interests exist.

References

1. Pfaller MA, Andes DR, Diekema DJ, Horn DL, Reboli AC, Rotstein C, Franks B, Azie NE. 2014. Epidemiology and outcomes of invasive candidiasis due to non-albicans species of

Candida in 2,496 patients: data from the Prospective Antifungal Therapy (PATH) registry 2004-2008. PLoS One 9:e101510.

2. Pfaller MA, Diekema DJ. 2007. Epidemiology of invasive candidiasis: a persistent public health problem. Clin Microbiol Rev 20:133-63.

3. Quindos G, Marcos-Arias C, San-Millan R, Mateo E, Eraso E. 2018. The continuous changes in the aetiology and epidemiology of invasive candidiasis: from familiar *Candida albicans* to multiresistant *Candida auris*. Int Microbiol 21:107-119.

4. Abe M, Kinjo Y, Ueno K, Takatsuka S, Nakamura S, Ogura S, Kimura M, Araoka H, Sadamoto S, Shinozaki M, Shibuya K, Yoneyama A, Kaku M, Miyazaki Y. 2018. Differences in Ocular Complications Between *Candida albicans* and Non-albicans *Candida* Infection Analyzed by Epidemiology and a Mouse Ocular Candidiasis Model. Front Microbiol 9:2477.

5. Leeyaphan C, Bunyaratavej S, Foongladda S, Rujitharanawong C, Maneeprasopchoke P, Surawan T, Muanprasat C, Matthapan L. 2016. Epidemiology, Clinical Characteristics, Sites of Infection and Treatment Outcomes of Mucocutaneous Candidiasis Caused by Non-albicans Species of *Candida* at a Dermatologic Clinic. J Med Assoc Thai 99:406-11.

6. Quindos G. 2014. Epidemiology of candidaemia and invasive candidiasis. A changing face. Rev Iberoam Micol 31:42-8.

7. Patel PK, Erlandsen JE, Kirkpatrick WR, Berg DK, Westbrook SD, Loudon C, Cornell JE, Thompson GR, Vallor AC, Wickes BL, Wiederhold NP, Redding SW, Patterson TF. 2012. The Changing Epidemiology of Oropharyngeal Candidiasis in Patients with HIV/AIDS in the Era of Antiretroviral Therapy. AIDS Res Treat 2012:262471.

- 639 8. Hachem R, Hanna H, Kontoyiannis D, Jiang Y, Raad I. 2008. The changing epidemiology
640 of invasive candidiasis: *Candida glabrata* and *Candida krusei* as the leading causes of
641 candidemia in hematologic malignancy. *Cancer* 112:2493-9.
- 642 9. Redding SW, Kirkpatrick WR, Dib O, Fothergill AW, Rinaldi MG, Patterson TF. 2000. The
643 epidemiology of non-albicans *Candida* in oropharyngeal candidiasis in HIV patients. *Spec*
644 *Care Dentist* 20:178-81.
- 645 10. Abi-Said D, Anaissie E, Uzun O, Raad I, Pinzcowski H, Vartivarian S. 1997. The
646 epidemiology of hematogenous candidiasis caused by different *Candida* species. *Clin*
647 *Infect Dis* 24:1122-8.
- 648 11. Bravo Ruiz G, Lorenz A. 2021. What do we know about the biology of the emerging fungal
649 pathogen of humans *Candida auris*? *Microbiol Res* 242:126621.
- 650 12. Du H, Bing J, Hu T, Ennis CL, Nobile CJ, Huang G. 2020. *Candida auris*: Epidemiology,
651 biology, antifungal resistance, and virulence. *PLoS Pathog* 16:e1008921.
- 652 13. Satoh K, Makimura K, Hasumi Y, Nishiyama Y, Uchida K, Yamaguchi H. 2009. *Candida*
653 *auris* sp. nov., a novel ascomycetous yeast isolated from the external ear canal of an
654 inpatient in a Japanese hospital. *Microbiol Immunol* 53:41-4.
- 655 14. Lee WG, Shin JH, Uh Y, Kang MG, Kim SH, Park KH, Jang HC. 2011. First three reported
656 cases of nosocomial fungemia caused by *Candida auris*. *J Clin Microbiol* 49:3139-42.
- 657 15. Lockhart SR, Etienne KA, Vallabhaneni S, Farooqi J, Chowdhary A, Govender NP,
658 Colombo AL, Calvo B, Cuomo CA, Desjardins CA, Berkow EL, Castanheira M, Magobo
659 RE, Jabeen K, Asghar RJ, Meis JF, Jackson B, Chiller T, Litvintseva AP. 2017.
660 Simultaneous Emergence of Multidrug-Resistant *Candida auris* on 3 Continents

Confirmed by Whole-Genome Sequencing and Epidemiological Analyses. Clin Infect Dis 64:134-140.

16. Chow NA, de Groot T, Badali H, Abastabar M, Chiller TM, Meis JF. 2019. Potential Fifth Clade of *Candida auris*, Iran, 2018. Emerg Infect Dis 25:1780-1781.

17. Proctor DM, Dangana T, Sexton DJ, Fukuda C, Yelin RD, Stanley M, Bell PB, Baskaran S, Deming C, Chen Q, Conlan S, Park M, Program NCS, Welsh RM, Vallabhaneni S, Chiller T, Forsberg K, Black SR, Pacilli M, Kong HH, Lin MY, Schoeny ME, Litvintseva AP, Segre JA, Hayden MK. 2021. Integrated genomic, epidemiologic investigation of *Candida auris* skin colonization in a skilled nursing facility. Nat Med 27:1401-1409.

18. Horton MV, Johnson CJ, Kernien JF, Patel TD, Lam BC, Cheong JZA, Meudt JJ, Shanmuganayagam D, Kalan LR, Nett JE. 2020. *Candida auris* Forms High-Burden Biofilms in Skin Niche Conditions and on Porcine Skin. mSphere 5.

19. Uppuluri P. 2020. *Candida auris* Biofilm Colonization on Skin Niche Conditions. mSphere 5.

20. Osei Sekyere J. 2018. *Candida auris*: A systematic review and meta-analysis of current updates on an emerging multidrug-resistant pathogen. Microbiologyopen 7:e00578.

21. Taori SK, Khonyongwa K, Hayden I, Athukorala GDA, Letters A, Fife A, Desai N, Borman AM. 2019. *Candida auris* outbreak: Mortality, interventions and cost of sustaining control. J Infect 79:601-611.

22. Arensman K, Miller JL, Chiang A, Mai N, Levato J, LaChance E, Anderson M, Beganovic M, Dela Pena J. 2020. Clinical Outcomes of Patients Treated for *Candida auris* Infections in a Multisite Health System, Illinois, USA. Emerg Infect Dis 26:876-880.

- 683 23. AlJindan R, AlEraky DM, Mahmoud N, Abdalhamid B, Almustafa M, AbdulAzeez S,
684 Borgio JF. 2020. Drug Resistance-Associated Mutations in *ERG11* of Multidrug-Resistant
685 *Candida auris* in a Tertiary Care Hospital of Eastern Saudi Arabia. J Fungi (Basel) 7.
- 686 24. Carolus H, Pierson S, Munoz JF, Subotic A, Cruz RB, Cuomo CA, Van Dijck P. 2021.
687 Genome-Wide Analysis of Experimentally Evolved *Candida auris* Reveals Multiple Novel
688 Mechanisms of Multidrug Resistance. mBio 12.
- 689 25. Chow NA, Munoz JF, Gade L, Berkow EL, Li X, Welsh RM, Forsberg K, Lockhart SR,
690 Adam R, Alanio A, Alastruey-Izquierdo A, Althawadi S, Arauz AB, Ben-Ami R, Bharat A,
691 Calvo B, Desnos-Ollivier M, Escandon P, Gardam D, Gunturu R, Heath CH, Kurzai O,
692 Martin R, Litvintseva AP, Cuomo CA. 2020. Tracing the Evolutionary History and Global
693 Expansion of *Candida auris* Using Population Genomic Analyses. mBio 11.
- 694 26. Chowdhary A, Prakash A, Sharma C, Kordalewska M, Kumar A, Sarma S, Tarai B, Singh
695 A, Upadhyaya G, Upadhyay S, Yadav P, Singh PK, Khillan V, Sachdeva N, Perlin DS,
696 Meis JF. 2018. A multicentre study of antifungal susceptibility patterns among 350
697 *Candida auris* isolates (2009-17) in India: role of the *ERG11* and *FKS1* genes in azole
698 and echinocandin resistance. J Antimicrob Chemother 73:891-899.
- 699 27. Healey KR, Kordalewska M, Jimenez Ortigosa C, Singh A, Berrio I, Chowdhary A, Perlin
700 DS. 2018. Limited *ERG11* Mutations Identified in Isolates of *Candida auris* Directly
701 Contribute to Reduced Azole Susceptibility. Antimicrob Agents Chemother 62.
- 702 28. Li J, Coste AT, Liechti M, Bachmann D, Sanglard D, Lamothe F. 2021. Novel *ERG11* and
703 *TAC1b* mutations associated with azole resistance in *Candida auris*. Antimicrob Agents
704 Chemother doi:10.1128/AAC.02663-20.

- 705 29. Munoz JF, Gade L, Chow NA, Loparev VN, Juieng P, Berkow EL, Farrer RA, Litvintseva
706 AP, Cuomo CA. 2018. Genomic insights into multidrug-resistance, mating and virulence
707 in *Candida auris* and related emerging species. Nat Commun 9:5346.
- 708 30. Rybak JM, Sharma C, Doorley LA, Barker KS, Palmer GE, Rogers PD. 2021. Delineation
709 of the Direct Contribution of *Candida auris* *ERG11* Mutations to Clinical Triazole
710 Resistance. Microbiol Spectr 9:e0158521.
- 711 31. Williamson B, Wilk A, Guerrero KD, Mikulski TD, Elias TN, Sawh I, Cancino-Prado G,
712 Gardam D, Heath CH, Govender NP, Perlin DS, Kordalewska M, Healey KR. 2022.
713 Impact of Erg11 Amino Acid Substitutions Identified in *Candida auris* Clade III Isolates on
714 Triazole Drug Susceptibility. Antimicrob Agents Chemother 66:e0162421.
- 715 32. Rybak JM, Doorley LA, Nishimoto AT, Barker KS, Palmer GE, Rogers PD. 2019.
716 Abrogation of Triazole Resistance upon Deletion of *CDR1* in a Clinical Isolate of *Candida*
717 *auris*. Antimicrob Agents Chemother 63.
- 718 33. Kim SH, Iyer KR, Pardeshi L, Munoz JF, Robbins N, Cuomo CA, Wong KH, Cowen LE.
719 2019. Genetic Analysis of *Candida auris* Implicates Hsp90 in Morphogenesis and Azole
720 Tolerance and Cdr1 in Azole Resistance. mBio 10.
- 721 34. Kim SH, Iyer KR, Pardeshi L, Munoz JF, Robbins N, Cuomo CA, Wong KH, Cowen LE.
722 2019. Erratum for Kim et al., "Genetic Analysis of *Candida auris* Implicates Hsp90 in
723 Morphogenesis and Azole Tolerance and Cdr1 in Azole Resistance". mBio 10.
- 724 35. Rybak JM, Munoz JF, Barker KS, Parker JE, Esquivel BD, Berkow EL, Lockhart SR, Gade
725 L, Palmer GE, White TC, Kelly SL, Cuomo CA, Rogers PD. 2020. Mutations in *TAC1B*: a
726 Novel Genetic Determinant of Clinical Fluconazole Resistance in *Candida auris*. mBio 11.

- 727 36. Wasi M, Khandelwal NK, Moorhouse AJ, Nair R, Vishwakarma P, Bravo Ruiz G, Ross
728 ZK, Lorenz A, Rudramurthy SM, Chakrabarti A, Lynn AM, Mondal AK, Gow NAR, Prasad
729 R. 2019. ABC Transporter Genes Show Upregulated Expression in Drug-Resistant
730 Clinical Isolates of *Candida auris*: A Genome-Wide Characterization of ATP-Binding
731 Cassette (ABC) Transporter Genes. Front Microbiol 10:1445.
- 732 37. Demers EG, Stajich JE, Ashare A, Occhipinti P, Hogan DA. 2021. Balancing Positive and
733 Negative Selection: *In Vivo* Evolution of *Candida lusitanae* *MRR1*. mBio 12.
- 734 38. Liu Z, Myers LC. 2017. *Candida albicans* Swi/Snf and Mediator complexes differentially
735 regulate Mrr1-induced *MDR1* expression and fluconazole resistance. Antimicrob Agents
736 Chemother 61.
- 737 39. Kannan A, Asner SA, Trachsel E, Kelly S, Parker J, Sanglard D. 2019. Comparative
738 Genomics for the Elucidation of Multidrug Resistance in *Candida lusitanae*. mBio 10.
- 739 40. Demers EG, Biermann AR, Masonjones S, Crocker AW, Ashare A, Stajich JE, Hogan DA.
740 2018. Evolution of drug resistance in an antifungal-naïve chronic *Candida lusitanae*
741 infection. Proc Natl Acad Sci U S A 115:12040-12045.
- 742 41. Schubert S, Popp C, Rogers PD, Morschhauser J. 2011. Functional dissection of a
743 *Candida albicans* zinc cluster transcription factor, the multidrug resistance regulator Mrr1.
744 Eukaryot Cell 10:1110-21.
- 745 42. Schubert S, Rogers PD, Morschhauser J. 2008. Gain-of-function mutations in the
746 transcription factor *MRR1* are responsible for overexpression of the *MDR1* efflux pump in
747 fluconazole-resistant *Candida dubliniensis* strains. Antimicrob Agents Chemother
748 52:4274-80.

- 749 43. Schneider S, Morschhauser J. 2015. Induction of *Candida albicans* drug resistance genes
750 by hybrid zinc cluster transcription factors. Antimicrob Agents Chemother 59:558-69.
- 751 44. Schubert S, Barker KS, Znaldi S, Schneider S, Dierolf F, Dunkel N, Aid M, Boucher G,
752 Rogers PD, Raymond M, Morschhauser J. 2011. Regulation of efflux pump expression
753 and drug resistance by the transcription factors Mrr1, Upc2, and Cap1 in *Candida*
754 *albicans*. Antimicrob Agents Chemother 55:2212-23.
- 755 45. Morschhauser J, Barker KS, Liu TT, Bla BWJ, Homayouni R, Rogers PD. 2007. The
756 transcription factor Mrr1p controls expression of the *MDR1* efflux pump and mediates
757 multidrug resistance in *Candida albicans*. PLoS Pathog 3:e164.
- 758 46. Mayr EM, Ramirez-Zavala B, Kruger I, Morschhauser J. 2020. A Zinc Cluster
759 Transcription Factor Contributes to the Intrinsic Fluconazole Resistance of *Candida auris*.
760 mSphere 5.
- 761 47. Spivak ES, Hanson KE. 2018. *Candida auris*: an Emerging Fungal Pathogen. J Clin
762 Microbiol 56.
- 763 48. Jin L, Cao Z, Wang Q, Wang Y, Wang X, Chen H, Wang H. 2018. *MDR1* overexpression
764 combined with *ERG11* mutations induce high-level fluconazole resistance in *Candida*
765 *tropicalis* clinical isolates. BMC Infect Dis 18:162.
- 766 49. Liston SD, Whitesell L, Kapoor M, Shaw KJ, Cowen LE. 2020. Enhanced Efflux Pump
767 Expression in *Candida* Mutants Results in Decreased Manogepix Susceptibility.
768 Antimicrob Agents Chemother 64.
- 769 50. Wirsching S, Moran GP, Sullivan DJ, Coleman DC, Morschhauser J. 2001. *MDR1*-
770 mediated drug resistance in *Candida dubliniensis*. Antimicrob Agents Chemother
771 45:3416-21.

- 772 51. You L, Qian W, Yang Q, Mao L, Zhu L, Huang X, Jin J, Meng H. 2017. *ERG11* Gene
773 Mutations and *MDR1* Upregulation Confer Pan-Azole Resistance in *Candida tropicalis*
774 Causing Disseminated Candidiasis in an Acute Lymphoblastic Leukemia Patient on
775 Posaconazole Prophylaxis. Antimicrob Agents Chemother 61.
- 776 52. Wirsching S, Michel S, Morschhauser J. 2000. Targeted gene disruption in *Candida*
777 *albicans* wild-type strains: the role of the *MDR1* gene in fluconazole resistance of clinical
778 *Candida albicans* isolates. Mol Microbiol 36:856-65.
- 779 53. Berkow EL, Manigaba K, Parker JE, Barker KS, Kelly SL, Rogers PD. 2015. Multidrug
780 Transporters and Alterations in Sterol Biosynthesis Contribute to Azole Antifungal
781 Resistance in *Candida parapsilosis*. Antimicrob Agents Chemother 59:5942-50.
- 782 54. Souza AC, Fuchs BB, Pinhati HM, Siqueira RA, Hagen F, Meis JF, Mylonakis E, Colombo
783 AL. 2015. *Candida parapsilosis* Resistance to Fluconazole: Molecular Mechanisms and
784 *In Vivo* Impact in Infected *Galleria mellonella* Larvae. Antimicrob Agents Chemother
785 59:6581-7.
- 786 55. Grossman NT, Pham CD, Cleveland AA, Lockhart SR. 2015. Molecular mechanisms of
787 fluconazole resistance in *Candida parapsilosis* isolates from a U.S. surveillance system.
788 Antimicrob Agents Chemother 59:1030-7.
- 789 56. Hampe IAI, Friedman J, Edgerton M, Morschhauser J. 2017. An acquired mechanism of
790 antifungal drug resistance simultaneously enables *Candida albicans* to escape from
791 intrinsic host defenses. PLoS Pathog 13:e1006655.
- 792 57. Hiller D, Sanglard D, Morschhauser J. 2006. Overexpression of the *MDR1* gene is
793 sufficient to confer increased resistance to toxic compounds in *Candida albicans* (vol 50,
794 pg 1365, 2006). Antimicrobial Agents and Chemotherapy 50:2591-2591.

- 795 58. Biermann AR, Demers EG, Hogan DA. 2021. Mrr1 regulation of methylglyoxal catabolism
796 and methylglyoxal-induced fluconazole resistance in *Candida lusitanae*. *Mol Microbiol*
797 115:116-130.
- 798 59. Chakraborty S, Karmakar K, Chakravorty D. 2014. Cells Producing Their Own Nemesis:
799 Understanding Methylglyoxal Metabolism. *Iubmb Life* 66:667-678.
- 800 60. Tian S, Rong C, Nian H, Li F, Chu Y, Cheng S, Shang H. 2018. First cases and risk factors
801 of super yeast *Candida auris* infection or colonization from Shenyang, China. *Emerg*
802 *Microbes Infect* 7:128.
- 803 61. Sayeed MA, Farooqi J, Jabeen K, Mahmood SF. 2020. Comparison of risk factors and
804 outcomes of *Candida auris* candidemia with non-*Candida auris* candidemia: A
805 retrospective study from Pakistan. *Med Mycol* 58:721-729.
- 806 62. Shastri PS, Shankarnarayan SA, Oberoi J, Rudramurthy SM, Wattal C, Chakrabarti A.
807 2020. *Candida auris* candidaemia in an intensive care unit - Prospective observational
808 study to evaluate epidemiology, risk factors, and outcome. *J Crit Care* 57:42-48.
- 809 63. Rudramurthy SM, Chakrabarti A, Paul RA, Sood P, Kaur H, Capoor MR, Kindo AJ, Marak
810 RSK, Arora A, Sardana R, Das S, Chhina D, Patel A, Xess I, Tarai B, Singh P, Ghosh A.
811 2017. *Candida auris* candidaemia in Indian ICUs: analysis of risk factors. *J Antimicrob*
812 *Chemother* 72:1794-1801.
- 813 64. Ruiz-Gaitan A, Martinez H, Moret AM, Calabuig E, Tasias M, Alastruey-Izquierdo A,
814 Zaragoza O, Mollar J, Frasquet J, Salavert-Lleti M, Ramirez P, Lopez-Hontangas JL,
815 Peman J. 2019. Detection and treatment of *Candida auris* in an outbreak situation: risk
816 factors for developing colonization and candidemia by this new species in critically ill
817 patients. *Expert Rev Anti Infect Ther* 17:295-305.

- 818 65. Al-Rashdi A, Al-Maani A, Al-Wahaibi A, Alqayoudhi A, Al-Jardani A, Al-Abri S. 2021.
819 Characteristics, Risk Factors, and Survival Analysis of *Candida auris* Cases: Results of
820 One-Year National Surveillance Data from Oman. J Fungi (Basel) 7.
- 821 66. Caceres DH, Rivera SM, Armstrong PA, Escandon P, Chow NA, Ovalle MV, Diaz J,
822 Derado G, Salcedo S, Berrio I, Espinosa-Bode A, Varon C, Stuckey MJ, Marino A,
823 Villalobos N, Lockhart SR, Chiller TM, Prieto FE, Jackson BR. 2020. Case-Case
824 Comparison of *Candida auris* Versus Other *Candida* Species Bloodstream Infections:
825 Results of an Outbreak Investigation in Colombia. Mycopathologia 185:917-923.
- 826 67. Khan Z, Ahmad S, Benwan K, Purohit P, Al-Obaid I, Bafna R, Emara M, Mokaddas E,
827 Abdullah AA, Al-Obaid K, Joseph L. 2018. Invasive *Candida auris* infections in Kuwait
828 hospitals: epidemiology, antifungal treatment and outcome. Infection 46:641-650.
- 829 68. Pandya N, Cag Y, Pandak N, Pekok AU, Poojary A, Ayoade F, Fasciana T, Giammanco
830 A, Caskurlu H, Rajani DP, Gupta YK, Balkan, II, Khan EA, Erdem H. 2021. International
831 Multicentre Study of *Candida auris* Infections. J Fungi (Basel) 7.
- 832 69. Rossow J, Ostrowsky B, Adams E, Greenko J, McDonald R, Vallabhaneni S, Forsberg K,
833 Perez S, Lucas T, Alroy KA, Jacobs Slifka K, Walters M, Jackson BR, Quinn M,
834 Chaturvedi S, Blog D, New York Candida auris Investigation W. 2021. Factors Associated
835 With *Candida auris* Colonization and Transmission in Skilled Nursing Facilities With
836 Ventilator Units, New York, 2016-2018. Clin Infect Dis 72:e753-e760.
- 837 70. Wang XJ, Ma SB, Liu ZF, Li H, Gao WY. 2019. Elevated levels of alpha-dicarbonyl
838 compounds in the plasma of type II diabetics and their relevance with diabetic
839 nephropathy. Journal of Chromatography B-Analytical Technologies in the Biomedical
840 and Life Sciences 1106:19-25.

- 841 71. McLellan AC, Thornalley PJ, Benn J, Sonksen PH. 1994. Glyoxalase system in clinical
842 diabetes mellitus and correlation with diabetic complications. Clin Sci (Lond) 87:21-9.
- 843 72. Lu J, Randell E, Han Y, Adeli K, Krahn J, Meng QH. 2011. Increased plasma
844 methylglyoxal level, inflammation, and vascular endothelial dysfunction in diabetic
845 nephropathy. Clin Biochem 44:307-11.
- 846 73. Odani H, Shinzato T, Usami J, Matsumoto Y, Brinkmann Frye E, Baynes JW, Maeda K.
847 1998. Imidazolium crosslinks derived from reaction of lysine with glyoxal and
848 methylglyoxal are increased in serum proteins of uremic patients: evidence for increased
849 oxidative stress in uremia. FEBS Lett 427:381-5.
- 850 74. Karg E, Papp F, Tassi N, Janaky T, Wittmann G, Turi S. 2009. Enhanced methylglyoxal
851 formation in the erythrocytes of hemodialyzed patients. Metabolism 58:976-82.
- 852 75. Lapolla A, Flamini R, Lupo A, Arico NC, Rugiu C, Reitano R, Tubaro M, Ragazzi E,
853 Seraglia R, Traldi P. 2005. Evaluation of glyoxal and methylglyoxal levels in uremic
854 patients under peritoneal dialysis. Ann N Y Acad Sci 1043:217-24.
- 855 76. Mukhopadhyay S, Ghosh A, Kar M. 2008. Methylglyoxal increase in uremia with special
856 reference to snakebite-mediated acute renal failure. Clin Chim Acta 391:13-7.
- 857 77. Brenner T, Fleming T, Uhle F, Silaff S, Schmitt F, Salgado E, Ulrich A, Zimmermann S,
858 Bruckner T, Martin E, Bierhaus A, Nawroth PP, Weigand MA, Hofer S. 2014.
859 Methylglyoxal as a new biomarker in patients with septic shock: an observational clinical
860 study. Critical Care 18.
- 861 78. Juarez P, Jeannot K, Plesiat P, Llanes C. 2017. Toxic Electrophiles Induce Expression of
862 the Multidrug Efflux Pump MexEF-OprN in *Pseudomonas aeruginosa* through a Novel
863 Transcriptional Regulator, CmrA. Antimicrobial Agents and Chemotherapy 61.

- 864 79. Ozyamak E, de Almeida C, de Moura AP, Miller S, Booth IR. 2013. Integrated stress
865 response of *Escherichia coli* to methylglyoxal: transcriptional readthrough from the
866 nemRA operon enhances protection through increased expression of glyoxalase I. Mol
867 Microbiol 88:936-50.
- 868 80. Lee C, Shin J, Park C. 2013. Novel regulatory system nemRA-gloA for electrophile
869 reduction in *Escherichia coli* K-12. Mol Microbiol 88:395-412.
- 870 81. Mostofa MG, Ghosh A, Li ZG, Siddiqui MN, Fujita M, Tran LP. 2018. Methylglyoxal - a
871 signaling molecule in plant abiotic stress responses. Free Radic Biol Med 122:96-109.
- 872 82. Kosmachevskaya OV, Shumaev KB, Topunov AF. 2017. Signal and regulatory effects of
873 methylglyoxal in eukaryotic cells (review). Applied Biochemistry and Microbiology 53:273-
874 289.
- 875 83. Maeta K, Izawa S, Okazaki S, Kuge S, Inoue Y. 2004. Activity of the Yap1 transcription
876 factor in *Saccharomyces cerevisiae* is modulated by methylglyoxal, a metabolite derived
877 from glycolysis. Mol Cell Biol 24:8753-64.
- 878 84. Aguilera J, Prieto JA. 2004. Yeast cells display a regulatory mechanism in response to
879 methylglyoxal. FEMS Yeast Res 4:633-41.
- 880 85. Roy A, Hashmi S, Li Z, Dement AD, Cho KH, Kim JH. 2016. The glucose metabolite
881 methylglyoxal inhibits expression of the glucose transporter genes by inactivating the cell
882 surface glucose sensors Rgt2 and Snf3 in yeast. Mol Biol Cell 27:862-71.
- 883 86. Anonymous. 2016. Correction for The glucose metabolite methylglyoxal inhibits
884 expression of the glucose transporter genes by inactivating the cell surface glucose
885 sensors Rgt2 and Snf3 in yeast. Mol Biol Cell 27:3178-3179.

- 886 87. Maeta K, Izawa S, Inoue Y. 2005. Methylglyoxal, a metabolite derived from glycolysis,
887 functions as a signal initiator of the high osmolarity glycerol-mitogen-activated protein
888 kinase cascade and Calcineurin/Crz1-mediated pathway in *Saccharomyces cerevisiae*.
889 Journal of Biological Chemistry 280:253-260.
- 890 88. Takatsume Y, Izawa S, Inoue Y. 2007. Modulation of Spc1 stress-activated protein kinase
891 activity by methylglyoxal through inhibition of protein phosphatase in the fission yeast
892 *Schizosaccharomyces pombe*. Biochem Biophys Res Commun 363:942-7.
- 893 89. Takatsume Y, Izawa S, Inoue Y. 2006. Methylglyoxal as a signal initiator for activation of
894 the stress-activated protein kinase cascade in the fission yeast *Schizosaccharomyces*
895 *pombe*. J Biol Chem 281:9086-92.
- 896 90. Welsh RM, Bentz ML, Shams A, Houston H, Lyons A, Rose LJ, Litvintseva AP. 2017.
897 Survival, Persistence, and Isolation of the Emerging Multidrug-Resistant Pathogenic
898 Yeast *Candida auris* on a Plastic Health Care Surface. J Clin Microbiol 55:2996-3005.
- 899 91. Rutala WA, Kanamori H, Gergen MF, Sickbert-Bennett EE, Weber DJ. 2019.
900 Susceptibility of *Candida auris* and *Candida albicans* to 21 germicides used in healthcare
901 facilities. Infect Control Hosp Epidemiol 40:380-382.
- 902 92. Jabeen K, Mal PB, Tharwani A, Hashmi M, Farooqi J. 2020. Persistence of *Candida auris*
903 on latex and nitrile gloves with transmission to sterile urinary cathetersdouble dagger.
904 Med Mycol 58:128-132.
- 905 93. Eyre DW, Sheppard AE, Madder H, Moir I, Moroney R, Quan TP, Griffiths D, George S,
906 Butcher L, Morgan M, Newnham R, Sunderland M, Clarke T, Foster D, Hoffman P,
907 Borman AM, Johnson EM, Moore G, Brown CS, Walker AS, Peto TEA, Crook DW, Jeffery

908 KJM. 2018. A *Candida auris* Outbreak and Its Control in an Intensive Care Setting. N Engl
909 J Med 379:1322-1331.

910 94. Hoehamer CF, Cummings ED, Hilliard GM, Morschhauser J, Rogers PD. 2009. Proteomic
911 analysis of Mrr1p- and Tac1p-associated differential protein expression in azole-resistant
912 clinical isolates of *Candida albicans*. Proteomics Clin Appl 3:968-78.

913 95. Karababa M, Coste AT, Rognon B, Bille J, Sanglard D. 2004. Comparison of gene
914 expression profiles of *Candida albicans* azole-resistant clinical isolates and laboratory
915 strains exposed to drugs inducing multidrug transporters. Antimicrob Agents Chemother
916 48:3064-79.

917 96. Rogers PD, Barker KS. 2003. Genome-wide expression profile analysis reveals
918 coordinately regulated genes associated with stepwise acquisition of azole resistance in
919 *Candida albicans* clinical isolates. Antimicrob Agents Chemother 47:1220-7.

920 97. Silva AP, Miranda IM, Guida A, Synnott J, Rocha R, Silva R, Amorim A, Pina-Vaz C,
921 Butler G, Rodrigues AG. 2011. Transcriptional profiling of azole-resistant *Candida*
922 *parapsilosis* strains. Antimicrob Agents Chemother 55:3546-56.

923 98. Kannan A, Asner SA, Trachsel E, Kelly S, Parker J, Sanglard D. 2020. Erratum for
924 Kannan et al., "Comparative Genomics for the Elucidation of Multidrug Resistance in
925 *Candida lusitanae*". mBio 11.

926 99. Rognon B, Kozovska Z, Coste AT, Pardini G, Sanglard D. 2006. Identification of promoter
927 elements responsible for the regulation of *MDR1* from *Candida albicans*, a major
928 facilitator transporter involved in azole resistance. Microbiology (Reading) 152:3701-
929 3722.

- 930 100. Ramirez-Zavala B, Mogavero S, Scholler E, Sasse C, Rogers PD, Morschhauser J. 2014.
931 SAGA/ADA complex subunit Ada2 is required for Cap1- but not Mrr1-mediated
932 upregulation of the *Candida albicans* multidrug efflux pump *MDR1*. Antimicrob Agents
933 Chemother 58:5102-10.
- 934 101. Mogavero S, Tavanti A, Senesi S, Rogers PD, Morschhauser J. 2011. Differential
935 requirement of the transcription factor Mcm1 for activation of the *Candida albicans*
936 multidrug efflux pump *MDR1* by its regulators Mrr1 and Cap1. Antimicrob Agents
937 Chemother 55:2061-6.
- 938 102. Hiller D, Stahl S, Morschhauser J. 2006. Multiple cis-acting sequences mediate
939 upregulation of the *MDR1* efflux pump in a fluconazole-resistant clinical *Candida albicans*
940 isolate. Antimicrob Agents Chemother 50:2300-8.
- 941 103. Harry JB, Oliver BG, Song JL, Silver PM, Little JT, Choiniere J, White TC. 2005. Drug-
942 induced regulation of the *MDR1* promoter in *Candida albicans*. Antimicrob Agents
943 Chemother 49:2785-92.
- 944 104. Gupta V, Kohli A, Krishnamurthy S, Puri N, Aalamgeer SA, Panwar S, Prasad R. 1998.
945 Identification of polymorphic mutant alleles of *CaMDR1*, a major facilitator of *Candida*
946 *albicans* which confers multidrug resistance, and its *in vitro* transcriptional activation. Curr
947 Genet 34:192-9.
- 948 105. Iyer KR, Camara K, Daniel-Ivad M, Trilles R, Pimentel-Elardo SM, Fossen JL, Marchillo
949 K, Liu Z, Singh S, Munoz JF, Kim SH, Porco JA, Jr., Cuomo CA, Williams NS, Ibrahim
950 AS, Edwards JE, Jr., Andes DR, Nodwell JR, Brown LE, Whitesell L, Robbins N, Cowen
951 LE. 2020. An oxindole efflux inhibitor potentiates azoles and impairs virulence in the
952 fungal pathogen *Candida auris*. Nat Commun 11:6429.

106. Dunkel N, Blass J, Rogers PD, Morschhauser J. 2008. Mutations in the multi-drug resistance regulator *MRR1*, followed by loss of heterozygosity, are the main cause of *MDR1* overexpression in fluconazole-resistant *Candida albicans* strains. *Mol Microbiol* 69:827-40.
107. Rodrigues CF, Rodrigues ME, Henriques M. 2019. *Candida* sp. Infections in Patients with Diabetes Mellitus. *J Clin Med* 8.
108. Pyrgos V, Ratanavanich K, Donegan N, Veis J, Walsh TJ, Shoham S. 2009. *Candida* bloodstream infections in hemodialysis recipients. *Med Mycol* 47:463-7.
109. Zhang MM, Ong CL, Walker MJ, McEwan AG. 2016. Defence against methylglyoxal in Group A *Streptococcus*: a role for Glyoxylase I in bacterial virulence and survival in neutrophils? *Pathog Dis* 74.
110. Rachman H, Kim N, Ulrichs T, Baumann S, Pradl L, Nasser Eddine A, Bild M, Rother M, Kuban RJ, Lee JS, Hurwitz R, Brinkmann V, Kosmiadi GA, Kaufmann SH. 2006. Critical role of methylglyoxal and AGE in mycobacteria-induced macrophage apoptosis and activation. *PLoS One* 1:e29.
111. Prantner D, Nallar S, Richard K, Spiegel D, Collins KD, Vogel SN. 2021. Classically activated mouse macrophages produce methylglyoxal that induces a TLR4- and RAGE-independent proinflammatory response. *J Leukoc Biol* 109:605-619.
112. Jimenez-Lopez C, Collette JR, Brothers KM, Shepardson KM, Cramer RA, Wheeler RT, Lorenz MC. 2013. *Candida albicans* induces arginine biosynthetic genes in response to host-derived reactive oxygen species. *Eukaryot Cell* 12:91-100.
113. Aranda A, del Olmo ML. 2004. Exposure of *Saccharomyces cerevisiae* to acetaldehyde induces sulfur amino acid metabolism and polyamine transporter genes, which depend

- on Met4p and Haa1p transcription factors, respectively. Appl Environ Microbiol 70:1913-22.
114. Lucau-Danila A, Lelandais G, Kozovska Z, Tanty V, Delaveau T, Devaux F, Jacq C. 2005. Early expression of yeast genes affected by chemical stress. Mol Cell Biol 25:1860-8.
115. Lelandais G, Tanty V, Geneix C, Etchebest C, Jacq C, Devaux F. 2008. Genome adaptation to chemical stress: clues from comparative transcriptomics in *Saccharomyces cerevisiae* and *Candida glabrata*. Genome Biol 9:R164.
116. Staub RE, Quistad GB, Casida JE. 1998. Mechanism for benomyl action as a mitochondrial aldehyde dehydrogenase inhibitor in mice. Chemical Research in Toxicology 11:535-543.
117. Fitzmaurice AG, Rhodes SL, Lulla A, Murphy NP, Lam HA, O'Donnell KC, Barnhill L, Casida JE, Cockburn M, Sagasti A, Stahl MC, Maidment NT, Ritz B, Bronstein JM. 2013. Aldehyde dehydrogenase inhibition as a pathogenic mechanism in Parkinson disease. Proceedings of the National Academy of Sciences of the United States of America 110:636-641.
118. Casida JE, Ford B, Jinsmaa Y, Sullivan P, Cooney A, Goldstein DS. 2014. Benomyl, aldehyde dehydrogenase, DOPAL, and the catecholaldehyde hypothesis for the pathogenesis of Parkinson's disease. Chem Res Toxicol 27:1359-61.
119. Leiphon LJ, Picklo MJ, Sr. 2007. Inhibition of aldehyde detoxification in CNS mitochondria by fungicides. Neurotoxicology 28:143-9.
120. Jakab A, Balla N, Ragyak A, Nagy F, Kovacs F, Sajtos Z, Toth Z, Borman AM, Pocsi I, Baranyai E, Majoros L, Kovacs R. 2021. Transcriptional Profiling of the *Candida auris*

Response to Exogenous Farnesol Exposure. mSphere doi:10.1128/mSphere.00710-21:e0071021.

121. Nagy F, Vitalis E, Jakab A, Borman AM, Forgacs L, Toth Z, Majoros L, Kovacs R. 2020. *In vitro* and *in vivo* Effect of Exogenous Farnesol Exposure Against *Candida auris*. Front Microbiol 11:957.

122. Shirtliff ME, Krom BP, Meijering RA, Peters BM, Zhu J, Scheper MA, Harris ML, Jabra-Rizk MA. 2009. Farnesol-induced apoptosis in *Candida albicans*. Antimicrob Agents Chemother 53:2392-401.

123. Hasim S, Vaughn EN, Donohoe D, Gordon DM, Pfiffner S, Reynolds TB. 2018. Influence of phosphatidylserine and phosphatidylethanolamine on farnesol tolerance in *Candida albicans*. Yeast 35:343-351.

124. Machida K, Tanaka T, Fujita K, Taniguchi M. 1998. Farnesol-induced generation of reactive oxygen species via indirect inhibition of the mitochondrial electron transport chain in the yeast *Saccharomyces cerevisiae*. J Bacteriol 180:4460-5.

125. Fairn GD, MacDonald K, McMaster CR. 2007. A chemogenomic screen in *Saccharomyces cerevisiae* uncovers a primary role for the mitochondria in farnesol toxicity and its regulation by the Pkc1 pathway. J Biol Chem 282:4868-4874.

126. Yadav SK, Singla-Pareek SL, Ray M, Reddy MK, Sopory SK. 2005. Methylglyoxal levels in plants under salinity stress are dependent on glyoxalase I and glutathione. Biochem Biophys Res Commun 337:61-7.

127. Nahar K, Hasanuzzaman M, Alam MM, Fujita M. 2015. Glutathione-induced drought stress tolerance in mung bean: coordinated roles of the antioxidant defence and methylglyoxal detoxification systems. AoB Plants 7.

- .021 128. Melvin P, Bankapalli K, D'Silva P, Shivaprasad PV. 2017. Methylglyoxal detoxification by
.022 a DJ-1 family protein provides dual abiotic and biotic stress tolerance in transgenic plants.
.023 Plant Mol Biol 94:381-397.
- .024 129. Vemanna RS, Babitha KC, Solanki JK, Amarnatha Reddy V, Sarangi SK, Udayakumar
.025 M. 2017. Aldo-keto reductase-1 (*AKR1*) protect cellular enzymes from salt stress by
.026 detoxifying reactive cytotoxic compounds. Plant Physiol Biochem 113:177-186.
- .027 130. Mahmud JA, Hasanuzzaman M, Khan MIR, Nahar K, Fujita M. 2020. beta-Aminobutyric
.028 Acid Pretreatment Confers Salt Stress Tolerance in *Brassica napus* L. by Modulating
.029 Reactive Oxygen Species Metabolism and Methylglyoxal Detoxification. Plants (Basel) 9.
- .030 131. Rohman MM, Islam MR, Monsur MB, Amiruzzaman M, Fujita M, Hasanuzzaman M. 2019.
.031 Trehalose Protects Maize Plants from Salt Stress and Phosphorus Deficiency. Plants
.032 (Basel) 8.
- .033 132. Veena, Reddy VS, Sopory SK. 1999. Glyoxalase I from *Brassica juncea*: molecular
.034 cloning, regulation and its over-expression confer tolerance in transgenic tobacco under
.035 stress. Plant J 17:385-95.
- .036 133. Shanks RM, Caiazza NC, Hinsa SM, Toutain CM, O'Toole GA. 2006. *Saccharomyces*
.037 *cerevisiae*-based molecular tool kit for manipulation of genes from gram-negative
.038 bacteria. Appl Environ Microbiol 72:5027-36.
- .039 134. Grahl N, Demers EG, Crocker AW, Hogan DA. 2017. Use of RNA-Protein complexes for
.040 genome editing in non-albicans *Candida* Species. mSphere 2:00218-17.
- .041 135. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment
.042 and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol 37:907-915.

- .043 136. Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for
.044 assigning sequence reads to genomic features. *Bioinformatics* 30:923-30.
- .045 137. Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for
.046 differential expression analysis of digital gene expression data. *Bioinformatics* 26:139-40.
- .047 138. Basenko EY, Pulman JA, Shanmugasundram A, Harb OS, Crouch K, Starns D,
.048 Warrenfeltz S, Aurrecochea C, Stoeckert CJ, Jr., Kissinger JC, Roos DS, Hertz-Fowler
.049 C. 2018. FungiDB: An Integrated Bioinformatic Resource for Fungi and Oomycetes. *J*
.050 *Fungi (Basel)* 4.
- .051 139. Stajich JE, Harris T, Brunk BP, Brestelli J, Fischer S, Harb OS, Kissinger JC, Li W, Nayak
.052 V, Pinney DF, Stoeckert CJ, Jr., Roos DS. 2012. FungiDB: an integrated functional
.053 genomics database for fungi. *Nucleic Acids Res* 40:D675-81.

.054

.055

Figure Legends

Figure 1. Mrr1a regulates expression of *MGD1* and *MDR1* in *C. auris* isolate B11221. A) Growth curves of B11221 WT (blue) and its *mrr1a*Δ (red), *mrr1b*Δ (green), and *mrr1c*Δ (purple) derivatives in YPD + 10 mM MG. Data shown represent the mean ± SD for three independent experiments. **B-C)** qRT-PCR assessment of *MGD1* (**B**) and *MDR1* (**C**) expression in B11221 WT (blue) and *mrr1a*Δ (red) cultures grown to exponential phase in YPD at 37°C. Data shown represent the mean ± SD for three independent experiments. Ratio paired t-test was used for statistical evaluation; * p < 0.05. **D)** Volcano plot of all quantified genes in B11221 WT vs *mrr1a*Δ in the control condition. Each point represents a single gene; blue points indicate genes significantly more highly expressed in WT; red points indicate genes significantly more highly expressed in *mrr1a*Δ. Numbers adjacent to each colored point indicate the log₂FC in *mrr1a*Δ versus WT.

Figure 2. MG and BEN both lead to a vast transcriptional response in *C. auris* B11221, which includes upregulation of *MDR1* and *MGD1*. A-B) qRT-PCR analysis for expression of *MGD1* (**A**) and *MDR1* (**B**) in exponential-phase cultures of B11221 WT (blue) or *mrr1a*Δ (red) treated with MG or BEN as indicated. Data shown represent the mean ± SD for three independent experiments. Ratio paired t-test was used for statistical evaluation; ns p > 0.05, * p < 0.05, ** p < 0.01. **C-D)** Volcano plots of all quantified genes in B11221 WT treated with either MG (**C**) or BEN (**D**). Each point represents a single gene; magenta points indicate genes that were significantly upregulated compared to the control condition, teal points indicate genes that were significantly downregulated compared to the control condition. *MDR1* and *MGD1* are

shown along with the two most up- and down- regulated genes in each condition. **E-F)** Scatter plots of the average CPMs of all quantified genes in *mrr1a*Δ vs. B11221 WT treated with MG (**E**) or BEN (**F**). Each point represents a single gene. Points below the dotted line indicate genes that were more highly expressed in the WT, and points above the dotted line indicated genes that were more highly expressed in the *mrr1a*Δ mutant. *MDR1* and *MGD1* are shown with red dots for reference.

Figure 3. *MDR1* and *MGD1* are among the genes significantly more highly expressed in isolate B11221 compared to isolate AR0390. **A)** Growth curves of B11221 (blue) and AR#0390 (orange) in YPD + 10 mM MG. Data shown represent the mean ± SD for three independent experiments. **B-C)** qRT-PCR assessment of *MGD1* (**B**) and *MDR1* (**C**) expression in B11221 (blue) and AR0390 (orange) grown to exponential phase in YPD at 37°C. Data shown represent the mean ± SD for three independent experiments. Ratio paired t-test was used for statistical evaluation; * $p < 0.05$, **** $p < 0.0001$. **D)** Volcano plot of all quantified genes, matched by syntenic ortholog, in B11221 and AR0390 in the control condition (YPD). Each point represents a single gene; blue points indicate genes significantly more highly expressed in B11221; orange points indicate genes significantly more highly expressed in AR0390. **E-F)** qRT-PCR expression analysis for *MGD1* (**E**) and *MDR1* (**F**) in *C. lusitaniae* U04 *mrr1*Δ (grey) and its derivatives expressing *CauMRR1a*^{N647T} (dark blue) or *CauMRR1a* (brown). Data shown represent the mean ± SD for three independent experiments. One-way ANOVA was used for statistical evaluation; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. **G)** Growth curves of *C. lusitaniae* U04 *mrr1*Δ (grey) and its derivatives expressing *CauMRR1a*^{N647T} (dark blue) or *CauMRR1a* (brown) in YPD + 15 mM MG. One representative experiment of three independent

experiments is shown; error bars represent the standard deviation of technical replicates within the experiment.

Figure 4. MG induces expression of *MGD1* and *MDR1* in *C. auris* isolates B11221 and AR0390, but *C. auris MRR1a* is not inducible by MG when heterologously expressed in *C. lusitaniae*. A-B) qRT-PCR analysis for expression of *MGD1* (A) and *MDR1* (B) in exponential-phase cultures of B11221 (blue) or AR0390 (orange) treated with MG as indicated. Data shown represent the mean $\pm$ SD for three independent experiments. Ratio paired t-test was used for statistical evaluation; ns $p > 0.05$, * $p < 0.05$, ** $p < 0.01$. (C-F) qRT-PCR analysis for expression of *MGD1* (C,E) and *MDR1* (D,F) in exponential-phase cultures of *C. lusitaniae* U04 *mrr1* Δ (grey) and its derivatives expressing *CauMRR1a*^{N647T} (dark blue) or *CauMRR1a* (brown) treated with 5 mM MG for 15 min (C,D) or 25 μ g/mL BEN for 30 min (E,F). Data shown represent the mean $\pm$ SD for three independent experiments. Ratio paired t-test was used for statistical evaluation; ns $p > 0.05$, * $p < 0.05$, ** $p < 0.01$.

Figure 5. MG induces and represses common pathways across B11221 and AR0390. A) Venn diagram of genes with syntenic orthologs between B11221 and AR0390 that were significantly induced (indicated by "up" arrows) or repressed (indicated by "down" arrows) by MG in either or both strains. **B)** Scatter plot of the log₂FC of genes significantly induced by MG in AR0390 vs the log₂FC of genes induced by MG in B11221. Only genes with syntenic orthologs between the two strains are shown. Each point represents a single gene; points above the dotted line indicate genes which exhibited a greater Log₂FC in AR0390, and points below the dotted line indicate genes which exhibited a greater log₂FC in B11221. *MGD1* and *MDR1* are indicated

with red dots for reference. **C)** Graphic summary of major groups of genes that were significantly up- or down-regulated in response to MG in both B11221 and AR0390. Genes in bold text were also up- or down-regulated in response to BEN in B11221.

Figure S1. The *mrr1a*Δ mutant has a growth defect in high concentrations of MG, but not at 5 mM MG or in the YPD control. Growth curves of B11221 WT (blue) and its *mrr1a*Δ (red), *mrr1b*Δ (green), and *mrr1c*Δ (purple) derivatives in YPD (left), or YPD supplemented with 5 mM (middle), or 15 mM (right) MG. Data shown represent the mean ± SD for three independent experiments.

Figure S2. The transcriptional response of *mrr1a*Δ to either MG or BEN is overall similar to that of the B11221 WT parent strain. Volcano plots of all quantified genes in the *mrr1a*Δ mutant treated with either MG (**A**) or BEN (**B**). Each point represents a single gene; magenta points indicate genes that were significantly upregulated compared to the control condition, teal points indicate genes that were significantly downregulated compared to the control condition. *MDR1* is shown along with the two most up- and down- regulated genes in each condition.

Figure S3. *C. auris* strain AR0390 has a growth advantage over B11221 in YPD but loses that advantage in the presence of increasing concentrations of MG. Growth curves of B11221 (blue) and AR0390 (orange) in YPD (left), or YPD supplemented with 5 mM (middle), or 15 mM (right) MG. Data shown represent the mean ± SD for three independent experiments.

Figure S4. *C. lusitaniae* strains complemented with *CauMRR1a*^{N647T} or *CauMRR1a* do not differ in growth from the *mrr1*Δ parent at MG concentrations below 15 mM. Growth curves of *C. lusitaniae* U04 *mrr1*Δ (grey) and its derivatives expressing *CauMRR1a*^{N647T} (dark blue) or *CauMRR1a* (brown) in YPD (left) or YPD supplemented with 5 mM (middle), or 10 mM (right) MG. Data shown represent the mean ± SD for three independent experiments.

Figure S5. Treatment with 5 mM MG leads to the differential expression of more genes in AR0390 than in B11221. Volcano plot of all quantified genes in AR0390 treated with MG. Each point represents a single gene; magenta points indicate genes that were significantly upregulated compared to the control condition, teal points indicate genes that were significantly downregulated compared to the control condition. *MDR1* and *MGD1* are shown for reference.

Figure 1. Mrr1a regulates expression of *MGD1* and *MDR1* in *C. auris* isolate B11221. **A)** Growth curves of B11221 WT (blue) and its *mrr1aΔ* (red), *mrr1bΔ* (green), and *mrr1cΔ* (purple) derivatives in YPD + 10 mM MG. Data shown represent the mean $\pm$ SD for three independent experiments. **B-C)** qRT-PCR assessment of *MGD1* (**B**) and *MDR1* (**C**) expression in B11221 WT (blue) and *mrr1aΔ* (red) cultures grown to exponential phase in YPD at 37°C. Data shown represent the mean $\pm$ SD for three independent experiments. Ratio paired t-test was used for statistical evaluation; * $p < 0.05$. **D)** Volcano plot of all quantified genes in B11221 WT vs *mrr1aΔ* in the control condition. Each point represents a single gene; blue points indicate genes significantly more highly expressed in WT; red points indicate genes significantly more highly expressed in *mrr1aΔ*. Numbers adjacent to each colored point indicate the log₂FC in *mrr1aΔ* versus WT.

Figure 2. MG and BEN both lead to a vast transcriptional response in *C. auris* B11221, which includes upregulation of *MDR1* and *MGD1*. **A-B** qRT-PCR analysis for expression of *MGD1* (**A**) and *MDR1* (**B**) in exponential-phase cultures of B11221 WT (blue) or *mrr1Δ* (red) treated with MG or BEN as indicated. Data shown represent the mean $\pm$ SD for three independent experiments. Ratio paired t-test was used for statistical evaluation; ns $p > 0.05$, * $p < 0.05$, ** $p < 0.01$. **C-D** Volcano plots of all quantified genes in B11221 WT treated with either MG (**C**) or BEN (**D**). Each point represents a single gene; magenta points indicate genes that were significantly upregulated compared to the control condition, teal points indicate genes that were significantly downregulated compared to the control condition. *MDR1* and *MGD1* are shown along with the two most up- and down- regulated genes in each condition. **E-F** Scatter plots of the average CPMs of all quantified genes in *mrr1Δ* vs. B11221 WT treated with MG (**E**) or BEN (**F**). Each point represents a single gene. Points below the dotted line indicate genes that were more highly expressed in the WT, and points above the dotted line indicated genes that were more highly expressed in the *mrr1Δ* mutant. *MDR1* and *MGD1* are shown with red dots for reference.

Figure 3. *MDR1* and *MGD1* are among the genes significantly more highly expressed in isolate B11221 compared to isolate AR0390. **A)** Growth curves of B11221 (blue) and AR#0390 (orange) in YPD + 10 mM MG. Data shown represent the mean $\pm$ SD for three independent experiments. **B-C)** qRT-PCR assessment of *MGD1* (**B**) and *MDR1* (**C**) expression in B11221 (blue) and AR0390 (orange) grown to exponential phase in YPD at 37°C. Data shown represent the mean $\pm$ SD for three independent experiments. Ratio paired t-test was used for statistical evaluation; * $p < 0.05$, **** $p < 0.0001$. **D)** Volcano plot of all quantified genes, matched by syntenic ortholog, in B11221 and AR0390 in the control condition (YPD). Each point represents a single gene; blue points indicate genes significantly more highly expressed in B11221; orange points indicate genes significantly more highly expressed in AR0390. **E-F)** qRT-PCR expression analysis for *MGD1* (**E**) and *MDR1* (**F**) in *C. lusitaniae* U04 *mrr1*Δ (grey) and its derivatives expressing *CauMRR1a*^{N647T} (dark blue) or *CauMRR1a* (brown). Data shown represent the mean $\pm$ SD for three independent experiments. One-way ANOVA was used for statistical evaluation; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. **G)** Growth curves of *C. lusitaniae* U04 *mrr1*Δ (grey) and its derivatives expressing *CauMRR1a*^{N647T} (dark blue) or *CauMRR1a* (brown) in YPD + 15 mM MG. One representative experiment of three independent experiments is shown; error bars represent the standard deviation of technical replicates within the experiment.

Figure 4. MG induces expression of *MGD1* and *MDR1* in *C. auris* isolates B11221 and AR0390, but *C. auris* *MRR1a* is not inducible by MG when heterologously expressed in *C. lusitanae*. A-B) qRT-PCR analysis for expression of *MGD1* (A) and *MDR1* (B) in exponential-phase cultures of B11221 (blue) or AR0390 (orange) treated with MG as indicated. Data shown represent the mean $\pm$ SD for three independent experiments. Ratio paired t-test was used for statistical evaluation; ns $p > 0.05$, * $p < 0.05$, ** $p < 0.01$. (C-F) qRT-PCR analysis for expression of *MGD1* (C,E) and *MDR1* (D,F) in exponential-phase cultures of *C. lusitanae* U04 *mrr1Δ* (grey) and its derivatives expressing *CauMRR1a*^{N647T} (dark blue) or *CauMRR1a* (brown) treated with 5 mM MG for 15 min (C,D) or 25 μ g/mL BEN for 30 min (E,F). Data shown represent the mean $\pm$ SD for three independent experiments. Ratio paired t-test was used for statistical evaluation; ns $p > 0.05$, * $p < 0.05$, ** $p < 0.01$.

Figure 5. MG induces and represses common pathways across B11221 and AR0390. **A)** Venn diagram of genes with syntenic orthologs between B11221 and AR0390 that were significantly induced (indicated by "up" arrows) or repressed (indicated by "down" arrows) by MG in either or both strains. **B)** Scatter plot of the log₂FC of genes significantly induced by MG in AR0390 vs the log₂FC of genes induced by MG in B11221. Only genes with syntenic orthologs between the two strains are shown. Each point represents a single gene; points above the dotted line indicate genes which exhibited a greater Log₂FC in AR0390, and points below the dotted line indicate genes which exhibited a greater log₂FC in B11221. *MGD1* and *MDR1* are indicated with red dots for reference. **C)** Graphic summary of major groups of genes that were significantly up- or down-regulated in response to MG in both B11221 and AR0390. Genes in bold text were also up- or down-regulated in response to BEN in B11221.

Figure S1. The *mrr1aΔ* mutant has a growth defect in high concentrations of MG, but not at 5 mM MG or in the YPD control. Growth curves of B11221 WT (blue) and its *mrr1aΔ* (red), *mrr1bΔ* (green), and *mrr1cΔ* (purple) derivatives in YPD (left), or YPD supplemented with 5 mM (middle), or 15 mM (right) MG. Data shown represent the mean $\pm$ SD for three independent experiments.

Figure S2. The transcriptional response of *mrr1a*Δ to either MG or BEN is overall similar to that of the B11221 WT parent strain. Volcano plots of all quantified genes in the *mrr1a*Δ mutant treated with either MG (**A**) or BEN (**B**). Each point represents a single gene; magenta points indicate genes that were significantly upregulated compared to the control condition, teal points indicate genes that were significantly downregulated compared to the control condition. *MDR1* is shown along with the two most up- and down- regulated genes in each condition.

Figure S3. *C. auris* strain AR0390 has a growth advantage over B11221 in YPD but loses that advantage in the presence of increasing concentrations of MG. Growth curves of B11221 (blue) and AR0390 (orange) in YPD (left), or YPD supplemented with 5 mM (middle), or 15 mM (right) MG. Data shown represent the mean $\pm$ SD for three independent experiments.

Figure S4. *C. lusitaniae* strains complemented with *CauMRR1a*^{N647T} or *CauMRR1a* do not differ in growth from the *mrr1Δ* parent at MG concentrations below 15 mM. Growth curves of *C. lusitaniae* U04 *mrr1Δ* (grey) and its derivatives expressing *CauMRR1a*^{N647T} (dark blue) or *CauMRR1a* (brown) in YPD (left) or YPD supplemented with 5 mM (middle), or 10 mM (right) MG. Data shown represent the mean $\pm$ SD for three independent experiments.

Figure S5. Treatment with 5 mM MG leads to the differential expression of more genes in AR0390 than in B11221. Volcano plot of all quantified genes in AR0390 treated with MG. Each point represents a single gene; magenta points indicate genes that were significantly upregulated compared to the control condition, teal points indicate genes that were significantly downregulated compared to the control condition. *MDR1* and *MGD1* are shown for reference.

Table S1. Select genes significantly upregulated by MG and/or BEN in the B11221 background.

Sulfur compound assimilation and biosynthesis						
Locus Tag	Gene Name	Predicted function	WT MG Log ₂ FC	WT BEN Log ₂ FC	<i>mrr1a</i> Δ MG Log ₂ FC	<i>mrr1a</i> Δ BEN Log ₂ FC
CJI97_001242	<i>AGP3</i>	Serine transporter	1.06	0.80	1.36	0.42
CJI97_002494	<i>DUG1</i>	Glutathione hydrolase	1.42	-0.02	1.38	-0.46
CJI97_001665	<i>CYS3</i>	Peroxisomal cystathionine beta-lyase	1.37	0.42	1.39	0.30
CJI97_001939	<i>CFD1</i>	Role in Fe-S cluster assembly	-0.02	1.17	0.26	1.31
CJI97_001514	<i>CIA1</i>	Role in protein maturation by Fe-S cluster transfer	0.40	1.12	0.34	1.20
CJI97_004156	<i>DRE2</i>	Cytosolic Fe-S protein assembly protein	1.09	1.08	1.15	0.98
CJI97_001503	<i>ECM4</i>	Cytoplasmic glutathione S-transferase	0.48	1.55	0.33	1.73
CJI97_001382	<i>ECM17</i>	Sulfite reductase beta subunit	1.14	0.76	1.45	0.33
CJI97_001705	<i>GCS1</i>	Gamma-glutamylcysteine synthetase	0.78	2.00	1.06	1.75
CJI97_003892	<i>GLR1</i>	Glutathione reductase	-0.04	1.17	0.06	1.11
CJI97_005081	<i>GSH2</i>	Glutathione synthase	1.07	1.41	0.98	1.13
CJI97_003274	<i>GTO1</i>	Cytoplasmic glutathione S-transferase	1.39	3.93	1.93	4.27
CJI97_001739	<i>JLP1</i>	Sulfonate dioxygenase	0.38	1.44	0.42	1.32
CJI97_002076	<i>MET1</i>	Uroporphyrin-3 C-methyltransferase	1.18	1.55	1.99	1.33
CJI97_002761	<i>MET2</i>	Homoserine acetyltransferase	2.20	0.49	2.31	0.30
CJI97_004689	<i>MET8</i>	Dehydrogenase, ferrochelatase	1.83	0.30	1.65	0.65
CJI97_003625	<i>MET10</i>	Sulfite reductase	1.01	0.94	1.31	0.86
CJI97_003066	<i>MET14</i>	Adenylylsulfate kinase	1.08	0.27	1.50	0.61
CJI97_005391	<i>MET16</i>	3'-phosphoadenylylsulfate reductase	1.63	1.44	1.95	1.29
CJI97_003613	<i>MUP1</i>	High affinity methionine permease	1.05	0.36	0.94	-0.06
CJI97_004493	<i>MUP3</i>	L-methionine transmembrane transporter	0.31	1.65	0.20	1.59
CJI97_004842	N/A	Role in Fe-S cluster assembly	-0.08	1.00	0.08	0.60
CJI97_005600	N/A	Cystathionine gamma-synthase	1.56	0.34	1.45	0.52
CJI97_001635	N/A	Glutathione S-conjugate transporter	0.05	1.10	0.12	0.90
CJI97_000433	<i>SPE2</i>	S-adenosylmethionine decarboxylase	1.11	0.19	1.14	0.05
CJI97_000171	<i>SRX1</i>	Sulfiredoxin	2.21	3.07	2.03	3.47
CJI97_003300	<i>STR2</i>	Cystathionine gamma-synthase	1.15	0.21	1.11	0.15
CJI97_001014	<i>SUL2</i>	Sulfate transporter	1.98	1.42	3.01	1.17
CJI97_003257	<i>TES1</i>	Acyl-CoA thioesterase	1.27	0.59	1.06	0.38
CJI97_001516	<i>TRR1</i>	Thioredoxin reductase	0.40	1.60	0.48	1.82
CJI97_000545	<i>TRX1</i>	Thioredoxin	-0.33	1.20	-0.82	2.05
Xenobiotic/Drug Transport						
Locus Tag	Gene Name	Predicted function	WT MG Log ₂ FC	WT BEN Log ₂ FC	<i>mrr1a</i> Δ MG Log ₂ FC	<i>mrr1a</i> Δ BEN Log ₂ FC
CJI97_002597	<i>AMF1</i>	MFS family transporter	0.35	1.28	0.71	1.29

CJI97_000167	<i>CDR1</i>	ABC family multidrug transporter	0.30	1.88	0.40	1.76
CJI97_000479	<i>CDR4</i>	ABC family multidrug transporter	1.34	1.13	1.27	1.14
CJI97_004181	<i>ERC1</i>	Xenobiotic transmembrane transporter	1.47	1.46	1.96	1.44
CJI97_004982	<i>ESBP6</i>	MFS membrane transporter	3.97	0.66	3.90	0.20
CJI97_002850	<i>FLU1</i>	Multidrug efflux pump of the plasma membrane	0.18	1.25	0.35	1.17
CJI97_002639	<i>MCH2</i>	MFS membrane transporter	1.36	0.04	1.24	-0.01
CJI97_000609	<i>MCH4</i>	MFS membrane transporter	2.13	0.47	2.14	0.33
CJI97_004042	<i>MDR1</i>	Plasma membrane MDR/MFS multidrug efflux pump	3.83	5.60	3.65	6.65
CJI97_000797	N/A	MFS membrane transporter	3.29	1.68	3.17	0.98
CJI97_005702	N/A	ABC family multidrug transporter	1.22	1.94	1.10	2.42
CJI97_005706	N/A	ABC family multidrug transporter	0.86	1.93	0.78	1.99
CJI97_005256	<i>QDR3</i>	MFS membrane transporter	2.83	-1.94	2.63	-2.88
CJI97_005513	<i>ROA1</i>	PDR-subfamily ABC transporter	2.86	0.05	3.58	0.01
CJI97_001481	<i>SNQ2</i>	ABC family multidrug transporter	1.50	2.56	1.34	2.19
CJI97_001817	<i>VBA1</i>	MFS transporter	0.68	1.18	0.64	0.95

Amino acid biosynthesis, excluding sulfur-containing amino acids

Locus Tag	Gene Name	Predicted function	WT MG Log ₂ FC	WT BEN Log ₂ FC	<i>mrr1Δ</i> MG Log ₂ FC	<i>mrr1Δ</i> BEN Log ₂ FC
CJI97_000687	<i>ARG1</i>	Argininosuccinate synthase	4.72	0.81	4.63	0.55
CJI97_004654	<i>ARG3</i>	Ornithine carbamoyltransferase	4.77	1.52	4.74	0.91
CJI97_002308	<i>ARG4</i>	Argininosuccinate lyase	1.75	0.39	1.67	0.21
CJI97_001846	<i>ARG5,6</i>	Arginine biosynthetic enzyme	1.50	0.60	1.46	0.19
CJI97_005293	<i>ARG8</i>	Acetylornithine aminotransferase	1.58	0.23	1.67	0.17
CJI97_002234	<i>ARO1</i>	Pentafunctional arom enzyme	1.49	-0.06	1.32	-0.20
CJI97_000465	<i>ARO2</i>	Chorismate synthase	2.40	0.53	2.37	0.12
CJI97_001597	<i>ARO3</i>	3-deoxy-D-arabinoheptulosonate-7-phosphate synthase	1.74	-0.12	1.64	-0.02
CJI97_003954	<i>ARO4</i>	3-deoxy-D-arabinoheptulosonate-7-phosphate synthase	2.50	0.36	2.38	0.18
CJI97_003913	<i>ARO7</i>	Chorismate mutase	1.51	0.08	1.30	0.37
CJI97_001973	<i>ASN1</i>	Asparagine synthetase	2.32	-0.20	2.06	-0.39
CJI97_001997	<i>BAT21</i>	Branched chain amino acid aminotransferase	2.51	0.24	2.61	0.10
CJI97_002013	<i>CPA2</i>	Carbamoyl-phosphate synthase subunit	1.78	0.15	1.83	-0.37
CJI97_005329	<i>HIS1</i>	ATP phosphoribosyl transferase	3.40	0.97	3.27	0.57
CJI97_003017	<i>HIS3</i>	Imidazoleglycerol-phosphate dehydratase	3.20	1.31	2.97	1.46
CJI97_003537	<i>HIS4</i>	Phosphoribosyl-AMP cyclohydrolase, phosphoribosyl-ATP diphosphatase, and histidinol dehydrogenase	2.86	0.94	2.73	0.33
CJI97_003604	<i>HIS5</i>	Histidinol-phosphate aminotransferase	2.72	0.73	2.60	0.49
CJI97_003946	<i>HIS7</i>	Imidazole glycerol phosphate synthase	2.39	0.63	2.34	0.55
CJI97_003721	<i>HOM2</i>	Aspartate-semialdehyde dehydrogenase	1.52	0.23	1.40	0.20
CJI97_003292	<i>HOM3</i>	L-aspartate 4-P-transferase	3.04	0.61	3.08	0.35

CJI97_003178	<i>ILV1</i>	Threonine dehydratase	2.06	0.53	1.93	0.28
CJI97_002682	<i>ILV2</i>	Acetolactate synthase	2.59	0.23	2.40	-0.17
CJI97_000020	<i>ILV3</i>	Dihydroxyacid dehydratase	2.77	0.75	2.60	0.37
CJI97_003514	<i>ILV5</i>	Ketol-acid reductoisomerase	1.99	-0.29	2.04	-0.50
CJI97_004523	<i>ILV6</i>	Regulatory subunit of acetolactate synthase	1.50	0.30	1.25	0.31
CJI97_004671	<i>LEU1</i>	3-isopropylmalate dehydratase	1.45	-0.06	1.30	-0.31
CJI97_001329	<i>LEU4</i>	2-isopropylmalate synthase	4.41	0.44	4.13	-0.12
CJI97_003280	<i>LYS1</i>	Saccharopine dehydrogenase	2.89	0.33	2.81	0.11
CJI97_003346	<i>LYS2</i>	Alpha-aminoacidopate reductase, large subunit	2.47	0.38	2.24	-0.31
CJI97_002417	<i>LYS4</i>	Homoaconitase	3.08	1.06	2.92	0.50
CJI97_002151	<i>LYS5</i>	Phosphopantetheinyl transferase	2.70	0.24	2.55	0.29
CJI97_001920	<i>LYS9</i>	Saccharopine dehydrogenase	1.13	0.07	0.96	-0.06
CJI97_003796	<i>LYS22</i>	Homocitrate synthase	2.91	0.07	2.82	-0.33
CJI97_003176	<i>SER1</i>	3-phosphoserine aminotransferase	2.11	0.16	2.05	0.15
CJI97_003156	<i>SER2</i>	Phosphoserine phosphatase	1.29	0.14	1.34	0.34
CJI97_000320	<i>SER33</i>	Enzyme of amino acid biosynthesis	1.05	-0.29	1.03	-0.51
CJI97_003863	<i>SHM1</i>	Mitochondrial serine hydroxymethyltransferase	1.30	-0.11	1.22	-0.01
CJI97_001157	<i>THR1</i>	Homoserine kinase	1.82	0.41	1.75	0.25
CJI97_000379	<i>TRP2</i>	Anthranilate synthase	1.05	0.42	0.91	0.08
CJI97_003979	<i>TRP3</i>	Indole-3-glycerol-phosphate synthase, anthranilate synthase	1.33	-0.02	1.27	0.12
CJI97_003855	<i>TRP4</i>	Enzyme of amino acid biosynthesis	1.25	0.06	1.17	-0.05
CJI97_003424	<i>TRP5</i>	Tryptophan synthase	2.45	0.28	2.25	0.17

Redox homeostasis and stress response

Locus Tag	Gene Name	Predicted function	WT MG Log ₂ FC	WT BEN Log ₂ FC	<i>mrr1Δ</i> MG Log ₂ FC	<i>mrr1Δ</i> BEN Log ₂ FC
CJI97_001841	<i>CAT1</i>	Catalase	0.54	2.49	0.67	2.38
CJI97_003095	<i>CIP1</i>	Oxidoreductase	0.51	8.04	1.33	8.70
CJI97_002824	<i>FDH3</i>	Oxidoreductase and zinc ion binding activity	-0.61	1.14	1.22	1.22
CJI97_000658	<i>MGD1</i>	NAD(H)-linked methylglyoxal oxidoreductase	1.35	2.46	-0.46	0.66
CJI97_002022	N/A	Quinone oxidoreductase	0.36	1.98	0.53	2.18
CJI97_002187	N/A	Oxidoreductase	-0.04	1.07	-0.07	0.97
CJI97_004869	N/A	Oxidoreductase	0.33	1.58	0.16	1.57
CJI97_004704	<i>POS5</i>	Mitochondrial NADH kinase	1.41	-0.46	1.54	-0.59
CJI97_004613	<i>PST3</i>	Flavodoxin-like protein	0.41	6.37	0.38	6.71
CJI97_000530	<i>SOD2</i>	Mitochondrial superoxide dismutase	-0.26	1.02	-0.39	1.41
CJI97_001282	<i>YAH1</i>	Oxidoreductase	1.01	0.99	0.87	1.07
CJI97_002560	<i>YCF1</i>	Glutathione S-conjugate transporter	4.58	0.85	0.12	0.90
CJI97_004612	<i>YCP4</i>	Flavodoxin-like protein	0.63	5.35	0.60	5.61

Significantly induced genes were determined using a cutoff of Log₂FC ≥ 1.00 and p-value < 0.05.

Table S2. Select genes significantly downregulated by MG and/or BEN in the B11221 background
Ergosterol biosynthesis

Locus Tag	Gene Name	Predicted function	WT MG Log ₂ FC	WT BEN Log ₂ FC	<i>mrr1a</i> Δ MG Log ₂ FC	<i>mrr1a</i> Δ BEN Log ₂ FC
CJI97_000262	<i>ERG1</i>	Squalene epoxidase	-0.85	-1.43	-0.68	-1.38
CJI97_003811	<i>ERG3</i>	C-5 sterol desaturase	-0.96	-1.11	-0.70	-0.73
CJI97_005423	<i>ERG6</i>	Delta(24)-sterol C-methyltransferase	-1.01	-1.53	-0.78	-1.09
CJI97_005634	<i>ERG10</i>	Acetyl-CoA acetyltransferase	-0.93	-1.04	-1.07	-0.77
CJI97_005638	<i>ERG10</i>	Acetyl-CoA acetyltransferase	-1.48	-0.63	-0.71	-1.63
CJI97_001156	<i>ERG11</i>	Lanosterol 14- α -demethylase	-0.24	-1.40	-0.13	-1.09

Metal acquisition, including regulation

Locus Tag	Gene Name	Predicted function	WT MG Log ₂ FC	WT BEN Log ₂ FC	<i>mrr1a</i> Δ MG Log ₂ FC	<i>mrr1a</i> Δ BEN Log ₂ FC
CJI97_002517	<i>CTR1</i>	Copper transporter	-1.13	-0.26	-1.24	0.01
CJI97_000015	<i>FRE7</i>	Ferric reductase	-1.23	-0.11	-1.43	-0.40
CJI97_003972	<i>FRE8</i>	Iron/copper reductase	-1.03	-1.18	-0.90	-1.30
CJI97_004532	<i>FRP1</i>	Ferric reductase	-1.41	-1.39	-0.85	-1.66
CJI97_001154	N/A	Ferric or cupric reductase	-0.57	-1.79	-0.97	-2.16
CJI97_004566	N/A	Ferric or cupric reductase	-1.43	-1.46	-1.51	-1.81
CJI97_005148	N/A	Ferric or cupric reductase	-1.56	-0.18	-1.81	-0.50
CJI97_002299	N/A	High affinity iron transporter for intravacuolar stores of iron	-0.19	-1.79	-0.16	-1.54
CJI97_001085	N/A	Transporter of ferrochrome siderophores	-0.06	-1.10	-0.57	-1.83
CJI97_001117	N/A	Transporter of ferrochrome siderophores	-1.90	-2.52	-1.67	-2.30
CJI97_001762	N/A	Transporter of ferrochrome siderophores	-1.94	-2.41	-1.87	-2.31
CJI97_004100	N/A	Transporter of ferrochrome siderophores	-0.26	-1.47	-0.21	-1.70
CJI97_004165	N/A	Transporter of ferrochrome siderophores	0.58	-1.14	0.83	-0.62
CJI97_001499	<i>SEF1</i>	Zn2-Cys6 transcription factor, regulates iron uptake	-1.13	0.31	-1.10	0.02
CJI97_000010	<i>ZRT2</i>	Zinc transporter	-0.38	-2.50	-0.80	-2.32

Carbohydrate metabolism and biosynthesis

Locus Tag	Gene Name	Predicted function	WT MG Log ₂ FC	WT BEN Log ₂ FC	<i>mrr1a</i> Δ MG Log ₂ FC	<i>mrr1a</i> Δ BEN Log ₂ FC
CJI97_003911	<i>DAC1</i>	N-acetylglucosamine-6-phosphate deacetylase	-0.35	-2.02	0.20	-2.05
CJI97_005247	<i>FBP1</i>	Fructose-1,6-bisphosphatase	-1.33	-1.11	-1.06	-0.96
CJI97_003057	<i>GLC3</i>	1,4-glucan branching enzyme	-1.05	-1.86	-0.95	-1.62
CJI97_001045	<i>GSY1</i>	Glycogen synthase	-1.17	-1.40	-1.10	-1.25
CJI97_003909	<i>HXK1</i>	N-acetylglucosamine kinase	-0.53	-1.15	-0.36	-1.37
CJI97_005579	<i>MAE1</i>	Mitochondrial malic enzyme	-1.44	-1.41	-1.36	-1.49
CJI97_000695	<i>MDH1</i>	Mitochondrial malate dehydrogenase	-1.23	-0.73	-1.09	-0.51
CJI97_003910	<i>NAG1</i>	Glucosamine-6-phosphate deaminase	-0.28	-1.76	0.03	-1.42

CJI97_001805	N/A	Role in beta-1,6 glucan biosynthesis	-1.16	-0.30	-0.95	-0.84
CJI97_000696	<i>NTH1</i>	Neutral trehalase	-1.07	-1.46	-1.20	-1.43
CJI97_002722	<i>PCK1</i>	Phosphoenolpyruvate carboxykinase	-2.03	-1.92	-1.68	-1.77
CJI97_002654	<i>PMM1</i>	Phosphomannomutase	-1.05	-1.33	-1.12	-0.73

Glucose transport, including regulation

Locus Tag	Gene Name	Predicted function	WT MG Log ₂ FC	WT BEN Log ₂ FC	<i>mrr1a</i> Δ MG Log ₂ FC	<i>mrr1a</i> Δ BEN Log ₂ FC
CJI97_000584	<i>HGT16</i>	MFS glucose transporter	1.38	-1.96	1.28	-2.14
CJI97_002713	<i>HGT17</i>	MFS glucose transporter	0.76	-3.05	-0.60	-2.46
CJI97_005108/ CJI97_005109	<i>HGT19</i>	MFS glucose/myo-inositol transporter	-1.71	-2.78	-1.95	-2.14
CJI97_001793	N/A	MFS glucose transporter	-2.00	-6.23	-2.27	-4.76
CJI97_001794	N/A	MFS glucose transporter	0.85	-2.17	-0.21	-1.41
CJI97_002023	N/A	MFS glucose transporter	-1.57	-5.00	-1.58	-5.13
CJI97_002024	N/A	MFS glucose transporter	-2.02	-2.80	-1.69	-2.70
CJI97_002448	<i>RGT1</i>	Transcriptional repressor of glucose transport	-1.26	-1.50	-1.29	-1.56
CJI97_005617	<i>SHA3</i>	Ser/thr kinase involved in glucose transport	-1.14	-0.65	-0.87	-1.09

Significantly repressed genes were determined using a cutoff of Log₂FC ≤ -1.00 and p-value < 0.05.

Table S3. Top 20 genes with predicted functions differentially expressed between B11221 and AR0390 in the control condition

Genes more highly expressed in B11221						
B11221 Locus Tag	AR0390 Locus Tag	Gene Name	Predicted Function	Log ₂ FC (B11221 vs AR0390)	B11221 average (CPM)	AR0390 average (CPM)
CJI97_004624	B9J08_004828	<i>MGD2</i>	NAD(H)-linked methylglyoxal oxidoreductase	11.29	1657	0.66
CJI97_000658	B9J08_000656	<i>MGD1</i>	NAD(H)-linked methylglyoxal oxidoreductase	8.53	4335	11.7
CJI97_004768	B9J08_004684	N/A	Role in histone deacetylation	7.87	51.9	0.22
CJI97_000946	B9J08_000928	<i>AQY1</i>	Aquaporin water channel, osmotic shock resistance	7.33	581	3.60
CJI97_004767	B9J08_004685	N/A	Curved DNA-binding protein	5.36	279	6.78
CJI97_003833	B9J08_003761	N/A	DNA topoisomerase	5.31	96.8	2.43
CJI97_002880	B9J08_002824	N/A	DNA replication licensing factor required for pre-replication complex assembly	4.49	316	14.1
CJI97_004042	B9J08_003981	<i>MDR1</i>	Plasma membrane MDR/MFS multidrug efflux pump	4.42	190	8.88
CJI97_002740	B9J08_002688	<i>FDH1</i>	Formate dehydrogenase	4.33	131	6.50
CJI97_004770	B9J08_004682	<i>ECM42</i>	Ornithine acetyltransferase	4.24	99.9	5.28
Genes more highly expressed in AR0390						
Locus Tag	AR0390 Locus Tag	Gene Name	Predicted Function	Log ₂ FC (B11221 vs AR0390)	B11221 average (CPM)	AR0390 average (CPM)
CJI97_001302	B9J08_001303	<i>BDF1</i>	Essential chromatin-binding bromodomain protein	-10.21	0.48	563
CJI97_004515	B9J08_004451	N/A	ALS family protein	-9.61	1.98	1546
CJI97_004556	B9J08_005565	<i>PRD1</i>	Proteinase	-7.29	0.14	22.6
CJI97_001865	B9J08_002322	<i>BMH1</i>	Role in morphology	-7.01	12.7	1639
CJI97_002974	B9J08_002900	<i>RMD9</i>	Mitochondrial protein with a predicted role in respiratory growth	-6.86	6.41	745
CJI97_003073	B9J08_003002	N/A	Iron permease	-6.51	36.8	3350
CJI97_002817	B9J08_002762	<i>INO1</i>	Inositol-1-phosphate synthase	-5.48	28.7	1281
CJI97_004514	B9J08_004450	<i>THI13</i>	Thiamin pyrimidine synthase	-4.95	2.34	72.4
CJI97_004654	B9J08_004798	<i>ARG3</i>	Ornithine carbamoyltransferase	-4.95	0.70	21.5
CJI97_000838	B9J08_000820	<i>SAM4</i>	S-adenosylmethionine-homocysteine methyltransferase	-4.43	1.93	41.5

Differentially expressed genes were determined using a cutoff of Log₂FC ≥ 1.00 or ≤ -1.00 and p-value < 0.05.

Table S4. Minimum inhibitory concentration (MIC) of fluconazole (FLZ) for *C. lusitaniae* U04 *mrr1*Δ and isogenic strains complemented with either *CauMRR1a*^{N647T} or *CauMRR1a* from *C. auris*

<i>C. lusitaniae</i> strain	FLZ MIC (μg/mL)	
	24h	48h
U04 <i>mrr1</i> Δ	4	4
U04 <i>mrr1</i> Δ + <i>CauMRR1a</i> ^{N647T} clone # 1	16	16
U04 <i>mrr1</i> Δ + <i>CauMRR1a</i> ^{N647T} clone # 2	16	16
U04 <i>mrr1</i> Δ + <i>CauMRR1a</i> ^{N647T} clone # 8	16	16
U04 <i>mrr1</i> Δ + <i>CauMRR1a</i> clone # 4	4	4
U04 <i>mrr1</i> Δ + <i>CauMRR1a</i> clone # 5	4	4
U04 <i>mrr1</i> Δ + <i>CauMRR1a</i> clone # 7	4	4

Table S5. Comparison of select genes induced by MG in B11221 and/or AR0390

B11221 Locus Tag	AR0390 Locus Tag	Gene Name	Predicted Function	B11221 Log ₂ FC	AR0390 Log ₂ FC
CJI97_005163	B9J08_005078	<i>AAH1</i>	Adenine deaminase, purine salvage and nitrogen catabolism	0.29	1.10
CJI97_002719	B9J08_002666	<i>AGP2</i>	Amino acid permease	0.45	1.16
CJI97_001242	B9J08_001362	<i>AGP3</i>	Serine transporter; sulfur assimilation	1.06	0.00
CJI97_004654	B9J08_004798	<i>ARG3</i>	Ornithine carbamoyltransferase	4.77	3.02
CJI97_003828	B9J08_003754	<i>ARG11</i>	Ornithine transporter of the mitochondrial inner membrane	0.94	1.70
CJI97_003954	B9J08_003882	<i>ARO4</i>	3-deoxy-D-arabinoheptulosonate-7-phosphate synthase; aromatic amino acid biosynthesis	2.50	2.22
CJI97_001997	B9J08_002453	<i>BAT21</i>	Branched chain amino acid aminotransferase	2.51	2.56
CJI97_005185	B9J08_005101	<i>BNA1</i>	3-hydroxyanthranilic acid dioxygenase; NAD biosynthesis	1.54	-0.33
CJI97_000479	B9J08_000479	<i>CDR4</i>	ABC transporter superfamily	1.34	1.01
CJI97_003095	B9J08_003024	<i>CIP1</i>	Oxidoreductase	0.51	1.30
CJI97_004156	B9J08_004088	<i>DRE2</i>	Cytosolic Fe-S protein assembly protein	1.09	2.52
CJI97_004181	B9J08_004118	<i>ERC1</i>	Xenobiotic transmembrane transporter	1.47	0.00
CJI97_002824	B9J08_002769	<i>FDH3</i>	Oxidoreductase and zinc ion binding activity	1.25	0.90
CJI97_005329	B9J08_005247	<i>HIS1</i>	ATP phosphoribosyl transferase; histidine biosynthesis	3.40	2.90
CJI97_001933	B9J08_002388	<i>HST6</i>	ABC transporter related to mammalian P-glycoproteins	0.66	1.18
CJI97_003449	B9J08_003374	<i>ICL1</i>	Isocitrate lyase; glyoxylate cycle enzyme	0.79	1.33
CJI97_000020	B9J08_000013	<i>ILV3</i>	Dihydroxyacid dehydratase	2.77	2.73
CJI97_004268	B9J08_004204	<i>JEN1</i>	Lactate transporter	1.84	-0.24
CJI97_001329	B9J08_001277	<i>LEU4</i>	2-isopropylmalate synthase	4.41	4.31
CJI97_001920	B9J08_002375	<i>LYS9</i>	Saccharopine dehydrogenase; lysine biosynthesis	1.13	0.52
CJI97_004689	B9J08_004763	<i>MET8</i>	Bifunctional dehydrogenase and ferrochelatase; siroheme biosynthesis	1.83	0.94
CJI97_003625	B9J08_003552	<i>MET10</i>	Sulfite reductase; sulfur amino acid metabolism	1.01	0.90
CJI97_000409	B9J08_000409	<i>MET13</i>	Methionine biosynthesis protein	0.04	1.65
CJI97_003066	B9J08_002995	<i>MET14</i>	Adenylylsulfate kinase; sulfur metabolism	1.08	0.93
CJI97_005391	B9J08_005307	<i>MET16</i>	3'-phosphoadenylylsulfate reductase; sulfur amino acid metabolism	1.63	1.75
CJI97_004042	B9J08_003981	<i>MDR1</i>	Plasma membrane MDR/MFS multidrug efflux pump	3.83	5.27
CJI97_000658	B9J08_000656	<i>MGD1</i>	NAD(H)-linked methylglyoxal oxidoreductase	1.35	2.04
CJI97_004624	B9J08_004828	<i>MGD2</i>	NAD(H)-linked methylglyoxal oxidoreductase	-0.06	1.60
CJI97_002730	B9J08_002677	<i>MIS12</i>	Mitochondrial C1-tetrahydrofolate synthase precursor	1.05	0.87
CJI97_004904	B9J08_004548	<i>NAR1</i>	Cytosolic iron-sulfur protein assembly machinery protein	0.84	2.13
CJI97_003488	B9J08_003413	<i>OPT7</i>	Oligopeptide transporter, may transport GSH or related compounds	1.09	0.90
CJI97_001481	B9J08_001125	<i>SNQ2</i>	Putative ABC transporter superfamily	1.50	2.25
CJI97_003300	B9J08_003225	<i>STR2</i>	Cystathionine gamma-synthase; sulfur compound metabolism	1.15	1.32

CJI97_001014	B9J08_000995	<i>SUL2</i>	Sulfate transporter	1.98	2.03
CJI97_004677	B9J08_004775	<i>TPO3</i>	Polyamine transporter, MFS-MDR family	0.91	1.24
CJI97_002495	B9J08_001834	<i>TPO4</i>	Spermidine transporter	0.48	1.49
CJI97_003424	B9J08_003349	<i>TRP5</i>	Tryptophan synthase	2.45	2.52
CJI97_002560	B9J08_001899	<i>YCF1</i>	Glutathione S-conjugate transporter	4.58	4.57
CJI97_005451	B9J08_005368	<i>YDJ1</i>	Type I HSP40 co-chaperone	0.25	1.20

Significantly induced genes were determined using a cutoff of Log₂FC ≥ 1.00 and p-value < 0.05.

Table S6. Comparison of select genes repressed by MG in B11221 and/or AR0390

B11221 Locus Tag	AR0390 Locus Tag	Gene Name	Predicted Function	B11221 Log₂FC	AR0390 Log₂FC
CJI97_001435	B9J08_001171	<i>ADH5</i>	Alcohol dehydrogenase	-1.54	-1.34
CJI97_002591	B9J08_001930	<i>AOX1</i>	Alternative oxidase, cyanide-resistant respiration	-1.31	-0.01
CJI97_004799	B9J08_004653	<i>ARP2</i>	Component of the Arp2/3 complex	-0.79	-1.11
CJI97_001469	B9J08_001137	<i>ARP3</i>	Protein with Myo5p-dependent localization to cortical actin patches at hyphal tip	-0.55	-1.05
CJI97_000089	B9J08_000084	<i>ATP14</i>	Mitochondrial F1F0 ATP synthase subunit	-0.45	-1.02
CJI97_002664	B9J08_002610	<i>ATP17</i>	Mitochondrial ATPase complex subunit	-0.33	-1.25
CJI97_003777	B9J08_003702	<i>CDG1</i>	Cysteine dioxygenases, role in conversion of cysteine to sulfite	-2.12	-2.65
CJI97_004336	B9J08_004273	<i>COX6</i>	Cytochrome c oxidase	-0.64	-1.03
CJI97_002184	B9J08_001993	<i>COX11</i>	Cytochrome oxidase assembly protein	-0.43	-1.43
CJI97_004817	B9J08_004635	<i>COX19</i>	Cytochrome c oxidase assembly protein	-0.39	-1.16
CJI97_002517	B9J08_001856	<i>CTR1</i>	Copper transporter	-1.13	-2.07
CJI97_005423	B9J08_005340	<i>ERG6</i>	Delta(24)-sterol C-methyltransferase, converts zymosterol to fecosterol, ergosterol biosynthesis	-1.01	-0.46
CJI97_005321	B9J08_005239	<i>FBA1</i>	Fructose-bisphosphate aldolase	-0.72	-1.13
CJI97_005247	B9J08_005163	<i>FBP1</i>	Fructose-1,6-bisphosphatase, key gluconeogenesis enzyme	-1.33	-1.48
CJI97_000015	B9J08_000008	<i>FRE7</i>	Ferric reductase	-1.23	-0.86
CJI97_003972	B9J08_004052	<i>FRE8</i>	Iron/copper reductase	-1.03	-0.23
CJI97_004532	B9J08_004468	<i>FRP1</i>	Ferric reductase	-1.41	-0.53
CJI97_002942	B9J08_002886	<i>GIT3</i>	Glycerophosphocholine permease	-1.18	-0.13
CJI97_003057	B9J08_002986	<i>GLC3</i>	1,4-glucan branching enzyme	-1.05	-0.93
CJI97_004438	B9J08_004375	<i>GPM1</i>	Phosphoglycerate mutase	-0.53	-1.40
CJI97_001045	B9J08_001025	<i>GSY1</i>	Glycogen synthase	-1.17	-1.36
CJI97_005108/ CJI97_005109	B9J08_005025	<i>HGT19</i>	MFS glucose/myo-inositol transporter	-1.71	-0.98
CJI97_002699	B9J08_002646	<i>HXT5</i>	Sugar transporter	0.13	-1.13
CJI97_002817	B9J08_002762	<i>INO1</i>	Inositol-1-phosphate synthase	-0.36	-1.41
CJI97_001658	B9J08_001652	<i>JAC1</i>	ATPase activator activity	-0.25	-1.04
CJI97_005579	B9J08_005497	<i>MAE1</i>	Mitochondrial malic enzyme	-1.44	-1.10
CJI97_000695	B9J08_000694	<i>MDH1</i>	Mitochondrial malate dehydrogenase	-1.23	-1.26
CJI97_002683	B9J08_002630	<i>MEP1</i>	Ammonium permease	-1.81	-1.38
CJI97_003663	B9J08_003590	<i>MIG2</i>	Transcription factor involved in glucose repression	-1.17	-1.44
CJI97_002101	B9J08_002556	<i>MIX14</i>	Role in aerobic respiration and mitochondrial intermembrane space localization	-0.20	-1.38
CJI97_002993	B9J08_002919	<i>MLS1</i>	Malate synthase, glyoxylate cycle enzyme	-0.84	-1.11
CJI97_001141	B9J08_001463	<i>MYO1</i>	Component of actomyosin ring at neck of newly emerged bud	-0.98	-1.22
CJI97_001117	B9J08_001487	N/A	Transporter of ferrochrome siderophores	-1.90	-1.06
CJI97_001762	B9J08_001547	N/A	Transporter of ferrochrome siderophores	-1.94	-1.07
CJI97_000596	B9J08_000675	N/A	Adhesin-like protein	-1.12	0.15

CJI97_002126	B9J08_002582	N/A	Adhesin-like protein	-1.48	1.00
CJI97_003987	B9J08_004037	N/A	Adhesin-like protein	-1.19	-0.35
CJI97_004240	B9J08_004176	N/A	Secreted lipase	-1.69	-0.78
CJI97_001776	B9J08_001533	N/A	NAD-aldehyde dehydrogenase	-1.57	-1.55
CJI97_003161	B9J08_003088	N/A	NAD-aldehyde dehydrogenase	-1.76	-1.89
CJI97_001793	B9J08_002250	N/A	MFS glucose transporter	-2.00	-1.41
CJI97_002024	B9J08_002481	N/A	MFS glucose transporter	-2.02	-1.52
CJI97_004566	B9J08_004886	N/A	Protein similar to ferric reductases and cupric reductases	-1.43	-1.43
CJI97_005148	B9J08_005064	N/A	Protein similar to ferric reductases and cupric reductases	-1.56	-1.84
CJI97_000696	B9J08_000695	<i>NTH1</i>	Neutral trehalase	-1.07	-0.98
CJI97_002722	B9J08_002669	<i>PCK1</i>	Phosphoenolpyruvate carboxykinase	-2.03	-2.01
CJI97_002521	B9J08_001860	<i>PGK1</i>	Phosphoglycerate kinase	-0.77	-1.19
CJI97_001140	B9J08_001464	<i>PHO84</i>	High-affinity phosphate transporter	-2.74	-2.72
CJI97_001580	B9J08_002202	<i>PHO89</i>	Phosphate permease	-1.48	-0.73
CJI97_001697	B9J08_001613	<i>PHO100</i>	Putative inducible acid phosphatase	-1.38	-1.60
CJI97_004666	B9J08_004786	<i>PIR1</i>	1,3-beta-glucan-linked cell wall protein	-0.44	-1.12
CJI97_002654	B9J08_002600	<i>PMM1</i>	Phosphomannomutase, enzyme of O- and N-linked mannosylation	-1.05	-0.94
CJI97_002321	B9J08_002130	<i>PUT1</i>	Putative proline oxidase	-2.49	-1.79
CJI97_004379	B9J08_004317	<i>PUT2</i>	Putative delta-1-pyrroline-5-carboxylate dehydrogenase	-1.01	-1.45
CJI97_002448	B9J08_001787	<i>RGT1</i>	Transcriptional repressor involved in the regulation of glucose transporter genes	-1.26	-1.62
CJI97_002974	B9J08_002900	<i>RMD9</i>	Mitochondrial protein with a predicted role in respiratory growth	-0.25	-1.09
CJI97_004415	B9J08_004352	<i>SAH1</i>	S-adenosyl-L-homocysteine hydrolase	-1.46	-1.22
CJI97_004940	B9J08_004512	<i>SCO1</i>	Copper transporter	-0.79	-1.06
CJI97_001499	B9J08_001107	<i>SEF1</i>	Zn2-Cys6 transcription factor, regulates iron uptake	-1.13	-0.81
CJI97_005617	B9J08_005567	<i>SHA3</i>	Ser/thr kinase involved in glucose transport	-1.14	0.10
CJI97_002536	B9J08_001875	<i>TPI1</i>	Triose-phosphate isomerase	-0.36	-1.42
CJI97_003198	B9J08_003126	<i>QCR8</i>	Ubiquinol cytochrome c reductase	-0.22	-1.12
CJI97_002481	B9J08_001820	<i>QCR10</i>	Ubiquinol-cytochrome-c reductase	-0.31	-1.01
CJI97_003997	B9J08_004027	<i>WOR1</i>	Transcription factor of white-opaque phenotypic switching in <i>C. albicans</i>	-1.22	-0.33

Significantly repressed genes were determined using a cutoff of $\text{Log}_2\text{FC} \leq -1.00$ and $p\text{-value} < 0.05$.

Table S7. Strains used in this study

Strain	Lab #	Species	Parent	Relevant Characteristics or Genotype	Source
Fungal Strains					
AR0390	DH2777	<i>C. auris</i>		Clinical isolate, clade I	(1)
B11221	DH3880	<i>C. auris</i>		Clinical isolate, clade III	(2)
<i>mrr1a</i> Δ	DH3881	<i>C. auris</i>	B11221	<i>mrr1a</i> Δ::caSAT1	(2)
<i>mrr1b</i> Δ	DH3882	<i>C. auris</i>	B11221	<i>mrr1b</i> Δ::caSAT1	(2)
<i>mrr1c</i> Δ	DH3883	<i>C. auris</i>	B11221	<i>mrr1c</i> Δ::caSAT1	(2)
U04 <i>mrr1</i> Δ	DH3306	<i>C. lusitaniae</i>	U04	<i>mrr1</i> Δ::NAT1	(3)
U04 <i>mrr1</i> Δ + <i>CauMRR1a</i> ^{N647T} clone #1	DH3914	<i>C. lusitaniae</i>	U04 <i>mrr1</i> Δ	<i>CauMRR1a</i> ^{N647T} -HygB	This study
U04 <i>mrr1</i> Δ + <i>CauMRR1a</i> ^{N647T} clone #2	DH3915	<i>C. lusitaniae</i>	U04 <i>mrr1</i> Δ	<i>CauMRR1a</i> ^{N647T} -HygB	This study
U04 <i>mrr1</i> Δ + <i>CauMRR1a</i> ^{N647T} clone #8	DH3916	<i>C. lusitaniae</i>	U04 <i>mrr1</i> Δ	<i>CauMRR1a</i> ^{N647T} -HygB	This study
U04 <i>mrr1</i> Δ + <i>CauMRR1a</i> clone #4	DH3917	<i>C. lusitaniae</i>	U04 <i>mrr1</i> Δ	<i>CauMRR1a</i> -HygB	This study
U04 <i>mrr1</i> Δ + <i>CauMRR1a</i> clone #5	DH3918	<i>C. lusitaniae</i>	U04 <i>mrr1</i> Δ	<i>CauMRR1a</i> -HygB	This study
U04 <i>mrr1</i> Δ + <i>CauMRR1a</i> clone #7	DH3919	<i>C. lusitaniae</i>	U04 <i>mrr1</i> Δ	<i>CauMRR1a</i> -HygB	This study
Plasmids in <i>E. coli</i> (DH5α)					
pMQ30 ^{MRR1-L1191H+Q1197*}	DH3829	<i>E. coli</i>		<i>MRR1</i> ^{L1191H+Q1197*} -HygB complementation, Gent ^R	(5)
pMQ30 ^{CauMRR1a} ^{N647T}	DH3912	<i>E. coli</i>		<i>CauMRR1a</i> ^{N647T} -HygB complementation, Gent ^R	This study
pMQ30 ^{CauMRR1a}	DH3913	<i>E. coli</i>		<i>CauMRR1a</i> -HygB complementation, Gent ^R	This study

Table S8. Oligonucleotides and primers used in this study

Name	Description	Sequence	Reference
ED222	<i>C. auris</i> ACT1 qRT Fwd	5' – GAA GGA GAT CAC TGC TTT AGC C – 3'	This study
ED223	<i>C. auris</i> ACT1 qRT Rev	5' – GAG CCA CCA ATC CAC ACA G – 3'	This study
ED224	<i>C. auris</i> MDR1 qRT Fwd	5' – GAA GTA TGA TGG CGG GTG – 3'	This study
ED225	<i>C. auris</i> MDR1 qRT Rev	5' – CCC AAG AGA GAC GAG CCC – 3'	This study
AB126	<i>C. auris</i> MGD1 qRT Fwd	5' – TTC CCC TGA AAT GGA TTT GA – 3'	This study
AB127	<i>C. auris</i> MGD1 qRT Rev	5' – GTC TTG GAG CCA TAG TAA CC – 3'	This study
AB130	Amplify <i>C. auris</i> MRR1a for heterologous complementation, Fwd	5' – CTT CAA CTC CGC AAC ACC TGG AAA CTT CAT TAC TAA AGA TGA TGG TAT CTT CGA AAG ATC – 3'	This study
AB131	Amplify <i>C. auris</i> MRR1a for heterologous complementation, Fwd	5' – CTT TAC CAG TAA AGT ATC CTT GCC AAA TTT CGT TCC ATA ATT ACA CAT CAA GCA TCT CTT C – 3'	This study
ED125	Forward upstream of MRR1 to validate complement	5' – GAA AAA GAA GCC AGC AGA CC – 3'	(5)
ED126	Reverse downstream of MRR1 to validate complement	5' – GGG TAA AGC CAT TGC AGA C – 3'	(5)
ACT1-F	<i>C. lusitaniae</i> ACT1 qRT Fwd	5' – GTA TCG CTG AGC GTA TGC AA – 3'	(6)
ACT1-R	<i>C. lusitaniae</i> ACT1 qRT Rev	5' – GAT GGA TGG TCC AGA CTC GT – 3'	(6)
ED058	<i>C. lusitaniae</i> MDR1 qRT Fwd	5' – TCC ATC CAT GGG TCC ATT ATT C – 3'	(3)
ED059	<i>C. lusitaniae</i> MDR1 qRT Rev	5' – CTC AAC ACA AGG AAA GCA CAT C – 3'	(3)
AB039	<i>C. lusitaniae</i> MGD1 qRT Fwd	5' – CGC AGA AAT CCC TAA AGT AAA T – 3'	(5)
AB040	<i>C. lusitaniae</i> MGD1 qRT Rev	5' – TAC CCT TTG CTT CGT TCT T – 3'	(5)
NAT1 crRNA	crRNA for NAT1; used to complement <i>C. auris</i> MRR1a alleles into <i>C. lusitaniae</i>	5' – GGG AAA ACC TTA GTC AAT GG – 3'	(5)

References

1. Pathirana RU, Friedman J, Norris HL, Salvatori O, McCall AD, Kay J, Edgerton M. 2018. Fluconazole-Resistant *Candida auris* Is Susceptible to Salivary Histatin 5 Killing and to Intrinsic Host Defenses. *Antimicrob Agents Chemother* 62.
2. Mayr EM, Ramirez-Zavala B, Kruger I, Morschhauser J. 2020. A Zinc Cluster Transcription Factor Contributes to the Intrinsic Fluconazole Resistance of *Candida auris*. *mSphere* 5.

3. Demers EG, Biermann AR, Masonjones S, Crocker AW, Ashare A, Stajich JE, Hogan DA. 2018. Evolution of drug resistance in an antifungal-naive chronic *Candida lusitanae* infection. *Proc Natl Acad Sci U S A* 115:12040-12045.
4. Demers EG, Stajich JE, Ashare A, Occhipinti P, Hogan DA. 2021. Balancing Positive and Negative Selection: *In Vivo* Evolution of *Candida lusitanae* *MRR1*. *mBio* 12.
5. Biermann AR, Demers EG, Hogan DA. 2021. Mrr1 regulation of methylglyoxal catabolism and methylglyoxal-induced fluconazole resistance in *Candida lusitanae*. *Mol Microbiol* 115:116-130.
6. Asner SA, Giulieri S, Diezi M, Marchetti O, Sanglard D. 2015. Acquired Multidrug Antifungal Resistance in *Candida lusitanae* during Therapy. *Antimicrob Agents Chemother* 59:7715-22.