Title: Mycorrhizal fungi stabilize bacterial growth and diversity in water-limited soils

Running Title: Mycorrhizae stabilize bacteria in water-limited soils

Rachel Hestrin^{1*}, Megan Kan¹, Marissa Lafler¹, Jessica Wollard¹, Jeffrey A. Kimbrel¹, Prasun Ray², Steven Blazewicz¹, Rhona Stuart¹, Kelly Craven³, Mary Firestone⁴, Erin Nuccio¹, Jennifer Pett-Ridge^{1,5*}

¹*Lawrence Livermore National Laboratory, Physical and Life Science Directorate, Livermore, CA 94550 USA*

²*Department of Natural Resources, University of Maryland Eastern Shore, Princess Anne, MD 21853 USA*

³*Noble Research Institute, Plant Biology Division, Ardmore, OK 73401 USA*

⁴*University of California Berkeley, Department of Environmental Science Policy and Management, Berkeley, CA 94720 USA*

⁵*University of California Merced, Life & Environmental Sciences Department, Merced, CA 95343 USA*

*Correspondence to:

Rachel Hestrin, 925-409-7954, hestrin1@llnl.gov

Jennifer Pett-Ridge, 925-337-7618, pettridge2@llnl.gov

Competing Interests: The authors declare no competing interests.

Abstract

Drought disrupts soil microbial activity and many biogeochemical processes. Although mycorrhizal fungi are known to impact plant functions and nutrient cycling during drought, their effects on drought-exposed soil microbial communities are not well resolved. We used H₂¹⁸O quantitative stable isotope probing (qSIP) to investigate microbial response to water limitation in the hyphospheres of two distinct mycorrhizal lineages (*Rhizophagus irregularis* and *Serendipita bescii*) grown with the bioenergy model grass *Panicum hallii*. In the absence of mycorrhizal inocula, water limitation resulted in significantly lower bacterial growth rates and lower diversity in the actively growing bacterial community. Water limitation also decoupled growth from CO₂ efflux, resulting in a threefold lower growth efficiency. In contrast, both mycorrhizal lineages appeared to protect bacterial communities exposed to water limitation: bacterial growth rates, growth efficiency, and the diversity of the active bacterial community were similar in water-replete and water-limited soils inoculated with either fungus. Several of the bacterial taxa that responded positively to mycorrhizal inocula in water-limited soil belong to lineages that are considered drought-susceptible. Together, these results indicate that mycorrhizal-bacterial interactions can stabilize bacterial communities in water-limited environments. As drought frequency and severity increase, these synergistic relationships may support ecosystem resilience to moisture stress.

Keywords: *Panicum hallii*, mycorrhizal fungi, bacteria, drought, quantitative stable isotope probing, plant-microbe interactions, ¹⁸O-water

Introduction

Drought is a destabilizing stressor that alters plant productivity¹, microbial community composition²⁻⁴, microbial biomass^{5,6}, greenhouse gas emissions^{7,8}, and many critical biogeochemical processes⁹. Plant-microbial mutualisms mitigate plant drought response and aid in post-drought recovery through a variety of mechanisms^{10,11}. In particular, mycorrhizal fungi—which associate with the majority of terrestrial root systems¹²—can support plants during drought by facilitating water transport¹³, soil aggregation^{14,15}, root growth¹⁶, plant nutrient uptake¹⁷, photosynthesis^{18,19}, and stomatal conductance¹⁸⁻²⁰. While mycorrhizal fungi may also influence soil microbial communities in water-limited environments, multipartite feedbacks during and after drought remain poorly quantified¹¹.

There is increasing evidence that the mycorrhizal hyphosphere—the region of soil that surrounds mycorrhizal fungi—is a hotspot for multipartite interactions. These interactions may be synergistic or antagonistic, with significant effects on resource exchange^{21,22}, stress tolerance²³, and organismal abundance and activity^{24,25}. Recent work indicates that specific bacterial lineages are more abundant in the presence of arbuscular mycorrhizal (AM) fungi²⁶⁻²⁸ and that fungal traits shape the composition of mycorrhizal-associated bacterial communities²⁸⁻³⁰. Together, these observations suggest that mycorrhizal-bacterial interactions have important ecological functions. Still, relatively little is known about how mycorrhizal fungi influence bacterial communities exposed to environmental stressors such as moisture limitation. A better understanding of these relationships would help us predict microbial response to global change,

harness beneficial plant-microbial relations for sustainable agriculture, and assess other applications of mycorrhizal-microbial mutualisms.

In situ investigation of mycorrhizal-bacterial interactions presents immense methodological challenges. The mycorrhizal hyphosphere is small, difficult to sample, and may not exert a detectable effect on microbial community composition or activity when assessed at a ‘bulk’ scale²⁸. Furthermore, a substantial proportion of soil DNA may represent inactive organisms or extracellular ‘relic’ DNA³¹⁻³². Stable isotope probing (SIP) coupled with 16S rRNA gene profiling can distinguish active organisms from inactive ones by tracing isotope incorporation into newly-synthesized microbial DNA^{33,34}. With quantitative SIP (qSIP), we can estimate growth rates of amplicon sequence variants (ASVs) based on shifts in DNA buoyant density caused by heavy isotope incorporation^{35,36}. This enables sensitive detection of actively growing taxa, even in samples where a substantial quantity of DNA belongs to dead or dormant organisms. In challenging experimental systems like the mycorrhizal hyphosphere, qSIP has the potential to provide novel insight into taxon-specific interactions.

In this study, we sought to investigate how two mycorrhizal lineages—the AM fungus *Rhizophagus irregularis* and the Sebaciniales fungus *Serendipita bescii*—mediate bacterial response to water limitation in a marginal soil planted with the switchgrass bioenergy model *Panicum hallii* (Hall’s panicgrass). For the purpose of this study we define ‘mycorrhiza’ as a symbiotic relationship between a fungus and a plant root, where the fungus acquires C from the living plant in exchange for mineral nutrients or other resources¹². Both AM and Sebaciniales fungi associate with a wide range of plant species, including switchgrass³⁷⁻³⁹, and support plant

function during drought⁴⁰⁻⁴². However, *R. irregularis* has a reduced genome and is dependent upon host-provided C^{43,44} and microbial transformation of nutrient sources into bioavailable forms^{45,46}. In contrast, *S. bescii* and other Serendipita lineages are facultative symbionts with a broad enzymatic repertoire that enables direct resource acquisition from both living plants and detritus⁴⁷⁻⁴⁹. We hypothesized that both fungi would mitigate moisture stress, but that distinct bacterial communities would respond to each fungus. Our results demonstrate that mycorrhizal fungi have a stabilizing effect on bacterial communities in water-limited soil.

Materials and Methods

Soil collection and characterization

Soil used for this study was a Pond Creek fine sandy loam, classified as a superactive, thermic Pachic Argiustoll⁵⁰. Soil was collected from a pasture in Caddo County, OK (35.072417/-98.303667) where switchgrass is endemic, on traditional land of the Anadarko (Nadaco) tribe and the Caddo Nation of Oklahoma. The soil is considered ‘marginal’ due to its high sand content (69%), low C, N, and P content (< 0.4%, < 0.04%, and < 6 ppm, respectively); its pH is ~5 (ref. 51). Surface soil (0-20 cm) was collected in May 2019, transported to Livermore, CA, sieved to 2 mm, and stored at 4 °C. A soil moisture retention curve was generated from air-dried soil using a pressure plate apparatus (WP4C, METER Environment, Pullman, WA) and the nonlinear fitting program SWRC-Fit to apply a Brooks and Corey model to the data⁵².

Mycorrhizal inoculum

Spores of *R. irregularis* (formerly *Glomus intraradices*) isolate DAOM-197198 were purchased from Premier Tech (Rivière-du-Loup, Quebec, Canada). *S. bescii* (sourced from the Noble Research Institute) was grown in modified Melin-Norkran's (MMN) broth. Bentonite clay particles were mixed with MMN broth, inoculated with *S. bescii*, and incubated at 24 °C for 8 weeks according to methods described by Ray et al. (ref. 53). Uninoculated bentonite clay particles were also mixed with MMN broth and incubated under the same conditions.

Greenhouse water limitation experiment

P. hallii seeds collected from the Edwards Plateau in central Texas were grown to seed, scarified, surface sterilized, stratified at 4 °C for one week, and germinated in petri plates. After 5 days, germinated seedlings were transferred into planting cones (Ray Leach Cone-tainers, Steuwe & Sons, Tangent, OR, USA) filled with double-autoclaved sand. Five days after transfer into cones, ~500 *R. irregularis* spores or 500 µL *S. bescii* inoculum were injected into the sand to inoculate the *P. hallii* seedlings. A subset of seedlings was left uninoculated. Eight weeks after inoculation, seedlings were transplanted into 960 cm³ containers (Anderson Plant Bands, Steuwe & Sons) filled with a 50:50 (v:v) mixture of 'live' soil and double-autoclaved sand (CEMEX Lapis Lustre Specialty Sand, #2/12) packed to 1.7 g mL⁻¹ bulk density at 15% moisture. Additional inoculum (consisting of 500 *R. irregularis* spores or 5 g bentonite clay particles coated with *S. bescii*) were placed near the roots of seedlings as they were transplanted. Five g of uncoated bentonite clay particles were placed near the roots of previously uninoculated plants and plants that had been inoculated with *R. irregularis*. Each microcosm contained a 25 µm mesh hyphal ingrowth core (2.3 cm diameter) filled with 75 g of the same 'live' soil (no sand) mixed with 0.5 g finely milled switchgrass biomass as mycorrhizal bait. The multi-phase mycorrhizal inoculation procedure

was intended to give *R. irregularis* and *S. bescii* a colonization advantage over other native mycorrhizal lineages or endophytes present in the soil.

After assembly, each microcosm was covered with 150 g double-autoclaved sand to prevent cross-contamination of mycorrhizal inocula. All microcosms were watered with a total of 20 mL ultrapure H₂O during the first week to facilitate plant and mycorrhizal establishment. Half of the microcosms were watered on a weekly or bi-weekly schedule to maintain soil moisture at approximately 15% (based on gravimetric measurements), a level that would not restrict plant growth. The other half of the microcosms were not watered for the remainder of the experiment and declined to 5% gravimetric soil moisture by 3 months. The moisture content of the soil surrounding the roots (i.e., not directly within the hyphal ingrowth cores) in 1 microcosm per treatment was monitored continuously with a volumetric water probe (ECH₂O EC-5, METER Group, Inc., Pullman, WA, USA). Another microcosm per treatment was weighed weekly to assess whole-microcosm gravimetric changes. Three replicate microcosms were maintained for each of the 6 treatment combinations (3 mycorrhizal inocula x 2 moisture regimes). Average daytime and nighttime temperatures were 27 °C and 24 °C, respectively, with a photoperiod of 16 h. After 3 months, the microcosms were destructively harvested. Soil from hyphal ingrowth cores was homogenized, flash frozen in liquid N₂ and stored at -80 °C for DNA extractions or at room temperature for qSIP assays (see Fig. 1a for experimental design).

Soil characteristics

Soil moisture content following the 3-month harvest was determined by mass difference after drying soil for 48 h at 105 °C. Total soil C and N contents were measured with a Costech ECS 4010 Elemental Analyzer (Costech Analytical Technologies Inc., Valencia, CA).

Soil H₂¹⁸O qSIP assay

Three days after the 3-month harvest, soil from 3 hyphal ingrowth cores per treatment was amended with either H₂¹⁶O or H₂¹⁸O (Fig. 1a). Each soil was divided into three subsamples: one 3.0 g dry weight equivalent sample for an initial H₂¹⁶O qSIP timepoint (“T0”) and two 4.0 g dry weight equivalent samples for H₂¹⁶O and H₂¹⁸O 7-day qSIP timepoint (“T7”). This resulted in a total of 54 samples (18 T0 H₂¹⁶O samples, 18 T7 H₂¹⁶O samples, and 18 T7 H₂¹⁸O samples). Each subsample was air-dried to 0.047 gravimetric water content in a biosafety cabinet and then brought up to 22.13% soil moisture (60% field capacity) with either ultrapure water at natural isotopic abundance or water enriched with ¹⁸O (98.38 atom % H₂¹⁸O, Isotex, San Francisco, CA, USA). The final enrichment of the samples containing H₂¹⁸O was 78.76 ¹⁸O atom %. Standardization of soil moisture across treatments was necessary for qSIP calculations. The T0 subsample (amended with H₂¹⁶O) was immediately flash frozen in liquid N₂ and stored at -80 °C. Each of the T7 soil samples was stored in the dark inside a separate 16 oz glass jar with a tight-fitting lid containing a septum for gas sampling. After 7 days, the soils were flash frozen in liquid N₂ and stored at -80 °C.

CO₂ efflux measurement

15 mL headspace gas samples were collected at the beginning and end of the qSIP assay. CO₂ concentrations were measured on a gas chromatograph equipped with a thermal conductivity detector (GC-14A, Shimadzu, Columbia, MD).

DNA extraction and qPCR

DNA was extracted from each soil sample in quadruplicate with the Qiagen PowerSoil Pro kit and then pooled per sample prior to downstream analysis. To assess the relative abundance of mycorrhizal inocula, we quantified *R. irregularis* and *S. bescii* gene copy number with qPCR primers designed specifically for each fungal lineage^{53,54} (see Table S1 and SI text for additional information). All sample and standard curve qPCR reactions were performed in triplicate. We note that DNA can be unevenly distributed throughout fungal biomass and while not necessarily indicative of total biomass or activity is a relative indicator of inoculation success^{55,56}.

Density gradient fractionation of DNA

To separate isotopically enriched DNA from unenriched DNA, samples were subjected to a cesium chloride density gradient formed in an ultracentrifuge as previously described^{57,58} with the following minor modifications. For each sample, 5 µg of DNA in 150 µL 1xTE was mixed with 1.00 mL gradient buffer, and 4.60 mL CsCl stock (1.885 g mL⁻¹) with a final average density of 1.730 g mL⁻¹. Samples were loaded into 5.2 mL ultracentrifuge tubes and spun at 20 °C for 108 hours at 176,284 RCF_{avg} in a Beckman Coulter Optima XE-90 ultracentrifuge using a VTi65.2 rotor. Sample fractionation was automated using Lawrence Livermore National Laboratory's high-throughput SIP pipeline, which automates the fractionation and clean-up tasks for the density gradient SIP protocol. The content of each ultracentrifuge tube was fractionated

into 22 fractions (~236 μL each) using an Agilent Technologies 1260 isocratic pump to deliver water at 0.25 mL min^{-1} through a 25G needle inserted through the top of the ultracentrifuge tube. Each tube was mounted in a Beckman Coulter fraction recovery system with a side port needle inserted through the bottom. The side port needle was routed to an Agilent 1260 Infinity fraction collector, and fractions were collected in 96-well deep well plates. The density of each fraction was measured using a Reichart AR200 digital refractometer fitted with a prism covering to facilitate measurement from $5 \mu\text{L}$, as previously described⁵⁹. DNA in each fraction was purified and concentrated using a Hamilton Microlab Star liquid handling system programmed to automate glycogen/PEG precipitations³⁴. Washed DNA pellets were suspended in $40 \mu\text{L}$ of 1xTE and the DNA concentration of each fraction was quantified using a PicoGreen fluorescence assay.

DNA sequencing

For each sample, we sequenced the 16S rRNA gene in unfractionated DNA as well as DNA fractions containing $> 1.0 \text{ ng } \mu\text{L}^{-1}$, resulting in 6-11 sequenced fractions per sample. The V4 region of the 16S rRNA gene was amplified with the 515F/806R primer pair⁶⁰, processed and barcoded through the Illumina V2 PE150 sample preparation kit, and sequenced on an MiSeq v2 platform in 3 runs (Illumina, Inc., San Diego, CA; see SI text and Tables S2 & S3 for additional information about taxonomic assignment and ASV recovery).

Quantitative stable isotope probing (qSIP) analysis

We calculated taxon-specific growth using the ^{18}O qSIP approach^{33,35}. Briefly, the qSIP mathematical model estimates taxon-specific ^{18}O incorporation into DNA based on the shift in

DNA density following exposure to natural abundance H_2O (H_2^{16}O) or ‘heavy’ H_2^{18}O . We determined baseline densities for each taxon in the absence of an isotopic tracer (i.e., H_2^{16}O), because even without tracer assimilation, DNA density varies by GC content⁶¹. For each ASV, we calculated median ^{18}O atom percent excess (APE) and gross growth rate. To increase confidence in these calculations, we first filtered out ASVs that were not recovered from all T0 and T7 replicates in a particular treatment.

Since ^{18}O incorporation could have occurred at any point during the qSIP assay, ^{18}O APE represents an integrated measure of activity. We note that a taxon’s ^{18}O APE value may be high due to an increase in abundance (i.e., cells incorporate ^{18}O into DNA during replication; new cell production increases total population size and average DNA ^{18}O content) or population turnover (i.e., DNA from older unenriched cells degrades, while new cells grow and incorporate ^{18}O into DNA). During turnover, DNA ^{18}O APE can increase while population-level abundance either remains similar (if mortality matches growth rate) or declines (if mortality outpaces growth). Therefore, a high ^{18}O APE value is not necessarily indicative of a net increase in abundance.

Statistical analyses

We performed all statistical analyses in R⁶². We assessed differences in mycorrhizal gene copy number with a Wilcoxon rank sum test with a Benjamini-Hochberg correction for multiple comparisons. We assessed differences in soil moisture, soil C and N content, Inverse Simpson’s diversity index, median ASV ^{18}O APE, and CO_2 efflux with a Tukey’s HSD test comparing the means of all treatments. First, we created a linear model with a fixed effect for treatment. If the raw data did not meet the assumptions of normality, it was log-transformed prior to further

analysis. We calculated growth efficiency by dividing gross growth by CO₂ efflux per treatment⁶³. We used ‘vegan’ to perform a principal coordinates analysis and permutational multivariate analysis of variance (PERMANOVA) on the weighted UniFrac distance between communities present under different treatments and qSIP timepoints^{64,65}. We used ‘metagenomeSeq’⁶⁶ with a wrench normalization for sparse data to identify significant differences in ASV relative abundance at T0 ($p < 0.05$ and $\log_2$ fold change > 2 or < -2). To compare taxon-specific growth between two treatments, we calculated ¹⁸O-based ‘growth ratios’ by dividing an ASV’s ¹⁸O APE in one treatment by its ¹⁸O APE in the second treatment. We used a Wilcoxon signed rank test with a Benjamini-Hochberg correction for multiple comparisons to assess whether mean phylum ratios were significantly greater than or less than 1.

Results

Mycorrhizal abundance and soil characteristics following 3-month water manipulation

After 3 months of *P. hallii* growth, *R. irregularis* and *S. bescii* abundances (measured with strain-specific qPCR primers) were significantly higher in the hyphal ingrowth cores of microcosms inoculated with each fungus, respectively (Fig. 1b; $p < 0.1$). Soil moisture was lower in the hyphal ingrowth cores of water-limited versus water-replete microcosms ($4.9\% \pm 0.85$ and $15.2\% \pm 0.57$, respectively, Fig. 1c; $p < 0.001$), and was not significantly associated with mycorrhizal abundance ($p > 0.05$). Matric suction was also lower in water-limited hyphal ingrowth core soils (Fig. S1). Total soil C and N did not vary significantly between treatments (Fig. S2; $p > 0.05$).

Moisture and mycorrhizal effects on microbial community structure

We assessed the effects of moisture regime and mycorrhizal inoculum on microbial community structure with both traditional and qSIP-enabled 16S rRNA gene profiling (Tables S4-S6 & Fig. 2a,b). The differences we detected in mycorrhizal-inoculated samples represent the influence of *R. irregularis* or *S. bescii* in soils containing a background microbial community. A PERMANOVA of weighted UniFrac distances between the ‘total’ communities (i.e., 16S rRNA gene profiles sequenced from unfractionated DNA) suggests that moisture regime and mycorrhizal inoculum explained 29.5% of variation in community structure at the end of the qSIP assay (Table S4 & Fig. 2a; $p < 0.05$). Although there was a small shift in community structure during the qSIP assay (6.3% of variance explained; Table S5; $p < 0.001$), moisture and mycorrhizal effects persisted throughout the assay (Table S5; $p < 0.05$).

When we used qSIP to filter out inactive ASVs (i.e., those that did not incorporate significant quantities of ^{18}O) prior to conducting a PERMANOVA, we found that moisture regime, mycorrhizal inoculum, and their interactive effect explained over 75% of the variation in the structure of the actively growing community (20.4%, 27.4%, and 27.5%, respectively; Table S6 & Fig. 2b; $p < 0.001$). This is more than twice the effect size observed in the ‘total’ community identified through traditional 16S rRNA analysis. qSIP-based filtering also highlighted differences in alpha diversity. While there was no relationship between moisture regime or mycorrhizal inoculum and the alpha diversity of the ‘total’ communities sequenced (Fig. 2c; $p > 0.05$), water limitation was associated with lower alpha diversity of the actively growing community in uninoculated soils ($p < 0.01$), no difference in *R. irregularis*-inoculated soils ($p > 0.05$), and higher alpha diversity in soils inoculated with *S. bescii* ($p < 0.01$).

We conducted differential abundance analysis to evaluate how water limitation influenced ASV relative abundance in the presence or absence of mycorrhizal inocula. Within each mycorrhizal treatment, some ASVs were more abundant in water-limited soils while others were less abundant (Fig. S3). In uninoculated soils, 56 ASVs were less abundant following water limitation while only 24 were more abundant, contributing to a twofold difference in the magnitude of cumulative $\log_2$ fold change (Fig. S3a,d). In contrast, within *R. irregularis*- and *S. bescii*-inoculated soils, the number of ASVs that were more abundant following water limitation was similar to the number that were less abundant, contributing to a similar magnitude of cumulative $\log_2$ fold change (Fig. S3b,c,d).

Moisture and mycorrhizal effects on microbial growth and efficiency

We used H_2^{18}O qSIP and CO_2 efflux measurements to quantify the effects of moisture regime and mycorrhizal inoculum on ASV ^{18}O incorporation, gross growth rates, and growth efficiency. Overall, DNA from soil amended with H_2^{18}O was denser than DNA from soil amended with H_2^{16}O (Fig. S4), indicating that many organisms incorporated ^{18}O into newly-synthesized DNA. Taxon-specific ^{18}O APE ranged from unenriched to 64.0 APE, with a median of 6.0. Between 35-59% of the total ASVs detected per treatment were significantly enriched (Table S3; lower 90% CI > 0); we refer to these as the ‘actively growing’ taxa.

In the absence of mycorrhizal inocula, microbial ASVs incorporated less ^{18}O in water-limited soils: median ASV ^{18}O APE was 2.14 compared to 9.06 in water-replete soils (Fig. 3a; $p < 0.05$). Gross growth was also lower in water-limited compared to water-replete soil (Fig. 3b). CO_2

efflux from uninoculated soils did not vary significantly between moisture regimes (Fig. 3c; $p > 0.05$). However, growth efficiency (gross growth divided by CO_2 efflux) was more than threefold lower in water-limited soil: only $8.4 (\pm 0.24) \mu\text{g DNA mg}^{-1} \text{CO}_2$ compared to $28.3 (\pm 2.16)$ in water-replete soils (Table S7). In contrast, microbial communities in mycorrhizal-inoculated soils were less affected by water limitation. Microbial ^{18}O APE was 6.56 compared to 7.45 in water-limited versus water-replete *R. irregularis*-inoculated soils (Fig. 3a; $p > 0.05$), and 5.67 compared to 4.85 in water-limited versus water-replete *S. bescii*-inoculated soils (Fig. 3a; $p > 0.05$). Gross growth was also similar under both moisture regimes in mycorrhizal-inoculated soils (Fig. 3b). CO_2 efflux from *S. bescii*-inoculated soils was higher than from *R. irregularis*-inoculated soils, but the difference was only significant between water-replete soils (Fig. 3c; $p < 0.05$). Compared to uninoculated soils, moisture effects on growth efficiency in mycorrhizal-inoculated soils were relatively small. Growth efficiency was $14.4 (\pm 0.8) \mu\text{g DNA mg}^{-1} \text{CO}_2$ compared to $20.2 (\pm 1.25)$ in water-limited versus water-replete soil inoculated with *R. irregularis*, and $12.5 (\pm 0.51)$ compared with $12.7 (\pm 0.75)$ in water-limited versus water-replete soils inoculated with *S. bescii* (Table S7).

Taxon-specific response to moisture regime and mycorrhizal inocula

Since compensatory dynamics may result in the increased activity of some organisms while others decline⁶⁷, we calculated ^{18}O -based ‘growth ratios’ for each actively growing ASV detected in both water-limited and water-replete soil. A ratio of greater than 1.0 suggests that the ASV sustained a higher growth rate in water-limited soil, a ratio of less than 1.0 suggests that the ASV sustained a lower growth rate in water-limited soil, and a ratio indistinguishable from 1.0 suggests that the ASV maintained a similar growth rate under both conditions. In the absence of

mycorrhizal inocula, most ASVs sustained lower growth rates in water-limited soil (Fig. 4a). When averaged at the phylum level, growth ratios were significantly less than 1 for 8 of the 10 most abundant phyla: Acidobacteria, Actinobacteria, Bacteroidetes, candidate division WPS-1, Gemmatimonadetes, Planctomycetes, Proteobacteria, and Verrucomicrobia (Fig. 4a; adjusted $p < 0.05$). In contrast, bacterial response to water limitation in mycorrhizal-inoculated soils was equivocal: many ASVs sustained lower growth rates, while many others sustained similar or greater growth rates. When averaged at the phylum level, this resulted in almost no significant effects of water limitation on microbial growth in mycorrhizal-inoculated soil (Fig 4b,c; adjusted $p > 0.05$ for most phyla). An exception to this pattern were the Acidobacteria, which sustained significantly higher growth rates in water-limited versus water-replete soils inoculated with *R. irregularis* (Fig. 4b; adjusted $p < 0.05$).

We also calculated ^{18}O -based growth ratios for ASVs detected in both uninoculated and mycorrhizal-inoculated water-limited soils. This allowed us to compare how mycorrhizal inocula influenced the growth of individual ASVs in water-limited soil. Most ASVs sustained higher growth rates in the presence of *R. irregularis* or *S. bescii* than in their absence (Fig. 5). Averaged at the phylum level, populations belonging to the phyla Acidobacteria, Actinobacteria, Bacteroidetes, candidate division WPS-1, Planctomycetes, Proteobacteria, and Verrucomicrobia were significantly more active in *R. irregularis*-inoculated soil (Fig. 5a,c; adjusted $p < 0.05$). In *S. bescii*-inoculated soils, populations from the phyla Actinobacteria and Proteobacteria were significantly more active than in uninoculated soil (Fig. 5b,d; adjusted $p < 0.05$). Some ASVs responded positively to both *R. irregularis* and *S. bescii*; others responded positively only to one

fungal lineage (Fig. 5c,d). Overall, the magnitude of positive growth response was greater in *R. irregularis*- than in *S. bescii*-inoculated soils.

Positive bacterial response to mycorrhizal inocula was also apparent in ASVs that did not grow under multiple conditions and therefore could not be compared via the ratios described above. For example, there were no Firmicutes ASVs with detectable growth rates in uninoculated water-limited soils (Fig. S5). However, several Firmicutes were active in mycorrhizal-inoculated soils regardless of moisture regime. More than double the number of Firmicutes ASVs were active in *R. irregularis*- compared to *S. bescii*-inoculated soils.

Discussion

Mycorrhizal fungi are well-known to improve plant drought tolerance^{16-20,40-42}, but their effects on bacterial communities in droughted soils are poorly quantified. With H₂¹⁸O qSIP, we show that mycorrhizal fungi protect bacterial communities from the suppressive and destabilizing effects of water limitation.

Water limitation suppresses bacterial growth and efficiency

Our findings suggest that water limitation suppresses taxon-specific bacterial growth rates and decouples growth from CO₂ efflux. Previous studies have shown that microbial carbon use efficiency is lower in water-limited soils^{68,69}. This is consistent with a microbial stress response, wherein cells prioritize essential metabolic activities over replication^{9,57}. Comparative transcriptomics analysis of drought-selected microbial communities has shown that there are trade-offs between stress tolerance and growth⁷⁰. Slowed growth helps bacteria persist in the

presence of antibiotics^{71,72}, which can accumulate in dry soils as microorganisms compete for limited resources⁷³. There is also evidence that lower microbial growth efficiency in dry soils may be due to reduced substrate diffusion^{63,74}. Our observations indicate that water limitation reduces microbial resource use efficiency and capacity to produce new biomass. Because microbial biomass is an important precursor to soil organic matter⁷⁵, this may impact C storage in droughted soils.

Growth suppression was apparent across a broad range of bacterial lineages present following water limitation, including taxa that are considered to be drought tolerant. Several previous studies have reported increases in monoderm (e.g., Actinobacteria, most Firmicutes, and some Chloroflexi) compared to diderm (e.g., most Acidobacteria, Bacteroidetes, Proteobacteria, and Verrucomicrobia) abundance and activity in dry soils^{2,9,76-78}. Monoderm physiology (i.e., thick cell walls and lack of an outer membrane) protects against oxidative damage under dry conditions^{9,76,77,79}. Monoderms can also outcompete other organisms through their capacity to produce antibiotics^{73,77,80} or utilize complex C substrates that remain available following water limitation^{4,73}. However, we did not find evidence that monoderm growth rates were higher than those of diderms in water-limited soil. Taxa belonging to the monoderm phylum Actinobacteria may remain metabolically active but replicate more slowly than other organisms⁸¹. This lifestyle could explain the decoupling of growth and CO₂ efflux that we observed in water-limited soils.

Mycorrhizal fungi stabilize bacterial response to soil moisture

The AM fungus *R. irregularis* and the Sebaciniales fungus *S. bescii* can substantially enhance plant performance in water-limited soils⁴⁰⁻⁴², thereby stabilizing plant communities during stress.

We found that both fungi also appeared to protect bacterial communities exposed to water limitation. Overall, bacterial growth and efficiency were similar in water-replete and water-limited soils inoculated with either fungus, and always higher in mycorrhizal-inoculated soils than in uninoculated soils. This ‘protective’ mycorrhizal effect extended broadly to the bacterial community, including many taxa that are considered drought-susceptible. While previous studies report a decrease in the relative abundances of Bacteroidetes, Planctomycetes, Verrucomicrobia, and many Proteobacteria and Acidobacteria in droughted soils^{2-4,82}, we found that in the presence of mycorrhizal inocula, many ASVs belonging to these phyla sustained similar growth rates in water-replete and water-limited soils. Mycorrhizal fungi could help bacteria tolerate water limitation through enhanced resource provisioning^{21,83-85}, soil structure^{14,15}, or biofilm formation⁸⁶. By supporting microbial diversity and activity in water-limited soil, mycorrhizal-bacterial synergies may counteract the destabilizing effect of environmental stress⁸⁷ and improve capacity for recovery.

In contrast to their synergistic effect in water-limited soils, *R. irregularis* and *S. bescii* appeared to suppress bacterial growth and diversity in water-replete soils. Previous reports show that mycorrhizal fungi can compete with other biota for N⁸⁸ and P^{24,25}, that putative bacterial predators are more abundant on extraradical mycorrhizal hyphae than in surrounding soil²⁸, and that soil filtrates containing bacteria inhibit mycorrhizal proliferation⁸⁹. Our results indicate that there is a trade-off between mycorrhizal and bacterial activity. While this trade-off may reduce maximum bacterial productivity under non-limiting conditions, it supports resilience under limiting conditions. Similar context-dependency is well-documented in other mutualisms⁹⁰. We

interpret the context-dependency of mycorrhizal-bacterial relationships as further indication of their stabilizing ecological effect.

Magnitude of microbial response is mycorrhizal lineage-dependent

While both mycorrhizal lineages stabilized bacterial response to soil moisture, *R. irregularis* elicited a stronger positive response than *S. bescii*—both with respect to the number of ASVs and the magnitude of their response. Previous studies show that distinct microbial consortia associate with different mycorrhizal lineages^{28,29}. Although empirical evidence remains sparse, different mycorrhizal exudate profiles, growth habits, and other functional traits may shape surrounding microbial community composition and activity^{45,91}, a phenomenon documented more extensively for root-microbial interactions⁹²⁻⁹⁴. Reduced bacterial growth rate and efficiency in *S. bescii*-compared to *R. irregularis*-inoculated soils may be due to *S. bescii*'s wider enzymatic repertoire, which could accelerate decomposition (as indicated by higher CO₂ efflux) or contribute to greater competition with other soil biota. It is also possible that greater abundance of *R. irregularis* compared to *S. bescii* contributed to the different magnitude of bacterial response. Together, these findings indicate that different mycorrhizal types elicit similar, but distinct bacterial response to water limitation, with significant effects on microbial community composition and growth rates.

H₂¹⁸O qSIP highlights microbial response

Previous studies report conflicting effects of drought on microbial biomass^{5,6,95,96}, activity^{5,6,97,98}, and community composition^{2-4,82}. These variable responses may be due to differences in drought severity, biota examined, and experimental methods employed^{10,99}. Previous drought exposure

can attenuate microbial response to additional water limitation². With qSIP, we detected a strong negative microbial response to water limitation in soils that had repeatedly experienced below-average rainfall during the preceding decade¹⁰⁰. This response was less apparent in our traditional microbial community analyses. For example, alpha diversity seemed unaffected by water limitation when assessed across ‘total community’ 16S rRNA gene profiles, but was significantly reduced in the actively growing community identified by qSIP. Similarly, compared to differential abundance analysis of 16S amplicons, qSIP identified tenfold more ASVs that were suppressed by water limitation (56 and 579, respectively). We attribute these differences to qSIP’s capacity to distinguish actively growing organisms from the rest of the microbial community.

Conclusion

As global precipitation patterns change and water limitation becomes more prevalent, it is important to understand biotic response and conditions that might mitigate associated declines in function. Mycorrhizal fungi support plant performance and alter biogeochemistry in droughted soils, but their effect on soil bacterial response to water limitation is poorly explored. With H₂¹⁸O qSIP, we show that mycorrhizal fungi protect bacterial communities from the suppressive and destabilizing effects of water limitation. Both the AM fungus *R. irregularis* and the Sebaciniales fungus *S. bescii* facilitated greater bacterial growth rates, growth efficiency, and the diversity of the active bacterial community in water-limited soil. While these divergent mycorrhizal lineages stimulated responses of differing magnitude, broad microbial patterns were similar, suggesting that the dominant underlying mechanisms are not limited to interactions between a small number of taxa. Remarkably, both *R. irregularis* and *S. bescii* prompted a

positive bacterial response despite the presence of a background microbial community in uninoculated soil. This finding is relevant for practical evaluation of mycorrhizal inoculants, whose ability to persist and elicit a positive effect in natural settings is not well-established. Together, our findings demonstrate that context-dependent mycorrhizal relationships support biotic resilience to moisture stress and may promote recovery once moisture is restored.

Acknowledgements

We thank Christine Hawkes for *P. hallii* seeds; Eric Slessarev for soil collection; Megan Foley, Anne Kakouridis, Benjamin Koch, and Alexa Nicolas for scientific discussions; Christina Ramon for procurement; Katerina Estera-Molina, Don Herman, Ilexis Chu Jacoby, Aaron Chew, Christina Fossum, Mengting Yuan, Tasnim Ahmed, Cynthia-Jeanette Mancilla, Melanie Rodriguez-Fuentes, David Sanchez, Laura Adame, Emily Kline, Madeline Moore, Jack Hagen, Anne Kakouridis, Ella Sieradzki, Nameer Baker, Sarah Baker, Alex Greenlon, Heejung Cho, Alexa Nicolas, Peter Weber, Vanessa Brisson, Eric Slessarev, Gareth Trubl, Dinesh Adhikari, Craig See, Maria Guerra, Rachel Neurath, and Evan Starr for their assistance with experimental set up, maintenance, and harvests; and Tibisay Perez and Whendee Silver for assistance with GC access and operation. This work was supported by the Genomic Science Program of the US Department of Energy Office of Biological and Environmental Research as part of the LLNL Biofuels SFA (SCW1039) and a sustainable bioenergy award (PI Firestone) to UC Berkeley (DE-SC0014079), Noble Research Institute, University of Oklahoma, Lawrence Berkeley National Laboratory, and Lawrence Livermore National Laboratory (SCW1555). Research at LLNL was performed under the auspices of the U.S. Department of Energy at Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.

497

498 **Competing Interests**

499 The authors declare no competing interests.

500

501 **References**

502 1. Leng G, Hall J. Crop yield sensitivity of global major agricultural countries to droughts and
503 the projected changes in the future. *Sci Total Environ* 2019; 654: 811–821.

504

505 2. Bouskill NJ, Lim HC, Borglin S, Salve R, Wood TE, Silver WL, et al. Pre-exposure to
506 drought increases the resistance of tropical forest soil bacterial communities to extended
507 drought. *ISME J* 2013; 7: 384–394.

508

509 3. Acosta-Martinez V, Cotton J, Gardner T, Moore-Kucera J, Zak J, Wester D, et al.
510 Predominant bacterial and fungal assemblages in agricultural soils during a record
511 drought/heat wave and linkages to enzyme activities of biogeochemical cycling. *Appl Soil
512 Ecol* 2014; 84: 69–82.

513

514 4. Yuste JC, Fernandez-Gonzalez AJ, Fernandez-Lopez M, Ogaya R, Penuelas J, Sardans J, et
515 al. Strong functional stability of soil microbial communities under semiarid Mediterranean
516 conditions and subjected to long-term shifts in baseline precipitation. *Soil Biol Biochem*
517 2014; 69: 223–233.

518

5. Hueso S, García C, Hernández T. Severe drought conditions modify the microbial community structure, size and activity in amended and unamended soils. *Soil Biol Biochem* 2012; 50: 167–173.
6. Alster CJ, German DP, Lu Y, Allison, SD. Microbial enzymatic responses to drought and to nitrogen addition in a southern California grassland. *Soil Biol Biochem* 2013; 64: 68–79.
7. Intergovernmental Panel on Climate Change (IPCC), Global Warming of 1.5°C, V. Masson Delmotte et al., Eds. (Cambridge Univ. Press, 2018).
8. O’Connell CS, Ruan L, Silver WL. Drought drives rapid shifts in tropical rainforest soil biogeochemistry and greenhouse gas emissions. *Nat Commun* 2018; 9: 1348.
9. Schimel J, Balser TC, Wallenstein M. Microbial stress-response physiology and its implications for ecosystem function. *Ecology* 2007; 88: 1386–1394.
10. Naylor D, Colemann-Derr D. Drought stress and root-associated bacterial communities. *Front Plant Sci* 2018; 8: 2223.
11. de Vries FT, Griffiths RI, Knight CG, Nicolitch O, Williams A. Harnessing rhizosphere microbiomes for drought-resilient crop production. *Science* 2020; 368: 270-274.

12. Smith SE, Read D. Mycorrhizal symbiosis. third ed. London: Academic Press; 2008. p. 145–190.
13. Kakouridis A, Hagen JA, Kan MP, Mambelli S, Feldman LJ, Herman DJ, et al. Routes to roots: direct evidence of water transport by arbuscular mycorrhizal fungi to host plants and identification of a hyphal transport pathway. bioRxiv <https://doi.org/10.1101/2020.09.21.305409>.
14. Rillig MC Mummey DL. Mycorrhizas and soil structure. New Phytol 2006; 171: 41-53.
15. Rillig MC, Lutgen ER, Ramsey PW, Klironomos JN, Gannon JE. Microbiota accompanying different arbuscular mycorrhizal fungal isolates influence soil aggregation. Pedobiologia 2005; 49: 251–259.
16. Gong M, You X, Zhang Q. Effects of *Glomus intraradices* on the growth and reactive oxygen metabolism of foxtail millet under drought. Ann Microbiol 2015; 65: 595–602.
17. Ruiz-Lozano JM. Arbuscular mycorrhizal symbiosis and alleviation of osmotic stress. New perspectives for molecular studies. Mycorrhiza 2003; 13: 309-317.
18. Morte A, Lovisolo C, Schubert A. Effect of drought stress on growth and water relations of the mycorrhizal association *Helianthemum almeriense*–*Terfezia clavervyi*. Mycorrhiza 2000; 10: 115–119.

19. Birhane E, Sterck F, Fetene M, Bongers F, Kuyper T. Arbuscular mycorrhizal fungi enhance photosynthesis, water use efficiency, and growth of frankincense seedlings under pulsed water availability conditions. *Oecologia* 2012; 169: 895–904.
20. Duan X, Neuman DS, Reiber JM, Green CD, Saxton AM, Augé RM. Mycorrhizal influence on hydraulic and hormonal factors implicated in the control of stomatal conductance during drought. *J Exp Bot* 1996; 47: 1541–1550.
21. Kaiser C, Kilburn MR, Clode PL, Fuchslueger L, Koranda M, Cliff JP, et al. Exploring the transfer of recent plant photosynthates to soil microbes: mycorrhizal pathway vs direct root exudation. *New Phytol* 2015; 205: 1537-1551.
22. Hodge A, Campbell CD, Fitter AH. An arbuscular mycorrhizal fungus accelerates decomposition and acquires nitrogen directly from organic matter. *Nature* 2001; 413: 297-299.
23. Medina A, Probanza A, Gutierrez Mañero FJ, Azcón R. Interactions of arbuscular-mycorrhizal fungi and *Bacillus* strains and their effects on plant growth, microbial rhizosphere activity (thymidine and leucine incorporation) and fungal biomass (ergosterol and chitin). *Appl Soil Ecol.* 2003; 22: 15–28.

24. Svenningsen NB, Watts-Williams SJ, Joner EJ. et al. Suppression of the activity of arbuscular mycorrhizal fungi by the soil microbiota. *ISME J* 2018; 12: 1296–1307.
25. Cruz-Paredes C, Svenningsen NB, Nybroe O, Kjølner R, Frøslev TG, Jakobsen I. Suppression of arbuscular mycorrhizal fungal activity in a diverse collection of non-cultivated soils. *FEMS Microbiol Ecol* 2019; 95: fiz020.
26. Nuccio EE, Hodge A, Pett-Ridge J, Herman DJ, Weber PK, Firestone MK. An arbuscular mycorrhizal fungus significantly modifies the soil bacterial community and nitrogen cycling during litter decomposition. *Environ Microbiol* 2013; 15: 1870–81.
27. Zhang L, Shi N, Fan J, Wang F, George TS, Feng G. Arbuscular mycorrhizal fungi stimulate organic phosphate mobilization associated with changing bacterial community structure under field conditions. *Environ Microbiol* 2018a; 20: 2639–51.
28. Emmet BD, Levesque-Tremblay V, Harrison MJ. Conserved and reproducible bacterial communities associate with extraradical hyphae of arbuscular mycorrhizal fungi. *ISME J* 2021; 15: 2276-2288.
29. Toljander JF, Artursson V, Paul LR, Jansson JK, Finlay RD. Attachment of different soil bacteria to arbuscular mycorrhizal fungal extraradical hyphae is determined by hyphal vitality and fungal species. *FEMS Microbiol Lett* 2006; 254: 34–40.

30. Yuan MM, Kakouridis A, Starr E, Nguyen N, Shi S, Pett-Ridge J, et al. Fungal-bacterial co-occurrence patterns differ between AMF and non-mycorrhizal fungi across space and time. *mBio* 2021; 12: e03509-20.
31. Carini P, Marsden P, Leff J, Morgan E, Strickland M, Fierer N. Relic DNA is abundant in soil and obscures estimates of soil microbial diversity. *Nat Microbiol* 2017; 2: 16242.
32. Lennon JT, Muscarella ME, Placella MA, Lehmkuhl BK. How, When, and Where Relic DNA Affects Microbial Diversity. *MBio* 2018; 9: e00637-18.
33. Hungate BA, Mau RL, Schwartz E, Caporaso JG, Dijkstra P, van Gestel N, et al. Quantitative microbial ecology through stable isotope probing. *Appl Environ Microbiol* 2015; 81: 7570–7581.
34. Neufeld JD, Vohra J, Dumont MG, Lueders T, Manefield M, Friedrich MW, Murrell JC. DNA stable-isotope probing. *Nat Protoc* 2007; 2: 860-866.
35. Koch BJ, McHugh TA, Hayer M, Schwartz E, Blazewicz SJ, Dijkstra P, et al. Estimating taxon-specific population dynamics in diverse microbial communities. *Ecosphere* 2018; 9: e02090–15.
36. Blazewicz SJ, Hungate BA, Koch BJ, Nuccio EE, Morrissey E, Brodie EL, Schwartz E, Pett-Ridge J, Firestone MK. Taxon-specific microbial growth and mortality patterns reveal

distinct temporal population responses to rewetting in a California grassland soil. ISME J
2020; 14: 1520-1532.

37. Kilronomos JN. Host-specificity and functional diversity among arbuscular mycorrhizal
fungi. In: Bell, C.R., Brylinski, M. and Johnson-Green, P., Eds., Proceedings of the 8th
International Symposium on Microbial Ecology, Atlantic Canada Society for Microbial
Ecology, Halifax, 2000; 845-851.

38. Ray P, Guo Y, Chi MH, Krom N, Saha MC, Craven KD. *Serendipita bescii* promotes winter
wheat growth and modulates the host root transcriptome under phosphorus and nitrogen
starvation. Environ Microbiol 2021; 23: 1876–1888.

39. Lee MR, Hawkes CV. Widespread co-occurrence of Sebaciniales and arbuscular mycorrhizal
fungi in switchgrass roots and soils has limited dependence on soil carbon or nutrients.
Plants, People, Planet 2021; 3: 614-626.

40. Ruiz-Lozano JM, Azcon R, Gomez M. Effects of arbuscular-mycorrhizal glomus species on
drought tolerance: physiological and nutritional plant responses. Appl Environ Microbiol
1995; 61: 456-460.

41. He F, Sheng M, Tang M. Effects of *Rhizophagus irregularis* on photosynthesis and
antioxidative enzymatic system in *Robinia pseudoacacia* L. under drought stress. Front Plant
Sci 2017; 8: 183.

42. Ghimire SR, Craven KD. Enhancement of switchgrass (*Panicum virgatum* L.) biomass production under drought conditions by the ectomycorrhizal fungus *Sebacina vermifera*. Appl Environ Microbiol 2011; 77: 7063-7067.
43. Tisserant E, Malbreil M, Kuo A, Kohler A, Symeonidi A, Balestrini R, et al. Genome of an arbuscular mycorrhizal fungus provides insight into the oldest plant symbiosis. Proc Natl Acad Sci USA 2013; 110: 20117-20122.
44. Kamel L, Keller-Pearson M, Roux C, Ané JM. Biology and evolution of arbuscular mycorrhizal symbiosis in the light of genomics. New Phytol 2017; 213: 531-536.
45. Zhang L, Feng G, Declerck S. Signal beyond nutrient, fructose, exuded by an arbuscular mycorrhizal fungus triggers phytate mineralization by a phosphate solubilizing bacterium. ISME J 2018; 12: 2339.
46. Hestrin R, Hammer EC, Mueller CW, Lehmann J. Synergies between mycorrhizal fungi and soil microbial communities increase plant nitrogen acquisition. Commun Biol 2019; 2: 233.
47. Zuccaro A, Lahrmann U, Guldener U, Langen G, Pfiffi S, Biedenkopf D, et al. Endophytic life strategies decoded by genome and transcriptome analyses of the mutualistic root symbiont *Piriformospora indica*. PLoS Pathog. 2011; 7: e1002290.

48. Ray P, Chi MH, Guo Y, Chen C, Adam C, Kuo A, et al. **Genome sequence of the plant growth promoting fungus *Serendipita vermifera* subsp. *bescii*: The first native strain from North America.** Phytobiomes J 2018; 2: 62-63.
49. Dias T, Pimentel V, Cogo AJD, Costa R, Bertolazi AA, Miranda C, et al. The free-living stage growth conditions of the endophytic fungus *Serendipita indica* may regulate its potential as plant growth promoting microbe. Front Microbiol 2020; 11: 562238.
50. Moffatt HH. Soil Survey of Caddo County, Oklahoma. United States Department of Agriculture Soil Conservation Service, Washington, D.C. 1973.
51. Sher Y, Baker NR, Herman NR, Fossum C, Hale L, Zhang XX, et al. Microbial extracellular polysaccharide production and aggregate stability controlled by Switchgrass (*Panicum virgatum*) root biomass and soil water potential. Soil Biol Biochem 2020; 143: 107742.
52. Seki K. SWRC fit - a nonlinear fitting program with a water retention curve for soils having unimodal and bimodal pore structure. Hydrol Earth Syst Sci Discuss 2007; 4: 407-437.
53. Ray P, Ishiga T, Decker SR, Turner GB, Craven KD. A novel delivery system for the root symbiotic fungus, *Sebacina vermifera*, and consequent biomass enhancement of low lignin COMT switchgrass lines. BioEnergy Res 2015; 8: 922–933.

54. Badri DV, Chaparro JM, Zhang R, Shen Q, Vivanco JM. Application of natural blends of phytochemicals derived from the root exudates of *Arabidopsis* to the soil reveal that phenolic-related compounds predominantly modulate the soil microbiome. *J Biol Chem* 2013; 288: 4502–4512.
55. Gamper HA, Young JP, Jones DL, Hodge A. Real-time PCR and microscopy: are the two methods measuring the same unit of arbuscular mycorrhizal fungal abundance? *Fungal Genet Biol* 2008; 45: 581-96.
56. Tellenbach C, Grünig CR, Sieber TN. Suitability of quantitative real-time PCR to estimate the biomass of fungal root endophytes. *Appl Environ Microbiol* 2010; 76: 5764-5772.
57. Blazewicz SJ, Schwartz E, Firestone MK. Growth and death of bacteria and fungi underlie rainfall-induced carbon dioxide pulses from seasonally dried soil. *Ecology* 2014; 95: 1162-1172.
58. Sieradzki ET, Koch BJ, Greenlon A, Sachdeva R, Malmstrom RR, Mau RL, et al. Measurement error and resolution in quantitative stable isotope probing: implications for experimental design. *mSystems* 2020; 5: e00151-20.
59. Buckley DH, Huangyutitham V, Hsu SF, Nelson TA. Stable isotope probing with ¹⁵N achieved by disentangling the effects of genome G+C content and isotope enrichment on DNA density. *Appl Environ Microbiol* 2007; 73: 3189-3195.

60. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, et al. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc Natl Acad Sci USA* 2011; 108: 4516-4522.
61. Schildkraut CL, Marmur J, Doty P. Determination of the base composition of deoxyribonucleic acid from its buoyant density in CsCl. *J Mol Biol* 1962; 4: 430–43.
62. R Core Team. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing; 2019.
63. Manzoni S, Taylor P, Richter A, Porporato A, Ågren GI. Environmental and stoichiometric controls on microbial carbon-use efficiency in soils. *New Phytol* 2012; 196: 79-91.
64. Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, OHara RB, et al. *vegan*: Community Ecology Package R package version 2.3-0. 2015. Available from <http://CRAN.R-project.org/package=vegan>.
65. Lozupone C, Lladser ME, Knights D, Stombaugh J, Knight R. UniFrac: an effective distance metric for microbial community comparison. *ISME J* 2011; 5: 169–172.
66. Paulson JN, Stine OC, Bravo HC, Pop M. Differential abundance analysis for microbial marker-gene surveys. *Nat Meth* 2013; 10: 1200-1202.

67. Wagg C, Dudenhöffer JH, Widmer F, van der Heijden MGA. Linking diversity, synchrony and stability in soil microbial communities. *Funct Ecol* 2018; 32: 1280–1292.
68. Tiemann LK, Billings SA. Changes in variability of soil moisture alter microbial community C and N resource use. *Soil Biol Biochem* 2011; 43: 1837–1847.
69. Domeignoz-Horta LA, Pold G, Liu XJA, Frey SD, Melillo JM, DeAngelis KM. Microbial diversity drives carbon use efficiency in a model soil. *Nat Commun* 2020; 11: 3684.
70. Malik AA, Martiny JBH, Brodie EL, Martiny AC, Treseder KK, Allison SD. Defining trait-based microbial strategies with consequences for soil carbon cycling under climate change. *ISME J* 2020; 14: 1–9.
71. Gefen O, Balaban NQ. The importance of being persistent: heterogeneity of bacterial populations under antibiotic stress. *FEMS Microbiol Rev* 2009; 33: 704–717.
72. Fridman O, Goldberg O, Ronin I, Shores N, Balaban NQ. Optimization of lag time underlies antibiotic tolerance in evolved bacterial populations. *Nature* 2014; 513: 418–421.
73. Bouskill NJ, Wood TE, Baran R, Hao Z, Ye Z, Bowen BP, et al. Belowground response to drought in a tropical forest soil. II. Change in microbial function impacts carbon composition. *Front Microbiol* 2016; 7: 323.

769

770 74. Stark JM, Firestone MK. Mechanisms for soil moisture effects on activity of nitrifying

771 bacteria. *Appl Environ Microbiol* 1995; 61: 218–221.

772

773 75. Liang C, Amelung W, Lehmann J, Kästner M. Quantitative assessment of microbial

774 necromass contribution to soil organic matter. *Glob Change Biol* 2019; 25: 3578–3590.

775

776 76. Sutcliffe IC. A phylum level perspective on bacterial cell envelope architecture. *Trends*

777 *Microbiol* 2010; 18: 464-470.

778

779 77. Xu L, Naylor D, Dong Z, Simmons T, Pierroz G, Hixson KK, et al. Drought delays

780 development of the sorghum root microbiome and enriches for monoderm bacteria. *Proc Natl*

781 *Acad Sci USA* 2018; 115: E4284-E4293.

782

783 78. Santos-Medellín C, Liechty Z, Edwards J, Nguyen B, Huang B, Weimer BC, Sundaresan V.

784 Prolonged drought imparts lasting compositional changes to the rice root microbiome. *Nat*

785 *Plants* 2021; 7, 1065–1077.

786

787 79. Tocheva EI, Ortega DR, Jensen GJ. Sporulation, bacterial cell envelopes and the origin of

788 life. *Nat Rev Microbiol* 2016; 14: 535–542.

789

80. Otoguro M, Yamamura H, Quintana ET. (2014) The Family Streptosporangiaceae. In:
Rosenberg E., DeLong E.F., Lory S., Stackebrandt E., Thompson F. (eds) The Prokaryotes.
Springer, Berlin, Heidelberg.
81. Nuccio EE, Starr E, Karaoz U, Brodie EL, Zhou J, Tringe SG, et al. Niche differentiation is
spatially and temporally regulated in the rhizosphere. ISME J 2020; 14: 999–1014.
82. Barnard RL, Osborne CA, Firestone MK. Responses of soil bacterial and fungal communities
to extreme desiccation and rewetting. ISME J 2013; 7: 2229–2241.
83. Guhr A, Borken W, Spohn M, Matzner E. Soil water redistribution by a saprotrophic fungus.
PNAS 2015; 112: 12647-14651.
84. Guhr A, Marzini C, Borken W, Poll C, Matzner E. Effect of water redistribution by two
distinct saprotrophic fungi on carbon mineralization and nitrogen translocation in dry soil.
Soil Biol Biochem 2016; 103: 380-387.
85. Drigo B, Pijl AS, Duyts H, Kielak AM, Gamper HA, Houtekamer MJ, et al. Shifting carbon
flow from roots into associated microbial communities in response to elevated atmospheric
CO₂. Proc Natl Acad Sci USA 2010; 107: 10938–10942.
86. Cruz AF, Ishii T. Arbuscular mycorrhizal fungal spores host bacteria that affect nutrient
biodynamics and biocontrol of soil-borne plant pathogens. Biol Open 2012; 1: 52-57.

87. Hernandez, D.J., David, A.S., Menges, E.S. *et al.* Environmental stress destabilizes microbial networks. ISME J 2021; 15: 1722–1734.
88. Bukovská P, Bonkowski M, Konvalinková T, Beskid O, Hujslová M, Püschel D et al. Utilization of organic nitrogen by arbuscular mycorrhizal fungi-is there a specific role for protists and ammonia oxidizers? Mycorrhiza 2018; 28: 269-283.
89. Leigh J, Fitter AH, Hodge A. Growth and symbiotic effectiveness of an arbuscular mycorrhizal fungus in organic matter in competition with soil bacteria. FEMS Microbiol Ecol 2011; 76: 428–438.
90. Bronstein JL. Conditional outcomes in mutualistic interactions. Trends Ecol Evol 1994; 9: 214–217.
91. Toljander JF, Lindahl BD, Paul LR, Elfstrand M, Finlay RD. Influence of arbuscular mycorrhizal mycelial exudates on soil bacterial growth and community structure. FEMS Microbiol Ecol 2007; 61: 295–304.
92. Zhalnina K, Louie KB, Hao Z, Mansoori N, Nunes da Rocha U, Shi S, et al. Dynamic root exudate chemistry and microbial substrate preferences drive patterns in rhizosphere microbial community assembly. Nat Microbiol 2018; 3: 470–480.

93. Chaparro JM, Badri DV, Bakker MG, Sugiyama A, Manter DK, Vivanco JM. Root exudation of phytochemicals in *Arabidopsis* follows specific patterns that are developmentally programmed and correlate with soil microbial functions. PLoS One 2013; 8: e55731.
94. Shi S, Richardson AE, O'Callaghan M, DeAngelis KM, Jones EE, Stewart A, et al. Effects of selected root exudate components on soil bacterial communities. FEMS Microbiol Ecol 2011; 77: 600–610.
95. Schaeffer SM, Homyak PM, Boot CM, Roux-Michollet D, Schimel JP. Soil carbon and nitrogen dynamics throughout the summer drought in a California annual grassland. Soil Biol Biochem 2017; 115: 54–62.
96. Fuchslueger L, Bahn M, Fritz K, Hasibeder R, Richter A. Experimental drought reduces the transfer of recently fixed plant carbon to soil microbes and alters the bacterial community composition in a mountain meadow. New Phytol 2014; 201: 916–927.
97. Davidson EA, Belk E, Boone RD. Soil water content and temperature as independent or confounded factors controlling soil respiration in a temperate mixed hardwood forest. Glob Change Biol 1998; 4: 217–227.

98. Herron PM, Stark JM, Holt C, Hooker T, Cardon ZG. Microbial growth efficiencies across a soil moisture gradient assessed using ¹³C-acetic acid vapor and ¹⁵N-ammonia gas. *Soil Biol Biochem* 2009; 41: 1262–69.

99. Schimel JP. Life in dry soils: Effects of drought on soil microbial communities and processes. *Annu Rev Ecol Evol Syst* 2018; 49: 409–432.

100. NOAA (National Oceanic and Atmospheric Administration). National Climatic Data Center. 2021. <https://www.ncdc.noaa.gov/>

Figures

Figure 1. Experimental design and effect of mycorrhizal inoculation and moisture

manipulation on mycorrhizal abundance and soil moisture after 3 months. Greenhouse

experiment with *Panicum hallii* (Hall's panicgrass) and two mycorrhizal inocula grown in a

marginal soil from central Oklahoma. **a** Illustration of microcosm experimental design and $H_2^{18}O$

stable isotope probing (SIP) incubation experiment conducted with soils collected from hyphal

ingrowth cores that excluded plant roots. *P. hallii* was inoculated with either *R. irregularis*, *S.*

bescii or left uninoculated (indicated in yellow, blue, or black, respectively) and grown for 3

months under water-replete or water-limited conditions (indicated in solid or dashed lines,

respectively; $n = 3$ replicates per treatment). Soils harvested at 3 months were mixed with either

enriched ($H_2^{18}O$) or natural abundance ($H_2^{16}O$) water and incubated for 7 days. **b** Abundance of

R. irregularis and *S. bescii* (DNA copies g^{-1} soil) in hyphal ingrowth cores after 3 months of

plant growth and moisture manipulation, measured with strain-specific qPCR primers. **c** Soil moisture after 3-monts. Bold lines represent median value; whiskers represent upper and lower quartiles (n = 3 replicates per treatment). Letters denote the results of a Wilcoxon rank sum test performed separately for each qPCR plot ($p < 0.1$) and a Tukey's HSD test for soil moisture comparisons ($p < 0.001$).

Figure 2. Structure and alpha diversity of total and active microbial communities based on traditional and H₂¹⁸O qSIP-filtered 16S rRNA gene profiling. Principal coordinates analysis (PCoA) of weighted UniFrac distances between ‘total’ microbial communities assessed with (a) traditional 16S rRNA gene analysis and (b) H₂¹⁸O qSIP-filtered ‘actively growing’ communities (i.e., ASVs that did not incorporate a significant quantity of ¹⁸O were removed prior to analysis). Moisture regime and mycorrhizal inoculum explained a total of 29.5% of the variation in ‘total’ community structure (p < 0.001; n = 6 replicates) and 75.2% of variation in actively growing community structure (p < 0.001; n = 6 replicates). Ellipses show treatment groupings in (a) and (b). c Inverse Simpson’s diversity index in total and actively growing communities. Letters denote the results of a Tukey’s HSD test (no significant differences between total communities; p < 0.01 for comparisons between active communities; n = 6 replicates). Bold lines represent median values; whiskers represent upper and lower quartiles. For all plots, uninoculated, *R. irregularis*-inoculated, and *S. bescii*-inoculated soils are represented in black, yellow, and blue, respectively. Water-replete and water-limited soils are represented in dark versus light color shades, respectively.

Figure 3. H_2^{18}O qSIP-based microbial activity in hyphal ingrowth core soils exposed to different moisture regimes and mycorrhizal inocula. a Median ^{18}O APE of all bacterial and archaeal ASVs present under each treatment. Uninoculated, *R. irregularis*-inoculated, and *S. bescii*-inoculated soils are represented in black, yellow, and blue, respectively. Water-replete and water-limited soils are represented in solid and dashed lines, respectively. Letters denote the results of a Tukey's HSD test comparing the means of all treatments ($p < 0.01$; $n = 3$ replicates). **b** Taxon-specific gross growth rates (ng DNA g^{-1} soil day^{-1}) based on ASV ^{18}O incorporation,

920 summed by phylum for each treatment. **c** CO₂ efflux (μg CO₂ mg⁻¹ soil C day⁻¹) from hyphal
 921 ingrowth core soils during H₂¹⁸O qSIP assay. Bold lines represent median values; whiskers
 922 represent upper and lower quartiles (p < 0.05; n = 6 replicates per treatment).
 923
 924
 925

Figure 4. Comparison of taxon-specific activity in water-limited versus water-replete hyphal ingrowth core soils within each mycorrhizal treatment. ¹⁸O atom percent excess (APE) ‘active growth’ ratios of ASVs present in both water-limited and water-replete soils that were either (a) uninoculated, (b) inoculated with *R. irregularis*, or (c) inoculated with *S. bescii*. Ratios were averaged at the phylum level. Ratios less than 1 (bars located to the left of the central line) indicate that water limitation suppressed growth. Ratios greater than 1 (bars located to the right of the central line) indicate that water limitation facilitated greater growth rates. Error bars represent the standard error. Asterisks denote phylum-level averages that are significantly

greater than or less than 1 (adjusted $p < 0.05$ based on Wilcoxon signed rank test and Benjamini-Hochberg correction for multiple comparisons). Only taxa that incorporated significant ^{18}O (lower 90% CI > 0) and belong to the 10 most abundant phyla are represented.

Figure 5. Comparison of taxon-specific activity in mycorrhizal-inoculated versus uninoculated hyphal ingrowth core soils following water limitation. ^{18}O atom percent excess (APE) ‘active growth’ ratios of bacterial ASVs present in water-limited soils that were either uninoculated or inoculated with *R. irregularis* or *S. bescii*, averaged at the phylum level (**a,b**) or

displayed by individual ASV (**c,d**). Ratios greater than 1 (to the right of the central line) indicate that mycorrhizal fungi supported greater bacterial growth rates. For phylum level averages, asterisks represent ^{18}O APE ratios that are significantly greater than 1 (adjusted $p < 0.05$ based on Wilcoxon signed rank test and Benjamini-Hochberg correction for multiple comparisons). Error bars represent the standard error within each bacterial phylum. Only taxa that incorporated significant ^{18}O (lower 90% CI > 0) and belong to the 10 most abundant phyla are represented. ASVs belonging to phyla that incorporated significantly more ^{18}O in mycorrhizal-inoculated soils are represented in **c,d**.